LMSM: a modular approach for identifying lncRNA related miRNA sponge modules in breast cancer

Junpeng Zhang[1,2,*,†], Taosheng Xu[3,†], Lin Liu[4], Wu Zhang[5], Chunwen Zhao[2], Sijing Li[2], Jiuyong Li[4], Nini Rao[1,*] and Thuc Duy Le[4,*]

[1]Center for Informational Biology, School of Life Science and Technology, University of Electronic Science and Technology of China, Chengdu, Sichuan 610054, China

[2]School of Engineering, Dali University, Dali, Yunnan 671003, China

[3]Institute of Intelligent Machines, Hefei Institutes of Physical Science, Chinese Academy of Sciences, Hefei, Anhui 230031, China

[4]School of Information Technology and Mathematical Sciences, University of South Australia, Mawson Lakes, SA 5095, Australia

[5]School of Agriculture and Biological Sciences, Dali University, Dali, Yunnan 671003, China

*To whom correspondence should be addressed. Email: zhangjunpeng_411@yahoo.com

Correspondence may also be addressed to Nini Rao. Email: raonn@uestc.edu.cn

Correspondence may also be addressed to Thuc Duy Le. Email: thuc.le@unisa.edu.au

[†]The authors wish it to be known that, in their opinion, the first two authors should be regarded as joint First Authors.

Abstract

Until now, existing methods for identifying lncRNA related miRNA sponge modules mainly rely on lncRNA related miRNA sponge interaction networks, which may not provide a full picture of miRNA sponging activities in biological conditions. Hence there is a strong need

of new computational methods to identify lncRNA related miRNA sponge modules. In this

work, we propose a framework, LMSM, to identify LncRNA related MiRNA Sponge

Modules from heterogeneous data. To understand the miRNA sponging activities in

biological conditions, LMSM uses gene expression data to evaluate the influence of the

shared miRNAs on the clustered sponge lncRNAs and mRNAs. We have applied LMSM to

the human breast cancer (BRCA) dataset from The Cancer Genome Atlas (TCGA). As a

result, we have found that the majority of LMSM modules are significantly implicated in

BRCA and most of them are BRCA subtype-specific. Most of the mediating miRNAs act as

crosslinks across different LMSM modules, and all of LMSM modules are statistically

significant. Multi-label classification analysis shows that the performance of LMSM modules

is significantly higher than baseline's performance, indicating the biological meanings of

LMSM modules in classifying BRCA subtypes. The consistent results suggest that LMSM is

robust in identifying lncRNA related miRNA sponge modules. Moreover, LMSM can be

used to predict miRNA targets. Finally, LMSM outperforms a graph clustering-based strategy

in identifying BRCA-related modules. Altogether, our study shows that LMSM is a

promising method to investigate modular regulatory mechanism of sponge lncRNAs from

heterogeneous data.

Author summary

Previous studies have revealed that long non-coding RNAs (lncRNAs), as microRNA

(miRNA) sponges or competing endogenous RNAs (ceRNAs), can regulate the expression

levels of messenger RNAs (mRNAs) by decreasing the amount of miRNAs interacting with

mRNAs. In this work, we hypothesize that the "tug-of-war" between RNA transcripts for

attracting miRNAs is across groups or modules. Based on the hypothesis, we propose a

framework called LMSM, to identify LncRNA related MiRNA Sponge Modules. Based on

the two miRNA sponge modular competition principles, significant sharing of miRNAs and

high canonical correlation between the sponge lncRNAs and mRNAs, LMSM is also capable

of predicting miRNA targets. LMSM not only extends the ceRNA hypothesis, but also

provides a novel way to investigate the biological functions and modular mechanism of

lncRNAs in breast cancer.

Introduction

Long non-coding RNAs (lncRNAs) are RNA transcripts with more than 200 nucleotides (nts)

in length [1]. More and more evidence has shown that lncRNAs play important functional

roles in many biological processes, including human cancers [2-4]. As a major class of non-

coding RNAs (ncRNAs), lncRNAs have attracted increasing interest from researchers in their

exploration of non-coding knowledge from the 'junk'.

Among the wide range of biological functions of lncRNAs, their role as competing

endogenous RNAs (ceRNAs) or miRNA sponges is in the limelight. As a family of small

ncRNAs (~18nts in length), miRNAs are important post-transcriptional regulators of gene

expression [5,6]. According to the ceRNA hypothesis [7], lncRNAs contain abundant miRNA

response elements (MREs) for competitively sequestering target mRNAs from miRNAs'

control. This regulation mechanism of lncRNAs when acting as miRNA sponges is highly

implicated in various human diseases [8], including breast cancer [9]. For example, lncRNA

H19, an imprinted gene is associated with breast cancer cell clonogenicity, migration and

mammosphere-forming ability. By sponging miRNA *let-7*, *H19* forms a *H19/let-7/LIN28*

reciprocal negative regulatory circuit to play a critical role in the breast cancer stem cell

maintenance [10].

To systematically investigate the functions of lncRNAs as miRNA sponges in human

cancer, a series of computational methods have been developed to infer lncRNA related

miRNA sponge interaction networks. The methods can be divided into three categories

according to the statistical or computational techniques employed: pair-wise correlation based

approach, partial association based approach, and mathematical modelling approach [11].

It is commonly known that to implement a specific biological function, genes tend to

cluster or connect in the form of modules or communities. Consequently, based on the

identified lncRNA related miRNA sponge interaction networks, several methods [12-17]

using graph clustering algorithms were developed to identify lncRNA related miRNA sponge

modules. For the identification of sponge lncRNA-mRNA pairs, most of existing methods

only consider pair-wise correlation of them. Since the lncRNA related miRNA sponge

interaction networks are created by simply putting together sponge lncRNA-mRNA pairs,

when the expression levels of each sponge lncRNA-mRNA pair are highly correlated, the

collective correlation between the set of sponge lncRNAs and the set of mRNAs in the same

identified module is not necessarily high. As we know, the pair-wise positive correlation

between the expression levels of a lncRNA and a mRNA pair is commonly used to identify

the sponge interactions between them. For the identification of lncRNA related miRNA

sponge modules, it is also necessary to investigate whether the clustered sponge lncRNAs and

mRNAs in a module have high collective positive correlation or not. Moreover, these

methods do not consider the influence of the shared miRNAs on the expression of the

clustered sponge lncRNAs and mRNAs. It is known that the "tug-of-war" between sponge

lncRNAs and mRNAs is mediated by miRNAs. Therefore, it is extremely important to

consider the influence of the shared miRNAs in identifying lncRNA related miRNA sponge

modules.

Recently, to study lncRNA, miRNA and mRNA-associated regulatory modules, Deng *et*

al. [18] and Xiao *et al*. [19] have proposed two types of joint matrix factorization methods to

identify mRNA-miRNA-lncRNA co-modules by integrating gene expression data and

putative miRNA-target interactions. However, it is still not clear how the shared miRNAs

influence the expression level of the sponge lncRNAs and mRNAs in a module.

To address the above issues, we firstly hypothesize that sponge lncRNAs form a group to

competitively release a group of target mRNAs from the control of the miRNAs shared by

the lncRNAs and mRNAs (details see Section Methods). We name this hypothesis the *miRNA*

sponge modular competition hypothesis in this paper. Then based on the hypothesis, we

propose a novel framework to identify LncRNA related MiRNA Sponge Modules (LMSM).

The framework firstly uses the WeiGhted Correlation Network Analysis (WGCNA) [20]

method to generate lncRNA-mRNA co-expression modules. Next, by incorporating matched

miRNA expression and putative miRNA-target interactions, LMSM applies three constraints

(see Section Methods) to obtain lncRNA related miRNA sponge modules (also called LMSM

modules in this paper). One of the constraints, high canonical correlation, is used to assess

whether the group of sponge lncRNAs and the group of mRNAs in the same module have a

high collective positive correlation or not. The other constraint, adequate sensitivity canonical

correlation conditioning on a group of miRNAs, is used to evaluate the influence of the

shared miRNAs on the clustered sponge lncRNAs and mRNAs.

To evaluate the LMSM approach, we apply it to matched miRNA, lncRNA and mRNA

expression data, and clinical information of breast cancer (BRCA) dataset from The Cancer

Genome Atlas (TCGA). The modular analysis results demonstrate that LMSM can help to

uncover modular regulatory mechanism of sponge lncRNAs in BRCA. LMSM is released

under the GPL-3.0 License, and is freely available through GitHub repository

(https://github.com/zhangjunpeng411/LMSM).

Materials and methods

A hypothesis on miRNA sponge modular competition

The ceRNA hypothesis [7] indicates that a pool of RNA transcripts (known as ceRNAs)

regulate each other's transcripts by competing for the shared miRNAs through MREs. Based

on this unifying hypothesis, a large-scale gene regulatory network including coding and non-

coding RNAs across the transcriptome can be formed, and it plays critical roles in human

physiological and pathological processes. However, by using MREs as letters of language,

the hypothesis only depicts the crosstalk between individual RNA transcript (e.g. coding

RNAs, lncRNAs, circRNAs or pseudogenes) and mRNA at the pair-wise interaction level

and the crosstalk between RNA transcripts and mRNAs at the module level is still an open

question.

There has been evidence showing that for the same transcriptional regulatory program,

biological process or signaling pathway, genes tend to form modules or communities to

coordinate biological functions [21]. These modules correspond to functional units in

complex biological systems, and they play important role in gene regulation. Based on these

findings, in this paper, we hypothesize that regarding miRNA sponging, the crosstalk

between different RNA transcripts is in the form of modular competition. We call the

hypothesis the *miRNA sponge modular competition hypothesis*.

As shown in Fig 1, based on our hypothesis, instead of having pair-wise competitions,

miRNA sponges form groups to compete at module level for common miRNAs. Here, a

miRNA sponge module consists of a competing group (other coding RNA group, pseudogene

group, circRNA group or lncRNA group) and a mRNA group. Here, other coding RNAs also

include mRNAs. From the perspective of modularity, the hypothesis at module level extends

the ceRNA hypothesis and provides a new channel to look into the functions and regulatory

mechanism of miRNA sponges or ceRNAs. Since the available resources of lncRNAs are

more abundant than those of other non-coding RNAs (e.g. circRNAs and pseudogenes), in

this paper, we focus on the competition between lncRNAs and mRNAs to validate and

demonstrate the proposed *miRNA sponge modular competition hypothesis*. Our goal is to

discover lncRNA related sponge modules, or LMSM modules. Here each LMSM module

contains a group of lncRNAs which compete collectively with a group of mRNAs for

sponging the same set of miRNAs.

148
Fig 1. An illustration of the miRNA sponge modular competition hypothesis. The four
types of miRNA sponges (other coding RNAs, lncRNAs, circRNAs or pseudogenes),
miRNAs and their target mRNAs are shown. Each miRNA sponge module consists of a

group of the same type of miRNA sponges, e.g. a group of lncRNAs and a group of target
mRNAs. In the same module, the group of miRNA sponges competes with the group of
target mRNAs for binding with a set of miRNAs. If the miRNA sponges win the competition,
the group of target mRNAs will be released from repression and they will be translated into
proteins. If the miRNA sponges lose the competition, the group of target mRNAs will be
post-transcriptionally repressed and degraded.

The LMSM framework

Overview of LMSM. As shown in Fig 2, the proposed LMSM framework comprises two

stages. In stage 1, the WGCNA method [20] is used for finding lncRNA-mRNA co-

expression modules from matched lncRNA and mRNA expression data. Then in stage 2,

LMSM identifies LMSM modules from the lncRNA-mRNA co-expression modules using

three criteria. That is, a co-expression module is considered as a LMSM module if the group

of lncRNAs and the group of mRNAs in the co-expression module: (1) have significant

sharing of miRNAs, (2) have high canonical correlation between their expression levels, and

(3) have adequate sensitivity canonical correlation conditioning on their shared miRNAs.

LMSM checks the criteria one by one, and once a co-expression module does not meet a

criterion, it is discarded and will not be checked for the next criterion. In the following, we

will describe the two stages in detail.

Identifying lncRNA-mRNA co-expression modules. For identifying lncRNA-mRNA co-

expression modules, we use the WGCNA method. WGCNA is a popular method for

identifying co-expressed genes across samples and it can be used to identify clusters of highly

co-expressed lncRNAs and mRNAs. In our task, we use the matched lncRNA and mRNA

expression data as input to the *WGCNA* R package [20] to identify lncRNA-mRNA co-

expression modules. We use the scale-free topology criterion for soft thresholding. The

coefficient of determination R^2 (the range is from 0 to 1) is used to quantify the goodness of

scale-free topology, and larger R^2 values mean better scale-free topology. Normally, the R^2

value larger than 0.8 in power law curve fit is ranked as good-level in the WGCNA method.

Therefore, the desired minimum scale free topology fitting index R^2 is set as 0.8 in this work.

180
Fig 2. Workflow of LMSM. Firstly, we use the WGCNA method to infer lncRNA-mRNA
co-expression modules from the matched lncRNA and mRNA expression. Then by using
miRNA expression data and putative miRNA-target interactions, we infer lncRNA related
miRNA sponge modules (LMSM) by applying three criteria: significant sharing of miRNAs
by the group of lncRNAs and the group of target mRNAs in the same co-expression module,
high canonical correlation between the lncRNA group and the target mRNA group, and
adequate sensitivity canonical correlation between the lncRNA group and the target mRNA
group conditioning on shared miRNAs. Each LMSM module must contain at least two
sponge lncRNAs and two target mRNAs.

Inferring lncRNA related miRNA sponge modules. To identify lncRNA related miRNA

sponge modules from the co-expression modules obtained in stage 1, we propose three

criteria (detailed below) by following the key tenet of our *miRNA sponge modular*

competition hypothesis. That is, a group of lncRNAs (acting as miRNA sponges) competes

with a group of mRNAs with respect to a set of miRNAs shared by the two groups.

The first criterion requires that the group of lncRNAs and the group of mRNAs in a

miRNA sponge module have a significant sharing of a set of miRNAs. LMSM uses a

hypergeometric test to assess the significance of the sharing of miRNAs between the group of

lncRNAs and the group of mRNAs in a co-expression module, based on putative miRNA-

target interactions. The *p*-value for the test is computed as:

200
$$p-value = 1 - \sum_{i_1=0}^{L_1-1} \frac{\binom{M_1}{i_1}\binom{N_1-M_1}{K_1-i_1}}{\binom{N_1}{K_1}}$$
(1)

In the equation, N_1 is the number of all miRNAs in the dataset, M_1 and K_1 denote the total

numbers of miRNAs interacting with the group of lncRNAs and the group of mRNAs in the

co-expression module respectively, and L_1 (e.g. 3) is the number of miRNAs shared by the

group of lncRNAs and the group of mRNAs in the co-expression module.

The second criterion is to assure that the sponge modular competition between the group

of lncRNAs and the group of mRNAs in a miRNA sponge module is strong enough. In

existing work, to identify lncRNA related mRNA sponge interactions, a principle followed is

that the expression level of a lncRNA and the expression level of a mRNA need to be

strongly and positively correlated. Following the same principle on strong positive correlation

in expression levels while considering our modular competition hypothesis, LMSM requires

the *collective* correlation between the expression levels of the group of lncRNAs and the

group of target mRNAs in the same module to be strong and positive. To assess the *collective*

correlation, we perform canonical correlation analysis [22] to obtain the canonical correlation

between the group of lncRNAs and the group of mRNAs in a co-expression module. Let the

two column vectors $X = (x_1, x_2, ..., x_m)^T$ and $Y = (y_1, y_2, ..., y_n)^T$ represent the group of

lncRNAs and the group of mRNAs in a co-expression module respectively. Σ_{XX}, Σ_{YY} and

Σ_{XY} are the variance or cross-covariance matrices calculated from the expression data of X

and Y. The canonical correlation analysis seeks the canonical vectors a ($a \in \mathbb{R}^m$) and b

($b \in \mathbb{R}^n$) which maximize the correlation of $corr(a^T X, b^T Y)$. The canonical correlation

between the group of lncRNAs and the group of mRNAs, denoted as $CC_{lncR\text{-}mR}$, is then

calculated as follows with the found canonical vectors:

$$CC_{lncR-mR} = corr(a^T X, b^T Y) = \frac{a^T \sum_{XY} b}{\sqrt{a^T \sum_{XX} a} \sqrt{b^T \sum_{YY} b}} \tag{2}$$

In this work, we use the *PMA* R package [23] to compute canonical correlation.

Finally, the third criterion adapted from the sensitivity correlation [24] is employed to

assess if the miRNAs shared by the group lncRNAs and the group of mRNAs in a module

have large enough influence on the modular competition between the two groups of RNAs.

To check according to this criterion, we incorporate miRNA expression data, and compute

$SCC_{lncR\text{-}mR}$ the sensitivity canonical correlation between the group of lncRNAs and the group

of mRNAs in a co-expression module as follows:

$$SCC_{lncR-mR} = CC_{lncR-mR} - PCC_{lncR-mR} \tag{3}$$

where $PCC_{lncR\text{-}mR}$ is the partial canonical correlation between the group of lncRNAs and the

group of mRNAs, i.e. the canonical correlation conditioning on the expression of their shared

miRNAs in the co-expression module, or the canonical correlation between the two groups of

RNAs when the influence of the shared miRNAs is eliminated. Therefore, from Eq. (3), we

see that $SCC_{lncR\text{-}mR}$ implies the correlation between the two groups of RNAs under the

influence of their shared miRNAs.

$PCC_{lncR\text{-}mR}$ in Eq. (4) can be calculated as:

238
$$PCC_{lncR-mR} = \frac{CC_{lncR-mR} - CC_{miR-mR}CC_{miR-lncR}}{\sqrt{1-CC_{miR-mR}^2}\sqrt{1-CC_{miR-lncR}^2}} \tag{4}$$

where $CC_{miR\text{-}mR}$ ($CC_{miR\text{-}lncR}$) is the canonical correlation between the set of miRNAs in the co-

expression module and the group of mRNAs (lncRNAs) in the co-expression module.

In this study, empirically, a lncRNA-mRNA co-expressed module with p-value < 0.05 for

the hypergeometric test of miRNA sharing (criterion 1), $CC_{lncR\text{-}mR} > 0.8$ for modular

competition strength assessment (criterion 2) and $SCC_{lncR\text{-}mR} > 0.1$ for miRNA influence

(criterion 3) is regarded as a lncRNA related miRNA sponge module (a LMSM module).

Evaluating statistical significance of LMSM modules

To evaluate the statistical significance of LMSM modules, we adapt the null model method

proposed in [25]. The null model method hypothesizes that the shared miRNAs do not affect

the correlation between two genes, i.e. the sensitivity correlation (the difference between

correlation and partial correlation) between two genes is 0, and has been successfully applied

to evaluate statistical significance of ceRNA interactions. Similar to [25], LMSM is also

adapted from the Sensitivity Correlation (SC) method [24]. Therefore, the null model method

can be applied to evaluate the statistical significance of LMSM modules. In our null model,

the null hypothesis is that the group of the shared miRNAs does not influence the canonical

correlation between the group of lncRNAs and the group of mRNAs, i.e. $SCC_{lncR\text{-}mR} = 0$. For

each LMSM module, a group of lncRNAs or a group of mRNAs corresponds to a gene, and a

group of the shared miRNAs corresponds to a miRNA in the null model. For obtaining more

precise p-values, the number of datasets sampled is set to 1E+06 for the null model. Since the

sampling procedure is computationally intensive, we use the pre-computed sets of covariance

matrices in *SPONGE* R package [25] to build our null model. Based on the constructed null

model, we can infer adjusted p-values (adjusted by Benjamini and Hochberg method [26]) for

each LMSM module. A LMSM module with adjusted p-value less than 0.05 is regarded as a

statistically significant module.

Application of LMSM in BRCA

BRCA enrichment analysis. Instead of performing Gene Ontology (GO) and Kyoto

Encyclopedia of Genes and Genomes Pathway (KEGG) enrichment analysis, to investigate

whether an identified LMSM module is functionally associated with BRCA, we focus on

conducting BRCA enrichment analysis by using a hypergeometric test. For a LMSM module,

the p-value for the test is calculated as:

269
$$p-value = 1 - \sum_{i_2=0}^{L_2-1} \frac{\binom{M_2}{i_2}\binom{N_2-M_2}{K_2-i_2}}{\binom{N_2}{K_2}} \tag{5}$$

where N_2 is the number of genes (lncRNAs and mRNAs) in the dataset, M_2 denotes the

number of BRCA genes in the dataset, K_2 represents the number of genes in the LMSM

module, and L_2 is the number of BRCA genes in the LMSM module. A LMSM module with

p-value < 0.05 is regarded as a BRCA-related module.

Module biomarker identification in BRCA. The module survival analysis can imply whether

the identified LMSM modules are good biomarkers of the metastasis risks of cancer patients

or not, and it can give us the hint whether the LMSM modules may be related to and

potentially affect the metastasis or survival of cancer patients. For each BRCA sample, we fit

the multivariate Cox model (proportional hazards regression model) [27] using the genes

(lncRNAs and mRNAs) in LMSM modules to compute its risk score. All the BRCA samples

are equally divided into the high risk and the low risk groups according to their risk scores.

The Log-rank test is used to evaluate the difference of each LMSM module between the high

and the low risk BRCA groups. Moreover, we also calculate the proportional hazard ratio

(HR) between the high and the low risk BRCA groups. In this work, the *survival* R package

[28] is utilized, and a LMSM module with Log-rank p-value < 0.05 and HR > 2 is regarded

as a module biomarker in BRCA.

Identification of BRCA subtype-specific modules. As is known, BRCA is a heterogeneous

disease with several molecular subtypes, and the choice of chemotherapy for each BRCA

subtype is different. This diversity indicates that the genetic regulation of each BRCA

subtype is specific. To identify BRCA subtype-specific modules, we firstly identify BRCA

molecular subtypes using the PAM50 classifier [29]. By using a 50-gene subtype predictor,

the PAM50 classifier classifies a BRCA sample into one of the five "intrinsic" subtypes:

Luminal A (LumA), Luminal B (LumB), HER2-enriched (Her2), Basal-like (Basal) or

Normal-like (Normal). In this work, we use the *genefu* R package [30] to predict molecular

subtypes of each BRCA sample in the dataset used in our study.

To identify BRCA subtype-specific LMSM modules, we firstly need to estimate the

enrichment scores of LMSM modules in BRCA samples. To calculate the enrichment score

of each LMSM module in BRCA samples, the gene set variation analysis (GSVA) method

[31] is used. To calculate the enrichment score, the GSVA method uses the Kolmogorov-

Smirnov (KS) like random walk statistic as follows:

$$v_{jk}(1) = \frac{\sum_{i=1}^{1} |r_{ij}|^{\tau} I(g(i) \in \gamma_k)}{\sum_{i=1}^{p} |r_{ij}|^{\tau} I(g(i) \in \gamma_k)} - \frac{\sum_{i=1}^{1} I(g(i) \notin \gamma_k)}{p - |\gamma_k|} \qquad (6)$$

where τ ($\tau = 1$ by default) is the weight of the tail in the random walk, r_{ij} is the normalized

expression-level statistics of the i-th gene in the j-th sample as defined in [31], γ_k is the k-th

LMSM module, $I(g(i) \in \gamma_k)$ is the indicator function on whether the i-th gene belongs to the

LMSM module γ_k, $|\gamma_k|$ is the number of genes in the k-th LMSM module, and p is the

number of genes in the dataset.

To transform the KS like random walk statistic into an enrichment score (*ES*, also called

GSVA score), we calculate the maximum deviation from zero of the random walk of the j-th

sample with respect to the k-th LMSM module in the following:

$$ES_{jk}^{\max} = v_{jk}[\arg \max_{1=1,\dots,p} (abs(v_{jk}(1)))] \qquad (7)$$

For each LMSM module γ_k, the formula generates a distribution of enrichment scores that is

bimodal (see the reference [31] for a more detailed description).

Based on the enrichment scores of LMSM modules in each BRCA sample, we further

identify two types of BRCA subtype-specific LMSM modules, up-regulated modules and

down-regulated modules. For one type of regulation pattern (up or down regulation), a

LMSM module is regarded to be specific to a BRCA subtype. For an up-regulated BRCA

subtype-specific LMSM module, the enrichment score of the LMSM module in the specific

BRCA subtype samples is significantly larger than the score in the other BRCA subtype

samples. For a down-regulated BRCA subtype-specific LMSM module, the enrichment score

of the LMSM module in the specific BRCA subtype samples is significantly smaller than the

score in the other BRCA subtype samples. For example, if a LMSM module γ_k is up-

regulated Basal-like specific, the enrichment scores of the LMSM module in Basal-like

samples should be significantly larger than those in Luminal A, Luminal B, HER2-enriched

and Normal-like samples. In this work, for each LMSM module, we use Welch's *t*-test [32] to

calculate the significance *p*-value for the difference of the average enrichment scores between

any two BRCA subtype samples. Given a BRCA subtype, a LMSM module is considered as

an up-regulated (or down-regulated) module specific to this BRCA subtype if the module's

average enrichment score in samples of the given subtype is higher (or smaller) than the

average enrichment score in samples of any other subtype and the significance *p*-value of the

Welch's *t*-test between the samples of this subtype and any other subtype is less than 0.05.

Performance of LMSM modules in classifying BRCA subtypes. In this section, to check the

biological relevance of the discovered LMSM modules, we conduct module classification of

BRCA subtypes. Here, classifying BRCA subtypes (LumA, LumB, Her2, Basal and Normal)

is a multi-class classification (also known as a special case of multi-label classification). To

understand the classification performance of the feature genes in each LMSM module, we

apply a state-of-the-art multi-label learning strategy called Binary Relevance (BR) [33]

implemented in the *utiml* R package [34] to conduct multi-label classification analysis. For

the BR strategy, we use the Support Vector Machine (SVM) classifier [35] with default

parameters implemented in *e1071* R package [36] as the base algorithm to build the multi-

label model. We select two commonly used multi-label classification measures: *Subset*

accuracy and *Hamming loss*, and conduct 10-fold cross-validation to evaluate the

performance of each LMSM module. In this work, *Subset accuracy* denotes the percentage of

correct predictions and *Hamming loss* is the fraction of wrong predictions to the total number

of predictions. Higher values of *Subset accuracy* and smaller values of *Hamming loss*

indicate better classification performance. In addition, for the evaluation, we use the baseline

method in [37], a commonly used multi-label classification method as the baseline for

comparison. The base algorithm of the baseline method is also the SVM classifier with

default parameters implemented in *e1071* R package [36].

Results

Heterogeneous data sources

We collect matched miRNA, lncRNA and mRNA expression data, and clinical data of BRCA

dataset from The Cancer Genome Atlas (TCGA, https://cancergenome.nih.gov/). A lncRNA

or mRNA without a corresponding gene symbol in the expression data of BRCA dataset is

removed. To obtain a unique expression value for replicates of miRNAs, lncRNAs or

mRNAs, we compute the average expression value of the replicates. As a result, we obtain

the matched expression data of 674 miRNAs, 12711 lncRNAs and 18344 mRNAs in 500

BRCA samples.

We retrieve putative miRNA-target interactions (including miRNA-lncRNA and miRNA-

mRNA interactions) from several high-confidence miRNA-target interaction databases and

use the combined database search results. Specifically, the putative miRNA-lncRNA

interactions are obtained from NPInter v3.0 [38] and the experimental module of DIANA-

LncBase v2.0 [39], and miRNA-mRNA interactions are from miRTarBase v8.0 [40],

TarBase v7.0 [41] and miRWalk v2.0 [42].

The BRCA related mRNAs are from DisGeNET v5.0 [43] and COSMIC v86 [44], and the

BRCA related lncRNAs are from LncRNADisease v2.0 [45], Lnc2Cancer v2.0 [46] and

MNDR v2.0 [47]. The ground truth of lncRNA related miRNA sponge interactions is

obtained by integrating the interactions from miRSponge [48], LncCeRBase [49] and

LncACTdb v2.0 [50].

Most of the mediating miRNAs act as crosslinks across LMSM modules

Following the steps shown in Fig 2, we have identified 17 LMSM modules (details can be

seen in S1 Data). The average size of the identified modules is 672.53 and the average

number of the shared miRNAs in a module is 232.82. In total, there are 549 unique miRNAs

mediating the 17 LMSM modules, and 90.16% (495 out of 549) miRNAs mediate at least

two LMSM modules (details can be seen in S2 Data). This result indicates that most of the

mediating miRNAs act as crosslinks across different LMSM modules.

LMSM modules are all statistically significant

376   In this section, by computing null-model-based p-values, we evaluate whether the identified

377   LMSM modules are statistically significant or not. As a result, the adjusted p-values for the

378   identified 17 LMSM modules (details can be seen in S3 Data) are all statistically significant

379   (adjusted p-value = 1.00E-06). This result demonstrates that LMSM modules are all

380   statistically significant.

Most of LMSM modules are implicated in BRCA

To investigate whether the identified LMSM modules are related to BRCA or not, we

conduct BRCA enrichment analysis and identify BRCA module biomarkers using the

methods described in Section Methods. For the BRCA enrichment analysis, we have

collected a list of 4819 BRCA genes (734 BRCA lncRNAs and 4085 BRCA mRNAs)

associated with the matched lncRNA and mRNA expression data (details in S4 Data). As

shown in Table 1, 10 out of 17 LMSM modules are functionally enriched in BRCA at a

significant level (p-value < 0.05). In Table 2, 15 out of 17 LMSM modules are regarded as

module biomarkers in BRCA at a significant level (Log-rank p-value < 0.05 and HR > 2).

Particularly, 90% (9 out of 10, excepting LMSM 14) of the BRCA-related LMSM modules

can act as module biomarker in BRCA. These results show that most of LMSM modules are

functionally implicated in BRCA.

Table 1. BRCA-related LMSM modules. L_2 is the number of BRCA genes in each LMSM
module, K_2 represents the number of genes in each LMSM module, the number of BRCA
genes in the dataset (M_2) is 4819, and the number of genes in the dataset (N_2) is 31055.

Module ID	L_2	K_2	p-value
LMSM 2	327	1338	0
LMSM 3	259	1340	7.34E-05
LMSM 4	78	392	1.14E-02
LMSM 5	89	449	8.07E-03
LMSM 6	88	370	1.97E-05
LMSM 8	275	880	0
LMSM 12	24	110	4.95E-02
LMSM 13	20	76	1.05E-02
LMSM 14	252	1004	8.88E-16
LMSM 16	48	182	1.11E-04

396 **Table 2. Survival analysis of LMSM modules in BRCA**. HRlow95 and HRup95 represent
397 the lower and upper of 95% confidence interval of HR, respectively.

Module ID	Chi-square	p-value	HR	HRlow95	HRup95
LMSM 1	170.37	0	10.75	5.88	19.65
LMSM 2	107.34	0	6.03	3.12	11.66
LMSM 3	90.62	0	5.43	2.94	10.01
LMSM 4	138.81	0	14.94	8.83	25.27
LMSM 5	148.49	0	8.64	4.63	16.13
LMSM 6	142.64	0	13.40	7.83	22.92
LMSM 7	161.91	0	13.97	8.01	24.36
LMSM 8	103.63	0	5.91	3.07	11.37
LMSM 10	144.86	0	8.63	4.74	15.71
LMSM 11	120.79	0	9.49	5.55	16.23
LMSM 12	49.31	2.19E-12	5.46	3.38	8.80
LMSM 13	60.08	9.10E-15	5.72	3.48	9.41

LMSM 15	83.26	0	12.00	7.46	19.32
LMSM 16	110.94	0	11.25	6.79	18.66
LMSM 17	106.96	0	9.14	5.42	15.41

LMSM modules are mostly BRCA subtype-specific

In this section, we firstly divide the 500 BRCA samples into five "intrinsic" subtypes
(Luminal A, Luminal B, HER2-enriched, Basal-like and Normal-like). The numbers of
LumA, LumB, Her2, Basal and Normal samples are 190, 155, 52, 85 and 18, respectively.
Then we calculate the enrichment scores of the identified 17 LMSM modules in the BRCA
subtype samples respectively (details in S5 Data).

As illustrated in Fig 3, out of the 17 LMSM modules, 4 and 6 modules are identified as
up-regulated and down-regulated BRCA subtype-specific LMSM modules, respectively. For
the up-regulated BRCA subtype-specific LMSM modules, the numbers of Basal-specific,
LumB-specific and Normal-specific modules are 1, 1 and 2, respectively. The numbers of
Basal-specific, LumB-specific and Normal-specific modules are 3, 1 and 2 respectively
among the down-regulated BRCA subtype-specific LMSM modules. In particular, only 1
module (LMSM 2) can act as both up-regulated and down-regulated BRCA subtype-specific
LMSM module. In total, the unique number of BRCA subtype-specific LMSM modules is 9,
indicating that most of LMSM modules are BRCA subtype-specific.

A

Fig 3. Heatmap of the enrichment scores of BRCA subtype-specific LMSM modules in five BRCA subtype samples. (A) Up-regulated BRCA subtype-specific LMSM modules. (B) Down-regulated BRCA subtype-specific LMSM modules.

The performance of LMSM modules is significantly higher than baseline's performance in classifying BRCA subtypes

For the identified 17 LMSM modules, the average *Subset accuracy* and *Hamming loss* in classifying BRCA subtypes is 0.7547 and 0.0892, respectively (details can be seen in S6 Data), The *Subset accuracy* and *Hamming loss* of the baseline are 0.3800 and 0.2480, respectively. By using Welch's *t*-test method, the *Subset accuracy* achieved using the 17 LMSM modules is significantly larger (better) than the *Subset accuracy* of the baseline (*p*-value < 2.20E-16), and the *Hamming loss* of the 17 LMSM modules is significantly smaller (better) than the *Hamming loss* of the baseline (*p*-value < 2.20E-16). The better performance than the baseline method indicates that LMSM modules are biological meaningful in classifying BRCA subtypes.

Several lncRNA-related miRNA sponge interactions are experimentally confirmed

For the ground truth used in the validation, we have collected 581 experimentally validated lncRNA-related miRNA sponge interactions associated with the matched lncRNA and mRNA expression data (details in S4 Data). After we merge the sponge lncRNA-mRNA pairs in the identified 17 LMSM modules, we have predicted 1471664 unique lncRNA-related miRNA sponge interactions (details at https://github.com/zhangjunpeng411/LMSM). For each LMSM module, the number of shared miRNAs, lncRNAs, mRNAs, predicted lncRNA-related miRNA sponge interactions can be seen in S7 Data.

As shown in Table 3, there are 4 LMSM modules (LMSM 2, LMSM 3, LMSM 5 and LMSM 8) containing 14 experimentally validated lncRNA-related miRNA sponge interactions in total. It is noted that all the lncRNAs and mRNAs in these confirmed lncRNA-related miRNA sponge interactions are BRCA-related genes, indicating they may have potentially involved in BRCA.

Table 3. Validated lncRNA-related miRNA sponge interactions.

Module ID	Validated lncRNA-related miRNA sponge interactions
LMSM 2	*H19*: *HMGA2, H19:IGF2, H19:ITGB1, H19*: *TGFB1, H19*: *VIM, H19:RUNX1, H19:CDH13, H19:KLF4, H19:TGFBI, H19:VDR*
LMSM 3	*LINC00152*: *MCL1*
LMSM 5	*NEAT1*: *EMP2*
LMSM 8	*LINC00324*: *BTG2, DLEU2*: *CCNE1*

LMSM is capable of predicting miRNA targets

LMSM use high-confidence miRNA-target interactions as seeds to predict miRNA-target interactions. A miRNA-mRNA or miRNA-lncRNA pair in a LMSM module has the potential to be a miRNA-target pair for the following reasons. Firstly, at sequence level, the sponge lncRNAs and mRNAs in each LMSM module have a significant sharing of miRNAs. Secondly, at expression level, the sponge lncRNAs and mRNAs in each LMSM module are

highly correlated. As a result, the sponge lncRNAs and mRNAs of each LMSM module have

a high chance to be target genes of the shared miRNAs. Thus, based on the identified LMSM

modules, we have predicted 2820524 unique miRNA-target interactions (including 2023304

miRNA-lncRNA   and   797220   miRNA-mRNA   interactions)   (details   at

https://github.com/zhangjunpeng411/LMSM). For each LMSM module, the numbers of

predicted miRNA-lncRNA interactions and miRNA-mRNA interactions can be seen in S7

Data.

In addition, we investigate the intersection of the miRNA-target interactions predicted by

LMSM with the other well-cited miRNA-target prediction methods. In terms of miRNA-

mRNA interactions, we select TargetScan v7.2 [51], DIANA-microT-CDS v5.0 [52],

starBase v3.0 [53] and miRWalk v3.0 [54] for investigation. We choose starBase v3.0 [53]

and DIANA-LncBase v2.0 [39] for investigation in terms of miRNA-lncRNA interactions.

As shown in the UpSet plot [55] of Fig 4A, the number of miRNA-mRNA interactions

identified by all the five methods is only 21842. However, the percentage of overlap between

LMSM and each of the other four methods achieves ~63.74% (1289620 out of 2023304). As

shown in Fig 4B, the number of miRNA-lncRNA interactions identified by all the three

methods is only 1160. Since the miRNA-lncRNA interactions are still limited, most of the

miRNA-lncRNA interactions (~93.90%, 748609 out of 797220) are individually predicted by

LMSM.

Fig 4. Overlaps and differences between predicted miRNA-target interactions by LMSM and other methods. (A) Predicted miRNA-mRNA interactions between LMSM and TargetScan, DIANA_microT_CDS, starBase, miRWalk. (B) Predicted miRNA-lncRNA interactions between LMSM and starBase, DIANA_LncBase. Each column corresponds to an exclusive intersection that includes the elements of the sets denoted by the dark or red circles, but not of the others. The overlap size between different methods denotes exclusive overlaps, i.e. the overlap set not in a subset of any other overlap set.

Comparison with graph clustering-based strategy

Graph clustering-based strategy [12-17] is an alternative approach to identifying lncRNA

related miRNA sponge modules. As there is no graph clustering-based strategy specifically

designed for finding lncRNA related miRNA sponge modules, so we create a baseline Graph

Clustering-based method (called GC in this paper) which uses well-known network

construction and graph clustering methods as described in the following. The GS method

includes two steps: i) identifying lncRNA related miRNA sponge interaction network, and ii)

identifying lncRNA related miRNA sponge modules from the identified network. In step 1,

we adapt the well-cited Sensitivity Correlation (SC) method [24] implemented in the

miRspongeR R package [56] to infer lncRNA related miRNA sponge interaction network. A

lncRNA-mRNA pair is considered as an interacting pair in the network if they have

significant sharing of the miRNAs, significant correlation and adequate sensitivity correlation.

We require that the pairs must share at least 3 miRNAs and their sensitivity correlation (the

difference between correlation and partial correlation) must be larger than 0.1. The

statistically significance of the miRNA sharing and positive correlations are tested using

hypergeometric test and Welch's *t*-test respectively, with a significant level at 0.05. In step 2,

we use the well-cited Markov cluster (MCL) algorithm [57] to infer lncRNA related miRNA

sponge modules. Here, each obtained cluster corresponds to a module. Each module should

contain at least 2 sponge lncRNAs and 2 target mRNAs. In total, by using the GC method, we

have obtained 108 lncRNA related miRNA sponge modules.

We compare LMSM and GC in terms of the percentage of BRCA-related modules, the

percentage of module biomarkers in BRCA, the classification performance (mean *Subset*

accuracy and mean *Hamming loss*) in classifying BRCA subtypes, and the number of

validated lncRNA-related miRNA sponge interactions. As shown in Table 4, the comparison

result indicates that LMSM always performs better than the GC method. The detailed results

of the GC method can be seen in S8 Data.

Table 4. Comparison results between LMSM and GC.

Method	%BRCA-related modules	%Module biomarkers	Mean *Subset accuracy*	Mean *Hamming loss*	#Validated interactions
LMSM	**58.82%**	**88.24%**	**0.7547**	**0.0892**	**14**
GC	32.41%	66.67%	0.6586	0.1319	2

LMSM is robust

To demonstrate the robustness of the LMSM workflow, we use the sparse group factor analysis (SGFA) method [58], instead of the WGCNA method to identify lncRNA-mRNA co-expression modules. The SGFA method is extended from the group factor analysis (GFA) method [59-61], and it can reliably infer biclusters (modules) from multiple data sources, and provide predictive and interpretable structure existing in any subset of the data sources. Given *B* biclusters to be identified, the SGFA method assigns each column (lncRNA or mRNA) or row (sample) a grade of membership (association) belonging to these biclusters. The range of the values of the associations is [-1, 1]. We use the absolute value of association (*AVA*) to evaluate the strength of lncRNAs and mRNAs belonging to a bicluster, and the cutoff of *AVA* is also set to 0.8. Specifically, we use the *GFA* R package [58] to identify lncRNA-mRNA co-expression modules. The parameter settings for inferring lncRNA-related miRNA sponge modules are the same.

By using the SGFA method, we have identified 51 LMSM modules (details can be seen in S1 Data). The average size of these LMSM modules is 277.63 and the average number of the shared miRNAs is 135.65. There are 490 unique miRNAs mediating the 51 LMSM modules, and 84.90% (416 out of 490) miRNAs mediate at least two LMSM modules (details can be seen in S2 Data). As the result obtained using the WGCNA method, the result with the SGFA method also implies that the mediating miRNAs mostly act as crosslinks across different LMSM modules. In addition, by using a null-model-based *p*-value computation method, the

identified 51 LMSM modules are also all statistically significant with adjusted *p*-value ≤

5.00E-06 (details can be seen in S3 Data).

As shown in S1 Table of S1 File, 3 out of the 51 LMSM modules are functionally

enriched in BRCA at a significant level (*p*-value < 0.05). Moreover, 49 out of the 51 LMSM

modules are regarded as module biomarkers in BRCA (see in S2 Table of S1 File). The

results indicate that most of LMSM modules are related to BRCA.

We also compute the enrichment scores of the identified 51 LMSM modules in the BRCA

subtype samples (details in S5 Data). As illustrated in S1 Fig of S1 File, out of the 51 LMSM

modules, 33 and 24 modules are regarded as up-regulated and down-regulated BRCA

subtype-specific LMSM modules, respectively. For the up-regulated BRCA subtype-specific

LMSM modules, the numbers of Basal-specific, Her2-specific, LumB-specific and Normal-

specific modules are 27, 2, 2 and 2, respectively. The numbers of Basal-specific, Her2-

specific, LumA-specific, LumB-specific and Normal-specific modules are 2, 3, 15, 3 and 1

respectively for the down-regulated BRCA subtype-specific LMSM modules. Particularly, 16

modules can act as both up-regulated and down-regulated BRCA subtype-specific LMSM

module. Overall, the unique number of BRCA subtype-specific LMSM modules is 41. This

result also indicates that the identified LMSM modules are mostly BRCA subtype-specific.

The average value of *Subset accuracy* and *Hamming loss* of the identified 51 LMSM

modules in classifying BRCA subtypes is 0.6921 and 0.1135, respectively (details can be

seen in S6 Data). In classifying BRCA subtypes, the baseline value of *Subset accuracy* and

Hamming loss is 0.3800 and 0.2480, respectively. By using Welch's *t*-test method, the value

of *Subset accuracy* for 51 LMSM modules is significantly larger (better) than the baseline

value of *Subset accuracy* (*p*-value < 2.20E-16), and the value of *Hamming loss* for 51 LMSM

modules is significantly smaller (better) than the baseline value of *Hamming loss* (*p*-value <

2.20E-16). The better performance than the baseline method also indicates that LMSM

modules are biological meaningful in classifying BRCA subtypes.

Moreover, we have predicted 605456 unique lncRNA-related miRNA sponge interactions

in the identified 51 LMSM modules (details at https://github.com/zhangjunpeng411/LMSM).

The number of the shared miRNAs, lncRNAs, mRNAs, predicted lncRNA-related miRNA

sponge interactions of each LMSM module can be seen in S7 Data. Since the experimentally

validated lncRNA-related miRNA sponge interactions are still limited, only 4 LMSM

modules containing 4 lncRNA-related miRNA sponge interactions (see S3 Table of S1 File)

are experimentally validated. All lncRNAs and mRNAs in the confirmed lncRNA-related

miRNA sponge interactions are also BRCA-related genes.

LMSM also has identified a large number of potential miRNA-target interactions

(1646449 in total, including 435345 miRNA-mRNA and 1211104 miRNA-lncRNA

interactions, details at https://github.com/zhangjunpeng411/LMSM). The number of

predicted miRNA-lncRNA interactions, predicted miRNA-mRNA interactions, putative

miRNA-lncRNA interactions and putative miRNA-mRNA interactions can be seen in S7

Data. As illustrated in S2 Fig of S1 File, the number of the miRNA-mRNA interactions

identified by all the five methods is 4897 and the number of the miRNA-lncRNA interactions

identified by all the three methods is 1149. Most of the identified miRNA-mRNA interactions

by LMSM (~58.55%, 254910 out of 435345) are also predicted by one of the other four

methods. In terms of the predicted miRNA-lncRNA interactions, ~94.23% (1141232 out of

1211104) miRNA-lncRNA interactions are also individually predicted by LMSM.

Finally, in terms of the percentage of BRCA-related modules, the percentage of module

biomarkers in BRCA, the classification performance (mean *Subset accuracy* and mean

Hamming loss) in classifying BRCA subtypes, and the number of validated lncRNA-related

miRNA sponge interactions, LMSM also generally performs better than the GC method (see

S4 Table of S1 File).

Altogether, the above results are consistent with those obtained using the WGCNA

method, indicating that our LMSM workflow is robust for studying lncRNA-related miRNA

sponge modules.

Discussion

The crosstalk between different RNA transcripts in a miRNA-dependent manner forms a

complex miRNA sponge interaction network and depicts a novel layer of gene expression

regulation. Until now, several types of RNA transcripts, e.g. lncRNAs, pseudogenes,

circRNAs and mRNAs, have been confirmed to act as miRNA sponges. Since lncRNAs are a

large class of ncRNAs and function in many aspects of cell biology, including human

cancers, we focus on identifying lncRNA related miRNA sponge modules in this work.

By integrating multiple data sources, previous studies mainly investigate the identification

of lncRNA related miRNA sponge interaction network. Based on the identified lncRNA

related miRNA sponge interaction network, they use graph clustering algorithms to further

infer lncRNA related miRNA sponge modules. Different from existing computational

methods on lncRNA related miRNA sponge modules, in this work, we propose a novel

method named LMSM to directly identify lncRNA related miRNA sponge modules from

heterogeneous data. It is noted that the LMSM method depends on our presented hypothesis

of miRNA sponge modular competition. In the hypothesis, miRNA sponges tend to form a

group to compete with a group of target mRNAs for binding with miRNAs.

We have applied the LMSM method to the BRCA dataset from TCGA. For the putative

miRNA-target interactions, we integrate high-confidence miRNA-target interactions from

several databases. The analysis results demonstrate that our LMSM method is useful in

identifying lncRNA related miRNA sponge modules, and it can help with understanding

regulatory mechanism of lncRNAs.

LMSM is a flexible method to investigate miRNA sponge modules in human cancer.

Firstly, any biclustering or clustering algorithm (e.g. the joint non-negative matrix

factorization methods presented by Deng *et al*. [18] and Xiao *et al*. [19]) can be plugged in

stage 1 of LMSM to identify lncRNA-mRNA co-expression modules. The only condition for

using these algorithms is that they can be used to identify biclusters or clusters from high-

dimensional expression data. Secondly, LMSM is a parametric model, and the parameter

settings of LMSM can be replaced according to the practical requirements of researchers. For

example, the threshold of the three metrics in stage 2 for identifying lncRNA related miRNA

sponge modules can be looser or stricter. Thirdly, LMSM can also be extended to study other

ncRNA (e.g. circRNA and pseudogene) related miRNA sponge modules. For instance, if we

change the matched lncRNA expression data and the miRNA-lncRNA interactions to

matched circRNA expression data and the miRNA-circRNA interactions respectively, the

pipeline of LMSM is to identify circRNA related miRNA sponge modules.

It is noted that each LMSM module contains many sponge lncRNAs and mRNAs, so it is

hard to experimentally validate such a module by follow-up wet-lab experiments. This is a

common issue of existing computational methods, including LMSM. We suggest that

biologists can select some sponge lncRNAs and mRNAs of interest in each LMSM module,

and then validate the modular competition between the selected sponge lncRNAs and target

mRNAs. We believe that LMSM is still useful in shortlisting high-confidence sponge

lncRNAs and mRNAs for experimental validation. For example, previous study [62] has

shown that lncRNA *MIR22HG* is functionally complementary to lncRNA *H19*. In the

identified LMSM module no. 2 (LMSM 2), lncRNA *H19* is experimentally validated to

compete with 10 target mRNAs (*HMGA2*, *IGF2*, *ITGB1*, *TGFB1*, *VIM*, *RUNX1*, *CDH13*,

KLF4, *TGFBI* and *VDR*). Thus, biologists can select 2 lncRNAs (*H19* and *MIR22HG*) and 10

target mRNAs (*HMGA2*, *IGF2*, *ITGB1*, *TGFB1*, *VIM*, *RUNX1*, *CDH13*, *KLF4*, *TGFBI* and

VDR) in LMSM 2 to validate the modular competition between them.

Taken together, based on the hypothesis of miRNA sponge modular competition, we

propose a new approach to identifying lncRNA related miRNA sponge modules by

integrating expression data and miRNA-target binding information. Our method not only

extends the ceRNA hypothesis, but also provides a novel way to investigate the biological

functions and modular mechanism of lncRNAs in BRCA. We believe that our method can be

also applied to other human cancer datasets assists in human cancer research.

Supporting information

S1 Data. The identified LMSM modules.

S2 Data. The distribution of the shared miRNAs in LMSM modules.

S3 Data. Statistically significant analysis results of LMSM modules.

S4 Data. BRCA-related genes and experimentally validated lncRNA related miRNA sponge interactions.

S5 Data. The enrichment scores of the identified LMSM modules in the BRCA subtype samples.

S6 Data. Classification analysis results of LMSM modules in classifying BRCA subtypes.

S7 Data. The number of shared miRNAs, lncRNAs, mRNAs, predicted interactions for each LMSM
module.

S8 Data. The results of a graph clustering-based strategy.

S1 File. Supporting file. Supplementary file.

References

1.   Derrien T, Johnson R, Bussotti G, Tanzer A, Djebali S, Tilgner H, et al. The GENCODE v7
catalog of human long noncoding RNAs: analysis of their gene structure, evolution, and
expression. Genome Res 2012; 22(9):1775-1789. doi: 10.1101/gr.132159.111.

2.   Fang Y, Fullwood MJ. Roles, functions, and mechanisms of long non-coding RNAs in cancer.
Genomics Proteomics Bioinformatics 2016; 14(1):42-54. doi: 10.1016/j.gpb.2015.09.006.

3.   Bhan A, Soleimani M, Mandal SS. Long noncoding RNA and cancer: a new paradigm.
Cancer Res 2017; 77(15):3965-81. doi: 10.1158/0008-5472.CAN-16-2634.

4.   Kopp F, Mendell JT. (2018) Functional classification and experimental dissection of long
noncoding RNAs. Cell 2018; 172(3):393-407. doi: 10.1016/j.cell.2018.01.011.

5.   Ambros V. The functions of animal microRNAs. Nature 2004; 431(7006):350-5. doi:
10.1038/nature02871.

6.   Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. Cell 2004;
116(2):281-97. doi: 10.1016/s0092-8674(04)00045-5.

7.   Salmena L, Poliseno L, Tay Y, Kats L, Pandolfi PP. A ceRNA hypothesis: the Rosetta Stone
of a hidden RNA language? Cell 2011; 146(3):353-8. doi: 10.1016/j.cell.2011.07.014.

8.   Tay Y, Rinn J, Pandolfi PP. The multilayered complexity of ceRNA crosstalk and competition.
Nature 2014; 505(7483):344-52. doi: 10.1038/nature12986.

9.   Zhou S, He Y, Yang S, Hu J, Zhang Q, Chen W, et al. The regulatory roles of lncRNAs in the
process of breast cancer invasion and metastasis. Biosci Rep 2018; 38(5):BSR20180772. doi:
10.1042/BSR20180772.

10. Peng F, Li TT, Wang KL, Xiao GQ, Wang JH, Zhao HD, et al. H19/let-7/LIN28 reciprocal
negative regulatory circuit promotes breast cancer stem cell maintenance. Cell Death Dis
2017; 8(1):e2569. doi: 10.1038/cddis.2016.438.

11. Le TD, Zhang J, Liu L, Li J. Computational methods for identifying miRNA sponge
interactions. Brief Bioinform 2017; 18(4):577-590. doi: 10.1093/bib/bbw042.

12. Shao T, Wu A, Chen J, Chen H, Lu J, Bai J, et al. Identification of module biomarkers from
the dysregulated ceRNA-ceRNA interaction network in lung adenocarcinoma. Mol Biosyst
2015; 11(11):3048-58. doi: 10.1039/c5mb00364d.

13. Zhang Y, Xu Y, Feng L, Li F, Sun Z, Wu T, et al. Comprehensive characterization of
lncRNA-mRNA related ceRNA network across 12 major cancers. Oncotarget 2016;
7(39):64148-67. doi: 10.18632/oncotarget.11637.

14. Zhang J, Le TD, Liu L, Li J. Inferring miRNA sponge co-regulation of protein-protein
interactions in human breast cancer. BMC Bioinformatics 2017; 18(1):243. doi:
10.1186/s12859-017-1672-2.

15. Wang H, Xu D, Huang H, Cui Y, Li C, Zhang C, et al. Detection of dysregulated competing
endogenous RNA modules associated with clear cell kidney carcinoma. Mol Med Rep 2018;
18(2):1963-72. doi: 10.3892/mmr.2018.9189.

16. Do D, Bozdag S. Cancerin: A computational pipeline to infer cancer-associated ceRNA
interaction networks. PLoS Comput Biol 2018; 14(7):e1006318. doi:
10.1371/journal.pcbi.1006318.

17. Zhang J, Liu L, Li J, Le TD. LncmiRSRN: identification and analysis of long non-coding
RNA related miRNA sponge regulatory network in human cancer. Bioinformatics 2018;
34(24):4232-40. doi: 10.1093/bioinformatics/bty525.

18. Deng J, Kong W, Wang S, Mou X, Zeng W. Prior knowledge driven joint NMF algorithm for
ceRNA co-module identification. Int J Biol Sci 2018; 14(13):1822-1833. doi:
10.7150/ijbs.27555.

19. Xiao Q, Luo J, Liang C, Cai J, Li G, Cao B. CeModule: an integrative framework for
discovering regulatory patterns from genomic data in cancer. BMC Bioinformatics 2019;
20(1):67. doi: 10.1186/s12859-019-2654-3.

20. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis.
BMC Bioinformatics 2008; 9:559. doi: 10.1186/1471-2105-9-559.

21. Segal E, Shapira M, Regev A, Pe'er D, Botstein D, Koller D, et al. Module networks:
identifying regulatory modules and their condition-specific regulators from gene expression
data. Nat Genet 2003; 34(2):166-76. doi: 10.1038/ng1165.

22. Hotelling H. Relations between two sets of variates. Biometrika 1936; 28(3/4):321-377. doi:
10.2307/2333955.

23. Witten DM, Tibshirani R, Hastie T. A penalized matrix decomposition, with applications to
sparse principal components and canonical correlation analysis. Biostatistics 2009; 10(3):515-
34. doi: 10.1093/biostatistics/kxp008.

24. Paci P, Colombo T, Farina L. Computational analysis identifies a sponge interaction network
between long non-coding RNAs and messenger RNAs in human breast cancer. BMC Syst
Biol 2014; 8:83. doi: 10.1186/1752-0509-8-83.

25. List M, Dehghani Amirabad A, Kostka D, Schulz MH. Large-scale inference of competing
endogenous RNA networks with sparse partial correlation. Bioinformatics 2019; 35(14):i596-
i604. doi:10.1093/bioinformatics/btz314.

26. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful
approach to multiple testing. J R Stat Soc Ser B (Methodological) 1995; 57(1): 289-300. doi:
10.2307/2346101.

27. Andersen P, Gill R. Cox's regression model for counting processes, a large sample study. Ann
Stat 1982; 10(4):1100-20. doi:10.1214/aos/1176345976.

28. Therneau TM, Grambsch PM. Modeling survival data: extending the Cox model. Springer-
Verlag, New York, 2000.

29. Parker JS, Mullins M, Cheang MC, Leung S, Voduc D, Vickery T, et al. Supervised risk
predictor of breast cancer based on intrinsic subtypes. J Clin Oncol 2009; 27(8):1160-7. doi:
10.1200/JCO.2008.18.1370.

30. Gendoo DM, Ratanasirigulchai N, Schröder MS, Paré L, Parker JS, Prat A, et al. Genefu: an
R/Bioconductor package for computation of gene expression-based signatures in breast cancer.
Bioinformatics 2016; 32(7):1097-9. doi: 10.1093/bioinformatics/btv693.

31. Hänzelmann S. GSVA: gene set variation analysis for microarray and RNA-seq data. BMC
Bioinformatics 2013; 14:7. doi: 10.1186/1471-2105-14-7.

32. Welch BL. The generalisation of student's problems when several different population
variances are involved. Biometrika 1947; 34(1-2):28-35. doi: 10.1093/biomet/34.1-2.28.

33. Tsoumakas G, Katakis I, Vlahavas I. Mining Multi-Label Data. In O. Maimon and L. Rokach,
editors, Data Mining and Knowledge Discovery Handbook, chapter 34, pages 667–685.
Springer-Verlag, 2 edition, 2010. ISBN 0387244352. doi:10.1007/978-0-387-09823-4_34.

34. Rivolli A, de Carvalho AC. The utiml package: Multi-label classification in R. The R Journal
2018; 10(2): 24-37. doi:10.32614/RJ-2018-041.

35. Chang CC, Lin CJ. LIBSVM: a library for support vector machines. ACM T Intel Syst Tec.
2011; 2(3):1-27. doi:10.1145/1961189.1961199.

36. Meyer D, Dimitriadou E, Hornik K, Weingessel A, Leisch F. e1071: Misc functions of the
department of statistics, probability theory group (Formerly: E1071), TU Wien, R package
version, 1.7-3. 2019. https://CRAN.R-project.org/package=e1071.

37. Metz J, de Abreu LF, Cherman EA, Monard MC. On the estimation of predictive evaluation
measure baselines for multi-label learning. In 13th Ibero-American Conference on AI, pages
189–198, Cartagena de Indias, Colombia, 2012. doi:10.1007/978-3-642-34654-5_20.

38. Hao Y, Wu W, Li H, Yuan J, Luo J, Zhao Y, et al. NPInter v3.0: an upgraded database of
noncoding RNA-associated interactions. Database (Oxford) 2016. doi:
10.1093/database/baw057.

39. Paraskevopoulou MD, Vlachos IS, Karagkouni D, Georgakilas G, Kanellos I, Vergoulis T, et
al. DIANA-LncBase v2: indexing microRNA targets on non-coding transcripts. Nucleic Acids
Res 2016; 44(Database issue):D231-D238. doi: 10.1093/nar/gkv1270.

40. Huang HY, Lin YC, Li J, Huang KY, Shrestha S, Hong HC, et al. miRTarBase 2020: updates
to the experimentally validated microRNA-target interaction database. Nucleic Acids Res
2019. doi: 10.1093/nar/gkz896.

41. Vlachos IS, Paraskevopoulou MD, Karagkouni D, Georgakilas G, Vergoulis T, Kanellos I, et
al. DIANA-TarBase v7.0: indexing more than half a million experimentally supported
miRNA: mRNA interactions. Nucleic Acids Res 2015; 43(Database issue):D153-D159. doi:
10.1093/nar/gku1215.

42. Dweep H, Gretz N. miRWalk2.0: a comprehensive atlas of microRNA-target interactions. Nat
Methods 2015; 12(8):697. doi: 10.1038/nmeth.3485.

43. Piñero J, Bravo À, Queralt-Rosinach N, Gutiérrez-Sacristán A, Deu-Pons J, Centeno E, et al.
DisGeNET: a comprehensive platform integrating information on human disease-associated
genes and variants. Nucleic Acids Res 2017; 45(Database issue):D833-D839. doi:
10.1093/nar/gkw943.

44. Forbes SA, Beare D, Boutselakis H, Bamford S, Bindal N, Tate J, et al. COSMIC: somatic
cancer genetics at high-resolution. Nucleic Acids Res 2017; 45(Database issue):D777-D783.
doi: 10.1093/nar/gkw1121.

45. Bao Z, Yang Z, Huang Z, Zhou Y, Cui Q, Dong D. LncRNADisease 2.0: an updated database
of long non-coding RNA-associated diseases. Nucleic Acids Res 2019; 47(Database
issue):D1034-D1037. doi: 10.1093/nar/gky905.

46. Gao Y, Wang P, Wang Y, Ma X, Zhi H, Zhou D, et al. Lnc2Cancer v2.0: updated database of
experimentally supported long non-coding RNAs in human cancers. Nucleic Acids Res 2019;
47(Database issue):D1028-D1033. doi: 10.1093/nar/gky1096.

47. Cui T, Zhang L, Huang Y, Yi Y, Tan P, Zhao Y, et al. MNDR v2.0: an updated resource of
ncRNA-disease associations in mammals. Nucleic Acids Res 2018; 46(Database issue):D371-
D374. doi: 10.1093/nar/gkx1025.

48. Wang P, Zhi H, Zhang Y, Liu Y, Zhang J, Gao Y, et al. miRSponge: a manually curated
database for experimentally supported miRNA sponges and ceRNAs. Database (Oxford) 2015.
doi: 10.1093/database/bav098.

49. Pian C, Zhang G, Tu T, Ma X, Li F. LncCeRBase: a database of experimentally validated
human competing endogenous long non-coding RNAs. Database (Oxford) 2018. doi:
10.1093/database/bay061.

50. Wang P, Li X, Gao Y, Guo Q, Wang Y, Fang Y, et al. LncACTdb 2.0: an updated database of
experimentally supported ceRNA interactions curated from low- and high-throughput
experiments. Nucleic Acids Res 2019; 47(Database issue):D121-D127. doi:
10.1093/nar/gky1144.

51. Agarwal V, Bell GW, Nam JW, Bartel DP. Predicting effective microRNA target sites in
mammalian mRNAs. Elife 2015; 4. doi: 10.7554/eLife.05005.
52. Paraskevopoulou MD, Georgakilas G, Kostoulas N, Vlachos IS, Vergoulis T, Reczko M, et al.
DIANA-microT web server v5.0: service integration into miRNA functional analysis
workflows. Nucleic Acids Res 2013; 41(Web Server issue):W169-73. doi:
10.1093/nar/gkt393.
53. Li JH, Liu S, Zhou H, Qu LH, Yang JH. starBase v2.0: decoding miRNA-ceRNA, miRNA-
ncRNA and protein-RNA interaction networks from large-scale CLIP-Seq data. Nucleic
Acids Res 2014; 42(Database issue):D92-7. doi: 10.1093/nar/gkt1248.
54. Sticht C, De La Torre C, Parveen A, Gretz N. miRWalk: An online resource for prediction of
microRNA   binding   sites.   PLoS   One   2018;   13(10):e0206239.   doi:
10.1371/journal.pone.0206239.
55. Conway JR, Lex A, Gehlenborg N. UpSetR: an R package for the visualization of intersecting
sets   and   their   properties.   Bioinformatics   2017;   33(18):2938-2940.   doi:
10.1093/bioinformatics/btx364.
56. Zhang J, Liu L, Xu T, Xie Y, Zhao C, Li J, et al. miRspongeR: an R/Bioconductor package
for the identification and analysis of miRNA sponge interaction networks and modules. BMC
Bioinformatics 2019; 20(1):235. doi:10.1186/s12859-019-2861-y.
57. Enright AJ, Van Dongen S, Ouzounis CA. An efficient algorithm for large-scale detection of
protein families. Nucleic Acids Res. 2002; 30(7):1575–1584. doi:10.1093/nar/30.7.1575.
58. Bunte K, Leppäaho E, Saarinen I, Kaski S. Sparse group factor analysis for biclustering of
multiple   data   sources.   Bioinformatics   2016;   32(16):2457-63.   doi:
10.1093/bioinformatics/btw207.
59. Klami A, Virtanen S, Leppäaho E, Kaski S. Group factor analysis. IEEE Trans Neural Netw
Learn Syst 2015; 26(9):2136-2147. doi: 10.1109/TNNLS.2014.2376974.
60. Suvitaival T, Parkkinen JA, Virtanen S, Kaski S. Cross-organism toxicogenomics with group
factor analysis. Syst Biomed 2014; 2(4):71-80. doi: 10.4161/sysb.29291.
61. Virtanen S, Klami A, Khan S, Kaski S. Bayesian group factor analysis. In: Lawrence,N. and
Girolami,M. (eds), Proc. of the 15th International Conference on Artificial Intelligence and
Statistics, 2012; pp. 1269-1277.
62. Wang P, Ning S, Zhang Y, Li R, Ye J, Zhao Z, et al. Identification of lncRNA-associated
competing triplets reveals global patterns and prognostic markers for cancer. Nucleic Acids
Res 2015; 43(7):3478-89. doi: 10.1093/nar/gkv233.