Genomic architecture of artificially and sexually selected traits in a wild cervid

Running title: Genomic basis of sexually selected traits

S. J. Anderson[1]*, S. D. Côté [2], J. H. Richard[2], A. B. A. Shafer[1,3]*

[1] Environmental and Life Sciences Graduate Program, Trent University, K9J 7B8

Peterborough, Canada.

[2] Département de biologie, Centre d'études nordiques & *Chaire de recherche*

industrielle CRSNG en aménagement intégré des ressources de l'île d'Anticosti,

Université Laval, G1V 0A6, Québec, QC, Canada

[3] Forensics Department, Trent University, K9J 7B8 Peterborough, Canada.

* Corresponding authors: sanderson@trentu.ca, aaronshafer@trentu.ca

17
18
19
20
21   **Keywords**
22   Quantitative traits, population genetics, gene enrichment, polygenic traits

Abstract

Characterization of the genomic architecture of fitness-related traits such as body

size and male ornamentation in mammals provides tools for conservation and

management: as both indicators of quality and health, these traits are often subject to

sexual and artificial selective pressures. Here we performed high-depth whole genome

re-sequencing on pools of individuals representing the phenotypic extremes in our study

system for antler and body size in white-tailed deer (*Odocoileus virginianus*). Samples

were selected from a tissue repository containing phenotypic data for 4,466 male white-

tailed deer from Anticosti Island, Quebec, with four pools representing the extreme

phenotypes for antler and body size in the population, after controlling for age. Our

results revealed a largely panmictic population, but detected highly diverged windows

between pools for both traits with high shifts in allele frequency (mean allele frequency

difference of 14% for and 13% for antler and body SNPs in outlier windows). These

regions often contained putative genes of small-to-moderate effect consistent with a

polygenic model of quantitative traits. Genes in outlier antler windows had known direct

or indirect effects on growth and pathogen defence, while body genes, overall GO

terms, and transposable element analyses were more varied and nuanced. Through

qPCR analysis we validated both a body and antler gene. Overall, this study revealed

the polygenic nature of both antler morphology and body size in free-ranging white-

tailed deer and identified target loci for additional analyses.

Introduction

Quantifying the genomic architecture underlying phenotypes in natural populations provides insights into the evolution of quantitative traits (Stinchcombe & Hoekstra, 2008). Some quantitative traits are correlated with metrics of fitness, and as such are particularly important because they might directly influence population viability (Kardos & Shafer, 2018). This relationship between genomic architecture and quantitative traits, however, is not easy to empirically identify (Barghi et al., 2020), and often has unclear and unpredictable responses to selection (Bünger et al., 2005). Outside of a few traits in well studied systems such as horn morphology in Soay sheep (Johnston et al., 2011, 2013) and migration timing in Pacific salmon (Prince et al., 2017), there are few studies that have successfully identified quantitative trait loci (QTL) in wild vertebrate populations with effects on fitness-related traits.

Genome-wide association scans (GWAS) have the potential to identify links between genomic regions and fitness relevant traits that can directly inform management and have become more prominent in non-model organisms (Ellegren 2014; Santure & Grant 2018). The variations of expressed phenotypes that are observed in wild populations are influenced by demographic, environmental and epigenetic factors (Ellegren & Sheldon 2008), which leads to difficulties when trying to quantify the direct influence of genetic polymorphisms, and more generally differentiate the effects of drift from selection (Kardos & Shafer 2018). Most traits also appear to be composed of many genes of small effect, and heritability for complex traits are often modulated by genes outside of core pathways and coding regions (Robinson et al., 2013; Santure et al., 2013; Boyle et al., 2017; Barghi et al. 2020). Population genetic

modeling allows disentangling phenotypic plasticity and responses to environmental

change, while expanding our understanding of the evolutionary processes in wild

populations (Santure & Grant 2018). Where traits directly linked to reproductive success

are also those humans target via harvesting (Coltman et al., 2003; Kuparinen & Festa-

Bianchet, 2017), there is a need to understand the genomic architecture to predict

evolutionary consequences (Kardos & Luikart, 2015), considering that the two modes of

selection (natural and artificial) might not always act in concert, and can vary spatially

and temporally (Edeline et al. 2007).

For traits relevant to fitness, including those under sexual selection, we might

expect directional selection towards the dominant or desired phenotype. In long term

studies of wild populations, however, there is often a relatively high degree of observed

genetic variation underlying sexually selected traits (Johnston et al., 2013; Berenos et

al., 2015; Malenfant et al., 2018), with the proposed mechanism for this genetic

variation being balancing selection (Mank, 2017; but see Kruuk et al. 2007). For

example, Johnston et al. (2013) showed that variation is maintained at a single large

effect locus through a trade-off between observed higher reproductive success for large

horns and increased survival for smaller horns in Soay sheep. These trade-offs,

creating signatures consistent with balancing selection, are also observed in cases

where social dominance structures exist as a result of male-male competition for female

mate acquisition, where alternative mating strategies have been suggested as a means

for younger or subordinate males to achieve greater mating success (Foley et al.,

2015).

Emerging approaches to investigate the genomic architecture of complex

phenotypes

The methodological approach of analyzing extreme phenotypes seeks to sample

individuals representing the extreme ends of the spectrum for any observable

phenotype, instead of randomly sampling individuals from the entire distribution (Perez-

Gracia et al., 2002; Li et al., 2011; Emond et al., 2012; Barnett et al., 2013; Gurwitz &

McLeod, 2013; Kardos et al., 2016). Pooled sequencing (pool-seq) uses the

resequencing of pooled DNA from individuals within a population and is a cost-effective

alternative to the whole genome sequencing of every individual (Anand et al., 2016) with

lower coverage options emerging (Tilk et al. 2019). Individual identity is lost through the

pooling of DNA, but the resulting allelic frequencies are representative of the population

and can be used to conduct standard population genomic analyses, including outlier

detection (Schlötterer et al., 2014). Pool-seq methods have been applied with the

purposes of identifying genes of moderate-to-large effect loci in a variety of taxa

including colour morphs in butterflies and birds, abdominal pigmentation in *Drosophila*,

and horn size of wild Rocky Mountain bighorn sheep (Kardos et al, 2015; Endler et al.,

2017; Neethiraj et al., 2017).

Here, we explored the genic basis for phenotypic variation by sampling the

extreme phenotypes in a non-model big game species, the white-tailed deer

(*Odocoileus virginianus*, WTD). Two traits are of particular interest in WTD; body size

and antler size, which have a degree of observable variation (Hewitt 2011) and are

connected to individual reproductive success (DeYoung et al., 2009; Newbolt et al.,

2016; Jones et al., 2018). Heritability for antler and body measurements are moderate

to high for antler features and body size (Michel et al., 2016; Williams et al., 1994;

Jamieson et al., 2020). Large antlers and body size are also sought after by hunters,

both as trophies and food sources throughout North America. Efforts to compare

harvested to non-harvested (i.e. naturally shed) antlers suggest the potential for artificial

selection with those harvested by hunters being larger in size (Schoenebeck & Petersen

2014 but see Ditchkoff et al. 2000), which is similar to what has been documented in

other cervids (Ramanzin & Sturaro, 2014; Pozo et al., 2016; Balčiauskas et al., 2017;

García-Ferrer et al., 2019). Our objective was to identify genomic windows associated

with variation in antler and body size phenotypes in WTD, under the hypothesis that

many genes of small to moderate effect underly these phenotypes, but core pathways

should be related to these phenotypes. We performed whole genome re-sequencing on

pools representing the extreme distribution of antler and body size phenotypes in a wild,

but intensively monitored WTD population.

Materials and Methods

Study area

Anticosti Island (49°N, 62°W; 7,943 km^2) is located in the Gulf of St. Lawrence, Québec

(Canada) at the northeastern limit of the white-tailed deer range (Figure 1). The island is

within the balsam fir-white birch bioclimatic region with a maritime sub-boreal climate

characterized by cool and rainy summers (630mm/year), and long and snowy winters

(406 cm/year; Environment Canada, 2006; Simard et al., 2010). The deer population

was introduced in 1896 with ca. 220 animals and rapidly increased. Today, densities are

>20 deer/km^2 and can exceed 50 deer/km^2 locally (Potvin & Breton 2005) with an

estimated population >160,000 individuals.

Sample Collection and Modeling

We collected tissue samples stored in 95% ethanol and phenotypic data on 4,466 male

deer harvested by hunters from September to early December, 2002–2014 on Anticosti

Island. The sample and measurements were completed as part of a large effort led by

the NSERC Industrial Research Chair in integrated resource management of Anticosti

Island focussed on collecting data on the body condition of deer on the island. We used

cementum layers in incisor teeth to age individuals (Hamlin et al., 2000). Two metrics of

antler size were selected: the number of antler tines or points (>2.5 cm) and beam

diameter (measured at the base; ±0.02 cm) (Simard et al., 2014). We used one metric

for body size: body length which is correlated to other metrics such as body weight and

hind foot length (Bundy et al., 1991). Because antler and body size of male cervids are

correlated to age (Solberg et al., 2004; Nilsen & Solberg, 2006), we controlled for age

before ranking males according to antler and body size. We used linear models to

assess the relationship between age and each metric separately. We computed an

antler and body size index based on the average rank of each individual's residual

variation for the phenotypic metrics. The top and bottom 150 individuals from each

group were selected from the available database for DNA extraction, with only deer

having equal representation for large and small phenotypes from the same given year

were included in the final pool in an effort to limit temporal variation. This created four

groups: large antler (LA), small antler (SA), large body size (LB), and small body size

(SB) (Table S1).

DNA Extraction and Genome Sequencing

We isolated DNA from tissue using the Qiagen DNeasy Blood & Tissue Kit. The

concentration of each DNA extract was determined using a Qubit dsDNA HS Assay Kit

(Life Technologies, Carlsbad, CA, USA). We aimed to sequence each pool to 50X as

per recommendations of Schlötterer et al. (2014). Equal quantities of DNA (100

ng/sample) were combined into representative pools for LA (n=48), and SA (n=48), LB

(n=54), and SB (n=61) for a desired final concentration of 20 ng/ul combined DNA for

each pool. Sequencing was conducted at The Centre for Applied Genomics (Toronto,

ON, Canada) on an Illumina HiSeqX with 150 bp pair-end reads (Table S2); meaning

each individual on average should have at least one chromosome sampled.

Genome Annotation

A newly generated draft WTD genome constructed from long (PacBio) and short-read

(Illumina) data was used as a reference (NCBI PRJNA420098; Accession No.

JAAVWD000000000). We performed a full genome annotation by masking repetitive

elements throughout the genome using a custom WTD database developed through

repeat modeler v1.0.11 (Smit & Hubley, 2015) in conjunction with repeatmasker v4.0.7

(Smit, Hubley & Green, 2015) using the NCBI database for artiodactyla, without

masking low complexity regions. The masked genome was then annotated using the

MAKER2 v2.31.9 pipeline (Holt & Yandell, 2011). We used a three-stage process (Laine

et al., 2016) utilizing publicly available white-tailed deer EST and Protein sequences

available through NCBI for initial training with SNAP v2013-11-29 (Korf, 2004). A

resulting hmm file was generated from the GFF output and was used as evidence for

the MAKER2 prediction software, again using SNAP. Evidence from the prior SNAP

trails in gff format were used as evidence for the generation of a training data set for the

AUGUSTUS v3.3.2 gene prediction software. The final annotation was completed using

MAKER2 and AUGUSTUS. Gene IDs were generated using blastp v2.9.0 (Johnson et

al., 2008) on the WTD annotation protein transcripts, restricting the blast search to

human protein annotations in the uniport database (parameters -max_hsps 1 -

max_target_seqs 1 -outfmt 6 -taxids 9606).

Mapping and characterization of SNPs

We performed initial quality filtering for all reads using fastqc and trimmed reads for

quality and adaptors using the default Trimmomatic v.0.36 settings (Bolger et al. 2014).

Reads were then aligned to the unmasked WTD reference genome with BWA-mem

v0.7.17 (Li, 2013). We used samtools v1.10 (Li et al., 2009) to merge and sort all

aligned reads into four files for each representative pool. We then filtered for duplicates

using Picard v2.20.6 (Broad Institute, 2020), obtained uniquely mapped reads with

samtools, and conducted local realignment using GATKv3.8 (Van der Auwera et al.

2013) prior to calling SNPs. Using samtools we called SNPs with mpileup (parameters =

-B -q20). We filtered out the masked regions and indels with 5bp flanking regions and

removed all scaffolds <= 50 kb.

Genome wide differences were calculated between large and small phenotypes

for antler and body size. We used a sliding window approach with window sizes of 1000

bp with a step size of 500 bp and minimum covered fraction of 0.8 to test for allele

frequency differences using Fisher's exact test (FET) and fixation index (F_{ST}) from the

Popoolation2 software suite (Kofler et al., 2011). A minimum and maximum coverage

were determined based off the mode depth +/- half the mode as per Kurland et al

(2019,) and a minimum overall count of the minor allele of 3 for each pool was specified

(Figure S1). Note F_{ST} windows were correlated to FET P-values (Figure S2). We

corrected for multiple testing by using an FDR correction (see Benjamini & Hochberg

1995) and set a conservative α (1.0e-7). Manhattan plots were generated using a

custom R script to plot the distribution of outliers by position and scaffold throughout the

entire WTD genome. Bedtools v2.27.1 was used to characterize each window being

within 25 kb up/downstream of a gene (i.e. regulatory). Numerous candidate genes for

body size and head gear have been identified in ruminants (Bouwman et al., 2018; Ker

et al., 2018; Wang et al., 2019) and were also complied into an *a priori* list (Table S4).

For these *a priori* genes and outlier windows we identified highly differentiated SNPs

using the modified chi-square test from Spitzer et al. (2020) that accounts for

overdispersion.

Transposable Elements

We used consensus transposable element (TE) sequences from the repbase database

for cow to repeat mask the WTD reference genome, and subsequently merged this

masked genome with the repbase TE reference sequences (Bao et al., 2015). We used

the repbase database consensus sequences to re-mask the genome due to more

robust information for TE identities, family and order required in this analysis. Trimmed

reads for both antler and body size phenotypes were aligned to the TE-WTD merged

reference genome using bwa-mem as stated in previous methods. We used the

recommended workflow for the software suite popoolationTE2 v1.10.04 (Kofler et al.,

2016) to create a list of predicted TE insertions. From this we calculated TE frequency

differences between large and small phenotypes as well as proximity to genic regions.

Here, we only examined TEs that were within 25-kb up or downstream from a gene and

within the 95th percentile for absolute frequency differences to allow for more features to

be assessed subsequently.

GO Pathways

To identify shared gene pathways among outlier regions we used an analysis of gene

ontology (GO) terms. The program Gowinda v1.12 (Kofler & Schlötterer, 2012) was

used to determine GO term enrichment, while also accounting for biases in gene length

but requires input of individual SNPs. A gtf version of our annotation was created by

removing duplicate genes and retaining only the longest gene – resulting in 15,395

unique genes. All SNPs in outlier windows that were also within 25-kb of a gene were

included in this analysis, and compared to all SNPs in every qualifying window for antler

and body size phenotypes independently. We used the program REVIGO (Supek et al.,

2011) to remove redundant GO terms and to visualize semantic similarity-based

scatterplots. Results from Gowinda with p-values <0.05 were used with REVIGO to

generate plots of significant GO terms for biological processes of antler and body size

phenotypes.

Validation of outliers

We selected three SNPs for qPCR validation that had significant chi-square values and

passed quality control with respect to primer design. Custom genotyping assays using

rhAMP chemistry (Integrated DNA Technologies) were designed and genotyped on the

QuantStudio 3 (Thermo Fisher Scientific). Oligo sequences and reaction parameters are

provided in Table S3. Using the phenotypic category as the binary response variable,

we ran a logistic regression treating the genotypic data as additive (e.g. 0-2 LA/LB

alleles per locus, per individual).

Results

Phenotypes

The distribution of phenotypes from all pools of individuals is shown in Figure 2. We

only selected the top 150 individuals at the tail ends of the distribution for

measurements used in our antler and body size rankings which are representative of

the extreme phenotypes. Artist renderings and the distribution of measurements for the

number of antler points, beam diameter, and body length between the groups of

individuals representing each extreme phenotype pool (LA, SA, LB, SB) are shown in

Figure 2a. There were no differences in mean age between the pools (Figure 2b).

Detecting genetic variants and their genomic regions

The total reads generated from resequencing are observed in Table S2 (BioProject ID

PRJNA576136). The WTD genome that has a N50 of 17 MB, 727 scaffolds >50 kb

(98.85% of genome), and BUSCO completeness of 91% across 303 orthologs. The

genome annotation resulted in 20,750 genes. The mean allele frequency difference

across the antler outlier windows were calculated to be 14% with 5249 windows

meeting the FDR and α threshold ($p <= 1\mathrm{x}10^{-7}$); many of the top windows were adjacent

to genes with known functions (Tables 1 & 2). For the comparison of body size, the

mean allele frequency difference was 13% in 1350 windows meeting the FDR threshold.

From the list of 32 *a priori* candidate genes in the literature, we identified the

highest differentiated windows (Table 3); two genes had window values exceeding the

FDR ($p < 10^{-7}$) for the antler analysis (OLIG1, HMGA2), and three from the body

analysis (OLIG1, IGF1, TWIST1). For all individual SNPs within outlier windows and a-

priori genes, the modified chi-square test with FDR correction identified 5,184 and 1,166

SNPs (p < 0.01) from the antler and body analysis, respectively.

TE Insertions

We identified 19,160 TE insertions through the joint analysis of antler phenotype

sequences (both large and small). Of these TEs, we identified 958 insertions that fell

within the 95th percentile for the absolute difference between large and small

phenotypes for further analysis (>0.20%; Figure S3). Of the 95th percentile TE

insertions in the antler analysis, 92 were found to overlap with genes, with 283 within a

25-kb window up or downstream of genic regions. For body size, 6,610 TE insertions

were identified with 332 insertions in the 95th percentile (>0.21%). Of these, 28

overlapped with genes, and 116 within 25-kb of a genic region.

Gene Ontology Annotations

The results from Gowinda comparing all SNPs within outlier windows to background

SNPs from all windows showed the top 10 enriched GO terms and gene counts (Figure

S4). Six statistically enriched GO terms (FDR < 0.05) from the antler analysis were:

GO:0004888 (transmembrane signaling receptor activity), GO:0004872 (receptor

activity), GO:0038023 (signaling receptor activity), GO:0004930 (G-protein coupled

receptor activity), GO:0022835 (transmitter-gated channel activity), GO:0022824

(transmitter-gated ion channel activity). There were no significant GO terms identified

through the body analysis following FDR corrections. We show term reductions for

significantly enriched GO terms (p-value <0.05) through the program REVIGO, with

clustered items relating to the semantic similarity in the antler pools (Figure 4 and

Figure S7).

Validation of Outliers

We genotyped putative antler loci (RIMS1, and SRP54) and one body loci (LIRF1) split

between pools which resulted in 80 and 70 individuals having complete genotype and

phenotypic data, respectively (Dryad Accession no. XXXXXX). The logistic regression

showed an effect of the number of antler alleles at the SRP54 locus on antler category

(ß = -0.99, p = 0.02) and the number of LIRF1 alleles on body category (ß = -0.94, p =

0.03); the RIMS1 locus frequency differences wereas not validated (p > 0.05).

Discussion

The study of sexually selected quantitative traits in the Anticosti Island WTD population

has provided novel insights into the underlying genomic architecture of phenotypic

variation. The extensive database with phenotypic measurements for these deer

(n=4,466) allowed us to select individuals from the wide range of the distribution (Figure

2 & Figure S1) that are representative of extreme phenotypes for the two antler and one

body size measurements. This sampling methodology is important to maximize the

additive genetic variance for each trait, increasing the power and the ability to detect

QTL (Kardos et al., 2016). Anticosti Island WTD form a putatively panmictic population,

where we would expect gene flow across the island (Fuller et al., 2020). Consistent with

this, genome-wide window F_{ST} was ~0.03 for both pools; however, we identified outlier

regions for both traits that are atypically differentiated for what would be expected in this

population (Fuller et al. 2020). The most divergent windows and TEs for antler and body

traits are widely dispersed throughout the WTD genome (Figure 3), which support a

polygenic model for these quantitative traits. This is consistent with the literature on

body size and ornaments in mammals (Visscher et al., 2007; Bouwman et al., 2018;

Wang et al., 2019).

Genomic architecture of antlers and body size

For both traits, there is evidence for divergent windows throughout the genome that

overlap with genic regions (Figure 3; Tables 1-3). Traditionally, variation in SNPs is

focussed on non-synonymous variants, but it is becoming evident that non-coding

regions impact phenotypic variation (Watanabe et al., 2019), often with clear

relationships to promoters and enhancer regions (Pagani & Baralle 2004; Zhang &

Lupski 2015; Foote et al., 2016). While some outliers are surely false positives (Whitlock

& Lottheros 2015), cattle GWAS studies typically identify 100s to 1000s of QTL (Cole et

al., 2011; Jiang et al., 2019). Our data suggest that it is the cumulative effects from

these variants, including TE insertions, and likely pleiotropic and epistatic interactions

that are ultimately driving the phenotypic variation in antler and body size in white-tailed

deer.

Antlers are the only completely regenerable organ found in mammals (Li et al.,

2014), a unique process that involves simultaneous exploitation of oncogenic pathways

and tumor suppressor genes, and the rapid recruitment of synaptic and blood vesicles

(Wang et al., 2019). Size of antlers might also be an honest signal of male quality

(Vanpé et.al, 2007). Accordingly, three of the most diverged windows (Figure 3) were in

close proximity to genes linked to biological processes related to development and

immune responses (Table 1). LGALS9 for example is a gene that produces Galactin-9,

a well studied protein expressed on tumour cells (Heusschen et.al., 2013). The function

of ruminant gammadelta T cells is defined by WC1.1 (Rogers et.al., 2005), and

suggests a role in host response to cell proliferation. Interestingly, MTMR2

(Myotubularin Related Protein 2) has been indicated as a candidate gene for litter size

in pelibuey sheep (Hernández-Montiel et.al., 2020) and plays a role in spermatogenesis

(Mruk & Cheng, 2011). Likewise, the qPCR validated SRP54 gene has been linked to

bone marrow failure syndromes and skeletal abnormalities (Carapito et al., 2017). Thus,

there are clear links to bone formation, oncogenic pathways, and sexual selection in the

antler outlier regions.

The comparison of extreme body size phenotype sequences also revealed an

array of genes of interest; however, it was more difficult than antlers to not engage in

storytelling (Pavlidis et al., 2012) as traits of this nature, for example human height,

involve thousands of SNPs and a multitude of biological pathways (Marouli et al., 2017).

Despite the high heritability of body size in this population (Jamieson et al., 2020), there

were fewer windows meeting our threshold, again consistent with many genes of small

effect (see Barghi et al., 2020). Similarly, the list of 32 *a priori* genes generally revealed

lower levels of variation (i.e. 29 of 32 genes had windows with $p > 10^{-7}$) between

phenotypes for these genes for both traits, despite their known effect; this is likely due

to their influence shaping between species, rather than within species trait variation.

That being said, observing three outlier genes in this non-exhaustive list does suggest

focussing on candidate genes still has value in the genomic era.

Transposable elements considerations

Traditionally, masking transposable elements reduces misalignments due to the

repetitive nature of their sequences and current limitations with mapping software

(Treangen & Salzberg 2012), but also misses a wealth of information that encompasses

a large portion of the genome. Our focus was the frequency differences of TE insertions

to properly mapped reference sequences, and how these varied between phenotypes.

This stems from increasing evidence showing that TE insertions impact gene

expression, and thus the variation for a given phenotype (Bourque et al., 2018). We

found 283 highly divergent TE insertions in genic regions from our antler analysis and

116 from the body size analysis. The insertion of these highly variable sequences has

the potential to impact gene function, and studies are now starting to emerge that both

validate insertions (Lerat et al., 2018) and show evidence for positive selection (Kofler et

al., 2011). This suggests the potential for TEs to modulate phenotypes within a

population and we feel this is an important and often overlooked pattern in similar

studies. Validating the TEs and characterizing their influence on neighbouring genes is

clearly warranted given the high frequency differences between pools.

Implications for understanding of artificial and sexual selection

Antler and body size are traits that are sexually selected for and linked to reproductive

success in WTD (Newbolt et al., 2016; Morina et al., 2018), and are traits desired by

hunters, managers, and farmers (e.g. antlers for velvet or large body for meat). We

have highly diverged genomic regions, and thousands genome-wide variants with allele

frequency differences between the phenotypic extremes for these traits. We suggest the

combined effects of these variants as being the genetic drivers behind observed

phenotypic variation, consistent with a polygenic model of quantitative traits (Barghi et

al., 2020). Additional lines of evidence, specifically RNA-seq studies, should look at the

regions identified here, but also consider interactions across the entire genomic,

epigenomic, and transcriptomic landscape.

We identified and validated two of three potential QTL relating to the variation in

phenotypes for WTD. While we applied the added stringency of accounting for

overdispersion in the chi-square test, our inability to validate one outlier suggests a

small effect size and thus requiring increased sample size (see Kardos et al. 2016), or a

true type I error. Alternatively, epistatic interactions might be at play (Knief et al. 2019),

but detection requires whole-genome sequencing of many individuals thereby defeating

the purpose of pooled sequencing. The high effective population size and homogeneity

of Anticositi Island white-tailed deer implies the need for small windows, and ultimately a

SNP-based analysis. Here, expanded qPCR sample sizes might still reveal the

predicted effect, and it is encouraging that approaches for low coverage pooled data are

emerging (Tilk et al. 2019). As we continue to identify and validate candidate QTL, it is

conceivable that a gene panel could be developed that could be used in breeding and

management programs (e.g. Quality Deer Management) or assess the effects of

artificial selection by trophy hunting; however until a reasonable amount of phenotypic

variation can be attributed to specific QTL, a gene targeted approach for management

and breeding of white-tailed deer will remain a challenge.

Acknowledgments

This work was supported by Natural Sciences and Engineering Research Council of

Canada Discovery Grant (ABAS and SDC); ComputeCanada Resources for Research

Groups (ABAS); Canadian Foundation for Innovation: John R. Evans Leaders Fund

(ABAS); The Symons Trust Fund for Canadian Studies (ABAS); Trent University start-

up funds [ABAS]; and Industrial Chairs and Collaborative Research and Development

Grants from the Natural Sciences and Engineering Research Council of Canada (SDC).

We thank the outfitters of Anticosti Island and the Ministère des Forêts, de la Faune et

des Parcs du Québec for logistical help associated with fieldwork, and all of the field

assistants and technicians who collected samples and made phenotypic measurements

throughout the duration of this project. We thank four anonymous reviewers for their

extensive comments and analytical suggestions on this manuscript.

References

Anand, S., Mangano, E., Barizzone, N., Bordoni, R., Sorosina, M., Clarelli, F., … De
Bellis, G. (2016). Next Generation Sequencing of Pooled Samples: Guideline for
Variants' Filtering. Scientific Reports, 6(1), 33735.
https://doi.org/10.1038/srep33735

Balčiauskas, L., Kazlauskas, M., & Balčiauskienė, L. (2017). European bison: changes
in species acceptance following plans for translocation. European Journal of
Wildlife Research, 63(1), 1–9. https://doi.org/10.1007/s10344-016-1066-1

Bao, W., Kojima, K. K., & Kohany, O. (2015). Repbase Update, a database of repetitive
elements in eukaryotic genomes. Mobile DNA, 6(1). https://doi.org/10.1186/s13100-
015-0041-9

Barghi, N., Hermisson, J., & Schlötterer, C. (2020). Polygenic adaptation: a unifying
framework to understand positive selection. Nature Reviews Genetics, 1–13.
https://doi.org/10.1038/s41576-020-0250-z

Barnett, I. J., Lee, S., & Lin, X. (2013). Detecting rare variant effects using extreme
phenotype sampling in sequencing association studies. Genetic Epidemiology,
37(2), 142–151. https://doi.org/10.1002/gepi.21699

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical
and powerful approach to multiple testing. Journal of the Royal Statistical Society
Series B, 57(1), 289–300

Bérénos, C., Ellis, P. A., Pilkington, J. G., Lee, S. H., Gratten, J., & Pemberton, J. M.
(2015). Heterogeneity of genetic architecture of body size traits in a free-living
population. Molecular Ecology, 24(8), 1810–1830.
https://doi.org/10.1111/mec.13146

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for
Illumina Sequence Data. Bioinformatics, btu170.

Bourque, G., Burns, K. H., Gehring, M., Gorbunova, V., Seluanov, A., Hammell, M., …
Feschotte, C. (2018). Ten things you should know about transposable elements.
Genome Biology, 19(1), 199. https://doi.org/10.1186/s13059-018-1577-z

Bouwman, A. C., Garrick, D. J., Reecy, J., & Van Tassell, C. P. (2018). Meta-analysis of
genome-wide association studies for cattle stature identifies common genes that
regulate body size in mammals. Nature Genetics, 50, 362–367
https://doi.org/10.1038/s41588-018-0056-5

Boyle, E. A., Li, Y. I., & Pritchard, J. K. (2017). An Expanded View of Complex Traits:
From Polygenic to Omnigenic. Cell, 169(7), 1177–1186.
https://doi.org/10.1016/j.cell.2017.05.038

Broad Institute (2020). Picard Tools – By Broad Institute. Github.
http://broadinstitute.github.io/picard/

Bundy, R. M., Robel, R. J., & Kemp, K. E. (1991). Whole Body Weights Estimated from
Morphological Measurements of White-Tailed Deer. Transactions of the Kansas
Academy of Science (1903-), 94(3/4), 95-100. https://doi.org/10.2307/3627856

Bünger, L., Lewis, R. M., Rothschild, M. F., Blasco, A., Renne, U., & Simm, G. (2005).
Relationships between quantitative and reproductive fitness traits in animals.
Philosophical Transactions of the Royal Society B: Biological Sciences, 360(1459),
1489–1502. https://doi.org/10.1098/rstb.2005.1679

Carapito, R., Konantz, M., Paillard, C., Miao, Z., Pichot, A., Leduc, M. S., … Bahram, S.
(2017). Mutations in signal recognition particle SRP54 cause syndromic
neutropenia with Shwachman-Diamond-like features. The Journal of Clinical
Investigation, 127(11), 4090–4103. https://doi.org/10.1172/JCI92876

Cole, J. B., Wiggans, G. R., Ma, L., Sonstegard, T. S., Lawlor, T. J., Crooker, B. A., …
Da, Y. (2011). Genome-wide association analysis of thirty one production, health,
reproduction and body conformation traits in contemporary U.S. Holstein cows.
BMC Genomics, 12(1), 408. https://doi.org/10.1186/1471-2164-12-408

Coltman, D. W., O'Donoghue, P., Jorgenson, J. T., Hogg, J. T., Strobeck, C., & Festa-
Bianchet, M. (2003). Undesirable evolutionary consequences of trophy hunting.
Nature, 426(6967), 655–658. https://doi.org/10.1038/nature02177

DeYoung, R. W., Demarais, S., Gee, K. L., Honeycutt, R. L., Hellickson, M. W., &
Gonzales, R. A. (2009). Molecular Evaluation of the White-tailed Deer (Odocoileus
Virginianus) Mating System. Journal of Mammalogy, 90(4), 946–953.
https://doi.org/10.1644/08-MAMM-A-227.1

Ditchkoff, S. S., Welch, E. R., & Lochmiller, R. L. (2000). Using cast antler
characteristics to profile quality of white-tailed deer Odocoileus virginianus
populations. Wildlife Biology, 6(1), 53–58. https://doi.org/10.2981/wlb.2000.031

Edeline, E., Carlson, S. M., Stige, L. C., Winfield, I. J., Fletcher, J. M., James, J. Ben, …
Stenseth, N. C. (2007). Trait changes in a harvested population are driven by a
dynamic tug-of-war between natural and harvest selection. Proceedings of the
National Academy of Sciences of the United States of America, 104(40), 15799–
15804. https://doi.org/10.1073/pnas.0705908104

Ellegren, H. (2014). Genome sequencing and population genomics in non-model
organisms. *Trends in Ecology & Evolution, 29*(1), 51–63.
https://doi.org/10.1016/j.tree.2013.09.008

Ellegren, H., & Sheldon, B. C. (2008). Genetic basis of fitness differences in natural
populations. Nature, 452(7184), 169–175. https://doi.org/10.1038/nature06737

Emond, M. J., Louie, T., Emerson, J., Zhao, W., Mathias, R. A., Knowles, M. R., …
Bamshad, M. J. (2012). Exome sequencing of extreme phenotypes identifies
DCTN4 as a modifier of chronic Pseudomonas aeruginosa infection in cystic
fibrosis. Nature Genetics, 44(8), 886–889. https://doi.org/10.1038/ng.2344

Endler, L., Betancourt, A. J., Nolte, V., & Schlötterer, C. (2016). Reconciling differences
in pool-GWAS between populations: A case study of female abdominal
pigmentation in Drosophila melanogaster. Genetics, 202(2), 843–855.
https://doi.org/10.1534/genetics.115.183376

Environment Canada. (2006). Climate normals and averages, daily data reports of Port-
Menier's station from 1995 to 2005. Canada's National Climate Archive. Retrieved
from http://climate.weatheroffice.gc.ca

Foley, A. M., DeYoung, R. W., Hewitt, D. G., Hellickson, M. W., Gee, K. L., Wester, D.
B., … Miller, K. V. (2015). Purposeful wanderings: mate search strategies of male
white-tailed deer. Journal of Mammalogy, 96(2), 279–286.
https://doi.org/10.1093/jmammal/gyv004

Foote, A. D., Vijay, N., Ávila-Arcos, M. C., Baird, R. W., Durban, J. W., Fumagalli, M., …
Wolf, J. B. W. (2016). Genome-culture coevolution promotes rapid divergence of
killer whale ecotypes. Nature Communications, 7(1), 11693.
https://doi.org/10.1038/ncomms11693

Fuller, J., Ferchaud, A. L., Laporte, M., Le Luyer, J., Davis, T. B., Côté, S. D., &
Bernatchez, L. (2020). Absence of founder effect and evidence for adaptive
divergence in a recently introduced insular population of white-tailed deer
(Odocoileus virginianus). Molecular Ecology, 29(1), 86–104.
https://doi.org/10.1111/mec.15317

García-Ferrer, D., Herrero, J., Jimeno-Brabo, P., Arnal, M. C., García-Serrano, A., &
Fernández de Luco, D. (2019). Can roe deer hunting be selective? A case study
from the Pyrenees. Galemys, Spanish Journal of Mammalogy, 31, 27–34.
https://doi.org/10.7325/galemys.2019.a3

Gurwitz, D., & McLeod, H. L. (2013). Genome-wide studies in pharmacogenomics:
harnessing the power of extreme phenotypes. Pharmacogenomics, 14(4), 337–
339. https://doi.org/10.2217/pgs.13.35

Hamlin, K. L., D. F. Pac, C. A. Sime, R. M. DeSimone, and G. L. Dusek. 2000.
Evaluating the accuracy of ages obtained by two methods for Montana ungulates.
Journal of Wildlife Management 64, 441–449

Hernández-Montiel, W., Martínez-Núñez, M. A., Ramón-Ugalde, J. P., Román-Ponce,
S. I., Calderón-Chagoya, R., & Zamora-Bustillos, R. (2020). Genome-wide
association study reveals candidate genes for litter size traits in pelibuey sheep.
Animals, 10(3), 434. https://doi.org/10.3390/ani10030434

Heusschen, R., Griffioen, A. W., & Thijssen, V. L. (2013, August 1). Galectin-9 in tumor
biology: A jack of multiple trades. Biochimica et Biophysica Acta - Reviews on
Cancer. Elsevier. https://doi.org/10.1016/j.bbcan.2013.04.006

Hewitt, D. G. (2011). Biology and management of white-tailed deer. CRC Press.
Retrieved from https://www.crcpress.com/Biology-and-Management-of-White-
tailed-Deer/Hewitt/p/book/9781439806517

Holt, C., & Yandell, M. (2011). MAKER2: an annotation pipeline and genome-database
management tool for second-generation genome projects. BMC Bioinformatics,
12(1), 491. https://doi.org/10.1186/1471-2105-12-491

Jamieson A, Anderson SJ, Fuller J, Côté SD, Northrup JM, Shafer ABA (2020)
Heritability estimates of antler and body traits in white-tailed deer (Odocoileus
virginianus) from genomic-relatedness matrices. Journal of Heredity doi:
10.1093/jhered/esaa023

Jiang, J., Ma, L., Prakapenka, D., VanRaden, P. M., Cole, J. B., & Da, Y. (2019). A
Large-Scale Genome-Wide Association Study in U.S. Holstein Cattle. Frontiers in
Genetics, 10, 412. https://doi.org/10.3389/fgene.2019.00412

Johnson, M., Zaretskaya, I., Raytselis, Y., Merezhuk, Y., McGinnis, S., & Madden, T. L.
(2008). NCBI BLAST: a better web interface. Nucleic Acids Research, 36(Web
Server), W5–W9. https://doi.org/10.1093/nar/gkn201

Johnston SE, McEwan J, Pickering NK et al. (2011) Genome-wide associa-tion
mapping identifies the genetic basis of discrete and quantitative variation in sexual
weaponry in a wild sheep population. Molecular Ecology, 20, 2555–2566.

Johnston, S. E., Gratten, J., Berenos, C., Pilkington, J. G., Clutton-Brock, T. H.,
Pemberton, J. M., & Slate, J. (2013). Life history trade-offs at a single locus
maintain sexually selected genetic variation. Nature, 502(7469), 93–95.
https://doi.org/10.1038/nature12489

Jones, P. D., Strickland, B. K., Demarais, S., Wang, G., & Dacus, C. M. (2018). Nutrition
and ontogeny influence weapon development in a long-lived mammal. Canadian
Journal of Zoology, 96(9), 955–962. https://doi.org/10.1139/cjz-2017-0345

Kardos, M., & Shafer, A. B. A. (2018). The Peril of Gene-Targeted Conservation. Trends
in Ecology & Evolution, 33(11), 827–839. https://doi.org/10.1016/j.tree.2018.08.011

Kardos, M., Husby, A., Mcfarlane, S. E., Qvarnstrom, A., & Ellegren, H. (2016). Whole-
genome resequencing of extreme phenotypes in collared flycatchers highlights the
difficulty of detecting quantitative trait loci in natural populations. Molecular Ecology
Resources, 16(3), 727–741. https://doi.org/10.1111/1755-0998.12498

Kardos, M., Luikart, G., Bunch, R., Dewey, S., Edwards, W., McWilliam, S., … Kijas, J.
(2015). Whole-genome resequencing uncovers molecular signatures of natural and
sexual selection in wild bighorn sheep. Molecular Ecology, 24(22), 5616–5632.
https://doi.org/10.1111/mec.13415

Ker, D. F. E., Wang, D., Sharma, R., Zhang, B., Passarelli, B., Neff, N., … Yang, Y. P.
(2018). Identifying deer antler uhrf1 proliferation and s100a10 mineralization genes
using comparative RNA-seq. Stem Cell Research & Therapy, 9(1), 292.
https://doi.org/10.1186/s13287-018-1027-6

Knief U, Bossu CM, Saino N et al. (2019) Epistatic mutations under divergent selection
govern phenotypic variation in the crow hybrid zone. Nat Ecol Evol: 3: 570–576.

Kofler, R. & Schlötterer, C. (2012). Gowinda: unbiased analysis of gene set enrichment
for genome-wide association studies. Bioinformatics 28(15):2084-2085. doi:
10.1093/bioinformatics/bts315

Kofler, R., Gómez-Sánchez, D., & Schlötterer, C. (2016). PoPoolationTE2: Comparative
Population Genomics of Transposable Elements Using Pool-Seq. Molecular
Biology and Evolution, 33(10), 2759–2764. https://doi.org/10.1093/molbev/msw137

Kofler, R., Orozco-terWengel, P., De Maio, N., Pandey, R. V., Nolte, V., Futschik, A., …
Schlötterer, C. (2011). PoPoolation: A Toolbox for Population Genetic Analysis of
Next Generation Sequencing Data from Pooled Individuals. PLoS ONE, 6(1),
e15925. https://doi.org/10.1371/journal.pone.0015925

Kofler, R., Pandey, R. V., & Schlötterer, C. (2011). PoPoolation2: Identifying
differentiation between populations using sequencing of pooled DNA samples
(Pool-Seq). Bioinformatics, 27(24), 3435–3436.
https://doi.org/10.1093/bioinformatics/btr589

Korf, I. (2004). Gene finding in novel genomes. BMC Bioinformatics, 5(1), 59.
https://doi.org/10.1186/1471-2105-5-59

Kruuk, L. E. B., Slate, J., Pemberton, J. M., Brotherstone, S., Guinness, F., & Clutton-
Brock, T. (2002). Antler size in red deer: Heritability and selection but no evolution.
Evolution, 56(8), 1683–1695. https://doi.org/10.1111/j.0014-3820.2002.tb01480.x

Kuparinen, A., & Festa-Bianchet, M. (2017). Harvest-induced evolution: insights from
aquatic and terrestrial systems. Philosophical Transactions of the Royal Society B:
Biological Sciences, 372(1712), 20160036. https://doi.org/10.1098/rstb.2016.0036

Laine, V. N., Gossmann, T. I., Schachtschneider, K. M., Garroway, C. J., Madsen, O.,
Verhoeven, K. J. F., … Groenen, M. A. M. (2016). Evolutionary signals of selection
on cognition from the great tit genome and methylome. Nature Communications,
7(1), 10474. https://doi.org/10.1038/ncomms10474

Lerat, E., Goubert, C., Guirao-Rico, S., Merenciano, M., Dufour, A., Vieira, C., &
González, J. (2019). Population-specific dynamics and selection patterns of
transposable element insertions in European natural populations. Molecular
Ecology, 28(6), 1506–1522. https://doi.org/10.1111/mec.14963

Li, C., Zhao, H., Liu, Z., & McMahon, C. (2014). Deer antler – A novel model for
studying organ regeneration in mammals. The International Journal of Biochemistry
& Cell Biology, 56, 111–122. https://doi.org/10.1016/j.biocel.2014.07.007

Li, D., Lewinger, J. P., Gauderman, W. J., Murcray, C. E., & Conti, D. (2011). Using
extreme phenotype sampling to identify the rare causal variants of quantitative

traits in association studies. Genetic Epidemiology, 35(8), 790–799.
https://doi.org/10.1002/gepi.20628

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. Retrieved from http://arxiv.org/abs/1303.3997

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., … Durbin, R.
(2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics,
25(16), 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Li, J., Dani, J. A., & Le, W. (2009). The role of transcription factor Pitx3 in dopamine
neuron development and Parkinson's disease. Current Topics in Medicinal
Chemistry, 9(10), 855–859. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/19754401

Malenfant, R. M., Davis, C. S., Richardson, E. S., Lunn, N. J., & Coltman, D. W. (2018).
Heritability of body size in the polar bears of Western Hudson Bay. Molecular
Ecology Resources, 18(4), 854–866. https://doi.org/10.1111/1755-0998.12889

Mank, J. E. (2017). Population genetics of sexual conflict in the genomic era. Nature
Reviews Genetics, 18(12), 721–730. https://doi.org/10.1038/nrg.2017.83

Marouli, E., Graff, M., Medina-Gomez, C., Lo, K. S., Wood, A. R., Kjaer, T. R., … Lettre,
G. (2017). Rare and low-frequency coding variants alter human adult height.
Nature, *542*(7640), 186–190. https://doi.org/10.1038/nature21039

Michel, E. S., Demarais, S., Strickland, B. K., Smith, T., & Dacus, C. M. (2016). Antler
characteristics are highly heritable but influenced by maternal factors. The Journal
of Wildlife Management, 80(8), 1420–1426. https://doi.org/10.1002/jwmg.21138

Morina, D. L., Demarais, S., Strickland, B. K., & Larson, J. E. (2018). While males fight,
females choose: male phenotypic quality informs female mate choice in mammals.
Animal Behaviour, 138, 69–74. https://doi.org/10.1016/J.ANBEHAV.2018.02.004

Mruk, D. D., & Cheng, C. Y. (2011, January 15). The myotubularin family of lipid
phosphatases in disease and in spermatogenesis. Biochemical Journal. NIH Public
Access. https://doi.org/10.1042/BJ20101267

Neethiraj, R., Hornett, E. A., Hill, J. A., & Wheat, C. W. (2017). Investigating the
genomic basis of discrete phenotypes using a Pool-Seq-only approach: New
insights into the genetics underlying colour variation in diverse taxa. Molecular
Ecology, 26(19), 4990–5002. https://doi.org/10.1111/mec.14205

Newbolt, C. H., Acker, P. K., Neuman, T. J., Hoffman, S. I., Ditchkoff, S. S., & Steury, T.
D. (2016). Factors Influencing Reproductive Success in Male White-Tailed Deer.
Journal of Wildlife Management 81(2), 206-217 https://doi.org/10.1002/jwmg.21191

Nilsen, E. B., and E. J. Solberg. (2006). Patterns of hunting mortality in Norwegian
moose (Alces alces) populations. European Journal of Wildlife Research 52:153–
163

Ozato K, Shin DM, Chang TH, Morse HC (2008). TRIM family proteins and their
emerging roles in innate immunity. Nat. Rev. Immunol. 8 (11): 849–60.
doi:10.1038/nri2413

Pagani, F., & Baralle, F. E. (2004). Genomic variants in exons and introns: identifying
the splicing spoilers. Nature Reviews Genetics, 5(5), 389–396.
https://doi.org/10.1038/nrg1327

Park, H. J., Lee, D. H., Park, S. G., Lee, S. C., Cho, S., Kim, H. K., … Park, B. C.
(2004). Proteome analysis of red deer antlers. PROTEOMICS, 4(11), 3642–3653.
https://doi.org/10.1002/pmic.200401027

Pavlidis, P., Jensen, J. D., Stephan, W., & Stamatakis, A. (2012). A Critical Assessment
of Storytelling: Gene Ontology Categories and the Importance of Validating
Genomic Scans. Molecular Biology and Evolution, 29(10), 3237–3248.
https://doi.org/10.1093/molbev/mss136

Perez-Gracia, J.P., Ruiz-Ilundain, M.G, Garcia-Ribas, I., & Carrasco, E.M. (2002). The
role of extreme phenotype selection studies in the identification of clinically relevant
genotypes in cancer research. Cancer, 95(7), 1605–1610.
https://doi.org/10.1002/cncr.10877

Potvin, F., & Breton, L. (2005). From the Field: Testing 2 aerial survey techniques on
deer in fenced enclosures—visual double-counts and thermal infrared sensing.
Wildlife Society Bulletin, 33(1), 317–325. https://doi.org/10.2193/0091-
7648(2005)33[317:FTFTAS]2.0.CO;2

Pozo, R. A., Schindler, S., Cubaynes, S., Cusack, J. J., Coulson, T., & Malo, A. F.
(2016). Modeling the impact of selective harvesting on red deer antlers. The
Journal of Wildlife Management, 80(6), 978–989.
https://doi.org/10.1002/jwmg.21089

Prince, D. J., O'Rourke, S. M., Thompson, T. Q., Ali, O. A., Lyman, H. S., Saglam, I. K.,
… Miller, M. R. (2017). The evolutionary basis of premature migration in Pacific
salmon highlights the utility of genomics for informing conservation. Science
Advances, 3(8), e1603198. https://doi.org/10.1126/sciadv.1603198

Ramanzin, M., & Sturaro, E. (2014). Habitat quality influences relative antler size and
hunters' selectivity in roe deer. European Journal of Wildlife Research, 60(1), 1–10.
https://doi.org/10.1007/s10344-013-0744-5

Robinson, M.R. et al. (2013) Partitioning of genetic variation across the genome using multimarker methods in a wild bird population. Mol. Ecol. 22, 3963–3980

Rogers, A. N., Vanburen, D. G., Hedblom, E., Tilahun, M. E., Telfer, J. C., & Baldwin, C. L. (2005). Function of ruminant γδ T cells is defined by WC1.1 or WC1.2 isoform expression. In Veterinary Immunology and Immunopathology (Vol. 108, pp. 211–217). Vet Immunol Immunopathol. https://doi.org/10.1016/j.vetimm.2005.08.008

Santure, A. W., & Garant, D. (2018). Wild GWAS-association mapping in natural populations. Molecular Ecology Resources, 18(4), 729–738. https://doi.org/10.1111/1755-0998.12901

Santure, A. W., De Cauwer, I., Robinson, M. R., Poissant, J., Sheldon, B. C., & Slate, J. (2013). Genomic dissection of variation in clutch size and egg mass in a wild great tit (Parus major) population. Molecular Ecology, 22(15), 3949–3962. https://doi.org/10.1111/mec.12376

Schlötterer, C., Tobler, R., Kofler, R., & Nolte, V. (2014). Sequencing pools of individuals — mining genome-wide polymorphism data without big funding. Nature Reviews Genetics, 15(11), 749–763. https://doi.org/10.1038/nrg3803

Schoenebeck, C. W., & Peterson, B. C. (2014, June 1). Evaluation of hunter antler-size selection through an age-specific comparison of harvested and naturally cast antler metrics. Journal of Fish and Wildlife Management. U.S. Fish and Wildlife Service. https://doi.org/10.3996/032013-jfwm-022

Simard, M. A., T. Coulson, A. Gingras, and S. D. Côté. (2010). Influence of density and climate on the population dynamics of a large herbivore under harsh environmental conditions. Journal of Wildlife Management, 74, 1671–1685

Simard, M.-A., J. Huot, S. de Bellefeuille and S. D. Côté. (2014). Influences of habitat composition, plant phenology, and population density on autumn indices of body condition in a northern white-tailed deer population. Wildlife Monographs 187, 1-28

Smit, AFA, Hubley, R & Green, P. (2019) RepeatMasker Open-4.0. 2013-2015. Retrieved from http://www.repeatmasker.org

Smit, AFA, Hubley, R. (2019) RepeatModeler Open-1.0. 2008-2015 retrieved from http://www.repeatmasker.org

Solberg, E. J., A. Loison, J.-M. Gaillard, and M. Heim. (2004). Lasting effects of conditions at birth on moose body mass. Ecography 27:677–687

Stinchcombe, J. R. & Hoekstra, H. E. (2008). Combining population genomics and quantitative genetics: Finding the genes underlying ecologically important traits. *Heredity*, 100(2), 58–170 https://doi.org/10.1038/sj.hdy.6800937

Supek, F., Bošnjak, M., Škunca, N., & Šmuc, T. (2011). REVIGO Summarizes and Visualizes Long Lists of Gene Ontology Terms. PLoS ONE, 6(7), e21800. https://doi.org/10.1371/journal.pone.0021800

Tilk, S., Bergland, A., Goodman, A., Schmidt, P., Petrov, D., & Greenblum, S. (2019). Accurate allele frequencies from ultra-low coverage Pool-seq samples in evolve-and-resequence experiments. G3: Genes, Genomes, Genetics, 9(12), 4159–4168. https://doi.org/10.1534/g3.119.400755

Treangen, T. J., & Salzberg, S. L. (2012). Repetitive DNA and next-generation sequencing: computational challenges and solutions. Nature Reviews Genetics, 13(1), 36–46. https://doi.org/10.1038/nrg3117

Vanpé, C., Gaillard, J. M., Kjellander, P., Mysterud, A., Magnien, P., Delorme, D., … Hewison, A. J. M. (2007). Antler size provides an honest signal of male phenotypic quality in roe deer. American Naturalist, 169(4), 481–493. https://doi.org/10.1086/512046

Visscher PM, Macgregor S, Benyamin B et al. (2007) Genome partitioning of genetic variation for height from 11,214 sib- ling pairs. American Journal of Human Genetics, 81, 1104–1110

Wang, Y., Zhang, C., Wang, N., Li, Z., Heller, R., Liu, R., … Qiu, Q. (2019). Genetic basis of ruminant headgear and rapid antler regeneration. Science, 364(6446), eaav6335. https://doi.org/10.1126/science.aav6335

Watanabe, K., Stringer, S., Frei, O., Mirkov, M., de Leeuw, C., Polderman, T. J. C., … Posthuma, D. (2019). A global overview of pleiotropy and genetic architecture in complex traits. Nature Genetics, 51(9), 1339–1348. https://doi.org/10.1038/s41588-019-0481-0

Whitlock, M. C., & Lotterhos, K. E. (2015). Reliable Detection of Loci Responsible for Local Adaptation: Inference of a Null Model through Trimming the Distribution of F ST. The American Naturalist, 186(S1), S24–S36. https://doi.org/10.1086/682949

Williams, J. D., Krueger, W. F., & Harmel, D. H. (1994). Heritabilities for antler characteristics and body weight in yearling white-tailed deer. Heredity, 73(1), 78–83. https://doi.org/10.1038/hdy.1994.101

Zhang, F., & Lupski, J. R. (2015). Non-coding genetic variants in human disease: Figure 1. Human Molecular Genetics, 24(R1), R102–R110. https://doi.org/10.1093/hmg/ddv259

Data Accessibility

Raw sequence data FastQ files are available on the Sequence Read Archive

(Accession: PRJNA576136). All bioinformatic and analytical code available on GitLab

(https://gitlab.com/WiDGeT_TrentU/PoolSeq). Raw data table have been uploaded to

Dryad (Dryad Accession no. XXXXXX)

Author Contributions

S.J.A., S.D.C., and A.B.A.S. designed the study; S.D.C., J.H.R., and S.J.A. coordinated

data collection and sample curation; S.J.A. performed research and analyzed data;

S.J.A. wrote the manuscript with input from J.H.R., S.D.C., and A.B.A.S.

Scaffold	Window Position	*p*	Gene	Function
ref0001370	16015000	2.66E-37	LGALS9	The galectins are a family of beta-galactoside-binding proteins implicated in modulating cell-cell and cell-matrix interactions.
ref0001836	5479500	1.24E-34	MTMR2	This gene is a member of the myotubularin family of phosphoinositide lipid phosphatases.
ref0000845	16534000	5.97E-33	WC1.1	The function of ruminant gammadelta T cells is defined by WC1.1. This suggests a role in host response to cell proliferation.
ref0002391	7908000	1.75E-29	ITLN2	ITLN2 (Intelectin 2) is likely involved include carbohydrate binding.
ref0001924	2123500	1.52E-28	TRIM64	TRIM proteins are involved in pathogen-recognition and by regulation of transcriptional pathways in host defence (Ozato et.al., 2008).
ref0000845	16751500	1.14E-27	WC1.1	See above.
ref0002391	7907500	1.79E-27	ITLN2	See above.
ref0000505	9079500	1.37E-26	ATP5F1A	This gene encodes a subunit of mitochondrial ATP synthase.
ref0000881	8059500	1.91E-26	MYH9	This gene encodes a conventional non-muscle myosin.
ref0002644	7312000	2.23E-26	RIMS1	The protein encoded by this gene is a RAS gene superfamily member that regulates synaptic vesicle exocytosis.

Scaffold	Window Position	p	Gene	Function
ref0002232	15310000	1.71E-31	FLVCR2	This gene encodes a transmembrane protein that is a calcium transporter.
ref0000892	4172500	4.82E-29	SLC7A3	SLC7A3 appears to be involved in amino acid transmembrane transporter activity and basic amino acid transmembrane transporter.
ref0002547	2917000	2.98E-23	KRR1	KRR1 (KRR1 Small Subunit Processome Component Homolog) appears to be involved win cytosol and Gene Expression.
ref0000845	16369000	1.26E-20	WC1.1	The function of ruminant gammadelta T-cells is defined by WC1.1. This suggests a role in host response to cell proliferation.
ref0000438	13320500	1.64E-20	Uncharacterized	Function unknown.
ref0002813	16028000	8.08E-20	DOCK1	This gene encodes a member of the dedicator of cytokinesis protein family.
ref0002788	19947500	2.45E-19	SYK	This gene encodes a member of the family of non-receptor type Tyr protein kinase that is involved in coupling activated immunoreceptors to downstream signaling events that mediate diverse cellular responses, including proliferation, differentiation, and phagocytosis.
ref0000529	431000	4.31E-19	TRDV1	T cell receptors recognize foreign antigens which have been processed as small peptides and bound to major histocompatibility complex (MHC) molecules at the surface of antigen presenting cells (APC).
ref0002645	24851500	8.41E-19	Uncharacterized LOC110125691	Detected in other ruminants, function unknown.
ref0002547	2916500	1.27E-18	KRR1	See above.

Gene Name	Scaffold	Antler (p)	Body (p)	Function
OLIGI1	ref0002192	4.08E-14	5.59E-10	Among its related pathways are Neural Crest Differentiation and Neural Stem Cells and Lineage-specific Markers.
TWIST1	ref0000889	4.29E-01	4.27E-10	This gene encodes a basic helix-loop-helix (bHLH) transcription factor that plays an important role in embryonic development.
HMGA2	ref0000505	1.57E-11	3.28E-01	HMG proteins function as architectural factors and are essential components of the enhancesome.
IGF1	ref0000881	1.40E-04	6.40E-12	The protein encoded by this gene is similar to insulin in function and structure and is a member of a family of proteins involved in mediating growth and development.