Sex-specific effects of an IgE polymorphism on immunity, susceptibility to infection and reproduction in a wild rodent

Klara M Wanelik[1*,] Mike Begon[1], Janette E Bradley[2], Ida M Friberg[3], Joseph A Jackson[3], Christopher H Taylor[2], Steve Paterson[1]

[1] Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, UK

[2] School of Life Sciences, University of Nottingham, Nottingham, UK

[3] School of Environment and Life Sciences, University of Salford, Salford, UK

* Corresponding author

Current address: Department of Zoology, University of Oxford, Oxford, UK

Email: klara.wanelik@zoo.ox.ac.uk

Abstract

The genotype of an individual is an important predictor of their immune function, and subsequently, their ability to control or avoid infection and ultimately contribute offspring to the next generation. However, the same genotype, subjected to different environments, can also result in different outcomes. The sexes represent two such different environments. Sexual dimorphism is widespread across the animal kingdom. Despite this, very little is known about the importance of sex for the expression of genotype in the context of health and disease, particularly in natural populations. We identified a synonymous polymorphism in the high-affinity Immunoglobulin E (IgE) receptor (GC and non-GC haplotypes) that has sex-specific effects on immune gene expression, susceptibility to infection and reproductive success of individuals in a natural population of field voles (*Microtus agrestis*). We found that the effect of the GC haplotype on the expression of genes affecting inflammation displayed a significant interaction with sex. While males with the GC haplotype had upregulated pro-inflammatory genes, in particular the pro-inflammatory cytokine *Il33,* females had upregulated anti-inflammatory genes, in particular the anti-inflammatory cytokine inhibitor *Socs3.* Furthermore we found that the effect of the GC haplotype on the probability of infection with a common microparasite*, Babesia microti,* displayed a significant interaction with sex. While males with the GC haplotype did not differ significantly in their susceptibility to infection, females with the GC haplotype were more likely to be infected. Finally, we found that the effect of the GC haplotype on reproductive success also displayed a significant interaction with sex. While males with the GC haplotype had a lower reproductive success, females with the GC haplotype did not differ in the number of offspring they produced. To our knowledge, this is the first time that a polymorphism with sex-specific

effects across all three levels (immune gene expression, susceptibility to infection and

reproductive success) has been documented in a natural population.

<u>Introduction</u>

In order for an individual to control or avoid infection and ultimately contribute offspring to

the next generation, they must have a well-functioning immune system [1]. An

individual's immune function is in part determined by their genotype (e.g. [2–4]).

However, the effect of any genotype will depend on the context in which it occurs.

Sexual dimorphism is widespread in the animal kingdom, with males and females

representing different contexts or environments within which genes must act. Sex can

therefore have an important effect on the phenotypic expression of genotype (genotype

by sex interactions), with the same genotype having either (i) effects in the same

direction but of different magnitude in the sexes, (ii) an effect in one sex but not the

other, or in the most extreme case (iii) opposing effects in the sexes. However, the

majority of evidence for genotype by sex interactions, in the context of health and

disease, comes from laboratory experiments conducted on model organisms [5,6].

Evidence for genotype by sex interactions in natural populations is limited. Studies of

natural populations have explored the effects of genotype on immune phenotype, and

have observed consequences for susceptibility to infection. Most notably, variability in

the genes of the Major Histocompatibility Complex (MHC) has been associated with

resistance to intestinal nematodes in domestic sheep [2] and with resistance to malaria,

hepatitis, and AIDS in humans [7–9]. The role of variability elsewhere in the genome, for

shaping immune phenotype, has also been studied [3,4]. However, in the majority of

these studies, genotype by sex interactions are not considered. In some studies, this is

because their low sample sizes make it difficult to test for data-hungry genotype by sex

interactions [10]. In some studies, sex is considered a confounding variable which is

controlled for, in order not to hide any subtle genetic associations [2]. Other studies

focus on a single sex for the sake of simplicity [11].

The few studies that do consider genotype by sex interactions do not tell the whole story.

While there are examples of gene variants with sex-specific effects on human immunity

and/or health in the medical literature (reviewed in [12]), longitudinal studies involving

large numbers of participants are still rare, making it difficult to explore long-term sex-

specific effects. The study of wild animals provides us with an opportunity to (more easily

and cheaply) follow large numbers of individuals, exposed to a challenging environment

and with varying genetic backgrounds, throughout their lives. This allows us to make a

more complete assessment of the impacts of genotype throughout the life of an

individual, whether male or female. For example, Graham et al 2010 [13] found evidence

for heritable variation in immunity associated with sex-specific effects on Soay sheep

reproduction. However, we know of no documented example of a polymorphism with

sex-specific effects on immunity, susceptibility to infection *and* reproduction in a natural

population. Here, we use a wild rodent, the field vole (*Microtus agrestis*), as a model in

which to do this. Wild rodents, in particular, offer an opportunity to quickly follow large

numbers of individuals throughout their lives, given their short lifespans. They also offer

the opportunity to draw on the immunological and genetics resources developed for

laboratory rodents, while providing a much more realistic ecological model of human

populations [14,15].

The sexes differ not only in their behaviour and morphology, but less conspicuously, in

their physiology, biochemistry and in their gene expression. Driven in part by sex

hormones [16], sex-biases in the expression and coexpression of many autosomal

genes have now been documented, suggesting that males and females differ not only in

their regulation of single genes [17–19], but whole networks of interacting genes [20,21].

The same genotype then, could be expressed as different immune phenotypes, and

subsequently have very different consequences for the susceptibility to infection and

reproductive success of males and females. Furthermore, males and females differ in

the environments they inhabit and the selective pressures they face, all of which may

also interact with genotype. One might therefore expect genotype by sex interactions to

be even more important in natural populations. We argue then that by neglecting to

consider sex, particularly in studies of natural populations, there is a danger of diluting

the effect of genetic variation if for example an effect is present in only one sex, or

worse, overlooking the effect of genetic variation completely if opposing effects in males

and females cancel each other out.

In humans, immunoglobulin E (IgE) mediated responses are associated with defence

against helminths [22] and with allergy [23]. They are controlled by the high-affinity IgE

receptor, FCER1, which is found on the surface of various immune effector cells e.g.

mast cells, basophils and eosinophils [24]. Naturally occurring polymorphisms in FCER1

are known to affect an individual's serum IgE levels, with consequences for their

susceptibility to infection [25,26] and their risk of developing inflammatory disease [27–

30]. Furthermore, sex differences in serum IgE levels [31] and the incidence of IgE-

mediated inflammatory disease [32] have been documented in humans, suggesting that

any polymorphism in this pathway is likely to experience different contexts in males and

females. Indeed, one study found evidence for a sex-specific effect of a polymorphism in

the *Fcer1a* gene (the alpha chain of FCER1) on susceptibility to systematic lupus

erythematosus (a chronic inflammatory disease) [33].

In a previous study of a natural population *M. agrestis*, we found that males carrying the

GC haplotype of the *Fcer1a* gene expressed the transcription factor GATA3 at a lower

level than males carrying non-GC haplotypes [3]. GATA3 is a biomarker of tolerance to

macroparasites in mature males in our population (macroparasite infection gives rise to

increased expression of GATA3, which gives rise to improved body condition and

survival [11]). Here, we explore the effects of this GC haplotype further in males, as well

as extending our analysis to females. In doing so, we provide the first example of a

polymorphism with sex-specific effects not only on immune gene expression, but

susceptibility to infection, and ultimately, on reproductive success in a natural population.

Results

We sampled a natural population of *M. agrestis* in Kielder Forest, Northumberland,

United Kingdom, over three years (2015-2017) and across seven different sites. Our

study involved a cross-sectional component (n = 317 destructively sampled voles) and a

longitudinal component (n = 850 marked individuals monitored through time, with n =

2,387 sampling points). We tested the consequences of the GC haplotype of the *Fcer1a*

gene using both cross-sectional and longitudinal components of our study. Two other

major haplotypes were present in our study population: AC haplotype and AT haplotype.

The GC haplotype has sex-specific effects on inflammation

In humans, naturally occurring polymorphisms in the *Fcer1a* gene have previously been

linked to inflammatory disease [27–30]. Therefore, we used the cross-sectional

component of our study to test the effects of the GC haplotype on inflammation in males

and females. Differential gene expression (DGE) analysis performed on 53 males and 31

females assayed by RNASeq, showed that the identity of top-responding inflammation-

associated (pro- and anti-inflammatory) genes differed between the sexes. In males, the

top-responding gene was the pro-inflammatory cytokine, *Il33* (log fold change (logFC) =

2.76, p = 0.00000015, q = 0.00048; Table S1) while in females it was the anti-

inflammatory suppressor of cytokine signalling *Socs3* (logFC = 1.07, p = 0.000061, q =

0.05; Table S2)*.* Looking at the ranking of each top-responding gene in the opposite sex

strengthens the case for distinct effects in males and females, with *Il33* ranked markedly

lower in females (rank = 8224/12904, logFC = 0.29, p = 0.70, q = 1.00) and *Socs3*

markedly lower in males (rank = 10886/12904, logFC = -0.05, p = 0.84, q = 1.00).

To test whether these sex-specific effects on inflammation were limited to these genes or

were more widespread, we performed a gene set enrichment analysis (GSEA) which

looked at the rankings of all pro-/anti-inflammatory genes. This analysis showed that

both males and females upregulated their expression of both pro- and anti-inflammatory

genes. However, males showed a stronger signal for pro-inflammatory genes (pro-

inflammatory genes: p = 0.007; anti-inflammatory genes: p = 0.04; Fig 1A, upper panel)

while females showed a stronger signal for anti-inflammatory genes (anti-inflammatory

genes: p = 0.001; pro-inflammatory genes: p = 0.04; Fig 1A, lower panel).

Finally, to confirm these sex-specific effects on inflammation, we used an independent

dataset for males and females whose spleens were stimulated with an immune agonist

and assayed by Q-PCR (for a panel of 18 immune genes). From this *ex vivo* assay, one

can gain insight into the types of immune response that could be made to a pathogen *in*

vivo (see Methods for details). We found that the effect of the GC haplotype on the

stimulated expression levels of *Il17a*, a pro-inflammatory cytokine, displayed a significant

interaction with sex (genotype by sex interaction in follow-up LMM, LRT $p = 0.001$).

While females with the GC haplotype expressed lower levels of *Il17a* (estimate = -0.12,

95% CI = -0.19 – -0.04), males with the GC haplotype did not differ in their expressed

levels of *Il17a* (estimate = 0.03, 95% CI includes zero = -0.02 – 0.08; Table S3 & S4).

The GC haplotype has the same effects on oxidative stress in both sexes

Inflammation and oxidative stress are closely related and often lead to pathology [34]. To

test whether the effects of the GC haplotype on inflammation may be having knock-on

effects on oxidative stress, we again used the cross-sectional component of our study.

We tested the effect of the GC haplotype on antioxidant superoxide dismutase 1 (SOD1)

enzymatic activity in males and females. SOD1 is one of the main antioxidative enzymes

within the antioxidative enzyme system [35]. We found that (irrespective of sex)

individuals with the AT haplotype had lower SOD1 activity when compared with other

haplotypes, including the GC haplotype ($p = 0.04$; Table S5).

The GC haplotype has sex-specific effects on the susceptibility to infection

We then used the longitudinal component of our study to test the effects of the GC

haplotype for the probability of infection with common parasites known to infect our study

population: two micro-parasites *Babesia microti* and *Bartonella* spp. and a range of

macro-parasites (ticks, fleas and cestodes). We found an effect of the GC haplotype on

the probability of infection with *B. microti* which displayed a significant interaction with

sex (genotype by sex interaction, LRT *p* = 0.048; Table S6). While females with the GC

haplotype were more likely to be infected with *B. microti* (odds ratio = 1.25, 95% CI =

0.44 – 2.07 ; Fig 1B, lower panel), males with the GC haplotype did not differ in their

susceptibility to infection (odds ratio = 0.14, 95% CI includes zero = -0.76 – 1.03; Fig 1B,

upper panel). Howevere, we found no effect of the haplotype (interactive or not) on the

probability of infection with the other parasites in our population (Tables S7, S8 & S12)

The GC haplotype has sex-specific effects on reproductive success

We genotyped both cross-sectional and longitudinal voles for 114 SNPs and used this

dataset to construct a pedigree. We estimated each individuals' reproductive success by

counting the number offspring it produced according to this pedigree. We then tested the

effects of the GC haplotype on this measure of reproductive success. We tried to run a

single model with both sexes (as above) but resulting residual variances differed

significantly between the sexes, which reflects the fact that male and female

reproduction are inherently different traits. This made it impossible to formally test for a

genotype by sex interaction and so instead we ran a separate model for each sex. We

found that males with the GC haplotype had an average of 2.2 fewer offspring than

males without the haplotype (*p* = 0.04; Table S9; Fig. 1C, upper panel). Despite a larger

sample size and lower variability in reproductive success, females with the GC haplotype

did not differ significantly in their number of offspring (genotype term did not appear in

best model; Table S10; Fig 1C, lower panel). This is suggestive of a significant genotype

by sex interaction.

Figure 1 Sex-specific effects of GC haplotype (hGC). Upper panel: males. Lower panel:

females: (A) Immune gene expression: Barcode plots indicating the position of pro- and

anti-inflammatory genes in a ranked list of log fold changes (logFC) for individuals with

(hGC+) vs. without (hGC-) the haplotype, showing that males with the haplotype have a

pro-inflammatory bias, whereas females have an anti-inflammatory bias. (B)

Susceptibility to infection: Association between hGC and the probability of infection with

Babesia microti, showing that only females with the haplotype have an increased

susceptibility to infection. (C) Reproduction: Association between hGC and reproductive

success, showing that only males with the haplotype have lower reproductive success

(error bars represent ± standard error; * indicates $p < 0.05$; ** indicates $p < 0.01$).

Discussion

In this study, we describe the first example of a polymorphism with sex-specific effects

linking together immune gene expression, susceptibility to infection and ultimately

reproductive success in a natural population. Here, we have found evidence for

opposing effects in the sexes (unstimulated immune gene expression), and an effect

present in one sex but not the other (stimulated immune gene expression, susceptibility

to infection and reproductive success), what could be called qualitative rather than

quantitative sex differences. However, we cannot rule out that, in the case of

susceptibility to infection and reproductive success, there was a smaller effect present in

the opposite sex (in the same or opposite direction) which our analysis did not have

sufficient power to detect.

The polymorphism in question lies in the high-affinity receptor for IgE. In humans,

naturally occurring polymorphisms in this receptor have previously been linked to

inflammatory disease [27–30]. Our work is consistent with this. More specifically though,

our results suggest opposing effects of this polymorphism on the inflammatory

phenotype of males and females. While males with the GC haplotype showed a pro-

inflammatory bias, females with the haplotype showed an anti-inflammatory bias. Again,

this is consistent with previous work. A previous study on another polymorphism in the

same gene, *Fcer1a*, also found evidence for sex-specific effects on inflammatory

disease [33]. In order to fully understand this genotype by sex interaction, it is important

to consider background differences in the inflammatory phenotypes of males and

females. Females in this study were pregnant for a large proportion of their lives (57% of

post-juvenile females were pregnant or lactating at the time of capture in the longitudinal

component of our study). In mammals, including humans, pregnancy has been shown to

be a largely anti-inflammatory state, with pregnant females suppressing inflammation.

This is to protect the foetus from attack by the mother's immune system [36]. While

testosterone in males has an anti-inflammatory effect [37], it also drives males to use

more space [38] and to be more aggressive [39]. These behaviours increase their rates

of contact with other individuals, and with their environment, and hence their likelihood of

infection and subsequent immune stimulation. We suggest that the polymorphism

appears may be exaggerating background sex differences in inflammatory activity. Sex

hormones have been implicated in the differential expression of some autosomal genes

in males and females [16], and could be driving these opposing effects.

Consistent with the association between IgE-mediated responses and defence against

helminths [22], in our previous study, we showed that males with the GC haplotype had

a lower level of an immunological marker of tolerance to ticks, fleas and adult cestodes

(i.e., they were less tolerant) [3]. Here, we use data on infection incidence to show that,

in a different context, the same haplotype is also associated with resistance to a

microparasite, *B. microti*, in females. Although observational studies of this kind are not

able to identify causal relationships, a realistic scenario is that females with the GC

haplotype are more likely to be infected with *B. microti* (i.e., they are less resistant) as a

result of a lower pro-inflammatory to anti-inflammatory cytokine ratio. Pro-inflammatory

cytokines (e.g., IL-6, IFN-γ, and TNF-α) may help to resist *B. microti* infection [40]. A lack

or imbalance of these cytokines may hamper this resistance. The panel of parasites that

we have measured is not exhaustive and previous studies have highlighted the

important role of species interactions in the parasite community in driving infection risk in

this study population [41]. *B. microti* may therefore represent a community of parasites,

to which this haplotype affects resistance in females.

There are major challenges associated with modelling the predictors of reproductive

success. As well as being non-normal and zero-inflated, distributions of reproductive

success are highly skewed – with an uneven distribution of success among individuals.

In polygynous species, like *M. agrestis*, reproductive skew is higher in males due to

intense competition for females. While we were able to account for non-normality and

zero-inflation using a quasi-Poisson error distribution, it is perhaps not surprising that we

were unable to fit a single model for reproductive success (and to formally test for a

genotype by sex interaction) due to unequal residual variances for males and females

(heteroscedasticity). Instead we ran a separate model for each sex. Nevertheless, we

argue that our data are suggestive of a significant genotype by sex interaction. While we

found a significant negative effect of the GC haplotype on reproductive success in

males, no significant effect was found in females despite a larger sample size and lower

reproductive skew (or variance) which should equate to higher statistical power.

A realistic scenario is that the larger cost incurred by males with the GC haplotype due to

inflammation reduced their reproductive value, as reproducing and mounting a pro-

inflammatory response are both costly activities [42]. Another realistic scenario is that

the positive effect of the GC haplotype on levels of oxidative stress (as indicated by a

tendency for higher SOD1 activity) may have had a more detrimental effect in males,

damaging sperm and reducing fertilising success. Sperm competition can be an

important feature in the reproduction of microtines, e.g., meadow vole (*M.*

pennsylvanicus) [43], with males having to produce many sperm of high quality in order

to successfully outcompete other males. Sperm are also highly susceptible to damage

by reactive oxygen species (ROS) [44]. In addition, the heightened level of oxidative

stress may have impaired olfactory sexual signals, e.g., excretion of major urinary

proteins (MUPs) making males less attractive to females and negatively impacting on

mating success [45].

Despite the immune system playing an important role in determining the health and

reproductive fitness of an individual, natural selection has not converged on a single

immune optimum. Instead, individuals in natural populations vary widely in their

response to infection. Understanding why immunoheterogeneity is maintained is a key

question in eco-immunology. Antagonistic selection, where a mutation is beneficial in

some environmental contexts and harmful in others, is thought to be one mechanism by

which balancing selection is generated, and genetic variation in immunity can be

maintained within natural populations [13]. In samples collected between 2008-2010, we

previously identified the three major haplotypes at this locus, GC, AC and AT, at

frequencies of 0.12, 0.76 and 0.07 respectively [3]. Seven years on, these frequencies

have remained relatively unchanged (0.08, 0.81 and 0.10). The fact that the GC

haplotype remains in the population, despite its detrimental effects on male reproductive

success and female infection susceptibility, suggests that it may be under balancing

selection.

In order to be under balancing selection though, the GC haplotype would need to confer

some advantage. We were not able to find evidence for any such advantage in this

study, but we can speculate on one. Microtines show cyclical population dynamics,

alternating between a "peak" phase (associated with good conditions and high vole

densities) and a "crash" phase (poor conditions and low vole densities). Selection

pressures acting during these two phases will be very different. We found that GC

haplotype was associated with a pro-inflammatory bias in males, indicating higher

investment in promoting inflammation. Thus, one possible scenario is that higher

investment in promoting inflammation is disadvantageous in the "peak" phase when the

risk of parasitism is low due to dilution effects, and it pays not to waste energy mounting

a strong immune response that is unnecessary. However, it might be advantageous in

the "crash" phase when conditions are poor, and the risk of parasitism is high due to the

time lagged transmission of parasite infectious stages that have accumulated during the

peak phase. Vole numbers were extremely low in our study area between 2012 and

2013 (indicating a "crash phase"). In 2014, vole numbers began to increase and by the

beginning of 2015 (the start of our study) had reached a peak. Although vole numbers

declined throughout our study period, they never reached the extremely low numbers

recorded in 2012-2013 (X. Lambin, personal communication, 2016). This suggests that

we sampled in the region of a "peak" phase, when this investment was disadvantageous,

and that if we sampled during a true "crash" phase we might detect an advantage to the

GC haplotype in males. Our future work will investigate this possibility further.

There is a growing interest in human genomic (or precision) medicine, with the potential

to use a patient's genotypic information to personalise their treatment. What we have

shown here demonstrates that considering genotype in isolation can be misleading, as

the same polymorphism can have different outcomes not only for the immune gene

expression, but the susceptibility to infection and ultimate reproductive success of males

and females.

Materials and Methods

We studied *M. agrestis* in Kielder Forest, Northumberland, UK, using live-trapping of individual animals from natural populations. Trapping was performed from 2015-2017 across seven different sites, each a forest clear-cut. Access to the sites was provided by the Forestry Commission. At each site, 150-197 Ugglan small mammal traps (Grahnab, Gnosjo, Sweden) were laid out in a grid spaced approximately 3-5 m apart. Our study was divided into longitudinal and cross-sectional components.

Ethics statement

All animal procedures were performed with approval from the University of Liverpool Animal Welfare Committee and under a UK Home Office licence (PPL 70/8210 to S.P.).

Longitudinal data

Every other week, traps were checked every day, once in the morning and once in the evening. Newly trapped field voles were injected with a Passive Integrated Transponder (PIT) tag (AVID, Uckfield, UK) for unique identification. This approach allowed us to build up a longitudinal record for voles that were caught on multiple occasions. A total of 850 voles were individually marked in this way. We also took a small tissue sample from the tail, for genotyping of individuals and a drop of blood from the tail which we put into 500 ml of RNAlater (Fisher Scientific, Loughborough, UK) for use in parasite detection (see below).

Parasite detection

We quantified infections by microparasites (*Babesia microti* and *Bartonella* spp.) in blood

samples taken from longitudinal animals using SYBR green based two-step reverse

transcription quantitative PCR (Q-PCR) targeted at pathogen ribosomal RNA genes.

Expression values were normalized to two host endogenous control genes. Blood

samples were derived from tail bleeds. RNA was extracted from blood samples stored in

RNAlater at -70°C using the Mouse RiboPure Blood RNA Isolation Kit (ThermoFisher,

Waltham, Massachusetts), according to manufacturer's instructions, and DNAse treated.

It was then converted to cDNA using the High-Capacity RNA-to-cDNA™ Kit

(ThermoFisher), according to manufacturer's instructions. *B. microti* primer sequences

targeting the 18S ribosomal RNA gene were as follows CTACGTCCCTGCCCTTTGTA

(forward primer sequence) and CCACGTTTCTTGGTCCGAAT (reverse primer

sequence). *Bartonella* spp. primer sequences targeting the 16S ribosomal RNA gene

were as follows GATGAATGTTAGCCGTCGGG (forward primer sequence) and

TCCCCAGGCGGAATGTTTAA (reverse primer sequences). Assays were pipetted onto

384 well plates by a robot (Pipetmax; Gilson, Middleton, Wisconsin) using a custom

programme and run on a QuantStudio 6-flex Real-Time PCR System (ThermoFisher) at

the machine manufacturers default real-time PCR cycling conditions. Reaction size was

10 µl, incorporating 1 µl of template (diluted 1/20) and PrecisionFAST qPCR Master Mix

(PrimerDesign, Chandler's Ford, UK) with low ROX and SYBR green and primers at the

machine manufacturer's recommended concentrations. Alongside the pathogen assays

we ran assays for two host genes (*Ywhaz* and *Actb*) that acted as endogenous controls.

For the pathogen assays we used as a calibrator sample a pool of DNA extracted from

154 blood samples from different *M. agrestis* at our study sites in 2015 and 2016; these

DNA extractions were carried out using the QIAamp UCP DNA Micro Kit (Qiagen,

Hilden, Germany) following manufacturer's instructions. For the *Ywhaz* and *Actb* assays

the calibrator sample was the same cDNA calibrator sample as described below for

other host gene expression assays. Relative expression values (normalised to the two

host endogenous control genes and presumed to relate to the expression of pathogen

ribosomal RNA genes and in turn to the force of infection) used in analyses below are

RQ values calculated by the QuantStudio 6-flex machine software according to the $\Delta\Delta$Ct

method, indexed to the calibrator samples. Melting curves and amplification plots were

individually inspected for each well replicate to confirm specific amplification. We

validated our diagnostic results by comparing our PCR RQ values to independent data

for a subset of voles from the cross-sectional component of our study with mapped

genus-level pathogen reads from RNASeq analysis of blood samples ($n = 44$), finding

the two data sets strongly corroborated each other (*Bartonella* spp., $r^2 = 0.81$, $p < 0.001$;

B. microti, $r^2 = 0.81$, $p < 0.001$).

Cross-sectional data

For the cross-sectional component of the study (which ran from 2015-2016 only),

animals were captured and returned to the laboratory where they were killed by a rising

concentration of CO_2, followed by exsanguination. As our UK Home Office licence did

not allow us to sample overtly pregnant females in this way, there are fewer females

than males present in this dataset. This component of the study allowed us to take a

more comprehensive set of measurements, and to culture cells and perform stimulatory

assays on them. A small tissue sample was taken from the ear of cross-sectional

animals for genotyping (see below).

 Parasite detection

After culling, the fur of cross-sectional animals was examined thoroughly under a

binocular dissecting microscope to check for ectoparasites, which were recorded as

direct counts of ticks and fleas. Guts of cross-sectional animals were transferred to 70%

ethanol, dissected and examined under a microscope for gastrointestinal parasites.

Direct counts of cestodes (the dominant endohelminths) were recorded.

SOD1 measurement

Given the well-established link between inflammation, oxidative stress and pathology

[34], we measured antioxidant enzymatic activity in blood samples taken from cross-

sectional animals. We chose to measure superoxide dismutase 1 (SOD1) because it is

one of the main antioxidative enzymes within the antioxidative enzyme system [35], and

is clearly linked to changes in immune gene expression in laboratory mice [46]. Assays

were carried out using the Cayman Superoxide Dismutase kit and following the

manufacturer's instructions except where otherwise indicated below. Blood pellets from

centrifuged cardiac bleeds were stored at -70°C and thawed on ice prior to assay. A 20

µl aliquot from each pellet was lysed in 80 µl of ultrapure water and centrifuged (10000 G

at 4°C for 15 minutes) and 40 µl of the supernatant added to a 1.6 × volume of ice-cold

chloroform/ethanol (37.5/62.5 (v/v)) (inactivating superoxide dismutase 2). This mixture

was then centrifuged (2500 G at 4°C for 10 minutes) and the supernatant removed and

immediately diluted 1:200 in kit sample buffer. A seven-point SOD activity standard was

prepared (0, 0.005, 0.010, 0.020, 0.030, 0.040, 0.050 U/ml) and the assay conducted on

a 96 well microplate with 10 µl of standard or sample, 200 µl of kit radical detector

solution and 20 µl of xanthine oxidase (diluted as per manufacturer's instructions) per

well. Plates were then covered with a protective film, incubated at room temperature for

30 mins on an orbital shaker and read at 450 nm on a VERSAmax tuneable absorbance

plate reader (Molecular Devices, San Jose, California), subtracting background and

fitting a linear relationship to the standard curve in order to estimate SOD activity in

unknown samples.

Splenocyte cultures

Spleens of cross-sectional animals were removed, disaggregated, and splenocytes

cultured under cell culture conditions equivalent to those used in [47]. Unstimulated

splenocytes, taken from 84 cross-sectional animals collected between July and October

2015, were initially used to assay expression by RNASeq (see below). We exposed

splenocytes from the remaining cross-sectional animals to stimulation with anti-CD3

antibodies (Hamster Anti-Mouse CD3e, Clone 500A2 from BD Pharmingen, San Diego,

California) and anti-CD28 antibodies (Hamster Anti-Mouse CD28, Clone 37.51 from

Tombo Biosciences, Kobe, Japan) at concentrations of 2 μg/ml and of 1 μg/ml

respectively for 24 hr. Costimulation with anti-CD3 and anti-CD28 antibodies was used

to selectively promote the proliferation of T cells [48]. We assumed that this would reflect

the potential response of T-cell populations *in vivo*. Stimulated splenocytes were used to

assay expression by Q-PCR.

RNASeq

Full details of the methods used for RNA preparation and sequencing can be found in

[3]. Briefly, samples were sequenced on an Illumina HiSeq4000 platform. High-quality

reads were mapped against a draft genome for *M. agrestis* (GenBank Accession no.

LIQJ00000000) and counted using featureCounts [49]. Genes with average log counts

per million, across all samples, of less than one were considered unexpressed and

removed from the data (n = 8,410). Following filtering, library sizes were recalculated,

data were normalized and MDS plots were generated to check for any unusual patterns

in the data.

Q-PCR

We used SYBR green based Q-PCR to measure the expression levels of a panel of 18

genes (Table S11) in splenocytes stimulated with anti-CD3 and anti-CD28 antibodies

from our cross-sectional animals. We did this, in part, to validate our RNASeq results in

an independent dataset. We used the observed expression profile as a measure of the

potential responsiveness of the immune system to an inflammatory stimulation *in vivo*.

The choice of our panel of genes was informed by (i) known immune-associated

functions in mice, combined with (ii) significant sensitivity to environmental or intrinsic

host variables in our previous studies [11,47] or in a recent DGE analysis of RNASeq

data (not reported here).

Primers (20 sets, including 2 endogenous control genes) were designed *de novo* and

supplied by PrimerDesign (16 sets) or designed *de novo* in-house (4 sets) and validated

(to confirm specific amplification and 100 ± 10% PCR efficiency under assay conditions).

All PrimerDesign primer sets were validated under our assay conditions before use. The

endogenous control genes (*Ywhaz* and *Sdha*) were selected as a stable pairing from our

previous stability analysis of candidate control genes in *M. agrestis* splenocytes [47]. We

extracted RNA from splenocytes conserved in RNAlater using the RNAqueous Micro

Total RNA Isolation Kit (ThermoFisher), following the manufacturer's instructions. RNA

extracts were DNAse treated and converted to cDNA using the High-Capacity RNA-to-

cDNA™ Kit (ThermoFisher), according to manufacturer's instructions, including reverse

transcription negative (RT-) controls for a subsample. SYBR green-based assays were

pipetted onto 384 well plates by a robot (Pipetmax, Gilson) using a custom programme

and run on a QuantStudio 6-flex Real-Time PCR System (ThermoFisher) at the machine

manufacturers default real-time PCR cycling conditions. Reaction size was 10 µl,

incorporating 1 µl of template and PrecisionFAST qPCR Master Mix with low ROX and

SYBR green (PrimerDesign) and primers at the machine manufacturer's recommended

concentrations. We used two standard plate layouts for assaying, each of which

contained a fixed set of target gene expression assays and the two endogenous control

gene assays (the same sets of animals being assayed on matched pairs of the standard

plate layouts). Unknown samples were assayed in duplicate wells and calibrator samples

in triplicate wells and no template controls for each gene were included on each plate.

Template cDNA (see above) was diluted 1/20 prior to assay. The calibrator sample

(identical on each plate) was created by pooling cDNA derived from across all

splenocyte samples. Samples from different sampling groups were dispersed across

plate pairs, avoiding confounding of plate with the sampling structure. Gene relative

expression values used in analyses are RQ values calculated by the QuantStudio 6-flex

machine software according to the $\Delta\Delta$Ct method, indexed to the calibrator sample.

Melting curves and amplification plots were individually inspected for each well replicate

to confirm specific amplification.

Longitudinal and cross-sectional data

Genotyping

We genotyped both cross-sectional and longitudinal animals for 346 single nucleotide

polymorphisms (SNPs) in 127 genes. See [3] for details of the approach used to select

these SNPs. Our list included two synonymous and tightly linked (r^2 = 0.50) SNPs in the

gene *Fcer1a* (the alpha chain of the high-affinity receptor for IgE; see Fig S1 for location)

which we had previously identified as a candidate tolerance gene in a natural population

of *M. agrestis.* In this previous work, we had identified three haplotypes at this locus

present in our population: GC, AC and AT, at frequencies of 0.12, 0.76 and 0.07

respectively. We had also identified the GC haplotype as being of particular interest,

given its significantly lower expression level of the transcription factor GATA3 (a

biomarker of tolerance to macroparasites in our population) compared to the other

haplotypes [3]. We concluded that this haplotype marked a causal (non-synonymous)

mutation. DNA was extracted from a tail sample (longitudinal component) or an ear

sample (cross-sectional component) taken from the animal using DNeasy Blood and

Tissue Kit (Qiagen). Genotyping was then performed by LGC Biosearch Technologies

(Hoddesdon, UK; http://www.biosearchtech.com) using the KASP SNP genotyping

system. This included negative controls (water) and duplicate samples for validation

purposes. The resulting SNP dataset was used for two purposes (i) genotyping

individuals within the locus of interest, and (ii) pedigree reconstruction (see below).

Pedigree reconstruction

We used a subset of our SNP dataset to reconstruct a pedigree for both cross-sectional

and longitudinal animals using the R package *Sequoia* [50]. SNPs which violated the

assumptions of Hardy-Weinberg equilibrium were removed from the dataset. For pairs of

SNPs in high linkage disequilibrium (most commonly within the same gene), the SNP

with the highest minor allele frequency (MAF) was chosen. A minimum MAF cut-off of

0.1 and call rate of > 0.7 was then applied, and any samples for which more than 50% of

SNPs were missing were removed. This resulted in a final dataset including 114 SNPs -

a sufficient number for very good performance of parentage assignment [50].

Life history information, namely sex and month of birth, was inputted into *Sequoia* where

possible. Juvenile voles weighing less than 12 g on first capture were assigned a birth

date 2 weeks prior to capture. Juvenile voles weighing between 12 and 15 g on first

capture were assigned a birth date 4 weeks prior to capture. Finally, adult voles breeding

on first capture, were assigned a birth date 6 weeks prior to capture (minimum age at

first breeding) [51,52]. Adult voles not breeding on first capture could not be assigned a

birth date, as it was not known whether they had previously bred or not. Virtually all

samples (99%) were assigned a sex, and approximately half (54%) were assigned a

birth month. As we sampled individuals from across seven different clear-cut areas of the

forest, each several kilometres apart, these were assumed to be independent, closed

populations with negligible dispersal. Site-specific pedigrees were therefore generated.

We assessed the accuracy of our reconstructed pedigrees by checking whether

predicted parent-offspring pairs met expectations given the biology of *M. agrestis*. As

expected, the majority of predicted parent-offspring pairs (87%) were born in the same

year. We also expected parents and offspring to overlap in space. Again, as expected,

the majority of predicted parent-offspring pairs (92%) were, at some point, found along

the same transects (horizontal or vertical). We also inspected log10 likelihood ratios

(LLRs) for parent pairs as recommended in the user manual for *Sequoia*. Almost all

LLRs were positive (n = 698/720 or 97% of LLRs) indicating confidence in our

assignments. Individuals with vs. without our haplotype of interest did not differ in their

probability of appearing in a pedigree (χ^2 = 0.09; d.f. = 1; p = 0.76). For each individual

that ended up in a pedigree i.e. with one or more relatives recorded (n = 652; Site COL =

3; Site BLB = 125 ; Site GRD = 204; Site CHE = 137; Site RAV = 16; Site SCP = 90;

Site HAM = 77), the number of offspring was counted to provide a measure of their

reproductive success. Few individuals were first trapped as juveniles, with the majority

trapped as adults which had already recruited into the population. Our measure of

reproductive success then, more closely resembles the number of recruited (rather than

newborn) offspring per individual. Half of individuals present in our pedigrees (n = 325)

were found to have no offspring. We expect the majority of these to be true zeros

(representing actual reproductive failure) as we generally sampled a large proportion of

the total population within clear-cuts. We also minimised the chance of false zeros by

excluding those individuals (e.g. at the periphery of a study grid) which did not end up in

a pedigree because we identified no relatives (including offspring), likely because we

had not sampled in the right place.

Statistical analyses

Not all individuals appeared in all datasets, therefore sample sizes (reported in Table 1)

vary between analyses. All analyses were performed in R statistical software version

3.5.2 [53].

Differential gene expression analysis

DGE analysis was performed on filtered and normalized count data using the R package

edgeR [54], the aim being to identify individual genes which were differentially expressed

between those individuals with and without a copy of the GC haplotype. Only those

individuals for which haplotype could be inferred with certainty could be included (n = 53

males and $n = 31$ females). Samples from different sexes were analysed together. To

test whether the top-responding genes differed between the sexes, we included a

dummy variable with four levels (males with haplotype, males without haplotype, females

with haplotype, females without haplotypes) and inspected the contrasts of interest

(males with versus without haplotypes; females with versus without haplotype). As this

was an exploratory analysis, used in conjunction with more targeted measurements of

immune gene expression (see below), model specification was kept as simple as

possible.

Having confirmed a sex-specific effect of the GC haplotype on the expression of some

inflammatory genes (see Results) in the initial DGE analysis, we ran separate DGE

analyses for males and females and tested more broadly, for enrichment of pro- and

anti-inflammatory genes in our results; more specifically, the gene ontology terms,

'positive regulation of inflammatory response' (GO:0050729; $n = 143$) and 'negative

regulation of inflammatory response' (GO:0050728; $n = 149$). The aim here was to

answer the question: are pro- or anti-inflammatory genes more highly ranked relative to

other genes, when we compare individuals with and without the GC haplotype, and does

this also differ between the sexes? This GSEA was performed using the R package

limma [55], and genes were ranked on log fold change.

Haplotype association analyses

Following the exploratory DGE analysis, the GC haplotype was tested for associations

with (i) gene expression assayed by Q-PCR to validate these results, (ii) macro- and

microparasite infection to test whether the GC haplotype predicted an individual's

susceptibility to infection, and (iii) reproductive success to test whether the GC haplotype

was associated with any reproductive costs. Given the well-established link between

inflammation and oxidative stress [56,57], we also tested for an association between the

GC haplotype and SOD1 activity.

For all analyses, this was initially attempted using the R package hapassoc [58] in order

to maximise sample size (see below). Hapassoc infers haplotypes on the basis of data

from single SNPs, and allows likelihood inference of trait associations with resulting SNP

haplotypes and other attributes. It adopts a generalized linear model framework and

estimates parameters using an expectation–maximization algorithm. Hapassoc models

assumed an additive genetic model. If the haplotype combination of an individual cannot,

with certainty, be inferred from its genotyping data (i) because it is heterozygous at two

or more markers or (ii) because it has missing data for a single marker, the approach

implemented in hapassoc is to consider all possible haplotype combinations for that

individual. Standard errors accounting for this added uncertainty are calculated using the

Louis' method [59]. We compared the GC haplotype against the other two major

haplotypes (AC and AT). Another haplotype, the GT haplotype, was identified in the

population but this was present at such low frequencies (frequency < 0.01 among

individuals for which haplotype could be inferred with certainty) that it was omitted from

all analyses. Results reported in the text for macroparasites and SOD1 activity come

from these hapassoc models.

However, there are some restrictions on model specification within hapassoc (e.g.

random terms cannot be included, limited choice of error distributions), so this was

followed up with regression models for some analyses. These regression models only

included those individuals whose haplotype combination could be inferred with certainty.

Results reported in the text for gene expression assayed by Q-PCR, microparasites and

reproductive success come from these regression models. As in the DGE analysis,

genotype was coded as the presence or absence of the GC haplotype. Regression

models were run using the R package lme4 [60] or glmmADMB [61,62].

Table 1 provides a summary of (hapassoc or regression) model specifications. As we

found evidence for sex-specific effect in the initial DGE analysis, with the exception of

the model for reproductive success (see below), all models included a genotype by sex

interaction which was retained if it improved model fit. All models also accounted for

other biological and technical covariates (the choice of which was informed by our

previous work, the literature and/or our experimental design). Full regression models

(including all covariates and interactions of interest) were reduced using backward

stepwise deletion of non-significant terms to minimize Akaike's information criterion

(AIC), following the drop1 function in the lme4 package. This was not possible for

hapassoc models. Summary tables including all estimates, standard errors and z-

statistics are included in the Supplementary information. Associated significance values

are also included in these tables, except for LMMs and GLMMs, for which significance

values were based on likelihood ratio tests (LRTs) and are reported in the main text.

Throughout, residuals from regression models were checked for approximate normality

and homoscedasticity, and all covariates were tested for independence using variance

inflation factors (all VIFs < 3).

685 **Table 1.** Model specifications including, for each model: covariates included in the full

686 model, datasets used and sample sizes.

	DGE analysis	Haplotype association analyses			
			Response variable		
		Immune gene expression assayed by Q-PCR	Parasite infection	Reproductive success	SOD1 activity
Covariates					
Snout-vent length		Y			Y
Eye lens weight		Y			Y
Reproductive status		Y	Y		Y
Body condition		Y	Y		Y
Birth month				Y	
Culled or not				Y	
Site		Y	Y	Y	Y
Year		Y	Y	Y	Y
Season		Y	Y		Y
Assay plate		Y (*Il17a* LMM)			
Dataset[a]	C	C	C (Macroparasites)	C + L	C
			L (Microparasites)		
Sample size	♀31 ♂53	*Il17a* LMM: ♀73 ♂220	Macroparasites: ♀82 ♂235	♀419	♀81
			B. microti: ♀1075 ♂1247	♂232	♂227
			Bartonella spp.: ♀1283 ♂1104		

688 [a] C, cross-sectional; L, longitudinal.
689

Association between GC haplotype and immune gene expression assayed by Q-

PCR

As we ran a total of 18 hapassoc models (one model per gene), the Benjamini and

Hochberg method of correction was applied to all *p*-values [63]. Resulting *q*-values

(FDR-corrected *p*-values) are reported, alongside original *p*-values (Table S3). A

hapassoc model including a genotype by sex interaction was tested against a null model

without an interaction term. A Gamma error distribution and log link were used. Other

covariates considered potential drivers of immune gene expression and included in both

models, were informed by our previous work [47]. These included snout-vent length

(SVL), eye lens weight (categorised into seven intervals; SVL and eye lens weight

represent the combined influence of age and historical growth trajectory), reproductive

status (males were considered to be reproductively active if they had descended testes;

females if they were pregnant or had perforate vaginas) and body condition (estimated

by regressing body weight against life history stage, SVL and its quadratic term). Site,

year and season (four levels, designated as spring [March & April], early summer [May

and June], late summer [July and August) and autumn [September and October]) were

included to account for any spatial and/or temporal autocorrelation. All covariates were

included as fixed effects.

In order to confirm these results, a linear mixed effects model (LMM) was run for a single

immune gene for which expression appeared to be associated with genotype ($q =$

0.037). The LMM included random terms for site and season, as well as assay plate

number (Table 1). The latter was included to account for nonindependence due to

immunological assaying structure. All other covariates were the same as the hapassoc

model. A Yeo-Johnson transformation (with $\lambda = -2$) was used to achieve more normal

and homoscedastic residuals [64].

Association between GC haplotype and parasite infection

The three macroparasite measurements taken from cross-sectional animals (counts of

ticks, fleas and cestodes) were log-transformed ($\log_{10} [x +1]$) and summarised as a

single principal component (explaining 39% of total variation; Table S12) to avoid

difficulties in interpretation due to multiple testing. See [3] and [11] for full details of this

approach. This combined measure of macroparasite burden was modelled using a

hapassoc model with a Gaussian error distribution.

Microparasite infection status was assessed multiple times for the majority of individuals

in the longitudinal component of the study (mean = 2.8; range = 1-11). Due to these

repeated measures, microparasite infection could not be modelled using hapassoc.

Instead, to test for an association between the GC haplotype and the probability of an

individual being infected with a microparasite, we ran a GLMM with a a binary response

(infected or not), log link and a random term for individual. Other covariates, considered

potential drivers of both macro and microparasite infection were, again, informed by our

previous work [3,11,65]. These included body condition, reproductive status, year,

season and site (Table 1). Season and site were included as random terms in the GLMM

for microparasites, as was individual identity, to account for nonindependence due to

repeat sampling.

Association between GC haplotype and reproductive success

Our measure of reproductive success was zero-inflated (50% zeros). This is consistent

with a previous study of the closely-related common vole (*Microtus arvalis*) [66]. A

Poisson error distribution was therefore deemed inappropriate and it could not be

modelled using hapassoc. Instead, to test for an association between the GC haplotype

and reproductive success, we ran a GLM with a quasi-Poisson error distribution. This

distribution has been previously used to model predictors of reproductive success in

other organisms, and accounts for the overdispersion caused by excess zeros.

We only recorded a reproductive failure (i.e. zero reproductive success) for those

individuals with some other familial relationship in our pedigree, purposefully omitting

those individuals (e.g. at the periphery of a study grid) for which we may have recorded

no relatives (including offspring) simply because we had not sampled in the right place.

Therefore, we expect all (or most) of our zeros to represent actual reproductive failure.

For this reason, zero-inflated models were deemed inappropriate.

As detailed in the Results, we tried to run a single quasi-Poisson GLM with both sexes

but resulting residual variances differed significantly between the sexes (F test to

compare variances of two samples; $p = 0.02$), making it impossible to formally test for a

genotype by sex interaction. Instead, we ran a separate model for each sex. We

included birth month as a covariate in this model, given that autumn-born voles (of the

closely-related common vole) have been shown to have a lower chance to reproduce

than spring-born voles [66]. Other covariates included in this model were, whether or not

an individual was culled for the cross-sectional component of this study (again, reducing

the opportunity to reproduce), site and the year in which an individual was most

frequently captured (Table 1). All covariates were included as fixed effects. Only a single

female was trapped in one of the sites (COL) and consequently caused convergence

problems in the female model. This female was therefore omitted.

Association between GC haplotype and SOD1 activity

SOD1 activity was modelled using a hapassoc model with a Gaussian error distribution.

Other covariates, considered potential drivers of SOD1 activity, were informed by the

literature. Previous studies on wild rodents have shown that antioxidant levels increase

with both reproductive effort [67] and with age [68]. Studies on birds have also shown

that improvements in body condition are often accompanied by increases in antioxidant

activity, for example in response to supplemental feeding [69]. We therefore included

SVL, eye lens weight, reproductive status and body condition as covariates in our model.

As in other models, we included site, year and season to account for spatial and/or

temporal autocorrelation in our data (Table 1). All covariates were included as fixed

effects.

Acknowledgements

We would like to thank all those involved in obtaining and processing samples from the

field: Rebecca Turner, Lukasz Lukomski, Stephen Price, Sarah Gore, Ed Parker, Maria

Capstick, Noelia Dominguez Alvarez, Susan Withenshaw, William Foster, Ann Lowe and

Benoit Poulin. We would also like to thank the Forestry Commission for access to the

study sites and the Centre for Genomic Research at the University of Liverpool for

sequencing samples. This research was funded by the Natural Environment Research

Council (NERC) award NE/L013452/1 to S.P., M.B., J.A.J. and J.E.B.

Competing interests

The authors declare no competing interests.

References

1.   Møller AP, Saino N. Immune response and survival. Oikos. 2004;104: 299–304.

doi:10.1111/j.0030-1299.2004.12844.x

2.   Paterson S, Wilson S, Pemberton J. Major histocompatibility complex variation

associated with juvenile survival and parasite resistance in a large unmanaged

ungulate population (*Ovis aries L.*). Proc Natl Acad Sci USA. 1998;95: 3714–

3719. doi:10.1073/pnas.95.7.3714

3.   Wanelik KM, Begon M, Birtles RJ, Bradley JE, Friberg IM, Jackson JA, et al. A

candidate tolerance gene identified in a natural population of field voles (*Microtus*

agrestis). Mol Ecol. 2018;27: 1044–1052. doi:10.1111/mec.14476

800 4. Turner AK, Begon M, Jackson JA, Bradley JE, Paterson S. Genetic Diversity in

801   Cytokines Associated with Immune Variation and Resistance to Multiple

802   Pathogens in a Natural Rodent Population. PLoS Genet. 2011;7: e1002343.

803   doi:10.1371/journal.pgen.1002343

804 5. Mackay TFC, Anholt RRH. Of Flies and Man: *Drosophila* as a Model for Human

805   Complex Traits. Annu Rev Genomics Hum Genet. 2006;7: 339–367.

806   doi:10.1146/annurev.genom.7.080505.115758

807 6. Hayes KS, Hager R, Grencis RK. Sex-dependent genetic effects on immune

808   responses to a parasitic nematode. BMC Genomics. 2014;15. doi:10.1186/1471-

809   2164-15-193

810 7. Hill AVS, Allsopp CEM, Kwiatkowski D, Anstey NM, Twumasi P, Rowe PA, et al.

811   Common West African HLA antigens are associated with protection from severe

812   malaria. Nature. 1991;352: 595–600. doi:10.1038/352595a0

813 8. Carrington M, Nelson GW, Martin MP, Kissner T, Vlahov D, Goedert JJ, et al. HLA

814   and HIV-1: Heterozygote Disadvantage and B*35-Cw*04 Disadvantage. Science.

815   1999;283: 1748–1752.

816 9. Thursz MR, Thomas HC, Greenwood BM, Hill A V. Heterozygote advantage for

817   HLA class-II type in hepatitis B virus infection. Nat Genet. 1997;17: 11–12.

818   doi:10.1038/ng0997-11

819 10. Ober C, Loisel DA, Gilad Y. Sex-specific genetic architecture of human disease.

820   Nat Rev Genet. 2008;9: 911–922. doi:10.1038/nrg2415

821 11. Jackson JA, Hall AJ, Friberg IM, Ralli C, Lowe A, Zawadzka M, et al. An

822   immunological marker of tolerance to infection in wild rodents. PLoS Biol.

823   2014;12: e1001901. doi:10.1371/journal.pbio.1001901

824 12. Patsopoulos NA, Tatsioni A, Ioannidis JPA. Claims of sex differences: An

empirical assessment in genetic associations. J Am Med Assoc. 2007;298: 880–

893. doi:10.1001/jama.298.8.880

13. Graham AL, Hayward AD, Watt KA, Pilkington JG, Pemberton JM, Nussey DH.

Fitness correlates of heritable variation in antibody responsiveness in a wild

mammal. Science. 2010;330: 662–665. doi:10.1126/science.1194878

14. Turner AK, Beldomenico PM, Bown K, Burthe SJ, Jackson JA, Lambin X, et al.

Host-parasite biology in the real world: the field voles of Kielder. Parasitology.

2014;141: 997–1017. doi:10.1017/S0031182014000171

15. Turner AK, Paterson S. Wild rodents as a model to discover genes and pathways

underlying natural variation in infectious disease susceptibility. Parasite Immunol.

2013;35: 386–395. doi:10.1111/pim.12036

16. Mayne BT, Bianco-Miotto T, Buckberry S, Breen J, Clifton V, Shoubridge C, et al.

Large scale gene expression meta-analysis reveals tissue-specific, sex-biased

gene expression in humans. Front Genet. 2016;7: 1–14.

doi:10.3389/fgene.2016.00183

17. Rinn JL, Snyder M. Sexual dimorphism in mammalian gene expression. Trends

Genet. 2005;21: 298–305. doi:10.1016/j.tig.2005.03.005

18. Yang X, Schadt EE, Wang S, Wang H, Arnold AP, Ingram-Drake L, et al. Tissue-

specific expression and regulation of sexually dimorphic genes in mice. Genome

Res. 2006;16: 995–1004. doi:10.1101/gr.5217506

19. Ellegren H, Parsch J. The evolution of sex-biased genes and sex-biased gene

expression. Nat Rev Genet. 2007;8: 689–698. doi:10.1038/nrg2167

20. Sutherland B, Prokkola J, Audet C, Bernatchez L. Sex-specific and conserved

gene co-expression networks underlie phenotypes 2 within the paleopolyploid

Salvelinus genus (Family Salmonidae). bioRxiv. 2018; 305680.

850   doi:10.1101/305680

851 21. Van Nas A, Guhathakurta D, Wang SS, Yehya N, Horvath S, Zhang B, et al.

852   Elucidating the role of gonadal hormones in sexually dimorphic gene

853   coexpression networks. Endocrinology. 2009;150: 1235–1249.

854   doi:10.1210/en.2008-0563

855 22. Gounni AS, Lamkhioued B, Ochiai K, Tanaka Y, Delaporte E, Capron A, et al.

856   High-affinity IgE receptor on eosinophils is involved in defence against parasites.

857   Nature. 1994;367: 183–186. doi:10.1038/368473a0

858 23. Tomassini M, Tsicopoulos A, Tai PC, Gruart V, Tonnel AB, Prin L, et al. Release

859   of granule proteins by eosinophils from allergic and nonallergic patients with

860   eosinophilia on immunoglobulin-dependent activation. J Allergy Clin Immunol.

861   1991;88: 365–375. doi:10.1016/0091-6749(91)90099-A

862 24. Daeron M, Nimmerjahn F, editors. Current Topics in Microbiology and

863   Immunology: Fc Receptors. Immunology. Springer; 2014.

864 25. Weidinger S, Gieger C, Rodriguez E, Baurecht H, Mempel M, Klopp N, et al.

865   Genome-wide scan on total serum IgE levels identifies FCER1A as novel

866   susceptibility locus. PLoS Genet. 2008;4: e1000166.

867   doi:10.1371/journal.pgen.1000166

868 26. Granada M, Wilk JB, Tuzova M, Strachan DP, Weidinger S, Albrecht E, et al. A

869   genome-wide association study of plasma total IgE concentrations in the

870   Framingham Heart Study. J Allergy Clin Immunol. 2012;129: 840–845.

871   doi:10.1016/j.jaci.2011.09.029

872 27. Hasegawa M, Nishiyama C, Nishiyama M, Akizawa Y, Mitsuishi K, Ito T, et al. A

873   Novel –66T/C Polymorphism in FcεRI α-Chain Promoter Affecting the

874   Transcription Activity: Possible Relationship to Allergic Diseases. J Immunol.

2003;171: 1927–1933. doi:10.4049/jimmunol.171.4.1927

28. Potaczek DP, Sanak M, Mastalerz L, Setkowicz M, Kaczor M, Nizankowska E, et

al. The α-chain of high-affinity receptor for IgE (FcεRIα) gene polymorphisms and

serum IgE levels. Allergy. 2006;61: 1230–1233. doi:10.1111/j.1398-

9995.2006.01195.x

29. Zhou J, Zhou Y, Lin L, Wang J, Peng X, Li J, et al. Association of polymorphisms

in the promoter region of FCER1A gene with atopic dermatitis, chronic uticaria,

asthma, and serum immunoglobulin E levels in a Han Chinese population. Hum

Immunol. 2012;73: 301–305. doi:10.1016/j.humimm.2011.12.001

30. Niwa Y, Potaczek DP, Kanada S, Takagi A, Shimokawa N, Ito T, et al. FcεRIα

gene (FCER1A) promoter polymorphisms and total serum IgE levels in Japanese

atopic dermatitis patients. Int J Immunogenet. 2010;37: 139–141.

doi:10.1111/j.1744-313X.2010.00901.x

31. Weiss LA, Pan L, Abney M, Ober C. The sex-specific genetic architecture of

quantitative traits in humans. Nat Genet. 2006;38: 218–222. doi:10.1038/ng1726

32. Chen W, Mempel M, Schober W, Behrendt H, Ring J. Gender difference, sex

hormones, and immediate type hypersensitivity reactions. Allergy. 2008;63: 1418–

1427. doi:10.1111/j.1398-9995.2008.01880.x

33. Yang J, Lu MM, Lu YW, Feng CC, Leng RX, Pan HF, et al. Sex-specific

differences in the relationship between the single-nucleotide polymorphism

rs2298804 of FCER1A and the susceptibility to systemic lupus erythematosus in a

Chinese Han population. Clin Exp Dermatol. 2013;38: 410–416.

doi:10.1111/ced.12035

34. Biswas SK. Does the Interdependence between Oxidative Stress and

Inflammation Explain the Antioxidant Paradox? Oxid Med Cell Longev.

2016;2016: 17–19. doi:10.1155/2016/5698931

35.   Sun Y, Oberley LW, Elwell JH, Sierra-Rivera E. Antioxidant enzyme activities in

normal and transformed mouse liver cells. Int J Cancer. 1989;44: 1028–1033.

doi:10.1002/ijc.2910440615

36.   Wegmann TG, Lin H, Guilbert L, Mosmann TR. Bidrectional cytokine interactions

in the maternal-fetal relationship: is successful pregnancy a TH2 phenomenon?

Immunol Today. 1993;14: 353–356. doi: 10.1016/0167-5699(93)90235-D

37.   Bianchi VE. The Anti-Inflammatory Effects of Testosterone. J Endocr Soc. 2019;3:

91–107. doi:10.1210/js.2018-00186

38.   Davis S, Abbasi B, Shah S, Telfer S, Begon M, Davis S. Spatial analyses of

wildlife contact networks. J R Soc Interface. 2014;12. doi:10.1098/rsif.2014.1004

39.   Martínez-Sanchis S, Arnedo MT, Salvador A, Moya-Albiol L, González-Bono E.

Effects of Chronic Administration with High Doses of Testosterone Propionate on

Behavioral and Physiological Parameters in Mice with Differing Basal

Aggressiveness. Aggress Behav. 2003;29: 173–189. doi:10.1002/ab.10050

40.   Djokic V, Akoolo L, Parveen N. *Babesia microti* infection changes host spleen

architecture and is cleared by a Th1 immune response. Front Microbiol. 2018;9:

85. doi:10.3389/fmicb.2018.00085

41.   Telfer S, Lambin X, Birtles R, Beldomenico P, Burthe S, Paterson S, et al.

Species interactions in a parasite community drive infection risk in a wildlife

population. Science. 2010;330: 243–246. doi:10.1126/science.1190333

42.   Sheldon BC, Verhulst S. Ecological immunology: costly parasite defences and

trade-offs in evolutionary ecology. Trends Ecol Evol. 1996;11: 317–321.

43.   DelBarco-Trillo J, Ferkin MH. Male mammals respond to a risk of sperm

competition conveyed by odours of conspecific males. Nature. 2004;431: 446–

449. doi:10.1038/nature02845

44.   Losdat S, Richner H, Blount JD, Helfenstein F. Immune activation reduces sperm

quality in the great tit. PLoS One. 2011;6: e22221.

doi:10.1371/journal.pone.0022221

45.   Garratt M, Pichaud N, Glaros EN, Kee AJ, Brooks RC. Superoxide dismutase

deficiency impairs olfactory sexual signaling and alters bioenergetic function in

mice. Proc Natl Acad Sci USA. 2014;111: 8119–8124.

doi:10.1073/pnas.1322282111

46.   Marikovsky M, Ziv V, Nevo N, Harris-Cerruti C, Mahler O. Cu/Zn Superoxide

Dismutase Plays Important Role in Immune Response. J Immunol. 2003;170:

2993–3001. doi:10.4049/jimmunol.170.6.2993

47.   Jackson JA, Begon M, Birtles R, Paterson S, Friberg IM, Hall A, et al. The

analysis of immunological profiles in wild animals: A case study on

immunodynamics in the field vole, *Microtus agrestis*. Mol Ecol. 2011;20: 893–909.

doi:10.1111/j.1365-294X.2010.04907.x

48.   Frauwirth KA, Thompson CB. Activation and inhibition of lymphocytes by

costimulation. J Clim Invest. 2002;109: 295–299. doi:10.1172/JCI14941

49.   Liao Y, Smyth GK, Shi W. featureCounts: An efficient general purpose program

for assigning sequence reads to genomic features. Bioinformatics. 2014;30: 923–

930. doi:10.1093/bioinformatics/btt656

50.   Huisman J. Pedigree reconstruction from SNP data: parentage assignment,

sibship clustering and beyond. Mol Ecol Resour. 2017;17: 1009–1024.

doi:10.1111/1755-0998.12665

51.   Begon M, Telfer S, Burthe S, Lambin X, Smith MJ, Paterson S. Effects of

abundance on infection in natural populations: field voles and cowpox virus.

Epidemics. 2009;1: 35–46. doi:10.1016/j.epidem.2008.10.001

52. Burthe SJ, Lambin X, Telfer S, Douglas A, Beldomenico P, Smith A, et al.

Individual growth rates in natural field vole, *Microtus agrestis*, populations

exhibiting cyclic population dynamics. Oecologia. 2010;162: 653–661.

doi:10.1007/s00442-009-1495-6

53. R Core Team. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-

project.org/. 2018.

54. Robinson MD, McCarthy DJ, Smyth GK. edgeR: A Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics.

2010;26: 139–140. doi:10.1093/bioinformatics/btp616

55. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. Limma powers

differential expression analyses for RNA-sequencing and microarray studies.

Nucleic Acids Res. 2015;43: e47. doi:10.1093/nar/gkv007

56. Reuter S, Gupta S, Chaturvedi M, Aggarwal B. Oxidative stress, inflammation,

and cancer: How are they linked? Free Radic Biol Med. 2011;49: 1603–1616.

doi:10.1016/j.freeradbiomed.2010.09.006.Oxidative

57. Collins AR. Oxidative DNA damage, antioxidants, and cancer. BioEssays.

1999;21: 238–246. doi:10.1002/(SICI)1521-1878(199903)21:3<238::AID-

BIES8>3.0.CO;2-3

58. Burkett K, Graham J, McNeney B. hapassoc: Software for likelihood inference of

trait associations with SNP haplotypes and other attributes. J Stat Softw. 2006;16:

1–19.

59. Louis T. Finding the observed information matrix when using the EM Algorithm. J

R Stat Soc B. 1982;44: 226–233.

60.   Bates D, Machler M, Bolker B, Walker S. Fitting linear mixed-effects models using

976   lme4. J Stat Softw. 2015;67: 1–48.

977   61.   Skaug H, Fournier D, Bolker B, Magnusson A, Nielsen A. Generalized Linear

978   Mixed Models using "AD Model Builder". R package version 0.8.3.3. 2016.

979   62.   Fournier DA, Skaug HJ, Ancheta J, Ianelli J, Magnusson A, Maunder MN, et al.

980   AD Model Builder: Using automatic differentiation for statistical inference of highly

981   parameterized complex nonlinear models. Optim Methods Softw. 2012;27: 233–

982   249. doi:10.1080/10556788.2011.597854

983   63.   Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and

984   powerful approach to multiple testing. J R Stat Soc Ser B. 1995;57: 289–300.

985   64.   Yeo I-K, Johnson RA. A new family of power transformations to improve normality

986   or symmetry. Biometrika. 2000;87: 954–959.

987   65.   Taylor CH, Wanelik KM, Friberg IM, Lowe A, Hall AJ, Ralli C, et al. Physiological,

988   but not fitness, effects of two interacting haemoparasitic infections in a wild

989   rodent. Int J Parasitol. 2018;48: 463–471. doi:10.1016/j.ijpara.2017.11.006

990   66.   Wang M, Ebeling M, Hahne J. Relevance of body weight effects for the population

991   development of common voles and its significance in regulatory risk assessment

992   of pesticides in the European Union. Environ Sci Eur. 2019;31–54.

993   doi:10.1186/s12302-019-0240-y

994   67.   Garratt M, Vasilaki A, Stockley P, McArdle F, Jackson M, Hurst JL. Is oxidative

995   stress a physiological cost of reproduction? An experimental test in house mice.

996   Proc R Soc B Biol Sci. 2011;278: 1098–1106. doi:10.1098/rspb.2010.1818

997   68.   Hindle AG, Lawler JM, Campbell KL, Horning M. Muscle aging and oxidative

998   stress in wild-caught shrews. Comp Biochem Physiol B Biochem Mol Biol.

999   2010;155: 427–434. doi:10.1016/j.cbpb.2010.01.007

1000   69.   Wilcoxen TE, Horn DJ, Hogan BM, Hubble CN, Huber SJ, Flamm J, et al. Effects

1001   of bird-feeding activities on the health of wild birds. Conserv Physiol. 2015;3: 1–

1002   13. doi:10.1093/conphys/cov058

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024