**Effects of an IgE receptor polymorphism acting on immunity, susceptibility to infection and reproduction in a wild rodent**

Klara M Wanelik[1*,] Mike Begon[1], Janette E Bradley[2], Ida M Friberg[3], Joseph A Jackson[3], Christopher H Taylor[2], Steve Paterson[1]

[1] Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, UK

[2] School of Life Sciences, University of Nottingham, Nottingham, UK

[3] School of Environment and Life Sciences, University of Salford, Salford, UK

* Corresponding author

Current address: Department of Biology, University of Oxford, Oxford, UK

**Email:** klara.wanelik@biology.ox.ac.uk

**Abstract**

The genotype of an individual is an important predictor of their immune function, and subsequently, their ability to control or avoid infection and ultimately contribute offspring to the next generation. However, the same genotype, subjected to different intrinsic and/or extrinsic environments can also result in different phenotypic outcomes, which can be missed in controlled laboratory studies. Natural wildlife populations, which capture both genotypic and environmental variability, provide an opportunity to more fully understand the phenotypic expression of genetic variation. We identified a synonymous polymorphism in the high-affinity Immunoglobulin E (IgE) receptor (GC and non-GC haplotypes) that has sex-dependent effects on immune gene expression, susceptibility to infection and reproductive success of individuals in a natural population of field voles (*Microtus agrestis*). We found that the effect of the GC haplotype on the expression of immune genes differed between sexes. While males with the GC haplotype had upregulated more pro-inflammatory genes, including the cytokine *Il33,* females had upregulated more anti-inflammatory genes, including the cytokine inhibitor *Socs3.* Furthermore we found an effect of the GC haplotype on the probability of infection with a common microparasite*, Babesia microti,* in females - with females carrying the GC haplotype being more likely to be infected. Finally, we found an effect of the GC haplotype on reproductive success in males - with males carrying the GC haplotype having a lower reproductive success. This is a rare example of a polymorphism whose consequences we are able to follow across immunity, infection and reproduction for both males and females in a natural population.

## Introduction

In order for an individual to control or avoid infection and ultimately contribute offspring to the next generation, they must have a well-functioning immune system [1]. An individual's immune function is in part determined by their genotype (e.g. [2–4]). Individuals within a population differ from each other in the genotypes they carry, with the potential for these different genotypes to affect the activity of different immune components, and thereby influence susceptibility to infection and reproduction and survivorship. Indeed, variability in immune responses to infection is well-characterised. The phenotypic expression of a genotype is dependent on the environment, including the sex of an individual (intrinsic environment) and exposure to infection (extrinsic environment). Since capturing genotypic and environmental variability in controlled laboratory studies is problematic, natural wildlife populations have been proposed as models in which to understand the phenotypic expression of genetic variation and its consequences for susceptibility to infection and reproduction.

Studies of natural populations have explored the effects of genotype on immune phenotype, and have observed consequences for susceptibility to infection. Most notably, variability in the genes of the Major Histocompatibility Complex (MHC) has been associated with resistance to intestinal nematodes in domestic sheep [2] and with resistance to malaria, hepatitis, and AIDS in humans [5–7]. The role of variability elsewhere in the genome, for shaping immune phenotype, has also been studied [3,4]. However, it remains challenging to follow the consequences of a genotypic effect for immunity, infection and reproduction and to account for any sex-dependent expression of a genotype. This is because of the difficulty in obtaining phenotypic data across

immune, infection and reproductive traits, especially for large enough sample sizes to

test for data-hungry genotype by sex interactions [8]. In many cases, sex, and other

environmental factors are considered as a confounding variable to be controlled for in

order not to hide any subtle genetic associations [2]. Other studies focus on a single sex

for the sake of simplicity [9]. More recently, however, there has been a growing body of

large-scale field studies of natural populations able to apply genetic and immunological

methods to follow large numbers of individuals, exposed to a challenging environment

and with varying genetic backgrounds, throughout their lives. This allows us to make a

more complete assessment of the impacts of genotype throughout the life of an

individual, whether male or female. For example, Graham et al. 2010 [10] found

evidence for heritable variation in immunity associated with sex-dependent effects on

Soay sheep reproduction. However, we know of no documented example of a

polymorphism affecting immunity, susceptibility to infection and reproduction in a natural

population investigated in males and females. Here, we use a wild rodent, the field vole

(*Microtus agrestis*), as a model in which to do this. Wild rodents, in particular, offer an

opportunity to quickly follow large numbers of individuals throughout their lives, given

their short lifespans. They also offer the opportunity to draw on the immunological and

genetics resources developed for laboratory rodents, while providing a much more

realistic ecological model of human populations [11,12].

Immunoglobulin E (IgE) mediated responses are associated with defence against

helminths [13] and with allergy [14]. They are controlled by the high-affinity IgE receptor,

FCER1, which is found on the surface of various immune effector cells e.g. mast cells,

basophils and eosinophils [15]. In humans, naturally occurring polymorphisms in FCER1

are known to affect an individual's serum IgE levels, with consequences for their

susceptibility to infection [16,17] and their risk of developing inflammatory disease [18–

21]. Furthermore, sex differences in serum IgE levels [22] and the incidence of IgE-

mediated inflammatory disease [23] have been documented in humans, suggesting that

any polymorphism in this pathway is likely to experience different contexts in males and

females. Indeed, one study found evidence for a polymorphism in the *Fcer1a* gene (the

alpha chain of FCER1) whose association with susceptibility to systematic lupus

erythematosus (a chronic inflammatory disease) differed between males and females

[24].

In a previous study of a natural population of *M. agrestis*, we found that males carrying

the GC haplotype of the *Fcer1a* gene expressed the transcription factor GATA3 at a

lower level than males carrying non-GC haplotypes [3]. GATA3 is a biomarker of

tolerance to macroparasites in mature males in our population (macroparasite infection

gives rise to increased expression of GATA3, which gives rise to improved body

condition and survival [9]). Here, we explore the effects of this GC haplotype further in

both males and females to analyse the effect of genotype acting across immune

expression, infection susceptibility and reproductive success.

**Results**

We sampled a natural population of *M. agrestis* in Kielder Forest, Northumberland,

United Kingdom, over three years (2015-2017) and across seven different sites. Our

study involved a cross-sectional component ($n = 317$ destructively sampled voles) and a

longitudinal component ($n = 850$ marked individuals monitored through time, with $n =$

2,387 sampling points). We tested the consequences of the GC haplotype of the *Fcer1a*

gene using both cross-sectional and longitudinal components of our study. As well as

the GC haplotype (present at a frequency of 0.08), three other haplotypes were present

in our study population: AC haplotype, AT haplotype and GT haplotype, present at

frequencies of 0.81, 0.10 and 0.01 respectively. The two SNPs composing the haplotype

were found to be tightly linked ($r^2 = 0.50$; D' = 0.70).

**The GC haplotype has effects on inflammation that differ between sexes**

In humans, naturally occurring polymorphisms in the *Fcer1a* gene have previously been

linked to inflammatory disease [18–21]. Therefore, we used the cross-sectional

component of our study to test the effects of the GC haplotype on inflammation in males

and females. Differential gene expression (DGE) analysis performed on unstimulated

splenocytes taken from 53 males and 31 females assayed by RNASeq, showed that the

identity of top-responding genes differed between the sexes. In males, the top-

responding immune gene was the cytokine, *Il33* (log fold change (logFC) = 2.76, *p* =

0.00000015, *q* = 0.00048; Appendix 1—table 2) while in females it was the suppressor of

cytokine signalling *Socs3* (logFC = 1.07, *p* = 0.000061, *q* = 0.05; Appendix 1—table 3).

Looking at the ranking of each top-responding gene in the opposite sex strengthens the

case for differing effects in males and females, with *Il33* ranked markedly lower in

females (rank = 8224/12904, logFC = 0.29, *p* = 0.70, *q* = 1.00) and *Socs3* markedly

lower in males (rank = 10886/12904, logFC = -0.05, *p* = 0.84, *q* = 1.00). *Il33* is commonly

associated with the anti-helminthic response [25] and *Socs3* with regulation of the

immune response more broadly [26]. Given the link between *Il33* and the antihelminthic

response (and more generally, IgE-mediated responses and the antihelminthic

response), we repeated the DGE analysis while controlling for cestode burden, but this

had little effect on our results (same top-responding immune genes; see Appendix 1—

table 4 & 5), suggesting that these effects were not driven by differences in cestode

infection.

Both *Il33* and *Socs3* also share an association with the inflammatory response [26,27].

While *Il33* positively regulates this response (appearing in the gene set GO:0050729),

*Socs3* negatively regulates it (GO:0050728). To test whether these effects on the

inflammatory response were limited to these genes or were more widespread, we

performed a gene set enrichment analysis (GSEA) which looked at the rankings of all

genes present in each of these gene sets (GO:0050729 and GO:0050728). This analysis

showed that both males and females upregulated their expression of both gene sets.

However, males showed a stronger signal for genes which positively regulate the

inflammatory response (GO:0050729: $p = 0.007$; GO:0050728: $p = 0.04$; Figure 1A,

upper panel) while females showed a stronger signal for genes which negatively

regulate the inflammatory response (GO:0050728: $p = 0.001$; GO:0050728: $p = 0.04$;

Figure 1A, lower panel).

To further explore these effects on the inflammatory response, we used an independent

dataset for males and females whose spleens were stimulated with two immune agonists

and assayed by Q-PCR (for a panel of 18 immune genes with limited redundancy; see

Methods for how these genes were selected). From this *ex vivo* assay, one can gain

insight into the types of immune response that could be made to a pathogen *in vivo* (see

Methods for details). Although our panel of genes did not include *Il33* or *Socs3*, it did

include other genes associated with the inflammatory response including *Il17a, Ifng*, *Il1b*,

*Il6* and *Tnfa*. We found that the effect of the GC haplotype on the stimulated expression

levels of *Il17a*, a pro-inflammatory cytokine involved in the antibacterial response,

displayed a significant interaction with sex (genotype by sex interaction in follow-up

LMM, LRT $p$ = 0.001). While females with the GC haplotype expressed lower levels of

*Il17a* (estimate = -0.12, 95% CI = -0.19 – -0.04), males with the GC haplotype did not

differ in their expressed levels of *Il17a* (estimate = 0.03, 95% CI includes zero = -0.02 –

0.08; Appendix 1—table 6 & 7).

**The GC haplotype affects oxidative stress**

Inflammation and oxidative stress are closely related and often lead to pathology [28]. To

test whether the effects of the GC haplotype on inflammation may be having knock-on

effects on oxidative stress, we again used the cross-sectional component of our study.

We tested the effect of the GC haplotype on antioxidant superoxide dismutase 1 (SOD1)

enzymatic activity in males and females. SOD1 is one of the main antioxidative enzymes

within the antioxidative enzyme system [29]. We found that (irrespective of sex)

individuals with the AT haplotype had lower SOD1 activity when compared with other

haplotypes, including the GC haplotype ($p$ = 0.04; Appendix 1—table 8).

**The GC haplotype affects susceptibility to *Babesia* infection in females**

We then used the longitudinal component of our study to test the effects of the GC

haplotype on the probability of infection with common parasites known to infect our study

population: two micro-parasites *Babesia microti* and *Bartonella* spp. and a range of

macroparasites (ticks, fleas and cestodes). We found an effect of the GC haplotype on

the probability of infection with *B. microti* which displayed a marginally significant

interaction with sex (genotype by sex interaction, LRT $p$ = 0.048; Appendix 1—table 9).

While females with the GC haplotype were more likely to be infected with *B. microti* (log

odds ratio = 1.25, 95% CI = 0.44 – 2.07; Figure 1B, lower panel), males with the GC

haplotype did not differ in their susceptibility to infection (log odds ratio = 0.14, 95% CI

includes zero = -0.76 – 1.03; Figure 1B, upper panel). However, we found no effect of

the haplotype (interactive or not) on the probability of infection with the other parasites in

our population (Appendix 1—table 11 & 17).

**The GC haplotype reduces male reproductive success**

We genotyped both cross-sectional and longitudinal voles for 114 SNPs and used this

dataset to construct a pedigree. We estimated each individuals' reproductive success by

counting the number offspring it produced according to this pedigree. We then tested the

effects of the GC haplotype on this measure of reproductive success. We tried to run a

single model with both sexes (as above) but resulting residual variances differed

significantly between the sexes, which reflects the fact that male and female

reproduction are inherently different traits. This made it impossible to formally test for a

genotype by sex interaction and so instead we ran a separate model for each sex. We

found that males with the GC haplotype had an average of 2.2 fewer offspring than

males without the haplotype ($p$ = 0.04; Appendix 1—table 12; Figure 1C, upper panel).

Despite a larger sample size and lower variability in reproductive success, females with

the GC haplotype did not differ significantly in their number of offspring (genotype term

did not appear in best model; Appendix 1—table 13; Figure 1C, lower panel). This is

suggestive of a significant genotype by sex interaction. We ran the same models

including microparasite variables, but this had little effect on our results (see Appendix

1—table 14 & 15), suggesting that these effects were not driven by differences in

microparasite infection.

## Discussion

In this study, we describe a polymorphism in the high-affinity receptor for IgE (*Fcer1a*) with effects that act across immune gene expression, oxidative stress, susceptibility to infection and ultimately reproductive success in a natural population. This begins to reveal how genotypic effects can have multiple effects on different phenotypic or life-history traits for organisms living in the natural environment. Interestingly, effects often differed between sexes, with evidence for opposing effects in the sexes (unstimulated immune gene expression), and an effect present in one sex but not the other (stimulated immune gene expression, susceptibility to infection and reproductive success). However, we cannot rule out that, in the case of susceptibility to infection and reproductive success, there was a smaller effect present in the opposite sex (in the same or opposite direction) which our analysis did not have sufficient power to detect.

In humans, naturally occurring polymorphisms in high-affinity IgE receptor have previously been linked to inflammatory disease [18–21]. Our work is consistent with this and suggests differing effects of this polymorphism on the inflammatory phenotype of males and females. While males with the GC haplotype showed a pro-inflammatory bias, females with the haplotype showed an anti-inflammatory bias. A previous study on another polymorphism in the human ortholog of *Fcer1a* also found evidence for sex-dependent effects on inflammatory disease [24]. In order to fully understand this genotype by sex interaction, it is important to consider background differences in the inflammatory phenotypes of males and females. Females in this study were pregnant for a large proportion of their lives (57% of post-juvenile females were pregnant or lactating at the time of capture in the longitudinal component of our study). In mammals, including

humans, pregnancy has been shown to be a largely anti-inflammatory state, with

pregnant females suppressing inflammation. This is to protect the foetus from attack by

the mother's immune system [30]. While testosterone in males has an anti-inflammatory

effect [31], it also drives males to use more space [32] and to be more aggressive [33].

These behaviours might increase their rates of contact with other individuals, and with

their environment, and hence their likelihood of infection and subsequent immune

stimulation. We suggest that the polymorphism may be exaggerating background sex

differences in inflammatory activity. Sex hormones have been implicated in the

differential expression of some autosomal genes in males and females [34], and could

be driving these opposing effects. Some of the differences in immune phenotype that we

observed may also be driven by difference in parasite infection (although we accounted

for cestode burden in a follow-up analysis, we cannot rule this out).

The effect of *Fcer1a* polymorphism on inflammatory responses may also, in turn, affect

individual susceptibility to infection. This is consistent with a previous association found

in this system between IgE-mediated responses and defence against helminths [13].

Although we did not find an association between macroparasite burden (including

cestode burden) and the GC haplotype in this study, in our previous study, we showed

that males with the GC haplotype had a lower level of an immunological marker of

tolerance to ticks, fleas and adult cestodes (i.e., they were less tolerant) [3]. Here, we

use data on infection incidence to show that, in a different context, the same haplotype is

also associated with resistance to a microparasite, *B. microti*, in females. Although

observational studies of this kind are not able to identify causal relationships, a realistic

scenario is that females with the GC haplotype are more likely to be infected with *B.*

*microti* (i.e., they are less resistant) as a result of a lower pro-inflammatory to anti-

inflammatory cytokine ratio. Pro-inflammatory cytokines (e.g., IL-6, IFN-γ, and TNF-α)

may help to resist *B. microti* infection [35]. A lack or imbalance of these cytokines may

hamper this resistance. The panel of parasites that we have measured is not exhaustive

and previous studies have highlighted the important role of species interactions in the

parasite community in driving infection risk in this study population [36]. *B. microti* may

therefore represent a community of parasites, to which this haplotype affects resistance

in females.

A fitness consequence of the GC haplotype was observed in reduced male reproductive

success. A realistic scenario is that the larger cost incurred by males with the GC

haplotype due to inflammation reduced their reproductive value, as reproducing and

mounting a pro-inflammatory response are both costly activities [37]. Another realistic

scenario is that the positive effect of the GC haplotype on levels of oxidative stress (as

indicated by a tendency for higher SOD1 activity) may have had a more detrimental

effect in males, damaging sperm and reducing fertilising success. Sperm competition

can be an important feature in the reproduction of microtines, e.g., meadow vole (*M.*

*pennsylvanicus*) [38], with males having to produce many sperm of high quality in order

to successfully outcompete other males. Sperm are also highly susceptible to damage

by reactive oxygen species (ROS) [39]. In addition, the heightened level of oxidative

stress may have impaired olfactory sexual signals, e.g., excretion of major urinary

proteins (MUPs) making males less attractive to females and negatively impacting on

mating success [40].

Despite the immune system playing an important role in determining the health and

reproductive fitness of an individual, natural selection has not converged on a single

immune optimum. Instead, individuals in natural populations vary widely in their

response to infection. Understanding why immunoheterogeneity is maintained is a key

question in eco-immunology. Antagonistic selection, where a mutation is beneficial in

some environmental contexts and harmful in others, is thought to be one mechanism by

which balancing selection is generated, and genetic variation in immunity can be

maintained within natural populations [10]. In samples collected between 2008-2010, we

previously identified four haplotypes at this locus, GC, AC, AT and GT, at frequencies of

0.12, 0.76, 0.07 and 0.05 respectively [3]. Seven years on, these frequencies have

remained relatively unchanged (0.08, 0.81, 0.10 and 0.01). The fact that the GC

haplotype remains in the population, despite its detrimental effects on male reproductive

success and female infection susceptibility, suggests that it may be under balancing

selection.

In order to be under balancing selection though, the GC haplotype would need to confer

some advantage. We were not able to find evidence for any such advantage in this

study, but we can speculate on one. Microtines show cyclical population dynamics,

alternating between a "peak" phase (associated with good conditions and high vole

densities) and a "crash" phase (poor conditions and low vole densities). Selection

pressures acting during these two phases will be very different. We found that GC

haplotype was associated with a pro-inflammatory bias in males, indicating higher

investment in promoting inflammation. Thus, one possible scenario is that higher

investment in promoting inflammation is disadvantageous in the "peak" phase when the

risk of parasitism is low due to dilution effects, and it pays not to waste energy mounting

a stronger immune response that is unnecessary. However, it might be advantageous in

the "crash" phase when conditions are poor, and the risk of parasitism is high due to the

time lagged transmission of parasite infectious stages that have accumulated during the

peak phase. Vole numbers were extremely low in our study area between 2012 and

2013 (indicating a "crash phase"). In 2014, vole numbers began to increase and by the

beginning of 2015 (the start of our study) had reached a peak. Although vole numbers

declined throughout our study period, they never reached the extremely low numbers

recorded in 2012-2013 (X. Lambin, personal communication, 2016). This suggests that

we sampled in the region of a "peak" phase, when this investment was disadvantageous,

and that if we sampled during a true "crash" phase we might detect an advantage to the

GC haplotype in males.

There is a growing interest in human genomic (or precision) medicine, with the potential

to use a patient's genotypic information to personalise their treatment. What we have

shown here demonstrates that considering genotype in isolation can be misleading, as

the same polymorphism can have different outcomes not only for the immune gene

expression, but the susceptibility to infection and ultimate reproductive success of males

and females.

**Materials and Methods**

We studied *M. agrestis* in Kielder Forest, Northumberland, UK, using live-trapping of

individual animals from natural populations. Trapping was performed from 2015-2017

across seven different sites, each a forest clear-cut. Access to the sites was provided by

the Forestry Commission. At each site, 150-197 Ugglan small mammal traps (Grahnab,

Gnosjo, Sweden) were laid out in a grid spaced approximately 3-5 m apart. Our study

was divided into longitudinal and cross-sectional components.

**Ethics statement**

All animal procedures were performed with approval from the University of Liverpool

Animal Welfare Committee and under a UK Home Office licence (PPL 70/8210 to S.P.).

**Longitudinal data**

Every other week, traps were checked every day, once in the morning and once in the

evening. Newly trapped field voles were injected with a Passive Integrated Transponder

(PIT) tag (AVID, Uckfield, UK) for unique identification. This approach allowed us to build

up a longitudinal record for voles that were caught on multiple occasions. A total of 850

voles were individually marked in this way. We also took a small tissue sample from the

tail, for genotyping of individuals and a drop of blood from the tail which we put into 500

µl of RNAlater (Fisher Scientific, Loughborough, UK) for use in parasite detection (see

below).

***Parasite detection***

We quantified infections by microparasites (*Babesia microti* and *Bartonella* spp.) in blood

samples taken from longitudinal animals using SYBR green based two-step reverse

transcription quantitative PCR (Q-PCR) targeted at pathogen ribosomal RNA genes.

Expression values were normalized to two host endogenous control genes: *Ywhaz* and

*Actb*. Blood samples were derived from tail bleeds. RNA was extracted from blood

samples stored in RNAlater at -70°C using the Mouse RiboPure Blood RNA Isolation Kit

(ThermoFisher, Waltham, Massachusetts), according to manufacturer's instructions, and

DNAse treated. It was then converted to cDNA using the High-Capacity RNA-to-cDNA™

Kit (ThermoFisher), according to manufacturer's instructions. *B. microti* primer

sequences targeting the 18S ribosomal RNA gene were as follows

CTACGTCCCTGCCCTTTGTA (forward primer sequence) and

CCACGTTTCTTGGTCCGAAT (reverse primer sequence). *Bartonella* spp. primer

sequences targeting the 16S ribosomal RNA gene were as follows

GATGAATGTTAGCCGTCGGG (forward primer sequence) and

TCCCCAGGCGGAATGTTTAA (reverse primer sequences). Assays were pipetted onto

384 well plates by a robot (Pipetmax; Gilson, Middleton, Wisconsin) using a custom

programme and run on a QuantStudio 6-flex Real-Time PCR System (ThermoFisher) at

the machine manufacturers default real-time PCR cycling conditions. Reaction size was

10 µl, incorporating 1 µl of template (diluted 1/20) and PrecisionFAST qPCR Master Mix

(PrimerDesign, Chandler's Ford, UK) with low ROX and SYBR green and primers at the

machine manufacturer's recommended concentrations. Alongside the pathogen assays

we ran assays for two host genes (see above) that acted as endogenous controls. For

the pathogen assays we used as a calibrator sample a pool of DNA extracted from 154

blood samples from different *M. agrestis* at our study sites in 2015 and 2016; these DNA

extractions were carried out using the QIAamp UCP DNA Micro Kit (Qiagen, Hilden,

Germany) following manufacturer's instructions. For the *Ywhaz* and *Actb* assays the

calibrator sample was the same cDNA calibrator sample as described below for other

host gene expression assays. Relative expression values (normalised to the two host

endogenous control genes and presumed to relate to the expression of pathogen

ribosomal RNA genes and in turn to the force of infection) used in analyses below are

RQ values calculated by the QuantStudio 6-flex machine software according to the ΔΔCt

method, indexed to the calibrator samples. Melting curves and amplification plots were

individually inspected for each well replicate to confirm specific amplification. We

validated our diagnostic results by comparing our PCR RQ values to independent data

for a subset of voles from the cross-sectional component of our study with mapped

genus-level pathogen reads from RNASeq analysis of blood samples ($n = 44$), finding

the two data sets strongly corroborated each other (*Bartonella* spp., Spearman's $\rho =$

0.79, $p < 0.001$, $n = 44$; *B. microti*, Spearman's $\rho = 0.88$, $p < 0.001$, $n = 43$).

**Cross-sectional data**

For the cross-sectional component of the study (which ran from 2015-2016 only),

animals were captured and returned to the laboratory where they were killed by a rising

concentration of $CO_2$, followed by exsanguination. As our UK Home Office licence did

not allow us to sample overtly pregnant females in this way, there are fewer females

than males present in this dataset. This component of the study allowed us to take a

more comprehensive set of measurements, and to culture cells and perform stimulatory

assays on them. A small tissue sample was taken from the ear of cross-sectional

animals for genotyping (see below).

***Parasite detection***

After culling, the fur of cross-sectional animals was examined thoroughly under a

binocular dissecting microscope to check for ectoparasites, which were recorded as

direct counts of ticks and fleas. Guts of cross-sectional animals were transferred to 70%

ethanol, dissected and examined under a microscope for gastrointestinal parasites.

Direct counts of cestodes (by far the dominant endohelminths in biomass) were

recorded.

***SOD1 measurement***

Given the well-established link between inflammation, oxidative stress and pathology

[28], we measured antioxidant enzymatic activity in blood samples taken from cross-

sectional animals. We chose to measure superoxide dismutase 1 (SOD1) because it is

one of the main antioxidative enzymes within the antioxidative enzyme system [29], and

is clearly linked to changes in immune gene expression in laboratory mice [41]. Assays

were carried out using the Cayman Superoxide Dismutase kit and following the

manufacturer's instructions except where otherwise indicated below. Blood pellets from

centrifuged cardiac bleeds were stored at -70°C and thawed on ice prior to assay. A 20

µl aliquot from each pellet was lysed in 80 µl of ultrapure water and centrifuged (10000 G

at 4°C for 15 minutes) and 40 µl of the supernatant added to a 1.6 × volume of ice-cold

chloroform/ethanol (37.5/62.5 (v/v)) (inactivating superoxide dismutase 2). This mixture

was then centrifuged (2500 G at 4°C for 10 minutes) and the supernatant removed and

immediately diluted 1:200 in kit sample buffer. A seven-point SOD activity standard was

prepared (0, 0.005, 0.010, 0.020, 0.030, 0.040, 0.050 U/ml) and the assay conducted on

a 96 well microplate with 10 µl of standard or sample, 200 µl of kit radical detector

solution and 20 µl of xanthine oxidase (diluted as per manufacturer's instructions) per

well. Plates were then covered with a protective film, incubated at room temperature for

30 mins on an orbital shaker and read at 450 nm on a VERSAmax tuneable absorbance

plate reader (Molecular Devices, San Jose, California), subtracting background and

fitting a linear relationship to the standard curve in order to estimate SOD activity in

unknown samples.

***Splenocyte cultures***

Spleens of cross-sectional animals were removed, disaggregated, and splenocytes

cultured under cell culture conditions equivalent to those used in [42]. Unstimulated

splenocytes, taken from 84 cross-sectional animals collected between July and October

2015, were initially used to assay expression by RNASeq (see below). We exposed

splenocytes from the remaining cross-sectional animals to stimulation with anti-CD3

antibodies (Hamster Anti-Mouse CD3e, Clone 500A2 from BD Pharmingen, San Diego,

California) and anti-CD28 antibodies (Hamster Anti-Mouse CD28, Clone 37.51 from

Tombo Biosciences, Kobe, Japan) at concentrations of 2 µg/ml and of 1 µg/ml

respectively for 24 hr. Costimulation with anti-CD3 and anti-CD28 antibodies was used

to selectively promote the proliferation of T cells [43,44]. We assumed that this would

reflect the potential response of T-cell populations *in vivo*. Stimulated splenocytes were

used to assay expression by Q-PCR.

**_RNASeq_**

Full details of the methods used for RNA preparation and sequencing can be found in

[3]. Briefly, samples were sequenced on an Illumina HiSeq4000 platform. High-quality

reads were mapped against a draft genome for *M. agrestis* (GenBank Accession no.

LIQJ00000000) and counted using featureCounts [45]. Genes with average log counts

per million, across all samples, of less than one were considered unexpressed and

removed from the data ($n = 8,410$). Following filtering, library sizes were recalculated,

data were normalized and MDS plots were generated to check for any unusual patterns

in the data. The mean library size was 19 million paired-end reads (range = $3 - 71$

million paired-end reads).

**_Q-PCR_**

We used SYBR green based Q-PCR to measure the expression levels of a panel of 18

genes (Appendix 1—table 16) in splenocytes stimulated with anti-CD3 and anti-CD28

antibodies (T cell stimulators) from our cross-sectional animals. We did this, in part, to

validate our RNASeq results in an independent dataset. We used the observed

expression profile as a general measure of the potential responsiveness of the immune

system to an inflammatory stimulation *in vivo*. Although *Fcer1a* is not expressed by T-

cells themselves, polymorphism in this gene could be acting indirectly on T-cells through

various pathways, including via cytokine signalling, following expression of *Fcer1a* by

other cells. The choice of our panel of genes was informed by (i) known immune-

associated functions in mice, combined with (ii) significant sensitivity to environmental or

intrinsic host variables in our previous studies [9,42] or in a recent DGE analysis of

RNASeq data (not reported here), and (iii) the aim of limited redundancy, with each gene

representing a different immune pathway.

Primers (20 sets, including 2 endogenous control genes) were designed *de novo* and

supplied by PrimerDesign (16 sets) or designed *de novo* in-house (4 sets) and validated

(to confirm specific amplification and 100 ± 10% PCR efficiency under assay conditions).

All PrimerDesign primer sets were validated under our assay conditions before use. The

endogenous control genes (*Ywhaz* and *Sdha*) were selected as a stable pairing from our

previous stability analysis of candidate control genes in *M. agrestis* splenocytes [42]. We

extracted RNA from splenocytes conserved in RNAlater using the RNAqueous Micro

Total RNA Isolation Kit (ThermoFisher), following the manufacturer's instructions. RNA

extracts were DNAse treated and converted to cDNA using the High-Capacity RNA-to-

cDNA™ Kit (ThermoFisher), according to manufacturer's instructions, including reverse

transcription negative (RT-) controls for a subsample. SYBR green-based assays were

pipetted onto 384 well plates by a robot (Pipetmax, Gilson) using a custom programme

and run on a QuantStudio 6-flex Real-Time PCR System (ThermoFisher) at the machine

manufacturers default real-time PCR cycling conditions. Reaction size was 10 µl,

incorporating 1 µl of template and PrecisionFAST qPCR Master Mix with low ROX and

SYBR green (PrimerDesign) and primers at the machine manufacturer's recommended

concentrations. We used two standard plate layouts for assaying, each of which

contained a fixed set of target gene expression assays and the two endogenous control

gene assays (the same sets of animals being assayed on matched pairs of the standard

plate layouts). Unknown samples were assayed in duplicate wells and calibrator samples

in triplicate wells and no template controls for each gene were included on each plate.

Template cDNA (see above) was diluted 1/20 prior to assay. The calibrator sample

(identical on each plate) was created by pooling cDNA derived from across all

splenocyte samples. Samples from different sampling groups were dispersed across

plate pairs, avoiding confounding of plate with the sampling structure. Gene relative

expression values used in analyses are RQ values calculated by the QuantStudio 6-flex

machine software according to the $\Delta\Delta$Ct method, indexed to the calibrator sample.

Melting curves and amplification plots were individually inspected for each well replicate

to confirm specific amplification.

**Longitudinal and cross-sectional data**

***Genotyping***

We genotyped both cross-sectional and longitudinal animals for 346 single nucleotide

polymorphisms (SNPs) in 127 genes. See [3] for details of the approach used to select

these SNPs. Our list included two synonymous and tightly linked ($r^2 = 0.50$; D' = 0.70)

SNPs in the gene *Fcer1a* (the alpha chain of the high-affinity receptor for IgE) on

scaffold 582 (CADCXT010006977 in ENA accession GCA_902806775; see Appendix

1—table 1 for genomic location information) which we had previously identified as a

candidate tolerance gene in a natural population of *M. agrestis.* In this previous work, we

had identified four haplotypes at this locus present in our population: GC, AC, AT and

GT, at frequencies of 0.12, 0.76, 0.07 and 0.05 respectively. We had also identified the

GC haplotype as being of particular interest, given its significantly lower expression level

of the transcription factor GATA3 (a biomarker of tolerance to macroparasites in our

population) compared to the other haplotypes [3]. We concluded that this haplotype

tagged a causal mutation in the coding sequence, or in the up or downstream regulatory

regions. DNA was extracted from a tail sample (longitudinal component) or an ear

sample (cross-sectional component) taken from the animal using DNeasy Blood and

Tissue Kit (Qiagen). Genotyping was then performed by LGC Biosearch Technologies

(Hoddesdon, UK; http://www.biosearchtech.com) using the KASP SNP genotyping

system. This included negative controls (water) and duplicate samples for validation

purposes. The resulting SNP dataset was used for two purposes (i) genotyping

individuals within the locus of interest, and (ii) pedigree reconstruction (see below).

***Pedigree reconstruction***

We used a subset of our SNP dataset to reconstruct a pedigree for both cross-sectional

and longitudinal animals using the R package *Sequoia* [46]. SNPs which violated the

assumptions of Hardy-Weinberg equilibrium were removed from the dataset. For pairs of

SNPs in high linkage disequilibrium (most commonly within the same gene), the SNP

with the highest minor allele frequency (MAF) was chosen. A minimum MAF cut-off of

0.1 and call rate of > 0.7 was then applied, and any samples for which more than 50% of

SNPs were missing were removed. This resulted in a final dataset including 114 SNPs -

a sufficient number for very good performance of parentage assignment [46].

Life history information, namely sex and month of birth, was inputted into *Sequoia* where

possible. Juvenile voles weighing less than 12 g on first capture were assigned a birth

date 2 weeks prior to capture. Juvenile voles weighing between 12 and 15 g on first

capture were assigned a birth date 4 weeks prior to capture. Finally, adult voles breeding

on first capture, were assigned a birth date 6 weeks prior to capture (minimum age at

first breeding) [47,48]. Adult voles not breeding on first capture could not be assigned a

birth date, as it was not known whether they had previously bred or not. Virtually all

samples (99%) were assigned a sex, and approximately half (54%) were assigned a

birth month. As we sampled individuals from across seven different clear-cut areas of the

forest, each several kilometres apart, these were assumed to be independent, closed

populations with negligible dispersal. Site-specific pedigrees were therefore generated.

We assessed the accuracy of our reconstructed pedigrees by checking whether

predicted parent-offspring pairs met expectations given the biology of *M. agrestis*. As

expected, the majority of predicted parent-offspring pairs (87%) were born in the same

year. We also expected parents and offspring to overlap in space. Again, as expected,

the majority of predicted parent-offspring pairs (92%) were, at some point, found along

the same transects (horizontal or vertical). We also inspected log10 likelihood ratios

(LLRs) for parent pairs as recommended in the user manual for *Sequoia*. Almost all

LLRs were positive ($n$ = 698/720 or 97% of LLRs) indicating confidence in our

assignments. Individuals with vs. without our haplotype of interest did not differ in their

probability of appearing in a pedigree ($\chi^2$ = 0.09, d.f. = 1, $p$ = 0.76). For each individual

that ended up in a pedigree i.e. with one or more relatives recorded ($n = 652$; Site COL =

3; Site BLB = 125 ; Site GRD = 204; Site CHE = 137; Site RAV = 16; Site  SCP = 90;

Site HAM = 77), the number of offspring was counted to provide a measure of their

reproductive success. Few individuals were first trapped as juveniles, with the majority

trapped as adults which had already recruited into the population. Our measure of

reproductive success then, more closely resembles the number of recruited (rather than

newborn) offspring per individual. Half of individuals present in our pedigrees ($n = 325$)

were found to have no offspring. We expect the majority of these to be true zeros

(representing actual reproductive failure) as we generally sampled a large proportion of

the total population within clear-cuts. We also minimised the chance of false zeros by

excluding those individuals (e.g. at the periphery of a study grid) which did not end up in

a pedigree because we identified no relatives (including offspring), likely because we

had not sampled in the right place.

**Statistical analyses**

Not all individuals appeared in all datasets, therefore sample sizes (reported in Table 1)

vary between analyses. All analyses were performed in R statistical software version

3.5.2 [49].

*Differential gene expression analysis*

DGE analysis was performed on filtered and normalized count data using the R package

edgeR [50], the aim being to identify individual genes which were differentially expressed

between those individuals with and without a copy of the GC haplotype (i.e. a dominant

model). Only those individuals for which haplotype could be inferred with certainty could

be included ($n$ = 53 males and $n$ = 31 females; none of which were known to have two

copies of the GC haplotype, hence the choice of a dominant model). Samples from

different sexes were analysed together. To test whether the top-responding genes

differed between the sexes, we included a dummy variable with four levels (males with

haplotype, males without haplotype, females with haplotype, females without

haplotypes) and inspected the contrasts of interest (males with versus without

haplotypes; females with versus without haplotype). As this was an exploratory analysis,

used in conjunction with more targeted measurements of immune gene expression (see

below), model specification was kept as simple as possible and no covariates were

initially included. However, in a follow-up DGE analysis we controlled for a key parasite

variable (cestode burden).

Having confirmed a sex-dependent effect of the GC haplotype on the expression of

some inflammatory genes (see Results) in the initial DGE analysis, we ran separate

DGE analyses for males and females and tested more broadly, for enrichment of pro-

and anti-inflammatory genes in our results; more specifically, the gene ontology terms,

'positive regulation of inflammatory response' (GO:0050729; $n$ = 143) and 'negative

regulation of inflammatory response' (GO:0050728; $n$ = 149). The aim here was to

answer the question: are pro- or anti-inflammatory genes more highly ranked relative to

other genes, when we compare individuals with and without the GC haplotype, and does

this also differ between the sexes? This GSEA was performed using the R package

limma [51], and genes were ranked on log fold change.

***Haplotype association analyses***

Following the exploratory DGE analysis, the GC haplotype was tested for associations

with (i) gene expression assayed by Q-PCR to validate these results, (ii) macro- and

microparasite infection to test whether the GC haplotype predicted an individual's

susceptibility to infection, and (iii) reproductive success to test whether the GC haplotype

was associated with any reproductive costs. Given the well-established link between

inflammation and oxidative stress [52,53], we also tested for an association between the

GC haplotype and SOD1 activity.

For all analyses, this was initially attempted using the R package hapassoc [54] in order

to maximise sample size (see below). Hapassoc infers haplotypes on the basis of data

from single SNPs, and allows likelihood inference of trait associations with resulting SNP

haplotypes and other attributes. It adopts a generalized linear model framework and

estimates parameters using an expectation–maximization algorithm. Hapassoc models

assumed an additive genetic model. If the haplotype combination of an individual cannot,

with certainty, be inferred from its genotyping data (i) because it is heterozygous at two

or more markers or (ii) because it has missing data for a single marker, the approach

implemented in hapassoc is to consider all possible haplotype combinations for that

individual. Standard errors accounting for this added uncertainty are calculated using the

Louis' method [55]. We compared the GC haplotype against the other two major

haplotypes (AC and AT). Another haplotype, the GT haplotype, was identified in the

population but this was present at such low frequencies (frequency $\leq 0.01$ among

individuals for which haplotype could be inferred with certainty) that it was omitted from

all analyses. Results reported in the text for macroparasites and SOD1 activity come

from these hapassoc models.

However, there are some restrictions on model specification within hapassoc (e.g.

random terms cannot be included, limited choice of error distributions), so this was

followed up with regression models for some analyses. As in the DGE analysis, these

regression models only included those individuals whose haplotype combination could

be inferred with certainty.  Results reported in the text for gene expression assayed by

Q-PCR, microparasites and reproductive success come from these regression models.

Again, as in the DGE analysis, genotype was coded as the presence or absence of the

GC haplotype (i.e. a dominant model). Regression models were run using the R package

lme4 [56] or glmmADMB [57,58].

Table 1 provides a summary of (hapassoc or regression) model specifications. As we

found evidence for a sex-dependent effect in the initial DGE analysis, with the exception

of the model for reproductive success (see below), all models included a genotype by

sex interaction which was retained if it improved model fit. All models also accounted for

other biological and technical covariates (the choice of which was informed by our

previous work, the literature and/or our experimental design). Full regression models

(including all covariates and interactions of interest) were reduced using backward

stepwise deletion of non-significant terms to minimize Akaike's information criterion

(AIC), following the drop1 function in the lme4 package. This was not possible for

hapassoc models. Summary tables including all estimates, standard errors and $z$-

statistics are included in Appendix 1. Associated significance values are also included in

these tables, except for LMMs and GLMMs, for which significance values were based on

likelihood ratio tests (LRTs) and are reported in the main text. Throughout, residuals

from regression models were checked for approximate normality and homoscedasticity,

and all covariates were tested for independence using variance inflation factors (all VIFs

< 3).

***Association between GC haplotype and immune gene expression assayed by Q-***

***PCR***

As we ran a total of 18 hapassoc models (one model per gene), the Benjamini and

Hochberg method of correction was applied to all *p*-values [59]. Resulting *q*-values

(FDR-corrected *p*-values) are reported, alongside original *p*-values (Appendix 1—table

6). A hapassoc model including a genotype by sex interaction was tested against a null

model without an interaction term. A Gamma error distribution and log link were used.

Other covariates considered potential drivers of immune gene expression and included

in both models, were informed by our previous work [42]. These included snout-vent

length (SVL), eye lens weight (categorised into seven intervals; SVL and eye lens weight

capture the combined influence of age and historical growth trajectory), reproductive

status (males were considered to be reproductively active if they had descended testes;

females if they were pregnant or had perforate vaginas) and body condition (estimated

by regressing body weight against life history stage, SVL and its quadratic term). Site,

year and season (four levels, designated as spring [March & April], early summer [May

and June], late summer [July and August] and autumn [September and October]) were

included to account for any spatial and/or temporal autocorrelation. All covariates were

included as fixed effects. We did not use a multidimensional approach (such as principal

component analysis) because of limited redundancy in our panel of genes.

In order to confirm these results, a linear mixed effects model (LMM) was run for a single

immune gene for which expression appeared to be associated with genotype (*q* =

0.037). The LMM included random terms for site and season, as well as assay plate

number (Table 1). The latter was included to account for nonindependence due to

immunological assaying structure. All other covariates were the same as the hapassoc

model. A Yeo-Johnson transformation (with $\lambda$ = -2) was used to achieve more normal

and homoscedastic residuals [60].

***Association between GC haplotype and parasite infection***

The three macroparasite measurements taken from cross-sectional animals (counts of

ticks, fleas and cestodes) were log-transformed ($\log_{10}$ [$x$ +1]) and summarised as a

single principal component (explaining 39% of total variation; Appendix 1—table 17) to

avoid difficulties in interpretation due to multiple testing. See [3] and [9] for full details of

this approach. This combined measure of macroparasite burden was modelled using a

hapassoc model with a Gaussian error distribution.

Microparasite infection status was assessed multiple times for the majority of individuals

in the longitudinal component of the study (mean = 2.8; range = 1-11). Due to these

repeated measures, microparasite infection could not be modelled using hapassoc.

Instead, to test for an association between the GC haplotype and the probability of an

individual being infected with a microparasite, we ran a GLMM with a binary response

(infected or not), log link and a random term for individual. Other covariates, considered

potential drivers of both macro and microparasite infection were, again, informed by our

previous work [3,9,61]. These included body condition, reproductive status, year, season

and site (Table 1). Season and site were included as random terms in the GLMM for

microparasites, as was individual identity, to account for nonindependence due to repeat

sampling.

***Association between GC haplotype and reproductive success***

Our measure of reproductive success was zero-inflated (50% zeros). This is consistent

with a previous study of the closely-related common vole (*Microtus arvalis*) [62]. A

Poisson error distribution was therefore deemed inappropriate and it could not be

modelled using hapassoc. Instead, to test for an association between the GC haplotype

and reproductive success, we ran a GLM with a quasi-Poisson error distribution. This

distribution has been previously used to model predictors of reproductive success in

other organisms, and accounts for the overdispersion caused by excess zeros.

We only recorded a reproductive failure (i.e. zero reproductive success) for those

individuals with some other familial relationship in our pedigree, purposefully omitting

those individuals (e.g. at the periphery of a study grid) for which we may have recorded

no relatives (including offspring) simply because we had not sampled in the right place.

Therefore, we expect all (or most) of our zeros to represent actual reproductive failure.

For this reason, zero-inflated models were deemed inappropriate.

As detailed in the Results, we tried to run a single quasi-Poisson GLM with both sexes

but resulting residual variances differed significantly between the sexes (F test to

compare variances of two samples; $p = 0.02$), making it impossible to formally test for a

genotype by sex interaction. Instead, we ran a separate model for each sex. We

included birth month as a covariate in this model, given that autumn-born voles (of the

closely-related common vole) have been shown to have a lower chance to reproduce

than spring-born voles [62]. Other covariates included in this model were, whether or not

an individual was culled for the cross-sectional component of this study (again, reducing

the opportunity to reproduce), site and the year in which an individual was most

frequently captured (Table 1). In a follow-up analysis, we also controlled for *Babesia*

*microti* and *Bartonella* spp. infection status. More specifically, we included the proportion

of samples taken from an individual that were *Babesia*-positive, and the proportion of

samples taken from an individual that were *Bartonella*-positive. All covariates were

included as fixed effects. Only a single female was trapped in one of the sites (COL) and

consequently caused convergence problems in the female model. This female was

therefore omitted.

***Association between GC haplotype and SOD1 activity***

SOD1 activity was modelled using a hapassoc model with a Gaussian error distribution.

Other covariates, considered potential drivers of SOD1 activity, were informed by the

literature. Previous studies on wild rodents have shown that antioxidant levels increase

with both reproductive effort [63] and with age [64]. Studies on birds have also shown

that improvements in body condition are often accompanied by increases in antioxidant

activity, for example in response to supplemental feeding [65]. We therefore included

SVL, eye lens weight, reproductive status and body condition as covariates in our model.

As in other models, we included site, year and season to account for spatial and/or

temporal autocorrelation in our data (Table 1). All covariates were included as fixed

effects.

**Acknowledgements**

We would like to thank all those involved in obtaining and processing samples from the

field: Rebecca Turner, Lukasz Lukomski, Stephen Price, Sarah Gore, Ed Parker, Maria

Capstick, Noelia Dominguez Alvarez, Susan Withenshaw, William Foster, Ann Lowe and

Benoit Poulin. We would also like to thank the Forestry Commission for access to the

study sites and the Centre for Genomic Research at the University of Liverpool for

sequencing samples. This research was funded by the Natural Environment Research

Council (NERC) award NE/L013452/1 to S.P., M.B., J.A.J. and J.E.B.

**Competing interests**

The authors declare no competing interests.

**Data availability**

RNASeq data have been deposited in the European Nucleotide Archive (study

accession number PRJEB51626). Longitudinal phenotypic data are available from the

NERC EDS Environmental Information Data Centre (doi:10.5285/e5854431-6fa4-4ff0-

aa02-3de68763c952). Cross-sectional phenotypic data are available from the University

of Liverpool's Research Data Catalogue (doi:10.17638/datacat.liverpool.ac.uk/1850).

Genotype data are also available from the University of Liverpool's Research Data

Catalogue (doi:10.17638/datacat.liverpool.ac.uk/1849).

**References**

1.    Møller AP, Saino N. Immune response and survival. Oikos. 2004;104: 299–304.

doi:10.1111/j.0030-1299.2004.12844.x

2.    Paterson S, Wilson S, Pemberton J. Major histocompatibility complex variation

associated with juvenile survival and parasite resistance in a large unmanaged

ungulate population (*Ovis aries* L.). Proc Natl Acad Sci USA. 1998;95: 3714–

3719. doi:10.1073/pnas.95.7.3714

3.     Wanelik KM, Begon M, Birtles RJ, Bradley JE, Friberg IM, Jackson JA, et al. A

candidate tolerance gene identified in a natural population of field voles (*Microtus*

*agrestis*). Mol Ecol. 2018;27: 1044–1052. doi:10.1111/mec.14476

4.     Turner AK, Begon M, Jackson JA., Bradley JE, Paterson S. Genetic Diversity in

Cytokines Associated with Immune Variation and Resistance to Multiple

Pathogens in a Natural Rodent Population. PLoS Genet. 2011;7: e1002343.

doi:10.1371/journal.pgen.1002343

5.     Hill AVS, Allsopp CEM, Kwiatkowski D, Anstey NM, Twumasi P, Rowe PA, et al.

Common West African HLA antigens are associated with protection from severe

malaria. Nature. 1991;352: 595–600. doi:10.1038/352595a0

6.     Carrington M, Nelson GW, Martin MP, Kissner T, Vlahov D, Goedert JJ, et al. HLA

and HIV-1: Heterozygote Disadvantage and *B\*35-Cw\*04* Disadvantage. Science.

1999;283: 1748–1752.

7.     Thursz MR, Thomas HC, Greenwood BM, Hill A V. Heterozygote advantage for

HLA class-II type in hepatitis B virus infection. Nat Genet. 1997;17: 11–12.

doi:10.1038/ng0997-11

8.     Ober C, Loisel DA, Gilad Y. Sex-specific genetic architecture of human disease.

Nat Rev Genet. 2008;9: 911–922. doi:10.1038/nrg2415

9.     Jackson JA, Hall AJ, Friberg IM, Ralli C, Lowe A, Zawadzka M, et al. An

immunological marker of tolerance to infection in wild rodents. PLoS Biol.

2014;12: e1001901. doi:10.1371/journal.pbio.1001901

10.    Graham AL, Hayward AD, Watt KA, Pilkington JG, Pemberton JM, Nussey DH.

Fitness correlates of heritable variation in antibody responsiveness in a wild

mammal. Science. 2010;330: 662–665. doi:10.1126/science.1194878

11.    Turner AK, Beldomenico PM, Bown K, Burthe SJ, Jackson JA, Lambin X, et al.

Host-parasite biology in the real world: the field voles of Kielder. Parasitology.

2014;141: 997–1017. doi:10.1017/S0031182014000171

12.  Turner AK, Paterson S. Wild rodents as a model to discover genes and pathways

underlying natural variation in infectious disease susceptibility. Parasite Immunol.

2013;35: 386–395. doi:10.1111/pim.12036

13.  Gounni AS, Lamkhioued B, Ochiai K, Tanaka Y, Delaporte E, Capron A, et al.

High-affinity IgE receptor on eosinophils is involved in defence against parasites.

Nature. 1994;367: 183–186. doi:10.1038/368473a0

14.  Tomassini M, Tsicopoulos A, Tai PC, Gruart V, Tonnel AB, Prin L, et al. Release

of granule proteins by eosinophils from allergic and nonallergic patients with

eosinophilia on immunoglobulin-dependent activation. J Allergy Clin Immunol.

1991;88: 365–375. doi:10.1016/0091-6749(91)90099-A

15.  Daeron M, Nimmerjahn F, editors. Current Topics in Microbiology and

Immunology: Fc Receptors. Immunology. Springer; 2014.

16.  Weidinger S, Gieger C, Rodriguez E, Baurecht H, Mempel M, Klopp N, et al.

Genome-wide scan on total serum IgE levels identifies FCER1A as novel

susceptibility locus. PLoS Genet. 2008;4: e1000166.

doi:10.1371/journal.pgen.1000166

17.  Granada M, Wilk JB, Tuzova M, Strachan DP, Weidinger S, Albrecht E, et al. A

genome-wide association study of plasma total IgE concentrations in the

Framingham Heart Study. J Allergy Clin Immunol. 2012;129: 840–845.

doi:10.1016/j.jaci.2011.09.029

18.  Hasegawa M, Nishiyama C, Nishiyama M, Akizawa Y, Mitsuishi K, Ito T, et al. A

Novel –66T/C Polymorphism in FcεRI α-Chain Promoter Affecting the

Transcription Activity: Possible Relationship to Allergic Diseases. J Immunol.

2003;171: 1927–1933. doi:10.4049/jimmunol.171.4.1927

19.    Potaczek DP, Sanak M, Mastalerz L, Setkowicz M, Kaczor M, Nizankowska E, et

al. The α-chain of high-affinity receptor for IgE (FcεRIα) gene polymorphisms and

serum IgE levels. Allergy. 2006;61: 1230–1233. doi:10.1111/j.1398-

9995.2006.01195.x

20.    Zhou J, Zhou Y, Lin L hui, Wang J, Peng X, Li J, et al. Association of

polymorphisms in the promoter region of FCER1A gene with atopic dermatitis,

chronic uticaria, asthma, and serum immunoglobulin E levels in a Han Chinese

population. Hum Immunol. 2012;73: 301–305. doi:10.1016/j.humimm.2011.12.001

21.    Niwa Y, Potaczek DP, Kanada S, Takagi A, Shimokawa N, Lto T, et al. FcεRIα

gene (FCER1A) promoter polymorphisms and total serum IgE levels in Japanese

atopic dermatitis patients. Int J Immunogenet. 2010;37: 139–141.

doi:10.1111/j.1744-313X.2010.00901.x

22.    Weiss LA, Pan L, Abney M, Ober C. The sex-specific genetic architecture of

quantitative traits in humans. Nat Genet. 2006;38: 218–222. doi:10.1038/ng1726

23.    Chen W, Mempel M, Schober W, Behrendt H, Ring J. Gender difference, sex

hormones, and immediate type hypersensitivity reactions. Allergy. 2008;63: 1418–

1427. doi:10.1111/j.1398-9995.2008.01880.x

24.    Yang J, Lu MM, Lu YW, Feng CC, Leng RX, Pan HF, et al. Sex-specific

differences in the relationship between the single-nucleotide polymorphism

rs2298804 of FCER1A and the susceptibility to systemic lupus erythematosus in a

Chinese Han population. Clin Exp Dermatol. 2013;38: 410–416.

doi:10.1111/ced.12035

25.    Liew FY, Pitman NI, McInnes IB. Disease-associated functions of IL-33: The new

kid in the IL-1 family. Nat Rev Immunol. 2010;10: 103–110. doi:10.1038/nri2692

26.  Carow B, Rottenberg ME. SOCS3, a major regulator of infection and

inflammation. Front Immunol. 2014;5: 58. doi:10.3389/fimmu.2014.00058

27.  Cayrol C, Girard JP. IL-33: An alarmin cytokine with crucial roles in innate

immunity, inflammation and allergy. Curr Opin Immunol. 2014;31: 31–37.

doi:10.1016/j.coi.2014.09.004

28.  Biswas SK. Does the Interdependence between Oxidative Stress and

Inflammation Explain the Antioxidant Paradox? Oxid Med Cell Longev.

2016;2016: 5698931. doi:10.1155/2016/5698931

29.  Sun Y, Oberley LW, Elwell JH, Sierra-Rivera E. Antioxidant enzyme activities in

normal and transformed mouse liver cells. Int J Cancer. 1989;44: 1028–1033.

doi:10.1002/ijc.2910440615

30.  Wegmann TG, Lin H, Guilbert L, Mosmann TR. Bidrectional cytokine interactions

in the maternal-fetal relationship: is successful pregnancy a TH2 phenomenon?

Immunol Today. 1993;14: 353–356. doi: 10.1016/0167-5699(93)90235-D

31.  Bianchi VE. The Anti-Inflammatory Effects of Testosterone. J Endocr Soc. 2019;3:

91–107. doi:10.1210/js.2018-00186

32.  Davis S, Abbasi B, Shah S, Telfer S, Begon M, Davis S. Spatial analyses of

wildlife contact networks. J R Soc Interface. 2014;12: 20141004.

doi:10.1098/rsif.2014.1004

33.  Martínez-Sanchis S, Arnedo MT, Salvador A, Moya-Albiol L, González-Bono E.

Effects of Chronic Administration with High Doses of Testosterone Propionate on

Behavioral and Physiological Parameters in Mice with Differing Basal

Aggressiveness. Aggress Behav. 2003;29: 173–189. doi:10.1002/ab.10050

34.  Mayne BT, Bianco-Miotto T, Buckberry S, Breen J, Clifton V, Shoubridge C, et al.

Large scale gene expression meta-analysis reveals tissue-specific, sex-biased

gene expression in humans. Front Genet. 2016;7: 183.

doi:10.3389/fgene.2016.00183

35.  Djokic V, Akoolo L, Parveen N. *Babesia microti* infection changes host spleen

architecture and is cleared by a Th1 immune response. Front Microbiol. 2018;9:

85. doi:10.3389/fmicb.2018.00085

36.  Telfer S, Lambin X, Birtles R, Beldomenico P, Burthe S, Paterson S, et al.

Species interactions in a parasite community drive infection risk in a wildlife

population. Science. 2010;330: 243–246. doi:10.1126/science.1190333

37.  Sheldon BC, Verhulst S. Ecological immunology: costly parasite defences and

trade-offs in evolutionary ecology. Trends Ecol Evol. 1996;11: 317–321.

38.  DelBarco-Trillo J, Ferkin MH. Male mammals respond to a risk of sperm

competition conveyed by odours of conspecific males. Nature. 2004;431: 446–

449. doi:10.1038/nature02845

39.  Losdat S, Richner H, Blount JD, Helfenstein F. Immune activation reduces sperm

quality in the great tit. PLoS One. 2011;6: e22221.

doi:10.1371/journal.pone.0022221

40.  Garratt M, Pichaud N, Glaros EN, Kee AJ, Brooks RC. Superoxide dismutase

deficiency impairs olfactory sexual signaling and alters bioenergetic function in

mice. Proc Natl Acad Sci USA. 2014;111: 8119–8124.

doi:10.1073/pnas.1322282111

41.  Marikovsky M, Ziv V, Nevo N, Harris-Cerruti C, Mahler O. Cu/Zn Superoxide

Dismutase Plays Important Role in Immune Response. J Immunol. 2003;170:

2993–3001. doi:10.4049/jimmunol.170.6.2993

42.  Jackson JA, Begon M, Birtles R, Paterson S, Friberg IM, Hall A, et al. The

analysis of immunological profiles in wild animals: A case study on

immunodynamics in the field vole, *Microtus agrestis*. Mol Ecol. 2011;20: 893–909.

doi:10.1111/j.1365-294X.2010.04907.x

43.  Frauwirth KA, Thompson CB. Activation and inhibition of lymphocytes by

costimulation. Cancer Res. 2002;109: 295–299.

doi:10.1172/JCI200214941.Mounting

44.  Wanelik K, Begon M, Arriero E, Bradley J, Friberg I, Jackson J, et al.

Transcriptome-wide analysis reveals different categories of immune response in a

wild rodent. Sci Rep. 2020;10: 7444. doi:10.1038/s41598-020-64307-7

45.  Liao Y, Smyth GK, Shi W. featureCounts: An efficient general purpose program

for assigning sequence reads to genomic features. Bioinformatics. 2014;30: 923–

930. doi:10.1093/bioinformatics/btt656

46.  Huisman J. Pedigree reconstruction from SNP data: parentage assignment,

sibship clustering and beyond. Mol Ecol Resour. 2017;17: 1009–1024.

doi:10.1111/1755-0998.12665

47.  Begon M, Telfer S, Burthe S, Lambin X, Smith MJ, Paterson S. Effects of

abundance on infection in natural populations: field voles and cowpox virus.

Epidemics. 2009;1: 35–46. doi:10.1016/j.epidem.2008.10.001

48.  Burthe SJ, Lambin X, Telfer S, Douglas A, Beldomenico P, Smith A, et al.

Individual growth rates in natural field vole, *Microtus agrestis*, populations

exhibiting cyclic population dynamics. Oecologia. 2010;162: 653–661.

doi:10.1007/s00442-009-1495-6

49.  R Core Team. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-

project.org/. 2018.

50.  Robinson MD, McCarthy DJ, Smyth GK. edgeR: A Bioconductor package for

945  differential expression analysis of digital gene expression data. Bioinformatics.

946  2010;26: 139–40. doi:10.1093/bioinformatics/btp616

947 51. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. Limma powers

948  differential expression analyses for RNA-sequencing and microarray studies.

949  Nucleic Acids Res. 2015;43: e47. doi:10.1093/nar/gkv007

950 52. Reuter S, Gupta S, Chaturvedi M, Aggarwal B. Oxidative stress, inflammation,

951  and cancer: How are they linked? Free Radic Biol Med. 2011;49: 1603–1616.

952  doi:10.1016/j.freeradbiomed.2010.09.006.Oxidative

953 53. Collins AR. Oxidative DNA damage, antioxidants, and cancer. BioEssays.

954  1999;21: 238–246. doi:10.1002/(SICI)1521-1878(199903)21:3<238::AID-

955  BIES8>3.0.CO;2-3

956 54. Burkett K, Graham J, McNeney B. hapassoc: Software for likelihood inference of

957  trait associations with SNP haplotypes and other attributes. J Stat Softw. 2006;16:

958  1–19.

959 55. Louis T. Finding the observed information matrix when using the EM Algorithm. J

960  R Stat Soc B. 1982;44: 226–233.

961 56. Bates D, Machler M, Bolker B, Walker S. Fitting linear mixed-effects models using

962  lme4. J Stat Softw. 2015;67: 1–48.

963 57. Skaug H, Fournier D, Bolker B, Magnusson A, Nielsen A. Generalized Linear

964  Mixed Models using "AD Model Builder". R package version 0.8.3.3. 2016.

965 58. Fournier DA, Skaug HJ, Ancheta J, Ianelli J, Magnusson A, Maunder MN, et al.

966  AD Model Builder: Using automatic differentiation for statistical inference of highly

967  parameterized complex nonlinear models. Optim Methods Softw. 2012;27: 233–

968  249. doi:10.1080/10556788.2011.597854

969 59. Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and

970   powerful approach to multiple testing. J R Stat Soc Ser B. 1995;57: 289–300.

971 60. Yeo I-K, Johnson RA. A new family of power transformations to improve normality

972   or symmetry. Biometrika. 2000;87: 954–959.

973 61. Taylor CH, Wanelik KM, Friberg IM, Lowe A, Hall AJ, Ralli C, et al. Physiological,

974   but not fitness, effects of two interacting haemoparasitic infections in a wild

975   rodent. Int J Parasitol. 2018;48: 463–471. doi:10.1016/j.ijpara.2017.11.006

976 62. Wang M, Ebeling M, Hahne J. Relevance of body weight effects for the population

977   development of common voles and its significance in regulatory risk assessment

978   of pesticides in the European Union. Environ Sci Eur. 2019;31: 54.

979   doi:10.1186/s12302-019-0240-y

980 63. Garratt M, Vasilaki A, Stockley P, McArdle F, Jackson M, Hurst JL. Is oxidative

981   stress a physiological cost of reproduction? An experimental test in house mice.

982   Proc R Soc B Biol Sci. 2011;278: 1098–1106. doi:10.1098/rspb.2010.1818

983 64. Hindle AG, Lawler JM, Campbell KL, Horning M. Muscle aging and oxidative

984   stress in wild-caught shrews. Comp Biochem Physiol B Biochem Mol Biol.

985   2010;155: 427–434. doi:10.1016/j.cbpb.2010.01.007

986 65. Wilcoxen TE, Horn DJ, Hogan BM, Hubble CN, Huber SJ, Flamm J, et al. Effects

987   of bird-feeding activities on the health of wild birds. Conserv Physiol. 2015;3:

988   cov058. doi:10.1093/conphys/cov058

**Figure legends and main tables**

**Figure 1.** Effects of GC haplotype (hGC). Upper panel: males. Lower panel: females: (A) Unstimulated immune gene expression: Barcode plots showing enrichment of the GO terms GO:0050729 (pro-inflammatory genes) and GO:0050728 (anti-inflammatory genes) in unstimulated splenocytes taken from individuals with (hGC+) vs. without (hGC-) the haplotype, showing that males with the haplotype have a pro-inflammatory bias, whereas females have an anti-inflammatory bias. In each plot, x-axis shows log fold change (logFC) in hGC+ vs. hGC-, black bars represent genes annotated with the GO terms and the worm shows relative enrichment. (B) Susceptibility to infection: Association between hGC and the odds of infection with *Babesia microti,* showing that females with the haplotype have an increased susceptibility to infection. (C) Reproduction: Association between hGC and reproductive success, showing that males with the haplotype have lower reproductive success (error bars represent ± standard error; * indicates $p < 0.05$; ** indicates $p < 0.01$).

**Table 1.** Model specifications including, for each main model: covariates included in the full model, datasets used and sample sizes (F = included as a fixed effect; R = included as a random effect).

| | DGE analysis | Haplotype association analyses | | | |
|---|---|---|---|---|---|
| | | | **Response variable** | | |
| | | Immune gene expression assayed by Q-PCR | Parasite infection | Reproductive success | SOD1 activity |
| **Covariates** | | | | | |
| Snout-vent length | | F | | | F |
| Eye lens weight | | F | | | F |
| Reproductive status | | F | F | | F |
| Body condition | | F | F | | F |
| Birth month | | | | F | |
| Culled or not | | | | F | |
| Site | | F; R (*Il17a* LMM) | F (Macro); R (Micro) | F | F |
| Year | | F | F | F | F |
| Season | | F; R (*Il17a* LMM) | F (Macro); R (Micro) | | F |
| Individual ID | | | R (Micro) | | |
| Assay plate | | R (*Il17a* LMM) | | | |
| **Dataset[a]** | C | C | C (Macro) L (Micro) | C + L | C |
| **Sample size** | ♀31 ♂53 | *Il17a* LMM: ♀73 ♂220 | Macro: ♀82 ♂235 *B. microti*: ♀1075 ♂1247 *Bartonella* spp.: ♀1283 ♂1104 | ♀419 ♂232 | ♀81 ♂227 |

[a] C, cross-sectional; L, longitudinal.

1024   **Appendix 1**
1025
1026   **Appendix 1—table 1.** Position of SNPs and other key features in the *Fcer1a* gene.
1027   Features lie in scaffold CADCXT010006977 within assembly GCA_902806775.

| Feature | Start | End |
|---|---|---|
| *Fcer1a* gene | 10745528 | 10745528 |
| Exon 5 | 10745528 | 10745881 |
| Exon 4 | 10746818 | 10747076 |
| Exon 3 | 10748609 | 10748864 |
| Exon 2 | 10750044 | 10750065 |
| Exon 1 | 10750465 | 10750642 |
| 5' UTR | 10750520 | 10750642 |
| CDS | 10750465 | 10750520 |
| CDS | 10750044 | 10750065 |
| CDS | 10748609 | 10748864 |
| CDS | 10746818 | 10747076 |
| CDS | 10745729 | 10745881 |
| 3' UTR | 10745528 | 10745729 |
| SNP 2 | 10748718 | 10748719 |
| SNP 1 | 10746846 | 10746847 |

1028

1029   **Appendix 1—table 2.** Top 10 annotated genes which were differentially expressed
1030   between males with vs. without the GC haplotype, including associated log fold changes
1031   (logFC), *p*-values and *q*-values (FDR-corrected *p*-values).
1032

| Gene | Protein | LogFC | *p*-value | *q*-value |
|---|---|---|---|---|
| *Snai3* | Snail family zinc finger 3 | 1.698 | $1.220 \times 10^{-9}$ | $1.574 \times 10^{-5}$ |
| *Pla2g4c* | Phospholipase A2, group IVC (cytosolic, calcium-independent) | 2.904 | $6.386 \times 10^{-9}$ | $4.120 \times 10^{-5}$ |
| *Il33* | Interleukin 33 | 2.756 | $1.474 \times 10^{-7}$ | $4.755 \times 10^{-4}$ |
| *Mmp13* | Matrix metallopeptidase 13 | 2.571 | $4.427 \times 10^{-7}$ | $1.142 \times 10^{-3}$ |
| *Uba7* | Ubiquitin-like modifier activating enzyme 7 | 0.771 | $5.979 \times 10^{-7}$ | $1.286 \times 10^{-3}$ |
| *Robo4* | Roundabout guidance receptor 4 | 0.892 | $3.738 \times 10^{-6}$ | $4.385 \times 10^{-3}$ |
| *Ttn* | Titin | 1.665 | $8.995 \times 10^{-6}$ | $7.254 \times 10^{-3}$ |
| *Flnc* | Filamin C, gamma | 1.554 | $1.245 \times 10^{-5}$ | $9.447 \times 10^{-3}$ |
| *Crb2* | Crumbs family member 2 | 1.327 | $1.567 \times 10^{-5}$ | $1.063 \times 10^{-2}$ |
| *Muc16* | Mucin 16 | 1.937 | $1.971 \times 10^{-5}$ | $1.156 \times 10^{-2}$ |

1033
1034
1035   **Appendix 1—table 3.** Annotated genes which were differentially expressed ($q \leq 0.1$)
1036   between females with vs. without the GC haplotype, including associated log fold
1037   changes (logFC), *p*-values and *q*-values (FDR-corrected *p*-values).

1038
1039

| Gene | Protein | LogFC | p-value | q-value |
|---|---|---|---|---|
| Tldc1 | TBC/LysM associated domain containing 1 | 1.489 | $2.502 \times 10^{-8}$ | $3.228 \times 10^{-4}$ |
| Peg3 | Paternally expressed 3 | 1.440 | $5.369 \times 10^{-5}$ | 0.049 |
| Socs3 | Suppressor of cytokine signalling 3 | 1.071 | $6.061 \times 10^{-5}$ | 0.052 |

1040
1041
1042 **Appendix 1—table 4.** Top 10 annotated genes which were differentially expressed
1043 between males with vs. without the GC haplotype when controlling for cestode burden,
1044 including associated log fold changes (logFC), *p*-values and *q*-values (FDR-corrected *p*-
1045 values).
1046

| Gene | Protein | LogFC | p-value | q-value |
|---|---|---|---|---|
| Il33 | Interleukin 33 | 3.091 | $2.473 \times 10^{-10}$ | $3.191 \times 10^{-6}$ |
| Snai3 | Snail family zinc finger 3 | 1.672 | $5.259 \times 10^{-9}$ | $3.393 \times 10^{-5}$ |
| Pla2g4c | Phospholipase A2, group IVC (cytosolic, calcium-independent) | 2.883 | $1.562 \times 10^{-8}$ | $6.721 \times 10^{-5}$ |
| Uba7 | Ubiquitin-like modifier activating enzyme 7 | 0.809 | $2.477 \times 10^{-8}$ | $7.990 \times 10^{-5}$ |
| Mmp13 | Matrix metallopeptidase 13 | 2.606 | $4.090 \times 10^{-7}$ | $8.796 \times 10^{-4}$ |
| Robo4 | Roundabout guidance receptor 4 | 0.892 | $8.160 \times 10^{-6}$ | $9.572 \times 10^{-3}$ |
| Ttn | Titin | 1.599 | $2.532 \times 10^{-5}$ | $1.951 \times 10^{-2}$ |
| Flnc | Filamin C, gamma | 1.493 | $2.721 \times 10^{-5}$ | $1.951 \times 10^{-2}$ |
| Crb2 | Crumbs family member 2 | 1.261 | $3.512 \times 10^{-5}$ | $2.151 \times 10^{-2}$ |

1047
1048
1049 **Appendix 1—table 5.** Annotated genes which were differentially expressed ($q \le 0.1$)
1050 between females with vs. without the GC haplotype when controlling for cestode burden,
1051 including associated log fold changes (logFC), *p*-values and *q*-values (FDR-corrected *p*-
1052 values).
1053
1054

| Gene | Protein | LogFC | p-value | q-value |
|---|---|---|---|---|
| Tldc1 | TBC/LysM associated domain containing 1 | 1.489 | $2.890 \times 10^{-8}$ | $3.730 \times 10^{-4}$ |
| Socs3 | Suppressor of cytokine signalling 3 | 1.083 | $4.133 \times 10^{-5}$ | 0.041 |
| Peg3 | Paternally expressed 3 | 1.411 | $6.375 \times 10^{-5}$ | 0.055 |
| Ppp1r3c | Protein phosphatase 1, regulatory | 1.662 | $8.798 \times 10^{-5}$ | 0.071 |

| | | | |
|---|---|---|---|
| (inhibitor) subunit 3C | | | |

**Appendix 1—table 6.** Significance values from hapassoc models for expression of 18 genes (assayed by Q-PCR) in splenocytes. Splenocytes were stimulated with anti-CD3 and anti-CD28 antibodies in order to promote the proliferation of T cells. *q*-values (FDR-corrected *p*-values) are reported alongside original *p*-values for the genotype by sex interaction.

| Gene | *p*-value | *q*-value |
|---|---|---|
| Cd4 | 0.124 | 0.822 |
| Cd8a | 0.744 | 0.866 |
| Foxp3 | 0.499 | 0.845 |
| Gata3 | 0.563 | 0.845 |
| Il10 | 0.866 | 0.866 |
| Mpo | 0.173 | 0.822 |
| Tbx21 | 0.650 | 0.866 |
| Tgfb1 | 0.271 | 0.822 |
| Ifng | 0.399 | 0.822 |
| Il17a | 0.002 | 0.037 |
| Il1b | 0.749 | 0.866 |
| Il6 | 0.282 | 0.822 |
| Ms4a1 | 0.281 | 0.822 |
| Orai1 | 0.353 | 0.822 |
| Tnfa | 0.858 | 0.866 |
| Il2 | 0.411 | 0.822 |
| Apobr | 0.857 | 0.866 |
| Arg1 | 0.552 | 0.845 |

**Appendix 1—table 7.** Estimates, standard errors and *z*-statistics from best LMM for Yeo-Johnson-transformed *Il17a* expression levels.

| | Estimate | SE | *z* |
|---|---|---|---|
| (Intercept) | 0.107 | 0.023 | 4.61 |
| Genotype | -0.115 | 0.037 | -3.10 |
| Sex Male | -0.055 | 0.019 | -2.89 |
| Year 2016 | 0.084 | 0.018 | 4.60 |
| Genotype x Sex Male | 0.143 | 0.044 | 3.25 |

**Appendix 1—table 8.** Effect sizes, standard errors, *z*-statistics and associated significance from Gaussian hapassoc model for SOD1 activity.

| | Estimate | SE | *z* | *p*-value |
|---|---|---|---|---|
| (Intercept) | 0.614 | 1.514 | 0.406 | 0.685 |
| hAT | -0.453 | 0.223 | -2.032 | 0.042 |
| Reproductive status Active | -0.112 | 0.266 | -0.422 | 0.673 |
| Sex Male | 0.031 | 0.250 | 0.124 | 0.901 |
| SVL | 0.016 | 0.017 | 0.945 | 0.345 |
| Lens weight 2 | 0.568 | 0.531 | 1.070 | 0.284 |

| | | | | |
|---|---|---|---|---|
| Lens weight 3 | 0.440 | 0.549 | 0.801 | 0.423 |
| Lens weight 4 | 0.445 | 0.587 | 0.758 | 0.448 |
| Lens weight 5 | 0.415 | 0.653 | 0.636 | 0.525 |
| Lens weight 6 | 0.301 | 0.752 | 0.400 | 0.689 |
| Lens weight 7 | -0.307 | 1.095 | -0.280 | 0.779 |
| Body condition | 0.020 | 0.030 | 0.677 | 0.498 |
| Site CHE | 1.429 | 0.431 | 3.316 | 0.001 |
| Site COL | 2.675 | 1.099 | 2.434 | 0.015 |
| Site GRD | 1.476 | 0.437 | 3.379 | 0.001 |
| Site RAV | 0.904 | 0.918 | 0.984 | 0.325 |
| Site SCP | 1.107 | 0.419 | 2.643 | 0.008 |
| Year 2016 | 0.593 | 0.242 | 2.457 | 0.014 |
| Season Early Summer | -0.392 | 0.389 | -1.007 | 0.314 |
| Season Late Summer | -0.098 | 0.266 | -0.369 | 0.712 |
| Season Spring | -0.969 | 0.383 | -2.529 | 0.011 |

1069

**Appendix 1—table 9.** Effect sizes, standard errors and *z*-statistics from best Binomial
GLMM for probability of infection with *Babesia microti.*

| | Estimate | SE | z |
|---|---|---|---|
| (Intercept) | -2.214 | 0.377 | -5.87 |
| Genotype | 1.254 | 0.416 | 3.02 |
| Sex Male | 0.957 | 0.229 | 4.19 |
| Year 2016 | 1.260 | 0.277 | 4.55 |
| Year 2017 | 1.047 | 0.302 | 3.46 |
| Reproductive status Active | 0.754 | 0.144 | 5.24 |
| Body condition | 0.044 | 0.016 | 2.76 |
| Genotype x Sex Male | -1.116 | 0.615 | -1.82 |

1073

**Appendix 1—table 10.** Effect sizes, standard errors and *z*-statistics from best Binomial
GLMM for probability of infection with *Bartonella* spp.

| | Estimate | SE | z |
|---|---|---|---|
| (Intercept) | 0.236 | 0.369 | 0.64 |
| Year 2016 | 0.740 | 0.135 | 5.49 |
| Year 2017 | 2.280 | 0.175 | 13.01 |

1077
1078

**Appendix 1—table 11.** Effect sizes, standard errors, *z*-statistics and associated
significance from Gaussian hapassoc for macroparasite infection summarised by a
single principal component.

| | Estimate | SE | z | *p*-value |
|---|---|---|---|---|
| (Intercept) | 0.050 | 0.331 | 0.152 | 0.879 |
| hAC | -0.102 | 0.130 | -0.786 | 0.432 |
| hAT | -0.108 | 0.158 | -0.682 | 0.495 |
| Sex Male | -0.271 | 0.126 | -2.146 | 0.032 |
| Season Early Summer | -0.103 | 0.182 | -0.567 | 0.571 |

| | | | | |
|---|---|---|---|---|
| Season Late Summer | 0.012 | 0.132 | 0.090 | 0.929 |
| Season Spring | 0.327 | 0.176 | 1.854 | 0.064 |
| Reproductive status Active | -0.222 | 0.119 | -1.868 | 0.062 |
| Body condition | -0.019 | 0.014 | -1.328 | 0.184 |
| Year 2016 | -0.473 | 0.124 | -3.810 | 0.000 |
| Site CHE | 0.714 | 0.226 | 3.157 | 0.002 |
| Site COL | 0.762 | 0.575 | 1.327 | 0.185 |
| Site GRD | 0.707 | 0.229 | 3.090 | 0.002 |
| Site RAV | 1.100 | 0.481 | 2.288 | 0.022 |
| Site SCP | 0.747 | 0.220 | 3.389 | 0.001 |

1083
1084
1085 **Appendix 1—table 12.** Effect sizes, standard errors, *z*-statistics and associated
1086 significance from best quasi-Poisson GLM for reproductive success in males.
1087

| | **Estimate** | **SE** | ***z*** | ***p*-value** |
|---|---|---|---|---|
| (Intercept) | 1.743 | 0.243 | 7.171 | $1.05 \times 10^{-11}$ |
| Genotype | -0.484 | 0.237 | -2.042 | 0.042 |
| Year 2016 | 0.592 | 0.185 | 3.193 | 0.002 |
| Year 2017 | -0.437 | 0.199 | -2.193 | 0.029 |
| Birth month | -0.358 | 0.048 | -7.449 | $1.97 \times 10^{-12}$ |
| Culled Yes | 0.422 | 0.150 | 2.809 | 0.005 |

1088
1089
1090 **Appendix 1—table 13.** Effect sizes, standard errors, *z*-statistics and associated
1091 significance from best quasi-Poisson GLM for reproductive success in females.
1092

| | **Estimate** | **SE** | ***z*** | ***p*-value** |
|---|---|---|---|---|
| (Intercept) | 1.137 | 0.264 | 4.298 | $2.15 \times 10^{-5}$ |
| Year 2016 | 0.520 | 0.211 | 2.466 | 0.014 |
| Year 2017 | -0.118 | 0.215 | -0.549 | 0.584 |
| Birth month | -0.298 | 0.047 | -6.391 | $4.42 \times 10^{-10}$ |

1093
1094
1095 **Appendix 1—table 14.** Effect sizes, standard errors, *z*-statistics and associated
1096 significance from best quasi-Poisson GLM for reproductive success in males, when
1097 controlling for *Babesia microti* and *Bartonella* spp. infection.
1098

| | **Estimate** | **SE** | ***z*** | ***p*-value** |
|---|---|---|---|---|
| (Intercept) | 1.467 | 0.418 | 3.513 | $7.64 \times 10^{-4}$ |
| Genotype | -1.081 | 0.425 | -2.544 | 0.013 |
| Year 2016 | 0.238 | 0.279 | 0.854 | 0.396 |
| Year 2017 | -0.912 | 0.386 | -2.338 | 0.022 |
| Birth month | -0.464 | 0.091 | -5.090 | $2.69 \times 10^{-6}$ |
| Proportion of samples *Babesia*-positive | 0.849 | 0.276 | 3.076 | 0.003 |
| Proportion of samples | 0.997 | 0.440 | 2.268 | 0.026 |

| | | | | |
|---|---|---|---|---|
| *Bartonella*-positive | | | | |

1099

1100 **Appendix 1—table 15.** Effect sizes, standard errors, *z*-statistics and associated
1101 significance from best quasi-Poisson GLM for reproductive success in females, when
1102 controlling for *Babesia microti* and *Bartonella* spp. infection.
1103

| | **Estimate** | **SE** | **z** | **p-value** |
|---|---|---|---|---|
| (Intercept) | 1.483 | 0.305 | 4.867 | $3.20 \times 10^{-6}$ |
| Birth month | -0.310 | 0.073 | -4.229 | $4.38 \times 10^{-5}$ |
| Proportion of samples *Babesia*-positive | 0.569 | 0.247 | 2.308 | 0.0226 |

1104

1105 **Appendix 1—table 16.** Panel of 18 genes for which expression levels in splenocytes
1106 stimulated with anti-CD3 and anti-CD28 antibodies were measured using two-step
1107 reverse transcription quantitative PCR (Q-PCR).

| **Gene** | **Protein** |
|---|---|
| *Cd4* | T-cell surface glycoprotein CD4 |
| *Cd8a* | T-cell surface glycoprotein CD8 alpha chain |
| *Foxp3* | Forkhead box protein P3 |
| *Gata3* | GATA binding protein 3 |
| *Il10* | Interleukin-10 |
| *Mpo* | Myeloperoxidase |
| *Tbx21* | T-box transcription factor TBX21 |
| *Tgfb1* | Transforming growth factor beta 1 |
| *Ifng* | Interferon gamma |
| *Il17a* | Interleukin-17a |
| *Il1b* | Interleukin-1 beta |
| *Il6* | Interleukin-6 |
| *Ms4a1* | B-lymphocyte antigen CD20 |
| *Orai1* | Calcium release-activated calcium channel protein 1 |
| *Tnfa* | Tumor necrosis factor alpha |
| *Il2* | Interleukin-2 |
| *Apobr* | Apolipoprotein B receptor |
| *Arg1* | Arginase-1 |

1108

1109 **Appendix 1—table 17.** Loadings from principal component analysis summarising
1110 infection by macroparasites (ticks, fleas and cestodes)

| **Macroparasite** | **PC1** |
|---|---|
| Cestodes | -0.481 |
| Ticks | -0.592 |
| Fleas | -0.646 |

1111

## (A) IMMUNITY

## (B) INFECTION

## (C) REPRODUCTION