Genomic epidemiology of vancomycin resistant *Enterococcus faecium*

(VR*Efm*) in Latin America: Revisiting the global VRE population structure

Rafael Rios[1], Jinnethe Reyes[1,2], Lina P. Carvajal[1], Sandra Rincon[1], Diana

Panesso[1,2,3], Paul J. Planet[4,5], Aura M. Echeverri[1], An Dinh[2,3], Sergios-Orestis

Kolokotronis [5,6], Apurva Narechania[5], Truc T. Tran[2,3], Jose M. Munita[2,3,7,8], Barbara

E. Murray[2,3,9], Cesar A. Arias[1,2,3,8,9] and Lorena Diaz[1,2,7*]

[1] Molecular Genetics and Antimicrobial Resistance Unit, International Center for

Microbial Genomics, Universidad El Bosque, Bogotá, Colombia.

[2] Center for Antimicrobial Resistance and Microbial Genomics, McGovern Medical

School, University of Texas Health Science Center, Houston, TX, USA.

[3] Division of Infectious Diseases, Department of Internal Medicine, McGovern

Medical School, University of Texas Health Science Center, Houston, Texas, USA

[4] Department of Pediatrics, Perelman School of Medicine, University of

Pennsylvania & Children's Hospital of Philadelphia, Philadelphia, PA, USA

[5] Institute for Comparative Genomics, American Museum of Natural History, New

York, NY, USA

[6] Department of Epidemiology and Biostatistics, School of Public Health, SUNY

Downstate Health Sciences University, Brooklyn, NY, USA

[7] Millennium Initiative for Collaborative Research On Bacterial Resistance

(MICROB-R), Santiago, Chile

[8] Genomics and Resistant Microbes Group, Facultad de Medicina Clinica Alemana,

Universidad del Desarrollo, Santiago, Chile

[9] Department of Microbiology and Molecular Genetics, McGovern Medical School,

University of Texas Health Science Center, Houston, Texas, USA

[*]Correspondence

Lorena Diaz, PhD

Molecular Genetics and Antimicrobial Resistance Unit, International Center for

Microbial Genomics, Universidad El Bosque

Av Cra 9 131a-02, Bogotá, Colombia.

Tel: +57 6489000 ext 1131

Email: diazsandra@unbosque.edu.co

Abstract

The prevalence of vancomycin-resistant *Enterococcus faecium* varies across geographical regions yet little is known about its population structure in Latin America. Here, we provide a complete genomic characterization of 55 representative Latin American VR*Efm* recovered from 1998-2015 in 5 countries. We found that VR*Efm* population in the region is structured into two main clinical clades without geographical clustering. To place our regional findings in context, we reconstructed the global population structure of VR*Efm* by including 285 genomes from 36 countries from 1946-2017. Our results differ from previous studies showing an early branching of animal related isolates and a further split of clinical isolates into two sub-clades, all within clade A. The overall phylogenomic structure was highly dependent on recombination (54% of the genome) and the split between clades A and B is estimated to have occurred more than 3585 years BP. Furthermore, while the branching of animal isolates and clinical clades was predicted to have occur ~894 years BP, our molecular clock calculations suggest that the split within the clinical clade occurred around ~371 years BP. By including isolates from Latin America, we present novel insights into the population structure of VR*Efm* and revisit the evolution of this pathogen.

Introduction

Enterococci are predominantly non-pathogenic gastrointestinal commensal bacteria that occasionally cause human infections. Among them, *Enterococcus faecalis* and *Enterococcus faecium* represent the species that account for most clinically relevant infections. In particular, *E. faecium* has been able to adapt to the hospital environment, emerging during the last few decades as a leading cause of health-care infections worldwide, and becoming the most challenging species to treat[1,2].

Genome plasticity, the presence of multiple antibiotic resistance determinants, and the lack of therapeutic options have contributed to the adaptation of *E. faecium* to hospital environments[3,4]. Moreover, high recombination rates and the acquisition of mobile elements in the genome of *E. faecium* also have driven this evolutionary process[5]. In addition, the enrichment of virulence determinants, such as surface proteins and phosphotransferase systems (particularly PTS[clin], a putative factor found to contribute to the intestinal colonization in a murine model) seems to provide an advantage to the hospital adaptive process[3,6]. Furthermore, functional gene groups, such as those involved in galactosamine metabolism, bile hydrolysis and phosphorus utilization, are also abundant in *E. faecium* clinical strains compared to non-clinical isolates, suggesting that specific metabolic factors have also facilitated adaptation[7].

In terms of antibiotic resistance, one of the most relevant antibiotic resistance traits acquired by enterococci is resistance to vancomycin due to the *van* gene clusters[8]. Furthermore, vancomycin-resistant *E. faecium* (VR*Efm*) frequently exhibits

resistance to ampicillin and high-level resistance to aminoglycosides[9,10]. Indeed,

the World Health Organization (WHO) has categorized VR*Efm* as a priority agent

for which the finding of new and effective therapeutic strategies is imperative[11].

VR*Efm* is widely distributed in hospitals around the world, with the prevalence

varying according to geographical location. In US hospitals, VR*Efm* is an important

clinical pathogen, particularly in immunosuppressed and critically-ill patients[1,12].

The National Health-Care Safety Network described that 82% of *E. faecium*

recovered from bloodstream infections in the US were vancomycin-resistant,

whereas only 9.8% of *E. faecalis* were resistant to vancomycin[12]. In Europe,

prevalence rates of VR*Efm* vary widely by country, but according to the European

Centre for Disease and Control (ECDC) 2016 report, overall prevalence across

European countries was 30%[13]. Although data regarding VR*Efm* in Latin America

are scarce, a few studies have shed some light on the current situation. A

prospective multicentre study focusing on 4 countries in northern South America

(i.e. Colombia, Ecuador, Peru and Venezuela) found an overall prevalence of

VR*Efm* in clinical enterococcal isolates of 31%[14]. More recently, another study

performed in Brazil reported a VR*Efm* prevalence close to 60%[15].

Tracking the population structure of *E. faecium* using conventional bacterial typing

techniques has been challenging[16]. Although wide genetic variability has been

observed among *E. faecium* strains causing clinical infections, a previously

described lineage (designated clonal complex CC17 by multi locus sequence

typing [MLST]), was initially recognized as globally distributed[17]. However, the

classification of this lineage by MLST has some important drawbacks when

analysing the population structure of *E. faecium*, since high rates of recombination

in the MLST loci often occurs in these organisms[18]. Additionally, some strains are

not type able by MLST due to the lack of the locus *pts*[19] leading to major

discrepancies compared to whole-genome sequencing (WGS) when it is used for

typing purposes[20].

Whole-genome-based comparative phylogenomic analyses using *E. faecium*

recovered from different geographical regions have identified two clades,

designated A and B. Clade A mostly contains isolates recovered in clinical settings

(including those from CC17)[21], while clade B encompasses organisms isolated in

community settings, usually from healthy individuals [3,20,22–24]. A further subdivision

has been described within clade A, which groups isolates from animal origin in a

subclade (designated as A2), separating them from those recovered from human

infections or colonization (subclade A1).

However, these analyses have been performed mostly with US and European

isolates, lacking geographical diversity particularly in areas such as Latin America.

Indeed, studies on the molecular epidemiology of VR*Efm* isolates from Latin

America are sparse, with one study suggesting that the CC17 lineage

predominates[14]. Furthermore, studies analysing the population structure of VR*Efm*

in the region using high-resolution, WGS-based phylogenomic comparative

methods are limited. Here, we sought to characterize the population structure of

VR*Efm* lineages in a collection of isolates recovered between 1998-2015 in

prospective multicentre studies performed in selected Latin-American

hospitals[14,25,26] and revisit the global population structure and evolutionary history

of VR*Efm*.

Results

Genomic characterization of Latin American VR*Efm* clinical isolates

From a collection of 207 VR*Efm* clinical isolates obtained between 1998 and 2015

in five Latin American countries (Colombia, Ecuador, Venezuela, Peru and

Mexico), we selected 55 representative isolates for WGS. We included the first

VR*Efm* (ERV1) reported in Colombia as the representative of 23 isolates with

identical PFGE banding pattern, recovered from an outbreak in 1998-1999 and

affecting 23 patients in a single teaching hospital[25]. Five isolates (out of 7

available) were selected from a national surveillance in Colombia during 2001-

2002, which included 15 tertiary hospitals among 5 cities[26] and 16 (out of 35

available) were chosen from a subsequent surveillance study (2006-2008)

performed in Colombia, Ecuador, Venezuela and Peru and the selected isolates

were chosen based on their different banding patterns[14]. The remaining 33 isolates

were obtained from sporadic isolates and outbreaks that occurred in Colombia and

Mexico (2002-2014). In order to characterize the VR*Efm* lineages circulating in

Latin America, we reconstructed their phylogenetic history based on 1,674 genes

(groups of orthologous sequences; hereafter referred to as orthogroups) present in

more than 90% of the genome sequences (core genome) from a total of 6735

orthogroups (pan-genome) using a Bayesian approach (Figure 1A). We observed a

split into two main clades (Clade I and Clade II, marked in red and green,

respectively). Clade I included all the ST412 isolates, while Clade II had all the

ST17 isolates from our sample. We observe that the emergence of VR*Efm* in

Colombia was associated with Clade II, including the first VR*Efm* (described in

1998) and representatives from the first national surveillance (2001 to 2002).

Additionally, ST412 was reported in 2005 and, since then, ST17 and ST412 seem

to be the most prevalent STs in the country. Our previous results showed that Peru

had the highest prevalence of VRE*fm* (48%) and our PFGE and MLST results

suggested higher diversity in Peruvian lineages compared to Colombia, Ecuador

and Venezuela with a predominant circulation of ST412[14]. Indeed, the

representative VR*Efm* isolates of the circulating lineages in Peru collected in the

two-year period (2006-2007) exhibited a marked genomic variability (Figure 1A and

B).

The resistome and virulome of Latin American VR*Efm*

In order to characterize antibiotic resistance determinants, we built resistome

profiles by detecting acquired resistance genes and mutations known to confer

resistance to linezolid, ciprofloxacin and daptomycin. All the VR*Efm* isolates from

our collection were resistant to vancomycin (MIC_{90} >256 μg/ml) and teicoplanin

(MIC_{90} 64 μg/ml) (Figure 1B). The presence of *vanA* was confirmed in all isolates

by PCR assays. Consistently, we confirmed the presence of the entire *vanA* cluster

in 54 out of the 55 sequenced genomes. Of note, the genome of ERV69 lacked the

two-component regulatory system *vanSR*, although still exhibiting MICs of >256

µg/ml and 64 µg/ml for vancomycin and teicoplanin, respectively. The deletion of

the two-component regulatory system has been previously reported[27].

High-level resistance to ampicillin was consistently found in all 55 *E. faecium*

isolates, a phenotype that was corroborated using comparisons of the PBP5

protein sequence using a machine-learning prediction model. This approach was

based on the amino acid changes present in the PBP5 protein across susceptible

and resistant isolates (see details in Methods).

High-level resistance to gentamicin was identified in 31% of the isolates of our

collection and, within the sequenced representatives, the presence of *aac(6´)-*

aph(2") was detected in 49% of the genome sequences. High-level resistance to

streptomycin was identified in 39% of the Latin American VRE*fm* isolates with a

high prevalence of the *ant(6)-Ia* gene (89%; n=49) in the sequenced genomes.

Fluoroquinolone resistance is very common in *E. faecium*. All isolates in our

collection were fluoroquinolone-resistant and we were able to predict the presence

of substitutions in GyrA and ParC associated with this phenotype. The most

common substitution in GyrA was Ser84Arg (67%; n=37). All isolates exhibited

Ser82Arg (53%; n=29) or Ser82Ile (47%; n=26) substitutions in ParC.

The *cat* gene conferring resistance to chloramphenicol was present only in the

three Peruvian genomes. Interestingly, Peruvian isolates had the highest

resistance to this antibiotic (21%). All the isolates from this collection were

susceptible to linezolid; however, we detected the various genetic elements

previously associated with linezolid resistance. The gene, *optrA*, was detected in

one genome of a Colombian linezolid-susceptible isolate (ERV138). Also, we

identified the presence of *cfrB,* a recently reported variant of *cfr*[28], in a Mexican

isolate (ERV275). We predicted tetracycline resistance owing to the presence of

tetM (43.6%; n=24), *tetL* (16.3%; n=9) and *tetS* (1.8%; n=1) in the sequenced

genomes, but resistance to this group of antibiotics was not tested phenotypically.

Substitutions in LiaS (Thr120Ala) and LiaR (Trp73Cys), which have been strongly

associated with daptomycin resistance and tolerance[29,30], were present in three

VR*Efm* isolates, recovered before daptomycin was available in the region. Of note,

the three isolates showed MICs between 2-4 µg/ml, considered now as

"daptomycin-susceptible dose-dependent", by the Clinical & Laboratory Standards

Institute (CLSI)[31].

Latin American VRE isolates also harboured a high proportion of putative virulence

determinants (Figure 2). The vast majority had gene clusters related to pilus

formation, adhesins and microbial surface components recognizing adhesive

matrix molecules (MSCRAMMS). Interestingly, the notable exception was the

Clade I isolates, which often lacked *fms22*, *swpC* and *hyl*$_{Efm}$. Our results suggest

that the "virulome" of Latin-American VRE is similar to those from other regions in

the world[32].

Global Phylogenetic Reconstructions of Latin American VRE

To place the genetic lineages of VR*Efm* isolates circulating in Latin America into a

global context, we performed a WGS-based phylogenomic analysis. We included

285 *E. faecium* genomes (VRE and non-VRE) from the publicly available NCBI

collection aiming to incorporate a diverse set of sequences for comparisons. The

included isolates were from colonizing, commensal, animal and clinical sources

and were collected between 1946-2017 from Europe, North America, Asia, Africa

and Australia (Supplementary Table 1). We constructed a pangenome (29,503

orthogroups) and core genome (978 orthogroups). Using the core genome, we built

a phylogenomic tree of the species to show the evolutionary relationships among

isolates based on the variation of their genomic sequences. Figure 3 shows that,

as previously reported, we found a clear split into two main clades corresponding

to the previously designated clades A and B[3,22,24]. All Latin American isolates from

our clinical collection were in clade A. We compared the genomic characteristics

among the two main clades and found similar findings as published previously

(Supplementary table 2)[3]. However, our data showed that the core genome was

larger in clade B as compared to clade A (1,466 vs 1,182 orthogroups,

respectively).

Considering the relevance of *E. faecium* as a cause of hospital-associated

infections and that all Latin American isolates were grouped within clade A, we

sought to dissect the population structure of this clade when adding the genomes

of these isolates. Our first approach was based on a core genome (>90%

reconstruction), which contained 1,226 orthogroups and the isolate Com15, from

clade B, as the outgroup to root the tree. We observed two major subclades. The

first was composed of 52 genomes, most of which were from animal sources (57%,

n=30), related to the previously described subclade A2[3]. The second lineage

harboured 273 genomes, with 91% (n=228) corresponding to isolates obtained

from clinical sources (Supp. Figure 2A), and related to the subclade A1[3].

Previous studies have shown contradictory distributions of the subclades A1 and

A2 within clade A[20]; suggesting that clade A2 is not in fact a clade, but rather

corresponds to the paraphyletic early branching lineages of clade A. To further

clarify the issue, we performed a phylogenomic analysis accounting for

recombination events within clade A. We used the variants found from paired

alignments of each genome against the chromosome of reference Aus0085 and

built a whole-genome multiple sequence alignment (WGMSA) of all genomes in the

clade. We used this alignment to create a maximum likelihood tree, which is

required for determining recombinant regions using ClonalFrameML[33]. The

average amount of recombination found in the 303 genomes belonging to clade A

was 19,539pb (Supp. figure 2C). The total recombinant regions found across

clinical isolates encompassed 1.6 Mb (54% of the length of WGMSA). Interestingly,

the exclusion of recombinant regions considerably altered the structure of the tree,

and showed 7 early-branching subclades that included 73 genomes (mostly from

animal sources) rather than a split into clades A1 and A2.

Following these animal-related early branches, we observed a split into two main

subclades (Supp. Figure 2B). Overall, these subclades were related to clinical

sources, exhibiting a high similarity in terms of prevalence of antibiotic resistance

and virulence determinants (Supplementary table 3). We refer to them as clinically-

related clades I and II (CRS-I and CRS-II), containing 101 and 124 genomes

respectively. Latin American genomes from our collection were split between these

two CRS, showing that Clade I and Clade II (derived from the analysis of Latin

American VR*Efm,* see above) belonged to CRS-I and CRS-II, respectively. Of

note, the genomes from our collection were distributed almost equally between

CRS-I (49%) and CRS-II (51%). Furthermore, despite the inclusion of a few

outbreak isolates and that VR*Efm* from Latin America originated in different

periods, cities and countries, our phylogenetic reconstruction showed 11

conserved clusters with four or more isolates from the same country (Figure 4). In

particular, three clusters had only Colombian genomes with the number of SNPs

among them, within the regions not showing recombination, ranging between 36

and 160. We also found clusters among isolates from Brazil (n=3), USA (n=3),

Denmark (n=1) and Sweden (n=1). The Danish cluster is situated in the animal-

associated branches, and their genomes were closely related (with an average of

43 SNPs among them). Of note, two of the USA clusters were related to each other

and to 5 other isolates, four of them from the UK and one from Colombia in our

collection (172 SNPs difference on average among them).

In CRS-I, there were 23 different STs, with ST412 and 78 the most frequent (34%

and 11%, respectively) (Figure 4). Importantly, we did not find a strong correlation

between MLST and the phylogenomic analysis, as isolates belonging to the same

ST were not all clustered in the same clades, and were distributed in different

groups in the phylogeny. In particular, 56% (n=9) of genomes from ST78 were in

CRS-I, while 37% (n=6) were in CRS-II. To further dissect this discrepancy, we

performed a phylogenetic reconstruction using only the sequences of the 7 MLST

loci and compared it against the phylogeny of Clade A. Our results showed that

many isolates from ST17, ST18, ST78, ST203, ST412 were in different clusters

and even formed subclades in the non-recombination reconstruction

(Supplementary Figure 3).

In relation to antibiotic resistance determinants, we found important differences

comparing the presence/absence of genomic elements associated with antibiotic

resistance between the CRSs and the animal branches. Indeed, the animal-

associated branches exhibited a lower prevalence of elements associated with

glycopeptide (34.2%), aminoglycoside (21.9%), ampicillin (9.5%) and

fluoroquinolone resistance (2.7%) compared to the CRS isolates, which harboured

these determinants in 78%, 85%, 95% and 99% of isolates, respectively. In

contrast, similar frequencies of determinants coding for resistance to macrolides

(>98%), tetracyclines (between 50-63%) and oxazolidinones (between 2-12%)

were found between animal and clinical clades (Supp. Table 3). Within the

subclades of clade A, only 9% of isolates within the animal-associated branches

exhibited resistance to ampicillin (7 out of 71 complete PBP5 sequences), while

99% of the clinically related subclades (100% in CRS-I and 98% in CRS-II)

harboured the predicted *pbp5-R* allele[34,35]. Mutations associated with

fluoroquinolone resistance were also much more highly prevalent in clinical clades

(>98% for CRSs) vs animal branches (2.7%; p<0.001).

Genes encoding putative surface adhesin proteins (e.g., *acm, scm, esp, sgrA,*

fms6 and *fms22*) and two of the pilus-forming clusters were significantly more

common in the CRSs, (p-values below 0.001 in all cases) compared to animal

isolates (Supp. Table 3). We next compared the presence/absence of putative

mobile elements between animal branches vs. CRSs. On average, the number of

insertion sequences in the former were 5.7, whereas the clinical subclades had 6.9

(6.76 CRS-I and 7.06 for CRS-II). Of note, *rep17* was notoriously overrepresented

in the CRSs (Supp. Table 3), located in the plasmid pRUM, which is a

representative member of rep17 family and has been associated with the

toxin/antitoxin system Txe/Axe[36].

Rates of evolution across the whole population of *E. faecium*

Using the sampling date of isolates within clade A, we performed molecular clock

analyses on the entire clade A and its subgroups (animal branches, CRS-I and

CRS-II). We found that the oldest split within clade A likely occurred ~3,585 years

ago (y.a.) (95% High Posterior Density Interval [HPDI]: [2626, 4690]). The

separation of the clinical subclades from the animal branches is predicted to have

occurred ~894 y.a. (95% HPDI: [649, 1171]) (Supplementary Figure 3). The most

recent split between CRS-I and CRS-II was dated ~371 y. a (95% HPDI: [272,

488]) (Supplementary Figure 4). The substitution rate across the clade A genomes

was 3.91E-7 (95% HPDI: [2.78E-7, 5E-7]), which translates to 0.53 SNPs per year

(using only non-recombinant regions or 1.17 SNPs if the WGSA is used). The

substitution rates within each subgroup of genomes were 3.02E-7 (95% HPDI:

[2,78E-7, 3,46E-7]) for animal branches, 4.7E-7 (95% HPDI: [4,01E-7, 4,98E-7]) for

CRS-I and 4.63E-7 (95% HPDI: [3.92E-7, 4.98E-7]) for CRS-II. These rates are

equivalent to 0.41, 0.64 and 0.63 SNPs per year for animal branches, CRS-I and

CRS-II, respectively. Our results support that clinically related clades are evolving

faster than those of the animal branches.

Discussion

Our results indicate that VR*Efm* is widely present in Latin America but that their

frequency and population structure seem to vary from country to country. As

multicentre analyses of VR*Efm* in the Latin American region are rare, our study is

unique in its dissection of the population structure of VR*Efm* in the region. Unlike

previous studies, we found two distinct populations of clinically-related isolates of

VR*Efm*. This subpopulation separation was also seen in our analyses of the global

population of *E. faecium*. The causes for the splitting of the population structure of

VRE (CRS-I and CRS-II) are not clear, but the findings were consistent when

analysing the population structure in the presence or absence of recombinant

regions. Such a separation suggests that these lineages have been expanding

through Latin American countries and highlights the importance of establishing

genomic surveillance studies for these multidrug-resistant organisms. Furthermore,

the distribution of the Latin American isolates across the tree does not suggest a

particular dominance of a specific lineage circulating in the region or country,

suggesting that the presence of VR*Efm* in Latin America is associated with multiple

introductions of VR*Efm* lineages that are circulating globally. Interestingly, some

South American countries such as Brazil (no isolates available for this study) have

reported VR*Efm* since 1997[37], and their prevalence appears to be increasing

exhibiting a shift from *E. faecalis* to VR*Efm* since 2007[15]. Of interest, ST412

isolates reported in some regions of Brazil[38,39] have also been detected in some

Caribbean countries[40] and this sequence type was also identified in our collection

in Colombia, Peru and Venezuela since 2005[14], suggesting wide dissemination of

this genetic lineage.

Our VR*Efm* phylogenomic analysis, which includes a highly diverse sample

collection and excludes recombinant regions from the genome, questions the

presence of a single animal clade. Our results suggest that the animal isolates

represent multiple lineages that diverged prior to the emergence of the clinical

subclades in the clade A[3]. Importantly, animal-associated branches have

significantly lower predicted ampicillin resistance, fluoroquinolone resistance

mutations, virulence elements and average number of insertion sequences,

similarly to what has previously reported[41]. Furthermore, the amount of

recombination in clade A genomes was greater than previous results. Importantly,

this difference (54% vs 44% found in previous studies[18,42]) could be due to the fact

that previous analyses were based on the alignment of SNPs from a core genome

and neither included non-coding regions nor invariant sites to identify the

recombinant DNA. Over the recombinant regions, we found partial sequences in 5

out of the 7 loci used by MLST (*ddl, gyd, purK, gdh* and *adk*), corroborating the

notion that the current *E. faecium* MLST scheme has major limitations to describe

the population structure of VR*Efm*. Interestingly, the exclusion of recombinant

regions considerably altered the structure of the tree, dissolving the animal-related

clade into a paraphyletic group and reducing the length of the branches across the

tree (Supplementary Figure 2). Additionally, we found a lack of concordance

between MLST classification and the clades. The discrepancy is likely explained by

the presence of recombinant regions in the MLST genes, low variation in some of

the loci, and the absence of *pst* in many isolates[19,20,43]

Previous studies estimated that the separation between clades A and B occurred

2776 ± 818 y.a.[3], a time frame that is similar to our results. However, the previously

reported split between animal branches and the clinically-related subclades was

reported to occur 74 ± 30 y. a., which is much more recent than what we found.

Our findings showed at least a tenfold lower mutation rate from what has been

previously reported[3,18]. This finding could be associated with the larger genomic

region used in our analysis and the increase in the diversity of the sampled

genomes. Indeed, dating of the splits between the animal-associated branches and

the clinically-related subclades, and the lower mutation rates across clade A

correlates with lower number of SNPs per year. It has been estimated that the

Enterococcus as a genus arose around 500 million years ago[44] and ancient

isolates of *E. faecium* have been found in permafrost over 20,000 y.a.[45], supporting

our findings that a more ancient branching between Clade A and B could have

occurred.

Our study could be subject to sampling bias due to small sample size of genomes

from Latin America, but we attempted to include as many and as diverse genomes

as possible from our collection, based on phenotypic characteristics and PFGE

typing of the strains. Also, we included all publicly available genomes from the

region, provided that the associated demographic information was complete

(source, year of sampling and geographical location), which also reiterates the low

number of previously sequenced genomes of *E. faecium* in Latin America at the

moment of sample selection. Nonetheless, our results supporting the existence of

two clinical subclades were maintained even after the inclusion of genomes from

other continents; that is, our conclusion holds beyond sample size, further

indicating that the population structure of the clinical related isolates is divided into

two main lineages within clade A.

Conclusions

We provide comprehensive insights into the genomic epidemiology of VR*Efm* using

available isolates from Latin America where previous studies are lacking. Our

results indicate that the population structure of VR*Efm* in the region is diverse and

can be grouped into two main lineages (Clades I and II) that belong to the

previously reported clade A. A novel global reconstruction of the *E. faecium*, using

a wide and diverse sample of isolates from 36 countries and obtained from clinical,

animal, environmental and commensal samples, corroborates previous reports that

recombination plays a major role in the evolution of this species. Our analyses also

indicate, contrary to previous results, that animal-associated genomes are not

monophyletic, and are instead a diverse collection of early-branching clades that

diverged prior to the emergence of the human clinical clade and its two subclades

(CRSI and CRSII).

The complex evolutionary dynamics of VR*Efm* highlight the importance of

employing phylogenomic approaches when studying the population structure of a

highly evolved hospital-associated pathogen.

Acknowledgments

This work was founded by Universidad El Bosque, grant PCI 2016-8865 to L.D.;

grants from the National Institutes of Health K24-AI121296 and R01-AI134637 to

C.A.A. and grant FONDECYT regular Project No. 1171805, Ministry of Education,

Government of Chile and the Millennium Science Initiative, Ministry of Economy,

Development and Tourism, Chile to J.M.M.

Methods

***Enterococcus faecium* isolates**

A total of 207 vancomycin-resistant Latin American *E. faecium* clinical isolates

have been collected between 1998 and 2014 including those belonging to the first

outbreak of VRE infections in Colombia and isolates collected in two multicentre

surveillances[14,25,26]. Isolates were recovered from patients in Colombia (n=177,

86%) Peru (n=14, 7%), Venezuela (n=6, 3%), Ecuador (n=5, 2%) and Mexico (n=5,

2%). The most common sources included blood (22%), urine (18%) and stools

(10%). For all the isolates, species (*E. faecium*) confirmation and the susceptibility

profiles determination were performed by PCR assays[46] and agar dilution,

respectively[31].

Whole genome sequencing

From our VR*Efm* characterized strains collection, we selected 55 representative

isolates based on distinct PFGE banding patterns. We included the first VRE

reported in Colombia as the representative of an outbreak of 23 infections at a

teaching hospital in 1998-1999[25]. Five isolates were selected from a national

surveillance in Colombia during 2001-2002, which included 15 tertiary hospitals in

5 cities[26] and 16 chosen from surveillance performed in Colombia, Ecuador,

Venezuela and Peru in 2006-2008[14]. The remaining 33 isolates were sent to our

lab for the confirmation of resistance or outbreak studies in 2005-2014. All selected

isolates were recovered from clinical samples including blood (32%), urine (13%),

faeces (13%), surgical wound (10%), pleural liquid (5%), peritoneal liquid (5%) and

other sources (22%). The isolates were subjected to whole genome sequencing on

the Illumina platform. Briefly, genomic DNA was extracted from overnight cultures

using the kit DNeasy Blood & Tissue Kit (Qiagen) after a lysozyme treatment. DNA

libraries were prepared using Nextera XT kit (illumina) and sequenced on a MiSeq

instrument using a 300pb paired-end strategy. The obtained paired-end reads were

trimmed for quality and used for assemblies using SPAdes[47].

Global *E. faecium* genomic characterization

To place the population structure of Latin American VR*Efm* into global context, we

included 285 *E. faecium* genomes from the publicly available collection available at

NCBI. We aimed to incorporate a diverse set of sequences, including colonizing,

commensal, animal and clinical sources recovered between 1946 and 2017 in

Europe, North America, Asia, Africa, and Australia (Supplementary Table 1).

Accordingly to the source, the *E. faecium* genomes were grouped into different

categories: **i**) isolates from stools or rectal swabs of hospitalized patients (n=59), **ii**)

organisms from hospitalized patients (n=196), recovered from sources other than

faeces, including blood (n=113), urine (n=18) and other sources (n=65), **iii**) stools

from healthy individuals not in hospital settings (n=13), **iv**) animal isolates (n=47),

obtained from different animals, including pets, wild and farm animals, and **v**)

"others" (n=25), which included isolates recovered from food products, water, soil,

among other non-human and non-animal sources.

All sequences (340 *E. faecium* genomes) were annotated using RAST[48]. The

sequence type (ST) was determined by MLST tools

(https://github.com/tseemann/mlst) and verified against PubMLST[49]. Genomic

characterization was performed to identify genetic elements associated with

resistance using BLASTX[50] searches against the ResFinder database[51].

Additionally, we specifically interrogated the genomes for substitutions in GyrAB

and ParCE proteins associated to fluoroquinolone resistance, and mutations in

genes encoding 23S rRNA and L3 and L4 proteins associated with linezolid

resistance. Detection of mobile elements was performed with BlastN[50]. Search for

rep families genes[52,53] and insertion sequences (IS) was carried out with BLASTN

searches and compared to the ISFinder database[54]. Identification of virulence

elements was performed with BLASTX against a set of potential virulence proteins

in enterococci[4,55]. Identification of CRISPR and *cas*-systems was done using

CRISPRfinder[56] and BLASTX searches using Cas system proteins[57] as templates.

All BLASTX hits were selected if they had an identity percentage higher or equal to

95% and a coverage of at least 80% of the target sequence. For BLASTN

searches, hits were selected if they had an identity percentage higher than 90%

and a coverage of at least 80% of the target sequence. To identify statistically

significant differences across proportions of the evaluated characteristics among

pairs of clades found, a Z-test was performed (α=0.01).

499   Ampicillin resistance prediction based on PBP5 sequences

The ampicillin resistance prediction model for *E. faecium* isolates consisted on a

random forest built upon a dataset of 250 PBP5 sequences from isolates with

known MIC of ampicillin (62 from susceptible isolates [MIC$\leq$8 µg/ml] and 188

belonging to resistant ones [MIC $\geq$ 16 µg/ml][Supplementary Table 4]). The model

was based on a multiple sequence alignment using the sequence of the PBP5 from

Com15 (GenBank accession: WP_002314979.1) isolate as reference (based on

previous studies of correlation of the amino acid sequence of this protein with the

resistant phenotype[34,35]) with 110 positions harbouring amino acid changes

(Supplementary Table 4). These positions were used to create a random forest

model with 100 decision trees; using a training set of 42 isolates (17 susceptible

and 25 resistant with a range of MIC values). Based on this training set, forty

amino acid changes were selected for the classification based on their

discrimination power using recursive elimination process of those with lower score.

Next, the model was tested on the whole dataset of PBP5 sequences and had a

100% specificity with 96% sensitivity, which resulted in 6 cases of major errors

were the isolate was resistant but predicted to be susceptible.

Phylogenetic analysis

We built a phylogenetic tree based on the core genome of 55 representative

genomes from our collection, including the genome Com15 as outgroup. The core

genome was obtained with Roary[58] and each of the orthogroups was aligned with

MUSCLE v3.8[59]. A Maximum Likelihood (ML) guide tree was built with RAxML

8.2.11[60] using a GTR+Γ model. Using Bayesian approach, we estimated a

Maximum Clade Credibility (MCC) tree based on 20 million trees in BEASTv1.8[61].

We employed a constant population size, a GTR+Γ+I substitution model, default

prior probability distributions, and a chain length of 100 million steps with a burn-in

of 10 million and a 5000-step thinning and the ML as starting tree.

The phylogenetic tree for the whole population of *E. faecium* included all the

genomes (n=340) and two outgroups (*Enterococcus durans* BDGP3 [GenBank

accession: CP022930.1] and *Enterococcus hirae* ATCC 9790 [CP003504.1]). This

tree was based on the core genome (genes present in at least 90% of the studied

genomes) obtained with Roary, each orthogroup was individually aligned with

MUSCLE and then concatenated to obtain a matrix. The alignment matrix was

used for Bayesian phylogenetic reconstruction with BEAST. Model parameters

were the same as above with a chain length of 300 million steps, a burn-in of 80

million steps, and a random starting tree.

The second phylogenetic reconstruction included the genomes grouped into the

clade corresponding to the previously designed Clade A[3]. We realized pairwise

comparisons of the assemblies with Mummer 3.23[62] against the reference genome

Aus0085 (CP006620.1). The identified variants and the reference sequence were

used to create a multiple whole genome alignment and, with it, we built a guide

tree with RAxML[60] using the abovementioned parameters. This guide tree was

used later to obtain the recombinant regions in the alignment with

ClonalFrameML[33] for each isolate. Those regions were further removed from the

alignment and then used to produce a MCC tree with BEAST. The same run

parameters as above were used with a 50-million step burn-in.

Finally, a strict molecular clock analysis was performed on clade A strains. We

dated the tips on the isolates accordingly to the sampling year. The analysis was

done with the non-recombinant regions of the whole genome alignment as matrix

and the MCC from the second analysis, as a guide tree. The analysis had a 300

million length chain and a burn-in of 30 million to obtain ESS numbers above 200.

All MCC trees were computed with a 0.3 posterior clade probability cut-off and

mean heights. To estimate the evolution rates across subclades, further

subgrouping of the isolates was performed and a similar molecular clock analysis

without guide tree were performed for each group using 100 million chain length

and 10% burn in. All BEAST runs were performed on the CIPRES Science

gateway servers[63].

Data Availability

559   All genomic data is available at GenBank database, accession numbers for the

560   sequenced genomes are listed in Supplementary Table 3. The datasets generated

561   during and/or analysed during the current study are available from the

562   corresponding author on reasonable request.

Ethics declarations

We declare no ethical competing interest. In our study, we did not perform any

experiments with animals or higher invertebrates, neither performed experiments

on humans and/or the use of human tissue samples. Our data have been

originated from bacteria, not linked to clinical information, collected in previous

studies and following full ethical approvals. Also, additional genomic data that we

included for the analysis are available on public repositories (NCBI and published

articles).

References

1.   Arias, C. a. & Murray, B. E. The rise of the Enterococcus: beyond

vancomycin resistance. *Nat. Rev. Microbiol.* **10,** 266–278 (2012).

2.   Cattoir, V. & Giard, J.-C. Antibiotic resistance in *Enterococcus faecium*

clinical isolates. *Expert Rev. Anti. Infect. Ther.* **12,** 239–248 (2014).

3.   Lebreton, F. *et al.* Emergence of epidemic multidrug-resistant Enterococcus

faecium from animal and commensal strains. *MBio* **4,** 1–10 (2013).

4.   Gao, W., Howden, B. P. & Stinear, T. P. Evolution of virulence in

Enterococcus faecium, a hospital-adapted opportunistic pathogen. *Curr.*

Opin. Microbiol. **41,** 76–82 (2018).

5.   van Hal, S. J. *et al.* Evolutionary dynamics of Enterococcus faecium reveals

complex genomic relationships between isolates with independent

emergence of vancomycin resistance. *Microb. genomics* **2,** (2016).

6.   Zhang, X. *et al.* Identification of a Genetic Determinant in Clinical

Enterococcus faecium Strains That Contributes to Intestinal Colonization

During Antibiotic Treatment. *J. Infect. Dis.* **207,** 1780–1786 (2013).

7.   Kim, E. B. & Marco, M. L. Nonclinical and Clinical Enterococcus faecium

Strains, but Not Enterococcus faecalis Strains, Have Distinct Structural and

Functional Genomic Features. *Appl. Environ. Microbiol.* **80,** 154–165 (2014).

8.   Courvalin, P. Vancomycin Resistance in Gram-Positive Cocci. *Clin. Infect.*

Dis. **42,** 25–34 (2006).

9.   Rubinstein, E. & Keynan, Y. Vancomycin-resistant enterococci. *Crit. Care*

Clin. **29,** 841–852 (2013).

10.   O'Driscoll, T. & Crank, C. W. Vancomycin-resistant enterococcal infections:

Epidemiology, clinical manifestations, and optimal management. *Infect. Drug*

Resist. **8,** 217–230 (2015).

11.   Tacconelli, E. *et al.* Discovery, research, and development of new antibiotics:

the WHO priority list of antibiotic-resistant bacteria and tuberculosis. *Lancet*

Infect. Dis. **18,** (2018).

12. Weiner, L. M. *et al.* Antimicrobial-Resistant Pathogens Associated with

Healthcare-Associated Infections: Summary of Data Reported to the National

Healthcare Safety Network at the Centers for Disease Control and

Prevention, 2011-2014. *Infect. Control Hosp. Epidemiol.* **37,** 1288–1301

(2016).

13. European Centre for Disease Prevention and Control. *Surveillance of*

antimicrobial resistance in Europe 2016. Annual report of the European

Antimicrobial REsistance Surveillance Network (EARS-Net). (2017).

doi:10.2900/296939

14. Panesso, D. *et al.* Molecular epidemiology of vancomycin-resistant

Enterococcus faecium: A prospective, multicenter study in South American

hospitals. *J. Clin. Microbiol.* **48,** 1562–1569 (2010).

15. Sacramento, A. G. *et al.* Changed epidemiology during intra and interhospital

spread of high-risk clones of vanA-containing Enterococcus in Brazilian

hospitals. *Diagn. Microbiol. Infect. Dis.* **88,** 348–351 (2017).

16. van Schaik, W. & Willems, R. J. L. Genome-based insights into the evolution

of enterococci. *Clin. Microbiol. Infect.* **16,** 527–532 (2010).

17. Top, J., Willems, R. & Bonten, M. Emergence of CC17 *Enterococcus*

faecium : from commensal to hospital-adapted pathogen. *FEMS Immunol.*

Med. Microbiol. **52,** 297–308 (2008).

18.   Howden, B. P. *et al.* Genomic Insights to Control the Emergence of

Vancomycin-Resistant Enterococci. *MBio* **4,** 1–9 (2013).

19.   Carter, G. P. *et al.* Emergence of endemic MLST non-typeable vancomycin-

resistant Enterococcus faecium. *J. Antimicrob. Chemother.* **71,** 3367–3371

(2016).

20.   Raven, K. E. *et al.* A decade of genomic history for healthcare-associated

Enterococcus faecium in the United Kingdom and Ireland. *Genome Res.* **26,**

1388–1396 (2016).

21.   Leavis, H. L., Bonten, M. J. & Willems, R. J. Identification of high-risk

enterococcal clonal complexes: global dispersion and antibiotic resistance.

Curr. Opin. Microbiol. **9,** 454–460 (2006).

22.   Galloway-Peña, J., Roh, J. H., Latorre, M., Qin, X. & Murray, B. E. Genomic

and SNP analyses demonstrate a distant separation of the hospital and

community-associated clades of enterococcus faecium. *PLoS One* **7,** (2012).

23.   van Schaik, W. *et al.* Pyrosequencing-based comparative genome analysis

of the nosocomial pathogen Enterococcus faecium and identification of a

large transferable pathogenicity island. *BMC Genomics* **11,** 239 (2010).

24.   Palmer, K. L., Schaik, W. Van, Willems, R. J. L. & Gilmore, M. S.

Enterococcal Genomics. *E-Book* (2014).

25.   Panesso, D. *et al.* First Characterization of a Cluster of VanA-Type

Colombia. *Emerg. Infect. Dis.* **8,** 961–965 (2002).

26.   Arias, C. A. *et al.* Multicentre surveillance of antimicrobial resistance in

enterococci and staphylococci from Colombian hospitals, 2001-2002. *J.*

Antimicrob. Chemother. **51,** 59–68 (2003).

27.   Sung, K., Khan, S. A. & Nawaz, M. S. Genetic diversity of Tn 1546 -like

elements in clinical isolates of vancomycin-resistant enterococci. **31,** 549–

554 (2008).

28.   Deshpande, L. M. *et al.* Detection of a New cfr -Like Gene , cfr(B), in

Enterococcus faecium Isolates Recovered from Human Specimens in the

United States as Part of the SENTRY Antimicrobial Surveillance Program.

Antimicrob. Agents Chemother. **59,** 6256–6261 (2015).

29.   Diaz, L. *et al.* Whole-genome analyses of Enterococcus faecium isolates with

diverse daptomycin MICs. *Antimicrob. Agents Chemother.* **58,** 4527–4534

(2014).

30.   Munita, J. M. *et al.* Correlation between mutations in liaFSR of Enterococcus

faecium and MIC of daptomycin: revisiting daptomycin breakpoints.

Antimicrob. Agents Chemother. **56,** 4354–9 (2012).

31.   Clinical Laboratory Standards Institute. *M100. Performance Standards for*

Antimicrobial Susceptibility Testing, 29th Edition. (2019).

32.   Freitas, A. R., Tedim, A. P., Novais, C., Coque, T. M. & Peixe, L. Distribution

of putative virulence markers in Enterococcus faecium: towards a safety

profile review. *J. Antimicrob. Chemother.* 1–14 (2017).

doi:10.1093/jac/dkx387

33. Didelot, X. & Wilson, D. J. ClonalFrameML: Efficient Inference of

Recombination in Whole Bacterial Genomes. *PLoS Comput. Biol.* **11,** 1–18

(2015).

34. Pietta, E., Montealegre, M. C., Roh, J. H., Cocconcelli, P. S. & Murray, B. E.

Enterococcus faecium PBP5-S/R, the Missing Link between PBP5-S and

PBP5-R. *Antimicrob. Agents Chemother.* **58,** 6978–6981 (2014).

35. Galloway-Peña, J. R., Rice, L. B. & Murray, B. E. Analysis of PBP5 of early

U.S. isolates of Enterococcus faecium: Sequence variation alone does not

explain increasing ampicillin resistance over time. *Antimicrob. Agents*

Chemother. **55,** 3272–3277 (2011).

36. Grady, R. & Hayes, F. Axe-Txe, a broad-spectrum proteic toxin-antitoxin

system specified by a multidrug-resistant, clinical isolate of Enterococcus

faecium. *Mol. Microbiol.* **47,** 1419–1432 (2003).

37. Zanella, R. C. *et al.* First Confirmed Case of a Vancomycin-Resistant

Enterococcus faecium with vanA Phenotype from Brazil: Isolation from a

Meningitis Case in São Paulo. *Microb. Drug Resist.* **5,** 159–162 (2009).

38. Alves, G. da S., Pereira, M. F., Bride, L. de L., Nunes, A. P. F. & Schuenck,

R. P. Clonal dissemination of vancomycin-resistant Enterococcus faecium

ST412 in a Brazilian region. *Brazilian J. Infect. Dis.* **21,** 656–659 (2017).

39. da Silva, L. P. P., Pitondo-Silva, A., Martinez, R. & da Costa Darini, A. L.

Genetic features and molecular epidemiology of Enterococcus faecium

isolated in two university hospitals in Brazil. *Diagn. Microbiol. Infect. Dis.* **74,**

267–271 (2012).

40.   Akpaka, P. E. *et al.* Genetic characteristics and molecular epidemiology of

vancomycin-resistant Enterococci isolates from Caribbean countries. *PLoS*

One **12,** 1–11 (2017).

41.   Torres, C. *et al.* Antimicrobial Resistance in Enterococcus spp . of animal

origin. *Microbiol. Spectr.* **6,** (2018).

42.   De Been, M., Van Schaik, W., Cheng, L., Corander, J. & Willems, R. J.

Recent recombination events in the core genome are associated with

adaptive evolution in Enterococcus faecium. *Genome Biol. Evol.* **5,** 1524–

1535 (2013).

43.   Been, M. De *et al.* Core Genome Multilocus Sequence Typing Scheme for

High- Resolution Typing of Enterococcus faecium. *J. Clin. Microbiol.* **53,**

3788–3797 (2015).

44.   Lebreton, F. *et al.* Tracing the Enterococci from Paleozoic Origins to the

Hospital. *Cell* 1–13 (2017). doi:10.1016/j.cell.2017.04.027

45.   Goncharov, A. *et al.* Draft Genome Sequence of Enterococcus faecium

Strain 58m , Isolated from Intestinal Tract Content of a Woolly Mammoth ,

Mammuthus primigenius. *Genome Announc.* **4,** 15–16 (2016).

46.   Dutka-Malen, S., Evers, S. & Courvalin, P. Detection of glycopeptide

resistance genotypes and identification to the species level of clinically

relevant enterococci by PCR. *J. Clin. Microbiol.* **33,** 24–27 (1995).

47.   Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its

708   Applications to Single-Cell Sequencing. *J. Comput. Biol.* **19,** 455–477 (2012).

709   48.   Overbeek, R. *et al.* The SEED and the Rapid Annotation of microbial

710   genomes using Subsystems Technology (RAST). *Nucleic Acids Res.* **42,**

711   D206–D214 (2014).

712   49.   Jolley, K. a & Maiden, M. C. J. BIGSdb: Scalable analysis of bacterial

713   genome variation at the population level. *BMC Bioinformatics* **11,** 595 (2010).

714   50.   Camacho, C. *et al.* BLAST plus: architecture and applications. *BMC*

715   *Bioinformatics* **10,** 1 (2009).

716   51.   Zankari, E. *et al.* Identification of acquired antimicrobial resistance genes. *J.*

717   *Antimicrob. Chemother.* **67,** 2640–2644 (2012).

718   52.   Jensen, L. B. *et al.* A classification system for plasmids from enterococci and

719   other Gram-positive bacteria. *J. Microbiol. Methods* **80,** 25–43 (2010).

720   53.   Lozano, C. *et al.* Expansion of a plasmid classification system for gram-

721   positive bacteria and determination of the diversity of plasmids in

722   Staphylococcus aureus strains of human, animal, and food origins. *Appl.*

723   *Environ. Microbiol.* **78,** 5948–5955 (2012).

724   54.   Siguier, P., Perochon, J., Lestrade, L., Mahillon, J. & Chandler, M. ISfinder:

725   the reference centre for bacterial insertion sequences. *Nucleic Acids Res.*

726   **34,** D32–D36 (2006).

55.   Sillanpää, J., Prakash, V. P., Nallapareddy, S. R. & Murray, B. E. Distribution

of genes encoding MSCRAMMs and pili in clinical and natural populations of

Enterococcus faecium. *J. Clin. Microbiol.* **47,** 896–901 (2009).

56. Grissa, I., Vergnaud, G. & Pourcel, C. CRISPRFinder: a web tool to identify

clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res.*

35, W52-7 (2007).

57. Makarova, K. S. *et al.* Evolution and classification of the CRISPR–Cas

systems. *Nat. Rev. Microbiol.* **9,** 467–477 (2011).

58. Page, A. J. *et al.* Roary: Rapid large-scale prokaryote pan genome analysis.

Bioinformatics **31,** 3691–3693 (2015).

59. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and

high throughput. *Nucleic Acid Res.* **32,** 1792–1797 (2004).

60. Stamatakis, A. RAxML version 8: A tool for phylogenetic analysis and post-

analysis of large phylogenies. *Bioinformatics* **30,** 1312–1313 (2014).

61. Drummond, A. J., Suchard, M. a, Xie, D. & Rambaut, A. Bayesian P

hylogenetics with BEAUti and the BEAST 1 . 7. *Mol. Biol. Evol.* **29,** 1969–

1973 (2012).

62. Kurtz, S. *et al.* Versatile and open software for comparing large genomes.

Genome Biol. **5,** R12 (2004).

63. Miller, M. A., Pfeiffer, W. & Schwartz, T. The CIPRES science gateway:

enabling high-impact science for phylogenetics researchers with limited

resources. *Proc. 1st Conf. Extrem. Sci. Eng. Discov. Environ. Bridg. from*

Extrem. to campus beyond 1–8 (2012). doi:10.1145/2335755.2335836

Author contributions

R.R. performed experiments, carried out all statistical analyses, analysed results

and wrote draft of the manuscript, L.D. and C.A.A conceived the study, analysed

the results, and drafted and reviewed the manuscript, J.R. and D.P. conceived the

study, interpreted data and analysed the results, P.J.P and SO.K. conceived

experiments and provided key experimental suggestions, B.E.M. T.T.T and J.M.M

interpreted and analyse data and helped to write the manuscript, L.P.C., S.R,

A.M.E., A.D. and A.N. performed experiments and analysed data. All authors

contributed to improve the manuscript and gave approval of the final version prior

to submission.

Additional Information

763 C.A.A has received grants funded by Merck Pharmaceuticals, MeMed Diagnostics

764 Ltd and Entasis Therapeutics. B.E.M has received grants funded by Cubist/Merck,

765 Forest/Actavis and is consultant of Paratek and Cempra.

766 The other authors declare no competing interests.

 Figure Legends

Figure 1. (A) Bayesian phylogenomic tree from the core genome and genomic

characterization of resistance elements of 55 representative Latin American VR*Efm*

strains; the presence of a genetic element is marked as a black box in the

corresponding column of the isolate. (B) Phenotypic resistance profile of 207 clinical

isolates of VR*Efm* from our Latin American collection for vancomycin (VAN),

teicoplanin (TEI), ampicillin (AMP), chloramphenicol (CHL), ciprofloxacin (CIP),

linezolid (LNZ), high-level resistance to gentamicin (HLR-GE) and high-level

resistance to streptomycin (HLR-STR).

Figure 2. Bayesian phylogenomic tree from the core genome and genomic

characterization of virulence factors of 55 representative Latin American VR*Efm*

strains, the presence of a genetic element is marked as a black box in the

corresponding column of the isolate.

Figure 3. Bayesian phylogenomic tree from the core genome of 340 genomes

sampled from 36 countries between 1946 and 2017 and from different sources. Blue

branches showed the genomes grouped within clade B, while brown branches show

isolates from clade A. The outer coloured rings (from inner to outer) indicate the

source of each isolate, the region from which it was sampled and its relationship

through MLST typing (if possible) to Clonal Complex 17. Labels show the isolates

originating from our Latin American collection.

Figure 4. Bayesian phylogenomic tree from the non-recombinant regions of the 303

Clade A genomes. Branches highlighted in orange represent genomes from the

animal early branches. Branches highlighted in pink show genomes from clinical

related isolates. Red and green branches show the genomes from clinically related

subclades (CRS) I and II, respectively. Annotation rings (from inner to outer) show

the sequence type (ST) of the isolate (only the five most prevalent STs in the sample

are shown), the isolation year, the region from which the isolate was sampled and if

the region was Latin America, the exact country from where it was obtained. The last

ring shows which isolates were recovered from blood.

Prevalent Sequence types

- 17
- 18
- 78
- 412
- 736

Regions

- Europe
- North America
- Asia
- Australia
- Africa

Latin American Countries

- Argentina
- Brazil
- Chile
- Colombia
- Ecuador
- Mexico
- Peru
- Venezuela

- Animal related
- Clinical related subclade I
- Clinical related subclade II
- Isolates from blood

Tree scale: 0.0001