Genetic Deletion of TrpV1 and TrpA1 Does not Alter Avoidance of or Patterns of Brainstem

Activation to Citric Acid in Mice

Tian Yu

Courtney E. Wilson

Jennifer M. Stratford

Thomas E Finger

Rocky Mountain Taste & Smell Center, Department of Cell and Developmental Biology,

University of Colorado School of Medicine, Aurora, CO 80045

ACKNOWLEDGEMENTS:

The authors thank Mei Li for histological preparations and for comparison of antibody staining

properties. We also thank NIDCD for supporting this investigation (RO1 DC012931).

ABSTRACT:

Exposure of the oral cavity to acidic solutions evokes not only a sensation of sour, but also of

sharp or tangy. Acidic substances potentially stimulate both taste buds and acid-sensitive

mucosal free nerve endings. Mice lacking taste function (P2X2/P2X3 double-KO mice) refuse

acidic solutions similarly to wildtype mice and intraoral infusion of acidic solutions in these KO

animals evokes substantial c-Fos activity within orosensory trigeminal nuclei as well as of the

nucleus of the solitary tract (nTS) (Stratford et. al 2017). This residual acid-evoked, non-taste

activity includes areas that receive inputs from trigeminal and glossopharyngeal peptidergic

(CGRP-containing) nerve fibers that express TrpA1 and TrpV1 both of which are activated by

low pH. We compared avoidance responses in wildtype (WT) and TrpA1/V1 double KO

(TRPA1/V1$^{Dbl-/-}$) mice in brief-access behavioral assay (lickometer) to 1, 3, 10, 30 mM citric

acid, along with 100 μM SC45647 and H_2O. Both WT and TRPA1/V1$^{Dbl-/-}$ show similar

avoidance, including to higher concentrations of citric acid (10 and 30 mM; pH 2.62 and pH 2.36

respectively), indicating that neither TrpA1 nor TrpV1 is necessary for the acid avoidance

behavior in animals with an intact taste system. Similarly, induction of c-Fos in the nTS and

dorsomedial spinal trigeminal nucleus was similar in the WT and TRPA1/V1$^{Dbl-/-}$animals. Taken

together these results suggest non-TrpV1 and non-TrpA1 receptors underlie the residual

responses to acids in mice lacking taste function.

INTRODUCTION

When acidic solutions (e.g. lemon juice) are taken into the mouth, we describe the

resulting perception as "sour" and consider it to be an aversive basic taste. However, the

sensations arising from acidic substances in the oral cavity are a compound sensation of sour (a

taste originating from taste buds) and direct activation of acid-sensitive general mucosal afferents

of the oropharynx including the trigeminal, glossopharyngeal and superior laryngeal nerves.

Indeed, a dictionary definition of sour (thefreedictionary.com) includes descriptors such 'sharp,'

'tart', or 'tangy' which are not taste modalities, but rather chemesthetic ones corresponding to

activation of acid-sensitive mucosal nerves likely including polymodal nociceptors. Thus,

aversion to acidic solutions might be due to activation of these nociceptors rather than of the

taste system alone.

In mice lacking a functional taste system (P2X2/3-dbl-KO mice), the chorda tympani

nerve, which provides only intragemmal innervation to taste buds (Silverman and Kruger 1990,

Ohman-Gault, Huang et al. 2017), shows no responses to any tastants including acids although

tactile and thermal responses remain (Finger, Danilova et al. 2005). Consistent with this, these

mice show no preference for sweeteners or avoidance of bitter substances in brief-access taste

tests. Despite the apparent lack of taste responses to acids, the P2X2/3-dbl-KO mice do exhibit

normal avoidance of citric acid in similar brief access tests (Hallock, Tatangelo et al. 2009). This

avoidance may be mediated not by taste, which is non-functional in these mice, but by acid-

responsive fibers in the trigeminal, glossopharyngeal or laryngeal nerves which do show

residual low level activity in the P2X2/3-dbl-KO mice (Ohkuri, Horio et al. 2012).

Polymodal nociceptors respond to acidification as well as other potentially painful stimuli

(Bessou and Perl 1969). Many small caliber polymodal nociceptors that innervate the oral cavity

(Kichko, Neuhuber et al. 2018, Wu, Arris et al. 2018) express one or both of the pH-sensitive

transient receptor potential (Trp) channels, TrpA1 (Wang, Chang et al. 2011) and TrpV1

(Tominaga, Caterina et al. 1998). TrpA1 has been implicated especially in responsiveness to

weak acids capable of penetrating cell membranes to produce intracellular acidification (Wang,

Chang et al. 2011). TrpV1 is implicated in responses of the glossopharyngeal and vagus nerves

to acidification of the oral and pharyngeal epithelium (Arai, Ohkuri et al. 2010). Gating of either

of these channels by acids can directly activate the nociceptive fiber resulting in a noxious

sensation suitable for driving avoidance behavior. Ablation of the ganglion cells expressing

TrpV1 in mice lacking sour taste receptors does result in loss of acid-avoidance behavior (Zhang,

Jin et al. 2019) but it is not clear whether this is due to loss of TrpV1 itself or other ion-sensing

mechanisms of these neurons. We tested whether genetic deletion of both TrpA1 and TrpV1

affected either acid-induced avoidance behaviors or the pattern of neuronal activation in the

brainstem taste (nuc. solitary tract, nTS) or trigeminal oral (dorsomedial trigeminal nucleus,

DMSp5) sensory nuclei.

Materials and Methods

Animals

TrpA1V1-double KO mice, on a C57BL6 background (TRPA1/V1$^{Dbl-/-}$) were a generous gifts from Diana Bautista, University of California Berkeley (Gerhold and Bautista 2008). In these mice, deletion of TrpA1 had been accomplished by elimination of residues 901–951 including most of exon 23 – a region encoding the putative pore and part of the sixth transmembrane domain (Bautista, Jordt et al. 2006). The VR1 gene was disrupted (Caterina, Leffler et al. 2000) by deleting part of the fifth and all of the sixth putative transmembrane domains, together with the intervening pore-loop region. The TRPA1/V1$^{Dbl-/-}$ mice were genotyped by PCR to ensure quality control of the breeding. C57BL6/J mice were purchased from The Jackson Laboratory (Bar Harbor, ME). For brief access lickometer behavioral assays, 2-6 months old male (n=9) and female (n=6) TRPA1/V1$^{Dbl-/-}$ mice and 2-10 month old male (n=8) and female (n=3) C57BL6/J mice were used. For immunohistochemical staining, we used TRPA1/V1$^{Dbl-/-}$ (2 to 8 mo, n=10, 7 male and 3 female)) and C57BL6/J mice (4 to 7 mo, n=6, all male). In an addition to the C57 cases, we utilized 2 mice (1 male, 1 female) of the line B6.Cg-Tg(Fos-tTA,Fos-EGFP*)1Mmay/J which carry 2 transgenes associated with c-Fos: cfos-tTA and cfos-shEGFP. In these mice, on a C57BL6/J background, the randomly inserted transgenes utilize the c-Fos promotor to drive expression of respectively tTA and a short-lived (two-hour half-life) GFP. We did not find close correlation between GFP expression and c-Fos immunostaining and so the GFP results were ignored in our analysis. Nonethelss, the c-Fos counts from these animals were entirely consistent with those obtained from the wildtype C57BL6/J mice and so were included in those results.

All animal procedures were performed in accordance with NIH guidelines and were approved by

the Institutional Animal Care and Use Committee at the University of Colorado School of

Medicine.

Brief access preference tests

Davis Rig Lickometers (MS-160; Dilog Instruments and Systems) were used for training

and brief-access testing of mice. The animals underwent a 4-day water training before acid

testing. Mice were placed on water deprivation 20 hours before the first day of training or acid

testing. On the first day of the 4-day training, water-deprived animals were allowed access to a

single water spout for a 30-min period. In the following three days of water training, 2-4 bottles

of water were given in random sequences for 15 min. Mice had 5 seconds to lick at each spout

after their first lick before the door closed and the bottle was switched to the next position with

an inter-trial interval of 7.5 s. The mice were considered well-trained if they lick more than 30

times consistently during each trial for the first 15 trials. No mice lost more than 15% of body

weight during any phase of the experimental procedure. All mice were given 2 day recovery

period with ad libitum access to water after every 4 day block of testing.

For the acid testing days, six bottles of tastants including 4 concentrations of acetic or

citric acid (1 mM, 3 mM, 10 mM and 30 mM), 1 concentration of artificial sweetener SC45647

(100 μM or 300μM) and H_2O were presented in a testing block with a set up similar to the

training session. Testing periods lasted for 15 min, with the opportunity for the mice to sample

from the 6 solutions a total of 30 times. Licks were measured using InstaCal software. Each

block of 6 tastant trials featured each solution in random order. Only data from completed blocks

were used in the calculation of preference. Preference was calculated by averaging the licks per

tastant block of each solution relative to those for water (i.e., a 'lick ratio'). The same mice that

were assessed for citric acid preference were then tested for acetic acid preference again with the

same paradigm. The solutions used in acetic acid testing were as follows: 100 or 300 μM

SC45647, H_2O, 1, 3, 10 and 30 mM acetic acid. Testing for each series of tastants was repeated

at least twice. Because there was no statistically significant difference in licking across each

testing day for each mouse [$F_{(1,2)} = 0.42$, $p = 0.66$], the number of licks for each tastant were

averages across these testing days for each mouse.

Citric acid stimulation for c-Fos via intraoral cannula

Both bilateral intraoral cannulae implantation and stimulation methods were adopted

from (Stratford and Thompson 2016). Briefly, mice were anesthetized with an intramuscular

injection of a combination of medetomidine hydrochloride (Domitor; 0.4 mg/kg; Pfizer) and

ketamine hydrochloride (40 mg/kg; Bioniche Pharma). Intraoral cannulae were inserted via a

midline incision immediately caudal to the pinnae. Then a sterile 2-3 cm stainless steel needle

(19 gauge; Hamilton) with a flared end polyethylene tube (50 gauge; Becton Dickinson) was

inserted from behind the pinna and guided subcutaneously into the oral cavity, lateral to the first

maxillary molar. The needle was then withdrawn, leaving the polyethylene cannula and washer

in place in the rear of the oral cavity. These mice were given 4 days for recovery prior to

training.

For training and testing, liquids were delivered into intraoral cannula via a 5-cc syringe

connected to a syringe pump (Model R99-E, Razel Scientific Instruments). To train the mice for

the acclimation of liquid stimulation through the cannulae, they were water deprived for 23 hour

/ day, and given 3 ml of deionized water as a constant pulsatile flow of 0.1 ml/min through one

of the two intraoral cannulae over the course of 30 min. We trained all the mice with deionized

water for 2 days followed by stimulation with either deionized water (n=4 [all M] for C57BL6/J,

n=5 [2 F, 3 M] for TrpA1V1-dbl KO), or 30 mM citric acid (n=4 [1 F, 3 M] for C57BL6/J, n=5

[1F, 4M] for TrpA1V1-dbl KO). Animals exposed to the stimulus were left undisturbed for an

additional 45 min prior to sacrifice.

c-Fos immunohistochemistry

Seventy-five minutes after onset of taste stimulation through cannula, animals were

deeply anesthetized with Fatal-Plus® (50 mg/kg intraperitoneally; MWI), and then perfused

transcardially with saline followed by 4% paraformaldehyde (PFA) in 0.1 M pH 7.2 phosphate

buffer (PB). The brains were post fixed for 3 hours at room temperature, and then cryo-protected

overnight in 20% sucrose in PB at 4 °C. After cyroprotection, brainstems were cut and embedded

in OCT, frozen and sectioned at 40 μm using a cryostat. Free-floating sections were collected in

PBS.

For immunostaining, sections were washed in PBS 3 times and then processed for antigen

retrieval in sodium citrate (pH 6) at 85 °C for 10 min. After tissues cooled, non-specific protein

binding was blocked in a medium consisting of 2% normal donkey serum (Jackson

ImmunoResearch) in antibody medium (AB medium: 0.3% TritonX100, 0.15 M NaCl, and 1%

BSA in PB) for 1 hour at room temperature.

We utilized Mouse c-Fos antibody (1:1000, PhosphoSolutions; Cat#: 309-cFos; RRID

RRID:AB_2632380; Lot#: GS117P) and Rabbit P2X2 antibody (1:1000, Alomone Labs;

Cat#:APR003; RRID AB_2040054; Lot# APR003AN1002). These antibodies were diluted in

AB medium accordingly and were used to incubate the sections for 4 days at 4 °C. After 3 10-

min washes in PBS, sections were transferred to secondary antibody cocktail (Alexa Fluor 568

donkey anti mouse, 1:500; Alexa Fluor 488 donkey anti rabbit, 1:500; NeuroTrace Nissl

640/660, 1:500; all from Life Technologies) for 2-hour incubation at room temperature.

Following another 3 10-min washes in PBS, free-floating tissue sections were mounted onto

Microscope Slides (Tanner scientific, #TNR WHT90AD), and then coverslipped using

Fuoromount-G (Southern Biotech).

According to the manufacturer's data sheet, the PhosphoSolutions Mouse c-Fos antibody

shows a broad band of reactivity in Western blots of HeLa cells centered at 50kDa. We tested

whether this antibody stains similarly in fixed mouse brain tissues by comparing

immunoreactivity of this antibody to the one we have utilized previously: we allowed a wildtype

mouse to drink 150 mM MSG for 30min and perfused the mouse with 4% buffered

paraformaldehyde after an additional 45 mins. The olfactory bulb & brainstem were removed,

cryoprotected in 20% sucrose and frozen free-floating sections were cut at 40 µm.

Representative sections were incubated in sodium citrate buffer pH6 at 85 °C for 10 min. After

cooling to room temperature for 20 min, the sections were rinsed 3 X 5min in PBS, then

incubated with 2%NDS+AB media for 1hr at room temperature. The free-floating sections then

were incubated with a mixture of primary antibodies: rabbit a-cFos (lot.D00148958) 1:500

/ mouse a-cFos (lot. GS418y) 1:1000 for 4 nights at 4 °C. After 3 X 10 min rinses in PBS,

sections were incubated with a mixture of secondary antibodies: A488 donkey-a-rabbit 1:500/

A568 donkey-a-mouse 1:500 / Nissl 1:500 for 2 hrs at room temperature. The sections then were

rinsed in PBS followed by 0.05M PB prior to coverslipping. The distribution of labeled cells

within and around the nTS was similar to that observed in single label cases and nearly all cells

were labeled by both antibodies. To quantitatively analyze the degree of co-localization, we

counted cells labeled by one or both antibodies in the olfactory bulb where the density of labeled

cells permits quantitative assessment. In randomly selected fields through the granule cells layer

of 3 sections, we counted the number of single and double-label cells. Of 153 labeled cells, 149

were double-labeled; 4 were labeled by only the PhosphoSolutions antiserum and none were

labeled only by the rabbit antibody. We conclude that the 2 antibodies label nearly identical

populations and that results from the PhosphoSolutions antibody should be comparable to those

obtained previously with the rabbit antibody.

c-Fos activity determination

Brainstem sections were observed under an Olympus BX41microscope. Representative

levels (r1, r2, r3, r4, i1, i2) of the nucleus of the solitary tract (nTS) were chosen under a 10X

magnification objective according to Stratford et al. (Stratford, Thompson et al. 2017) and

photographed at 20X magnification using CellSense software with a XM10 camera. Boundaries

of the nTS were drawn using P2X2 and Nissl staining as reference and according to Stratford et

al. (2017) in ImageJ. Boundaries of the DMSp5 were not distinct based on the images but the

approximate boundaries were drawn using nTS boundaries and other brainstem nuclei as

reference (Corson, Aldridge et al. 2012). The DMSp5 was taken as the area between the lateral

edge of nTS and the spinal trigeminal nucleus.

The area of nTS was further divided into 6 parts following the system described in

Stratford & Finger (Stratford and Finger 2011) using a horizontal line dividing the area into

dorsal and ventral tiers and two vertical lines dividing the medial-lateral extent into thirds. The

six parts of nTS were named DM, DI, DL, VM, VI, VL, and the cFos signals in these areas as

well as DMSp5 were counted using ImageJ Cell counter plugin. The identity, stimulus and

genotype of all cases were blinded to the person performing the cell counts.

Statistical Analysis

Behavioral data are presented as group means with individual data points indicated.

Immunohistochemical data are presented as group means ± SD. Data were first analyzed for

normality using an Anderson-Darling test (AD). Because all data were deemed normally

distributed based on the results of the AD test, appropriate two- and three-way analysis of

variance (ANOVA)s were conducted (Statistica; StatSoft, Tulsa, OK). Tukey's honest significant

difference tests were used to assess statistically significant (p < 0.05) main effects or interactions

(see Results for details).

RESULTS

Behavioral Assessment

For the acetic acid experiments, behavioral preferences were not statistically different

between males and females across each genotype (WT and TRPA1/V1[Dbl-/-]), (genotype x sex

interaction, $F(1, 14) = 0.00$, $p = 0.99$). However, behavioral preferences were statistically

different between different taste solutions, (main effect of taste, $F(4, 56) = 24.89$, $p < 0.0001$,

Fig. 1A).

For the citric acid experiments, behavioral preferences were not statistically different

between males and females, $F(1, 22) = 2.80$. $p = 0.11$. Moreover, the overall preference curves

for both WT and TRPA1/V1[Dbl-/-] animals appeared nearly identical, (taste x genotype interaction,

$F(4,88) = 0.34$, $p = 0.85$, Fig. 1B).

c-Fos Induction

Injection of 30 mM citric acid in the mouth via intraoral cannulae reliably induced c-Fos

within particular regions of the nTS as compared to similar injection of water. As calculated

previously, raw counts of c-Fos positive neurons in sub-compartments of the nTS were compared

between water-stimulated and citric acid-stimulated animals to produce a measure of citric acid-

specific c-Fos induction. In our previous study (Stratford, Thompson et al. 2017), acid-induced

c-Fos was highest in the central portion of rostral-intermediate nTS (r3-i2) and in ventrolateral

subdivisions in intermediate nTS (i1-i3). These regions did not show such activation in the

P2X2/P2X3 dbl KO mice which lack taste function. However, in P2X2/P2X3 dbl KO animals,

significant levels of activation to citric acid did remain in the DMSp5, which receives substantial

input from polymodal nociceptor (CGRP+) fibers of the trigeminal and glossopharyngeal nerves.

Accordingly, we especially focused on possible changes in c-Fos activation in this trigeminal-

recipient area. Fig.2 shows a representative image of c-Fos activation in relation to the lateral

nTS and the DMSp5 of TRPA1/V1$^{Dbl-/-}$ mice following stimulation with citric acid.

<u>'Raw' Tastant-Evoked c-Fos Counts</u>. For all animals (i.e. both WT and TRPA1/V1$^{Dbl-/-}$

mice), the total number of c-Fos positive cells evoked by citric acid oral stimulation was

significantly greater than water stimulation within the nTS overall (water: 39.48 ± 3.59; citric

acid: 71.03 ± 5.65, main effect of tastant: $F(1, 5) = 49.53$, $p < 0.001$; see Table 1). Moreover, the

total number of water-evoked and citric acid-evoked c-Fos positive cells were significantly

higher in WT as compared to TRPA1/V1$^{Dbl-/-}$ mice, regardless of tastant (For WT: 73.50 ± 7.17;

For TRPA1/V1$^{Dbl-/-}$: 58.25 ± 5.57; main effect of genotype: $F(1, 5) = 6.67$, $p < 0.05$. However,

there was no genotype x tastant interaction: $F(1, 82) = 0.05$, $p = 0.83$, see Table 1 and Fig. 3).

Finally, there was no significant three-way tastant x genotype x nTS level interaction (F (1, 82) =

0.28, p = 0.92), nor was there a four-way tastant x genotype x nTS level x nTS subregion

interaction (F (25, 410) = 0.55, p = 0.96).

Table 1. Average (+/-S.D.) c-fos positive cells per region in
the nTS and DMSp5 in WT and TrpA1/V1[Dbl-/-] mice following
water and citric acid stimulation.

	Water	Citric Acid
WT	5.28 ± 1.58	9.83 ± 1.59
TRPA1/V1[Dbl-/-]	13.72 ± 2.50	19.17 ± 2.54

<u>Citric Acid- Specific c-Fos Activation</u>. Because the number of c-Fos positive cells was

lower in TRPA1/V1[Dbl-/-] mice as compared to WT nice (regardless of tastant), we calculated citric

acid-specific activation for each region by taking the number of 'raw' citric acid-evoked c-Fos

positive cells minus the average number of water-evoked c-Fos positive cells for each region for

each genotype. Overall, citric-acid specific activation in the nTS did not differ between

genotypes (main effect of genotype: F (1,5) = 0.08, p = 0.78). Furthermore, as shown in Fig. 3,

the amount of citric acid-specific c-Fos was significantly different between nTS subregions and

levels regardless of genotype, interaction between nTS subregion x nTS level: F (25, 210) =

4.24, p < 0.001. In particular, citric acid evoked significant activity in the central subregion of

rostrointermediate nTS (r2-i2), in the lateral subregion of intermediate nTS (i1-2) and in the

DMSp5 of both WT and TRPA1/V1[Dbl-/-] mice as compared to all other nTS subregions (all p's <

0.05). This pattern is similar to that reported previously for the mixed background (Ola-

C57/BL6) controls in the previous study (Stratford, Thompson et al. 2017).

In summary, the TRPA1/V1[Dbl-/-] mice show no differences in the pattern of neural

activation by citric acid compared to the WT animals. This suggests that neither TrpA1 nor

TrpV1 plays a significant role in detection and avoidance of 30 mM concentrations of citric acid,

although this concentration is readily avoided by both TRPA1/V1$^{Dbl-/-}$ and WT animals.

DISCUSSION

Acidity is the underlying chemical feature of sour substances. When acids are taken into

the mouth, they stimulate sour-sensitive taste cells which depolarize, then release

neurotransmitter to activate the taste fibers innervating them. This signaling between taste cells

and nerve fibers requires functional P2X2 and P2X3 receptors; genetic deletion or

pharmacological blockage of these receptors essentially eliminates taste-mediated neural activity

(Finger, Danilova et al. 2005, Vandenbeuch, Anderson et al. 2013). Similarly, taste driven

acceptance of sweet and umami, and taste driven avoidance of bitter is lost in mice lacking

functional P2X2 and P2X3 receptors. In contrast, behavioral avoidance of intraoral acids remains

intact despite the loss of taste-related neural activity to these stimuli. Similarly, avoidance of

acids persists even after genetic deletion of Otop1, the ion channel receptor for H^+ underlying

sour detection (Teng, Wilson et al. 2019, Zhang, Jin et al. 2019). These findings strongly suggest

that not just taste, but another sensory modality drives the avoidance of acids in this context.

The oropharynx is innervated by chemically-sensitive free nerve endings arising from the

trigeminal, glossopharyngeal and vagus nerves. These nerves include populations of polymodal

nociceptors that express acid-sensitive ion channels including TrpA1 and TrpV1 (Tominaga,

Caterina et al. 1998, Wang, Chang et al. 2011), and the TrpV1-expressing fibers are necessary

for non-taste mediated avoidance of citric acid (Zhang, Jin et al. 2019). Yet these studies do not

demonstrate that TrpV1 itself is the necessary receptor. It is likely that other acid-sensitive

channels and receptors exist in these TrpV1-expressing fibers. Further, TrpV1 fully activates at a

pH around 5 at body temperature (Tominaga, Caterina et al. 1998), whereas avoidance of citric

acid begins at a pH near 3. It is likely that the tissue overlying the TrpV1 sensory terminals

provides some buffering of acids applied to the surface of the epithelium, but weak acids, such as

citric acid, effectively acidify the epithelium deep into the tissue (Richter, Caicedo et al. 2003),

well beyond the region in which the nerve terminals lie. Taken together, these results suggest

that TrpV1 itself may not be entirely responsible for responses to citric acid. Accordingly, we

tested whether either TrpV1 or TrpA1 channels contribute to either the behavioral avoidance

response, or the activation of brainstem neurons by citric acid. We found that neither the

behavior nor the pattern and degree of neural activation was altered by genetic deletion of these

channels.

If neither TrpV1 nor TrpA1 channels underlie avoidance of acidic substances in the

absence of taste, what other mechanisms might be responsible? Likely candidates are one or

more members of the ASIC (acid sensing ion channel) family, which are gated by decreases in

extracellular pH. While many members of this family open at pH values near 7, others activate

at lower pH values (Deval, Gasull et al. 2010). In particular, ASIC3 is expressed widely in

polymodal nociceptors including those expressing TrpV1 (Trigeminal ganglion clusters 7 & 10

in (Nguyen, Wu et al. 2017)) and plays a role in responses to pH5.0 (Price, McIlwrath et al.

2001). Further, amiloride, a non-specific blocker of ASIC channels, decreases the irritation of

citric acid (albeit at a much higher concentration) measured psychophysically (Dessirier,

O'Mahony et al. 2000), supporting the role of ASICs in this response. Conversely, inclusion of

amiloride, a blocker of ASICs, in orally-applied solutions in rats neither decreased the trigeminal

response to citric acid nor attenuated acid-induced c-Fos in brainstem trigeminal nuclei (Sudo,

Sudo et al. 2003). Whether ASICs or some other acid-responsive mechanisms play a role in non-

gustatory behavioral avoidance of weak acids then is unresolved.

334 Fig. 1: Brief access tests showing similar avoidance of acetic acid and citric acid for

335 TRPA1/V1$^{Dbl-/-}$ mice and wildtype controls of the same background. ANOVA shows no

336 significant difference between the groups.

Fig. 2. C-Fos staining (red dots) in nTS and DMsp5 of the TRPA1/V1$^{Dbl-/-}$ mouse at

approximately the R3 level comparing a mouse stimulated by 30 mM citric acid (A, B) to one

stimulated only by water (C, D). The number and location of c-fos positive cells is similar in

WT and TRPA1/V1$^{Dbl-/-}$ strains. Note the higher number of c-fos positive cell nuclei (red) in the

acid stimulated compared to the water-stimulated animal. Green staining shows the distribution

of P2X3-immunoreactive fibers which terminate largely in the dorsocentral part of the nTS. Most

acid-stimulated cells lie in the ventral tier of the nucleus and in the DMSp5 region ventrolateral

to the nTS. Blue counterstain from FluoroNissl.

WT Mice

KO Mice

Water Stimulated

WT Mice

		Medial	I	Lateral	DMSp5
r1	D	1.5	2.3	2.3	
	V	2.1	2.0	1.1	2.9
r2	D	3.3	2.9	2.9	
	V	2.1	1.0	1.9	3.0
r3	D	5.6	12.6	9.8	
	V	4.1	13.4	9.6	8.5
r4	D	4.3	16.6	10.9	
	V	4.5	14.8	12.9	10.6
i1	D	15.0	15.8	15.1	
	V	7.3	9.5	12.1	12.3
i2	D	11.2	14.2	7.8	
	V	5.5	8.2	12.2	18.2

KO Mice

		Medial	I	Lateral	DMSp5
r1	D	2.3	5.0	1.9	
	V	1.4	0.6	0.9	2.1
r2	D	2.4	8.0	5.4	
	V	1.6	2.6	4.0	5.1
r3	D	4.1	10.9	3.8	
	V	3.0	5.3	2.9	2.4
r4	D	1.6	14.3	6.5	
	V	1.5	11.9	4.5	5.3
i1	D	11.6	9.9	9.9	
	V	6.1	6.8	9.2	9.1
i2	D	9.5	8.6	4.6	
	V	5.8	6.3	6.7	9.9

Citric Acid Stimulated

WT Mice

		Medial	I	Lateral	DMSp5
r1	D	3.4	6.9	4.6	
	V	2.9	2.3	2.6	4.9
r2	D	4.5	9.8	3.5	
	V	5.0	5.8	4.0	2.8
r3	D	7.0	21.4	8.1	
	V	5.9	18.9	12.3	11.1
r4	D	9.6	27.9	13.3	
	V	8.3	24.6	20.3	22.6
i1	D	22.9	31.0	13.6	
	V	10.8	22.4	26.4	20.3
i2	D	13.3	22.9	11.6	
	V	9.5	16.6	28.1	26.4

KO Mice

		Medial	I	Lateral	DMSp5
r1	D	3.7	5.0	2.3	
	V	2.4	2.8	1.7	3.4
r2	D	5.9	13.4	5.3	
	V	6.4	6.6	4.4	5.2
r3	D	6.2	17.2	8.3	
	V	6.1	19.0	11.0	6.3
r4	D	9.5	23.1	13.7	
	V	7.0	24.8	13.4	10.9
i1	D	10.6	22.9	18.0	
	V	5.8	19.2	27.9	17.3
i2	D	10.0	12.3	7.5	
	V	8.3	15.1	22.6	17.3

Acid Specific (C.A. - water)

WT Mice

		Medial	I	Lateral	DMSp5
r1	D	1.9	4.6	2.4	
	V	0.8	0.3	1.5	2.0
r2	D	1.3	6.9	0.6	
	V	2.9	4.8	2.1	-0.3
r3	D	1.4	8.8	-1.6	
	V	1.8	5.5	2.6	2.6
r4	D	5.4	11.3	2.4	
	V	3.8	9.9	7.4	12.0
i1	D	7.9	15.3	-1.5	
	V	3.5	12.9	14.3	8.0
i2	D	2.1	8.7	3.8	
	V	4.0	8.5	16.0	8.2

KO Mice

		Medial	I	Lateral	DMSp5
r1	D	1.5	0.0	0.4	
	V	1.0	2.2	0.8	1.3
r2	D	3.5	5.4	-0.1	
	V	4.8	4.0	0.4	0.1
r3	D	2.1	6.3	4.5	
	V	3.1	13.7	8.1	3.9
r4	D	7.9	8.8	7.2	
	V	5.5	12.9	8.9	5.6
i1	D	-1.0	13.0	8.1	
	V	-0.3	12.4	18.7	8.2
i2	D	0.5	3.7	2.9	
	V	2.5	8.8	15.9	7.4

Fig. 3.Heat maps of c-Fos expression in nTS and DMSp5 regions comparing wildtype (left) and

KO (TRPA1/V1$^{Dbl-/-}$) mice, stimulated by intraoral injection of water (top row and by citric acid

(middle row). The numbers indicate the aaverage number of c-fos-positive neurons in that

compartment. See Suppl. Table 1 for means and standard deviations for each location for each

genotype. Bottom row shows acid-specific activation, i.e. citric acid counts minus the counts

produced by water alone. No significant differences occur either in the pattern or absolute level

of expression across subregions (see text for details).

.

Supplemenatry Table 1: Means and standard deviations of counts of c-fos positive cells in

DMSp5 and each subnucleus of the NTS for water and citric acid stimulation by intraoral

cannula for WT and TrpA1/V1$^{-/-}$ mice.

.

WT H2O		DM	DI	DL	VM	VI	VL	DMSp5
r1		1.5	2.3	2.3	2.1	2.0	1.1	2.9
	stdev	1.1	1.8	1.0	0.8	1.1	1.1	2.7
r2		3.3	2.9	2.9	2.1	1.0	1.9	3.0
	stdev	1.4	1.5	2.0	1.1	0.5	1.0	1.8
r3		5.6	12.6	9.8	4.1	13.4	9.6	8.5
	stdev	4.3	6.3	5.9	1.9	7.6	6.0	9.5
r4		4.3	16.6	10.9	4.5	14.8	12.9	10.6
	stdev	3.0	6.7	7.0	2.1	4.4	6.4	6.9
i1		15.0	15.8	15.1	7.3	9.5	12.1	12.3
	stdev	6.2	8.7	6.3	5.3	4.5	6.8	8.2
i2		11.2	14.2	7.8	5.5	8.2	12.2	18.2
	stdev	10.6	8.0	4.7	3.7	5.5	7.5	14.4

WT Cit Acid		DM	DI	DL	VM	VI	VL	DMSp5
r1		3.4	6.9	4.6	2.9	2.3	2.6	4.9
	stdev	1.8	1.8	3.3	2.6	1.3	1.6	3.4
r2		4.5	9.8	3.5	5.0	5.8	4.0	2.8
	stdev	1.6	3.4	2.0	2.7	4.3	2.1	2.9
r3		7.0	21.4	8.1	5.9	18.9	12.3	11.1
	stdev	3.6	8.3	3.4	2.7	2.6	6.8	9.4
r4		9.6	27.9	13.3	8.3	24.6	20.3	22.6
	stdev	3.4	9.4	3.3	3.9	6.0	7.4	14.3
i1		22.9	31.0	13.6	10.8	22.4	26.4	20.3
	stdev	10.7	9.6	7.3	4.8	4.9	7.8	8.5
i2		13.3	22.9	11.6	9.5	16.6	28.1	26.4
	stdev	9.0	4.8	5.4	5.4	7.5	9.8	18.8

368
369
370
371

TrpA1/V1$^{-/-}$ H2O	DM	DI	DL	VM	VI	VL	DMSp5
r1	2.3	5.0	1.9	1.4	0.6	0.9	2.1
stdev	3.4	4.6	2.0	1.5	0.7	1.7	2.9
r2	2.4	8.0	5.4	1.6	2.6	4.0	5.1
stdev	1.6	5.0	5.8	1.5	4.3	5.8	7.8
r3	4.1	10.9	3.8	3.0	5.3	2.9	2.4
stdev	2.3	4.8	1.9	2.7	5.2	2.7	2.1
r4	1.6	14.3	6.5	1.5	11.9	4.5	5.3
stdev	1.8	8.9	4.1	2.2	6.8	3.7	8.2
i1	11.6	9.9	9.9	6.1	6.8	9.2	9.1
stdev	6.1	5.9	8.0	3.4	5.2	7.2	11.6
i2	9.5	8.6	4.6	5.8	6.3	6.7	9.9
stdev	9.1	5.4	3.3	5.3	3.4	5.3	12.9

TrpA1/V1$^{-/-}$ Cit Acid	DM	DI	DL	VM	VI	VL	DMSp5
r1	3.7	5.0	2.3	2.4	2.8	1.7	3.4
stdev	2.8	2.0	1.1	0.8	2.4	1.7	2.2
r2	5.9	13.4	5.3	6.4	6.6	4.4	5.2
stdev	3.2	5.8	3.1	5.6	4.2	2.9	5.2
r3	6.2	17.2	8.3	6.1	19.0	11.0	6.3
stdev	3.9	8.0	5.3	4.4	9.2	8.7	4.7
r4	9.5	23.1	13.7	7.0	24.8	13.4	10.9
stdev	7.6	8.4	7.7	2.7	7.8	6.2	5.5
i1	10.6	22.9	18.0	5.8	19.2	27.9	17.3
stdev	5.3	18.3	9.2	3.2	9.8	12.0	10.9
i2	10.0	12.3	7.5	8.3	15.1	22.6	17.3
stdev	5.1	17.4	8.6	3.9	6.9	8.9	13.1

References

Arai, T., T. Ohkuri, K. Yasumatsu, T. Kaga and Y. Ninomiya (2010). "The role of transient
receptor potential vanilloid-1 on neural responses to acids by the chorda tympani,
glossopharyngeal and superior laryngeal nerves in mice." Neuroscience **165**(4): 1476-1489.

Bautista, D. M., S. E. Jordt, T. Nikai, P. R. Tsuruda, A. J. Read, J. Poblete, E. N. Yamoah, A. I.
Basbaum and D. Julius (2006). "TRPA1 mediates the inflammatory actions of environmental
irritants and proalgesic agents." Cell **124**(6): 1269-1282.

Bessou, P. and E. R. Perl (1969). "Response of cutaneous sensory units with unmyelinated fibers
to noxious stimuli." J Neurophysiol **32**(6): 1025-1043. .

Caterina, M. J., A. Leffler, A. B. Malmberg, W. J. Martin, J. Trafton, K. R. Petersen-Zeitz, M.
Koltzenburg, A. I. Basbaum and D. Julius (2000). "Impaired nociception and pain sensation in
mice lacking the capsaicin receptor." Science **288**(5464): 306-313.

Corson, J., A. Aldridge, K. Wilmoth and A. Erisir (2012). "A survey of oral cavity afferents to
the rat nucleus tractus solitarii." J Comp Neurol **520**(3): 495-527.

Dessirier, J. M., M. O'Mahony, M. Iodi-Carstens and E. Carstens (2000). "Sensory properties of
citric acid: psychophysical evidence for sensitization, self-desensitization, cross-desensitization
and cross-stimulus-induced recovery following capsaicin." Chem Senses **25**(6): 769-780.

Deval, E., X. Gasull, J. Noel, M. Salinas, A. Baron, S. Diochot and E. Lingueglia (2010). "Acid-
sensing ion channels (ASICs): pharmacology and implication in pain." Pharmacol Ther **128**(3):
549-558.

Finger, T. E., V. Danilova, J. Barrows, D. L. Bartel, A. J. Vigers, L. Stone, G. Hellekant and S.
C. Kinnamon (2005). "ATP signaling is crucial for communication from taste buds to gustatory
nerves." Science **310**(5753): 1495-1499.

Gerhold, K. A. and D. M. Bautista (2008). "TRPA1: irritant detector of the airways." J Physiol
586(14): 3303.

Hallock, R. M., M. Tatangelo, J. Barrows and T. E. Finger (2009). "Residual chemosensory
capabilities in double P2X2/P2X3 purinergic receptor null mice: intraoral or postingestive
detection?" Chem Senses **34**(9): 799-808.

Kichko, T. I., W. Neuhuber, G. Kobal and P. W. Reeh (2018). "The roles of TRPV1, TRPA1 and
TRPM8 channels in chemical and thermal sensitivity of the mouse oral mucosa." Eur J Neurosci
47(3): 201-210.

Nguyen, M. Q., Y. Wu, L. S. Bonilla, L. J. von Buchholtz and N. J. P. Ryba (2017). "Diversity
amongst trigeminal neurons revealed by high throughput single cell sequencing." PLoS One
12(9): e0185543.

Ohkuri, T., N. Horio, J. M. Stratford, T. E. Finger and Y. Ninomiya (2012). "Residual
chemoresponsiveness to acids in the superior laryngeal nerve in "taste-blind" (P2X2/P2X3
double-KO) mice." Chem Senses **37**(6): 523-532.

Ohman-Gault, L., T. Huang and R. Krimm (2017). "The transcription factor Phox2b
distinguishes between oral and non-oral sensory neurons in the geniculate ganglion." J Comp
Neurol **525**(18): 3935-3950.

Price, M. P., S. L. McIlwrath, J. Xie, C. Cheng, J. Qiao, D. E. Tarr, K. A. Sluka, T. J. Brennan,
G. R. Lewin and M. J. Welsh (2001). "The DRASIC cation channel contributes to the detection
of cutaneous touch and acid stimuli in mice." Neuron **32**(6): 1071-1083.

Richter, T. A., A. Caicedo and S. D. Roper (2003). "Sour taste stimuli evoke Ca2+ and pH
responses in mouse taste cells." J Physiol **547**(Pt 2): 475-483.

Silverman, J. D. and L. Kruger (1990). "Analysis of taste bud innervation based on
glycoconjugate and peptide neuronal markers." J Comp Neurol **292**(4): 575-584.

Stratford, J. M. and T. E. Finger (2011). "Central representation of postingestive chemosensory
cues in mice that lack the ability to taste." J Neurosci **31**(25): 9101-9110.

Stratford, J. M. and J. A. Thompson (2016). "MSG-Evoked c-Fos Activity in the Nucleus of the
Solitary Tract Is Dependent upon Fluid Delivery and Stimulation Parameters." Chemical Senses
41(3): 211-220.

Stratford, J. M., J. A. Thompson and T. E. Finger (2017). "Immunocytochemical organization
and sour taste activation in the rostral nucleus of the solitary tract of mice." J Comp Neurol
525(2): 271-290.

Sudo, S., M. Sudo, C. T. Simons, J. M. Dessirier, M. Iodi Carstens and E. Carstens (2003).
"Activation of neurons in trigeminal caudalis by noxious oral acidic or salt stimuli is not reduced
by amiloride." Brain Res **969**(1-2): 237-243.

Teng, B., C. E. Wilson, Y. H. Tu, N. R. Joshi, S. C. Kinnamon and E. R. Liman (2019). "Cellular
and Neural Responses to Sour Stimuli Require the Proton Channel Otop1." Curr Biol.

Tominaga, M., M. J. Caterina, A. B. Malmberg, T. A. Rosen, H. Gilbert, K. Skinner, B. E.
Raumann, A. I. Basbaum and D. Julius (1998). "The cloned capsaicin receptor integrates
multiple pain-producing stimuli." Neuron **21**(3): 531-543.

Vandenbeuch, A., C. B. Anderson, A. P. Ford, S. Smith, T. E. Finger and S. C. Kinnamon
(2013). "A selective P2X3, P2X2/3 receptor antagonist abolishes responses to all taste stimuli
in mice." Chem Senses **38**(Supp): 86.

Wang, Y. Y., R. B. Chang, S. D. Allgood, W. L. Silver and E. R. Liman (2011). "A TRPA1-
dependent mechanism for the pungent sensation of weak acids." J gen physiol **137**(6): 493-505.

Wu, P., D. Arris, M. Grayson, C. N. Hung and S. Ruparel (2018). "Characterization of sensory
neuronal subtypes innervating mouse tongue." <u>PLoS One</u> **13**(11): e0207069.

Zhang, J., H. Jin, W. Zhang, C. Ding, S. O'Keeffe, M. Ye and C. S. Zuker (2019). "Sour Sensing
from the Tongue to the Brain." <u>Cell</u> **179**(2): 392-402 e315.

452