Transcriptomic Profiling of Human Pluripotent Stem Cell- Derived Retinal Pigment Epithelium Over Time

Grace E. Lidgerwood [1,2,3,*], Anne Senabouth[4], Casey J.A. Smith-Anttila[5], Vikkitharan Gnanasambandapillai[4], Dominik C. Kaczorowski[4], Daniela Amann-Zalcenstein[5], Erica L. Fletcher[1], Shalin H. Naik[5,6], Alex W. Hewitt [2,3,7,#], Joseph E. Powell [4,8,#], Alice Pébay [1,2,3,#,*].

[1]Department of Anatomy and Neuroscience, The University of Melbourne, Parkville, VIC 3010, Australia

[2]Department of Surgery, The University of Melbourne, Parkville, VIC 3010, Australia

[3] Centre for Eye Research Australia, Royal Victorian Eye and Ear Hospital, East Melbourne, VIC 3002, Australia

[4] Garvan Weizmann Centre for Cellular Genomics, Garvan Institute of Medical Research, The Kinghorn Cancer Centre, Darlinghurst, NSW 2010, Australia

[5] Single Cell Open Research Endeavour (SCORE), The Walter and Eliza Hall Institute of Medical Research, Parkville, VIC 3052, Australia

[6] Immunology Division, The Walter and Eliza Hall Institute of Medical Research, Parkville, VIC 3052, Australia

[7] School of Medicine, Menzies Institute for Medical Research, University of Tasmania, Hobart, TAS 7005, Australia

[8] University of New South Wales Cellular Genomics Futures Institute, School of Medical Sciences, University of New South Wales, Sydney, NSW 2052, Australia

[#] Equal senior authors

*Corresponding authors: grace.lidgerwood@unimelb.edu.au (Lidgerwood GE), apebay@unimelb.edu.au (Pébay A).

Running Title *Lidgerwood et al/ Transcriptomic Profile of hPSC-derived RPE*

Total counts. Words: 8,203; References: 82; Figures: 8; Tables: 0; Supplementary Figures: 4; Supplementary Tables: 7; title: 100 characters; running title: 61 characters; keywords: 5; abstract: 248 words.

Abstract

Human pluripotent stem cell (hPSC)-derived progenies are immature versions of cells, presenting a potential limitation to the accurate modelling of disease associated with maturity or age. Hence, it is important to characterise how closely cells used in culture resemble their native counterparts. In order to select appropriate points in time for RPE cultures to reflect native counterparts, we characterised the transcriptomic profiles of hPSC-derived retinal pigment epithelium (RPE) cells from 1- and 12-month cultures. We differentiated the human embryonic stem cell line H9 into RPE cells, performed single cell RNA-sequencing of a total of 16,576 cells, and analysed the resulting data to assess the molecular changes of RPE cells across these two culture time points. Our results indicate the stability of the RPE transcriptomic signature, with no evidence of an epithelial – mesenchymal transition, and with maturing populations of RPE observed with time in culture. Assessment of gene ontology pathways revealed that as cultures age, RPE cells upregulate expression of genes involved in metal binding and antioxidant functions. This might reflect an increased ability to handle oxidative stress as cells mature. Comparison with native human RPE data confirmed a maturing transcriptional profile of RPE cells in culture. These results suggest that *in vitro* long-term culture of RPE cells allow the modelling of specific phenotypes observed in native mature tissue. Our work highlights the transcriptional landscape of hPSC-derived RPE as they age in culture, which provides a reference for native and patient- samples to be benchmarked against.

KEYWORDS: Human Embryonic Stem Cells; Human Pluripotent Stem Cells; Retinal Pigment Epithelium; Single Cell RNA Sequencing; Aging.

Introduction

The retinal pigment epithelium (RPE) is a monolayer of post-mitotic, pigmented polarized cells that is key to the health and function of photoreceptors and underlying vasculature. In particular, the RPE protects the retina against photo-oxidation and phagocytoses photoreceptor outer segments. The RPE is also essential to the immune privilege of the eye, as it physically contributes to the blood retina barrier and also expresses molecules repressing the migration of immune cells in the retina [1]. In the human retina, aging is associated with vision decline and delayed dark adaptation, both of which are direct consequences of tissue stress and retinal damage [2]. It is hypothesized that over time, oxidative stress leads to the death of retinal neurons; a decrease in RPE numbers; an accumulation of the toxic waste lipofuscin within the RPE; and an accumulation of basal toxic deposits called drusen underneath the RPE [2]. Together, these events contribute to a loss of homeostasis and low-grade inflammation within the retina [2]. Although it is clear that the RPE is key to the health of the retina, the precise molecular mechanisms responsible for its aging are not well understood.

Human pluripotent stem cells (hPSCs) have the ability to propagate indefinitely *in vitro* and give rise to any cell type in the body, including cells that form the retina. Various protocols have been described to differentiate hPSCs into RPE cells [3-8]. RPE cells are generally assayed after a few weeks of differentiation, at which stage they demonstrate similarity to their human native counterparts, in terms of morphology/expression of key proteins, functions and expression profiles, however with a profile closer to a foetal stage than adult stage [4, 9, 10]. Interestingly, the transcriptome profile of hPSC-derived RPE cells as they age in culture is unknown. To date, most RNA-sequencing (RNA-seq) studies of RPE cells have been performed on bulk samples. Yet, the ability to sequence individual cells provides a powerful tool to precisely uncover potential heterogeneity in cell population, especially as those develop and mature *in vitro*. Here, we used single cell RNA-seq (scRNA-seq) of hPSC-derived RPE cells maintained in culture for 1 and 12 months to assess the impact of time on the RPE transcriptome and whether genetic hallmarks of maturation can be observed over time. A short time of differentiation (1 month) was chosen as it represents a time routinely used in RPE *in vitro* assays [3]. A prolonged time of differentiation (12 months) was chosen as its characterisation could be subsequently used for comparison with other retinal cell differentiation methods, in particular of retinal organoids and

82   photoreceptors, for which differentiation and relative maturity are obtained after prolonged time

83   in culture and would thus be present at that later time point [11-14].

Results

Single cell RNA-sequencing profiled the transcriptomes of 16,576 cells

The human embryonic stem cell line H9 was differentiated to RPE following the protocol

described in the methods section. RPE cells from the same culture and original passaging were

isolated after 1 and 12 months of differentiation, dissociated to single cells and processed to

generate libraries for scRNA-seq analysis, in order to generate a transcriptional map of the RPE

cells reflecting time in culture (**Figure 1A**). The capture of the 1-month single cell library

detected 12,873 cells at the mean read depth of 40,499 reads per cell, while the 12-month capture

detected 4,850 cells at the mean read depth of 114,503 reads per cell (Supplementary Table 1).

Both datasets underwent cell-specific quality control, where 510 cells and 637 cells were

removed from the 1-month and 12-month datasets respectively. The remaining 16,576 cells were

retained for further analysis.

We compared variations in the transcriptomic profiles between the 1-month-old and 12-month-

old samples (**Figure 1B, C**) to identify potential changes in phenotypes upon aging of RPE cells

in vitro, analysing differential expression. A range of RPE markers were observed as conserved

between both points in time (**Figure 1D**). In particular, canonical RPE markers [15] associated

with extracellular structure organization (*CST3*, *EFEMP1*, *ITGAV, CRISPLD1*, *ITGB8*), melanin

biosynthesis (*PMEL*, *TTR*, *TYRP1*, *TYR*, *DCT*), lipid biosynthesis (*PTGDS*, *INPP5K),* visual

cycle (*LRAT, PLTP, ABHD2/RLBP1*, *RPE65*, *RGR*, *RPB1*, *BEST1*), and secretion (*SERPINF1*)

were present at both time points (Figure 1D).

Cluster analysis highlights twelve subpopulations of RPE cells

Cluster analysis was performed independently and identified 12 subpopulations in each sample

(Figure 1C, Supplementary Table 2). After the data were integrated with anchors identified with

a method described by [16], MetaNeighbor was used to match common subpopulations across

both samples [17], denoted as "Common". Clusters unique to 1-month and 12-month samples

were respectively denoted as "Young" and "Aged" (Figure 1C). In total, 18 subpopulations were

identified with six common subpopulations ("Common"; 8,484 cells; Supplementary Table 2),

six subpopulations exclusive to the 1-month dataset ("Young"; 5,758 cells; Supplementary Table

2) and six subpopulations exclusive to the 12-month dataset ("Aged"; 2,334 cells;

Supplementary Table 2). Cell counts per cluster (Figure 1C, Supplementary Table 2) and the top

conserved markers for each distinct cluster (**Figure 2A**, Supplementary Tables 3-5) were

identified. Clusters were visualised by Uniform Manifold Approximation and Projection

(UMAP) plot (Figure 1B). 3,070 cells were considered singletons as they had less connections

with similar cells (neighbors) relative to the rest of the cell population and they could not be

assigned to a subpopulation (cluster Zero in the 1-month dataset: 2,219 cells, 18% of all 1-

month-old cells) and cluster Zero in the 12-month dataset: 851 cells, 20% of all 12-month-old

cells, Supplementary Table 2). There were more singleton cells in the Young population as we

captured more cells from this group. We assessed the expression profile of genes characteristic

of progenitors (*BMP7*, *SOX4*) and canonical RPE genes [15] across all subpopulations, both in

terms of frequency and intensity of expression (**Figure 2B**, Supplementary Tables 3-5). Those

included genes linked to lipid biosynthesis (*INPP5K*, *PTGDS*); visual cycle (*LRAT, RGR*, *PLTP*,

RLBP1/CRALBP, RBP1/CRBP1, BEST1, *RPE65*); melanin biosynthesis (*DCT*, *TYRP1*, *TYR*,

TTR, *PMEL*); secretion (*ENPP2, VEGFA*, *SERPINF1*); phagocytic activity (*GULP1*); and

extracellular structure organization (*ITGAV*, *CRISPLD1*, *CST3*, *EFEMP1*). Most canonical RPE

genes were expressed across most populations, although many were found at lower levels in the

1-month-old cells, confirming the purity of the RPE cell cultures over time (Figure 2B). As these

transcripts are associated with stages of RPE maturity, our data suggests that all subpopulations

are of RPE lineage, potentially at various stages of differentiation and maturation. Differential

gene expression analysis and pathway enrichment were performed to characterise the molecular

signature of these subpopulations.

Most cells share common transcriptomic profiles suggestive of maturing RPE cells.

Of the cells examined, more than half (8,484 cells) were clustered into six "Common"

subpopulations that intersect the 1-month-old and 12-month-old cell cultures (Figure 1B, C),

indicating a large shared transcriptional profile between the two conditions. Some commonalities

and differences were observed between the "Common" samples arising from the 1-month-old

and 12-month-old cultures. A range of RPE markers was observed as conserved between both
points in time (Figure 2B). In particular, RPE markers associated with melanin biosynthesis
(*MITF, PMEL, TTR, TYR, TYRP1, DCT*), extracellular structure organization (*EFEMP1, CST3,*
CRISPLD1, ITGAV, ITGB8), secretion (*SERPINF1, VEGFA*), visual cycle (*RPE65, BEST1,*
RBP1, RLBP1, PLTP, RGR, LRAT), tight junctions (*TJP1*), phagocytic activity (*GULP1*) and
lipid biosynthesis (*PTGDS, CYP27A1, INPP5K, PLA2G16, PLCE1*) were conserved in all or in
some of the subpopulations (Supplementary Table 3). Expression of some genes related to RPE
maturity did not appear to differ between the 1-month-old and 12-month-old RPE, including
RBP1, TYRP1, and *SERPINF1,* suggesting some of the necessary retinoid-cycle binding proteins,
melanin biosynthesis and secretory proteins are expressed in early RPE development (Figure
2B). More heterogeneity was observed in the expression of RPE markers *RGR, PLTP, RLBP1,*
BEST1, ENPP2, VEGFA, and *TYR* in the 1-month-old RPE cells relative to the 12-month-old
RPE cells, where expression was generally high and stable (Figures 1D, 2B). Expressions of
LRAT and *RPE65* were predominantly observed in the 12-month-old RPE cells, with the
exception of small expression for RPE65 in the 1-month-old members of "Common"
Subpopulation Two. This suggests that the catabolic machinery for converting all-*trans*-retinol
into all-*trans*-retinyl ester (LRAT) and 11-*cis*-retinol (RPE65) for phototransduction is expressed
at low levels in the 1-month-old RPE samples but becomes more comprehensive as cells age in
culture (**Figure 2C, D**). Variations in the pattern of gene expressions were observed between
cells identified from the 1-month-old or 12-month-old cultures within each subpopulation
(**Figure 3A-C**). Genes involved in neural differentiation including *DCT, PAX6, SOX1* and *MDK*
exhibited notable differences between the 1-month-old and 12-month-old samples (Figure 3B),
as did genes involved in the extracellular matrix (ECM) formation and maintenance *CST3,*
EFEMP1, ITGAV and *CRISPLD1* (Figure 3C)*,* which were more heterogeneous in the 1-month-
old samples, suggesting transitional changes in ECM markers during early RPE differentiation.
Examples of genes characteristics of RPE, neural differentiation, and extracellular matrix are
illustrated in Figure 3A-C respectively.
"Common" Subpopulation One (2,282 cells) was characterised by 891 identified conserved
markers (p value cut-off of < 0.74) (Supplementary Table 3). Among the most highly conserved
markers was *NDUFA4L2*, a gene known to be associated with the macula retina [18]; the zinc
metalloenzyme gene *CA9*; and genes involved in pigment/melanin biosynthesis (*DCT, PMEL,*

MITF, *TYR,* and *TYRP1*). This subpopulation also expressed genes involved in early retinal

development including of the RPE and eye morphogenesis (*SOX4, EFEMP1*, *BMP7*, *VIM*,

GJA1, *PTN*) and in the retinoid cycle (*RPE65* and *RLBP1*). In addition, 79 ribosomal genes (32

RPS, 47 *RPL*) and 11 mitochondrially-encoded genes (*MT-*) were identified. Expression of

ribosomal genes and mitochondrially-encoded genes has been correlated with development and

maturation [19], including of the retina [20, 21]. This is supported by the GO analysis showing

an overrepresentation of pathways involved in mitochondrial and ribosomal functions; protein

biogenesis, transport, assembly and function and ATP biosynthesis and metabolism (**Figure 3D**,

Supplementary Table 3). Hence, together with the presence of RPE markers, the data suggests

this subpopulation comprises a highly metabolically active maturing RPE phenotype.

"Common" Subpopulation Two (2,204 cells) identified 1,077 conserved markers. Its 15 most

conserved markers were all known RPE markers. Many of the markers expressed are involved in

the generation of RPE or in their maturation and homeostasis. For instance, Cystatin C (encoded

by *CST3*) is abundantly produced by RPE cells [22] and its secretion diminishes with age [23].

DCT is expressed in the developing retina [15] and is important for the production of melanin

and to the RPE homeostasis [24, 25]. Its downregulation is associated with mature native RPE

[26]. We thus suggest this subpopulation is a mature functional RPE population.

"Common" Subpopulation Three (1,811 cells) identified 1,226 conserved markers. Its most

conserved markers were mostly known to be expressed in the RPE (*DCT*, *TTR*, *CST3*, *AQP1*,

FTH1 [27], *BEST1*), with some markers, such as *TFPI2*, known to promote survival and

maintenance of RPE cells [28]. The markers were similar to those of the "Common"

Subpopulation Two. Other markers identified are not necessarily RPE-specific such as *GNGT1*

which encodes for a protein found in rod outer segments could suggest that cells are not yet fully

committed. Together, those markers indicate cellular functions suggestive of functional and

maturing RPE cells.

"Common" Subpopulation Four (1,170 cells) identified 1,421 conserved markers. Many of

the most conserved markers of this subpopulation were not specifically linked to the RPE. For

instance, *TMSB4X* is linked to the cytoplasmic sequestering of NF-kB but has not yet been

associated to molecular events in RPE cells, whilst many others are associated with the

cytoskeleton, such as *TAGLN*, *TNNC1*, *CALD1*, *MYLK*, *TPM1*, *ACTA2*, *MYL9*. On the other

hand, the presence of markers known to be expressed by the RPE (including *TTR*, *BEST1*, *CST3*,

CSTV [29], *CRYAB*, *SERPINF1*, *PMEL, VEGFA*, *RBP1, RLBP1, TYR, TYRP1*) confirms the RPE
identity of the subpopulation. The presence of genes associated with early differentiation, such as
IGFBP5 (downregulated relative to other clusters, as observed upon RPE differentiation [30])
and *CRB2* [31], involved in RPE polarity [32], suggests an early stage of RPE maturity. This
pattern is suggestive of a differentiating RPE population.

"Common" Subpopulation Five (630 cells; 907 conserved markers) was characterised by
conserved RPE markers (*SERPINE2* [33], *SFRP5* [34]) with many being downregulated when
compared to the other populations (*AQP1*, *CST3, PTGDS*, *SERPINF1*, *BEST1*, *SMOC2*). Other
genes indicate an immaturity/ differentiation or proliferation of cells, such as *GAP43*, *DAAM1*,
CD44 [35], and *DUSP4*. Together, this suggests this subpopulation comprises cells in early
differentiation to RPE.

"Common" Subpopulation Six (387 cells; 1,664 conserved markers) was characterised by
markers associated with retinal cell types other than RPE [36] (such as *SPP1*, *CPODXL*, *STAC2*,
PCDH9) or found at low levels in the RPE (such as *CPAMD8* and *SFRP2* [37]). Only the marker
CRABP1 was highly conserved and upregulated, whilst other RPE markers were downregulated
(*SERPINF1*, *BEST1*, *RLBP1* or *RPE65*). This thus suggests a subpopulation of immature cells.

Finally, the analysis of the PANTHER GO-Slim Biological Processes conserved within the
"Common" subpopulations identified pathways that were predominantly involved in
mitochondrial, metabolic and ribosomal processes as well as purine biosynthesis, nucleotide
metabolism and protein biogenesis, localisation and transport (Figure 3D, Supplementary Table
3). Altogether, this data suggests that the common population to both time points is
heterogenous, with subpopulations representing different stages of RPE cell differentiation.

227 The Young Subpopulation is characterised by immature and differentiating cells.
Around a third of all cells (5,625 cells) clustered into six "Young" subpopulations
(Supplementary Table 4). Common RPE markers were conserved within a few "Young"
subpopulations (Figure 2A, B, Supplementary Table 4). These were associated with lipid
biosynthesis (*PTGDS*), visual cycle (*RGR*, *RLBP1, RBP1*, *BEST1*), melanin biosynthesis (*DCT*,
TYRP1, *TYR*, *TTR*, *PMEL*), phagocytic activity (*GULP1*), secretion (*ENPP2, VEGFA*,
SERPINF1), and extracellular structure organization (*ITGAV, CRISPLD1*, *CST3*, *EFEMP1*,
ITGB8).

"Young" Cluster Zero (2,219 cells; 20 conserved markers) was composed of singleton cells
which showed expressions of *SERPINF1*, *CST3*, *DCT*, *TSC22D4, IGFBP5,* and *RNASE1*. Most
have been reported in the retina (Courtesy of Human Protein Atlas, www.proteinatlas.org) [36]
and in the human RPE (*IGFBP5* [30]). PANTHER GO (biological process) analysis indicates
regulation of neuroblast proliferation, suggestive of progenitor cells (Supplementary Table 4).

"Young" Subpopulation One (1,873 cells; 58 conserved markers) was characterised by gene
expression of *CTNNB1*, *NOG*, *ATP1B1*, *GSTP1*, *CD63*, *HNRNPH1*. All play fundamental roles
for the homeostasis and functions of the RPE: *ATP1B1* encodes an apical Na^+/K^+ ATPase which
expression reduces with age and in AMD [38]; defects in *CTNNB1* are linked to abnormalities in
RPE development and pigmentation [39]; GSTP1 is a survival factor for RPE cells, which levels
increase as cells mature [40]; CD63 is a late endosome/exosome marker known to be released by
RPE cells [41] and *HNRNPH1* levels are associated with improved survival of RPE cells in
culture [42]. Hence, those markers indicate cellular functions suggestive of functional RPE cells.
It is interesting to note that known canonical RPE markers, such as *SERPINF1*, *RLBP1*, *TTR*,
PMEL, *CRYAB* were less expressed in this subpopulation than in all other subpopulations.
PANTHER GO (Biological Process) analysis highlights that this population is highly
metabolically active (Supplementary Table 4). This is suggestive of an earlier stage of
maturation of the RPE cells.

"Young" Subpopulation Two (144 cells; 16 conserved markers) was characterised by the
specific upregulation of genes including *CCL2*, *SFRP1* and *B2M*, which are implicated in the
negative regulation of epithelial cell proliferation. Downregulation of *CTSV* and *TMSB4X* -
linked to the cytoplasmic sequestering of NF-κB but not known to have associated molecular
events in RPE cells - together with the expression of *DCT*, known to be expressed during RPE
development [43], suggests an immature RPE population. This is supported by PANTHER GO
(Biological Process) analysis which highlights the regulation of differentiation and NF-kB
sequestration as enriched in this subpopulation (Supplementary Table 4).

"Young" Subpopulation Three (67 cells; 369 conserved markers) was characterised by the
expression of genes such as *TOP2A*, *PCLAF*, *PTTG1*, *ANLN*, *MKI67*, *RRM2*, *TPX2*, *PBK*.
Although all genes are found to be expressed in the retina [36], none are associated with a
specific RPE signature. However, they are associated with cell proliferation (*TOP2A*, *PCLAF,*
PTTG1, MKI67, *RRM2*, *TPX2*, *PBK)* and cellular rearrangements (*ANLN*), which have been

described as characteristics of immature RPE cells [44]. In particular, ANLN is reported to
promote maturity of intercellular adhesions (tight junctions and adherens junctions) in epithelial
cells [45]. TOP2A is associated with retinal development and proliferation [46], which combined
with expression of *PCLAF, PTTG1* and *MKI67* suggests a proliferating cell population. Low
expression of RPE markers (*RBP1*, *ENPP2, CRABP1*, *HNRNPH1*) further suggests the RPE
identity of the developing retinal cell subpopulation. Altogether, this expression profile suggests
an immature differentiating cell population.

Similarly, "Young" Subpopulation Four (871 cells; 49 conserved markers) was characterised
by expression of genes that are not traditionally associated with the RPE identity. The expression
of *CRYAB* (downregulated), *CRX*, *FTH1*, *TFPI2* (known to promote survival and maintenance of
RPE cells [28]) and *DCT*, expressed in the native RPE, suggests a differentiation to RPE, yet the
presence of the photoreceptor specific *GNGT1* could suggest an early differentiation step where
cells are not yet fully committed.

Interestingly, "Young" Subpopulation Five (584 cells; 99 conserved markers) displayed a
signature comprising mitochondrial and ribosomal -related transcripts with 9 mitochondrial-
related genes (MT-) and 14 ribosomal genes (RPS- or RPL-) marking this subpopulation. These
genes are ubiquitous and have not been specifically correlated to the retina or the RPE, however
they are known to facilitate fundamental processes of biology, including electron transfer and
energy provision, ribosome biogenesis and protein synthesis. PANTHER GO-Slim Biological
Process and GO (Biological Process) analysis confirmed that the conserved pathways within this
subpopulation are mostly related to the mitochondrial energetic metabolism, and nucleotide
metabolism (Supplementary Table 4). This subpopulation also expressed RPE markers, such as
BEST1, *VEGFA*, *ENPP2*, *TIMP3*, *TYRP1* as well as genes involved in early retinal development
including of the RPE and eye morphogenesis (*SOX11, PMEL*, *EFEMP1*, *BMP7*, *VIM*, *GJA1*,
PTN) [15, 47]. Hence the presence of high level of expression of mitochondrial-encoded genes
and ribosomal-related genes in RPE are suggestive of highly active cells with high protein
synthesis, which may imply this is a maturing RPE population.

Taken together, our results suggest that all subpopulations within the "Young" cohort are
immature cells developing to RPE cells.

The Aged Subpopulation is characterised by higher maturity of RPE cells

Less than 10% of all cells (2,334 cells) clustered into the "Aged" category, which comprised six subpopulations (Supplementary Table 5). All six identified "Aged" subpopulations were subjected to the same analysis; however, a statistical overrepresentation test returned no positive results for "Aged" Subpopulations Zero, One and Five (most likely owing to the low number of conserved genes identified). Only a few common RPE markers associated with lipid biosynthesis (*INPP5K*), visual cycle (*RLBP1*), melanin biosynthesis (*TTR*, *DCT*), secretion (*SERPINF1*, *VEGFA*), and extracellular structure organization (*CST3*, *CRISPLD1*), were conserved in some of the "Aged" subpopulations (Figure 2, Supplementary Table 5).

"Aged" Cluster Zero (851 cells; 6 conserved markers) was composed of singleton cells with cells expressing *CRISPLD1*, *PCCA*, *WFDC1*, *TTR*, *SH3BGRL3* and *TMSB4X* yet with lower average levels of expression per cell than those observed in all other subpopulations. Some of these genes have been found to be expressed in the RPE (*CRISPLD1*, *WFDC1* [48], *TTR*, *TMSB4X* [43]) and are associated with late RPE development (*CRISPLD1*, *TTR* [15]), whilst others have a wider expression pattern (*PCCA*, *SH3BGRL3*) and encode for proteins involved in more universal cellular events: mitochondrial protein PCCA plays roles in death/survival, SH3BGRL3 is involved in oxidoreduction, whilst *TMSB4X* encodes proteins of the cytoskeleton. Likewise, the "Aged" Subpopulation One (815 cells; 12 conserved markers) presented more cells expressing *HSD17B2*, *TPM1, MYL9, NDUFA4L2*, *BNIP3*, *CALD1*, *TTR*, *DCT*, *MT-CYB*, *FTH1*, *CRYAB* and *TMSB4X* but at lower average levels per cell than in all other subpopulations. Nine out of the twelve genes are known to be expressed in the RPE (*TTR*, *TMSB4X* [43], *CALD1* [49], *DCT* [43], *HSD17B2* [50], *NDUFA4L2* [15], *BNIP3* [27], *FTH1* [27], *CRYAB*). Some of these markers are associated with late RPE development (*TTR* [15]) whilst others are associated with a geographic localisation of cells within the retina (*HSD17B2* [51], *NDUFA4L2* [18]). *BNIP3* and *NDUFA4L2* encode for mitochondrial proteins playing roles in death/survival and electron transport, whilst *TPM1*, *MYL9*, *TMSB4X*, *CALD1* encode proteins of the cytoskeleton. A similar pattern of expression was observed in "Aged" Subpopulation Five (170 cells; 6 conserved markers) with *PCCA*, *WFDC1*, *SERPINE2*, *TMSB4X*, *TTR* being expressed in more cells with lower average levels than compared to all other subpopulations. *SPON2* was expressed in more cells and at higher levels and encodes for extracellular matrix proteins important for cell

adhesion. The close similarity of these three "Aged" subpopulations suggest a late RPE
phenotype, with higher levels of maturation, towards regionalisation of cells.

Similarly, "Aged" Subpopulations Two (46 cells; 22 conserved markers), Three (32 cells;
121 conserved markers) and Four (420 cells; 34 conserved markers) showed close similarities in
terms of gene expression profiles. The presence of known RPE markers, such as *DCT*
(Subpopulation Two, downregulated), *CALD1* (Subpopulation Two, downregulated), *TTR*
(Subpopulation Three, downregulated; Subpopulation Four), *SOX9* (Subpopulation Three,
downregulated), *RBP1* (Subpopulation Three, downregulated; Subpopulation Four), *SERPINF1*
(Subpopulation Three, downregulated), *VEGFA* (Subpopulation Four, downregulated), *TMSB4X*
(downregulated in all three subpopulations), *CRYAB* (downregulated in Subpopulations Two and
Three) confirms their RPE identity.

PANTHER GO- Slim Biological Process (Cluster Two) and PANTHER GO (Biological
Process) analysis (Clusters Two, Three and Four) were performed. "Aged" Subpopulations Two
and Four were similar, with high significance in pathways associated with response to metal ions
(particularly cadmium, copper, iron and zinc); response to stress, chemicals and toxins; and
neural crest fate commitment, which was the most significantly identified biological process in
the "Aged" Subpopulation Three, with a 92.9-fold enrichment. Interestingly, GO analysis
indicated that nuclear genes encoding metallothioneins (which are involved in metal binding), in
particular of zinc (*MT1E*, *MT1F*, *MT1G*, *MT2A*, *MT1X*), metal iron (*MT1E*, *MT1F, MT1G*, *DCT*,
MT2A, *MT1X*) and copper iron (*DCT*) as well as in oxidoreduction (*DCT*) were significantly
differentially expressed between the "Aged" subpopulations compared to all others, with an
overall increased expression per cell as cells age (Supplementary Table 5). Further analysis of
the metallothioneins *MT1E*, *MT1F, MT1G, MT1X* and *MT2A* across all subpopulations at the two
points in time confirmed a large increase in their mRNA expression in the 12-month-old culture
when compared to the 1-month-old cells (**Figure 4A, B**). On the other hand, *DCT* was reduced in
the 12-month-old culture compared to the 1-month-old cells, which is consistent with a mature
RPE profile [26] (Figure 4A, B). Altogether, this data suggests that the RPE cells of these
"Aged" subpopulations are increasing their handling of metals and antioxidant abilities, which
likely reflects a further maturation of the RPE cells.

RPE cells from different timepoints share common developmental trajectories

We investigated the pseudo-temporal transition of 1-month-old cells to 12-month-old cells using
trajectory analysis methods to identify trajectories originating from proliferative cells in
"Young" Subpopulation 3. *Monocle* 3 [52] identified a complex, branched development
trajectory that included cells from both timepoints (**Figure 5A, B**). Interestingly, the pseudo-
temporal ordering of cells across the trajectory did not correspond to timepoint (Supplementary
Figure 3A). To determine the nature of the trajectories, we used Moran's I test to identify 158
genes that were significantly associated with pseudotime (Supplementary Table 6). These genes
formed four co-expression modules that were further characterised with STRING
(Supplementary Table 6, Supplementary Figure 3B-F). As illustrated in Supplementary Figure 3,
Module One revealed genes whose expressions are involved in development (red) and Module
Four revealed genes involved in cell cycle (blue) and mitosis (red). The two other modules did
not show clear biological processes associations but also showed genes associated with neural
development. The differential expression of the residual gene trajectory- for the four modules
confirmed differences of the various clusters. In particular, all young subpopulations except
"Young Four" were found to show the highest expressions of genes of the proliferative Module
Four (Supplementary Figure 3B), confirming their immature proliferative nature.

Differentiated RPE cells are post-mitotic and fully committed to the RPE lineage, in large
part due to their cell-cell contact inhibition (see reviews by [53, 54]). The expression of the
proliferation marker *MKI67* and of other genes associated with proliferation (*TOP2A*, *TPX2,*
PTTG1, *PCLAF, RRM2*) was monitored to assess whether some progenitor populations remained
with proliferative potentials, and if those vary over time in culture. Our data indicates that only a
small portion of cells remained proliferative over time in culture. In particular, *MKI67* was
almost uniquely expressed in the immature "Young" Subpopulation 3, together with other cell
proliferation markers (*TOP2A*, *TPX2, PTTG1*, *PCLAF* and *RRM2*, **Figure 5C**, Supplementary
Table 4). Very few cells of the 12-month-old population were found to express markers
associated with proliferation (Figure 5C). This confirms that only a small subpopulation of
progenitor/ immature cells present early in culture is proliferative and disappears with time in
culture.

Pseudotime analysis of early retinal markers (*PAX6*, *RAX*) and RPE genes (*PMEL*, *RLPB1*,
RGR, *TYR, RBP1*, *RPE65*) were performed in all subpopulations to measure gene expression
trajectories (**Figure 5D, E**; Supplementary Figure 4). The early retinal markers *PAX* and *RAX,*

were expressed in similar manners across subpopulations, with minor variations between samples, with the exception of the "Aged Subpopulation Three" which showed higher levels of *PAX6* and absence of expression of *RAX* (Figure 5D, E; Supplementary Figure 4). The marker of pigmentation *TYR* was consistently expressed across all populations, whilst *PMEL* was downregulated in all aged subpopulations and in Common Subpopulations Five and Six (Figure 5D, E; Supplementary Figure 4). Similarly, fairly comparable expression levels of *RGR*, *RLBP1*, *RBP1* were observed across all subpopulations, with the exception of Aged Subpopulations Three and Four which showed higher levels of expression (Figure 5D, E; Supplementary Figure 4). These variations in gene expression are likely indicative of RPE cell maturation in culture.

RPE subpopulations contribute to paracrine signalling

RPE cells can modulate immune responses and differentiation of other retinal cell types, in part via paracrine signalling. *In vitro*, this is observable through the expression of specific secretion factors and receptors, known to play roles in immune or developmental events. For instance, in the retina, the chemokine CCL2 is implicated in monocyte infiltration following damage [55] and its secretion by RPE cells contributes to the regulation of the immunological response to inflammation [56]. *CCL2* was found to be faintly expressed in cells at both points of time (**Figure 6A**). *CCL2* was identified as a marker of the immature "Young" Subpopulation Two (Supplementary Table 2); and was also expressed, albeit at lower levels, across other "Aged" and "Common" populations (Figure 6A). Unsurprisingly, its cognate receptor *CCR2*, was not expressed in any subpopulations as CCL2 is most likely targeting monocytes and not endogenous cells of the retina (data not shown). *CCL2* was more uniformly expressed in the 12-month-sample, suggesting more homogeneity in its expression as cells mature.

Secreted frizzled-related proteins (SFRPs) are secreted ligands that modulate WNT signalling, binding to the receptors Frizzled (FZD1-10) which are also receptors for WNT proteins (WNTs). SFRPs are considered WNT antagonists, binding WNTs and FZDs [60]. Cells in culture from both points in time express mRNA for genes involved in the WNT signalling pathways, in particular *SFRPs 1, 2, 5*; *FZDs 1, 8*; *WNTs 2B, 3*, *4*, and *5A*, with more expression homogeneity in the 12-month-old samples for many of these (**Figure 6B**). *SFRP1* was expressed across all subpopulations with varying levels and was identified as a marker of the immature "Young" Subpopulation Two (Figure 6B, Supplementary Table 4). *SFRP2* was identified as a

marker of the immature "Common" Subpopulation Six and where it was mainly expressed,

whilst *SFRP5* was identified as a marker of the immature "Common" Subpopulation Five and

expressed across most subpopulations (Figure 6B, Supplementary Table 3). It is interesting to

note that *SFRP1* and *SFRP5* expressions were both downregulated in the "Aged" Subpopulation

Three (relative to "Aged" cells in other populations), reflective of a dynamic expression pattern

of *SFRPs* by RPE cells in culture (Figure 6B, Supplementary Table 3). Interestingly, 1-month-

old RPE cells had low levels of expression of *FZD1, FZD8*, *WNT2B, WNT3, WNT4* and *WNT5A*

with all other *FZDs* and *WNTs* undetectable (Figure 6B). The expression of all detectible *SFRPs*,

FZDs and *WNTs* was higher in the 12-month-old RPE cells (Figure 6B). This dynamic profile of

expression of molecules involved in WNT signalling indicates that RPE cells can signal via the

WNT pathway. This correlates with the known role of WNT signalling in retinal development

including the RPE [34, 57].

PEDF is another factor key to various retinal cell differentiation, maturation and survival,

including RPE and photoreceptors [58]. It is encoded by *SERPINF1*, highly expressed by RPE

cells, and was found as a marker of subpopulations within the Young, Common, and Aged

groups (Figure 1D, Supplementary Tables 3-5). However, the PEDF receptor *PNPLA2* was not

found to be expressed in the RPE cells, indicating that PEDF secreted from RPE largely acts in a

paracrine manner. Likewise, the pleiotropic factor *VEGFA* [59], was also identified as a marker

of subpopulations within the three groups, but no VEGF receptor mRNA (*VEGFR1-3*) was

detected in the samples, again indicating that the RPE-secreted VEGF likely acts on

neighbouring cells to maintain homeostasis (**Figure 6C**, Supplementary Tables 3-5). These

examples of paracrine molecules being secreted by RPE subpopulations illustrate the important

role played by RPE cells in the shaping of a developing retina, releasing bioactive factors

regulating events such as cell fate, differentiation, polarity and maturity and contributing to the

regulation of the immune environment of the retina.

***APOE* is a conserved marker of various RPE subpopulations**

Apolipoproteins E (APOE) are proteins involved in lipid metabolism including cholesterol and

are also regulated with complement activation in the RPE [60]. APOEs interact with the low-

density lipoprotein (LDL) receptors (LDLRs) and very-low-density lipoprotein receptors

(VLDLRs). *APOE* was highly expressed in cells across both points in time in culture (**Figure 6D**). *APOE* was identified as a conserved marker of the "Young" subpopulations One and Five, as well as all "Common" Subpopulations (Figure 6D, Supplementary Tables 3, 4). The levels of expression per cell was however different between subpopulations, with an upregulation in the "Young" Subpopulation Five, and a downregulation in the "Young" Subpopulation One as well as in the "Common" subpopulations One, Two, Three and Six (Figure 6D). Interestingly, within the "Common" subpopulation Four, cells arising from the 1-month-old culture expressed less *APOE* than cells arising from 1-month culture of all other "Common" subpopulations whilst the opposite was observed with the 12-month-old cells within this subpopulation (Supplementary Table 3). The opposite pattern was observed in the "Common" Subpopulation Five (Supplementary Table 3). The APOE receptors *LDLR*, *LRP1*, *LRP2* and *LRP8* were detected in the cell cultures, with increased levels in the 12-month-old samples when compared to the 1-month-old culture (Figure 6D). Interestingly, these receptors were mainly absent from all Young Subpopulations, and the Aged subpopulations showed higher levels of all APOE receptor mRNAs (Figure 6D). This data illustrates that the expression of *APOE* and associated receptors are dynamic within RPE subpopulations and with time in culture.

The complement pathway is not modulated with time in culture

RPE cells express many complement components in various retinal diseases, inflammation and/or aging [61]. The complement regulators *C1s*, *C1r*, and *C1QBP* which form the C1 complex were conserved markers in a few "Young" and "Common" subpopulations (all in "Common" Subpopulation Three; *C1s* and *C1r* in "Young" Subpopulation Five and "Common" Subpopulations One and Five; *C1r* and *C1QBP* in "Common" Subpopulation Two; *C1s* "Common" Subpopulation Four; *C1r* only "Common" Subpopulations Two and Six; *C1QBP* only in "Common" Subpopulations Two, Three and Six) (Supplementary Tables 3, 4). Other genes associated with the complement response (such as *CFH*, *CFB*, *CFHR1*, *CFHR3*, *C3*) were not identified as markers of any subpopulation (Supplementary Tables 3-5). This suggests that the complement components are not modulated with time in culture.

RPE cells do not undergo Epithelial Mesenchymal Transition

Finally, it is interesting to note that no markers of Epithelial Mesenchymal Transition (such as
SNAI1, *SNAI2, ZEB1*, *TWIST1*, *GSC*) were characterised in any examined subpopulation
(Supplementary Tables 3-5), demonstrating the stability of the cell culture over time with no
evidence of a transition to a mesenchymal phenotype.

The RPE subpopulations express native RPE markers with different patterns.

We compared the hPSC-derived RPE signature to that of fetal native RPE cells, which was
described by scRNA-seq in [15]. As observed in foetal native tissue, most subpopulations of
hPSC-derived RPE cells expressed *SERPINF1*, *BEST1*, *TYR*, *TTR*, and *RPE65*, and the more
immature subpopulations also expressed *MKI67* and *DCT* (**Figure 7A, B**). Genes commonly
expressed in native RPE cells were enriched in 12-month-old subpopulations and in Aged
subpopulations (such as *RPE65*, *LRAT*, *PLTP*, *RGR, RLBP1*, *LRAT*, *INPP5K*, *ITGB8*, *EFEMP1*,
ITGAV, *GULP1* and *VEGFA*), whilst other genes were found at similar levels in 1-month- and
12-month- cultures, and in Young, Common and Aged groups, including *TYRP1* (**Figure 7C-H**).
Others were significantly less expressed in 12-month- cultures, such as *PMEL*, *PTGDS*, *CST3,*
CRISPLD1 (**Figure 7E-G**). Some common RPE genes are expressed in the adult native RPE at
much higher levels than in fetal RPE [26]. Those include *TTR, RPE65, BEST1, CHRNA3, RBP1,*
MYRIP, TFPI2, PTGDS, SERPINF1, *DUSP4 GEM* and *CRX*. Similarly, downregulation of *DCT*,
SFRP5, *SILV*, *TYRP1*, *SLC6A15* is associated with mature native RPE [26]. We thus compared
the expression profiles of these genes in the RPE cultures overtime, in order to assess the
maturity of the cultured cells [26] (**Figure 8**). *CHRNA3*, *MYRIP, TFPI2*, *GEM*, and *CRX* - which
are upregulated in the adult RPE- were more highly expressed in the Aged population (Figure 8
A, B). However, *RBP1*, *PTGDS*, *DUSP4*, genes also highly expressed in the adult RPE were not
found at higher levels in the Aged or 12-month-old cultures than in the Young, or 1-month-old
culture (Figures 7E, 8 A, B). *DCT*, *SFRP5*, *SILV, PMEL, TYRP1*, *SLC6A15*, were generally
downregulated in the Aged subpopulations and in the 12-month-old "Common" subpopulations
when compared to the 1-month-old "Common" subpopulations (Figures 7B, D, 8 C, D).

Discussion

Here, we provide a dynamic profile of the transcriptome of hPSC-derived RPE cells over 12

months. Our data confirms expression of markers of RPE homeostasis and functions in hESC-

derived RPE [15, 26, 62, 63] and provides novel information on the timing of expression of those

markers. At both early and late points in time, we observed that hPSC-derived RPE cells

expressed genes associated with lipid metabolism, secretion, visual cycle, melanin synthesis,

phagocytic activity, metal binding and oxidoreductase activity. Based on expression of genes

known to be associated with levels of RPE maturity and on PANTHER GO analysis, the

identified 18 subpopulations regroup into populations from immature and progenitor cells, to

maturing RPE cells and functionally mature RPE cells (based on genes involved in RPE

functions). Within those, some subpopulations were comprised of highly metabolically active

cells. The pseudo-temporal analysis could not identify trajectories corresponding with points in

time, which is likely due to the experimental design of two time points only. Outside the scope of

this study, but of interest, intermediate culture timepoints could provide information on the

pseudo-temporal ordering of cells across time in culture.

An essential function of the RPE is photoprotection of the retina, which is accomplished by

different mechanisms. Those include absorbing radiation, binding and sequestering redox-active

metals such as iron, and scavenging free radicals and reactive oxygen species [64].

Metallothioneins are metal-binding proteins that are protective against oxidative stress.

Compared to the 12-month-old RPE, the 1-month-old RPE cells express less transcripts for the

metallothioneins *MT1E*, *MT1F*, *MT1G*, *MT2A*, *MT1X* and higher levels of the Dopachrome

Tautomerase *DCT*. This data suggests a variation in the handling of metals and in the antioxidant

abilities of RPE cells, which could be reflective of either a necessity to handle more oxidative

stress in an aging *in vitro* environment or a maturation of RPE cells towards a more mature and

protective phenotype [64]. The assessment of variations in other gene expressions between the

two points in time indicates a maturation profile of cells rather than an increased stress. Indeed,

DCT is known to be expressed in human retinal progenitor cells [15] and its expression is

regulated by the early RPE marker MITF. It is thus not surprising that as RPE cells mature,

MITF expression reduces [65] and subsequently reduces *DCT* expression, as is observed in

human fetal retina [15]. Similarly, *CTNNB1* regulates *MITF* and *OTX2* expression and

subsequently RPE differentiation [39]. Finally, *SOX11* is known to be expressed in early retinal

progenitor and early in differentiating RPE cells [15, 47], hence its downregulation as cell

culture ages further supports a maturation of RPE cells in culture. The "Young" Subpopulations Zero and One are characterised by the expression of *CST3*, which encodes for the cysteine proteinase inhibitor Cystasin. Interestingly, this protein is known to decrease in native RPE cells with aging [23]. Its presence in the cell population further strengthens the suggestion of a maturing RPE population over time. The high level of expression of mitochondrial and ribosomal -related genes in some cell populations likely indicates that cells are metabolically and transcriptionally active, necessitating energy and ribosomal activities for protein synthesis respectively. It could also suggest that ribosomes potentially contribute to extra-ribosomal functions in the RPE cells, such as cell development and maturation [19], as already reported for melanocyte development [66], retinal development [20, 21] and retinal degeneration [67]. Of note, the absence of markers of Epithelial Mesenchymal Transition highlights the robustness and ability of RPE cells to maintain their phenotype *in vitro* despite prolonged culture times.

The expression analysis in the various cell populations and over time in culture of ligands and receptors involved in retinal development and homeostasis demonstrates a dynamic profile of gene expression of hPSC-derived RPE cells. This highlights the importance of RPE in the development and homeostasis of the retina and not solely of the RPE. For instance, WNT signalling is known to be important to the early stages of RPE differentiation [39, 68]. Some *SFRPs* and *WNTs* are expressed in RPE cell subpopulations and the receptors *FZDs* are very lowly expressed in most subpopulations, which suggests that the ligand expression could be directed to paracrine signalling, playing roles in the development and homeostasis of the other neighbouring retinal cells. For instance, *SFRPs 1, 2* and *5*, found to be expressed in the hPSC-derived RPE cells, are associated with retinal development [34, 57] and known to promote the differentiation of retinal ganglion cells and photoreceptors [69], and axon guidance [70]. Similarly, high levels of *PEDF/SERPINF1* and of *VEGF* were found across many subpopulations, yet their respective receptors *PNPLA2* and *VEGFRs* were not found to be expressed in the cells, suggestive of a paracrine signalling mechanism between RPE and neighbouring cells. This is in accordance with a role of these growth factors in retinal cell biology and survival, including of photoreceptors [58, 71], retinal ganglion cells [72] or beyond the retina on neighbouring endothelial cells. Interestingly, *APOE* was also expressed in cells across both points in time in culture, with variations observed between subpopulations. In the RPE, APOE is involved in lipid metabolism including cholesterol and drusen content [60].

APOE also plays a striking role in melanogenesis, regulating the formation of functional PMEL
amyloid fibrils in RPE cells [73]. Hence, the variations observed in its expression levels and that
of its receptors could possibly be indicators of melanogenesis within RPE subpopulations.
Altogether, the expression analysis of ligands and receptors in RPE cells also hints at the
possible value of co-culturing RPE cells with retinal organoid cultures to further support retinal
differentiation, cell maturation and improve the *in vitro* modelling of retinal biology.

Our analysis also revealed that cells in culture can develop a transcriptomic profile more
closely related to the adult native RPE with higher levels of expression of some RPE genes and
lower levels of expression of others, as observed in their native counterpart. Altogether, this thus
strongly suggests that as hPSC-derived RPE cells mature with time in culture, they acquire
characteristics more closely resembling those of an adult RPE profile.

Conclusion

The novel insight into the underlying genetic architecture of hPSC-derived RPE cells at short and
long time points in culture conditions revealed a gradual differentiation and maturation process,
as well as a stable RPE phenotype over time. Most cells with a clear RPE signature are found in
the "Common" subpopulation, indicating that RPE cells are present from an early time point in
culture and maintain this identity with time. The clustering also reveals that whilst some
subpopulations expressed more genes associated with retinal and RPE biology, other RPE
subpopulations demonstrated increased expression in mitochondrial and/or ribosomal related
genes. Altogether, this data suggests that hPSC-derived RPE cells develop their characteristic
signature early in the differentiation process and continue to mature over time in culture. It also
warrants the use of hPSC-derived RPE cells for modelling of RPE biology at early and later
differentiation timings.

Materials and methods

Cell culture and differentiation of hESCs to RPE cells

The hESC line H9 (Wicell) was maintained on vitronectin-coated plates using StemFlex (Thermo Fisher), with medium changed every second day [74]. Cells were differentiated into RPE cells as previously described [3] with the following modifications. Briefly, hESCs were maintained in culture until 70-80% confluent at which stage StemFlex was replaced with Complete E6 (05946, Stem Cell Technologies, Vancouver, Canada) supplemented with N2 (17502048, Thermo Fisher Scientific, Waltham, MA, USA) to induce retinal differentiation, with thrice weekly media changes for 33 days. On Day 33, medium was replaced with RPEM [α-MEM (12571071, Thermo Fisher Scientific), 5% fetal bovine serum (26140079, Thermo Fisher Scientific), non-essential amino acids (11140050, Thermo Fisher Scientific), penicillin-streptomycin- glutamine (10378016, Thermo Fisher Scientific), N1 (N6530, Sigma-Aldrich, St Louis, MO, USA), taurine-hydrocortisone-triiodo-thyronin (in-house)] to promote RPE differentiation, with medium changed every second day. Cells were cultured for 32 days, after which point maximal pigmentation is routinely observed. Cells were harvested with an 8-minute exposure to 0.25% Trypsin-EDTA (25200056, Thermo Fisher Scientific) and inactivated with RPEM. Non-RPE contaminants (visible as unpigmented cells), were manually removed from the culture, which begin shedding off the culture plate after ~ 2 minutes. Cells were seeded at a density of 75,000 cells / cm^2 onto growth-factor-reduced Matrigel-coated tissue culture plates (Corning). Media was changed every second day, with the first sample of cells harvested after 30 days (D30) and the second sample harvested on day 367 (D367) for scRNA-seq analysis. (Figure 1A).

RPE cell harvest and single-cell preparation

RPE cells were dissociated to single cells using 0.25% Trypsin-EDTA for 8 min and inactivated with RPEM. Cells were centrifuged (300g, 1 min) to pellet and resuspended in a small volume of RPEM containing 0.1% v/v propidium iodide (PI, Sigma-Aldrich) to exclude non-viable cells. Single cell suspensions were passed through a 35 µm filter prior to sorting. A minimum of 60,000 live cells (PI-negative) were sorted on a BD FACSAria IIU (100 µm, 20psi) into culture

624   medium. Cells were centrifuged (300g, 5 min) and resuspended in PBS containing 0.04% BSA

625   to a concentration of ~800-1,000 cells/µl. Approximately 17,400 cells were loaded onto a 10X

626   chip for a target recovery of 10,000 cells. The two culture time points were captured separately

627   to prepare two separate 10X reactions.

Generation of single cell GEMs and sequencing libraries

To generate single cell gel beads in emulsion (GEMs), single cell suspensions were loaded onto

10X Genomics Single Cell 3′ Chips together with the reverse transcription master mix following

the manufacturer's protocol for the Chromium Single Cell 3′ v2 Library (PN-120233, 10X

Genomics, Pleasanton, CA, USA). For each sample, sequencing libraries were generated with

unique sample indices (SI), assessed by gel electrophoresis (Agilent D1000 ScreenTape Assay,

Santa Clara, CA, USA) and quantified with qPCR (Illumina KAPA Library Quantification Kit,

Roche, Pleasanton, CA, USA). Following pooling and normalization to 4 nM, libraries were

denatured and diluted to 1.6 pM for loading onto the sequencer. Libraries were sequenced on an

Illumina NextSeq 500 (NextSeq Control Software v2.2.0 / Real Time Analysis v2.4.11) using

NextSeq 500/550 High Output Kit v2.5 (150 Cycles) (20024907, Illumina, San Diego, CA,

USA) as follows: 26 bp (Read 1), 8 bp (i7 Index) and 98 bp (Read 2).

Preprocessing, mapping and quantification of scRNA-seq data

We used the *cellranger mkfastq* and *cellranger count* pipelines from the Cell Ranger Single Cell

Software Suite (version 3.0.2) by 10x Genomics (http://10xgenomics.com) for initial quality

control, sample demultiplexing, mapping and quantification of raw sequencing data. The

cellranger count pipeline was run with the following argument: "--expect-cells=10000", and

reads were mapped to the *Homo sapiens* reference genome (GRCh38, Annotation: Gencode

v29). Filtered count matrices were then used for downstream analyses in R.

Quality control and normalization

Using Seurat v3.1.3 [16], data from the two culture timepoints underwent quality control and

normalization separately. The following values were calculated for each cell: total number of

Unique Molecular Identifiers (UMIs), number of detected genes and proportion of mitochondrial
and ribosomal-related transcripts relative to total expression. Cells were removed from
subsequent analysis if the total UMI content of the cell exceeded the 3x median absolute
deviation (MAD) of this measurement across all cells in the sample. Manual thresholds were
derived from outlier peaks in the distributions of the number of detected genes, and fraction of
mitochondrial and ribosomal-associated transcripts to total expression (Supplementary Figure 1).
Cells were removed from the 1-month time point sample if the number of detected genes
exceeded the lower and upper limits of 220 and 5,000, while cells from the 12-month time point
sample were removed if the number of detected genes were less than 220. Cells from both
timepoints were removed if mitochondrial-associated transcripts more than 25% of total
expression, and/or ribosomal-associated transcripts accounted for more than 60% of total
expression (Supplementary Figure 1A-D, Supplementary Table 1). The confounding effect of
these mitochondrial and ribosomal-associated QC metrics in remaining cells were regressed out
during cell-cell normalization, using the SCTransform function from Seurat [75] (Supplementary
Table 7).

Integration, dimensionality reduction and clustering
The dimensionality of the data then was reduced with Principal Component Analysis (PCA).
Subsequently, the 30 most statistically significant principal components (PCs) were reduced to
two dimensions by Uniform Manifold Approximation Projection (UMAP). These values were
used to construct a Shared Nearest Neighbour (SNN) graph for each cell. The Louvain method
for community detection was then used to identify clusters in each dataset across a range of
resolutions from 0 to 1.5. The results for all resolutions were plotted using clustree [76], which
showed the stabilisation of cell population identities at the resolution of 0.6 in the 1-month
culture and 0.7 in the 12-month culture (Supplementary Figure 2A). Both timepoint datasets
combined into one dataset for comparative analysis with the integration workflow from Seurat
[77]. This workflow used canonical correlation analysis (CCA) to identify 22,023 anchors based
on 3,000 of the most variable genes. The anchors were then used to align both datasets. To
integrate the clusters across both timepoints, the unsupervised version of MetaNeighbor [17] was
used to evaluate the similarities between the 1-month clusters and 12-month clusters. Cluster
pairs that were reciprocal top hits, and received a mean Area Under the Receiver Operating

Characteristic (AUROC) score greater than 0.8 were merged into one cluster (Supplementary

Figure 2C).

Clustering characterization and analysis

Network analysis was performed on significant differentially-expressed genes using Reactome

functional interaction analysis [78, 79]. Differential expression (DE) analysis was performed

using the FindMarkers function based on the Likelihood Ratio Test adapted for single cell gene

expression [80]. Gene Ontology (GO) analysis [81, 82] was performed using a PANTHER

overrepresentation test (Fisher exact test, FDR < 0.05) against the *Homo sapien* genome

(PANTHER version 14.1 Released 2019-03-12). Some clusters had insufficient gene markers for

GO analysis (Aged Clusters 0, 1 and 5). Canonical RPE markers, gene expression profiles and

their associated GO analysis specific to each cluster are provided in Supplementary Tables 3-5.

Trajectory analysis

Trajectory analysis was performed with *Monocle 3* v0.2.4 [52]. Harmonized Pearson residuals

produced by the integration step underwent dimensionality reduction with UMAP, and the

resulting projection was used to initialize trajectory inference. The node closest to cells

expressing proliferative markers was selected as the root of the trajectory, and pseudotime values

were calculated. Gene expression dynamics across the trajectory were characterised with Moran

I's test, which was applied via the "graph_test" function using the following arguments:

neighbor_graph = "principal_graph", reduction_method = "UMAP", expression_family =

"quasiposson". Differentially expressed genes with an FDR of 0.05 or less were clustered into

co-expression modules using the "find_gene_modules" function, and resulting protein

interactions were characterised with STRING.

Ethical statements

709 The experimental work was approved by the Human Research Ethics Committee of the

710 University of Melbourne (1545484) with the requirements of the National Health & Medical

711 Research Council of Australia (NHMRC) in accordance with the Declaration of Helsinki.

Data availability

Code and usage notes are available at:

https://github.com/powellgenomicslab/RPE_scRNA_AgedStudy. This repository consists of code used to process raw sequencing data in FASTQ format to cell-gene expression tables via the Cell Ranger pipeline, and code used to perform the following analysis: quality control, normalization, dimensionality reduction, clustering, differential expression and integration.

Data records

Data is available at ArrayExpress: E-MTAB-8511. Files are raw FASTQ files, and a tab separated matrix of UMIs per gene for each cell passing quality control filtering. BAM files can be generated using the supplied repository to process the FASTQ files via Cell Ranger.

Authors' contributions

GEL: Conceptualization; Data curation; Formal analysis; Visualization; Funding acquisition; Investigation; Methodology; Project administration; Resources; Roles/Writing - original draft; Writing - review & editing. AS: Conceptualization; Data curation; Formal analysis; Visualization; Methodology; Investigation; Software; Roles/Writing - original draft; Writing - review & editing. CJASA, VG, DCK, DAZ: Conceptualization; Methodology; Investigation; Roles/ Writing - review & editing. ELF, SHN: Conceptualization; Methodology; Funding acquisition; Roles/ Writing - review & editing. AWH: Conceptualization; Funding acquisition; Methodology; Resources; Writing - review & editing, Supervision JEP: Conceptualization; Data curation; Formal analysis; Funding acquisition; Methodology; Resources; Software, Roles/Writing - original draft; Writing - review & editing, Supervision AP: Conceptualization; Formal analysis; Visualization; Funding acquisition; Methodology; Project administration; Resources; Roles/Writing - original draft; Writing - review & editing, Supervision.

Competing interests

The authors have declared no competing interests.

Acknowledgments

This work was supported by a National Health and Medical Research Council (NHMRC-
Australia) Practitioner Fellowship (AWH), Career Development Fellowship (JEP) and Senior
Research Fellowship (AP, 1154389), an Australian Research Council Future Fellowship (AP,
FT140100047), a NHMRC project grants (1138253 to ELF and AP; 1062820 to SHN, 1124812
to SHN), a NHMRC synergy grant (1181010 to ELF and AP), grants from the Macular Disease
Foundation Australia (AP, JEP, AWH), the Jack Brockhoff Foundation (GEL), the DHB
Foundation (GEL, AP), the Ophthalmic Research Institute of Australia (AP, AWH), Stem Cells
Australia – the Australian Research Council Special Research Initiative in Stem Cell Science
(SHN, AWH, JEP, AP), the TMG Family Fund (AP, GEL), a donation from Ms Jacqueline
Pascual, the University of Melbourne and Operational Infrastructure Support from the Victorian
Government.

Authors' ORCID IDs

754 0000-0002-1774-5944 (Lidgerwood, G.E.); 0000-0002-8798-9821 (Senabouth, A.); 0000-0003-
755 1391-7160 (Smith-Anttila, C.J.A.); 0000-0003-0306-1952 (Gnanasambandapillai, V); 0000-
756 0002-9205-1934 (Kaczorowski, D.C.); 0000-0003-4928-1846 (Amann-Zalcenstein, D.); 0000-
757 0001-9412-9523 (Fletcher, E.L.); 0000-0003-0299-3301 (Naik, S.H.); 0000-0002-5123-5999
758 (Hewitt, A.W.); 0000-0001-9031-6356 (Powell, J.E.); 0000-0002-7408-9453 (Pébay, A)

References

[1] Perez VL, Saeed AM, Tan Y, Urbieta M, Cruz-Guilloty F. The eye: A window to the soul of the immune system. J Autoimmun 2013;45:7-14.

[2] Xu H, Chen M, Forrester JV. Para-inflammation in the aging retina. Prog Retin Eye Res 2009;28:348-68.

[3] Lidgerwood GE, Morris AJ, Conquest A, Daniszewski M, Rooney LA, Lim SY, et al. Role of lysophosphatidic acid in the retinal pigment epithelium and photoreceptors. Biochim Biophys Acta Mol Cell Biol Lipids 2018;1863:750-61.

[4] Lidgerwood GE, Lim SY, Crombie DE, Ali R, Gill KP, Hernández D, et al. Defined medium conditions for the induction and expansion of human pluripotent stem cell-derived retinal pigment epithelium. Stem Cell Rev Rep 2016;12:179-88.

[5] Kamao H, Mandai M, Okamoto S, Sakai N, Suga A, Sugita S, et al. Characterization of
human induced pluripotent stem cell-derived retinal pigment epithelium cell sheets aiming for
clinical application. Stem Cell Reports 2014;2:205-18.

[6] Osakada F, Jin Z-B, Hirami Y, Ikeda H, Danjyo T, Watanabe K, et al. In vitro differentiation
of retinal cells from human pluripotent stem cells by small-molecule induction. J Cell Sci
2009;122:3169-79.

[7] Buchholz DE, Pennington BO, Croze RH, Hinman CR, Coffey PJ, Clegg DO. Rapid and
efficient directed differentiation of human pluripotent stem cells into retinal pigmented
epithelium. Stem Cells Transl Med 2013;2:384-93.

[8] Foltz LP, Clegg DO. Rapid, directed differentiation of retinal pigment epithelial cells from
human embryonic or induced pluripotent stem cells. J Vis Exp 2017:56274.

[9] Bennis A, Jacobs JG, Catsburg LAE, ten Brink JB, Koster C, Schlingemann RO, et al. Stem
cell derived retinal pigment epithelium: The role of pigmentation as maturation marker and gene
expression profile comparison with human endogenous retinal pigment epithelium. Stem Cell
Rev Rep 2017;13:659-69.

[10] Kokkinaki M, Sahibzada N, Golestaneh N. Human induced pluripotent stem-derived retinal
pigment epithelium (RPE) cells exhibit ion transport, membrane potential, polarized vascular
endothelial growth factor secretion, and gene expression pattern similar to native RPE. Stem
Cells 2011;29:825-35.

[11] Lidgerwood GE, Hewitt AW, Pébay A. Human pluripotent stem cells for the modelling of
diseases of the retina and optic nerve: toward a retina in a dish. Curr Opin Pharmacol
2019;48:114-9.

[12] Llonch S, Carido M, Ader M. Organoid technology for retinal repair. Dev Biol
2018;433:132-43.

[13] Gagliardi G, Ben M'Barek K, Goureau O. Photoreceptor cell replacement in macular
degeneration and retinitis pigmentosa: A pluripotent stem cell-based approach. Prog Retin Eye
Res 2019;71:1-25.

[14] Sridhar A, Hoshino A, Finkbeiner CR, Chitsazan A, Dai L, Haugan AK, et al. Single-cell
transcriptomic comparison of human fetal retina, hPSC-derived retinal organoids, and long-term
retinal cultures. Cell Rep 2020;30:1644-59.e4.

[15] Hu Y, Wang X, Hu B, Mao Y, Chen Y, Yan L, et al. Dissecting the transcriptome landscape

of the human fetal neural retina and retinal pigment epithelium by single-cell RNA-seq analysis.

PLoS Biol 2019;17:e3000365.

[16] Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al.

Comprehensive integration of single-cell data. Cell 2019;177:1888-902.e21.

[17] Crow M, Paul A, Ballouz S, Huang ZJ, Gillis J. Characterizing the replicability of cell types

defined by single cell RNA-sequencing data using MetaNeighbor. Nat Commun 2018;9:884.

[18] Tarau I-S, Berlin A, Curcio CA, Ach T. The cytoskeleton of the retinal pigment epithelium:

from normal aging to age-related macular degeneration. Int J Mol Sci 2019;20:3578.

[19] Zhou X, Liao W-J, Liao J-M, Liao P, Lu H. Ribosomal proteins: functions beyond the

ribosome. J Mol Cell Biol 2015;7:92.

[20] Barkić M, Crnomarković S, Grabušić K, Bogetić I, Panić L, Tamarut S, et al. The p53 tumor

suppressor causes congenital malformations in Rpl24-deficient mice and promotes their survival.

Mol Cell Biol 2009;29:2489-504.

[21] Watkins-Chow DE, Cooke J, Pidsley R, Edwards A, Slotkin R, Leeds KE, et al. Mutation of

the diamond-blackfan anemia gene Rps7 in mouse results in morphological and neuroanatomical

phenotypes. PLoS Genet 2013;9:e1003094.

[22] Paraoan L, Grierson I, Maden BE. Analysis of expressed sequence tags of retinal pigment

epithelium: cystatin C is an abundant transcript. Int J Biochem Cell Biol 2000;32:417-26.

[23] Kay P, Yang YC, Hiscott P, Gray D, Maminishkis A, Paraoan L. Age-related changes of

cystatin C expression and polarized secretion by retinal pigment epithelium: potential age-related

macular degeneration links. Invest Ophthalmol Vis Sci 2014;55:926-34.

[24] Guyonneau L, Murisier F, Rossier A, Moulin A, Beermann F. Melanocytes and

pigmentation are affected in dopachrome tautomerase knockout mice. Mol Cell Biol

2004;24:3396-403.

[25] Takeda K, Yokoyama S, Yasumoto K-I, Saito H, Udono T, Takahashi K, et al. OTX2

regulates expression of DOPAchrome tautomerase in human retinal pigment epithelium.

Biochem Biophys Res Commun 2003;300:908-14.

[26] Strunnikova NV, Maminishkis A, Barb JJ, Wang F, Zhi C, Sergeev Y, et al. Transcriptome

analysis and molecular signature of human retinal pigment epithelium. Hum Mol Genet

2010;19:2468-86.

[27] Strunnikova N, Zhang C, Teichberg D, Cousins SW, Baffi J, Becker KG, et al. Survival of

retinal pigment epithelium after exposure to prolonged oxidative injury: a detailed gene

expression and cellular analysis. Invest Ophthalmol Vis Sci 2004;45:3767-77.

[28] Tanaka Y, Utsumi J, Matsui M, Sudo T, Nakamura N, Mutoh M, et al. Purification,

molecular cloning, and expression of a novel growth-promoting factor for retinal pigment

epithelial cells, REF-1/TFPI-2. Invest Ophthalmol Vis Sci 2004;45:245-52.

[29] Alizadeh P, Smit-McBride Z, Oltjen SL, Hjelmeland LM. Regulation of cysteine cathepsin

expression by oxidative stress in the retinal pigment epithelium/choroid of the mouse. Exp Eye

Res 2006;83:679-87.

[30] Samuel W, Kutty RK, Vijayasarathy C, Pascual I, Duncan T, Redmond TM. Decreased

expression of insulin-like growth factor binding protein-5 during N-(4-

hydroxyphenyl)retinamide-induced neuronal differentiation of ARPE-19 human retinal pigment

epithelial cells: regulation by CCAAT/enhancer-binding protein. J Cell Physiol 2010;224:827-

36.

[31] Quinn PM, Buck TM, Mulder AA, Ohonin C, Alves CH, Vos RM, et al. Human iPSC-

derived retinas recapitulate the fetal CRB1 CRB2 complex formation and demonstrate that

photoreceptors and Müller glia are targets of AAV5. Stem Cell Reports 2019;12:906-19.

[32] Paniagua AE, Herranz-Martín S, Jimeno D, Jimeno ÁM, López-Benito S, Carlos Arévalo J,

et al. CRB2 completes a fully expressed Crumbs complex in the retinal pigment epithelium. Sci

Rep 2015;5:14504.

[33] Winokur PN, Subramanian P, Bullock JL, Arocas V, Becerra SP. Comparison of two

neurotrophic serpins reveals a small fragment with cell survival activity. Mol Vis 2017;23:372-

84.

[34] Chang JT, Esumi N, Moore K, Li Y, Zhang S, Chew C, et al. Cloning and characterization

of a secreted frizzled-related protein that is expressed by the retinal pigment epithelium. Hum

Mol Genet 1999;8:575-83.

[35] Liu NP, Roberts WL, Hale LP, Levesque MC, Patel DD, Lu CL, et al. Expression of CD44

and variant isoforms in cultured human retinal pigment epithelial cells. Invest Ophthalmol Vis

Sci 1997;38:2027-37.

[36] Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, et al.

Proteomics. Tissue-based map of the human proteome. Science 2015;347:1260419.

[37] Bennis A, Ten Brink JB, Moerland PD, Heine VM, Bergen AA. Comparative gene

expression study and pathway analysis of the human iris- and the retinal pigment epithelium.

PLoS One 2017;12:e0182983.

[38] Cao L, Liu J, Pu J, Milne G, Chen M, Xu H, et al. Polarized retinal pigment epithelium

generates electrical signals that diminish with age and regulate retinal pathology. J Cell Mol Med

2018;22:5552-64.

[39] Westenskow P, Piccolo S, Fuhrmann S. Beta-catenin controls differentiation of the retinal

pigment epithelium in the mouse optic cup by regulating Mitf and Otx2 expression.

Development 2009;136:2505-10.

[40] Lee W-H, Joshi P, Wen R. Glutathione S-transferase pi isoform (GSTP1) expression in

murine retina increases with developmental maturity. Adv Exp Med Biol 2014;801:23-30.

[41] Wang AL, Lukas TJ, Yuan M, Du N, Tso MO, Neufeld AH. Autophagy and exosomes in

the aged retinal pigment epithelium: possible relevance to drusen formation and age-related

macular degeneration. PLoS One 2009;4:e4160.

[42] Pasovic L, Utheim TP, Reppe S, Khan AZ, Jackson CJ, Thiede B, et al. Improvement of

storage medium for cultured human retinal pigment epithelial cells using factorial design. Sci

Rep 2018;8:5688.

[43] Cai H, Shin MC, Tezel TH, Kaplan HJ, Del Priore LV. Use of iris pigment epithelium to

replace retinal pigment epithelium in age-related macular degeneration: a gene expression

analysis. Arch Ophthalmol 2006;124:1276-85.

[44] Alge CS, Suppmann S, Priglinger SG, Neubauer AS, May CA, Hauck S, et al. Comparative

proteome analysis of native differentiated and cultured dedifferentiated human RPE cells. Invest

Ophthalmol Vis Sci 2003;44:3629-41.

[45] Wang D, Chadha GK, Feygin A, Ivanov AI. F-actin binding protein, anillin, regulates

integrity of intercellular junctions in human epithelial cells. Cell Mol Life Sci 2015;72:3185-200.

[46] Li Y, Hao H, Tzatzalos E, Lin R-K, Doh S, Liu LF, et al. Topoisomerase IIbeta is required

for proper retinal development and survival of postmitotic cells. Biol Open 2014;3:172-84.

[47] Usui A, Mochizuki Y, Iida A, Miyauchi E, Satoh S, Sock E, et al. The early retinal

progenitor-expressed gene Sox11 regulates the timing of the differentiation of retinal cells.

Development 2013;140:740-50.

[48] van Soest SS, de Wit GMJ, Essing AHW, ten Brink JB, Kamphuis W, de Jong PTVM, et al.

Comparison of human retinal pigment epithelium gene expression in macula and periphery

highlights potential topographic differences in Bruch's membrane. Mol Vis 2007;13:1608-17.

[49] Tian J, Ishibashi K, Honda S, Boylan SA, Hjelmeland LM, Handa JT. The expression of

native and cultured human retinal pigment epithelial cells grown in different culture conditions.

Br J Ophthalmol 2005;89:1510-7.

[50] Whitmore SS, Wagner AH, DeLuca AP, Drack AV, Stone EM, Tucker BA, et al.

Transcriptomic analysis across nasal, temporal, and macular regions of human neural retina and

RPE/choroid by RNA-Seq. Exp Eye Res 2014;129:93-106.

[51] Cai H, Fields MA, Hoshino R, Del Priore LV. Effects of aging and anatomic location on

gene expression in human retina. Front Aging Neurosci 2012;4:8.

[52] Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, et al. The single-cell

transcriptional landscape of mammalian organogenesis. Nature 2019;566:496-502.

[53] Stern J, Temple S. Retinal pigment epithelial cell proliferation. Exp Biol Med

2015;240:1079-86.

[54] Qiu TG. Transplantation of human embryonic stem cell-derived retinal pigment epithelial

cells (MA09-hRPE) in macular degeneration. NPJ Regen Med 2019;4:19.

[55] Chen M, Forrester JV, Xu H. Dysregulation in retinal para-inflammation and age-related

retinal degeneration in CCL2 or CCR2 deficient mice. PLoS One 2011;6:e22818.

[56] Holtkamp GM, De Vos AF, Peek R, Kijlsta A. Analysis of the secretion pattern of

monocyte chemotactic protein-1 (MCP-1) and transforming growth factor-beta 2 (TGF-beta2) by

human retinal pigment epithelial cells. Clin Exp Immunol 1999;118:35-40.

[57] Bovolenta P, Esteve P, Ruiz JM, Cisneros E, Lopez-Rios J. Beyond Wnt inhibition: new

functions of secreted Frizzled-related proteins in development and disease. J Cell Sci

2008;121:737-46.

[58] He X, Cheng R, Benyajati S, Ma J-X. PEDF and its roles in physiological and pathological

conditions: implication in diabetic and hypoxia-induced angiogenic diseases. Clin Sci

2015;128:805-23.

[59] Penn JS, Madan A, Caldwell RB, Bartoli M, Caldwell RW, Hartnett ME. Vascular

endothelial growth factor in eye disease. Prog Retin Eye Res 2008;27:331-71.

[60] Yang P, Skiba NP, Tewkesbury GM, Treboschi VM, Baciu P, Jaffe GJ. Complement-
mediated regulation of Apolipoprotein E in cultured human RPE cells. Invest Ophthalmol Vis
Sci 2017;58:3073-85.

[61] Toomey CB, Johnson LV, Bowes Rickman C. Complement factor H in AMD: Bridging
genetic associations and pathobiology. Prog Retin Eye Res 2018;62:38-57.

[62] Liao J-L, Yu J, Huang K, Hu J, Diemer T, Ma Z, et al. Molecular signature of primary
retinal pigment epithelium and stem-cell-derived RPE cells. Hum Mol Genet 2010;19:4229-38.

[63] May-Simera HL, Wan Q, Jha BS, Hartford J, Khristov V, Dejene R, et al. Primary cilium-
mediated retinal pigment epithelium maturation is disrupted in ciliopathy patient cells. Cell Rep
2018;22:189-205.

[64] Rodríguez-Menéndez S, García M, Fernández B, Álvarez L, Fernández-Vega-Cueto A,
Coca-Prados M, et al. The Zinc-Metallothionein redox system reduces oxidative stress in retinal
pigment epithelial cells. Nutrients 2018;10.

[65] Capowski EE, Simonett JM, Clark EM, Wright LS, Howden SE, Wallace KA, et al. Loss of
MITF expression during human embryonic stem cell differentiation disrupts retinal pigment
epithelium development and optic vesicle cell proliferation. Hum Mol Genet 2014;23:6332-44.

[66] McGowan KA, Li JZ, Park CY, Beaudry V, Tabor HK, Sabnis AJ, et al. Ribosomal
mutations cause p53-mediated dark skin and pleiotropic effects. Nat Genet 2008;40:963-70.

[67] Grewal R, Stepczynski J, Kelln R, Erickson T, Darrow R, Barsalou L, et al. Coordinated
changes in classes of ribosomal protein gene expression is associated with light-induced retinal
degeneration. Invest Ophthalmol Vis Sci 2004;45:3885-95.

[68] Leach LL, Buchholz DE, Nadar VP, Lowenstein SE, Clegg DO. Canonical/β-catenin Wnt
pathway activation improves retinal pigmented epithelium derivation from human embryonic
stem cells. Invest Ophthalmol Vis Sci 2015;56:1002-13.

[69] Esteve P, Trousse F, Rodríguez J, Bovolenta P. SFRP1 modulates retina cell differentiation
through a beta-catenin-independent mechanism. J Cell Sci 2003;116:2471-81.

[70] Rodriguez J, Esteve P, Weinl C, Ruiz JM, Fermin Y, Trousse F, et al. SFRP1 regulates the
growth of retinal ganglion cell axons through the Fz2 receptor. Nat Neurosci 2005;8:1301-9.

[71] Saint-Geniez M, Maharaj ASR, Walshe TE, Tucker BA, Sekiyama E, Kurihara T, et al.
Endogenous VEGF is required for visual function: evidence for a survival role on müller cells
and photoreceptors. PLoS One 2008;3:e3554.

[72] Nishijima K, Ng Y-S, Zhong L, Bradley J, Schubert W, Jo N, et al. Vascular endothelial

growth factor-A is a survival factor for retinal neurons and a critical neuroprotectant during the

adaptive response to ischemic injury. Am J Pathol 2007;171:53-67.

[73] van Niel G, Bergam P, Di Cicco A, Hurbain I, Lo Cicero A, Dingli F, et al. Apolipoprotein

E regulates amyloid formation within endosomes of pigment cells. Cell Rep 2015;13:43-51.

[74] Daniszewski M, Nguyen Q, Chy HS, Singh V, Crombie DE, Kulkarni T, et al. Single-cell

profiling identifies key pathways expressed by iPSCs cultured in different commercial media.

iScience 2018;7:30-9.

[75] Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq

data using regularized negative binomial regression. Genome Biol 2019;20:296.

[76] Zappia L, Oshlack A. Clustering trees: a visualization for evaluating clusterings at multiple

resolutions. Gigascience 2018;7.

[77] Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell transcriptomic

data across different conditions, technologies, and species. Nat Biotechnol 2018;36:411-20.

[78] Anders S, Huber W. Differential expression analysis for sequence count data. Genome Biol

2010;11:R106.

[79] Wu G, Feng X, Stein L. A human functional protein interaction network and its application

to cancer data analysis. Genome Biol 2010;11:R53.

[80] McDavid A, Finak G, Chattopadyay PK, Dominguez M, Lamoreaux L, Ma SS, et al. Data

exploration, quality control and testing in single-cell qPCR-based gene expression experiments.

Bioinformatics 2013;29:461-7.

[81] Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology:

tool for the unification of biology. The Gene Ontology Consortium. Nat Genet 2000;25:25-9.

[82] The Gene Ontology Consortium. The gene ontology resource: 20 years and still going

strong. Nucleic Acids Research 2019;47:D330-D8.

 Figure Legends

Figure 1: scRNA-seq transcriptome profiling of hPSC-derived RPE cells reveals 18

subpopulations.

(**A**) Schematic representations of the experimental flow. (**B**) UMAP of single cell- expression

profile from 16,576 cells, clustering into 18 subpopulations, split by condition (1-month- and 12-

month-old) and combined. (**C**) Cluster grouping represented by a Venn diagram, identifying 18

subpopulations, showing Young (red), Aged (green) and their common subpopulations (blue).

Cell numbers are indicated for each subpopulation. (**D**) UMAP of canonical RPE markers in 1-

month- and 12-month-old cultures, organised by cellular functions: extracellular structure

organization (*CST3*, *EFEMP1*, *ITGAV, CRISPLD1*, *ITGB8*); melanin biosynthesis (*PMEL*, *TTR*,

TYRP1, *TYR*, *DCT*), lipid biosynthesis (*PTGDS*, *INPP5K),* visual cycle (*LRAT, PLTP, RLBP1*,

RPE65, *RGR*, *RBP1*, *BEST1*) and secretion (*SERPINF1*). Levels of gene expression per cell

(percent expressed) are shown with colour gradients. Abbreviations: scRNA-seq, single cell

RNA sequencing; hPSC, human pluripotent stem cell; RPE, retinal pigment epithelium; UMAP,

Uniform Manifold Approximation and Projection for Dimension Reduction.

Figure 2: Characterisation of hPSC-derived RPE populations.

(**A**) Heatmap showing the top most conserved markers (gene transcripts are indicated on the left

side) in all individual cells of each 18 subpopulation (indicated on top, with colour matching

subpopulations of Figure 1C). The intensity of gene expression is indicated by colour variation.

(**B**) Dotplot representation of single cell- expression profile from 1-month-old and 12-month-old

cells for selected gene markers, representative of progenitor cells, or RPE with genes linked to

RPE functions. Populations arising from 1-month-old cultures are represented in orange and

those from 12-month-old cultures in black. Levels of gene expression per cell are shown with

colour gradients, frequency of cells expressing gene (percent expressed) are shown with size of

dots. Abbreviations: hPSC, human pluripotent stem cell; RPE, retinal pigment epithelium.

1006

Figure 3: Expression patterns of selected conserved markers and gene ontology pathway in

the hPSC-derived RPE cells.

(A-C) Violin plots of selected conserved markers in each "Common" subpopulation

characteristic of the RPE **(A),** neural differentiation **(B)** and extracellular matrix **(C)**. The plots

describe the distribution and relative expression of each transcript in each subpopulation, with

separation of cells belonging to the 1-month-old ("Control") and 12-month-old ("Aged")

cultures. (**D**) PANTHER GO-Slim (Biological Process) pathways associated with each of the

"Common" subpopulations (1-6), with fold enrichment per subpopulation. Abbreviations: hPSC,

human pluripotent stem cell; RPE, retinal pigment epithelium

1016

Figure 4: Aged RPE subpopulations likely increase their handling of metals and

antioxidant abilities.

(**A**) Feature plots and (**B**) ridge plots of expression profiles of key metallothioneins and *DCT*

across the 1-month-old and 12-month-old cells (**A**) and across all subpopulations (**B**). In (**A**), the

intensity of gene expression is indicated by colour gradient. Abbreviations: RPE, retinal pigment

epithelium.

1023

Figure 5: Only a few cells retain a proliferative profile.

(**A**) UMAP of single cell- expression profile split by conditions (1-month-old and 12-month-old)

and (**B**) trajectory analysis using Monocle 3, and for (**C**) markers associated with proliferation

(*MKI67*, *TOP2A*, *PCLAF*, *RRM2*, *TPX2*, *PTTG1*) across all cells. (**D, E**) Pseudotime analysis of

early retinal markers (*PAX6*, *RAX*) and RPE genes (*PMEL*, *RLBP1*, *RGR*, *TYR, RBP1*, *RPE65*)

across the 1-month-old and 12-month-old cells (**D**) and across all subpopulations (**E**). In (**C, D**),

the intensity of gene expression is indicated by colour gradient. Abbreviations: UMAP, Uniform

Manifold Approximation and Projection for Dimension Reduction; RPE, retinal pigment

epithelium.

1033

Figure 6: RPE subpopulations contribute to paracrine signalling.

UMAP of single cell- expression profile of markers for 1-month-old and 12-month-old cultures,

with associated violin plots across all populations for (**A**) *CCL2*; (**B**) *SFRPs*, *FZDs* and *WNTS*;

(**C**) *VEGFA;* (**D**) *APOE*, *LDLR* and *LRPs*. In the UMAPs, the intensity of gene expression is

indicated by colour gradient. Abbreviations: RPE, retinal pigment epithelium; UMAP, Uniform

Manifold Approximation and Projection for Dimension Reduction.

1040

Figure 7: hPSC-derived RPE subpopulations express native RPE markers with different

patterns.

Violin plots of selected markers representative of native RPE cells (obtained from [28]) across

all subpopulations and in the three main populations "Young", "Aged", "Common", represented

in different colours. Genes found in (**A**) all subpopulations (*SERPINF1*, *BEST1*, *TYR*, *TTR*,

RPE65); (**B**) in more immature subpopulations (*SFRP2*, *MKI67*, *DCT*); with variations

associated with (**C**) visual cycle (*PLTP, RGR, RLBP1, LRAT*); (**D**) melanin biosynthesis (*PMEL*,

TYRP1); (**E**) lipid biosynthesis (*PTGDS*, *INPP5K*); (**F**) extracellular structure organization

(*CST3*, *ITGB8*, *EFEMP1*, *ITGAV*, *CRISPLD1*); (**G**) phagocytic activity (*GULP1*), and secretion

(*VEGFA*). The plots describe the distribution and relative expression of each transcript in the

subpopulations. Abbreviations: hPSC, human pluripotent stem cell; RPE, retinal pigment

epithelium.

1053

Figure 8: Some hPSC-derived RPE subpopulations acquire a gene expression profile closer

to adult native RPE with time in culture.

Violin plots of selected markers representative of native adult RPE cells (obtained from [28])

across all subpopulations and in the three main populations "Young", "Aged", "Common",

represented in different colours. Genes found to be (**A**) upregulated in adult RPE (*CHRNA3,*

RBP1, *MYRIP*, *DUSP4, GEM, CRX* and *TFPI2*); or (**B**) downregulated in adult RPE (*SFRP5*,

SLC6A15). The plots describe the distribution and relative expression of each transcript in the

subpopulations. Abbreviations: hPSC, human pluripotent stem cell; RPE, retinal pigment

epithelium.

1063

Supplementary Figure Legends

Supplementary Figure 1: Filtering of cells based on quality control metrics.

(**A**) Total UMIs per cell. Adaptive thresholds depicted in red represent 3 MAD of total UMI

distribution across all cells. (**B**) Total features per cell. Thresholds represented in red were

manually set to 220 for both samples. Upper threshold was set to 5,000 for 1-month timepoint.

(**C**) Percentage of mitochondrial gene expression relative to total expression. Threshold was set

to 25% for both timepoints. (**D**) Percentage of ribosomal expression relative to total expression.

Threshold was set to 60% for both samples. (**E**) Relationship between total UMIs, detected genes

and percentage of mitochondrial gene expression. Histograms for total UMIs and detected

features are located on x and y margins of scatter plot, respectively. Scatter plot represents the
relationships between the two metrics and are coloured by percentage mitochondrial expression.
Abbreviations: UMIs, Unique Molecular Identifiers; 3 MAD, 3 Median Absolute Deviations.
1076

Supplementary Figure 2: Characterisation of stable cell subpopulations.
Graph-based clustering was performed at different resolutions ranging from 0 to 1.5, in
increments of 0.1 for (**A**) 1-month-old cells, (**B**) 12-month-old cells and (**C**) integrated
(combined) datasets using *clustree.* Regions of stability are represented by minimal branching.
(**D**) Integration of clusters identified in 1-month-old and 12-month-old cells via MetaNeighbor.
Heatmap represents degree of similarity of clusters as measured by AUROC values.
Abbreviations: AUROC, Area Under Receiver Operating Characteristic.
1084

Supplementary Figure 3: Residual Trajectory Genes Modules Analysis measured with
Monocle 3.
(**A**) Monocle 3 pseudotime represented in beeswarm plots. (**B**) Heatmap generated from
Monocle 3 Residual gene trajectory- differential expression for the four modules. (**C-F**) STRING
analysis of genes from Supplementary Table 6 for Modules (**C**) One (red: regulation of
development); (**D**) Two (red: regulation of synapse organisation); (**E**) Three (with no relevant
biological processes); and (**F**) Four (blue: cell cycle and red: mitosis).
1092

1093 **Supplementary Figure 4: Analysis of gene trajectory by pseudo-time analysis of early and**
1094 **mature RPE markers.**
1095 Gene expression for early and mature RPE markers using Monocle 3, with each subpopulation
1096 being colour coded. Trends of expression are represented by a black line within each panel.
1097 Abbreviations: RPE, retinal pigment epithelium.
1098

1099 ## Supplementary Tables

1100 **Supplementary Table 1. Quality control parameters.**

1101

1102 **Supplementary Table 2. Cluster analysis with number of cells per subpopulation.**

1103

1104 **Supplementary Table 3. Conserved cluster markers, PANTHER Gene Ontology pathways**
1105 **and differential expression for each subpopulation within the "Common" Subpopulations.**
1106 Cluster (pct.1) shows frequency of cells expressing a marker compared to all others (pct.2).
1107 avg_logFC: average level of expression per cell in subpopulations compared to all others.

1108

1109 **Supplementary Table 4. Conserved cluster markers and Gene Ontology pathways for each**
1110 **subpopulation within the "Young" Subpopulations.**
1111 Cluster (pct.1) shows frequency of cells expressing a marker compared to all others (pct.2).
1112 avg_logFC: average level of expression per cell in subpopulations compared to all others.

1113

1114 **Supplementary Table 5. Conserved cluster markers and Gene Ontology pathways for each**
1115 **subpopulation within the "Aged" Subpopulations.**
1116 Cluster (pct.1) shows frequency of cells expressing a marker compared to all others (pct.2).
1117 avg_logFC: average level of expression per cell in subpopulations compared to all others.

1118

1119 **Supplementary Table 6. Monocle3 residual trajectory Residual Trajectory.**
1120 Differential expression and genes modules.

1121

1122 **Supplementary Table 7. Details of cell filtration criteria.**
1123 Number of cells removed by each criterion in 1-month-old and 12-month-old cultures.

A Young clusters | Common clusters | Aged clusters

0 | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 | 6 | 0 | 1 | 2 | 3 | 4 | 5

Scaled expression (average log fold change)

B

Percentage expressed (%)

Progenitor
Lipid biosynthesis
Visual cycle
Melanin biosynthesis
Secretion
Phagocytosis
Extracellular structure organisation

1-month-old
12-month-old

Common clusters: 6, 6, 5, 5, 4, 4, 3, 3, 2, 2, 1, 1

Aged clusters: 5, 4, 3, 2, 1, 0

Young clusters: 5, 4, 3, 2, 1, 0

BMP7 *SOX4* *INPP5K* *PTGDS* *LRAT* *RGR* *PLTP* *RLBP1* *RBP1* *BEST1* *RPE65* *DCT* *TYRP1* *TYR* *TTR* *PMEL* *ENPP2* *VEGFA* *SERPINF1* *GULP1* *ITGAV* *CRISPLD1* *CST3* *EFEMP1*

A Cells ordered by Monocle 3 pseudotime

B Residual gene trajectory

C STRING analysis for Module One

D STRING analysis for Module Two

E STRING analysis for Module Three

F STRING analysis for Module Four