Müllerian mimicry of a quantitative trait despite contrasting levels of genomic divergence and

selection

Emma V. Curran[1*], Sean Stankowski[1], Carolina Pardo-Diaz[2], Camilo Salazar[2], Mauricio Linares[2], Nicola

J. Nadeau[1, 3*]

[1]Department of Animal and Plant Sciences, University of Sheffield, UK;

[2]Biology Program, Faculty of Natural Sciences and Mathematics, Universidad del Rosario, Bogota,

Colombia;

[3]The Smithsonian Tropical Research Institute, Panama City, Republic of Panama

*Corresponding authors: e.v.curran@sheffield.ac.uk, n.nadeau@sheffield.ac.uk

Running title: Müllerian mimicry of a quantitative trait

Abstract word count: 244

Total word count: 7750

Abstract

Hybrid zones, where distinct populations meet and interbreed, give insight into how differences between

populations are maintained despite gene flow. Studying clines in genetic loci and adaptive traits across

hybrid zones is a powerful method for understanding how selection drives differentiation within a single

species, but can also be used to compare parallel divergence in different species responding to a common

selective pressure. Here, we study parallel divergence of wing colouration in the butterflies *Heliconius*

erato and *H. melpomene*, which are distantly related Müllerian mimics that show parallel geographic

variation in both discrete variation in pigmentation, and quantitative variation in structural colour. Using

geographic cline analysis, we show that clines in these traits are positioned in the roughly the same

geographic region for both species, which is consistent with direct selection for mimicry. However, the

width of the clines varies markedly between species. This difference is explained in part by variation in

the strength of selection acting on colour traits within each species, but may also be influenced by

differences in the dispersal rate and total strength of selection against hybrids between the species.

Genotyping-by-sequencing also revealed weaker population structure in *H. melpomene*, suggesting the

hybrid zones may have evolved differently in each species; which may also contribute to the patterns of

phenotypic divergence in this system Overall, we conclude that multiple factors are needed to explain

patterns of clinal variation within and between these species, although mimicry has probably played a

central role.

Key words: Hybrid zones, quantitative trait variation, cline analysis, Müllerian mimicry, parallel

divergence

Introduction

Hybrid zones, where genetically differentiated populations are in contact and interbreed, have long been a valuable resource in understanding the evolutionary processes shaping taxonomic boundaries (Barton & Gale, 1993; Endler, 1977). Hybrid zones can form in continuously distributed populations, where different alleles are favoured at either end of an ecological gradient, a process called primary intergradation (Endler, 1977). Alternatively, they can form when previously isolated populations, which have become genetically differentiated in allopatry, come into secondary contact (Endler, 1977). Both scenarios can lead to the formation of sharp geographic clines in quantitative traits and the loci that underlie them. These clines reflect the balance between gene flow and divergent selection (Barton & Hewitt, 1985), and their study can therefore provide deep insight into potential targets of natural selection.

Cline theory provides a powerful framework for studying patterns of variation across hybrid zones, enabling key biological parameters, including the strength and nature of selection shaping variation, to be estimated (Barton & Hewitt, 1985). By fitting geographic cline models to many loci or quantitative traits, it is possible to understand how the nature and the relative strength of selection varies among them. For example, assuming selection is acting across a sharp environmental gradient, the cline centre is indicative of the geographic location where the direction of divergent selection switches. In this case, if clines are centred at the same location (henceforth referred to as cline coincidence), this indicates that a suite of traits and loci are all affected by a common selective agent, or multiple agents that coincide geographically (Barton and Hewitt 1985). Variation in cline width can be used to make inferences about the strength of selection acting on a locus or trait, with narrower clines indicating stronger selection, all else being equal. The overall shape of clines is also informative about the nature of selection shaping the cline. For example, if variation at a trait or locus is shaped only by direct selection, the cline is predicted to have a sigmoidal shape (Barton and Hewitt 1985; Barton and Gale 1993). However, if the strength of direct selection on each locus is outweighed by indirect selection from many loci in linkage disequilibrium (LD), the total selection affecting each locus in LD will be approximately equal (Kruuk, Baird, Gale, & Barton, 1999; J M Szymura & Barton, 1991). This can result in many clines with similar

centres and widths, and with steeper centres than would be expected from direct selection alone, referred

to as "stepped" clines (Barton & Hewitt, 1985; Vines et al., 2016)

Despite being primarily used to study patterns of trait and marker variation within a single species,

cline analysis may also be used to understand how closely related species are shaped by the same extrinsic

selection pressure (e.g. Mallet et al., 1990). For example, if multiple closely related and ecologically similar

species are distributed across the same habitat transition, local adaptation may cause similar traits to diverge

in concert. Although examples of parallel adaptation can demonstrate striking convergence, the extent of

trait divergence within each species, and extent of parallelism between them, may vary depending on a host

of factors. For example, at White Sands, New Mexico, three lizard species show strong divergence in their

dorsal colour, an adaptation that improves crypsis on different soil types (Rosenblum & Harmon, 2011).

However, the extent of colour divergence varies between the species for reasons that are not entirely clear

(Rosenblum & Harmon 2011). Cline analysis can be used to precisely quantify differences in the

geographical distribution and variation of putative adaptive traits between species. When combined with

genome-wide data, this can provide insight into factors influencing the degree of parallelism. Differences

between species in intrinsic factors, such as their dispersal rate, population density, variation in their past

demographic histories, and the genetic architecture of traits, could alter patterns of clinal variation in

adaptive traits between species that are otherwise subject to the same extrinsic selection pressures.

Here, we studied a case of parallel divergence in the Müllerian co-mimics *Heliconius erato* and

Heliconius melpomene. We examined clinal variation in two colour pattern traits: the yellow hindwing

bar and iridescence. Where the pair co-occur, they converge on almost identical patterns to share the cost

of educating predators of their distastefulness (Brown, 1981). Both species comprise many parapatric

colour pattern races, or subspecies, connected by hybrid zones (Mallet, 1993; Rosser, Dasmahapatra, &

Mallet, 2014). When different subspecies hybridise, their offspring can display novel or heterozygous

phenotypes (Arias et al., 2008; Mallet, 1989; Mallet, 1986). Predators are less likely to learn to avoid rare

phenotypes, causing frequency-dependent selection on colour patterns (Langham, 2004; Mallet & Barton,

1989). This maintains stable hybrid zones (Mallet, 1986; Rosser et al., 2014). The diverse colouration

seen in the *Heliconius* genus has been extensively studied, the vast majority of which is determined by a genetic 'tool kit' of five major-effect loci (S W Baxter, Johnston, & Jiggins, 2009; Kronforst et al., 2006; A. Martin et al., 2012; Nadeau, 2016; Nadeau et al., 2016; Reed et al., 2011; Westerman et al., 2018). Previous studies have found low levels of genetic differentiation between parapatric colour races, with a few diverged loci, mainly controlling colour pattern differences (Martin et al., 2013; Nadeau et al., 2014; Supple et al., 2013).

Near the Panamanian-Colombian border, there are co-occurring hybrid zones between subspecies of *H. erato* and *H. melpomene*, which differ in the presence of a yellow hindwing bar and in iridescent blue colouration (Mallet, 1986; Figure 1). Iridescence is produced by nano-structural ridges on the surface of wing scales, which are layered to produce constructive interference of blue light (Parnell et al., 2018). In a system so well-studied, little is known regarding selection on structural colour (Sweeney, Jiggins, & Johnsen, 2003). Divergence in this trait has not been previously studied. While the yellow bar is controlled by a single major-effect gene (Mallet, 1986; Nadeau, 2016), iridescence segregates as continuous variation, with conservative estimates suggesting it is controlled by around five additive genetic loci (Brien et al., 2018). While differences in pigment colouration across hybrid zones seem to be maintained by strong divergent selection, despite gene flow across the rest of the genome (Nadeau et al., 2014), it is unclear whether we would expect to see this in a more complex trait such as iridescence. Polygenic local adaptation may only require small allele frequency changes, but can also involve greater levels of covariance between loci (Le Corre & Kremer, 2012). The combined action of divergent selection and the build-up of statistical associations between loci can reduce effective migration rates across the genome (Flaxman, Wacholder, Feder, & Nosil, 2014; Kruuk et al., 1999). Therefore, an increased level of overall genome-wide differentiation, and population level genetic structure may be expected across hybrid zones over which quantitative variation is maintained, particularly if the trait is highly polygenic.

Here, we use geographic cline analysis to examine the selection regimes impacting variation and convergence of iridescence and other traits, both within and between the co-mimics *H. erato* and *H. melpomene*. Within species, we are primarily interested in understanding how the genetic basis of these

traits has influenced their divergence across the hybrid zone. Because iridescence is polygenic, it may be

more difficult for direct selection to maintain strong trait divergence along the cline compared with the

yellow bar trait, which has a simple genetic basis and is highly visible to selection. Between the species,

our aim is to compare the extent of parallel divergence in the co-mimics. Due to the strong existing

evidence that mimicry drives colour pattern convergence between *H. erato* and *H. melpomene*, our null

expectation would be that clines in colour traits should be very similar in position and shape. Any

deviations from this expectation would suggest that either selection is acting differently on iridescence in

each species, or that some other factor has affected the extent of divergence within each species.

Methods

Butterfly specimens

Heliconius melpomene and *Heliconius erato* individuals were collected from several sites in the Chocó-

Darien ecoregion between the Andes and the Pacific in Colombia, and part way across the isthmus of

Panama (Figure 1, SI Table S1). Wings were removed and stored in envelopes. Bodies were preserved in

NaCl saturated 20% dimethyl sulfoxide (DMSO) 0.25M EDTA.

Sequencing data

Restriction-associated DNA (RAD) sequence data were generated for 265 *H. erato* (SI Table S2), and

whole genome re-sequencing was carried out on 36 *H. melpomene* individuals (SI Table S3). Genomic

DNA was extracted from each individual using DNeasy Blood and Tissue Kits (Qiagen). Library

preparation and sequencing was carried out by Edinburgh Genomics (University of Edinburgh).

Single-digest RAD libraries were prepared using the *Pst1* restriction enzyme, with eight base-pair

barcodes and sequenced on the Illumina HiSeq 2500 platform (v4 chemistry), generating an average of

554,826 125 base paired-end reads per individual (see SI Table S2 for coverage and accession

information). We demultiplexed the pooled reads using the RADpools program in the RADtools package

version 1.2.4 (Baxter et al., 2011).

For the whole-genome sequencing, TruSeq Nano, gel-free libraries were prepared from genomic

DNA samples of 36 *H. melpomene* individuals and sequenced on Illumina HiSeq 2500 platform (v4

chemistry), generating an average of 31,484,363 125 base paired-end reads per individual (see SI Table

S3 for coverage and accession information).

Data processing and variant calling

We checked the quality of all the raw sequencing reads using FastQC (v 0.11.5) and removed any

remaining adapters using Trim Galore (v 0.4.1). We aligned the sequence data of all individuals, both

RAD sequenced and WGS, to their corresponding reference genomes, either *Heliconius melpomene*

version 2 (Davey et al., 2016) or *Heliconius erato* (Van Belleghem et al., 2017)*,* obtained from lepbase

(Challis, Kumar, Dasmahapatra, Jiggins, & Blaxter, 2016), using bowtie2 (v 2.3.2), with the local

alignment option, and the very-sensitive pre-set parameter options to improve accuracy of the alignment.

We used samtools (v 1.3.1) to sort and index the alignment files. We removed any duplicates that may

have arisen during library preparation using the MarkDuplicates program in Picard tools (v 1.92).

Single nucleotide polymorphism (SNP) datasets were generated using samtools mpileup (v 1.5) to

compute genotype likelihoods and bcftools (v 1.5) for variant calling. For a site to be a variant, the

probability that it was homozygous for the reference allele across all samples was required to be less than

0.05. Multiallelic sites, insertions and deletions were ignored. For *H. melpomene* we identified

30,027,707 SNPs and for *H. erato* we identified 5,088,449 SNPs. We removed SNPs with a phred quality

score lower than 30, that lacked sequence data in 50% or more of the individuals, that had a minor allele

frequency lower than 0.05 or that were private variants. We pruned based on linkage disequilibrium,

discarding SNPs within a 20kb window with $r^2 > 0.8$, using the bcftools plugin '+prune'. This reduced the

initial number of called SNPs down to 9,336,937 in *H. melpomene* and 159,405 in *H. erato*.

Population structure

To examine population structure, we estimated the ancestry of each individual using the software

NGSadmix (Skotte, Korneliussen, & Albrechtsen, 2013), which estimates the proportion of each genome

that can be attributed to predefined number of populations (*k*) using genotype likelihoods. For each

species, NGSadmix was run for a range of values of *k*, one to ten, each being replicated ten times with a

random seed. The value of *k* best describing the population structure was determined using the Δk

criterion (Evanno, Regnaut, & Goudet, 2005), implemented in CLUMPAK (Kopelman, Mayzel,

Jakobsson, Rosenberg, & Mayrose, 2015).

We carried out a principal components analysis (PCA) using PCAngsd (Meisner & Albrechtsen,

2018), which estimates a covariance matrix based on genotype likelihoods. We used eigenvector

decomposition to retrieve the principal components of genetic structure.

Population differentiation

To test the extent of genetic differentiation between the iridescent and non-iridescent subspecies, we

measured F_{ST} between all individuals from iridescent populations south of the hybrid zone, and all non-

iridescent individuals north of the hybrid zone, excluding the sampling site Jaqué, which was in the centre

of the hybrid zone in both species. In each species, the two non-iridescent colour pattern races were

collapsed into a single "non-iridescent" group, north of the hybrid zone, since our results show there is no

genetic structure between them based on race. SNP datasets were generated for each species, using

samtools mpileup and bcftools (v 1.5). In each species Hudson's F_{ST} estimator was calculated among

populations (Hudson, Slatkin, & Maddison, 1992):

192
$$F_{ST}^{Hudson} = 1 - \frac{Hw}{Hb} = \frac{p_1(1-p_1) + p_2(1-p_2)}{p_1(1-p_2) + p_2(1-p_1)}$$

Where *Hw* is the within-population heterozygosity, *Hb* is the between-population heterozygosity,

and p_1 and p_2 represent the allele frequencies in each population. This was calculated in *R* for every SNP

with a custom script. Average genome-wide F_{ST} was calculated as a ratio of averages, by averaging the

variance components, *Hw* and *Hb*, separately, as recommended by Bhatia et al. (2013) We also estimated

average genome-wide F_{ST} between all pairs of populations, including those in the hybrid zone, for each

species, and plotted pairwise F_{ST} against pairwise geographic distance.

Phenotypic measurements

Digital images of butterfly wings were taken with a Nikon D7000 DSLR camera fitted with an AF-S DX

Micro NIKKOR 40 mm f/2.8G lens (Nikon UK Ltd., Surrey, UK), mounted on an adjustable platform.

Standardised lighting conditions were achieved using two external natural daylight fluorescent lights,

mounted to illuminate at 45 degrees from incident, to maximise brightness of observed iridescent colour.

Photographs were taken with a shutter speed of 1/60 sec and an aperture of f/10. Each sample was

photographed with an X-Rite colorchecker passport (X-Rite, Inc., MI, USA) in shot. The Nikon raw

(.NEF) image files were converted to standard raw files (.DNG) using Adobe DNG converter (Adobe

Systems Inc., USA). The RGB channels in the images were then linearized using the neutral grey scale on

the colorchecker using GNU Image Manipulation Program, v2.8.

The mean RGB values from regions in the discal cell on the right forewing and the Cu_2 cell on the

right hindwing were measured (SI Figure S1A). If the wings on the right-hand side showed damage,

wings on the left-hand side were used. Wing regions were selected using the polygon selection tool in

ImageJ, version 1.50b (Abràmoff, Magalhães, & Ram, 2004), and mean RGB scores were measured using

the Color Histogram plugin. To minimise variation in blue colour due to age and wing wear, we excluded

individuals with extensive wing wear or damage.

We tested for repeatability (Whitlock & Schluter, 2009) of the RGB values on 26 individuals

photographed a second time under the same conditions on a different day, with a second set of RGB

measurements taken. These individuals were selected from regions in which varying levels of iridescence

is seen (20 individuals from Valle del Cauca, Colombia, and 6 individuals from Darién, Panama).

Variance among individuals was calculated by taking the difference between the group mean square and

the error mean square, and dividing it by the number of replicates. These components of variance were

extracted from a general linear model in R v3.2.3 (R Core Team, 2015). The fraction of total variance that

is due to true differences between individuals was then calculated by dividing the variance among

individuals by the total variance.

A measure of relative blue reflectance (blue score) was determined for each individual by taking

the mean blue channel value (B) and the mean red channel value (R) for both wing regions and

calculating:

$$BR = (B-R)/(B+R)$$

This gives a standardised score of how blue an individual is, with BR = 1 being the 'bluest', and

BR = -1 being the 'reddest' (Figure S1 B, C).

Estimation of 'yellow bar' allele frequencies

Allele frequencies for the yellow hindwing bar were estimated based on phenotype for both species. This

was done for all sampling sites in Colombia and Panama with five or more individuals. The 'yellow bar'

phenotype was scored categorically according to Mallet (1986), who showed that this phenotype

segregates in the same way for both *Heliconius erato* and *H. melpomene*. Variation in the yellow bar

across this hybrid zone is controlled by three alleles: The North Colombian yellow bar allele (Y), the

West Colombian yellow bar allele (y_{wc}) and the Central American yellow bar allele (y_{ca}). Individuals of

both species with a yellow bar on both sides of the wing (Figure 1A) have genotype $y_{ca}y_{ca}$. Individuals

lacking a yellow bar (Figure 1B) have genotype YY. Individuals with the "shadow bar" phenotype, where

the outline of the bar can be seen on the underside of the hindwing without any yellow pigment, and

without a bar on the upper side of the hindwing, have genotype Yy_{wc} or Yy_{ca}. Individuals with a yellow

bar on the underside of the hindwing (Figure 1C) have genotype $y_{wc}y_{ca}$ or $y_{wc}y_{wc}$. As two of the four

phenotypes can be produced by two different allele combinations we inferred the allele frequencies at

each locality for each species assuming Hardy-Weinberg equilibrium for the three alleles. The frequency

of Y could be directly observed from both its heterozygous and homozygous phenotypes. The frequency

of y_{ca} could be inferred from the frequency of its homozygous phenotype, allowing us to infer the

frequency of y_{wc}. We focus on the y_{wc} allele for the remainder of the paper, as this underlies the yellow

bar phenotype seen in the iridescent forms of both species, and appears to be lost across the same hybrid

zone over which iridescence is lost. This provides us with the opportunity to directly compare clines in

Mendelian and polygenic traits.

Geographic cline analysis

We used geographic cline analysis to model patterns of clinal variation in (*i*) the mean iridescence score,

(ii) frequency of the yellow bar allele (y_{wc}), and (iii) mean admixture, estimated using NGSadmix, across

both hybrid zones. We assumed two parental populations here, as our analyses of population structure

reveal two genetic clusters in each species, despite there being three overlapping colour pattern races.

Specifically, we fitted three alternative geographic cline models (Szymura & Barton, 1991; Szymura &

Barton, 1986) using ANALYSE v1.30 (Barton & Baird, 2002). Sampling sites with fewer than five

individuals were excluded from the cline analyses, leaving 529 *H. erato* and 126 *H. melpomene*. Blue

scores were normalised to a new range of 0 to 1 (Fig S1 B, C) as required by the software. Distances

between sampling sites were estimated using the great circle distance, calculated using the

'hzar.map.greatCircleDistance' function in the R package HZAR (Derryberry, Derryberry, Maley, &

Brumfield, 2014).

ANALYSE fits cline models to marker loci and/or quantitative trait data, and can be used to

compare the fit of three alternative cline models to either population means (used for iridescence and

admixture scores) or frequency data (used for the yellow bar allele). The simplest model is a sigmoid

cline described by a hyperbolic tangent (Szymura & Barton, 1986). The other two more complex models

are 'stepped' clines. They consist of a central sigmoid step flanked by two exponential tails that describe

the pattern of introgression from the centre into the foreign genepool; θ is the rate of decay, and the

strength of the barrier to gene flow, B, can be estimated as the ratio between the difference in the allele

frequency and the initial gradient in allele frequency with distance x at the edges of the central segment.

In the symmetrical 'Sstep' model, θ and B are equal on both sides. In the asymmetrical 'Astep' model, the

pattern of introgression is different on the left and right side.

ANALYSE uses the Metropolis algorithm to search the likelihood surface to find the ML solution

to the model. To ensure that the likelihood surface was thoroughly explored, independent runs were

conducted using a range of initial parameter estimates. After obtaining maximum likelihood (ML)

solutions for the three cline models, the most likely model was identified using Likelihood Ratio Tests.

As the minimum and maximum mean allele frequency or trait values ($p(z)_{min}$, $p(z)_{max}$) were allowed to

vary in the tails of the cline, the sigmoid, Sstep and Astep models were described by 2 (c, w), 4 (c, w, θ,

B) and 6 parameters (c, w, θ_0, θ_1, B_0, B_1), respectively.

After model selection, support limits were estimated for each parameter in the ML model. Starting

with the optimum fit, and constraining the values of all other parameters, the likelihood surface for

individual parameters were explored by making 10,000 random changes of their value. The range of

estimates that was within 2 log-likelihood units of the maximum estimate was taken as the support limit

for that parameter, and is approximately equivalent to a 95% confidence interval.

Coincidence of cline centres (c) and concordance of cline widths (w) were tested using the

composite likelihood method (Kawakami, Butlin, Adams, Paull, & Cooper, 2009; Phillips, Baird, &

Moritz, 2004). The method involves obtaining a composite ML score for a given parameter (ML_{comp}) and

comparing it with the sum of the ML estimates obtained for each profile (ML_{sum}). ML_{comp} was obtained by

constructing a log-likelihood profile (10 km intervals for c and w, between 0 km and 1000 km) with all

other parameters allowed to vary, summing the profiles, and obtaining the ML estimate; ML_{sum} was

obtained by summing the ML estimates from each profile. If clines are not coincident or concordant,

ML_{sum} is significantly smaller than ML_{comp}, as determined by a chi-squared test ($\alpha = 0.05$) with n-1

degrees of freedom, where n is the number of traits. One complication with this method for comparing

cline parameters is that the profiles for each trait must be built using the same model. Although the more

complex Sstep and Astep models are a significantly better fit than the sigmoid model, the parameters

estimates for the cline centre and cline width were similar regardless of the model fit (see results).

Therefore, all likelihood profiling was conducted with the sigmoid model.

To estimate the strength of selection acting on y_{wc}, the following equation was used from Barton

and Gale (1993):

303                              $$s^* = (1.782\sigma/\omega)^2$$

Where s^* is the difference in mean fitness between populations at the edge of the cline, and

populations at the centre. This demonstrates the mean strength of effective selection on loci underlying a

trait required to maintain a cline of width (w), given the dispersal distance per generation (σ). Dispersal

estimates were taken from Mallet et al. (1990) and Blum (2002).

Tests for concordance of clines using regression analysis

In addition to geographic cline analysis, we also used the regression procedure outlined in

Nürnberger et al. (1995) as a method for testing for the concordance of clines within and between the

species. Concordance of clines is predicted to result in a linear regression of population means or allele

frequencies between characters i and j. Alternatively, non-concordant clines should show a deviation from

linearity that can be described by a quadratic polynomial. We compared the fits of linear and quadratic

models to each pair of characters, including the mean admixture score, frequency of the y_{wc} alleles and

the mean iridescence score within a species using custom script in R. Because data were collected for

each species in the same location, we could also use this analysis to compare clines in the same traits

between *H. erato* and *H. melpomene*, with the exception of the admixture score because few sites

included genetic data for both species. Significance of the deviation from linearity was determined by

comparing the F-ratios of the quadratic and linear fits.

Results

Population Structure

We investigated population structure using genome-wide SNP data in the programs NGSadmix, to

estimate ancestry proportions from a varying number of genetic clusters (*K*), and PCAngsd to confirm

population clustering by principal components (PCA). This revealed different patterns of population

structure between the co-mimics. In *H. erato,* NGSadmix supported *K*=2 (SI Figure S3B), representing a

"Panama-like" and a "Colombia-like" genetic background (Figure 2B), with individuals of consistently

mixed ancestry found in the site closest to the centre of the iridescence cline. Introgression from

Panamanian populations could be detected in northern Colombian populations. The PCA supported this,

with three clusters separated by geography along the first axis of variation, representing the Colombian

populations, the Panamanian populations, and individuals with mixed ancestry and intermediate levels of

iridescence clustered between them (Figure 2C). PC1 explained 5.84% of genetic variation in *H. erato,*

with all subsequent eigenvectors explaining 0.7% or less of the genetic variation (SI Figure S4).

NGSadmix also supported *K*=2 for *H. melpomene* (although *K*=1 cannot be tested, SI Figure

S3D), but revealed a less straightforward population structure. While a "Colombia-like" genetic

background could be seen, Panamanian individuals showed mixed ancestry, with the exception of four

individuals from the site closest to the centre of the iridescence cline (Figure 2D). This is supported by the

PCA. PC1 explained 5.28% of genetic variation, separating Colombian and Panamanian individuals.

Individuals with intermediate levels of iridescence do not form an intermediate cluster between

Panamanian and Colombian individuals, as is seen in *H. erato* (Figure 2E). The percent of genetic

variation explained by PC1 and subsequent principal components show a more uniform distribution than

in *H. erato* (SI Figure S4) consistent with weaker population structure.

Given the support for two genetic clusters, we compared the levels of differentiation between

these populations using SNPs from individuals either side of the hybrid zone in southern Panama.

Genome-wide average Hudson's F_{ST} was estimated for each species, using the ratio of averages approach.

This revealed that genome-wide divergence across the hybrid zone is greater in *H. erato* (F_{ST}=0.188),

compared to *H. melpomene* (F_{ST}=0.0739). The difference in genetic structure is also apparent in the plots

of the pairwise genetic distance between sampling locations, plotted against their geographic distance. In

H. erato, within-race comparisons that span distances of 195 – 325 km show a range of F_{ST} values

between 0.063 – 0.129. However, between-race comparisons made over a similar range of distances (188

– 345 km) have substantially higher F_{ST} (0.226 – 0.271), suggesting that the genetic structure is much

stronger than would be expected based on geography alone (Figure 3). The pattern in *H. melpomene* is

very different, as the between-race comparisons span a similar range of F_{ST} values to the within-race

comparisons.

Phenotypic Variation

Strong phenotypic variation was observed across our range of sampling sites, with some difference

apparent between *H. erato* and *H. melpomene* (SI Figure S2). The West Colombian yellow bar allele (y_{wc})

was fixed in all Colombian sampling sites, apart from at some of the northernmost Colombian sampling

sites near Bahía Solano (BS; SI Figure S2 C, D.). In *H. melpomene*, the frequency of y_{wc} gradually

decreased, and persisted at comparable frequencies to the North Colombian yellow bar allele (Y) for ~200

km, before the Central American yellow bar allele (y_{ca}) became predominant (SI Figure S2D). In contrast,

in *H. erato* Y became the predominant allele, with y_{ca} approaching fixation towards the end of the transect

(SI Figure S2C).

In both species the blue score, used as a proxy measure for iridescence, decreased across the

transect (SI Figure S2 A,B). The colour measurements used to calculate the blue score were highly

repeatable ($p<0.001$ for both red and blue values in both wing patches measured, Table S5). The bluest *H.*

melpomene individuals were less blue than the bluest *H. erato* (SI Figure S2), which is consistent with

reflectance spectrometry data from *H. erato cyrbia* and *H. melpomene cythera* (Parnell et al., 2018)

Clinal variation within species

Cline fitting revealed that an asymmetrical stepped cline best described the variation in iridescence in *H. erato*, with a steeper right tail, which continually declines away from the cline centre. Neither stepped model was a significantly better fit than sigmoidal clines for the yellow bar in *H. erato*, and both colour traits in *H. melpomene* (Table 1; SI Table S6). For the admixture proportion, an asymmetrical stepped cline model was the best fit in *H. erato* (Table 1, SI Table S6), with a steeper right tail, similar to the iridescence cline, whereas the sigmoid model was the best fit for *H. melpomene.*

Overall, our analysis indicates that the clines in iridescence, yellow bar and the admixture were highly similar within both of the species. Likelihood profiling revealed that we could not reject the null hypothesis that both iridescence and the yellow bar clines had coincident centres and concordant widths within both species. In *H. erato*, the parameters for the cline in the admixture score were different from the clines in iridescence (both centre and width) and yellow bar (centre only) (Table 2), though the differences were quite subtle. For *H. melpomene*, likelihood profiling indicated that neither the width nor centre of the admixture cline differed from the phenotypic clines; inspection of the likelihood surfaces indicates that power to reject the hypothesis of coincidence and concordance were low, as the profiles were flat across a broad range of the parameter space (Figures 3, S5). This could be due to weaker/non-clinal population structure, or the sparse sampling of genomic data within *H*. *melpomene.*

The similarity of clines within the two species was supported by pairwise regression analysis, as the linear model was always a better description of the data than the quadratic polynomial (Figure S6). For *H. erato*, the linear model explained between 98% and 99% percent of the variation relationship between for all pairs of characters, and had a higher F-ratio than the polynomial quadratic, which explained a similar amount of the variation in the data (Table S7). The results were the same for *H. melpomene,* except that the quadratic fit was often a much poorer fit than the linear fit (Table S7).

 Comparison of clines between H. melpomene and H. erato

In contrast with the similar patterns of clinal variation within species, our profiling and regression

analyses revealed striking differences in the cline shape between the species. For both iridescence, and

yellow bar, the ML estimates of the cline width were roughly four times wider in *H. melpomene* than in

H. erato (Table 1, Figure 4)*.* For both traits, the peaks of the likelihood profiles did not overlap (Figure

5), with the difference being significant for iridescence ($p = 0.01$). Although the difference was not

significant for yellow bar, because the change in likelihood was not as dramatic across the profile for that

trait ($p = 0.19$), the regression analysis indicated that clines were not concordant as the quadratic model

was a far better fit to the regression of the frequency of y_{wc} between *H. melpomene* and *H. erato* (Figure

S6, Table S7). Both likelihood profile analysis and regression analysis indicated that the admixture clines

were concordant (Figures 3, S6), but again had low power to detect any difference due to the coarse

geographic sampling in both species.

In contrast with the difference in cline widths, clines for each trait tended to have highly similar

centres, indicating that they were positioned in roughly the same geographic area (Figure 4, Table 2). This

is clearly observed in the likelihood profiles for each trait, as the -Lnl tended to peak over a relatively

broad area, between 400 and 600 km along the transect (SI Figure S5). For all three traits the difference in

the location of the peak likelihoods was not significantly difference (P ranging from 0.22 - 0.62)

Given the differences in cline width between the species, we estimated the effective selection ($s*$)

on y_{wc} across the hybrid zone in both species using the ML estimates and support limits of cline width

(Table 1) and the dispersal estimates from Mallet et al. (1990) of 2.6 km for *H. erato* and 3.7 km for *H.*

melpomene. Selection estimates were 0.00203 (0.00102–0.00427) for *H. erato*, and 0.000213 (0.000165–

0.000303) for *H. melpomene*. Blum (2002) estimates higher dispersal for *H. erato*, 10 km, which increased

the value of *s** to 0.0300 (0.0151–0.0632). Given that the widths of the yellow bar and iridescence clines

were not different within each species, similar estimates were found for iridescence. For *H. melpomene*,

s*=0.000200 (0.000135–0.000300), for *H. erato*, *s**= 0.00208 (0.00167–0.00267) if the dispersal distance

is 2.6 km, and 0.0307 (0.0247–0.0395) if the dispersal distance is 10 km. However, it should be noted that

in the case of iridescence, $s*$ is the average strength of selection acting across loci controlling iridescence.

Discussion

Our analysis of parallel hybrid zones in the co-mimics *H. erato* and *H. melpomene* has revealed

similarities, as well as striking differences in colour trait divergence between the species. Consistent with

the predictions of the mimicry hypothesis, the clines in yellow bar and iridescence are highly coincident

within and between species, suggesting that they are maintained by the same selective pressure. In contrast,

the width of the clines in both colour traits vary substantially between the species, being far wider in *H.*

melpomene. The difference in cline widths is probably due, at least in part, to differences in the strength of

direct selection acting on colour variation between *H. melpomene*, and *H. erato.* However, differences in

population structure and levels of genomic differentiation indicate that species-specific factors, such as

different population histories, dispersal rates, and strength of reproductive isolation between the subspecies

may also contribute to the different cline widths between the species.

Comparing clines within species

Our geographic cline analysis revealed that clines in mean iridescence and the y_{wc} allele frequency

had highly similar centres and widths within both of the species. While this is predicted to result from

direct selection on a warning colour pattern, similar clines could also arise as a correlated response to

selection if traits have a shared genetic basis, or if the loci that underlie them are physically linked (Price

& Langen, 1992). We are able to rule out these explanations for the similarity of the clines in *H. erato*, as

these colour traits segregate independently in F_2 crosses made between iridescent and non-iridescent races

(Brien et al., 2018). As the colour pattern traits studied here very different genetic architectures, with the

yellow bar being controlled by a single major-effect locus (Joron et al., 2006; Mallet, 1986; Nadeau et al.,

2016), and iridescence being controlled by multiple genes (Brien et al., 2018), it is highly unlikely that the

clines in one of the colour traits could arise as a correlated response to selection acting on the other.

Another alternative explanation for the similar clines within the species, aside from direct selection

acting on each trait, is that the clines are maintained by a permeable, but genome-wide barrier to gene

flow between the subspecies. When reproductive isolation involves a large number of loci, or is very

strong, selection against unfit hybrid offspring can generate a barrier to gene flow that can impact the

spread of even neutral alleles across a hybrid zone (Barton & Gale, 1993). For a trait that is also under

direct selection, the importance of the overall barrier in shaping a cline depends on the strength of direct

selection acting on the trait relative to the strength of indirect selection resulting from selection at other

barrier loci. For example, if the strength of indirect selection acting on a trait is much greater than direct

selection, then the cline shape will be more informative about the overall barrier strength, and tell us

nothing about the strength of direct selection. In situations where this is the case, clines in the trait should

show a "stepped", rather than sigmoid shape. In *H. erato*, the two colour pattern clines are coincident, and

the best fitting cline model for variation in iridescence is stepped, which would indicate that indirect

selection plays some role in shaping the cline (Kruuk et al., 1999). In contrast, the simple sigmoidal cline

fits best for the yellow bar. Finally, while we do see stepped clines in *H. erato,* they are asymmetrical,

with a left tail closely resembling that of the sigmoidal cline, and a much steeper right tail. It is possible

that these tails reflect genuine asymmetry, due to hybrid zone movement, which has been predicted and

documented in these species (Blum, 2002; Mallet, 1986; Thurman, Szejner-Sigal, & McMillan, 2019).

Although it is difficult to determine the overall importance and indirect selection in shaping these clines,

it is unlikely that all of our results can be explained purely in terms of indirect selection.

Given the abundant evidence for the role of direct selection in shaping colour pattern variation across

the genus *Heliconius*, it is likely that it plays at least some role in shaping variation in these species.

Under a scenario where the colour pattern clines are maintained by a balance between migration and

divergent ecological selection, similar cline centres arise when both traits experience the same source of

selection, or when different ecological gradients change in approximately the same location (Barton &

Hewitt, 1985). In *Heliconius,* local warning colour patterns are maintained by predator-mediated positive

frequency-dependent selection, with rare colour morphs experiencing increased predation (Benson, 1972;

Dell'aglio, Stevens, & Jiggins, 2016; Langham, 2004; Mallet & Barton, 1989). The centre of colour

pattern clines could represent the location where the most effective warning pattern shifts to that of a

neighbouring subspecies. The coincidence of cline centres for iridescence and y_{wc}, which is observed in

both *H. melpomene* and *H. erato*, suggests that both traits contribute to the warning signal.

In *H. melpomene*, the width and centre of the admixture proportion cline was not significantly

different to the colour pattern clines. However, variation in admixture proportions had a poor fit to any of

the cline models (Figures 2, 3, S5), illustrated by the large confidence intervals (Table 1). This is in part

due to coarse sampling, but can also be explained by the less defined population structure in this species.

Clear phenotypic intermediates in the hybrid zone are not of mixed ancestry (Figures 3, 4). This suggests

that divergence in iridescence in this species is not tightly coupled with genome-wide differentiation. The

broad phenotypic clines that we see are not characteristic of the steep, stepped clines which result from

strong LD between selected loci and indirect selection (e.g. Szymura & Barton, 1991), and are more

likely due to weak selection and/or isolation-by-distance.

Comparing clines between species

Although patterns of clinal variation are very similar within species, we observed substantial

differences between *H. melpomene* and *H. erato*. This is in contrast to what is expected, based on the

strong existing evidence that colour pattern convergence in this pair of co-mimics is driven by Müllerian

mimicry - a common positive frequency-dependent selection pressure based on predator learning. The

coincidence of cline centres in colour pattern traits between the co-mimics is consistent with this

hypothesis, as it suggests that variation in both species is structured by the same agent of selection. While

the clines in yellow bar and iridescence are coincident between species, they are four times wider in *H.*

melpomene. This difference could result from (*i*) variation in the strength of direct or indirect selection

between the species, (*ii*) species-specific differences in the dispersal rate, (*iii*) different demographic

histories between the species, or a combination of these explanations.

First, the wider clines in *H. melpomene* could be a result of having a greater dispersal capability.

Direct estimates of dispersal are difficult in *Heliconius* butterflies due to most dispersal occurring soon

after adult eclosion (Mallet, 1986a). The most reliable estimates are thought to be those made using cline

theory and patterns of linkage disequilibrium (Blum, 2002; Mallet et al., 1990), including the only direct

comparison of *H. erato* and *H. melpomene* (Mallet et al., 1990). This study reports higher dispersal

distances in *H. melpomene*. However, our estimates of the selection coefficient s^* (Barton & Gale, 1993)

show that even if this higher dispersal rate is taken into account, colour pattern traits in *H. melpomene*

appear to be under much weaker selection. Other studies on parallel hybrid zones between neighbouring

races in this species pair show that *H. melpomene* tend to have wider clines than *H. erato*, but not to the

degree seen in the present study (Mallet et al., 1990; Salazar, 2012). *H. melpomene* displays less vivid

iridescence than its co-mimic, and the colour difference between iridescent and non-iridescent *H.*

melpomene is less pronounced than the colour difference between *H. erato* races (Parnell et al., 2018,

Figure 1). Hybrid phenotypes are therefore less distinct from the parental populations in *H. melpomene*

which could result in weaker selection against hybrid offspring.

The difference in divergence of one of the colour traits, namely iridescence, could also explain why

clines differ in shape between the co-mimics. The main predators of *Heliconius* butterflies are thought to

be birds of the tyrant flycatcher (Tyrannidae) and jacamar (Galbulidae) families (Jiggins, 2017), hence

bird predation is expected to be the main driver of mimicry and phenotypic convergence between species.

Previous work modelling bird visual systems has shown that birds can discriminate between the iridescent

blue in *H. erato* and *H. melpomene* (Parnell et al., 2018). However, iridescence in *H. melpomene* is not as

bright as in *H. erato*, which means that the visibility of the trait to selection may also vary between the

species. This may weaken the overall mimetic signal in *H. melpomene,* which may also influence the

strength of selection acting on the yellow bar.

Another possible factor that may explain the differences in cline widths between the species is

that they may have experienced very different demographic histories. The inclusion of genomic data in

our study revealed a striking difference in the level of population structure across these parallel hybrid

zones. Specifically, we found strong divergence across the *H. erato* hybrid zone in contrast with the very

weak structure across the *H. melpomene* hybrid zone. The defined population structure in *H. erato* is

typically associated with populations that have diverged in allopatry, followed by secondary contact. This

scenario can lead to genetic discontinuity and coincidence of clines in multiple traits (Barton, 1983),

along with strong genome-wide reproductive isolation, meaning that indirect selection can play a greater

role in the maintenance of geographic clines. It is also possible that strong selection acting on a

quantitative trait in *H. erato* could be responsible for the formation of a genome-wide barrier to gene flow

(Feder et al., 2012), although conservative estimates suggest iridescence is not polygenic enough to act as

such a barrier (Brien et al., 2018). In contrast, the relatively low genetic structure in *H. melpomene* is

usually associated with a primary intergradation scenario, where hybrid zones form due to divergent

selection acting across a strong environmental gradient. Because primary hybrid zones form in the face of

continuous gene flow, other barriers cannot evolve in isolation, meaning that direct selection on

phenotypic traits must alone overcome migration. This makes it much harder for sharp clines to become

established. Future studies of the historical demography of these species may shed more light on the role

of history in shaping phenotypic traits associated with mimicry.

Conclusions

Examples of parallel evolution are celebrated as some of the best evidence for the power of

natural selection in driving local adaptation. Most studies of parallel evolution tend to focus on

understanding the similarities between populations and species subject to the same selective pressures.

Despite the striking parallelism at the level of the phenotype, including simple and complex colour traits,

the Müllerian co-mimics *H. erato* and *H. melpomene* show striking differences in how trait variation is

structured across geography, which would not be apparent without detailed sampling and analysis across

their distribution. Although mimicry has almost certainly been the primary driver of parallel evolution in

this system, other factors are needed to explain patterns of phenotypic variation, both within and between

species. More focus on phenotypic differences may provide new insight into the processes underlying

parallel evolution, and may help us to understand the factors that limit adaptation in general.

Acknowledgements

Thanks to the governments of Panama and Colombia (ANLA- Permit 0530) for giving permission to

collect butterfly specimens. Thanks to the McMillan and Jiggins labs for providing access to samples.

Thanks also to Patricio Salazar, Juan Enciso, Juan Camilo Dumar, Melanie Brien, Carlos Arias, Agata

Surma and others in Panama and Colombia, in particular the residents of Jaqué, Darién, for help with

logistics and collecting in the field. Thanks to Roger Butlin for valuable comments on this manuscript.

This work was funded by the UK Natural Environment Research Council (NERC) through an

Independent Research Fellowship (NE/K008498/1) to NJN, and by The Royal Society through an

International Exchange Scheme grant. EVC was funded by the NERC doctoral training partnership,

ACCE. CS and CP were funded by COLCIENCIAS (Grant FP44842-5-2017).

References

Abràmoff, M. D., Magalhães, P. J., & Ram, S. J. (2004). Image processing with imageJ. *Biophotonics*
    International, *11*(7), 36–41. https://doi.org/10.1117/1.3589100

Arias, C. F., Muñoz, A. G., Jiggins, C. D., Mavárez, J., Bermingham, E., & Linares, M. (2008). A hybrid
    zone provides evidence for incipient ecological speciation in Heliconius butterflies. *Molecular*
    Ecology, *17*(21), 4699–4712. https://doi.org/10.1111/j.1365-294X.2008.03934.x

Barton, N H. (1983). Multilocus Clines. *Evolution*, *37*(3), 454–471. https://doi.org/10.2307/2408260

Barton, N H, & Hewitt, G. M. (1985). Analysis of Hybrid Zones. *Annual Review of Ecology and*

Systematics, Vol. 16, pp. 113–148. https://doi.org/10.1146/annurev.es.16.110185.000553

Barton, Nicholas H., & Baird, S. J. E. (2002). *Analyse. Version 1.30.* Retrieved from

http://archive.bio.ed.ac.uk/software/Mac/Analyse/index.html

Barton, Nicholas H., & Gale, K. S. (1993). Genetic analysis of hybrid zones. In R G Harrison (Ed.),

Hybrid zones and the evolutionary process (pp. 13–45). Retrieved from

https://books.google.co.uk/books?id=aFJFkVKskYIC&pg=PA13&source=gbs_toc_r&cad=4#v=one

page&q&f=false

Baxter, S W, Johnston, S. E., & Jiggins, C. D. (2009). Butterfly speciation and the distribution of gene

effect sizes fixed during adaptation. *Heredity*, *102*, 57–65. https://doi.org/10.1038/hdy.2008.109

Baxter, Simon W, Davey, J. W., Johnston, J. S., Shelton, A. M., Heckel, D. G., Jiggins, C. D., & Blaxter,

M. L. (2011). Linkage Mapping and Comparative Genomics Using Next-Generation RAD

Sequencing of a Non-Model Organism. *PLoS ONE*, *6*(4), e19315.

https://doi.org/10.1371/journal.pone.0019315

Benson, W. (1972). Natural Selection for Mullerian Mimicry in Heliconius erato in Costa Rica. *Science*,

176(4037), 936–939. Retrieved from http://www.sciencemag.org/content/176/4037/936.short

Bhatia, G., Patterson, N., Sankararaman, S., & Price, a. L. (2013). Estimating and interpreting F_{ST}: The

impact of rare variants. *Genome Research*, *23*(9), 1514–1521. https://doi.org/10.1101/gr.154831.113

Blum, M. J. (2002). Rapid movement of a Heliconius hybrid zone: evidence for phase III of Wright's

shifting balance theory? *Evolution; International Journal of Organic Evolution*, *56*(10), 1992–1998.

https://doi.org/10.1554/0014-3820(2002)056[1992:RMOAHH]2.0.CO;2

Brien, M. N., Enciso-Romero, J., Parnell, A. J., Salazar, P. A., Morochz, C., Chalá, D., … Nadeau, N. J.

(2018). Phenotypic variation in Heliconius erato crosses shows that iridescent structural colour is

sex-linked and controlled by multiple genes. *Interface Focus*, *9*(1), 20180047.

https://doi.org/10.1098/rsfs.2018.0047

Brown, K. S. (1981). The Biology of Heliconius and Related Genera. *Annual Review of Entomology*,

26(1), 427–457. https://doi.org/10.1146/annurev.en.26.010181.002235

Challis, R. J., Kumar, S., Dasmahapatra, K. K., Jiggins, C. D., & Blaxter, M. (2016). Lepbase: The

Lepidopteran genome database. *BioRxiv*, 056994. https://doi.org/10.1101/056994

Davey, J. W., Chouteau, M., Barker, S. L., Maroja, L., Baxter, S. W., Simpson, F., … Jiggins, C. D. (2016). Major Improvements to the Heliconius melpomene Genome Assembly Used to Confirm 10 Chromosome Fusion Events in 6 Million Years of Butterfly Evolution. *G3 Genes|Genomes|Genetics*, *6*(3), 695–708. https://doi.org/10.1534/g3.115.023655

Dell'aglio, D. D., Stevens, M., & Jiggins, C. D. (2016). Avoidance of an aposematically coloured butterfly by wild birds in a tropical forest. *Ecological Entomology*, *41*(5), 627–632. https://doi.org/10.1111/een.12335

Derryberry, E. P., Derryberry, G. E., Maley, J. M., & Brumfield, R. T. (2014). <scp>hzar</scp> : hybrid zone analysis using an R software package. *Molecular Ecology Resources*, *14*(3), 652–663. https://doi.org/10.1111/1755-0998.12209

Endler, J. A. (1977). *Geographic variation, speciation, and clines*. Princeton, New Jersey: Princeton University Press.

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: A simulation study. *Molecular Ecology*, *14*(8), 2611–2620. https://doi.org/10.1111/j.1365-294X.2005.02553.x

Feder, J. L., Gejji, R., Yeaman, S., & Nosil, P. (2012). Establishment of new mutations under divergence and genome hitchhiking. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *367*(1587), 461–474. https://doi.org/10.1098/rstb.2011.0256

Flaxman, S. M., Wacholder, A. C., Feder, J. L., & Nosil, P. (2014). Theoretical models of the influence of genomic architecture on the dynamics of speciation. *Molecular Ecology*, *23*(16), 4074–4088. https://doi.org/10.1111/mec.12750

Hudson, R. R., Slatkin, M., & Maddison, W. P. (1992). Estimation of levels of gene flow from DNA sequence data. *Genetics*, *132*(2), 583–589. https://doi.org/PMC1205159

Jiggins CD. (2017). *The ecology and evolution of Heliconius butterflies: a passion for diversity.* Oxford, UK.

Joron, M., Papa, R., Beltrán, M., Chamberlain, N., Mavárez, J., Baxter, S., … Jiggins, C. D. (2006). A

conserved supergene locus controls colour pattern diversity in Heliconius butterflies. *PLoS Biology*, *4*(10), 1831–1840. https://doi.org/10.1371/journal.pbio.0040303

Kawakami, T., Butlin, R. K., Adams, M., Paull, D. J., & Cooper, S. J. B. (2009). Genetic analysis of a chromosomal hybrid zone in the Australian morabine grasshoppers (Vandiemenella, viatica species group). *Evolution*, *63*(1), 139–152. https://doi.org/10.1111/j.1558-5646.2008.00526.x

Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A., & Mayrose, I. (2015). Clumpak: A program for identifying clustering modes and packaging population structure inferences across K. *Molecular Ecology Resources*, *15*(5), 1179–1191. https://doi.org/10.1111/1755-0998.12387

Kronforst, M. R., Young, L. G., Kapan, D. D., Mcneely, C., Neill, R. J. O., & Gilbert, L. E. (2006). Linkage of butterfly mate preference and wing color preference cue at the genomic location of wingless. *Proceedings of the National Academy of Sciences*, *103*(17), 6575–6580.

Kruuk, L. E. B., Baird, S. J. E., Gale, K. S., & Barton, N. H. (1999). A Comparison of Multilocus Clines Maintained by Environmental Adaptation or by Selection Against Hybrids. *Genetics*, *153*, 1959–1971.

Langham, G. M. (2004). Specialized Avian Predators Repeatedly Attack Novel Color Morphs of Heliconius Butterflies. *Evolution*, *58*(12), 2783–2787.

Le Corre, V., & Kremer, A. (2012). The genetic differentiation at quantitative trait loci under local adaptation. *Molecular Ecology*, *21*(7), 1548–1566. https://doi.org/10.1111/j.1365-294X.2012.05479.x

Mallet, J. (1989). The Genetics of Warning Colour in Peruvian Hybrid Zones of Heliconius erato and H. melpomene. *Proceedings of the Royal Society B: Biological Sciences*, *236*(1283), 163–185. https://doi.org/10.1098/rspb.1989.0019

Mallet, James. (1986a). Dispersal and gene flow in a butterfly with home range behavior: *Heliconius erato* (Lepidoptera, Nymphalidae). *Oecologia*, *68*(2), 210–217. https://doi.org/10.1007/BF00384789

Mallet, James. (1986b). Hybrid zones of Heliconius butterflies in Panama and the stability and movement of warning colour clines. *Heredity*, *56*(2), 191–202. https://doi.org/10.1038/hdy.1986.31

Mallet, James. (1993). Speciation, raciation, and color pattern evolution in Heliconius butterflies:

evidence from hybrid zones. In Richard G Harrison (Ed.), *Hybrid zones and the evolutionary*

process (pp. 226–260). New York: Oxford University Press.

Mallet, James, Barton, N., Gerardo Lamas, M., Jose Santisteban, C., Manuel Muedas, M., & Eeley, H.

(1990). Estimates of selection and gene flow from measures of cline width and linkage

disequilibrium in Heliconius hybrid zones. *Genetics*, *124*, 921–936.

Mallet, James, & Barton, N. H. (1989). Strong Natural Selection in a Warning-Color Hybrid Zone.

Evolution, *43*(2), 421–431. https://doi.org/10.2307/2409217

Martin, A., Papa, R., Nadeau, N. J., Hill, R. I., Counterman, B. A., Halder, G., … Reed, R. D. (2012).

Diversification of complex butterfly wing patterns by repeated regulatory evolution of a Wnt ligand.

Proceedings of the National Academy of Sciences, *109*(31), 12632–12637.

https://doi.org/10.1073/pnas.1204800109

Martin, S. H., Dasmahapatra, K. K., Nadeau, N. J., Slazar, C., Walters, J. R., Simpson, F., … Jiggins, C.

D. (2013). Genome-wide evidence for speciation with gene flow in Heliconius butterflies. *Genome*

Research, *23*, 1817–1828. https://doi.org/10.1101/gr.159426.113.

Meisner, J., & Albrechtsen, A. (2018). Inferring population structure and admixture proportions in low-

depth NGS data. *Genetics*, *210*(2), 719–731. https://doi.org/10.1534/genetics.118.301336

Nadeau, N. J. (2016). Genes controlling mimetic colour pattern variation in butterflies. *Current Opinion*

in Insect Science, Vol. 17, pp. 24–31. https://doi.org/10.1016/j.cois.2016.05.013

Nadeau, N. J., Pardo-Diaz, C., Whibley, A., Supple, M. A., Saenko, S. V., Wallbank, R. W. R., …

Jiggins, C. D. (2016). The gene cortex controls mimicry and crypsis in butterflies and moths.

Nature, *534*(7605), 106–110. https://doi.org/10.1038/nature17961

Nadeau, N. J., Ruiz, M., Salazar, P., Counterman, B., Medina, J. A., Ortiz-Zuazaga, H., … Papa, R.

(2014). Population genomics of parallel hybrid zones in the mimetic butterflies, H. melpomene and

H. erato. *Genome Research*, *24*(8), 1316–1333. https://doi.org/10.1101/gr.169292.113

Nürnberger, B., Barton, N., MacCallum, C., Gilchrist, J., & Appleby, M. (1995). Natural Selection on

Quantitative Traits in the Bombina hybrid zone. *Evolution*, *49*(6), 1224–1238.

Parnell, A. J., Bradford, J. E., Curran, E. V., Washington, A. L., Adams, G., Brien, M. N., … Nadeau, N.

J. (2018). Wing scale ultrastructure underlying convergent and divergent iridescent colours in

mimetic *Heliconius* butterflies. *Journal of The Royal Society Interface*, *15*(141), 20170948.

https://doi.org/10.1098/rsif.2017.0948

Phillips, B. L., Baird, S. J. E., & Moritz, C. (2004). When Vicars Meet: A Narrow Contact Zone between

Morphologically Cryptic Phylogeographic Lineages of the Rainforest Skink, Carlia rubrigularis.

Evolution, *58*(7), 1536–1548.

Price, T., & Langen, T. (1992). Evolution of correlated characters. *Trends in Ecology & Evolution*, *7*(9),

307–310. https://doi.org/10.1016/0169-5347(92)90229-5

Reed, R. D., Papa, R., Martin, A., Hines, H. M., Counterman, B. a, Pardo-Diaz, C., … McMillan, W. O.

(2011). Optix Drives the Repeated Convergent Evolution of Butterfly Wing Pattern Mimicry.

Science, *333*(6046), 1137–1141. https://doi.org/10.1126/science.1208227

Rosenblum, E. B., & Harmon, L. J. (2011). "Same same but different": Replicated ecological speciation

at white sands. *Evolution*, *65*(4), 946–960. https://doi.org/10.1111/j.1558-5646.2010.01190.x

Rosser, N., Dasmahapatra, K. K., & Mallet, J. (2014). Stable Heliconius butterfly hybrid zones are

correlated with a local rainfall peak at the edge of the Amazon basin. *Evolution*, *68*(12), 3470–3484.

https://doi.org/10.1111/evo.12539

Salazar, P. A. (2012). *Hybridization and the genetics of wing colour-pattern diversity in Heliconius*

butterflies. PhD thesis. 119.

Skotte, L., Korneliussen, T. S., & Albrechtsen, A. (2013). Estimating individual admixture proportions

from next generation sequencing data. *Genetics*, *195*(3), 693–702.

https://doi.org/10.1534/genetics.113.154138

Supple, M. A., Hines, H. M., Dasmahapatra, K. K., Lewis, J. J., Nielsen, D. M., Lavoie, C., …

Counterman, B. A. (2013). Genomic architecture of adaptive color pattern divergence and

convergence in Heliconius butterflies. *Genome Research*, *23*, 1248–1257.

https://doi.org/10.1101/gr.150615.112.1248

Sweeney, A., Jiggins, C., & Johnsen, S. (2003). Polarized light as a butterfly mating signal. *Nature*,

423(6935), 31–32. https://doi.org/10.1038/423031a

Szymura, J M, & Barton, N. H. (1991). The genetic structure of the hybrid zone between the fire-bellied

toads *Bombina bombina* and *B. variegata*: comparisons between transects and between loci.

Evolution, *45*(2), 237–261. https://doi.org/10.2307/2408943

Szymura, Jacek M, & Barton, N. H. . (1986). Genetic Analysis of a Hybrid Zone Between the Fire-

Bellied Toads, Bombina bombina and B. variegata, Near Cracow in Southern Poland. *Evolution*,

40(6), 1141–1159.

Thurman, T. J., Szejner-Sigal, A., & McMillan, W. O. (2019). Movement of a Heliconius hybrid zone

over 30 years: A Bayesian approach. *Journal of Evolutionary Biology*, *32*(9), 974–983.

https://doi.org/10.1111/jeb.13499

Van Belleghem, S. M., Rastas, P., Papanicolaou, A., Martin, S. H., Arias, C. F., Supple, M. A., … Papa,

R. (2017). Complex modular architecture around a simple toolkit of wing pattern genes. *Nature*

Ecology and Evolution, *1*(3), 1–12. https://doi.org/10.1038/s41559-016-0052

Vines, T. H., Dalziel, A. C., Albert, A. Y. K., Veen, T., Schulte, P. M., & Schluter, D. (2016). Cline

coupling and uncoupling in a stickleback hybrid zone. *Evolution*, *70*(5), 1023–1038.

https://doi.org/10.1111/evo.12917

Westerman, E. L., VanKuren, N. W., Massardo, D., Tenger-Trolander, A., Zhang, W., Hill, R. I., …

Kronforst, M. R. (2018). Aristaless Controls Butterfly Wing Color Variation Used in Mimicry and

Mate Choice. *Current Biology*, *0*(0), 3469–3474. https://doi.org/10.1016/J.CUB.2018.08.051

Whitlock, M. C., & Schluter, D. (2009). *The Analysis of Biological Data*. Greenwood Village, Colorado:

Roberts and Company Publishers.

Data Accessibility Statement

Sequence data have been deposited in the European Nucleotide Archive with the project number

PREJEB32848. Code for implementing the regression-based tests for concordance can be found at

https://github.com/seanstankowski/Heliconius_MS. Phenotype measurements deposited at Dryad: XXX

Author Contributions

737 EVC and NJN conceived and designed the study. EVC, NJN, CPD, CAS and ML carried out field work.

738 EVC generated and analysed the data with the help of SS. EVC wrote the manuscript, and all co-authors

739 revised the manuscript.

Figures and tables

Figure 1 – Sampled populations in Colombia and Panama. Sites are labelled with abbreviations (further
information about sites and collections are in Table S1). Photographs show the phenotypes of mimetic
races of *H. erato* and *H. melpomene* from Central America (**A**), North Colombia (**B**), and Western
Colombia (**C**). For each pictured phenotype, the wings on the left-hand side show the ventral wing
pattern, and the wings on the right-hand side show the dorsal wing pattern. Approximate ranges for the
mimetic race pairs are outlined with dashed lines (Rosser, Phillimore, Huertas, Willmott, & Mallet,
2012). Populations that are included in the phenotypic analysis only are shown in grey, populations that
are included in both the phenotypic and genetic analysis are shown in black.
753

Figure 2 – Population structure across the hybrid zones in *H. erato* (**B,C**) and in *H. melpomene* (**D, E**).
Sampling locations across the hybrid zone. Approximate centre of the iridescence cline in *H. erato*
indicated by a dashed line.(**A**). Individual admixture proportions estimated using NGSadmix, with $k = 2$
(**B, D**). Each vertical bar represents an individual, bar colour represents the estimated proportion of
ancestry derived from population 1 (dark grey) or population 2 (light grey). Horizontal bars indicate the
population of origin, colours match those on the map. Principal components analysis (**C, E**). Colour of
points indicate the population of origin, as shown on the map.

Figure 3 – Relationship between geographic distance and genetic differentiation (genome-wide average F_{ST}) between sampling sites in *H. erato* (**A**) and *H. melpomene* (**B**). Pairwise comparisons are colour-coded to indicate comparisons between populations of the same colour pattern race (blue), between populations of different colour pattern races (yellow), and comparisons where one population is from the hybrid zone centre (green).

777   **Figure 4** – The best fitting geographic clines (dashed lines) of iridescence (**A, B**; blue), the West Colombian
778   yellow bar allele frequency (y_{wc}; **C, D**; yellow), and admixture proportions (**E, F**; red), across a transect of
779   sampling sites (points) for *Heliconius melpomene* (**A, C, E**) and *Heliconius erato* (**B, D, F**). The transect
780   begins (at 0 km) in the Queremal (Qu) locality, in the Cauca Valley region of Colombia.

Figure 5 – Likelihood profiles for the cline width for mean iridescence, frequency of yellow bar and

mean admixture score for *H. melpomene* (narrow dashed line) and *H. erato* wide dashed line. Profiles

were constructed using a step size of 10 km with all of the model parameters free to vary.

Table 1 – Cline parameter estimates for variation in iridescence, y_{wc} allele frequency, and admixture proportion across transects for *H. erato* and *H. melpomene,* which begin at the Queremal (Qu) locality. ML estimates for sigmoid models (Sig), symmetrical stepped models (Sstep), and asymmetrical stepped models (Astep) were estimated for each trait. If a model is a significantly better fit as determined by a likelihood ratio test (details in Table S6) it is denoted with *. Parameters are log-likelihood (LnL) cline centre (*c*), width (*w*), barrier strength for either side of stepped models (B_0/w, B_1/w), the rate of exponential decay for either tail (θ_0, θ_1).

Species	Trait	Model	LnL	Centre (km)	Width (km)	pmin	pmax	B_0/w	θ_0	B_1/w	θ_1
Heliconius erato	Iridescence	Sig	-38.62	537.38 (534.97-539.92)	101.66 (89.66-113.50)	0.101 (0.096-0.106)	0.901 (0.889-0.914)	--	--	--	--
		Sstep	-34.12	549.48 (508.87-562.98)	46.11 (38.78-74.86)	0.084 (0.020-0.092)	0.917 (0.888-0.913)	63.63 (37.24-80.76)	0.012 (0.0003-0.9998)	--	--
		Astep*	-27.34	546.38 (547.28-550.91)	78.62 (69.46-90.07)	0.089 (0.021-0.0903)	0.90 (0.089-0.92)	40.63 (38.69-56.57)	0.005 (0.002-0.006)	9.56E+09 (2E+06 – 1E+10)	0.73 (0.001-0.999)
	y_{wc}	Sig	-9.29	530.53 (515.48-543.70)	102.87 (70.90-145.19)	0.056 (0.013-0.088)	1.000 (0.989-1.00)	--	--	--	--
		Sstep	-9.83	530.53 (510.32-543.20)	102.81 (66.99-167.62)	0.056 (0.011-0.087)	0.999 (0.989-1.000)	7.44E+09 (195247-1E+10)	0.35 (0.00-0.99)	--	--
		Astep	-6.59	536.09 (523.98-547.17)	98.19 (70.48-145.52)	1.00E-04 (0.000-0.087)	1.00 (0.989-1.000)	8.41 (4.17-13.34)	0.075 (0.052-0.144)	6.86E+09 (3886-1E+10)	0.51 (0.000-0.991)
	Admixture proportion	Sig	-25.83	523.60 (521.26-525.76)	171.36 (167.33-175.56)	0.000017 (0.000015-0.010190)	0.9995 (0.996-0.9998)	--	--	--	--
		Sstep	-20.53	524.28	165.84	0.000012	0.99	1.01	0.98	--	--

Species	Trait	Model									
				(520.34-527.59)	(163.23-169.74)	(0.00001-0.00078)	(0.996-1.000)	(0.83-1.54)	(0.932-0.999)		
		Astep*	-13.63	536.79 (522.65-537.43)	101.30 (97.54-110.43)	0.021 (0.018-0.023)	0.99 (0.9997-1.000)	4144656896 (6775875-778645878)	0.11 (0.003-0.999)	1.88 (0.994-37.975)	0.37 (0.00-0.873)
Heliconius melpomene	Iridescence	Sig	-5.82	504.27 (474.68-532.89)	466.31 (380.85-567.89)	0.14 (0.12-0.17)	0.75 (0.70-0.79)	--	--	--	--
		Sstep	-3.62	553.87 (468.98-557.76)	145.565 (126.873-172.654)	0.0001 (0.00001-0.0118)	0.82 (0.79-0.83)	2.10 (0.17-12.34)	0.021 (0.001-0.046)	--	--
		Astep	-3.40	572.18 (542.77-584.88)	454.48 (398.87-528.72)	0.0001 (0.00001-0.0087)	0.72 (0.70-0.73)	1.72 (1.23-17.35)	0.17 (0.000-0.65)	136571824 (943923-232068365)	0.89 (0.67-0.999)
	y_{wc}	Sig	-6.70	649.14 (597.78-666.74)	451.92 (378.88-513.53)	0.0001 (0.0001-0.14)	1.000 (0.999-1.000)	--	--	--	--
		Sstep	-6.70	649.17 (587.88-668.80)	451.78 (382.81-498.02)	0.0001 (0.0001-0.14)	1.000 (0.999-1.000)	7857362432 (4730095-9998937088)	0.78 (0.29-0.998)	--	--
		Astep	-6.05	511.27 (505.82-527.64)	156.12 (135.76-190.54)	0.0001 (0.0001-0.14)	1.00 (0.999-1.000)	0.70 (0.12-0.999)	0.078 (0.001-0.64)	2958797824 (4729519 - 8986493729)	0.999 (0.000-0.999)
	Admixture proportion	Sig	-0.062	313.45 (126.86-363.68)	39.30 (0.43-123.81)	0.13 (0.0001-0.17)	1.000 (0.900-1.000)	--	--	--	--
		Sstep	-0.062	306.36 (150.64-313.85)	122.47 (0.37-152.64)	0.13 (0.0001-0.16)	1.000 (0.900-1.000)	7427586048 (1567365-9867457635)	0.32 (0.28-0.999)	--	--
		Astep	-0.062	311.74 (147.74-343.76)	59.34 (0.52-76.78)	0.13 (0.0001-0.16)	1.000 (0.900-1.000)	1764016384 (863863-8223565965)	0.79 (0.67-0.999)	8084597760 (878456-9864222189)	0.51 (0.000-0.97)

Table 2 – Likelihood ratio tests for coincidence (*c*) and concordance (*w*) of iridescence, yellow bar, and admixture proportion clines. ΔML is the test statistic, d.f. is degrees of freedom. The combination of clines being compared is noted under Trait(s).

Trait(s)	*c*			*w*		
	ΔML	d.f.	*P*	ΔML	d.f.	*P*
H. erato						
iridescence, yellow bar	0.75	1	0.39	0	1	1.00
iridescence, admixture proportion	11.22	1	< 0.001	16.85	1	< 0.001
iridescence, yellow bar, admixture proportion	11.93	2	0.003	20.47	2	< 0.001
H. melpomene						
iridescence, yellow bar	0.80	1	0.37	0.004	1	0.95
iridescence, admixture proportion	0.22	1	0.64	0.008	1	0.93
iridescence, yellow bar, admixture proportion	1.02	2	0.60	0.01	2	0.99
Both species						
iridescence	0.62	1	0.43	6.42	1	0.01
yellow bar	0.59	1	0.44	1.70	1	0.19
admixture proportion	0.22	1	0.64	0	1	1.00