1 **von Willebrand Factor D and EGF Domains is an evolutionarily conserved and required**
2 **feature of blastemas capable of multi-tissue appendage regeneration**
3
4

5 Leigh N.D.[1,2], Sessa S.[1], Dragalzew A.C.[3], Payzin-Dogru D.[1], Sousa J.F.[3], Aggouras A.N.[1],
6 Johnson K.[1], Dunlap G.S.[1], Haas B.J.[2], Levin M.[4,5], Schneider I.[3], Whited J.L.[1,2,4]
7
8 Affiliations:
9
10 [1]Department of Stem Cell and Regenerative Biology, Harvard University, 7 Divinity Ave.
11 Cambridge, MA, 02138, USA
12
13 [2]Broad Institute of MIT and Harvard, 7 Cambridge Center, Cambridge, MA, 02142, USA
14
15 [3]Instituto de Ciências Biológicas, Universidade Federal do Pará, 66075-900 Belém, Brazil
16
17 [4]Allen Discovery Center at Tufts University, Tufts University, Medford, MA, 02155, USA
18
[5]Department of Biology, Tufts University, Medford, MA 02155, USA

Abstract
Regenerative ability varies tremendously across species. A common feature of regeneration of

appendages such as limbs, fins, antlers, and tails is the formation of a blastema--a transient

structure that houses a pool of progenitor cells that regenerate the missing tissue. We have

identified the expression of *von Willebrand Factor D and EGF Domains* (*vwde*) as a common

feature of blastemas capable of regenerating limbs and fins in a variety of highly regenerative

species. Further, *vwde* expression is tightly linked to the ability to regenerate appendages.

Functional experiments demonstrate a requirement for *vwde* in regeneration and indicate that

Vwde is a potent mitogen in the blastema. These data identify a key role for *vwde* in regenerating

blastemas and underscore the power of an evolutionarily-informed approach for identifying

conserved genetic components of regeneration.

Introduction

The underlying reasons why some animals have the ability to regenerate complex structures, while others cannot, remains an important and open question. This knowledge gap has led to intense study of how regeneration-competent species are able to perform complex multi-tissue regeneration, with a particular focus on the ability to regenerate paired appendages, such as limbs and fins. However, this has long been a pursuit without an understanding of whether this ability was present when paired appendages first evolved or was acquired by certain phylogenetic lineages (e.g. urodele amphibians).

Recent work regarding the evolutionary origins of regenerative capacity has indicated that the ability to regenerate paired appendages is an inherited feature of the fin-to-limb transition [1–4]. Evidence found in the fossil record [3,4], functional studies across species [2], and comparisons of gene expression profiles of regenerating tissue [1,2] support the notion that paired appendage regeneration is a feature lost by certain lineages and was not a newly derived capacity in highly regenerative lineages. This indicates that the amniote lineage (which includes humans) has lost regenerative tendencies in appendages over evolutionary time. Therefore, the ability to stimulate regeneration in non-regenerative species, potentially in a therapeutic context, may require the re-initiation of a core, evolutionary conserved program.

All species that are able to regenerate appendages share a conserved trait: the ability to form a blastema. The blastema is the morphological structure that forms at the amputation plane and houses the progenitor cells responsible for regeneration. Recent efforts have focused on elucidating the molecular definition of the blastema, with many of these efforts aimed at the axolotl limb blastema due to the ease of tissue acquisition and the ability to perform experimentation in the lab [5–13]. These studies provide a wealth of information about transcriptomic changes over time, cell types, and blastema-enriched genes. More recently,

sequencing efforts of non-model species have allowed for comparisons to the axolotl limb

blastema and indicate a core molecular signature that is shared between the blastemas of

distantly related species [1,2]. Due to these similarities and the common evolutionary origin of

limb regeneration capacity, we can use an evolutionarily-informed approach to understand what

constitutes a blastema and for identifying core features required for regeneration.

A recent approach to identify the unique gene expression in the axolotl limb blastema

compared blastema gene expression to a variety of homeostatic and embryonic tissues and

identified over 150 blastema-enriched genes [11]. These blastema-enriched genes may help to

explain the unique functions of the blastema, but the question remains as to whether these genes

represent a core program or are functionally required for regeneration. One of the most blastema-

enriched genes in this dataset was *von Willebrand Factor D and EGF Domains* (*vwde*), which to

date has not been functionally studied in any context.

We decided to apply an evolutionary framework to determine if *vwde* fit the description

of an evolutionary-conserved, blastema-enriched gene and if such an approach may help to

identify genes required for regeneration. We found that *vwde* expression is a common feature of

both fin and limb blastemas and was highly enriched in regenerating appendages as compared to

pre-amputation intact appendages. In addition, using the natural regeneration-competent and

regeneration-refractory periods during *Xenopus laevis* development, we observed that *vwde*

expression was tightly linked to the regeneration-component environment. This suggests that

vwde may be a critical factor in the regenerative niche. Finally, we found that *vwde* is

functionally required for axolotl limb regeneration, with transient knockdown of protein levels

resulting in aberrant regeneration. These data suggest that an evolutionarily-informed approach

can help to prioritize target genes and that genes that are blastema-enriched across different

species may prove to be critical factors in the ability to regenerate appendages.

Results

With the goal of identifying genes enriched to the regenerating blastema, a tissue-mapped

axolotl transcriptome was recently published [11]. Of particular interest are blastema-enriched

genes with high expression in the blastema and relative low expression in all other tissues

sampled. We found that *vwde* was highly enriched to the axolotl limb blastema (Figure 1A). This

analysis, however, was limited to one time point, the medium-bud blastema, and it did not

provide spatial information about the expression of *vwde* across the regenerating limb. To

understand the spatial and temporal regulation of *vwde*, we performed RNA *in situ*

hybridizations over a time course of axolotl limb regeneration. We found *vwde* expression to be

tightly tied to the presence of a blastema and expressed exclusively in the blastema and not the

overlying wound epidermis   (Figure 1B-E). Thus, *vwde* fits the description of an axolotl limb

blastema-enriched gene and we were interested in pursuing whether *vwde* may be a core

component of blastemas able to regenerate appendages.

We next sought to determine if *vwde* was present in a selection of deuterostomes,

including species with various regenerative abilities. Using a comparative genomics approach

[14], we found *vwde* to have orthologs across deuterostomes (though no ortholog was detected in

Ciona intestinalis), as well as a non-blastema-enriched paralogous gene in axolotl and other

species (Figure 1F, Supplementary Figure 1). We compared axolotl VWDE to proteins from

other species, and we found putative orthologs harboring predicted von Willebrand Factor D

domains and EGF-like domains. The number of EGF domains may be more variable across

species. However, since this gene has not yet been studied in-depth in any species, additional

experimental work may be required to fully characterize the expressed transcripts and proteins

for individual species. Using these identified orthologs, we moved forward to ask whether *vwde*

was a blastema-enriched gene during paired fin regeneration.

We explored the possibility that *vwde* could be a common feature of blastemas

responsible for regenerating paired fins, which share a deep homology with limbs [15], and

likely share an inherited gene regulatory program for regeneration [1,2]. We chose two highly

regenerative, but distantly related, fish species to determine if *vwde* expression was a conserved

feature of blastemas capable of regenerating paired appendages. These include a species in the

sister group to tetrapods, the Lungfish (*Lepidosiren paradoxa*), which is a lobe-finned fish, and

Polypterus senegalus, a ray-finned fish that is capable of regenerating after amputation through

skeletal elements that develop by endochondral ossification. We first inspected publicly available

transcriptome datasets of lungfish and *Polypterus* regenerating fins for the *vwde* orthologs we

previously identified (Figure 1F). The lungfish LG29893_g1_i1 contig was upregulated in

blastemas 21 days post-amputation (dpa) relative to uninjured fins [1], and the *Polypterus*

PS64836c0_g1_i1 contig was upregulated in 9 dpa blastemas relative to uninjured fins [2].

Assessment of expression levels via qPCR at various regeneration stages showed an upregulation

of *vwde* coinciding with blastema formation during lungfish fin regeneration (Supplemental

Figure 2A).

A similar pattern was seen for *Polypterus* fin regeneration, with expression reaching

highest levels at 5 dpa (Supplemental Figure 2B). Next, we assessed the spatial pattern of *vwde*

in histological sections of regenerating fins. Lungfish 21 dpa blastemas show distal

mesenchymal expression of *vwde* (Figure 2A). In 5 dpa *Polypterus* blastemas, expression is

observed distal to the amputation plane in mesenchymal cells but also in the epithelium,

suggesting that *Polpyterus* may use *vwde* in both compartments (Figure 2C). *In situ* hybridization

with control sense probes did not yield specific signal (Supplemental Figure 2C-D).

Histologically, these samples are similar to the medium-bud blastema time point in which we

identified *vwde* in the axolotl limb (Figure 2B, 2D). Together, these data indicate that *vwde* is

expressed in regenerating fins and limbs and that *vwde* expression is a conserved feature of

blastemas.

To further investigate *vwde* during regeneration, we took advantage of the regeneration-

competent and regeneration-refractory periods during *Xenopus laevis* tail development [16]. A

blastema forms in response to amputation during both distinct developmental stages, but only in

the regeneration-competent setting is full regeneration accomplished. This developmental feature

provides an ideal situation to compare regeneration-component versus regeneration-refractory

environments. We reasoned that finding factors that differentiate these two contexts may provide

clues for identifying the core requirements for successful regeneration. We probed for the

expression of *vwde* during the regeneration-competent and regeneration-refractory periods of

Xenopus laevis tail regeneration. Interestingly, we found that *vwde* expression was present in

tails prior to amputation in both the regeneration-competent and regeneration-refractory setting

(Figure 3A-B, 3G-H). We found robust *vwde* expression along the peripheral edge of the

amputation plane and near the blastema in regeneration-competent tails (Figure 3C-F). In

contrast, in the regeneration-refractory setting, *vwde* expression was restricted to the peripheral

edges of the amputation plane and was not detected near the blastema (Figure 3I-L). This

indicated a striking correlation between *vwde* expression and regeneration, providing evidence

that *vwde* may be an important factor in forming a pro-regenerative niche. These expression data

across a range of species indicate that *vwde* fits the profile of an evolutionarily-conserved,

regeneration-enriched gene and that *vwde* may play an important role in the blastema niche.

To investigate if *vwde* is required for regeneration, we performed morpholino-mediated

knockdown at its peak expression in the medium-bud limb blastema. We found a substantial

reduction in the length of the blastemas when Vwde was knocked down with two separate

translation-blocking morpholinos (Figure 4A-B). Fluorescent reporter constructs with *vwde*-

morpholino binding sites confirmed that both unique *vwde*-targeting morpholinos were capable

of blocking translation (Supplemental Figure 3). Due to the dramatic reduction in blastema

length, we investigated if *vwde* was important for blastema proliferation and/or cell survival. We

found that knockdown of Vwde substantially reduced blastema cell cycle entry (Figure 4C-D)

and did not alter cell survival compared to control limbs (Supplemental Figure 4). Due to the

observed delay in blastema growth, we questioned whether blastemas treated with translation-

blocking morpholinos were capable of recovering from the transient knockdown of Vwde and

produce fully regenerated limbs. We therefore performed the same Vwde morpholino-mediated

knockdown on a separate group of axolotls, and then allowed for the full course of regeneration

to complete, harvesting limbs more than eight weeks post-amputation. We observed that one-

time injection of Vwde-targeting morpholino caused substantial abnormalities in regenerated

limbs, suggesting an essential role for *vwde* during limb regeneration (Figure 4E-G). We found

defects in 4.2% (1/24) control limbs compared to 46% (13/28) of limbs treated with *vwde* MO1

and 25% (5/20) of limbs treated with *vwde* MO2 (Fisher's exact test $P < 0.05$) (Figure 4F, Table

1, Supplemental Figure 5). A second experiment yielded similar results, with defects at endpoint

in 27.6% (8/29) of control (*vwde* MO1 inverted) treated limbs compared to 47% (18/38) of limbs

treated with *vwde* MO1 (Fisher's exact test P < 0.05) (Figure 4G, Table 2, Supplemental Figure

6). We found a variety of defects, some of which are reminiscent of limb development

phenotypes where limited distal elements are present such as has been observed in *fgf4,8*-double-

knockout mice [17] and in the absence of *sonic hedgehog* (*ssh*) [18]. In addition, these

phenotypes also resemble the defective regenerative spike characteristic of *Xenopus* limb

regeneration [19]. Altogether, these data highlight the functional requirement for *vwde* during

limb regeneration.

176
Discussion

Recent work, most notably next generation sequencing, has led to a plethora of

information about the genes and cells that define the blastema [1,2,5–13]. However, it is difficult

to determine which genes may have functional relevance based purely on their expression. We

decided to investigate a single blastema-enriched gene, *vwde*, using an evolutionarily-informed

approach, assuming that a gene whose expression is enriched in blastemas of multiple, distantly-

related, species is likely a key factor during regeneration.

The *in vivo* assays used here place Vwde as a critical regulator of cell cycle entry during

axolotl limb regeneration. Proliferation is a complex, but fundamental, aspect of regeneration, as

there are many different cell types and potential origins of proliferative signals. Previous work

indicates that mitogenic signals are produced directly following amputation independent of the

nerve or wound epidermis [20,21], but are also provided by the nerve [22,23] or wound

epidermis [24]. There are thus multiple sources of mitogenic signals in the regenerating limb, but

it is unclear if mitogenic signals from multiple tissues are required simultaneously or perhaps in

a more stepwise fashion to maintain blastema proliferation. Our data indicate that Vwde may be

a blastema progenitor cell-derived mitogen, which adds to the potential sources of proliferative

signals in the regenerating limb. It has been previously postulated that nerve-derived signals are

required early on during blastema formation and growth, but a fibroblast-derived factor is

required for complete regeneration [25]. We speculate that *vwde*, which appears to be expressed

across the majority of cells in the blastema including likely fibroblasts, may provide one of the

essential fibroblast-derived factors required after the nerve has provided sufficient input. While

there is limited knowledge of fibroblast- or blastema cell-derived mitogens, *in vitro* cultures have

shown that blastema protein extracts are able to drive blastema cell proliferation [26]. More

generally, a global and temporally-based view of the cellular origins of mitogens and the cell

types that require these mitogens will provide a better understanding of what is driving

proliferation during different stages of regeneration.

In addition to the dramatic reduction in proliferation, we observed striking end point

phenotypes after transient knockdown of Vwde. The loss of distal elements and spike-like

phenotypes observed after Vwde knockdown suggest that Vwde plays a role in proximal-distal

determination in the regenerating limb. These phenotypes showing similarities to *ssh* and *fgf4,8*-

double-knockout mice, suggest that Vwde may be working similarly to—or in concert with—

FGFs during regeneration. Though many FGFs are epidermal factors during limb development in

mice and chick, FGFs are expressed in the mesenchyme during axolotl limb development [27]

and regeneration [28]. Thus, it may be that during axolotl limb regeneration, blastema-derived

factors are primarily responsible for proximal-distal patterning and that Vwde is working to

promote the formation of distal elements. Intriguingly, *vwde* has remained unexplored in highly

studied, but less regenerative species such as mouse and human, so whether *vwde* plays a role in

limb development in these species is unknown.

It is interesting to speculate on what has been lost in amniotes that prevents appendage

regeneration. One possibility is genes that are lost in amniotes and present in anamniotes can

explain differences in regenerative capacity [29]. However, the absence of a gene in amniotes is

not necessarily a prerequisite when considering which candidate genes might be responsible for

high regenerative capacity. Alternative scenarios include, but are not limited to, genes that have

lost ancestral pro-regenerative function or have altered expression domains/kinetics. *Vwde* may

fit the paradigm of a gene that is present in both regeneration-competent and regeneration-

incompetent species, but may exclusively be used in the blastema, a structure that cannot be

produced by most regeneration-incompetent species.

While the blastema is required for regeneration, wound healing and activation of

progenitor cells required for formation of the blastema must precede blastema formation. Based

on the expression profile, we do not expect *vwde* to be a driver of blastema formation, but more

likely a downstream effector once a blastema has been established. In most cases of amputation

in less regenerative species, the blastema is not able to form, and thus we suspect that a more

upstream or systemic factor may prevent blastema formation. While there may have initially

been one primary cause of the loss of regenerative ability, such as the rise of adaptive immunity

[30] or trade-offs associated with endothermy [31], it is likely that other aspects of the

regenerative response have now been lost due to their lack of utility. If *vwde* played a relatively

specialized function in the blastema and blastemas generally do not exist in less regenerative

species then the use for *vwde* decreases. This could explain why 42.7% of human genomes have

a predicted loss-of-function copy of *VWDE*, leading to speculation that *VWDE* is potentially

drifting towards inactivation in the human population [32]. While the blastema remains the

elusive feature required for appendage regeneration, this work illustrates that taking an

evolutionarily-informed approach can lead to identification of functionally important genes. This

also suggests that further work to understand the similarities between different species blastemas

may help to elucidate the core molecular program of the blastema.

Methods
242

Animal Experimentation

All axolotl experiments were performed in accordance with Brigham and Women's Hospital

Institutional Animal Care and Use Committee in line with Animal Experimentation Protocol

#04160. All animals were bred in house, but the colony was originally derived from animals

obtained from Ambystoma Genetic Stock Center (Lexington, KY, NIH grant P40-OD019794).

For amputations, animals were narcotized in 0.1% MS-222, confirmed to be fully narcotized by

pinch test, amputated mid-zeugopod and the bone was trimmed. Animals were allowed to

recover overnight in 0.5% sulfamerazine. For all functional experiments, all four limbs were

amputated and injected individually. Functional experiments were performed on animals ranging

from 3.8-8cm.

Polypterus senegalus and *Lepidosiren paradoxa* were maintained in individual tanks in a

recirculating freshwater system. Animals were anesthetized before amputations: *P. senegalus* in

0.1% MS-222 (Sigma) and *L. paradoxa* in 0.1% clove oil diluted in the system water. Experiments

and animal care were performed following animal care guidelines approved by the Animal Care

Committee at the Universidade Federal do Para (protocol no. 037-2015). Pectoral fins in both

species were bilaterally amputated. For *L. paradoxa* fins were amputated at approximately 1 cm

distance from the body, and for *P. senegalus*, fins were amputated across the fin endoskeleton.

Amputated fins (regenerating and uninjured) were used for histology, *in situ* hybridization and

qRT-PCR analysis.

Electroporation

Electroporation was performed while axolotls were narcotized in 0.1% tricaine and subsequently

immersed in ice cold 1x PBS using a NepaGene Super Electroporator NEPA21 Type II

electroporator. Settings for electroporation included: 3 poring pulses at 150 Volts with a pulse

length of 5 milliseconds, a pulse interval of 10 milliseconds, a decay rate of 0 %, and a positive

(+) polarity. Transfer pulse consisted of 5 pulses at 50 Volts with a pulse length of 50 milliseconds,

a pulse interval of 950 milliseconds, a decay rate of 0 %, and a positive (+) polarity.

qRT-PCR.

Total RNA from regenerating or uninjured pectoral fins was extracted using TRIzol reagent

(Thermo Fisher Scientific). Residual DNA removal and RNA cleanup were performed following

the RNeasy Mini Kit (Qiagen) protocol. cDNA was synthesized from 0.5 μg RNA using the

Superscript III First-Strand Synthesis Supermix (Thermo Fisher Scientific) with oligo-dT. For

qPCR, amplification reactions (10 μl) prepared with the GoTaq Probe qPCR Master Mix

(Promega) were run in a StepOnePlus Real-Time PCR System (Applied Biosystems). Gene-

specific oligos (Table 3) for qRT-PCR assays were designed using Primer 3.0

(http://bioinfo.ut.ee/primer3/) and used in a final concentration of 200 nM to each primer. Each

qPCR determination was performed with three biological and three technical replicates. Relative

mRNA expressions were calculated with the $2^{-\Delta\Delta CT}$ method [33], using *sdha* (*P. senegalus*) or

polrc1 (*L. paradoxa*) genes as endogenous control and the uninjured fin (mean ΔCT value of the

three biological replicates) as reference sample.

In Situ Hybridization

For *in situ* hybridization using axolotl samples a gene fragment from the 3' UTR was amplified

from blastema cDNA and cloned into the pGEM-T Easy vector and sequenced. Depending upon

orientation, T7 or Sp6 polymerase was used to transcribe the probe. Primers for *in situ* probes

against axolotl *vwde* (contig c1084387_g3_i1 from [11]) can be found in Table 3. Colorimetric

in situ hybridization in axolotl tissue harvested from animals with snout to tail lengths of 9.5-

11.5cm and was performed as previously described at protocols.io

(https://www.protocols.io/view/rna-in-situ-hybridization-p33dqqn).

For *in situ* hybridizations with fish samples, fins of *P. senegalus* (5 dpa and uninjured) and *L.*

paradoxa (21 dpa and uninjured) were amputated, embedded in TissueTek O.C.T compound

(Fisher Scientific), and maintained at −80°C until use. Frozen sections of 20 μm were obtained on

a Leica CM1850 UV cryostat, positioned on slides (Color Frost Plus/Thermo Fisher Scientific)

and fixed as previously described [1]. Riboprobe templates containing a gene-specific segment

(400-500 bp) and a T7 promoter sequence were produced by a 2-round PCR strategy (primers are

listed in Table 3). Riboprobes were synthesized with T7 RNA polymerase (Roche) and DIG-

labeling mix (Roche). Controls probes (sense riboprobes) were synthesized from a template

containing the T7 promoter in a reverse orientation. A total of 300 ng of DIG-labeled riboprobe

was used per slide during *in situ hybridization* performed as previously described [1]. Images were

obtained on a Nikon Eclipse 80i microscope and processed using the NIS-Element D4.10.1

program.

Whole mount RNA-FISH

Xenopus laevis eggs were obtained, fertilized, and cultured as embryos at 18 °C using standard methods as in [34]. All experimental procedures using *Xenopus laevis* were approved by the Institutional Animal Care and Use Committee (IACUC) and Tufts University Department of Laboratory Animal Medicine (DLAM) under protocol number M2017-53. Once embryos reached regeneration-competent (Stage 40) or regeneration-incompetent (Stage 46) stages, animals were anesthetized using 0.005% MS222 in 0.1X MMR and tails were amputated at the posterior third of the tail and allowed to regenerate for 24 hours. Embryos at both stages, which had not been amputated, were also collected as intact controls. Regenerating and intact control embryos were anesthetized in 0.005% MS222 and then fixed at 4℃, rocking overnight, in either 4% paraformaldehyde in 1X DEPC PBS or MEMPA buffer (0.1 M MOPS (pH 7.4), 2 mM EGTA, 1 mM MgSO4, 3.7% paraformaldehyde). We used a slightly modified whole-mount mouse protocol [35] using hybridization chain reaction v3.0 [36] with slight modifications. After overnight incubation, embryos were washed 3 times for 5 min in PBST and then taken through a methanol series on ice. This series consisted of 10 min washes on ice in ice cold 25%MeOH/75% PBST, 50%MeOH/50% PBST, 75%MeOH/25%PBST, 100%MeOH, and then finally stored in a fresh 100%MeOH solution. Dehydrated embryos were then stored at -20℃ until use. For *in situ*, embryos were subsequently rehydrated via a reverse methanols series, on ice, with 10 min washes of 75% MeOH/25% PBST, 50% MeOH/50% PBST, 25% MeOH/75% PBST, 100% PBST, and another final wash in 100% PBST. Embryos were then digested with proteinase K (10μg/mL) in DEPC PBS for 5 minutes at room temperature. Post-fixation was then performed in 4% PFA in 1X DEPC PBS for 20 minutes at room temperature. Next, three five minutes washes with PBST at room temperature was followed by 5 minutes at 37℃ in hybridization solution (50% formamide, 5x sodium chloride sodium citrate (SSC), 9 mM citric acid (pH 6.0), 0.1% Tween-20, 50μg/mL heparin, 1x Denhardt's solution, and 20% dextran sulfate). Samples were pre-hybridized by full immersion in hybridization solution without probes for 30 min at 37℃. Hybridization was performed overnight at 37℃ with samples immersed in hybridization solution containing twenty probe pairs against *vwde.L* (XM_018267342.1) diluted at 1:200 of 1 μM (hybridization chain reaction v3.0 RNA fluorescent in situ probes were ordered from Molecular Instruments (https://www.molecularinstruments.com/). The following day, samples were washed four times at 37℃ in probe wash buffer (50% formamide, 5X SSC, 9 mM citric acid (pH 6.0), 0.1% Tween-20,

and 50μg/mL heparin). Samples were then washed two times in 5X SSC at room temperature. Pre-

amplification was then performed at room temperature for 30 minutes in amplification buffer (5X

SSC, 0.1% Tween-20, 10% dextran sulfate). During pre-amplification, hairpin probes (ordered

from https://www.molecularinstruments.com/) compatible with *vwde.L* probe pairs were heated

individually at 95℃ for 30 seconds and then snap cooled for 30 min at room temperature in the

dark. After 30 minutes, probe pairs were added to amplification buffer at 1:50 (3 μM stock) and

this probe containing buffer was subsequently added to samples, ensuring that samples were fully

immersed. Incubation was performed overnight at room temperature. The next day, samples were

washed for 5 min in 5X SSCT, twice for 30 min in 5X SSCT, and a 5 min wash in 5X SSCT.

Samples were then stained with DAPI for 5 min in 1X PBS, washed for 5 min in 1X PBS, and then

stored in 1X PBS. Samples were then mounted in low melt agarose and imaged on a Zeiss LSM

880 Upright. A median 3x3 filter followed by maximum projection was applied to all images.

Morpholino design and administration
Morpholinos were designed and synthesized by GeneTools. Morpholino sequences can be found

in Table 3. About 1.25 μl of morpholino was injected in the blastema and electroporation was

performed as described in Electroporation. All morpholinos were 3' fluorescein conjugated to

allow for visualization. Morpholinos were reconstituted to 1 mM in 2X PBS and diluted to a

working concentration of 500 μM in 1X PBS prior to injection.

351
352
EdU staining
Stock solutions of 5-ethynyl-2'-deoxyuridine (EdU) dissolved in dimethyl sulfoxide were

prepared per manufacturer's instructions (Thermo Fisher). Axolotls (3-6cm tail to snout) were

narcotized in 0.1% tricaine at 7 days post amputation and control or Vwde-targeting morpholino

was injected and subsequently electroporated as described in Electroporation section of methods

into the blastema. At 9 dpa, intraperitoneal injections with 400 μM EdU in 0.7X PBS at a

volume of 20μL/g were performed. 18 hours later blastemas were harvested, fixed for 1-2h in

4% PFA and then taken through a sucrose gradient to 30% sucrose in 1x PBS. Tissue was then

embedded in OCT and frozen in a dry ice/ethanol bath. Sections were cut at 16 μm with a

cryostat, collected on Superfrost Plus slides (Fisher), and stored at -80℃. EdU staining was

performed with the Click-iT EdU Alexa Fluor 594 Imaging Kit per manufacturers instructions

(Thermo Fisher).

365
366 **TUNEL assay**

367 TUNEL assays were performed as previously described [11]

368
Skeletal preparations and scoring

Limbs were stained with Alcian blue/Alizarin red according to [37]. In brief, limbs were

incubated with rocking overnight in 95% ethanol and then rocking overnight an acetone. Limbs

were then incubated for at least 7 days in alcian blue/alizarin red at 37 °C. Limbs were then

cleared by incubation in 1% (wt/vol) KOH, followed by 1% (vol/vol) KOH/25% glycerol, 1%

KOH/50% glycerol, and 1% KOH/75% glycerol. Limbs were imaged in 1% KOH/75% glycerol.

Alcian blue stock was 0.3% alcian blue in 70% ethanol; alizarin red stock was 0.1% alizarin red

95% ethanol; the working solution was 5% alcian blue stock/5% alizarin red stock/5% glacial

acetic acid/volume in 70% ethanol.

Definitions for limbs after regeneration. Normal: All digits and carpals present, zeugopod and

stylopod intact. Spike: Single outgrowth from amputation plane without obvious turn at joint. Loss

of distal elements: Distal elements without obvious autopod. Oligodactyly: Loss or reduction in

size at least one digit. Syndactyly: Fusion of digits. Additional elements: Extra bones in stylopod

or zeugopod. For statistical analysis normal was compared to all of the above listed abnormalities.

Ortholog analysis

The following proteomes were downloaded from uniprot.org, human (*Homo sapiens,*

UP000005640, accessed 5/18/2019), zebrafish (*Danio rerio*, UP000000437, accessed

5/18/2019), mouse (*Mus musculus*, accessed 5/18/2019), amphioxus (*Branchiostoma floridae*,

accessed 8/27/2019), chick (*Gallus gallus*, accessed 8/27/2019), sea squirt (*Ciona intestinalis*,

accessed 8/27/2019), lamprey (*Petromyzon marinus*, accessed 8/27/2019), green anole (*Anolis*

carolinensis, 8/27/2019), frog (*Xenopus laevis*, UP000186698, accessed 7/11/2019), frog

(*Xenopus tropicalis*, UP000008143, accessed 7/11/2019). The South American lungfish

transcriptome was downloaded from https://www.ncbi.nlm.nih.gov/Traces/wgs/?val=GEHZ01

and converted to a putative reference protein using TransDecoder (version 5.3.0) like so:

`TransDecoder.LongOrfs -t`. The *Polpyterus* transcriptome can be found here:

https://www.ncbi.nlm.nih.gov/bioproject/480698 and converted with TransDecoder as referenced

above. The axolotl (*Ambystoma mexicanum*) predicted proteome was obtained from

https://data.broadinstitute.org/Trinity/SalamanderWeb/Axolotl.Trinity.CellReports2017.transdec

oder.pep.gz [11]. Cloning of axolotl *vwde* revealed a sequencing error in the axolotl

transcriptome which eliminated the first ~500bp of the sequence. We manually changed the

axolotl proteome to include this corrected version of *vwde* (Supplementary File 1).

To predict orthologs, we used OrthoFinder2.0 (version 2.3.3) [14]. Orthofinder was implemented

as follows:

`orthofinder -f /path/to/proteomes -M msa -A mafft  -T fasttree -t 20 -o /path/to/output/directory`

Protein domain diagrams

The R package, drawProteins [38] was used to draw protein domains for different species Vwde.

For all genes contained within Uniprot, these were downloaded directly with drawProteins. For

genes not available via Uniprot (https://www.uniprot.org/)[39] (e.g. *Polpyterus*, axolotl, and

Lungfish), the amino acid sequence of the protein was queried via Interpro with default settings

(https://www.ebi.ac.uk/interpro/)[40] and positions and domain annotations were extracted and

made into a matrix that matched the required structure for drawProteins. The Uniprot version of

mouse *vwde* (Uniprot ID: Q6DFV8) in the proteome used did not co EGF-like domains, so we

manually searched UCSC genome broswer to confirm this lack of EGF-like domains. This

revealed a full length Vwde (ENSMUST00000203074.2), which was then fed into Interpro and

domains were manually input into drawProteins.

Vwde knockdown confirmation

Two separate constructs to test the target specificity of each MO used. GFP was removed and

from pCAG-GFP (pCAG-GFP was a gift from Connie Cepko (Addgene plasmid # 11150 ;

http://n2t.net/addgene:11150 ; RRID:Addgene_11150)[41] and replaced with vectors containing

td-Tomato sequence and the morpholino binding site (Supplementary File 2). To confirm

knockdown we co-injected and electroporated into medium-bud blastemas the generated

constructs and the appropriate fluorescein-conjugated morpholino. Fluorescein fluorescence was

used to confirm injection efficiency and td-tomato expression was used to measure ability to

block translation.

Statistics

Nested one way ANOVA was used to determine significance between blastema lengths. Each limb was considered a technical replicate within one biological (i.e. animal) replicate. Nested t tests were used to determine significance in EdU and TUNEL experiments, again treating each limb as a technical replicate and placing limbs from the same animal within one biological replicate. Fisher's exact tests (control vs. treated) were used to determine significance of outgrowth phenotypes. Significant results were considered as $P < 0.05$.

432 Contributions

	NDL	SS	ACD	DPD	JFS	ANA	KJ	GSD	BJH	ML	IS	JLW
Wrote the paper	■											■
Initiation/conception	■								■			■
Corresponding author												■
Animal experiments/in situ	■	■	■	■	■	■	■	■		■		■
Orthology analysis	■											
Experimental design	■	■	■	■	■			■			■	■
Manuscript editing	■	■	■	■	■	■	■	■	■	■	■	■
Contributed resources											■	■

Acknowledgements

This work was supported by the Eunice Kennedy Shriver National Institute of Child and Human Development of the National Institutes of Health (5R03HD083434-02 to J.L.W.). N.D.L. was supported by Award Number F32HD092120 from the Eunice Kennedy Shriver National Institute of Child and Human Development of the National Institutes of Health. Portions of this research were conducted on the O2 High Performance Compute Cluster, supported by the Research Computing Group, at Harvard Medical School. See http://rc.hms.harvard.edu for more information. Support for work on lungfish and *Polypterus* was provided by CNPq Universal Program Grant 403248/2016-7 and CAPES/DAAD PROBRAL Grant 88881.198758/2018-01 to I.S, and a postdoctoral fellowship from CNPq to A.C.D. We thank William Ye, Adam Gramy, Sarah Lemire, and Bonney Couper-Kiablick for expert animal care and other members of the Whited lab for feedback and discussions. We also thank Dr. Mansi Srivastava and Dr. Ryan Walker for their insights on orthology and protein structure and Dr. James Monaghan and Timothy Duerr for sharing information on whole mount in situ protocols.

References

1. Nogueira, A.F., Costa, C.M., Lorena, J., Moreira, R.N., Frota-Lima, G.N., Furtado, C.,
Robinson, M., Amemiya, C.T., Darnet, S., and Schneider, I. (2016). Tetrapod limb and
sarcopterygian fin regeneration share a core genetic programme. Nat. Commun. *7*, 13364.

2. Darnet, S., Dragalzew, A.C., Amaral, D.B., Sousa, J.F., Thompson, A.W., Cass, A.N.,
Lorena, J., Pires, E.S., Costa, C.M., Sousa, M.P., *et al.* (2019). Deep evolutionary origin of
limb and fin regeneration. Proc. Natl. Acad. Sci. U. S. A. *116*, 15106–15115.

3. Fröbisch Nadia B., Bickelmann Constanze, and Witzmann Florian (2014). Early evolution
of limb regeneration in tetrapods: evidence from a 300-million-year-old amphibian.
Proceedings of the Royal Society B: Biological Sciences *281*, 20141550.

4. Fröbisch, N.B., Bickelmann, C., Olori, J.C., and Witzmann, F. (2015). Deep-time evolution
of regeneration and preaxial polarity in tetrapod limb development. Nature *527*, 231–234.

5. Voss, S.R., Palumbo, A., Nagarajan, R., Gardiner, D.M., Muneoka, K., Stromberg, A.J., and
Athippozhy, A.T. (2015). Gene expression during the first 28 days of axolotl limb
regeneration I: Experimental design and global analysis of gene expression. Regeneration
(Oxf) *2*, 120–136.

6. Knapp, D., Schulz, H., Rascon, C.A., Volkmer, M., Scholz, J., Nacu, E., Le, M.,
Novozhilov, S., Tazaki, A., Protze, S., *et al.* (2013). Comparative transcriptional profiling
of the axolotl limb identifies a tripartite regeneration-specific gene program. PLoS One *8*,
e61352.

7. Stewart, R., Rascon, C.A., Tian, S., Nie, J., Barry, C., Chu, L.F., Ardalani, H., Wagner, R.J.,
Probasco, M.D., Bolin, J.M., *et al.* (2013). Comparative RNA-seq analysis in the
unsequenced axolotl: the oncogene burst highlights early gene expression in the blastema.
PLoS Comput. Biol. *9*, e1002936.

8. Monaghan, J.R., Athippozhy, A., Seifert, A.W., Putta, S., Stromberg, A.J., Maden, M.,
Gardiner, D.M., and Voss, S.R. (2012). Gene expression patterns specific to the
regenerating limb of the Mexican axolotl. Biol. Open *1*, 937–948.

9. Monaghan, J.R., Epp, L.G., Putta, S., Page, R.B., Walker, J.A., Beachy, C.K., Zhu, W., Pao,
G.M., Verma, I.M., Hunter, T., *et al.* (2009). Microarray and cDNA sequence analysis of
transcription during nerve-dependent limb regeneration. BMC Biol. *7*, 1.

10. Wu, C.H., Tsai, M.H., Ho, C.C., Chen, C.Y., and Lee, H.S. (2013). De novo transcriptome
sequencing of axolotl blastema for identification of differentially expressed genes during
limb regeneration. BMC Genomics *14*, 434.

11. Bryant, D.M., Johnson, K., DiTommaso, T., Tickle, T., Couger, M.B., Payzin-Dogru, D.,
Lee, T.J., Leigh, N.D., Kuo, T.-H., Davis, F.G., *et al.* (2017). A Tissue-Mapped Axolotl De
Novo Transcriptome Enables Identification of Limb Regeneration Factors. Cell Rep. *18*,
762–776.

12. Leigh, N.D., Dunlap, G.S., Johnson, K., Mariano, R., Oshiro, R., Wong, A.Y., Bryant,
D.M., Miller, B.M., Ratner, A., Chen, A., *et al.* (2018). Transcriptomic landscape of the
blastema niche in regenerating adult axolotl limbs at single-cell resolution. Nat. Commun.
9, 5153.

13. Gerber, T., Murawala, P., Knapp, D., Masselink, W., Schuez, M., Hermann, S., Gac-Santel,
M., Nowoshilow, S., Kageyama, J., Khattak, S., *et al.* (2018). Single-cell analysis uncovers
convergence of cell identities during axolotl limb regeneration. Science *362*. Available at:
http://dx.doi.org/10.1126/science.aaq0681.

14. Emms, D.M., and Kelly, S. (2018). OrthoFinder2: fast and accurate phylogenomic
orthology analysis from gene sequences. biorxiv. Available at:
https://www.biorxiv.org/content/biorxiv/early/2018/11/08/466201.full.pdf.

15. Shubin, N., Tabin, C., and Carroll, S. (1997). Fossils, genes and the evolution of animal
limbs. Nature *388*, 639–648.

16. Beck, C.W., Christen, B., and Slack, J.M.W. (2003). Molecular pathways needed for
regeneration of spinal cord and muscle in a vertebrate. Dev. Cell *5*, 429–439.

17. Mariani, F.V., Ahn, C.P., and Martin, G.R. (2008). Genetic evidence that FGFs have an
instructive role in limb proximal-distal patterning. Nature *453*, 401–405.

18. Chiang, C., Litingtung, Y., Lee, E., Youngt, K.E., Cordent, J.L., Westphal, H., and Beachyt,
P.A. Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function.
Available at: https://www-nature-com.ezp-prod1.hul.harvard.edu/articles/383407a0.pdf.

19. Dent, J.N. (1962). Limb Regeneration in Larvae and Metamorphosing Individuals of the
South African Clawed Toad. Journal of Morphology *110*, 61–77.

20. Mescher, A.L., and Tassava, R.A. (1975). Denervation effects on DNA replication and
mitosis during the initiation of limb regeneration in adult newts. Dev. Biol. *44*, 187–197.

21. Johnson, K., Bateman, J., DiTommaso, T., Wong, A.Y., and Whited, J.L. (2017). Systemic
cell cycle activation is induced following complex tissue injury in axolotl. Dev. Biol.
Available at: https://www.ncbi.nlm.nih.gov/pubmed/29111100.

22. Farkas, J.E., Freitas, P.D., Bryant, D.M., Whited, J.L., and Monaghan, J.R. (2016).
Neuregulin-1 signaling is essential for nerve-dependent axolotl limb regeneration.
Development *143*, 2724–2731.

23. Brockes, J.P., and Kintner, C.R. (1986). Glial growth factor and nerve-dependent
proliferation in the regeneration blastema of Urodele amphibians. Cell *45*, 301–306.

24. Sugiura, T., Wang, H., Barsacchi, R., Simon, A., and Tanaka, E.M. (2016). MARCKS-like
protein is an initiating molecule in axolotl appendage regeneration. Nature *531*, 237–240.

25. Endo, T., Bryant, S.V., and Gardiner, D.M. (2004). A stepwise model system for limb

regeneration. Dev. Biol. *270*, 135–145.

26. Boilly, B., and Albert, P. (1990). In vitro control of blastema cell proliferation by extracts
from epidermal cap and mesenchyme of regenerating limbs of axolotls. Rouxs. Arch. Dev.
Biol. *198*, 443–447.

27. Purushothaman, S., Elewa, A., and Seifert, A.W. (2019). Fgf-signaling is
compartmentalized within the mesenchyme and controls proliferation during salamander
limb development. Elife *8*. Available at: http://dx.doi.org/10.7554/eLife.48507.

28. Nacu, E., Gromberg, E., Oliveira, C.R., Drechsel, D., and Tanaka, E.M. (2016). FGF8 and
SHH substitute for anterior-posterior tissue interactions to induce limb regeneration. Nature
533, 407–410.

29. Korotkova, D.D., Lyubetsky, V.A., Ivanova, A.S., Rubanov, L.I., Seliverstov, A.V.,
Zverkov, O.A., Martynova, N.Y., Nesterenko, A.M., Tereshina, M.B., Peshkin, L., *et al.*
(2019). Bioinformatics Screening of Genes Specific for Well-Regenerating Vertebrates
Reveals c-answer, a Regulator of Brain Development and Regeneration. Cell Rep. *29*,
1027–1040.e6.

30. Godwin, J.W., Pinto, A.R., and Rosenthal, N.A. (2017). Chasing the recipe for a pro-
regenerative immune system. Semin. Cell Dev. Biol. *61*, 71–79.

31. Hirose, K., Payumo, A.Y., Cutie, S., Hoang, A., Zhang, H., Guyot, R., Lunn, D., Bigley,
R.B., Yu, H., Wang, J., *et al.* (2019). Evidence for hormonal control of heart regenerative
capacity during endothermy acquisition. Science *364*, 184–188.

32. MacArthur, D.G., Balasubramanian, S., Frankish, A., Huang, N., Morris, J., Walter, K.,
Jostins, L., Habegger, L., Pickrell, J.K., Montgomery, S.B., *et al.* (2012). A systematic
survey of loss-of-function variants in human protein-coding genes. Science *335*, 823–828.

33. Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using
real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods *25*, 402–408.

34. Sive, H.L., Grainger, R.M., and Harland, R.M. (2010). Early Development of Xenopus
Laevis by Hazel L Sive, Robert M Grainger | Waterstones. Available at:
https://www.waterstones.com/book/early-development-of-xenopus-laevis/hazel-l-
sive//9780879699420 [Accessed November 12, 2019].

35. Choi, H.M.T., Calvert, C.R., Husain, N., Huss, D., Barsi, J.C., Deverman, B.E., Hunter,
R.C., Kato, M., Lee, S.M., Abelin, A.C.T., *et al.* (2016). Mapping a multiplexed zoo of
mRNA expression. Development *143*, 3632–3637.

36. Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J.,
Cunha, A., and Pierce, N.A. (2018). Third-generation in situ hybridization chain reaction:
multiplexed, quantitative, sensitive, versatile, robust. Development *145*. Available at:
http://dx.doi.org/10.1242/dev.165753.

37. Whited, J.L., Lehoczky, J.A., and Tabin, C.J. (2012). Inducible genetic system for the
axolotl. Proc. Natl. Acad. Sci. U. S. A. *109*, 13662–13667.

38. Brennan, P. (2018). drawProteins: a Bioconductor/R package for reproducible and
programmatic generation of protein schematics. F1000Res. *7*, 1105.

39. UniProt Consortium (2019). UniProt: a worldwide hub of protein knowledge. Nucleic Acids
Res. *47*, D506–D515.

40. Hunter, S., Apweiler, R., Attwood, T.K., Bairoch, A., Bateman, A., Binns, D., Bork, P.,
Das, U., Daugherty, L., Duquenne, L., *et al.* (2009). InterPro: the integrative protein
signature database. Nucleic Acids Res. *37*, D211–5.

41. Matsuda, T., and Cepko, C.L. (2004). Electroporation and RNA interference in the rodent
retina in vivo and in vitro. Proc. Natl. Acad. Sci. U. S. A. *101*, 16–22.

Figure Legends

Figure 1: *von Willebrand Factor D and EGF-like Domains* (*vwde*) **is a blastema-enriched**
gene that is found across deuterostomes. (A) *vwde* (contig c1084387_g3_i1) expression in
FPKM across tissues sampled from Bryant et al. Cell Reports 2017. Proximal and distal blastema
samples are combined. (B-E) RNA *in situ* hybridization for *vwde* at (B) wound healing, (C)
early-bud blastema, (D) medium-bud blastema, and (E) palette stage regenerating limbs. Black
arrows indicate *vwde* expression, scale bar is 100 μm. (F) OrthoFinder 2.0 phylogeny with
corresponding protein domain structure for putative Vwde orthologs. Protein domain pictures
were generated with drawProteins [38]. Species and Uniprot ID, transcriptome contig number, or
Ensembl ID are included. *Polypterus vwde* contained multiple splice isoforms and the closest
match to axolotl Vwde is shown here. Axolotl Vwde is denoted with (*) and other species Vwde
that are described in this manuscript are marked with (#). Orthologs to the Vwde studied in this
work are indicated with brackets, paralog is also denoted with brackets.

Figure 2: *vwde* **is enriched in the regenerating fin of Lungfish** (*L. paradoxa*) **and** *Polypterus*
(*P. senegalus*)**.** Expression pattern of *vwde* in the fin blastema tissues *of L. paradoxa* and *P.*
senegalus. Longitudinal histological sections of fins from *L. paradoxa* at 21 dpa (A-B), and from
P. senegalus at 5 dpa (C-D). (A and C) *In situ* hybridization using an anti-sense riboprobe to
vwde. (B and D) H&E staining on sequential sections. All panels show posterior view, dorsal to
the top. Dotted lines indicate amputation site (Scale bars, 1 mm in all panels).

Figure 3: *vwde* **expression is tightly linked with the regeneration-component environment.**
In situ hybridization chain reaction probing for *vwde* in (A-F) regeneration-competent *Xenopus*
laevis tails (A-B) prior to amputation, (C-D) blastema 24 hours post-amputation, and (E-F) the
peripheral edge of the amputation plane 24 hours post-amputation. (G-L) Regeneration-
refractory tails (G-H) prior to amputation, (I-J) blastema 24 hours post-amputation, and (K-L)
the peripheral edge of the amputation plane 24 hours post-amputation. White arrows indicate the
location of *vwde* expression. Scale bars are 100 μm.

Figure 4: Vwde is essential for limb regeneration. (A) Representative images of blastemas 16
days post-amputation (9 days post-morpholino administration) from control morpholino
(Standard control MO), *vwde*-targeting morpholino 1 (*vwde* MO1), and *vwde*-targeting
morpholino 2 (*vwde* MO2). Dotted line indicates amputation plane, blastemas are all tissue distal
to amputation plane. Scale bars at 1mm. (B) Quantification of blastema length 16 days-post
amputation. Median and quartiles noted with dotted lines, ** indicates $P< 0.01$, * is $P< 0.05$ by
nested T test. (C) Representative EdU stained sections of blastemas 10 dpa (3 days post-
electroporation) of control morpholino (*vwde* morpholino 1 inverted, *vwde* MO1 INV) and *vwde*-
targeting morpholino 1. Scale bars are 100μm (D) Quantification of percent of blastema cells
positive for EdU in control (*vwde* MO1 INV) and knockdown (*vwde* MO1). Each dot represents
a limb, 4-5 animals per group. Median and quartiles noted with dotted lines,** indicates $P< 0.01$
by nested T test. (E) Representative skeletal preparations of limbs after full regeneration after
knockdown of Vwde at 7 dpa. From left to right, normal forelimb, normal hindlimb, spike, and
loss of distal elements. Scale bars are 5 mm. (F) Donut plots of regenerative outcomes, pooled as
abnormal versus normal from experiment with standard control morpholino, *vwde* MO1 and
vwde MO2. Asterisk (*) indicates $P< 0.05$ by Fisher's exact test comparing control versus *vwde*
MO1 and control versus *vwde* MO2. (G) Donut plots of regenerative outcomes, pooled as
abnormal versus normal from experiment with *vwde* MO1 INV and *vwde* MO1. Asterisk (*)
indicates $P< 0.05$ by Fisher's exact test comparing outcomes from *vwde* MO1 INV compared to
vwde MO1.

618 Table 1. Phenotypic outcomes of morpholino-mediated knockdown in axolotl (Standard control
619 vs. *vwde* MO1, *vwde* MO2)

	Normal	Spike	Loss of distal elements	Oligodactyly	Polydactyly	Syndactyly	Additional elements
Control	23	0	0	0	0	1	0
vwde MO1	15	4	5	4	0	0	0
vwde MO2	11	0	0	3	1	1	0

622 Table 2. Phenotypic outcomes of morpholino-mediated knockdown in axolotl (Inverted control vs.
623 *vwde* MO1)

	Normal	Spike	Loss of distal elements	Oligodactyly	Polydactyly	Syndactyly	Additional elements
Vwde MO1 INV	30	0	4	4	0	0	0
Vwde MO1	20	6	9	1	0	1	1

634   Table 3. Oligos used for qPCR, morpholinos, and ISH probe templates.

Primer	Application	Sequence (5'- 3')
Ax_vwde_ISH_F	Riboprobe template	TGTGGAAAGAAACTTGTGCATCA
Ax_vwde_ISH_R	Riboprobe template	TTTAATCTGAAAATGGACCAGTAGATT
vwde MO1	Anti-sense morpholino	ATATCCCATACATCCTTGCGTTGGC
vwde MO2	Anti-sense morpholino	AGAAACCATCACAGTTCCTCACAGT
vwde MO1 INV	Sense (control) morpholino	CGGTTGCGTTCCTACATACCCTATA
Standard control MO	Control morpholino	CCTCTTACCTCAGTTACAATTTATA
Ps_Vwde-qPCR_F	qPCR	AGAATTCCTGTGACTGTGCGA
Ps_Vwde-qPCR_R	qPCR	TTCTGGTGTTGTTGGTGAGGG
Ps_Vwde-ISH_F	Riboprobe template	GGCCGCGGGCATGCGGAATAATGTGTGCT
Ps_Vwde-ISH_R	Riboprobe template	CCCGGGGCAGTCCAGTCTTCAGCAGTGTG
Lp_Vwde-qPCR_F	qPCR	TTCTTCTTGGAGACCCCTGAT
Lp_Vwde-qPCR_R	qPCR	GGTCTTGCTGGCTAGTGTCAG
Lp_Vwde-ISH_F	Riboprobe template	GGCCGCGGAGCTAACAGCCTGTGCAACAT
Lp_Vwde-ISH_R	Riboprobe template	CCCGGGGCATCAGGGGTCTCCAAGAAGAA
3'_T7 universal	Riboprobe template, 2nd-round PCR	AGGGATCCTAATACGACTCACTATAGGGCC CGGGGC
5'_T7 universal	Riboprobe template, 2nd-round PCR	GAGAATTCTAATACGACTCACTATAGGGCC GCGG

A

vwde

B	C	D	E
Wound healing	Early-bud	Medium-bud	Palette

vwde

F

Amphioxus C3YEN5_BRAFL		⎫
Urchin W4ZF50_STRPU		⎬ ortholog
Lamprey S4R7D9_PETMA		
Lamprey S4RMD6_PETMA		⎭
Axolotl c1047326_g1_i1		⎫
Chick F1NA43_CHICK		
Anolis G1KVZ1_ANOCA		⎬ paralog
Polypterus PS60682_c1_g1_i1		
Zebrafish A0A2R8RMM6_DANRE		⎭
Zebrafish A0A0R4IWA2_DANRE		⎫
Polypterus PS64836_c0_g1_i1	#	
Lungfish c29693_g1_i1	#	
Xenopus tropicalis F7EBV1_XENTR		
Xenopus laevis A0A1L8FVU5_XENLA	#	⎬ ortholog
Axolotl c1084387_g3_i1	*	
Human Q8N2E2 VWDE_HUMAN		
Mouse ENSMUST00000203074.2		
Chick A0A1D5PM68_CHICK		
Anolis G1KNT2_ANOCA		⎭

Domain Type
- EGF-like
- Ig-like
- VWFD

0   1000   2000

Amino acid number

Figure 2

L. paradoxa

P. senegalus

Figure 3

Figure 4

A

| Control MO | *vwde* MO1 | VWDE MO2 |

B

Blastema length AU relative to body length

** *

NS

Standard control | *vwde* MO1 | *vwde* MO2

C

| Control | *vwde* MO1 |

EdU
DAPI

D

% EdU+ of total

**

vwde MO1 INV | *vwde* MO1

E

| Normal forelimb | Normal hindlimb | Spike | Loss of distal elements |

F
Experiment 1

Standard Control | *vwde* MO1 | *vwde* MO2

Normal
Abnormal

Total=24 | Total=28 | Total=16

G
Experiment 2

vwde MO1 INV | vwde MO1

Normal
Abnormal

Total=38 | Total=38