Resist or perish: fate of a microbial population subjected to a periodic presence of antimicrobial

Loïc Marrec[1], Anne-Florence Bitbol[1*]

1 Sorbonne Université, CNRS, Laboratoire Jean Perrin (UMR 8237), F-75005 Paris, France

* anne-florence.bitbol@sorbonne-universite.fr

Abstract

The evolution of antimicrobial resistance can be strongly affected by variations of antimicrobial concentration. Here, we study the impact of periodic alternations of absence and presence of antimicrobial on resistance evolution in a microbial population, using a stochastic model that includes variations of both population composition and size, and fully incorporates stochastic population extinctions. We show that fast alternations of presence and absence of antimicrobial are inefficient to eradicate the microbial population and strongly favor the establishment of resistance, unless the antimicrobial increases enough the death rate. We further demonstrate that if the period of alternations is longer than a threshold value, the microbial population goes extinct upon the first addition of antimicrobial, if it is not rescued by resistance. We express the probability that the population is eradicated upon the first addition of antimicrobial, assuming rare mutations. Rescue by resistance can happen either if resistant mutants preexist, or if they appear after antimicrobial is added to the environment. Importantly, the latter case is fully prevented by perfect biostatic antimicrobials that completely stop division of sensitive microorganisms. By contrast, we show that the parameter regime where treatment is efficient is larger for biocidal drugs than for biostatic drugs. This sheds light on the respective merits of different antimicrobial modes of action.

Introduction

Antibiotics and antivirals allow many major infectious diseases to be treated. However, with the increasing use of antimicrobials, pathogenic microorganisms tend to become resistant to these drugs, which then become useless. Understanding the evolution of resistance is of paramount importance in order to fight the major public health issue raised by antimicrobial resistance [1,2].

The evolution of antimicrobial resistance often occurs in a variable environment, as antimicrobial is added and removed from a medium or given periodically to a patient [3,4]. This results into varying patterns of selection, which are known to have a dramatic effect on evolution in other contexts [5–9]. To address how variations of antimicrobial concentration impact resistance evolution, we investigate theoretically the *de novo* acquisition of resistance in a microbial population in the presence of alternations of phases of presence and absence of antimicrobial. This situation can represent, for example, a treatment where the concentration within the patient falls under the Minimum Inhibitory Concentration (MIC) between drug intakes [10], which is a realistic case [10,11].

We propose a general stochastic model that incorporates variations of both population composition and size, i.e. population genetics and population dynamics. Despite having a

common origin in stochastic birth, death and mutation events, and thus being intrinsically coupled, these phenomena are seldom considered together in theoretical studies [12]. However, it is particularly crucial to address both of them when studying the evolution of antimicrobial resistance, because the aim of an antimicrobial treatment is to eradicate a microbial population, or at least to substantially decrease its size, while the evolution of resistance corresponds to a change in the genetic makeup of the population. Our general model allows us to fully incorporate the stochasticity of mutation occurrence and establishment [13–17], as well as that of population extinction, whose practical importance was recently highlighted [18–20].

In this framework, we ask whether a microbial population subject to alternations of phases of presence and absence of antimicrobial develops resistance, which corresponds to treatment failure and to rescue of the microbial population by resistance [21, 22], or goes extinct, which corresponds to treatment success. In other words, we ask whether the microbial population resists or perishes.

We study both the impact of biocidal drugs, that kill microorganisms, and of biostatic drugs, that prevent microorganisms from growing. We show that fast alternations of phases with and without antimicrobial do not permit eradication of the microbial population before resistant mutants fix, unless the death rate with antimicrobial is large enough. Conversely, intermediate alternation speeds are effective for a wider range of antimicrobial modes of action, but the probability of population extinction and therefore of treatment success, which we fully quantify, is not one, because resistance can rescue the population, and this effect depends on the size of the microbial population. We find that the parameter range where antimicrobial treatment is efficient is larger for biocidal drugs than for biostatic drugs. However, we also show that biocidal and imperfect biostatic antimicrobials permit an additional mechanism of rescue by resistance compared to biostatic drugs that completely stop growth. This sheds light on the respective merits of different antimicrobial modes of action. Finally, we find a population size-dependent critical drug concentration below which antimicrobials cannot eradicate microbial populations.

Model and methods

We consider a microbial population with carrying capacity K, corresponding to the maximum population size that the environment can sustain, given the nutrients available. The division rate of each microorganism is assumed to be logistic, and reads $f(1 - N/K)$, where N represents the total population size, while the fitness f is the maximal division rate of the microorganism, reached when $N \ll K$. This model therefore incorporates population size variations, and allows us to include extinctions induced by the antimicrobial drug.

Mutations that confer antimicrobial resistance are often associated with a fitness cost, i.e. a slower reproduction [23–25], but this fitness cost can be compensated by subsequent mutations [26–29]. The acquisition of resistance is therefore often irreversible, even if the antimicrobial is removed from the environment [24, 26]. Thus motivated, we consider three types of microorganisms: sensitive (S) microorganisms, whose division or death rate is affected by antimicrobials, resistant (R) microorganisms, that are not affected by antimicrobials but that bear a fitness cost, and resistant-compensated (C) microorganisms that are not affected by antimicrobials and do not bear a fitness cost. In the absence of antimicrobial, their fitnesses (maximal division rates) are denoted by f_S, f_R and f_C, respectively, and their death rates by g_S, g_R and g_C. Values in the presence of antimicrobial are denoted by a prime, e.g. f'_S. Note that we include small but nonzero baseline death rates, which can model losses or the impact of the immune system *in vivo*, and allows for population evolution even at steady-state size.

Without loss of generality, we set $f_S = 1$ throughout. In other words, the maximum reproduction rate of S microorganisms, attained when population size is much smaller than the carrying capacity, sets our time unit. We further denote by μ_1 and μ_2 the mutation probabilities upon each division for the mutation from S to R and from R to C, respectively. In several actual cases, the effective mutation rate towards compensation is higher than the one towards the return to sensitivity, because multiple mutations can compensate for the initial cost of resistance [27, 28, 30]. Thus, we do not take into account back-mutations. Still because of the abundance of possible compensatory mutations, often $\mu_1 \ll \mu_2$ [27, 31]. We provide general analytical results as a function of μ_1 and μ_2, and we focus more on the limit $\mu_1 \ll \mu_2$, especially in simulations.

Our model thus incorporates both population dynamics and population genetics [7, 12, 32], and is more realistic than descriptions assuming constant population sizes [33], e.g. in the framework of the Moran process [13, 34]. Throughout, our time unit corresponds to a generation of sensitive microorganisms without antimicrobial in the exponential phase (reached when $N \ll K$).

The action of an antimicrobial drug can be quantified by its MIC, which corresponds the minimum concentration that stops the growth of a microbial population [24]. More precisely, the MIC corresponds to the concentration such that death rate and division rate are equal [18]: in a deterministic framework, above the MIC, the population goes extinct, while below it, it grows until reaching carrying capacity. We investigate the impact of periodic alternations of phases of absence and presence of antimicrobial, at concentrations both above and below the MIC. We consider both biostatic antimicrobials, which decrease the division rate of microorganisms ($f'_S < f_S$), and biocidal antimicrobials, which increase the death rate of microorganisms ($g'_S > g_S$) [18].

We start from a microbial population where all individuals are S (sensitive), without antimicrobial. Specifically, we generally start our simulations with 10 S microorganisms, thus including a phase of initial growth, which can model the development of an infection starting from the bottleneck at transmission [35]. Our results are robust to variations of this initial condition, since we mainly consider timescales longer than that of the initial growth of the population to its equilibrium size. Note however that if we started with a very small number of S microorganisms (i.e. 1 or 2), we would need to take into account rapid stochastic extinctions (see Fig. S3B).

Antimicrobial both drives the decrease of the population of sensitive microorganisms and selects for resistance. We ask whether resistance fully evolves *de novo*, leading to the C microorganisms taking over, or whether the microbial population goes extinct before this happens. The first case corresponds to treatment failure, and the second to treatment success. Hence, we are interested in the probability p_0 of extinction of the microbial population before C microorganisms fix in the population, i.e. take over. We also discuss the average time t_{fix} it takes for the population to fully evolve resistance, up to full fixation of the C microorganisms, and the mean time to extinction before the fixation of the C type t_{ext}.

We present both analytical and numerical results. Our analytical results are obtained using methods from stochastic processes, including the Moran process at fixed population size [13] and birth-death processes with time varying rates [36–39]. Our simulations employ a Gillespie algorithm [40, 41], and incorporate all individual stochastic division, mutation and death events with their exact rates (see Supporting Information, section 5 for details).

Results

Conditions for a periodic presence of perfect biostatic antimicrobial to eradicate the microbial population

Do periodic alternations of phases with and without antimicrobial allow the eradication of a microbial population, or does resistance develop? We first address this question in the case of a biostatic antimicrobial sufficiently above the MIC to completely stop the growth of S microorganisms. With such a "perfect" biostatic antimicrobial, the fitness of S microorganisms is $f'_S = 0$, while without antimicrobial, $f_S = 1$. Here, we assume that the death rate of S microorganisms is not affected by the antimicrobial, i.e. $g'_S = g_S$, but the case of a biocidal antimicrobial will be considered next.

A crucial point is how the duration of a phase with antimicrobial, which corresponds here to the half-period $T/2$ of alternations, compares to the average time τ_S needed for a population of S microorganisms to go extinct in the presence of antimicrobial. Indeed, if $T/2 \gg \tau_S$, one single phase with antimicrobial suffices to eradicate a microbial population in the absence of resistance. An exact first passage time calculation [33,42] (see Supporting Information, section 1.2, Eq. S7) yields $\tau_S = (1/g_S) \times \sum_{i=1}^{N}(1/i) \approx \log(N)/g_S$, where $N \gg 1$ represents the number of microorganisms when antimicrobial is first added, i.e. at $T/2$. If the phase before antimicrobial is added is much longer than the initial growth timescale of the population, i.e. if $T/2 \gg 1/(f_S - g_S)$ (see Supporting Information, section 1.3.1), N can be taken equal to the deterministic equilibrium population size $N = K(1 - g_S/f_S)$, obtained by setting the birth rate $f_S(1 - N/K)$ equal to the death rate g_S. Hence, $\tau_S \approx \log\left[K(1 - g_S/f_S)\right]/g_S$. Our simulation results in Fig. 1A display an abrupt increase in the probability p_0 that the microbial population goes extinct before developing resistance for $T = 2\tau_S$, in good agreement with our analytical prediction.

Fig 1. Periodic presence of a perfect biostatic antimicrobial. A: Probability p_0 that the microbial population goes extinct before resistance gets established versus alternation period T, for various carrying capacities K. Markers: simulation results, with probabilities estimated over $10^2 - 10^3$ realizations. Horizontal solid lines: analytical predictions from Eq. 1. Dashed lines: $T/2 = \tau_S$. **B and C:** Numbers of sensitive (S), resistant (R) and compensated (C) microorganisms versus time in example simulation runs for $K = 1000$, with $T = 20$ and $T = 1000$ respectively. In **B**, resistance takes over, while in **C**, extinction occurs shortly after antimicrobial is first added. Phases without (resp. with) antimicrobial are shaded in white (resp. gray). Parameter values: $f_S = 1$ without antimicrobial, $f'_S = 0$ with antimicrobial, $f_R = 0.9$, $f_C = 1$, $g_S = g_R = g_C = 0.1$, $\mu_1 = 10^{-5}$ and $\mu_2 = 10^{-3}$. All simulations start with 10 S microorganisms.

For fast alternations satisfying $T/2 \ll \tau_S$, the phases with antimicrobial are not long enough to eradicate the microbial population, yielding a systematic evolution of resistance, and thus a vanishing probability p_0 of extinction before resistance takes over. This prediction is confirmed by our simulation results in Fig. 1A, and an example of resistance evolution in this regime is shown in Fig. 1B. In the limit of very fast alternations, we expect an effective averaging of the fitness of S microorganisms, with $\tilde{f}_S = 0.5$. Thus, an R mutant whose lineage will take over the population (i.e. fix) appears after an average time $\tilde{t}_R^a = 1/(\tilde{N}\mu_1 g_S \tilde{p}_{SR})$ where $\tilde{N}\mu_1 g_S$ represents the total mutation rate in the population, with $\tilde{N} = K(1 - g_S/\tilde{f}_S)$, and where $\tilde{p}_{SR} = (1 - \tilde{f}_S/f_R)/[1 - (\tilde{f}_S/f_R)^{\tilde{N}}]$ is the probability that a single R mutant fixes in a population of S microorganisms with constant size $\tilde{N}$, calculated within the Moran model [13]. Subsequently, C mutants will appear and fix, thus leading to the full evolution of resistance in the population. The corresponding average total time t_{fix} of resistance evolution [33] obtained in our simulations agrees well with the analytical expression of $\tilde{t}_R^a$ for $T/2 \ll \tau_S$ (see Fig. S4C).

Conversely, if $T/2 \gg \tau_S$, the microbial population is eradicated by the first phase with antimicrobial, provided that no resistant mutant preexists when antimicrobial is added to the environment. Indeed, resistance cannot appear in the presence of a perfect biostatic antimicrobial since S microorganisms then cannot divide. Thus, in the absence of existing R mutants, extinction occurs shortly after time $T/2$ (see Fig. S4B), and the situation is equivalent to adding antimicrobial at $T/2$ and leaving it thereafter, as exemplified by Fig. 1C. Note that although unlikely, fixation of resistance in the absence of antimicrobial will end up happening by spontaneous fitness valley crossing if the first phase without antimicrobial is long enough. Specifically, this will occur if $T/2 \gg \tau_V$, where $\tau_V \approx (f_S - f_R)/(\mu_1 \mu_2 g_S)$ is the average valley crossing time by tunneling, which is the relevant process unless populations are very small [17, 33, 43, 44]. Accordingly, our simulation results in Fig. S4, which includes longer alternation periods than Fig. 1, feature three distinct regimes, and vanishing extinction probabilities are obtained for $T/2 \gg \tau_V$, as well as for $T/2 \ll \tau_S$.

Let us now focus on the regime where antimicrobial treatment can induce extinction of the microbial population, namely $\tau_S \ll T/2 \ll \tau_V$, and calculate the extinction probability p_0. A necessary condition for the population to be rescued by resistance [21] and avoid extinction is that at least one R mutant be present when antimicrobial is added. In the rare mutation regime $K\mu_1 \ll 1$, this occurs with probability $p_R = t_R^{app}/\tau_R^d = N\mu_1 g_S \tau_R^d$, where $t_R^{app} = 1/(N\mu_1 g_S)$ is the average time of appearance of a resistant mutant, while τ_R^d is the average lifetime of a resistant lineage (destined for extinction without antimicrobial), both calculated in a population of S individuals with fixed size $N = K(1 - g_S/f_S)$ [13, 33]. Importantly, the presence of R mutants does not guarantee the rescue of the microbial population, because small subpopulations of R microorganisms may undergo a rapid stochastic extinction. The probability $p_R^e(i)$ of such an extinction event depends on the number of R microorganisms present when antimicrobial is added, which is i with a probability denoted by $p_R^c(i)$. The probability p_0 that the microbial population is not rescued by resistance and goes extinct can then be expressed as:

$$p_0 = 1 - p_R \sum_{i=1}^{N-1} p_R^c(i)(1 - p_R^e(i)). \tag{1}$$

The probability $p_R^c(i)$ can be calculated within the Moran model since the population size is stable around $N = K(1 - g_S/f_S)$ before antimicrobial is added. Specifically, it can be expressed as the ratio of the average time $\tau_{R,i}^d$ the lineage spends in the state where i mutants exist to the total lifetime τ_R^d of the lineage without antimicrobial: $p_R^c(i) = \tau_{R,i}^d/\tau_R^d$ (see Supporting Information, section 3.1). Next, in order to calculate the probability $p_R^e(i)$ that the lineage of R mutants then quickly goes extinct, we

approximate the reproduction rate of the R microorganisms by $f_R(1-(S(t)+R(t))/K) \approx$
$f_R(1 - S(t)/K)$, where $S(t)$ and $R(t)$ are the numbers of S and R individuals at time
t. Indeed, early extinctions of R mutants tend to happen shortly after the addition of
antimicrobials, when $S(t) \gg R(t)$. Thus motivated, we further take the deterministic
approximation $S(t) = K(1 - g_S/f_S)e^{-g_St}$, while retaining a stochastic description for
the R mutants [36,37]. We then employ the probability generating function $\phi_i(z,t) = \sum_{j=0}^{\infty} z^j P(j,t|i,0)$, where i is the initial number of R microorganisms, which satisfies
$p_R^e(i) = \lim_{t\to\infty} P(0,t|i,0) = \lim_{t\to\infty} \phi_i(0,t)$. Solving the partial differential equation
governing the evolution of $\phi_i(z,t)$ (see Supporting Information, section 3.2) yields [38,39]

$$p_R^e(i) = \lim_{t\to\infty} \left[\frac{g_R \int_0^t e^{\rho(u)}du}{1 + g_R \int_0^t e^{\rho(u)}du} \right]^i , \tag{2}$$

with

$$\rho(t) = \int_0^t \left[g_R - f_R \left(1 - \frac{S(u)}{K} \right) \right] du . \tag{3}$$

Eq. 1 then allows us to predict the probability that the microbial population goes
extinct thanks to the first addition of antimicrobial. Fig. 1A demonstrates a very good
agreement between this analytical prediction and our simulation results in the rare
mutation regime $K\mu_1 \ll 1$, and Fig. S7 further demonstrates good agreement for each
separate term of Eq. 1 in this regime. For larger populations, the probability that the
microbial population is rescued by resistance increases, and the extinction probability
tends to zero for frequent mutations $K\mu_1 \gg 1$ because R mutants are then always
present in the population, in numbers that essentially ensure their survival (see Fig. 1A).

Biocidal antimicrobials and imperfect biostatic ones allow an extra mechanism of rescue by resistance

How does the mode of action of the antimicrobial impact our results? So far, we considered
a perfect biostatic antimicrobial that stops the growth of sensitive microorganisms but
does not affect their death rate. Let us now turn to the general case of an antimicrobial
that can affect both the division rate and the death rate of sensitive microorganisms,
and let us assume that we are above the MIC, i.e. $g_S' > f_S'$. In this section, we present
general calculations, but focus most of our discussion on purely biocidal antimicrobials,
which increase the death rate of sensitive microorganisms without affecting their growth
rate, and compare them to purely biostatic antimicrobials. Again, a crucial point is how
the duration $T/2$ of a phase with antimicrobial compares to the average time τ_S needed
for a population of S microorganisms to go extinct in the presence of antimicrobial (see
Eq. S6). Indeed, our simulation results in Figs. 2A and 2D display an abrupt change in
the probability that the microbial population goes extinct before developing resistance
for $T = 2\tau_S$.

For small periods $T/2 \ll \tau_S$, one phase with antimicrobial is not long enough to
eradicate the microbial population. However, the alternations may induce an overall
decrease in the population over multiple periods, then leading to extinction. This is
the case when the deterministic growth timescale $1/(f_S - g_S)$ is larger than the decay
timescale $1/(g_S' - f_S')$. Equivalently, in the limit of very fast alternations, there is
no nonzero stationary population size when $\tilde{f}_S = (f_S + f_S')/2 < \tilde{g}_S = (g_S + g_S')/2$,
yielding the same condition. For a biostatic drug such that $g_S' = g_S$, this situation
cannot happen if $g_S < f_S/2$, which is realistic since baseline death rates are usually
small. Conversely, for a biocidal drug such that $f_S' = f_S$, a systematic evolution of
resistance will occur if $g_S' < 2f_S - g_S$, while population decay over several periods and
extinction will occur if $g_S' > 2f_S - g_S$. These predictions are confirmed by the simulation

Fig 2. Periodic presence of a biocidal antimicrobial above the MIC. A:
Probability p_0 that the microbial population goes extinct before resistance gets
established versus alternation period T, for various carrying capacities K. Markers:
simulation results, with probabilities estimated over $10^2 - 10^3$ realizations. Horizontal
solid lines: analytical predictions from Eq. 4. Dashed lines: $T/2 = \tau_S$. **B** and **C:**
Numbers of sensitive (S), resistant (R) and compensated (C) microorganisms versus
time in example simulation runs for $K = 1000$, with $T = 8$ and $T = 1000$ respectively.
In **B**, resistance takes over, while in **C**, extinction occurs shortly after antimicrobial is
first added. Phases without (resp. with) antimicrobial are shaded in white (resp. gray).
Parameter values in **A**, **B** and **C**: $f_S = 1$, $f_R = 0.9$, $f_C = 1$, $g_S = 0.1$ without
antimicrobial, $g'_S = 1.1$ with antimicrobial, $g_R = g_C = 0.1$, $\mu_1 = 10^{-5}$ and $\mu_2 = 10^{-3}$.
All simulations start with 10 S microorganisms. **D**, **E** and **F**: same as **A**, **B** and **C**, but
with $g'_S = 2$. All other parameters are the same.

Fig 3. Resistance emergence in the presence of a biocidal antimicrobial above the MIC. A: Numbers of sensitive (S), resistant (R) and compensated (C) microorganisms versus time in an example simulation run for $K = 10^4$, with $T = 1000$. Here resistance takes over. Phases without (resp. with) antimicrobial are shaded in white (resp. gray). **B:** Zoom showing the emergence of resistance in this realization: an R mutant appears after antimicrobial is added (gray). At this time, the S population is decreasing due to the antimicrobial-induced high death rate, but the surviving S microorganisms are still able to divide. Parameter values and initial conditions are the same as in Fig. 2A, B and C.

results in Figs. 2A and D, respectively, and the two different cases are exemplified in Figs. 2B and E. Both of these regimes can arise, depending on the concentration of biocidal antimicrobial. Figs. 2A-C corresponds to concentrations just above the MIC, while Figs. 2D-F correspond to larger concentrations of bactericidal drugs, which can induce death rates equal to several times the birth rate [45, 46]. Note that in Fig. 2A, the extinction probability is not zero for small periods with $K = 10^2$: this is because stochastic extinctions can occur before resistance takes over for such a small equilibrium population size.

For slower alternations satisfying $T/2 \gg \tau_S$, the microbial population is eradicated by the first phase with antimicrobial, unless resistance rescues it. Extinction then occurs shortly after time $T/2$ (see Fig. S5B and examples in Fig. 2C and F). Importantly, with a biocidal antimicrobial or with an imperfect biostatic one, the microbial population can be rescued by resistance in two different ways: either if resistant bacteria are present when antimicrobial is added, or if they appear afterwards. This second case is exemplified in Fig. 3. It can happen because even at high concentration, such antimicrobials do not prevent S microorganisms from dividing, contrarily to a perfect biostatic one. Because of this, rescue by resistance can become more likely than with perfect biostatic antimicrobials. Note that, as in the perfect biostatic case, the spontaneous fixation of resistant mutants without antimicrobial will occur if $T/2 \gg \tau_V \approx (f_S - f_R)/(\mu_1\mu_2 g_S)$ (see Fig. S5).

Let us focus on the regime where the treatment can efficiently induce extinction, namely $\tau_S \ll T/2 \ll \tau_V$. The probability p_0 that the microbial population is not rescued by resistance and goes extinct can then be expressed as:

$$p_0 = \left[1 - p_R \sum_{i=1}^{N-1} p_R^c(i)(1 - p_R^e(i)) \right] \left[1 - p_R^a(1 - p_R^{e'}) \right] . \tag{4}$$

Apart from the last term, which corresponds to resistance appearing after antimicrobial is first added, Eq. 4 is identical to Eq. 1. The quantities p_R and $p_R^c(i)$ are the same

as in that case, since they only depend on what happens just before antimicrobial is added. While $p_R^e(i)$ is conceptually similar to the perfect biostatic case, it depends on f_S' and g_S', and its general calculation is presented in Section 3.2 of the Supporting Information. This leaves us with the new case where resistance appears in the presence of antimicrobial. In the rare mutation regime such that $N_{div}\mu_1 \ll 1$, it happens with probability $p_R^a = N_{div}\mu_1$, where

$$N_{div} = \int_0^{\tau_S} f_S \left(1 - \frac{S(t)}{K}\right) S(t)\, dt \qquad (5)$$

is the number of divisions that would occur in a population of S microorganisms between the addition of antimicrobial (taken as the origin of time here) and extinction. Employing the deterministic approximation for the number $S(t)$ of S microorganisms (see Eq. S21), the probability that the lineage of an R mutant that appears at time t_0 quickly goes extinct can be obtained in a similar way as for Eq. 2, yielding

$$p_R^{e'}(t_0) = \lim_{t \to \infty} \frac{g_R \int_{t_0}^t e^{\eta(u)}\,du}{1 + g_R \int_{t_0}^t e^{\eta(u)}\,du}, \qquad (6)$$

with

$$\eta(t) = \int_{t_0}^t \left[g_R - f_R \left(1 - \frac{S(u)}{K}\right) \right] du \ . \qquad (7)$$

We then estimate the probability $p_R^{e'}$ that the lineage of an R mutant that appears after the addition of antimicrobial quickly goes extinct by averaging $p_R^{e'}(t_0)$ over the time t_0 of appearance of the mutant, under the assumption that exactly one R mutant appears:

$$p_R^{e'} = \int_0^\infty p_R^{e'}(t_0)\ \wp_R^a(t_0)\, dt_0\,, \qquad (8)$$

with

$$\wp_R^a(t_0) = \frac{S(t_0)\left(1 - \frac{S(t_0)}{K}\right)}{\int_0^\infty S(t)\left(1 - \frac{S(t)}{K}\right) dt}\ . \qquad (9)$$

Eq. 4 then yields the probability that the microbial population goes extinct thanks to the first addition of antimicrobial. Fig. 2A demonstrates a very good agreement between this analytical prediction and our simulation results in the rare mutation regime $K\mu_1 \ll 1$, and Figs. S7A-B, S8 and S9 further demonstrate good agreement for each term involved in Eq. 4 in this regime.

The extinction probability p_0 depends on the size of the microbial population through its carrying capacity K and on the division and death rates with antimicrobial. Fig. 4 shows the decrease of p_0 with K, with p_0 reaching 0 for $K\mu_1 \gg 1$ since resistant mutants are then always present when antimicrobial is added. Moreover, Fig. 4 shows that p_0 depends on the antimicrobial mode of action, with large death rates favoring larger p_0 in the biocidal case, and with the perfect biostatic antimicrobial yielding the largest p_0. Qualitatively, the observed increase of p_0 as g_S' increases with a biocidal drug mainly arises from the faster decay of the population of S microorganisms, which reduces the probability p_R^a that an R mutant appears in the presence of antimicrobial. Furthermore, one can show that the extinction probability p_0 is larger for a perfect biostatic antimicrobial than for a perfect biocidal antimicrobial with $g_S' \to \infty$ (see Supporting Information, Section 3.4). Indeed, S microorganisms survive longer in the presence of a perfect biostatic drug, which reduces the division rate of the R mutants due to the logistic growth term, and thus favors their extinction. Such a competition effect is realistic if S microorganisms still take up resources (e.g. nutrients) even while they are not dividing.

Fig 4. Dependence of the extinction probability p_0 on population size and antimicrobial mode of action. The extinction probability p_0 is plotted versus carrying capacity K for the perfect biostatic drug (corresponding to Fig. 1) and two different concentrations of biocidal drugs yielding two different death rates g'_S (corresponding to Fig. 2). Markers correspond to simulation results, computed over 10^3 realizations. Solid lines correspond to our analytical predictions from Eqs. 1 and 4, respectively, which hold for $K \ll 1/\mu_1$. Parameter values and initial conditions are the same as in Figs. 1 and 2, respectively, and the period of alternations is $T = 10^3$, which is in the large-period regime.

Sub-MIC drug concentrations and stochastic extinctions

So far, we considered antimicrobial drugs above the MIC, allowing deterministic extinction in the absence of resistance for long enough drug exposure times. However, sub-MIC drugs can also have a major impact on the evolution of resistance, by selecting for resistance without killing large microbial populations, and moreover by facilitating stochastic extinctions in finite-sized microbial populations [18–20]. In the sub-MIC regime where $f'_S > g'_S$, the population has a nonzero deterministic equilibrium size $N' = K(1 - g'_S/f'_S)$ in the presence of antimicrobial. Nevertheless, stochastic extinctions can remain relatively fast, especially in the weakly-sub-MIC regime where f'_S is close to g'_S, and if K is not very large. The key point is whether resistance appears before the extinction time τ_S. The average time of appearance of an R mutant that fixes in a population of N' individuals in the presence of sub-MIC antimicrobial is $t^a_R = 1/(N'\mu_1 g'_S p'_{SR})$, where $p'_{SR} = [1 - f'_S g_R/(f_R g'_S)]/[1 - (f'_S g_R/(f_R g'_S))^{N'}]$ is the fixation probability of an R mutant in a population of S individuals with fixed size N' (see Supporting Information, Section 4, and Ref. [47]). Therefore, we expect resistance to take over and the extinction probability p_0 to be very small if $t^a_R \ll \tau_S$ below the MIC, even for large periods such that $\tau_S < T/2$.

Fig. 5 shows heatmaps of the probability p_0 that the microbial population goes extinct before resistance takes over, in the cases of biostatic and biocidal drugs, plotted versus the period of alternations T and the non-dimensional variable $\mathcal{R} = (g'_S - f'_S)/g'_S$, which increases with antimicrobial concentration and is zero at the minimum inhibitory concentration (MIC). In both cases, two main regions are apparent, one with $p_0 = 0$ and one where p_0 is close to one. The transition between them is well described by the solid line $T/2 = \tau_S$ such that the time spent with drug is equal to the extinction time τ_S of a population of sensitive microbes with drug, except for large periods, where the relevant transition occurs below the MIC ($\mathcal{R} < 0$) and is given by $t^a_R = \tau_S$ (dashed line), consistently with our analytical predictions.

Fig 5. Heatmaps of the extinction probability. Extinction probability p_0 versus alternation period T and $\mathcal{R} = (g'_S - f'_S)/g'_S$ with biostatic (**A**) or biocidal (**B**) antimicrobial. Heatmap: simulation data, each point computed over 10^3 realizations of simulation results, and linearly interpolated. Dashed white line: value of $\mathcal{R}$ such that $t^a_R = \tau_S$ (see main text). Solid white line: $T/2 = \tau_S$. Parameter values: $K = 10^3$, $\mu_1 = 10^{-5}$, $\mu_2 = 10^{-3}$, $f_S = 1$, $f_R = 0.9$, $f_C = 1$, $g_S = g_R = g_C = 0.1$, and (**A**) $g'_S = 0.1$ and variable f'_S or (**B**) $f'_S = 1$ and variable g'_S. Dotted line in **B**: $\mathcal{R} = (f_S - g_S)/(2f_S - g_S)$. All simulations start with 10 S microorganisms.

The ratio $\mathcal{R}$ enables us to make a quantitative comparison between biostatic and

biocidal drugs. Let us focus first on the transition $\tau_S = t_R^a$. Eq. S6 shows that the average time it takes for the sensitive microorganisms to spontaneously go extinct in the presence of antimicrobial can be written as $\tau_S(f_S', g_S') = \Phi(\mathcal{R})/g_S'$, where Φ is a non-dimensional function. Besides, the average fixation time of a R mutant in a population of S individuals can also be expressed as $t_R^a(f_S', g_S') = \Psi(\mathcal{R})/g_S'$, where Ψ is a non-dimensional function. Thus, the transition $\tau_S = t_R^a$ will be the same for biostatic and biocidal drugs at a given value of $\mathcal{R}$. Conversely, the transition $\tau_S = T/2$, i.e. $\Phi(\mathcal{R})/g_S' = T/2$, depends on g_S', and is thus different for biostatic and biocidal drugs at the same value of $\mathcal{R}$. Specifically, for a given value of $\mathcal{R}$, smaller periods T will suffice to get extinction after the first addition of antimicrobial for a biocidal drug than for a biostatic drug, because g_S' is increased by biocidal drugs, and hence τ_S is smaller in the biocidal case than in the biostatic case. This means that the parameter regime where treatment is efficient is larger for biocidal drugs than for biostatic drugs, as can be seen by comparing Fig. 5A and Fig. 5B. Significantly above the MIC, another difference is that biocidal drugs become efficient even for short periods $T/2 \ll \tau_S$ if their concentration is large enough to have $g_S' > 2f_S - g_S$, i.e. $\mathcal{R} > (f_S - g_S)/(2f_S - g_S)$ (see above, esp. Figs. 2D-E). Numerical simulation results agree well with this prediction (dotted line on Fig. 5B).

Importantly, the transition between large and small extinction probability when $\mathcal{R}$ (and thus the antimicrobial concentration) is varied strongly depends on population size (Figs. 6 and S6), and also depends on antimicrobial mode of action (Fig. 6). For small periods where the relevant transition occurs for $T/2 = \tau_S$, concentrations above the MIC ($\mathcal{R} > 0$) can actually be necessary to get extinction because one period may not suffice to get extinction, and moreover, the extinction threshold value $\mathcal{R}$ is not the same for biostatic and biocidal antimicrobials (see above and Figs. 6A-B). Conversely, for large periods where the relevant transition occurs for $t_R^a = \tau_S$, and extinction occurs upon the first addition of drug, the extinction threshold is always below the MIC ($\mathcal{R} < 0$) and it is the same for biostatic and biocidal antimicrobials (see above and Fig. 6C). In both cases, the larger the population, the larger the concentration required to get large extinction probabilities. For large periods (Fig. 6C), the transition occurs close to the MIC for large populations, but the smaller the population, the larger the discrepancy between the MIC and the actual transition, as predicted by our analytical estimate based on $t_R^a = \tau_S$ (see Fig. S6). This is because in small populations, stochastic extinctions of the population are quite fast at weakly sub-MIC antimicrobial. This is a form of inoculum effect, where the effective MIC depends on the size of the bacterial population [20]. In the large period regime (Fig. 6C), the extinction probability p_0 is well-predicted by Eqs. 1 and 4 for the $\mathcal{R}$ values such at most one R mutant can appear before the extinction of the population (as assumed in our calculation of p_R^a).

In Figs. 6A-B, transitions between small and large values of p_0 in simulated data are observed for smaller threshold values of $\mathcal{R}$ than predicted by $T/2 = \tau_S$ (this can also be seen in Fig. 5, where the solid white line is somewhat in the blue zone corresponding to large p_0). This is because we have employed the average extinction time τ_S, while extinction is a stochastic process. Thus, even if $T/2 < \tau_S$, upon each addition of antimicrobial, there is a nonzero probability that extinction actually occurs within the half-period. Denoting by p the probability that a given extinction time is smaller than $T/2$, the population will on average go extinct after $1/p$ periods, unless resistance fixes earlier. For instance, a population with carrying capacity $K = 10^2$ submitted to alternations with $T = 10^{2.5}$ is predicted to develop resistance before extinction if $\mathcal{R} < 0.055$. However, for $\mathcal{R} = -0.1$, simulations yield a probability $p_0 = 0.99$ of extinction before resistance takes over (see Fig. 6A). In this case, simulations yield $p = 0.3$, implying that extinction typically occurs in ~ 3 periods, thus explaining the large value of p_0. More generally, the probability distribution function of the extinction time can depend on various parameters, which can impact the discrepancy between

Fig 6. Dependence of the extinction transition on population size and antimicrobial mode of action. Extinction probability p_0 versus the ratio $\mathcal{R} = (g'_S - f'_S)/g'_S$ with biostatic or biocidal antimicrobial, for different carrying capacities K, either in the small-period regime, with $T = 10^{2.5}$ (**A** and **B**) or in the large-period regime, with $T = 10^5$ (**C**). Markers: simulation results, calculated over 10^3 realizations. Vertical dashed lines: predicted extinction thresholds, i.e. values of $\mathcal{R}$ such that $T/2 = \tau_S$ (**A** and **B**) or $t_R^a = \tau_S$ (**C**). Solid lines (**C**): Analytical estimates of p_0 from Eq. 1 (biostatic) or Eq. 4 (biocidal). For $K = 10^2$ and 10^3, the analytical predictions in the biostatic and biocidal case are confounded, while for $K = 10^4$ we used two shades of green to show the slight difference (light: biostatic, dark: biocidal). Parameter values: $\mu_1 = 10^{-5}$, $\mu_2 = 10^{-3}$, $f_S = 1$, $f_R = 0.9$, $f_C = 1$, $g_S = g_R = g_C = 0.1$, and $g'_S = 0.1$ (biostatic) or $f'_S = 1$ (biocidal). All simulations start with 10 S microorganisms.

the predicted and observed transitions. A more precise calculation would involve this distribution. Note that the distribution of extinction times is known to be exponential for populations with a quasi-stationary state [48, 49], but the present situation is more complex because there is no nonzero deterministic equilibrium population size below the MIC, and because the population size at the time when antimicrobial is added is far from the equilibrium value with antimicrobial. Nevertheless, our prediction based on the average extinction time τ_S yields the right transition shape (see Fig. 5) and the correct expectations for $T/2 \gg \tau_S$ and $T/2 \ll \tau_S$.

Discussion

The evolution of antimicrobial resistance often occurs in variable environments, as antimicrobial is added and removed from a medium or given periodically to a patient, e.g. in a treatment by the oral route [3, 4]. Alternations of phases of absence and

presence of antimicrobial induce a dramatic time variability of selection pressure on microorganisms, and can thus have a strong impact on resistance evolution. Using a general stochastic model which includes variations of both composition and size of the microbial population, we have shed light on the impact of periodic alternations of presence and absence of antimicrobial on the probability that resistance evolves *de novo* and rescues a microbial population from extinction. The majority of previous studies of periodic antimicrobial treatments [10, 50–55] neglect stochastic effects, while they can have a crucial evolutionary impact [13, 15], especially on population extinction [18, 20]. In addition, established microbial populations are structured, even within a single patient [56], and competition is local, which decreases their effective size, thus making stochasticity relevant. While a few previous studies did take stochasticity into account, some did not include logistic growth or compensation of the cost of resistance [36], while others made specific assumptions on treatments or epidemiology [57, 58], focused on numerical results with few analytical predictions [59], or assumed a constant population size [33]. The present model has the advantage of being quite general while fully accounting for stochasticity and finite-population effects.

We showed that fast alternations of presence and absence of antimicrobial are inefficient to eradicate the microbial population and strongly favor the establishment of resistance, unless the antimicrobial increases enough the death rate, which can occur for biocidal antimicrobials at high concentration [45, 46]. The corresponding criterion on the death rate g'_S of sensitive microorganisms with biocidal antimicrobial, namely $g'_S > 2f_S - g_S$, is generally more stringent than simply requiring drug concentrations to be above the MIC during the phases with biocidal antimicrobial, namely $g'_S > f_S$. Indeed, the population can re-grow without antimicrobial: in this regime, extinction occurs over multiple periods, and involves decaying oscillations. Conversely, for biostatic antimicrobials, as well as for biocidal ones at smaller concentrations, extinction has to occur within a single phase with antimicrobial, and thus the half-period $T/2$ has to be longer than the average extinction time τ_S, which we fully expressed analytically. Importantly, shorter periods suffice for biocidal antimicrobials compared to biostatic ones in order to drive a population to extinction upon the first addition of antimicrobial, at the same value of $\mathcal{R} = (g'_S - f'_S)/g'_S$. Hence, the parameter regime where treatment is efficient is larger for biocidal drugs than for biostatic drugs. If $T/2 > \tau_S$, the microbial population goes extinct upon the first addition of antimicrobial, unless it is rescued by resistance. We obtained an analytical expression for the probability p_0 that the population is eradicated upon the first addition of antimicrobial, assuming rare mutations. Rescue by resistance can happen either if resistant mutants preexist upon the addition of antimicrobial, or if they appear after antimicrobial is added to the environment, during the decay of the population. Importantly, the latter case is fully prevented by perfect biostatic antimicrobials that completely stop division of sensitive microorganisms. This sheds light on the respective merits of different antimicrobial modes of action. Finally, we showed that due to stochastic extinctions, sub-MIC concentrations of antimicrobials can suffice to yield extinction of the population, and we fully quantified this effect and its dependence on population size. Throughout, all of our analytical predictions were tested by numerical simulations, and the latter also allowed us to explore cases beyond the rare mutation regime, where resistance occurs more frequently.

This work opens many possible theoretical extensions. In particular, it will be very interesting to include effects such as antibiotic tolerance, which tend to precede resistance under intermittent antibiotic exposure [4], as well as to consider the possibility of concentrations above the mutant prevention concentration, such that resistant microbes are also affected by the drug [4, 54]. Another exciting extension would be to incorporate spatial structure [60–62] and environment heterogeneity, in particular drug concentration gradients. Indeed, static gradients can strongly accelerate resistance evolution [63–66],

and one may ask how this effect combines with the temporal alternation-driven one investigated here. Besides, it would be interesting to explicitly model horizontal gene transfer of resistance mutations, to include realistic pharmacodynamics and pharmacokinetics [10], and also to compare the impact of periodic alternations to that of random switches of the environment [5–9, 67–69].

Moreover, our results have consequences for actual experimental and clinical situations. First, several of our predictions can be tested experimentally, in particular the transition of extinction probability between the small-period and the large-period regimes, and the predicted values of this extinction probability for large periods. Second, the situation where the phases of absence and presence of antimicrobial have similar durations, which we considered here, is unfortunately clinically realistic. Indeed, a goal in treatment design is that the serum concentration of antimicrobial exceeds the MIC for at least 40 to 50% of the time [11]. Because bacteria divide on a timescale of about an hour in exponential growth phase, and because antimicrobial is often taken every 8 to 12 hours in treatments by the oral route, the alternation period lasts for a few generations in treatments: this is the same order of magnitude as the transition we found between the short-period and long-period regimes, meaning that this transition is relevant in clinical cases. Therefore, it would be very interesting to conduct precise measurements of both division rates and death rates [70] in actual infections in order to determine the relevant regime in each case. This is all the more important that in the short-period regime, we showed that only large concentrations of biocidal antimicrobials are efficient, while other antimicrobials systematically lead to the *de novo* evolution of resistance before eradication of the microbial population. Conversely, a broader spectrum of modes of action can be successful for longer periods of alternation of drug absence and presence. This constitutes a striking argument in favor of the development of extended-release antimicrobial formulations [71].

Despite the fact that only biocidal antimicrobials at high concentration are efficient for short alternation periods of absence and presence of drug, and the fact that the parameter regime where treatment is efficient is larger for biocidal drugs than for biostatic drugs, biostatic antimicrobials that fully stop division of sensitive microorganisms have a distinct advantage over drugs with other modes of action. Indeed, they prevent the emergence of resistant mutants when drug is present, which is all the more important that such resistant mutants are immediately selected for by the antimicrobial and are thus quite likely to rescue the microbial population and to lead to the fixation of resistance. This argues in favor of combination therapies involving a biostatic and a biocidal antimicrobial. Note however that the combined drugs need to be chosen carefully, because some of them have antagonistic interactions [72], depending on their mode of action.

Supporting information

S1 Appendix. Methodological details and further results. In S1 Appendix, we present additional details about our model and methods, as well as additional results.

Acknowledgments

The authors thank Claude Loverdo for inspiring discussions. LM acknowledges funding by a graduate fellowship from EDPIF.

References

1. World Health Organization. Antimicrobial resistance: global report on surveillance; 2014.

2. UK Review on Antimicrobial Resistance, chaired by Jim O'Neill, 2016;.

3. Lin WH, Kussell E. Complex Interplay of Physiology and Selection in the Emergence of Antibiotic Resistance. Curr Biol. 2016;26(11):1486–1493.

4. Levin-Reisman I, Ronin I, Gefen O, Braniss I, Shoresh N, Balaban NQ. Antibiotic tolerance facilitates the evolution of resistance. Science. 2017;355(6327):826–830.

5. Mustonen V, Lässig M. Molecular evolution under fitness fluctuations. Phys Rev Lett. 2008;100(10):108101.

6. Rivoire O, Leibler S. The Value of Information for Populations in Varying Environments. J Stat Phys. 2011;142:1124–1166.

7. Melbinger A, Vergassola M. The Impact of Environmental Fluctuations on Evolutionary Fitness Functions. Sci Rep. 2015;5:15211.

8. Desponds J, Mora T, Walczak AM. Fluctuating fitness shapes the clone-size distribution of immune repertoires. Proc Natl Acad Sci USA. 2016;113(2):274–279.

9. Wienand K, Frey E, Mobilia M. Evolution of a fluctuating population in a randomly switching environment. Phys Rev Lett. 2017;119(15):158301.

10. Regoes RR, Wiuff C, Zappala RM, Garner KN, Baquero F, Levin BR. Pharmacodynamic functions: a multiparameter approach to the design of antibiotic treatment regimens. Antimicrob Agents Chemother. 2004;48(10):3670–3676.

11. Jacobs MR. Optimisation of antimicrobial therapy using pharmacokinetic and pharmacodynamic parameters. Clin Microbiol Infect. 2001;7(11):589–596.

12. Melbinger A, Cremer J, Frey E. Evolutionary game theory in growing populations. Phys Rev Lett. 2010;105(17):178101.

13. Ewens WJ. Mathematical Population Genetics. Springer-Verlag; 1979.

14. Rouzine IM, Rodrigo A, Coffin JM. Transition between stochastic evolution and deterministic evolution in the presence of selection: general theory and application to virology. Microbiol Mol Biol Rev. 2001;65(1):151–185.

15. Fisher DS. Evolutionary Dynamics. In: Bouchaud JP, Mézard M, Dalibard J, editors. Les Houches, Session LXXXV, Complex Systems. Elsevier; 2007.

16. Patwa Z, Wahl LM. The fixation probability of beneficial mutations. J R Soc Interface. 2008;5(28):1279–1289.

17. Weissman DB, Desai MM, Fisher DS, Feldman MW. The rate at which asexual populations cross fitness valleys. Theor Pop Biol. 2009;75:286–300.

18. Coates J, Park BR, Le D, Simsek E, Chaudhry W, Kim M. Antibiotic-induced population fluctuations and stochastic clearance of bacteria. Elife. 2018;7.

19. Teimouri H, Kolomeisky AB. Theoretical investigation of stochastic clearance of bacteria: first-passage analysis. J R Soc Interface. 2019;16(152):20180765.

20. Alexander HK, MacLean RC. Stochastic bacterial population dynamics prevent the emergence of antibiotic resistance. BioRxiv; p. 1–24, `http://dx.doi.org/10.1101/458547`.

21. Martin G, Aguilee R, Ramsayer J, Kaltz O, Ronce O. The probability of evolutionary rescue: towards a quantitative comparison between theory and evolution experiments. Philos Trans R Soc Lond, B, Biol Sci. 2013;368(1610):20120088.

22. Alexander HK, Martin G, Martin OY, Bonhoeffer S. Evolutionary rescue: linking theory for conservation and medicine. Evol Appl. 2014;7(10):1161–1179.

23. Borman AM, Paulous S, Clavel F. Resistance of human immunodeficiency virus type 1 to protease inhibitors: selection of resistance mutations in the presence and absence of the drug. J Gen Virol. 1996;77 (Pt 3):419–426.

24. Andersson DI, Hughes D. Antibiotic resistance and its cost: is it possible to reverse resistance? Nat Rev Microbiol. 2010;8:260–271.

25. zur Wiesch PA, Kouyos R, Engelstadter J, Regoes RR, Bonhoeffer S. Population biological principles of drug-resistance evolution in infectious diseases. Lancet Infect Dis. 2011;11(3):236–247.

26. Schrag SJ, Perrot V, Levin BR. Adaptation to the fitness cost of antibiotic resistance in *E. coli*. Proc R Soc Lond B. 1997;264:1287–1291.

27. Levin BR, Perrot V, Walker N. Compensatory mutations, antibiotic resistance and the population genetics of adaptive evolution in bacteria. Genetics. 2000;154(3):985–997.

28. Paulander W, Maisnier-Patin S, Andersson DI. Multiple mechanisms to ameliorate the fitness burden of mupirocin resistance in Salmonella typhimurium. Mol Microbiol. 2007;64(4):1038–1048.

29. Moura de Sousa J, Sousa A, Bourgard C, Gordo I. Potential for adaptation overrides cost of resistance. Future Microbiol. 2015;10(9):1415–1431.

30. Hughes D, Andersson DI. Evolutionary consequences of drug resistance: shared principles across diverse targets and organisms. Nat Rev Genet. 2015;16(8):459–471.

31. Poon A, Davis BH, Chao L. The coupon collector and the suppressor mutation: estimating the number of compensatory mutations by maximum likelihood. Genetics. 2005;170(3):1323–1332.

32. Huang W, Hauert C, Traulsen A. Stochastic game dynamics under demographic fluctuations. Proc Natl Acad Sci USA. 2015;112(29):9064–9069.

33. Marrec L, Bitbol AF. Quantifying the impact of a periodic presence of antimicrobial on resistance evolution in a homogeneous microbial population of fixed size. J Theor Biol. 2018;457:190–198.

34. Moran PAP. Random processes in genetics. Mathematical Proceedings of the Cambridge Philosophical Society. 1958;54(1):60–71.

35. Abel S, Abel zur Wiesch P, Davis BM, Waldor MK. Analysis of Bottlenecks in Experimental Models of Infection. PLoS Pathog. 2015;11(6):e1004823.

36. Nissen-Meyer S. Analysis of effects of antibiotics on bacteria by means of stochastic models. Biometrics. 1966;22(4):761–780.

37. Bailey NTJ. The Elements of Stochastic Processes with Applications to the Natural Sciences. John Wiley and Sons; 1964.

38. Alexander HK, Bonhoeffer S. Pre-existence and emergence of drug resistance in a generalized model of intra-host viral dynamics. Epidemics. 2012;4:187–202.

39. Parzen E. Stochastic processes. SIAM, Philadelphia; 1999.

40. Gillespie DT. A general method for numerically simulating the stochastic time evolution of coupled chemical reactions. J Comput Phys. 1976;22:403–434.

41. Gillespie DT. Exact stochastic simulation of coupled chemical reactions. J Phys Chem. 1977;81:2340–2361.

42. Sekimoto K. Stochastic Energetics. Springer-Verlag; 2010.

43. Nowak MA, Komarova NL, Sengupta A, Jallepalli PV, Shih IM, Vogelstein B, et al. The role of chromosomal instability in tumor initiation. Proc Natl Acad Sci USA. 2002;99(25):16226–16231.

44. Weinreich DM, Chao L. Rapid evolutionary escape in large populations from local peaks on the Wrightian fitness landscape. Evolution. 2005;59:1175–1182.

45. Levin BR, Udekwu KI. Population dynamics of antibiotic treatment: a mathematical model and hypotheses for time-kill and continuous-culture experiments. Antimicrob Agents Chemother. 2010;54(8):3414–3426.

46. Khan DD, Lagerback P, Cao S, Lustig U, Nielsen EI, Cars O, et al. A mechanism-based pharmacokinetic/pharmacodynamic model allows prediction of antibiotic killing from MIC values for WT and mutants. J Antimicrob Chemother. 2015;70(11):3051–3060.

47. Traulsen A, Hauert C. Stochastic evolutionary game dynamics. In: Schuster HG, editor. Reviews of Nonlinear Dynamics and Complexity. vol. II. Wiley-VCH; 2009.

48. Grimm V, Wissel C. The intrinsic mean time to extinction: a unifying approach to analysing persistence and viability of populations. Oikos. 2004;105(3):501–511.

49. Ovaskainen O, Meerson B. Stochastic models of population extinction. Trends Ecol Evol. 2010;25(11):643 – 652.

50. Lipsitch M, Levin BR. The population dynamics of antimicrobial chemotherapy. Antimicrob Agents Chemother. 1997;41(2):363–373.

51. Wahl LM, Nowak MA. Adherence and drug resistance: predictions for therapy outcome. Proc Biol Sci. 2000;267(1445):835–843.

52. Schulz zur Wiesch P, Engelstadter J, Bonhoeffer S. Compensation of fitness costs and reversibility of antibiotic resistance mutations. Antimicrob Agents Chemother. 2010;54(5):2085–2095.

53. Meredith HR, Lopatkin AJ, Anderson DJ, You L. Bacterial temporal dynamics enable optimal design of antibiotic treatment. PLoS Comput Biol. 2015;11(4):e1004201.

54. Bauer M, Graf IR, Ngampruetikorn V, Stephens GJ, Frey E. Exploiting ecology in drug pulse sequences in favour of population reduction. PLoS Comput Biol. 2017;13(9):e1005747.

55. Hansen E, Woods RJ, Read AF. How to Use a Chemotherapeutic Agent When Resistance to It Threatens the Patient. PLoS Biol. 2017;15(2):e2001110.

56. van Marle G, Gill MJ, Kolodka D, McManus L, Grant T, Church DL. Compartmentalization of the gut viral reservoir in HIV-1 infected patients. Retrovirology. 2007;4:87.

57. Abel Zur Wiesch P, Kouyos R, Abel S, Viechtbauer W, Bonhoeffer S. Cycling empirical antibiotic therapy in hospitals: meta-analysis and models. PLoS Pathog. 2014;10(6):e1004225.

58. Ke R, Loverdo C, Qi H, Sun R, Lloyd-Smith JO. Rational Design and Adaptive Management of Combination Therapies for Hepatitis C Virus Infection. PLoS Comput Biol. 2015;11(6):e1004040.

59. Wu Y, Saddler CA, Valckenborgh F, Tanaka MM. Dynamics of evolutionary rescue in changing environments and the emergence of antibiotic resistance. J Theor Biol. 2014;340:222–231.

60. Bitbol AF, Schwab DJ. Quantifying the role of population subdivision in evolution on rugged fitness landscapes. PLoS Comput Biol. 2014;10(8):e1003778.

61. Nahum JR, Godfrey-Smith P, Harding BN, Marcus JH, Carlson-Stevermer J, Kerr B. A tortoise-hare pattern seen in adapting structured and unstructured populations suggests a rugged fitness landscape in bacteria. Proc Natl Acad Sci USA. 2015;112(24):7530–7535.

62. Cooper JD, Neuhauser C, Dean AM, Kerr B. Tipping the mutation-selection balance: Limited migration increases the frequency of deleterious mutants. J Theor Biol. 2015;380:123–133.

63. Zhang Q, Lambert G, Liao D, Kim H, Robin K, Tung C, et al. Acceleration of emergence of bacterial antibiotic resistance in connected microenvironments. Science. 2011;333(6050):1764–1767.

64. Greulich P, Waclaw B, Allen RJ. Mutational pathway determines whether drug gradients accelerate evolution of drug-resistant cells. Phys Rev Lett. 2012;109:088101.

65. Hermsen R, Deris JB, Hwa T. On the rapidity of antibiotic resistance evolution facilitated by a concentration gradient. Proc Natl Acad Sci USA. 2012;109:10775–10780.

66. Baym M, Lieberman TD, Kelsic ED, Chait R, Gross R, Yelin I, et al. Spatiotemporal microbial evolution on antibiotic landscapes. Science. 2016;353(6304):1147–1151.

67. Hufton PG, Lin YT, Galla T, McKane AJ. Intrinsic noise in systems with switching environments. Phys Rev E. 2016;93(5):052119.

68. Meyer I, Shnerb NM. Noise-induced stabilization and fixation in fluctuating environment. Sci Rep. 2018;8(1):9726.

69. Danino M, Kessler DA, Shnerb NM. Stability of two-species communities: Drift, environmental stochasticity, storage effect and selection. Theor Popul Biol. 2018;119:57 – 71.

70. Frenoy A, Bonhoeffer S. Death and population dynamics affect mutation rate estimates and evolvability under stress in bacteria. PLoS Biol. 2018;16(5):e2005056.

71. Gao P, Nie X, Zou M, Shi Y, Cheng G. Recent advances in materials for extended-release antibiotic delivery system. J Antibiot. 2011;64(9):625–634.

72. Bollenbach T, Quan S, Chait R, Kishony R. Nonoptimal Microbial Response to Antibiotics Underlies Suppressive Drug Interactions. Cell. 2009;139(4):707 – 718.