BIOLOGICAL SCIENCES – Applied Biological Sciences

RNA Sequencing by Direct Tagmentation of RNA/DNA Hybrids

Lin Di,[a,#] Yusi Fu,[a,1,#] Yue Sun,[a] Jie Li,[b] Lu Liu,[b] Jiacheng Yao,[b] Guanbo

Wang,[c,d] Yalei Wu,[e] Kaiqin. Lao,[e] Raymond W. Lee,[e] Genhua Zheng,[e]

Jun Xu,[f] Juntaek Oh,[f] Dong Wang,[f] X. Sunney Xie,[a,d,*] Yanyi Huang,[a,d,g,*]

and Jianbin Wang[b,*]

[a] Beijing Advanced Innovation Center for Genomics (ICG), Biomedical

Pioneering Innovation Center (BIOPIC), School of Life Sciences, and Peking-

Tsinghua Center for Life Sciences, Peking University, Beijing 100871, China.

[b] School of Life Sciences, and Tsinghua-Peking Center for Life Sciences,

Tsinghua University, Beijing 100084, China.

[c] School of Chemistry and Materials Science, Nanjing Normal University,

Nanjing, Jiangsu Province, China

[d] Institute for Cell Analysis, Shenzhen Bay Laboratory, Shenzhen, Guangdong

Province, China

[e] XGen US Co, South San Francisco, CA

[f] Department of Cellular and Molecular Medicine, Skaggs School of Pharmacy

and Pharmaceutical Sciences, University of California, San Diego, La Jolla,

CA 92093

[g] College of Engineering, Peking University, Beijing 100871, China

[#] These authors contributed equally to this work.

[1] Present address: Department of Molecular and Human Genetics, Baylor

College of Medicine, Houston, TX 77030, USA

* Corresponding authors: jianbinwang@tsinghua.edu.cn (J.W.),

yanyi@pku.edu.cn (Y.H.) and sunneyxie@pku.edu.cn (X.S.X.)

Keywords: single cell | RNA-seq | Tn5 transposase

Abstract

Transcriptome profiling by RNA sequencing (RNA-seq) has been widely used to characterize cellular status but it relies on second strand cDNA synthesis to generate initial material for library preparation. Here we use bacterial transposase Tn5, which has been increasingly used in various high-throughput DNA analyses, to construct RNA-seq libraries without second strand synthesis. We show that Tn5 transposome can randomly bind RNA/DNA heteroduplexes and add sequencing adapters onto RNA directly after reverse transcription. This method, Sequencing HEteRo RNA-DNA-hYbrid (SHERRY), is versatile and scalable. SHERRY accepts a wide range of starting materials, from bulk RNA to single cells. SHERRY offers a greatly simplified protocol, and produces results with higher reproducibility and GC uniformity compared with prevailing RNA-seq methods.

Significance Statement

RNA sequencing is widely used to measure gene expression in biomedical research; therefore, improvements in the simplicity and accuracy of the technology are desirable. All existing RNA sequencing methods rely on the conversion of RNA into double-stranded DNA through reverse transcription followed by second strand synthesis. The latter step requires additional enzymes and purification, and introduces sequence-dependent bias. Here, we show that Tn5 transposase, which randomly binds and cuts double-stranded DNA, can directly fragment and prime the RNA/DNA heteroduplexes generated by reverse transcription. The primed fragments are then subject to PCR amplification. This provides a new approach for simple and accurate RNA characterization and quantification.

Introduction

Transcriptome profiling through RNA sequencing (RNA-seq) has become routine in biomedical research since the popularization of next-generation sequencers and the dramatic fall in the cost of sequencing. RNA-seq has been widely used in addressing various biological questions, from exploring the pathogenesis of disease (1, 2) to constructing transcriptome maps for various species (3, 4). RNA-seq provides informative assessments of samples, especially when heterogeneity in a complex biological system (5, 6) or time-dependent dynamic processes are being investigated (7–9). A typical RNA-seq experiment requires a DNA library generated from mRNA transcripts. The commonly used protocols contain a few key steps, including RNA extraction, poly-A selection or ribosomal RNA depletion, reverse transcription, second strand cDNA synthesis, adapter ligation, and PCR amplification (10–12).

Although many experimental protocols, combining novel chemistry and processes, have recently been invented, RNA-seq is still a challenging technology to apply. On one hand, most of these protocols are designed for conventional bulk samples (11, 13–15), which typically contain millions of cells or more. However, many cutting-edge studies require transcriptome analyses of very small amounts of input RNA, for which most large-input protocols do not work. The main reason of this incompatibility is because the purification operations needed between the main experimental steps cause inevitable loss of the nucleic acid molecules.

On the other hand, many single-cell RNA-seq protocols have been invented in the past decade (16–19). However, for most of these protocols it is difficult to achieve both high throughput and high detectability. One type of single-cell RNA-seq approach, such as Smart-seq2 (17), is to introduce pre-amplification to address the low-input problem but such an approach is likely to introduce bias and to impair quantification accuracy. Another type of approach is to barcode each cell's transcripts and hence bioinformatically assign identity to the sequencing data that is linked to each cell and each molecule (20–23). However, the detectability and reproducibility of such approaches are still not ideal (24). An easy and versatile RNA-seq method is needed that works with input from single cells to bulk RNA.

Bacterial transposase Tn5 (25) has been employed in next generation sequencing, taking

advantage of the unique 'tagmentation' function of dimeric Tn5, which can cut double-

stranded DNA (dsDNA) and ligate the resulting DNA ends to specific adaptors.

Genetically engineered Tn5 is now widely used in sequencing library preparation for its

rapid processivity and low sample input requirement (26, 27). For general library

preparation, Tn5 directly reacts with naked dsDNA. This is followed by PCR amplification

with sequencing adaptors. Such a simple one-step tagmentation reaction has greatly

simplified the experimental process, shortening the workflow time and lowering costs.

Tn5-tagmentation has also been used for the detection of chromatin accessibility, high-

accuracy single-cell whole-genome sequencing, and chromatin interaction studies (28–

32). For RNA-seq, the RNA transcripts have to undergo reverse transcription and second

strand synthesis, before the Tn5 tagmentation of the resulting dsDNA (33, 34).

In this paper we present a novel RNA-seq method using Tn5 transposase to directly

tagment RNA/DNA hybrids to form an amplifiable library. We experimentally show that,

as an RNase H superfamily member (35), Tn5 binds to RNA/DNA heteroduplex similarly

as to dsDNA and effectively fragments and then ligates the specific amplification and

sequencing adaptor onto the hybrid. This method, named Sequencing HEteRo RNA-

DNA-hYbrid (SHERRY), greatly improves the robustness of low-input RNA-seq with a

simplified experimental procedure. We also show that SHERRY works with various

amounts of input sample, from single cells to bulk RNA, with a dynamic range spanning

six orders of magnitude. SHERRY shows superior cross-sample robustness and

comparable detectability for both bulk RNA and single cells compared with other

commonly used methods and provides a unique solution for small bulk samples that

existing approaches struggle to handle. Furthermore, this easy-to-operate protocol is

scalable and cost effective, holding promise for use in high-quality and high-throughput

RNA-seq applications.

Results

New RNA-seq strategy using RNA/DNA hybrid tagmentation. Because of its nucleotidyl transfer activity, transposase Tn5 has been widely used in recently developed DNA sequencing technologies. Previous studies (36, 37) have identified a catalytic site within its DDE domain (**Fig. S1A**). Indeed, when we mutated one of the key residues (D188E) (38) in pTXB1 Tn5, its fragmentation activity on dsDNA was notably impaired (**Fig. S1A, B**). Increased amounts of the mutated enzyme showed no improvement in tagmentation, verifying the important role of the DDE domain in Tn5 tagmentation. Tn5 is a member of the ribonuclease H-like (RNHL) superfamily, together with RNase H and MMLV reverse transcriptase (35, 39, 40); therefore, we hypothesized that Tn5 is capable of processing not only dsDNA but also RNA/DNA heteroduplex. Sequence alignment between these three proteins revealed a conserved domain (two Asps and one Glu) within their active sites, termed the RNase H-like domain (**Fig. 1A**). The two Asp residues (D97 and D188) in the Tn5 catalytic core were structurally similar to those of the other two enzymes (**Fig. 1B**). Moreover, divalent ions, which are important for stabilizing substrate and catalyzing reactions, also occupy similar positions in all three proteins (**Fig. 1B**) (39). We determined the nucleic acid substrate binding pocket of Tn5 according to charge distribution. We then docked double-stranded DNA and RNA/DNA heteroduplex into this predicted pocket and showed that the binding site had enough space for an RNA/DNA duplex (**Fig. S1C**). These structural similarities among Tn5, RNase H and MMLV reverse transcriptase and the docking results further supported the possibility that Tn5 can catalyze the strand transfer reaction on RNA/DNA heteroduplex (**Fig. 1C**).

To validate our hypothesis, we purified Tn5 using the pTXB1 plasmid and corresponding protocol. We prepared RNA/DNA hybrids using mRNA extracted from HEK293T cells. Using a typical dsDNA tagmentation protocol, we treated 15 ng of RNA/DNA hybrids with 0.6 µl pTXB1 Tn5 transposome. Fragment analysis of the tagmented RNA/DNA hybrids showed an obvious decrease (~1000bp) in fragment size compared with that of untreated control, validating the capability of Tn5 to fragment the hybrid (**Fig. 1D**).

Fig. 1 Tn5 tagmentation activity on double-stranded hybrids and the experimental process of SHERRY.
(**A**) RNase H-like (RNHL) domain alignment of Tn5 (TN5P_ECOLX), RNase H (RNH_ECOLI) and M-
MLV reverse transcriptase (POL_MLVMS). Active residues in the RNHL domains are labeled in bright
yellow. Orange boxes represent other domains. (**B**) Superposition of the RNHL active sites in these
three enzymes. PDB IDs are 1G15 (RNase H), 2HB5 (M-MLV) and 1MUS (Tn5). (**C**) Putative
mechanism of Tn5 tagmentation of an RNA/DNA hybrid. Crooked arrows represent nucleophilic
attacks. (**D**) Size distribution of mRNA/DNA hybrids with and without Tn5 tagmentation and after
amplification with index primers. (**E**) Workflow of sequencing library preparation by SHERRY. The input
can be a lysed single cell or extracted bulk RNA. After reverse transcription with oligo-dT primer, the
hybrid is directly tagmented by Tn5, followed by gap-repair and enrichment PCR. Wavy and straight
gray lines represent RNA and DNA, respectively. Dotted lines represent the track of extension step.

Based on the ability of the Tn5 transposome to fragment RNA/DNA heteroduplexes, we
propose SHERRY (Sequencing HEteRo RNA-DNA-hYbrid), a rapid RNA-seq library
construction method (**Fig. 1E**). SHERRY consists of three components: RNA reverse
transcription, RNA/cDNA hybrid tagmentation, and PCR amplification. The resulting
product is an indexed library that is ready for sequencing. Specifically, mRNA is reverse
transcribed into RNA/cDNA hybrids using $d(T)_{30}VN$ primer. The hybrid is then tagmented
by the pTXB1 Tn5 transposome, the adding of partial sequencing adaptors to fragment
ends. DNA polymerase then amplifies the cDNA into a sequencing library after initial end
extension. The whole workflow only takes approximately four hours with hand-on time
less than 30 minutes.

To test SHERRY feasibility, we gap-repaired the RNA/DNA tagmentation products
illustrated in **Fig. 1D** (red line) and then amplified these fragments with library construction
primers. Amplified molecules (**Fig. 1D**, blue line) were approximately 100~150 bp longer
than the tagmentation products, which matched the extra length of adaptors added by
gap-repair and index primer amplification. (**Fig. S2**). Thus, direct Tn5 tagmentation of
RNA/DNA hybrids offers a new strategy for RNA-seq library preparation.

Tn5 has ligation activity on tagmented RNA/DNA hybrids. To further investigate the
detailed molecular events of RNA/DNA hybrid tagmentation, we designed a series of
verification experiments. First, we wanted to verify that the transposon adaptor can be
ligated to the end of fragmented RNA (**Fig. 2A**). In brief, we prepared RNA/DNA hybrids
from HEK293T RNA by reverse transcription. After tagmentation with the Tn5
transposome, we purified the products to remove Tn5 proteins and free adaptors. We
assumed that Tn5 ligated the adaptor to the fragmented DNA. At the same time, if Tn5
ligated the adaptor (**Fig. 2A,** dark blue) to the RNA strand, the adaptor could serve as a
template in the subsequent extension step. After extension the DNA strand should have
a primer binding site on both 5′ and 3′ ends for PCR amplification. RNase H treatment
should not affect production of the sequencing library. If Tn5 failed to ligate the adaptor to
the RNA strand, neither strand of the heteroduplex would be converted into a sequencing
library.

After PCR amplification, we obtained a high quantity product regardless of RNase H
digestion, indicating successful ligation of the adaptor to the fragmented DNA.
Sequencing results from both reaction test conditions as well as from SHERRY
showed >90% mapping rate to the human genome with ~80% exon rate and nearly
12,000 genes detected, validating the transcriptome origin of the library (**Fig. 2B, Fig.
S3A**). The additional purification step after reverse transcription and/or RNase H digestion
before PCR amplification did not affect the results, probably because of the large amount
of starting RNA. We examined the sequencing reads with an insert size shorter than 100
bp (shorter than the sequence read length), and 99.7% of them contained adaptor
sequence. Such read-through reads directly proved ligation of the adaptor to the

fragmented RNA (**Fig. S3B and C**). In summary, we confirmed that Tn5 transposome can
tagment both DNA and RNA strands of RNA/DNA heteroduplexes.

206
Fig.2 Verification of Tn5 tagmentation of RNA/DNA heteroduplexes. (**A**) Procedures of two Ligation
Tests. Gray dotted box indicates negative results. The table below lists key experimental parameters
that are different from standard SHERRY. (**B**) Comparison of two Ligation Tests and standard SHERRY
with respect to mapping rate, exon rate and number of genes detected. Each test consisted of two
replicates of 200 ng HEK293T total RNA. (**C**) Strand Test procedures. (**D**) Comparison among three
Strand Tests and dU-SHERRY with respect to mapping rate, exon rate and number of genes detected.

213 Each test consisted of two replicates of 200 ng HEK293T total RNA.
214

Tagmented cDNA is the preferred amplification template. Next, we investigated whether RNA and DNA strands could be amplified to form the sequencing library (**Fig. 2C**). We replaced dTTP with dUTP during the reverse transcription and then purified the tagmented products to remove free dUTP and Tn5 proteins. Bst 2.0 Warmstart DNA polymerase was used for extension because it is able to use RNA as a primer and to process the dU bases in the template. The product fragments were then treated with either USER enzyme or RNase H to digest cDNA and RNA, respectively. We performed RT-PCR with the USER-digested product, to test the efficiency of converting tagmented RNA for library construction (Strand Test 1). To exclude interference from undigested DNA, we performed PCR amplification with the USER-digested fragments using dU-compatible polymerase (Strand Test 2). We also used dU-compatible PCR to test the efficiency of converting tagmented cDNA for library construction (Strand Test 3). For comparison, we included a control experiment with the same workflow as Strand Test 3 except that the RNase H digestion step was omitted (dU-SHERRY) to ensure that Tn5 can recognize substrates with dUTP.

Sequencing results of Strand Test 1 showed a low mapping rate and gene detection count that were only slightly higher than those of Strand Test 2. In contrast, Strand Test 3 demonstrated a similar exon rate and gene count to dU-SHERRY and SHERRY (**Fig. 2D, Fig. S3A**). Based on these results, we conclude that the tagmented cDNA contributes to the majority of the final sequencing library, likely because of inevitable RNA degradation during the series of reactions.

SHERRY for rapid one-step RNA-seq library preparation. We tested different reaction conditions to optimize SHERRY with 10 ng total RNA as input (**Fig. S4A**). We evaluated the impact of different crowding agents, different ribonucleotide modifications on transposon adaptors, and different enzymes for gap filling. We also included purification after certain steps to remove primer dimers and carry-over contaminations. Sequencing results showed little change in performance from most of these modifications, indicating that SHERRY is robust under various conditions.

We then compared the optimized SHERRY with NEBNext® Ultra™ II, a commercially

available kit, for bulk RNA library preparation. This NEBNext kit is one of the most

commonly used kits used for RNA-seq experiments, with 10 ng total RNA being its

minimum input limit. We therefore tested the RNA-seq performance with 10 ng and 200

ng total RNA inputs, each condition having three replicates. SHERRY demonstrated

comparable performances with NEBNext for both input levels (**Fig. S4B**). For the 10 ng

input tests, SHERRY produced more precise gene expression measurements across

replicates (**Fig. 3A**), probably because of the simpler SHERRY workflow.

Fig.3 Performance of SHERRY with large RNA input. (**A**) Coefficient of variation (CV) across three replicates was plotted against the mean value of each gene's FPKM (Fragments Per Kilobase of transcript per Million mapped reads). All experiments used HEK293T total RNA as input. (**B**) Genes detected by SHERRY in three replicates of 200 ng HEK293T or HeLa total RNA are plotted in Venn Diagrams. Numbers of common genes are indicated. (**C**) Common genes detected by SHERRY and NEBNext in the three replicates of 200 ng HEK293T or HeLa total RNA. (**D**) Distance heatmap of samples prepared by SHERRY or NEBNext for three replicates using 200 ng HEK293T or HeLa total RNA. The color bar indicates the Euclidian distance. (**E**) Correlation of gene expression fold-change identified by SHERRY and NEBNext. Involved genes are differentially expressed genes between HEK293T and HeLa detected by both methods.

Next, we compared the ability to detect differentially expressed genes between HEK293T

and HeLa cells using SHERRY and NEBNext. In all three replicates, SHERRY detected 11,269 genes in HEK293T cells and 10,774 genes in HeLa cells, with high precision (correlation coefficient R^2=0.999) (**Fig. 3B, Fig. S5A**). The numbers of detected genes and their read counts identified by SHERRY and NEBNext were highly concordant (**Fig. 3C, Fig.S5B**). This excellent reproducibility of SHERRY ensured the reliability of subsequent analyses. Then we plotted a heatmap of the distance matrix (**Fig. 3D**) between different cell types and library preparation methods. Libraries from the same cell type were clustered together as expected. Libraries from the same method also tended to cluster together, indicating internal bias in both methods.

We then used DESeq2 to detect differentially expressed genes (P-value <5×10^{-6}, |$\log_2$Fold change| >1). In general, the thousands of differentially expressed genes detected by both methods were highly similar (**Fig. S5C**) and their expression fold-change was highly correlated (correlation coefficient R^2=0.977) between SHERRY and NEBNext (**Fig. 3E**). Examination of the genes that showed differential expression in only one method, revealed the same trend of expression change in the data from the other method (**Fig. S5D and E**). We conclude that SHERRY provides equally reliable differential gene expression information as NEBNext, but with a much faster and less labor-intensive process, specifically saving around two hours hand-on time (**Fig. S7A**).

SHERRY using trace amounts of RNA or single cells. We next investigated whether SHERRY could construct RNA-seq libraries from single cells. First, we reduced the input to 100 pg total RNA, which is equivalent to RNA from about 10 cells. SHERRY results were high quality, with high mapping and exon rates and nearly 9,000 genes detected (**Fig. S6A**). 72% of these genes were detected in all three replicates, demonstrating good reproducibility (**Fig. 4A**). The expression of these genes showed excellent precision with R^2 ranging from 0.958 to 0.970. (**Fig. 4B**).

To further push the detection limit, we carried out single-cell SHERRY experiments (scSHERRY) using HEK293T cell line. In contrast to the experiments with purified RNA, scSHERRY required several optimizations to the standard protocol (**Fig. 4C, Fig. S6B**).

Although we found no positive effect by replacing betaine in the standard protocol with
other crowding agents during optimization, we found that addition of a crowding reagent
with a higher molecular weight improved the library quality from single cells. Therefore,
we used PEG8000 for the following scSHERRY experiments. For the extension step, the
use of Bst 3.0 or Bst 2.0 WarmStart DNA polymerases detected more genes than the use
of Superscript II or Superscript III reverse transcriptases. This is probably because of the
stronger processivity and strand displacement activity of Bst polymerases, and better
compatibility with higher reaction temperatures to open the secondary structure of RNA
templates. We also tried to optimize the PCR strategy because extensive amplification
can lead to strong bias. Compared to the continuous 28-cycle PCR, the incorporation of
a purification step after 10 cycles, or simply reducing the total cycle number to 15
increased the mapping rate and the number of genes detected. Therefore, we performed
15-cycle PCR without extra purification to better accommodate high-throughput
experiments.

Fig.4 Performance of SHERRY with micro-input samples. (**A**) Genes detected by SHERRY in three replicates with 100 pg HEK293T total RNA. (**B**) Correlations of normalized gene read counts between replicates with 100 pg HEK293T total RNA. (**C**) Gene number detected by scSHERRY under various experimental conditions in single HEK293T cells. Each condition involved 3–4 replicates. (**D**) The heatmap of R^2 calculated from scSHERRY and Smart-seq2 replicates, and slope deviation in a linear fitting equation for the two methods. (**E**) Normalized gene numbers with different GC content.

The optimized scSHERRY was capable of detecting 8,338 genes with a 50.17% mapping rate (**Fig. S6D**), and the gene read counts correlated well (correlation coefficient $R^2=0.600$) with Smart-seq2, the most prevalent protocol in the single-cell RNA amplification field. Besides, scSHERRY showed better reproducibility than Smart-seq2 (**Fig.4D**). Compared with Smart-seq2, the gene number and coverage uniformity of the scSHERRY-generated library was slightly inferior (**Fig. S6D, E**) because Smart-seq2 enriches for full-length transcripts via a preamplification step. However, this enrichment step of Smart-seq2 also introduced bias (**Fig. 4E**). We used 200 ng of HEK293T total RNA to construct a sequencing library using the NEBNext kit, expecting to capture as many genes as possible. Besides, the NEBNext protocol used short RNA fragments for reverse transcription and fewer cycles of PCR amplification, which should introduce less GC bias than the other protocols. We then compared the GC distribution of genes detected by scSHERRY or Smart-seq2 with NEBNext results. scSHERRY, which is free from second-strand synthesis and preamplification, produced a distribution similar to the standard. However, the library from Smart-seq2-amplified scRNA showed clear enrichment for genes with lower GC content. Genes with high GC content were less likely to be captured by Smart-seq2, which may cause biased quantitation results. Overall, compared with Smart-seq2, scSHERRY produced libraries of comparable quality and lower GC-bias. Moreover, the scSHERRY workflow spares pre-amplification and QC steps before tagmentation, saving around four hours (**Fig. S7A**), and the one tube strategy is promising for high throughput application.

Discussion

We found that the Tn5 transposome has the capability to directly fragment and tag RNA/DNA heteroduplexes and, therefore, we have developed a quick RNA amplification and library preparation method called SHERRY. The input for SHERRY could be RNA from single cell lysate or total RNA extracted from a large number of cells. Comparison of

SHERRY with the commonly used Smart-seq2 protocol for single-cell input or the
NEBNext kit for bulk total RNA input, showed comparable performance for input amount
spanning more than five orders of magnitude. Furthermore, the whole SHERRY workflow
from RNA to sequencing library consists of only five steps in one tube and takes about
four hours, with hands-on time of less than 30 min. Smart-seq2, requires twice this
amount of time and an additional library preparation step is necessary. The ten-step
NEBNext protocol is much more labor-intensive and time-consuming (**Fig. S7A**).
Moreover, the SHERRY reagent cost is five-fold less compared with that of the other two
methods (**Fig. S7B**). For single cells, the lower mapping rate of SHERRY compared to
Smart-seq2 could increase sequencing cost. However, both methods reached a plateau
of saturation curve with 2 million total reads (**Fig. S8**), which costs less than $5. Therefore,
SHERRY has strong competitive advantages over conventional RNA library preparation
methods and scRNA amplification methods.

In our previous experiments, we assembled a Tn5 transposome using home-purified
pTXB1 Tn5 and synthesized sequencer-adapted oligos. To generalize the SHERRY
method, and to confirm Tn5 tagmentation of RNA/DNA heteroduplexes, we tested two
commercially available Tn5 transposomes, Amplicon Tagment Mix (abbreviated as ATM)
from the Nextera XT kit (Illumina, USA) and TTE Mix V50 (abbreviated as V50) from the
TruPrep kit (Vazyme, China). We normalized the different Tn5 transposome sources
according to the enzyme processing 5 ng of genomic DNA to the same size under the
same reaction conditions (**Fig. S9A**). The tagmentation activity of our in-house pTXB1
Tn5 was 10-fold higher than V50 and 500-fold higher than ATM when using transposome
volume as the metric. The same units of enzyme were then used to process RNA/DNA
heteroduplexes prepared from 5 ng mRNA to confirm similar performance on such hybrids
(**Fig. S9B**). The RNA-seq libraries from all three enzymes showed consistent results,
demonstrating the robustness of SHERRY (**Fig. S9C**).

During DNA and RNA/DNA heteroduplex tagmentation, the Tn5 transposome reacted
with these two substrates in different patterns. We tagmented 5 ng DNA or mRNA/DNA
hybrids with 0.02 µl, 0.05 µl or 0.2 µl pTXB1 Tn5 transposome (**Fig. S10**). As the amount

of Tn5 increased, dsDNA was cut into overall shorter fragments. While for the hybrid, Tn5

cut the template 'one by one', because only hybrids above a certain size became shorter

and most were too short to be cut. We supposed that such phenomenon might attribute

to the different conformation of dsDNA and RNA/DNA hybrid, since diameter of the latter

is larger. Thus, the binding pocket or catalytic site of Tn5 would be tuned to accommodate

the hybrid strands and cause different tagmentation pattern.

Despite its ease-of-use and commercial promise, the library quality produced by SHERRY

may be limited by unevenness of transcript coverage. Unlike the NEBNext kit, which

fragments RNA before reverse transcription, or Smart-seq2, which performs pre-

amplification to enrich full-length cDNAs, SHERRY simply reverse transcribes full-length

RNA. Reverse transcriptase is well known for its low efficiency and when using polyT as

the primer for extension, it is difficult for the transcriptase to reach the 5′ end of the RNA

template. This can cause coverage imbalance across transcripts, making the RNA-seq

signal biased toward the 3′ end of genes (**Fig. S11A**). In an attempt to solve this problem,

we added TSO primer, the sequence and concentration of which was the same as Smart-

seq2 protocol (17), to the reverse transcription buffer to mimic the Smart-seq2 reverse

transcription conditions. The resulting hybrid was then tagmented and amplified following

standard SHERRY workflow. This produced much improved evenness across transcripts

(**Fig. S11A and B**), although some of the sequencing parameters dropped accordingly

(**Fig. S11C**). We believe that with continued optimization SHERRY will improve RNA-seq

performance.

Author contributions

Y.F., K.L., X.S.X., Y.H. and J.W. conceived the project; L.D., J.X., J.O. and D.W. performed

structural analysis; Y.S., J.L. and G.W. performed Tn5 purification and characterization;

L.D., Y.F. and L.L. conducted research; L.D., Y.F., L.L., J.Y., Y.W., R.L., G.Z., Y.H. and J.W.

analyzed the data; L.D., X.S.X., Y.H. and J.W. wrote the manuscript with the help from all

other authors.

Conflict of interest statement

XGen US Co has applied for a patent related to this work. X. Sunney Xie, Kaiqin. Lao and Yalei Wu are shareholders of XGen US Co. Kai Q. Lao, Yalei Wu, Raymond W. Lee and Genhua Zheng are employees of XGen US Co.

Acknowledgement

We thank Dr. Yun Zhang and BIOPIC sequencing platform at Peking University for the assistance of high-throughput sequencing experiments. This work was supported by National Natural Science Foundation of China (21675098, 21525521), Ministry of Science and Technology of China (2018YFA0800200, 2018YFA0108100, 2018YFC1002300), 2018 Beijing Brain Initiation (Z181100001518004), Beijing Advanced Innovation Center for Structural Biology, and Beijing Advanced Innovation Center for Genomics.

References
1. Y. Liu, Y. L. Lightfoot, N. Seto, C. Carmona-Rivera, E. Moore, R. Goel, L. O'Neil, P. Mistry, V. Hoffmann, S. Mondal, P. N. Premnath, K. Gribbons, S. Dell'Orso, K. Jiang, P. R. Thompson, H. W. Sun, S. A. Coonrod, M. J. Kaplan, Peptidylarginine deiminases 2 and 4 modulate innate and adaptive immune responses in TLR-7-dependent lupus. JCI Insight **3**, e124729 (2018)

2. E. Schmidt, I. Dhaouadi, I. Gaziano, M. Oliverio, P. Klemm, M. Awazawa, G. Mitterer, E. Fernandez-Rebollo, M. Pradas-Juni, W. Wagner, P. Hammerschmidt, R. Loureiro, C. Kiefer, N. R. Hansmeier, S. Khani, M. Bergami, M. Heine, E. Ntini, P. Frommolt, P. Zentis, U. A. Ørom, J. Heeren, M. Blüher, M. Bilban, and J. W. Kornfeld, LincRNA H19 protects from dietary obesity by constraining expression of monoallelic genes in brown fat. Nat. Commun. **9**, 3622 (2018).

3. O. Wurtzel, R. Sapra, F. Chen, Y. Zhu, B. A. Simmons, R. Sorek, A single-base resolution map of an archaeal transcriptome. Genome Res. **20**, 133-141 (2010).

4. A. A. Penin, A. V. Klepikova, A. S. Kasianov, E. S. Gerasimov, M. D. Logacheva, Comparative Analysis of Developmental Transcriptome Maps of Arabidopsis thaliana and Solanum lycopersicum. Genes **10**, 50 (2019).

5. P. Civita, S. Franceschi, P. Aretini, V. Ortenzi, M. Menicagli, F. Lessi, F. Pasqualetti, A. G. Naccarato, C. M. Mazzanti, Laser Capture Microdissection and RNA-Seq Analysis:

High Sensitivity Approaches to Explain Histopathological Heterogeneity in Human Glioblastoma FFPE Archived Tissues. Front Oncol. **9**, 482 (2019).

6. D. A. Jaitin, E. Kenigsberg, H. Keren-Shaul, N. Elefant, F. Paul, I. Zaretsky, A. Mildner, N. Cohen, S. Jung, A. Tanay, I. Amit, Massively Parallel Single-Cell RNA-Seq for Marker-Free Decomposition of Tissues into Cell Types. Science **343**, 776-779 (2014)

7. Y. Chen, Y. Zheng, Y. Gao, Z. Lin, S. Yang, T. Wang, Q. Wang, N. Xie, R. Hua, M. Liu, J. Sha, M. D. Griswold, J. Li, F. Tang, M. H. Tong, Single-cell RNA-seq uncovers dynamic processes and critical regulators in mouse spermatogenesis. Cell Res. **28**, 879-896 (2018).

8. M. Wang, X. Liu, G. Chang, Y. Chen, G. An, L. Yan, S. Gao, Y. Xu, Y. Cui, J. Dong, Y. Chen, X. Fan, Y. Hu, K. Song, X. Zhu, Y. Gao, Z. Yao, S. Bian, Y. Hou, J. Lu, R. Wang, Y. Fan, Y. Lian, W. Tang, Y. Wang, J. Liu, L. Zhao, L. Wang, Z. Liu, R. Yuan, Y. Shi, B. Hu, X. Ren, F. Tang, X. Y. Zhao, J. Qiao, Single-Cell RNA Sequencing Analysis Reveals Sequential Cell Fate Transition during Human Spermatogenesis. Cell Stem Cell **23**, 599-614 (2018).

9. H. Hochgerner, A. Zeisel, P. Lönnerberg, S. Linnarsson, Conserved properties of dentate gyrus neurogenesis across postnatal development revealed by single-cell RNA sequencing. Nat. Neurosci. **21**, 290-299 (2018).

10. U. Gubler, B. J. Hoffman, A simple and very efficient method for generating cDNA libraries. Gene **25**, 263-269 (1983).

11. A. Mortazavi, B. A. Williams, K. McCue, L. Schaeffer, B. Wold, Mapping and quantifying mammalian transcriptomes by RNA-Seq. Nat. Methods **5**, 621-628 (2008).

12. P. Cui, Q. Lin, F. Ding, C. Xin, W. Gong, L. Zhang, J. Geng, B. Zhang, X. Yu, J. Yang, S. Hu, J. Yu, A comparison between ribo-minus RNA-sequencing and polyA-selected RNA-sequencing. Genomics **96**, 259-265 (2010).

13. N. Cloonan, A. R. Forrest, G. Kolle, B. B. Gardiner, G. J. Faulkner, M. K. Brown, D. F. Taylor, A. L. Steptoe, S. Wani, G. Bethel, A. J. Robertson, A. C. Perkins, S. J. Bruce, C. C. Lee, S. S. Ranade, H. E. Peckham, J. M. Manning, K. J. McKernan, S. M. Grimmond, Stem cell transcriptome profiling via massive-scale mRNA sequencing. Nat. Methods **5**, 613-619 (2008).

14. C. D. Armour, J. C. Castle, R. Chen, T. Babak, P. Loerch, S. Jackson, J. K. Shah, J.

Dey, C. A. Rohl, J. M. Johnson, C. K. Raymond, Digital transcriptome profiling using selective hexamer priming for cDNA synthesis. Nat. Methods **6**, 647-649 (2009).

15. D. Parkhomchuk, T. Borodina, V. Amstislavskiy, M. Banaru, L. Hallen, S. Krobitsch, H. Lehrach, A. Soldatov, Transcriptome analysis by strand-specific sequencing of complementary DNA. Nucleic Acids Res. **37**, e123 (2009).

16. F. Tang, C. Barbacioru, Y. Wang, E. Nordman, C. Lee, N. Xu, X. Wang, J. Bodeau, B. B. Tuch, A. Siddiqui, K. Lao, M. A. Surani, mRNA-Seq whole-transcriptome analysis of a single cell. Nat. Methods **6**, 377-382 (2009).

17. S. Picelli, O. R. Faridani, A. K. Björklund, G. Winberg, S. Sagasser, R. Sandberg, Full-length RNA-seq from single cells using Smart-seq2. Nat. Protoc. 9, 171-181 (2014).

18. T. Hashimshony, N. Senderovich, G. Avital, A. Klochendler, Y. de Leeuw, L. Anavy, D. Gennert, S. Li, K. J. Livak, O. Rozenblatt-Rosen, Y. Dor, A. Regev, I. Yanai, CEL-Seq2: sensitive highly-multiplexed single-cell RNA-Seq. Genome Biol. **17**, 77 (2016).

19. Y. Fu, H. Chen, L. Liu, Y. Huang, Single Cell Total RNA Sequencing through Isothermal Amplification in Picoliter-Droplet Emulsion. Anal. Chem. **88**, 10795-10799 (2016).

20. G. X. Zheng, J. M. Terry, P. Belgrader, P. Ryvkin, Z. W. Bent, R. Wilson, S. B. Ziraldo, T. D. Wheeler, G. P. McDermott, J. Zhu, M. T. Gregory, J. Shuga, L. Montesclaros, J. G. Underwood, D. A. Masquelier, S. Y. Nishimura, M. Schnall-Levin, P. W. Wyatt, C. M. Hindson, R. Bharadwaj, A. Wong, K. D. Ness, L. W. Beppu, H. J. Deeg, C. McFarland, K. R. Loeb, W. J. Valente, N. G. Ericson, E. A. Stevens, J. P. Radich, T. S. Mikkelsen, B. J. Hindson, J. H. Bielas, Massively parallel digital transcriptional profiling of single cells. Nat. Commun. **8**, 14049 (2017).

21. T. M. Gierahn, M. H. Wadsworth, T. K. Hughes, B. D. Bryson, A. Butler, R. Satija, S. Fortune, J. C. Love, A. K. Shalek, Seq-Well: portable, low-cost RNA sequencing of single cells at high throughput. Nat. Methods **14**, 395-398 (2017).

22. J. Cao, J. S. Packer, V. Ramani, D. A. Cusanovich, C. Huynh, R. Daza, X. Qiu, C. Lee, S. N. Furlan, F. J.Steemers, A. Adey, R. H. Waterston, C. Trapnell, J. Shendure, Comprehensive single-cell transcriptional profiling of a multicellular organism. Science **357**, 661-667 (2017).

23. A. B. Rosenberg, C. M. Roco, R. A. Muscat, A. Kuchina, P. Sample, Z. Yao, L. T. Graybuck, D. J. Peeler, S. Mukherjee, W. Chen, S. H. Pun, D. L. Sellers, B. Tasic, G.

Seelig, Single-cell profiling of the developing mouse brain and spinal cord with split-pool barcoding. Science **360**, 176-182 (2018).

24. P. See, J. Lum, J. Chen, F. Ginhoux, A Single-Cell Sequencing Guide for Immunologists. Front Immunol. **9**, 2425 (2018).

25. I. Y. Goryshin, W. S. Reznikoff, Tn5 in vitro transposition. J. Biol. Chem. **273**, 7367–7374 (1998).

26. A. Adey, H. G. Morrison, Asan, X. Xun, J. O. Kitzman, E. H. Turner, B. Stackhouse, A. P. MacKenzie, N. C. Caruccio, X. Zhang, J. Shendure, Rapid, low-input, low-bias construction of shotgun fragment libraries by high-density in vitro transposition. Genome Biol. **11**, R119 (2010).

27. S. Picelli, A. K. Björklund, B. Reinius, S. Sagasser, G. Winberg, R. Sandberg, Tn5 transposase and tagmentation procedures for massively scaled sequencing projects. Genome Res. **24**, 2033-2040 (2014).

28. J. D. Buenrostro, P. G. Giresi, L. C. Zaba, H. Y. Chang, W. J. Greenleaf, Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. Nat. Methods **10**, 1213-1218 (2013).

29. C. Chen, D. Xing, L. Tan, H. Li, G. Zhou, L. Huang, X. S. Xie, Single-cell whole-genome analyses by Linear Amplification via Transposon Insertion (LIANTI). Science **356**, 189-194 (2017).

30. S. Rohrback, C. April, F. Kaper, R. R. Rivera, C. S. Liu, B. Siddoway, J. Chun, Submegabase copy number variations arise during cerebral cortical neurogenesis as revealed by single-cell whole-genome sequencing. Proc. Natl. Acad. Sci. U. S. A. **115**, 10804-10809 (2018).

31. L. Tan, D. Xing, C. H. Chang, H. Li, X. S. Xie, Three-dimensional genome structures of single diploid human cells. Science **361**, 924-928 (2018).

32. B. Lai, Q. Tang, W. Jin, G. Hu, D. Wangsa, K. Cui, B. Z. Stanton, G. Ren, Y. Ding, M. Zhao, S. Liu, J. Song, T. Ried, K. Zhao, Trac-looping measures genome structure and chromatin accessibility. Nat. Methods **15**, 741-747 (2018).

33. J. Gertz, K. E. Varley, N. S. Davis, B. J. Baas, I. Y. Goryshin, R. Vaidyanathan, S. Kuersten, R. M. Myers, Transposase mediated construction of RNA-seq libraries. Genome Res. **22**, 134-141 (2011).

34. S. Brouilette, S. Kuersten, C. Mein, M. Bozek, A. Terry, K. R. Dias, L. Bhaw-Rosun, Y.

Shintani, S. Coppen, C. Ikebe, V. Sawhney, N. Campbell, M. Kaneko, N. Tano, H. Ishida,

K. Suzuki, K. Yashiro, A simple and novel method for RNA-seq library preparation of single

cell cDNA analysis by hyperactive Tn5 transposase. Dev. Dyn. **241**, 1584-1590 (2012).

35. K. A. Majorek, S. Dunin-Horkawicz, K. Steczkiewicz, A. Muszewska, M. Nowotny, K.

Ginalski, J. M. Bujnicki, The RNase H-like superfamily: new members, comparative

structural analysis and evolutionary classification. Nucleic Acids Res. **42**, 4160-4179

(2014).

36. D. R. Davies, I. Y. Goryshin, W. S. Reznikoff, I. Rayment, Three-dimensional structure

of the Tn5 synaptic complex transposition intermediate. Science **289**, 77-85 (2000).

37. W. S. Reznikoff, Transposon Tn5. Annu. Rev. Genet. **42**, 269-286 (2008).

38. G. Peterson, W. Reznikoff, Tn5 transposase active site mutations suggest position of

donor backbone DNA in synaptic complex. J. Biol. Chem. **278**, 1904-1909 (2003).

39. M. Nowotny, S. A. Gaidamakov, R. J. Crouch, W. Yang, Crystal structures of RNase

H bound to an RNA/DNA hybrid: Substrate specificity and metal-dependent catalysis. Cell

121, 1005-1016 (2005)

40. D. Lim, G. G. Gregorio, C. Bingman, E. Martinez-Hackert, W. A. Hendrickson, S. P.

Goff, Crystal structure of the Moloney murine leukemia virus RNase H domain. J. Virol.

80, 8379-8389 (2006).

Supplementary Figures

Fig.S1 Structure of Tn5 and D188E mutation in Tn5. (**A**) Structure of pTXB1 Tn5 (PDB ID: 1MUS).

Left monomer marked domains in different colors. Right monomer marked residues of catalytical core

and putative substrate binding site in atom form. [Referred to D. R. Davies, I. Y. Goryshin, W. S.

Reznikoff, I. Rayment, Three-dimensional structure of the Tn5 synaptic complex transposition

intermediate. Science 289, 77-85 (2000).] Black box showed D188E mutation. (**B**) Size distribution of

genomic DNA with no treatment or tagmented by D188E mutant Tn5 or tagmented by pTXB1 Tn5. (**C**)

Model of docking double-stranded DNA (Left) or RNA/DNA heteroduplex (Right) in predicted substrate

binding sites of Tn5.

Fig.S2 Composition of products amplified from tagmented RNA/DNA hybrid. Gray wavy line and

straight line represent RNA and DNA separately. Dotted lines represent the track of extension step.

A

	Mapping Rate	Exonic Rate	Gene Detected
Ligation Test 1	92.20±0.20%	77.76±0.18%	11,825±7
Ligation Test 2	93.30±0.00%	78.98±0.10%	11,784±1
SHERRY	89.77±0.05%	80.32±0.04%	11,784±8
Strand Test 1	11.25±0.05%	54.92±0.06%	7,384±13
Strand Test 2	4.05±0.15%	61.21±0.32%	5,566±81
Strand Test 3	78.20±0.50%	77.38±0.10%	11,470±14
dU-SHERRY	90.90±0.10%	76.31±0.00%	11,335±40

576 **Fig.S3** Comparison within Ligation Tests and Strand Tests. (**A**) Sequencing indicators of Ligation Tests

577 and Strand Tests. Each test consisted of two replicates of 200 ng HEK293T total RNA. (**B-C**) Insert

578 size distribution of Ligation Tests. The colored bars marked reads which insert size is shorter than

579 100bp. Adaptors detected in these reads are counted.

B

	Mapping Rate	Exonic Rate	Gene Detected
SHERRY (10ng)	90.33±0.12%	76.73±0.12%	11,439±48
NEBNext Ultra II (10ng)	93.47±0.39%	87.25±0.20%	12,338±71
SHERRY (200ng)	89.77±0.05%	80.89±0.32%	11,784±10
NEBNext Ultra II (200ng)	93.17±0.17%	86.98±0.04%	12,374±10

580
581
582 **Fig.S4** Optimization of SHERRY and comparison with NEBNext. (**A**) Gene number detected by
583 SHERRY under various experimental conditions. Each condition consisted of three replicates of 10 ng
584 HEK293T total RNA. (**B**) Comparison of sequencing indicators between SHERRY and NEBNext with
585 10 ng and 200 ng HEK293T total RNA input. Each condition consisted of three replicates and down-
586 sampled to 2 million total reads.
587

Fig.S5 Functional comparison between SHERRY and NEBNext. (**A**) Correlation of normalized gene

counts among duplicates of SHERRY, which start from 200 ng HEK293T total RNA input. (**B**)

Correlation of normalized genes counts (average of three replicates) between SHERRY and NEBNext

within the two cell types. The input was 200 ng total RNA. (**C**) Differentially expressed genes of HeLa

and HEK293T detected by SHERRY and NEBNext kit (200 ng input) are plotted into Venn Diagram.

Colored area represents genes identified by both methods. Gene numbers are listed on corresponding

part. (**D**) Heatmap of differentially expressed genes detected by SHERRY while missed by NEBNext

kit. The Color bar indicates Z-score. (**E**) Heatmap of differentially expressed genes detected by

NEBNext kit while missed by SHERRY. The Color bar indicates Z-score.

A

	Mapping Rate	Exonic Rate	Gene Detected
Replicate1	64.60%	62.60%	8,955
Replicate2	61.80%	59.90%	8,783
Replicate3	64.80%	63.00%	8,733

B

		Mapping Rate	Exonic Rate	Gene Detected
Crowding Agent in Tagmentation	8% PEG8000 (m/v)	50.17±2.72%	44.28±6.98%	8,338±345
	20% PEG400 (v/v)	29.98±6.92%	56.53±2.13%	7,835±326
	20% Betaine (v/v)	67.63±2.51%	43.47±5.38%	2,846±367
Extension Enzyme	Bst 3.0	34.53±5.49%	60.57±3.30%	7,966±209
	Bst 2.0 Warmstart	34.45±3.62%	58.08±7.79%	7,790±321
	SuperScript™ II	33.25±6.67%	55.11±4.74%	5,065±307
	SuperScript™ III	33.20±13.68%	48.33±4.34%	4,803±550
PCR Cycle	28 cycle	24.43±7.65%	52.46±7.11%	7,590±778
	10 cycle + 18 cycle	50.17±2.72%	44.28±6.98%	8,338±345
	15 cycle	31.97±2.85%	53.94±9.83%	8,352±255

C

D

	Mapping Rate	Exonic Rate	Gene Detected
scSHERRY	50.17±2.72%	44.28±6.98%	8,338±345
Smart-seq2	91.23±0.46%	72.78±2.26%	9,761±577

E

Fig.S6 Optimization of micro-input SHERRY and comparison with Smart-seq2. (**A**) Sequencing indicators of SHERRY (n=3) starting from 100 pg HEK293T total RNA input. (**B**) Comparison of scSHERRY library quality under various experiment conditions. Each condition used single HEK293T cell as input and consisted of 3-4 replicates. (**C**) Correlation of normalized gene counts (average of three replicates) between scSHERRY and Smart-seq2. (**D**) Comparison of sequencing indicators between scSHERRY (n=3) and Smart-seq2 (n=4). Both of them used single HEK293T cells as input. (**E**) The coverage of GAPDH transcript calculated from scSHERRY and Smart-seq2.

609
610
611 **Fig.S7** Comparison of workflow and cost among Smart-seq2, SHERRY and NEBNext kit. **(A)** Workflow
612 of Smart-seq2, SHERRY and NEBNext kit. Length of arrow indicates time consumed for each step.
613 The human-shaped icon indicated hands-on time. Dotted box means this step is alternative. Gray line
614 connects corresponding key step in each method. **(B)** Cost list of Smart-seq2, SHERRY and NEBNext
615 kit.

616
617

618
619

620   **Fig.S8** Saturation curve of scSHERRY (n=3) and Smart-seq2 (n=4). Both methods used single

621   HEK293T cell as input. Dotted line indicated 2 million total reads.

622
623
624

	Mapping Rate	Exonic Rate	Gene Detected	Median Size
pTXB1-Tn5	88.80%	74.93%	11,416	113
	88.20%	75.02%	11,492	118
V50 (Vazyme)	88.90%	77.44%	11,501	149
	88.80%	76.89%	11,542	146
ATM (Illumina)	90.20%	79.29%	11,666	168
	90.40%	79.25%	11,612	166

627   **Fig.S9** Hybrid tagmentation activity of commercial Tn5. (**A**) Size distribution of genomic DNA with no

628   treatment or tagmented by different volumes of pTXB1 Tn5, V50 or ATM. The dotted black line

629   indicates peak of fragment size. (**B**) Size distribution of mRNA/DNA hybrid with no treatment or

630   tagmented by different volumes of pTXB1 Tn5, V50 or ATM. (**C**) Sequencing indicators of SHERRY

631   library constructed by three Tn5.

633
634
635 **Fig.S10** Titration of Tn5 transposome for tagmentation. (**A-B**) Size distribution of 5ng genomic DNA

636 or mRNA/DNA hybrid with no treatment or tagmented by different gradients of pTXB1 Tn5.

644
645
646 **Fig.S11** Coverage evenness optimization of SHERRY (10 ng HEK293T total RNA input). (**A**)

647 Normalized transcript coverage of standard SHERRY, SHERRY using TSO-RT method and NEBNext

648 kit. (**B**) The coverage of GAPDH transcript calculated from SHERRY and TSO-RT SHERRY. (**C**)

649 Comparison of sequencing indicators between SHERRY (n=3) and TSO-RT SHERRY (n=2).

Materials and Methods

Purification of pTXB1 Tn5 and D188E mutation

The pTXB1 cloning vector, which introduced hyperactive E54K and L372P mutation into wildtype Tn5, was acquired from Addgene. The pTXB1 Tn5 and its mutant were expressed and purified mainly according to the protocol published by Picelli S et al. [1] To construct D188E mutation into Tn5, pTXB1 vector was firstly amplified into two parts by two sets of primers. Mutagenesis primers used for the first part (3771-7979) which contained site 188 were 5'-GGCAGCATGATGAGCAACGTGATTGCGGTGTGC**GAA**CG TGAAGCGGATATTCATGC-3' and 5'-TATCAGCTCACTCAAAGG-3'. Amplified primers for the remaining part were 5'-GTATTACCGCCTTTGAGT-3' and 5'-CAATCACGTTGCTC ATCA-3'. The purified PCR products were then assembled into intact plasmid using Gibson Assembly Master Mix (NEB, Cat.No. E2611). The newly assembled plasmid was transformed into E.coli Trans5α chemically competent cells (Transgene, CD201-01). After growing overnight on LB medium plate, single colony was picked and shaken in SOC liquid medium for at least 9 hours. Plasmid was extracted by PurePlasmid Mini Kit (CWBIO, Cat.No.CW0500S) and confirmed carrying D188E mutation by Sanger sequencing. Then the plasmid was transformed into E.coli Transetta (DE3) chemically competent cells (Transgene, CD801-01) for further protein expression and purification.

Cell culture

HEK293T and HeLa cell lines are acquired from ATCC. Both of them were cultured in Dulbecco's Modified Eagle Medium (Gibco, Cat.No.11965092), supplemented with 10% fetal bovine serum (Gibco, Cat.No.1600044) and 1% penicillin-streptomycin (Gibco, Cat.No.15140122). The cell incubator (Thermo Scientific) was set at temperature of 37°C with 5% CO_2 injected.

Adherent cells were washed twice by DPBS (Gibco, Cat.No.14190136) and detached by 0.05% Trypsin-EDTA (Gibco, Cat.No.25300062) at 37°C for 4min. Then double volume of culture media was added to terminate trypnization. Cells were collected by centrifugation at 200g for 5min and resuspended for downstream experiment or passage cultivation.

Nucleic acids extraction and messenger RNA isolation

Genomic DNA was extracted using PureLink Genomic DNA Mini Kit (Invitrogen, Cat.No.K182002), and total RNA was extracted using RNeasy Mini Kit (Qiagen, Cat.No.74104). The resulting total RNA was then reacted with 10ul DNase I (NEB, Cat.No.M0303) to remove remaining DNA thoroughly, and concentrated by RNA Clean & Concentrator-5 kit (Zymo Research, Cat.No.R1015). The quality of extracted DNA and RNA was assessed by the Fragment Analyzer Automated CE System (AATI) and quantification was done by Qubit 2.0 (Invitrogen, Cat.No.Q33230/ Q32852).

We followed standard protocol of NEBNext Poly(A) mRNA Magnetic Isolation Module (NEB, Cat.No.E7490) to isolate messenger RNA from the purified total RNA and stored them at -80°C.

Single cell preparation

We used pipette tips to form some drops made up of PBS (containing 1% BSA) (Thermo Scientific, Cat.No.37525) on a clean petri dish. The cell resuspension was pipetted up and down gently to disperse into single cells and we took ~5µl of them diluted in one of the drops. The mouth pipette with 50µm inside diameter was then used to pick one cell in the drop and release it in another clean drop. The picked cell was passed by at least three clean drops in order to wash away any debris and confirm that only one cell was in the last drop. We then aspirated the cell with as little buffer as possible and blew it into 4µl lysis buffer [4 units of Recombinant RNase Inhibitor (Takara, Cat.No.2313), 2.5µM poly(T)$_{30}$VN primer (Sangon), 2.5mM dNTP (NEB, Cat.No.N0447) and 0.48% Triton X-100 (Sigma, Cat.No.T9284)]. Successful transfer was confirmed by blowing mouth pipette again in a clean drop and no cell was to be seen in visual field. Reaction was carried out at 72°C for 3min after violent vortex.

mRNA/DNA hybrid formation

Total RNA was reverse transcribed into mRNA/DNA hybrid mainly referring to Smart-seq2 protocol [2], but with several modifications: 1) The ISPCR part in Oligo-dT primer was removed; 2) TSO was omitted, but in TSO-RT SHERRY, it should be kept; 3) The reaction was performed at 42°C for 1.5h without cycling. If input was purified RNA, Triton X-100 was omitted. When inputting more than 10ng total RNA, we would slightly upregulate

amount of dNTP, poly(T)$_{30}$VN primer and Superscript II (Invitrogen, Cat.No.18064014).

Tn5 transposome in vitro assembly and tagmentation

Functional mosaic-end (ME) oligonucleotides (5'-CTGTCTCTTATACACATCT-3', 100µM)

was separately annealed with equal amounts of Adaptor A (5'-TCGTCGGCAGCGTCAG

ATGTGTATAAGAGACAG-3', 100µM) and Adaptor B (5'-GTCTCGTGGGCTCGGAGATG

TGTATAAGAGACAG-3', 100µM). The concentration of purified Tn5 was quantified by

Qubit Protein Assay Kit (Invitrogen, Cat.No.Q33212) and took around 100µg for

transposome assembly. We mixed the Tn5 transposase with annealed ME-Adaptor A or

B (20µM) in 45% glycerol (Sigma, Cat.No.G5516) thoroughly and incubated the mixture

at 30°C for one hour. These two resulting transposomes (assembled with ME-Adaptor

A/B) were then mixed together, ready for tagmentation or stored at -20°C. Specifically, to

assemble rCrArG Tn5 in **Fig.S4**, ribonucleotide modifications were made on the three

terminal bases at 3'-end of Adaptor A/B. And for rG Tn5, the last base at 3'-end of adaptors

was modified.

The dsDNA tagmentation was performed in 1xTD buffer [10mM Tris-Cl (pH 7.6,

ROCKLAND, Cat.No.MB-003), 5mM MgCl$_2$ (Invitrogen, Cat.No.AM9530G), 10% N,N-

Dimethylformamide (Sigma, Cat.No.D4551)]. The reaction was incubated at 55°C for

30min.

As for RNA/DNA hybrid, tagmentation was performed in buffer containing 10mM Tris-Cl

(pH 7.6), 5mM MgCl$_2$, 10% N,N-Dimethylformamide, 9% PEG8000 (VWR Life Science,

Cat.No.97061), 0.85mM ATP (NEB, Cat.No.P0756). In SHERRY library preparation, we

used 0.05 µl, 0.006 µl and 0.003 µl Tn5 transposome for input of 200 ng, 10 ng and 100

pg total RNA, respectively. scSHERRY also used 0.003 µl Tn5 transposome.

The transposome could be diluted in 1xTn5 dialysis buffer [50mM Hepes (pH 7.2,

Leagene, Cat.No.CC064), 0.1M NaCl (Invitrogen, Cat.No. AM9759), 0.1 mM EDTA

(Invitrogen, Cat.No.AM9260G), 1 mM DTT, 0.1% Triton X-100, 10% glycerol].The reaction

was incubated at 55°C for 30min.

Commercial Tn5 transposomes were available in Nextera XT DNA Library Prep Kit

(Illumina, Cat.No.FC-131-1024) and TruePrep DNA Library Prep Kit V2 for Illumina

(Vazyme, Cat.No.TD501).

SHERRY library preparation and sequencing

To construct scSHERRY library, the single cell tagmentation product was mixed well with 4 units of Bst 3.0 DNA Polymerase (NEB, Cat.No.M0374) and indexed common primers (Vazyme, Cat.No.TD202) in 1 x Q5 High-Fidelity Master Mix (NEB, Cat.No.M0492). Then index PCR was performed as follow: 72°C 15min, 98°C 30s, 10 cycles of [98°C 20s, 60°C 20s, 72°C 2min], 72°C 5min. The PCR product was purified with 0.85:1 ratio by VAHTS DNA Clean Beads (Vazyme, Cat.No.N411) and eluted in 30µl nuclease-free water (Invitrogen, Cat.No.AM9937) for another 18 cycles of PCR. When performing high-throughput experiment, each sample could be amplified by 15 cycles, then merged for beads purification and library quality check.

As for purified 10ng or 200ng total RNA input, the tagmentation product was firstly gap-filled with 100 units of Superscript II and 1 x Q5 High-Fidelity Master Mix at 42°C for 15min, then Superscript II was inactivated at 70°C for 15min. When inputting 100pg total RNA, the extension enzyme was replaced with 4 units of Bst 2.0 Warmstart DNA Polymerase (NEB, Cat.No.M0538). Correspondingly, the reaction temperature was upregulated to 72°C and inactivation was performed at 80°C for 20min. After that, indexed common primers were added to perform PCR. We performed 12, 15 and 25 cycles of PCR for input of 200 ng, 10 ng and 100 pg total RNA, respectively.

The resulting library was purified with 1:1 ratio by VAHTS DNA Clean Beads. Quantification was done by Qubit 2.0 and quality check was done by Fragment Analyzer Automated CE System. The sequencing platform we used was Illumina NextSeq 500 or HiSeq 4000.

NEBNext and SmartSeq2 library preparation

NEBNext RNA-Seq library preparation starting from 10ng and 200ng total RNA was performed using NEBNext Ultra II RNA Library Prep Kit for Illumina (NEB, Cat.No.E7770). Single cell SmartSeq2 library was constructed as previously reported [2].

Ligation Test and Strand Test

In Ligation Tests, tagmentation products from 200ng HEK293T total RNA were purified by

DNA Clean & Concentrator-5 (Zymo Research, Cat.No.D4013) and eluted with 20μl
nuclease-free water. After gap-filling with Superscript II, Ligation Test 1 was processed
directly to index PCR while product in Ligation Test 2 was digested by 12.5 units of RNase
H (NEB, Cat.No.M0297) at 37°C for 20min before index PCR.

In Strand Tests, we used dUTP (Thermo Scientific, Cat.No.R0133), dATP, dCTP, dGTP
(NEB, Cat.No.N0446) mix, each of them at equal concentration, to incorporate in cDNA
during reverse transcription step. 0.15μl Tn5 transposome was used to tagment the
resulting hybrid. Fragments were then column-purified and gap-filled by Bst 2.0 Warmstart
DNA Polymerase, and column purification was again applied. For Strand Test 1, 3 units
of USER enzyme (NEB, Cat.No.M5505) and 40units of recombinant rnase inhibitor was
added into elution products and incubated at 37°C for 20min for DNA strand digestion.
Indexed common primers added with digestion product in 1 x Q5 High-Fidelity Master Mix
were then reacted at 85°C for 30s, followed by 60°C for 2min and temperature went down
to 4°C slowly. After that, reverse transcription was performed with 200 units of Superscript
II added at 42°C for 30min, then transferred to index PCR program. For Strand Test 2,
the USER digestion product was directly performed index PCR with 1 x KAPA HiFi
HotStart Uracil+ ReadyMix (Kapa Biosystems, Cat.No.KK2801). Protocol of Strand Test
3 was almost same as Strand Test 2, except replacing USER enzyme with RNase H. For
dU-SHERRY, the USER enzyme digestion was omitted compared with Strand Test 2
workflow.

Docking model
To generate the substrate-transposon DNA-Tn5 structure model, a 32bp dsDNA or
DNA/RNA hybrid were generated by 3D-NuS (3-Dimensional Nucleic Acid Structures)
web server. Then the substrate dsDNA or DNA/RNA was manually docked to the
transposon DNA-Tn5 structure, PDB ID: 1MUS, based on charge and shape
complimentary.

Data analysis
Sequencing adaptors or poly(T/A) positioned at end of paired reads were recognized and
removed by Cutadapt v1.15 [3]. The trimmed reads which length was shorter than 20bp

were filtered. Remaining reads were down sampled to 2 million (except that library with 200ng total RNA input used for differential gene expression analysis was 10 million) total reads, and aligned with index built from human(hg38) genome and known transcript annotations by Tophat2 v2.1.1 [4]. The mapped reads were then used to calculate FPKM value for each known gene (annotation acquired from UCSC) by Cufflinks v2.2.1 with multi-mapped reads correction. Gene with FPKM more than 1 was considered to be detected. The exonic rate, duplicate rate and insert size of library were all calculated by Picard Tools v2.17.6.

General coverage across known transcripts was plotted by RSeQC v.2.6.4 [5]. For specific transcript, depth of mapped reads overlapped with transcript position was calculated by Samtools v1.3.1.

We used DESeq2 v1.22.2 [6] to perform differential gene expression analysis with raw count-matrix acquired by HTSeq v.0.11.0 [7]. Differentially expressed genes should meet following criteria: 1) FPKM value >1; 2) significant p-value $<5\times10^{-6}$; 3) absolute value of $\log_2$(Fold Change) >1. Counts in correlation plot were normalized mainly according to DESeq2 normalization method [6], which considered library size and library compensation. Correlation efficient R^2 and slope of linear fitting equations were calculated by least square method. The slope-skewness between two replicates was defined as $|k_1-1|+|k_2-1|$, k_1 or k_2 was slope when one of the replicates was conducted as X or Y axis.

GC content distribution of detected genes was plotted by custom Perl script. GC content was binned by 2% and gene number at each bin was normalized by dividing maximum gene number of one bin. And for each gene, only the longest transcript isoform was calculated.

Data deposition

The sequence reported in this paper has been deposited in the Genome Sequence Archive (accession no. CRA002081).

References

[1] S. Picelli, A. K. Björklund, B. Reinius, S. Sagasser, G. Winberg, R. Sandberg, Tn5 transposase and tagmentation procedures for massively scaled sequencing projects.

Genome Res. 24, 2033-2040 (2014).

[2] S. Picelli, O. R. Faridani, A. K. Björklund, G. Winberg, S. Sagasser, R. Sandberg, Full-
length RNA-seq from single cells using Smart-seq2. Nat. Protoc. 9, 171-181 (2014).

[3] M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing
reads. Embnet Journal 17, 200 (2011).

[4] D. Kim, G. Pertea, C. Trapnell, H. Pimentel, R. Kelley, S. L. Salzberg, TopHat2:
accurate alignment of transcriptomes in the presence of insertions, deletions and gene
fusions. Genome Biol. 14, R36 (2013).

[5] L. Wang, S. Wang, W. Li, RSeQC: quality control of RNA-seq experiments.
Bioinformatics 28, 2184-2185 (2012).

[6] M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 15, 550 (2014).

[7] S. Anders, P. T. Pyl, W. Huber, HTSeq—a Python framework to work with high-
throughput sequencing data. Bioinformatics 31, 166-169 (2015).

858
859