1 **Phosphate transporter PstSCAB of *Campylobacter jejuni* is essential for lactate-dependent**
2 **growth and colonization in chickens**
3

4 Ritam Sinha[1], Rhiannon M. LeVeque[1], Marvin Q. Bowlin[2], Michael J. Gray[2], and Victor J.
5 DiRita[1*]
6
7
8 [1]Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing,
9 Michigan, 48823
10
11 [2]Department of Microbiology, University of Alabama at Birmingham, Birmingham, Alabama,
12 35294
13
14
15 *Corresponding Author:

16 Mailing Address:

17 Department of Microbiology and Molecular Genetics,

18 Michigan State University, East Lansing, MI-48824

19 Email: diritavi@msu.edu.

42
43
44   Abstract
45

Campylobacter jejuni causes acute gastroenteritis world-wide and is transmitted primarily

through poultry, in which it is often a commensal member of the intestinal microbiota. Previous

RNASeq experiments showed that transcripts from an operon encoding a high affinity phosphate

transporter (PstSCAB) of *C. jejuni* were among the most abundant when grown in chickens.

Elevated levels of the *pstSCAB* mRNA were also identified in an RNASeq experiment from

human infection studies. In this study, we explore the role of PstSCAB in the biology and

colonization potential of *C. jejuni.* Our experimental results demonstrate that cells lacking

PstSCAB survive poorly in stationary phase, nutrient-limiting media, and under osmotic

conditions reflective of those in the chicken. Polyphosphate levels in the mutant cells were

elevated at stationary phase, consistent with alterations in expression of polyphosphate

metabolism genes. *C. jejuni* were highly attenuated in colonization of newly hatched chicks,

recovered at levels several orders of magnitude below wild type. Mutant and wild type grew

similarly in complex media but the *pstSCAB* mutant exhibited a significant growth defect in

minimal media supplemented with L-lactate, postulated as a carbon source *in vivo*. Poor growth

in lactate correlated with diminished expression of acetogenesis pathway genes previously

demonstrated as important for colonizing chickens. The phosphate transport system is thus

essential for diverse aspects of *C. jejuni* physiology and *in vivo* fitness and survival.

Importance:

C. jejuni causes millions of gastrointestinal infections annually worldwide. Poultry and poultry products are major sources of *C. jejuni* infection to human as the microbe is a commensal colonizer of the chicken gastrointestinal tract. Due to the emergence of multi-drug resistance in *C. jejuni*, there is need to identify alternative ways to control this pathogen. Genes encoding the high-affinity phosphate transporter PstSCAB were highly expressed during colonization of *C. jejuni* in chicken and human. In this study, we address the role this high-affinity phosphate transporter PstSCAB of *C. jejuni* on chicken colonization and for its general physiology. PstSCAB is required for colonization in chicken, metabolism and survival under different stress responses and during growth on lactate, a potential substrate for growth of *C. jejuni* in chickens. Our study highlights that PstSCAB may be an effective target to develop mechanisms to control the bacterial burden in both chicken and human.

109 **Introduction**
110

Campylobacter jejuni is a microaerophilic, Gram-negative bacterium that causes acute

gastroenteritis worldwide (1). Common campylobacteriosis symptoms are self-limiting diarrhea

(which may be watery to bloody), with nausea, fever and abdominal cramping in human. As *C.*

jejuni is a common member of the chicken gastrointestinal microbiota, poultry meat is a major

source of human campylobacteriosis, either through direct consumption of poultry or through

cross-contamination of other food (2, 3). As consumption of poultry meat worldwide over the

last 20 years has approximately doubled, (4) the threat of campylobacteriosis remains high.

To explore mechanisms by which *Campylobacter* colonizes the chicken gastrointestinal

tract, we used RNASeq to determine the transcriptome of *C. jejuni* strain 81-176 harvested from

ceca of seven day-old chickens after experimental oro-gastric inoculation on day-of-hatch (5).

Among the more highly abundant transcripts in this population were those encoding a predicted

phosphate transporter (*pstSCAB*), suggesting that phosphate transport is important for the *in vivo*

fitness of *C. jejuni*, and perhaps that inorganic phosphate plays a key role of chicken colonization

(5).

Phosphorus, or inorganic phosphate (Pi), is one of the most important macronutrients for

microbial growth. It is required for biosynthesis of cellular components such as nucleic acid,

phospholipids and proteins, and is critical for maintaining a cellular energy source in ATP. It is

also involved in regulating carbon and nitrogen metabolism (6). Uptake and storage of inorganic

phosphate, a growth-limiting nutrient, is heavily regulated in bacteria. Generally, in Gram-

negative bacteria, a two-component sensor-kinase/response regulator system controls phosphate

transport in response to levels of available inorganic phosphate. In *C. jejuni*, a two component

regulatory system, PhosSR, controls the high affinity phosphate transporter operon (*pstSCAB*),

as well as other genes, in phosphate limiting condition (7). PstSCAB comprises an ATP binding

cassette (ABC) transporter system in the inner membrane of many Gram-negative bacteria, with

PstS the periplasmic phosphate binding protein, PstAC comprising transmembrane protein

permeases and PstB an ATP binding and hydrolyzing subunit. PstS bound to Pi is stimulated to

associate with PstA/C transmembrane proteins. This triggers ATP hydrolysis in the PstB dimer

which induces the conformational changes that facilitate release of Pi into the cytosol (8). In

many bacteria a PstSCAB-associated protein, PhoU, acts to reduce expression of the Pho regulon

under conditions of phosphate sufficiency (6), but a PhoU homologue has not been reported in *C.*

jejuni.

Phosphate transporters contribute to virulence in other microbes, although the molecular

mechanisms by which that occurs varies across species and in any event are incompletely

understood (9). For example, inoculation into chickens of an avian pathogenic *E.coli* mutant

strain lacking *pst* led to fewer extra-intestinal lesions than did inoculation of wild type, and the

mutant also showed higher sensitivity to rabbit serum (though not to chicken serum), polymyxin

B, and acid shock (10). A *pstS* mutant of virulent EHEC and EPEC showed significant adherence

and colonization defects in intestinal epithelial cells as well as in animal models, resulting from

altering expression of adhesin proteins (9, 11). *Mycobacterium tuberculosis* contains two

different phosphate transporters which regulate replication in macrophages and provide

protection from antimicrobial stress response (12, 13).

Given the high abundance of transporter mRNA during *in vivo* growth of *C. jejuni*, it

seems likely that the transporter is a determinant of colonization. We analyzed a mutant lacking

pstSCAB for several growth phenotypes as well as chicken colonization. We demonstrate a

severe colonization defect in mutants lacking the *pstSCAB* operon, and we also uncover

metabolic deficiencies in the mutant strain – particularly its inability to thrive on L-lactate at

concentrations approximating those in the chicken cecum - that likely contribute to loss of *in*

vivo fitness.

159   **Results**
160
Mutagenesis of and characterization of the *C. jejuni pstSCAB* operon.   We constructed an

insertion allele using a cassette encoding resistance to kanamycin, inserted into the *pstS* gene of

C. jejuni strain 81-176 **(Fig.S1).** The resulting kanamycin resistant strain was used in qRT-PCR

experiments designed to analyze transcription of the *pstSCAB* locus. Consistent with the earlier

RNASeq study (5), qRT-PCR of the wild type locus using primers specific for the junctions

between open reading frames confirmed that the four genes are co-transcribed, a conclusion

further supported by the observation that insertion in the *pstS* gene, the most upstream gene of

the locus, resulted in loss of detectable downstream transcription (**Fig. 1**).

In the earlier RNASeq study abundance of the *pstS* mRNA was significantly higher than

distal genes in the operon (5). Quantitative RT-PCR experiments confirmed that mRNA

encoding *pstS* is more abundant than that of the *pstCAB* genes **(Fig. S2A)**. Differential stability

within a polycistronic mRNA can be controlled through a conserved internal RNA hairpin that

protects the transcript from RNA decay (14). The protective RNA structure is often present in the

open reading frame of the downstream gene to protect the essential upstream gene from decay

(14). Consistent with such a mechanism acting on the *pstSCAB* operon, we identified a

thermodynamically stable hairpin loop structure ($\Delta G = -27.70$) within the protein coding region of

pstC gene **(Fig. S2B and C**). We hypothesize this RNA structure in *pstC* protects *pstS* mRNA

from decay, leading to its elevated abundance relative to the other genes in the operon. Because

of polarity from the *pstS::kan* insertion, for further experiments we used a complementing

plasmid encoding the entire *pstSCAB* operon. In describing work with mutant and complemented

strains, the strain is generally designated by its genotype (*pstS::kan*), and the strain with the

mutation and carrying the complementing plasmid is designated *pstS::kan/C.*

Phosphate storage and metabolism gene products are altered in the *pstSCAB* mutant.

Survival of *C. jejuni* in stationary phase and nutrient limiting condition is dependent on

polyphosphate (poly-P) metabolism (15). *C. jejuni ppk1* and *ppk2* encode poly-P kinases, with

the former responsible for poly-P synthesis, using ATP generated from inorganic phosphate (16),

and the latter generating GTP from poly-P (17). In some microbes Ppk2 can produce poly-P

from GTP (17), but this activity has not been reported in *C. jejuni*. *C. jejuni* also encodes two

exo-polyphosphatases, Ppx1 and Ppx2, which degrade poly-P into inorganic phosphate and have

guanosine pentaphosphate hydrolase activity to generate guanosine tetraphosphate (ppGpp) (18).

With the regulatory gene product SpoT, ppGpp regulates the stringent response of *C.jejuni,*

contributing to survival in stationary phase (19). Another key gene of phosphate metabolism is

ppa, which encodes an inorganic pyrophosphatase that hydrolyses pyrophosphate to inorganic

phosphate (PPi→Pi)(19).

We investigated the effect of a *pstSCAB* mutation on these phosphate storage and stress

response mechanisms, examining expression in wild type, mutant, and complemented mutant

strains after 24 and 48 hours of growth in MHB. The gene largely responsible for poly-P

synthesis, *ppk1*, was unchanged in its level of expression in the mutant versus the wild type at

either time point. Expression of *ppk2,* which converts GDP to GTP using poly-P as a substrate -

and is reported in some microbes (although not in *C. jejuni*) to produce poly-P from GTP - was

elevated up to 10-fold in the mutant strain at 24 hours, returning to wild type expression levels

by 48 hours **(Fig. 2A)**. At 48 hours, expression levels of an exo-polyphosphatase gene (*ppx2*),

the inorganic pyrophosphatase gene *ppa,* and the stringent regulatory gene *spoT* were

significantly reduced in the mutant compared to wild type (**Fig. 2B**). As *ppx2* and *spoT* of

C.jejuni regulate synthesis of ppGpp, important for survival in stationary phase, reduced

expression of these two genes would likely result in decrease in the level of ppGpp in the

pstS::kan mutant strain.

At 24 hours, poly-P levels in the mutant were roughly half of those observed in the wild

type. At 48 hours, by which time the mutant had diminished in viability by several orders of

magnitude (see below, Fig. 3B), poly-P was elevated to about three times the levels observed in

wild type (**Fig. 2C and D)**. With elevated *ppk2* expression at 24 hours in the *pstS::kan* mutant,

decreased intracellular poly-P might be expected, as the Ppk2 enzyme consumes poly-P to

convert GDP to GTP. The higher levels of poly-P observed in the *pstS::kan* mutant compared to

wild type at 48 hours is readily explained by decreased expression of the exopolyphosphatase

gene *ppx2* in the mutant background, and is consistent with a similar report of a *pstS::kan* mutant

of *Pseudomonas aeruginosa* (20).

***pstS* is essential for stationary phase survival and osmotic stress response.** Expression of *pstS*

is elevated approximately four-hundred-fold in 0.08 mM inorganic phosphate (Pi) compared with

1.6 mM Pi, through activation of the PhoRS system at the more limiting phosphate concentration

(7). The PhoRS-regulated alkaline phosphatase gene of *C. jejuni* (*phoX*) is also induced at lower

phosphate concentration, providing Pi for transport into the cell by the PstSCAB system (21).

We hypothesized that the *pstS::kan* mutant strain would signal phosphate limitation to the cell,

even at a concentration (1.6 mM) that does not typically result in *phoX* expression. We measured

alkaline phosphatase activity in wild type and *pstS::kan* mutant cells grown in 0.08 mM and 1.6

mM phosphate. Supporting our hypothesis, the mutant expressed significantly elevated alkaline

phosphatase activity at both the lower (0.08 mM) and higher (1.6 mM) concentrations, whereas

wild type (and the complemented mutant) expressed high levels of alkaline phosphatase activity

only at the lower phosphate concentration (**Fig. 3A**). We conclude that, even at elevated

phosphate concentrations, the *pstS::kan* mutant strain senses phosphate limitation.

To assess the effect of loss of *pstSCAB* on growth of *C. jejuni,* we cultured wild type,

mutant, and complemented mutant strains in Mueller-Hinton broth (MHB), a complex medium

of protein hydrolysate and infusion. All grew to similar levels for 24 hours, peaking at 10^{10}

cfu/ml **(Fig. 3B).** By 48 hours, the wild type and complemented mutant strains had dropped

about two orders of magnitude in plating efficiency, while the mutant strain had dropped by over

five orders of magnitude in plating efficiency, suggesting a severe defect in survival in stationary

phase and consistent with the diminished levels of poly-P we detected in the cells at that time-

point. Assessing survivability in media depleted of nutrients provided a more direct examination

of the challenge posed by loss of *pstSCAB*. In this experiment the wild type strain was grown for

48h in MH broth to reach stationary phase, after which the growth media was collected by filter

sterilization. We inoculated mid-log phase wild type, *pstS::kan* and the complemented mutant in

the sterilized depleted media, with initial inoculum (5×10^8 CFU/ml), and assessed viable cells at

24 hours. In contrast to wild type and complemented mutant, which were still recoverable at high

numbers in depleted media, the mutant was completely unrecoverable **(Fig. 3C)** confirming that

loss of the phosphate uptake system causes a significant defect in growth physiology. Likely

contributing to this loss of fitness under limiting nutrient conditions **(Fig. 3D)** is the decreased

expression demonstrated **(Fig. 2B)** in of *ppx2* and *spoT,* regulators of ppGpp synthesis.

In addition to stationary phase survival, poly-P metabolism contributes to survival under

osmotic stress, based on analysis of strains carrying mutations in the genes described above (16).

To assess whether phosphate transport *per se* contributes to osmotic survival, we analyzed wild

type and *pstS::kan* mutant growth and survival at 1% and 1.5% NaCl, which approximate

concentrations within the chicken gastrointestinal tract (22). The *pstS::kan* mutant strain

demonstrated a significant growth defect (**Fig. 3E)** and lower plating efficiency than wild type

on 1% and 1.5% NaCl **(Fig. 3F and G).** The mutant strain was restored to wild type levels of

plating by complementing with *pstSCAB* in trans, confirming that loss of viability at elevated

osmolarity is due to loss of phosphate transport. Thus, similar to mutant strains lacking *ppx1* and

ppx2 (exo-polyphosphatases) (18), which are unable to degrade polyphosphate, the phosphate

transporter mutant survives poorly under osmotic stress conditions that are similar to those found

in the chicken gastrointestinal tract.

Loss of PstSCAB compromises *C. jejuni* chicken colonization and fitness   We investigated

whether elevated abundance of *pstS* mRNA in *C. jejuni* harvested from chick ceca, compared to

C. jejuni grown *in vitro* (5), correlates to actual requirement for *pstSCAB* during growth in the

chicken. We orally inoculated day-of-hatch chickens with two different doses (10^3 or 10^6 CFU)

of wild type *C. jejuni* 81-176, the *pstS::kan* mutant derivative, and the complemented mutant,

and enumerated *C. jejuni* loads in cecal contents on days three and seven after inoculation.

Irrespective of the inoculum dose, the mutant strain exhibited a large colonization defect at both

days three and seven compared with wild type **(Fig 4A,B,C and D)**, being recovered

approximately five orders of magnitude less than wild type when introduced at the lower

inoculum and by roughly two orders of magnitude when introduced at the higher inoculum. In all

cases the complemented strain colonized at levels similar to wild type.

In addition to individual colonization assays with wild type and *pstSCAB* mutant strains,

we carried out a competition assay between the wild type strain 81-176 and the *pstSCAB*

derivative, mixing the strains at 1:1 concentration in the inoculum and introducing the

combination at 10^6 CFU/ml by oral gavage to day-of-hatch chickens. We harvested cecal

contents on day seven post-infection and determined the ratios of the mutant and wild type

strains by measuring relative numbers of kanamycin resistant (*pstSCAB* mutant) and sensitive

(wild type) *C. jejuni* in the output **(Fig 4E)**. The wild type strain exhibited greater fitness in this

experiment than the mutant by 100-fold. The wild type strain also out-competed the mutant when

grown together in nutrient-rich media or in minimal media supplemented with 80 uM phosphate

(Fig. S4A and B). These findings indicate the essential nature of *pstSCAB* for chick colonization

and extend the earlier observation of high-level *in vivo* expression of this operon in the

transcriptome study (5).

The phosphate transporter is essential for lactate-dependent growth.   Lactate is present in

both D- and L- isomers at varying concentrations in the chicken gastrointestinal tract, generally

decreasing from the upper to the lower intestine and ceca (23). Although at high concentrations

lactate inhibits growth of *C. jejuni* (24) and reduces expression of some colonization traits (23),

many strains - including 81-176 - have pathways for uptake of and growth in lactate, including a

transporter for L-lactate encoded by *lctP* (CJJ81176_0113) and two NAD-independent

membrane-associated L-lactate dehydrogenase complexes (CJJ81176_0112-0110) and

(CJJ81176_1182), which convert L-lactate to pyruvate (25). Mechanisms for uptake and

assimilation of D-lactate are less understood (25).

Given the reduced fitness of the *pstSCAB* mutant in the cecum, we investigated the

growth of the mutant on this potentially relevant carbon source in that environment. We cultured

wild type, mutant, and complemented mutant strains on minimal media containing D- + L-lactate

(10 mM total lactate) in both low (80 micromolar) and high (1.6 millimolar) phosphate

concentrations. In both phosphate concentrations, the mutant grew to levels 100-1000 times

lower than wild type at 24 and 48 hours of growth (**Fig. 5A and B**). This contrasts to cells

cultured in media with pyruvate as the carbon source, where mutant and wild type grew to

similar levels in high phosphate media (1.6 mM) and differed by only an order of magnitude at

the lower phosphate concentration (80 µM; **Fig. S5A-B)**. When cultured on L-lactate alone, the

pstS::kan mutant strain was significantly limited compared to wild type, irrespective of

phosphate concentration, while on D-lactate alone, the mutant was diminished for growth during

initial exponential growth phase (irrespective of phosphate levels) but ultimately attained the

same level of growth as the wild type and the complemented mutant (**Fig. 6A-D).** Collectively

these data suggest that the phosphate transport mutant is challenged for growth on L-lactate in

particular; even when both D- and L- lactate are available *in vitro* the mutant is at a significant

disadvantage compared to wild type (**Fig. 5**). Given that D- and L-lactate may serve as growth

substrates for *C. jejuni* growth in the chick cecum (25), these data also contribute to explaining

the fitness disadvantage of the mutant *in vivo*.

Lactate-derived pyruvate is converted to acetate in *C. jejuni* via phosphotransacetylase

(Pta) and acetate kinase (AckA); this acetogenesis pathway is critical for establishing chicken

colonization (23). Genes controlled by this acetogenesis pathway may contribute to fitness in the

chick ceca including those encoding γ-glutamyl transferase (Ggt), a component of an amino acid

transporter complex (Peb1c), and a putative di/tri-peptide transporter (Cjj81176_0683) (23). The

higher L-lactate levels in the upper intestine, compared to those of the lower intestine and ceca,

lead to reduced expression of these genes through an undefined regulatory mechanism that

correlates to relatively lower levels of *C. jejuni* in the upper intestine (23). We hypothesized that

the poor growth on lactate of the *pstSCAB* mutant might also correlate with diminished

expression of acetogenesis-regulated colonization traits. To test this, we carried out quantitative

RT-PCR to measure transcript levels of *ggt*, *peb1c* and *Cjj81176_0683*. Expression of each was

reduced by two-to-four-fold in the mutant compared to wild type **(Fig 7).** This reduction in

expression is similar in magnitude to their reduced expression in the upper intestine where *C.*

jejuni colonizes poorly (23). Thus, we conclude that in addition to poor growth on L-lactate by

the mutant lacking PstSCAB, the strain also expresses potentially key colonization determinants

at levels apparently insufficient for cecal colonization.

Metabolomic analysis of wild type and *pstSCAB* mutant *C. jejuni.* To explore further the

growth physiology of wild type and *pstSCAB* mutant on lactate, we measured intracellular

metabolites using mass spectrometry. We observed no significant difference in intracellular

lactate or pyruvate between wild type and the *pstS::kan* mutant strain **(Fig S6A)** indicating that

the phosphate transport system has no effect on lactate uptake or its conversion to pyruvate. Of

metabolites we measured, adenine - and its deamination product hypoxanthine - were

significantly lower in the mutant compared with the wild type (**Fig. 8 B and C**), and this was

reflected in diminished ATP levels in the mutant as well. (**Fig. 8 A).** The low intracellular Pi

concentration in the *pstS::kan* mutant (as reflected in elevated alkaline phosphatase activity; **Fig.**

3A), correlated to decreased intracellular ATP levels and might account in part for its poor

fitness. This was confirmed with ATP assays on extracts of cells cultured for 24 hours in

minimal-L-lactate-media demonstrating intracellular ATP concentrations in the *pstS::kan* mutant

significantly below those of wild type, irrespective of phosphate concentration **(Fig 8A)**. *C.*

jejuni does not encode a purine/pyrimidine or nucleotide/nucleoside transporter (26), and indeed

adenine supplementation (1uM or 10uM) of L-lactate media did not restore growth or ATP levels

in the mutant (**Fig. S7).**

Discussion

The *pstSCAB* phosphate transporter locus of *C. jejuni* is among the more significantly expressed transcripts identified by RNASeq from microbes harvested from the chick cecum (5) and during infection of humans (27). A transcriptomic study of *C. jejuni* colonizing chicks using microarray analyses (28) did not report high level expression of *pstSCAB*, although there are significant differences in the way the RNASeq and microarray studies were carried out with regards to chicken inoculation and colonization. The RNASeq study (5) used an inoculum of 10^3 CFU of *C. jejuni*, and harvested cecal contents for RNA preparation after seven days of colonization. In contrast the microarray study used a much higher inoculum of 10^9 CFU and harvested cecal contents after only 12 hours of colonization (28). It is conceivable that these differences in inocula and time of colonization would result in different physiological outcomes in the *C. jejuni* recovered from the cecal contents.

Generally, elevated level of expression of the high-affinity phosphate transporter in bacteria is observed in a low-phosphate environment. For example, transcription of *pstS* in *C.jejun*i is significantly increased at 0.08 mM concentration compared to 1.6 mM (7). We determined phosphate concentrations within the chicken cecum as relatively high, 0.5- 0.8 mM/100mg of cecal content **(Fig. S3)**. Further, previous metabolite studies estimated that the Pi concentration in intestinal lumen of healthy adults are varied between 15-30 mM on a Western diet (29). At these *in vivo* levels of inorganic phosphate, we would not expect elevated expression of *pstSCAB* during colonization in chicks and humans, so it remains unclear what drives this outcome during host colonization. We hypothesize that, although present at high detectable levels, phosphate is nevertheless unavailable to *C. jejuni*, perhaps due to competition from other microbes. We previously demonstrated *in vivo* competition for zinc, resulting in poor

colonization of a *C. jejuni* mutant lacking the *znuA* high-affinity zinc transporter. In that case, the

colonization defect of the *znuA* mutant was mitigated in chicks reared in germ-free conditions

harboring a limited microbiota (30).

The *pstSCAB* mutant of *C. jejuni* showed a significant colonization defect on chicken at

both inocula we tested (10^3 and 10^6 CFU), with the most significant defect at day three after

infection. From these data, we hypothesize that *pstSCAB* enables *C. jejuni* to survive initial

conditions after reaching the chicken gut. We also observed a dose dependent colonization defect

of the mutant at day seven. A previous study reported similar colonization levels between wild

type *C. jejuni* and a mutant lacking *phoRS*, which encodes a regulator of *pstSCAB* expression (6).

That study used a different strain of *C. jejuni*, and the animal data were not shown, so we do not

know what infection dose was used or what time points were assessed post-inoculation. Another

study demonstrated that mutants lacking the alkaline phosphatase PhoX have diminished

capacity for colonizing chickens compared to wild type (21) and that this is a dose-dependent

outcome with somewhat greater colonization defect of the mutant observed after a low dose (10^3)

inoculum by comparison to a higher dose inoculum (10^5).

As a proxy for analyzing fitness of wild type and *pstSCAB* mutant in humans, we

measured growth of WT and *pstSCAB* mutant strain *in vitro* in minimal media with 10% human

fecal extract, in which the phosphate concentration was approximately 500 μM **(Fig. S8A)**. The

mutant strain grew in this medium after 24 hours, but nevertheless grew to levels 10x lower than

wild type and the complemented mutant (**Fig. S8B**). Taken together with previous human

transcriptomic data demonstrating elevated levels of *pst* messages during human infection (27),

we conclude that phosphate transporter of *C. jejuni* is required for fitness in both chickens and

humans.

The *pstSCAB* operon is transcribed as a full-length polycistronic mRNA, demonstrated

both in the earlier RNASeq study and in our qRT-PCR presented in this work. Trancripts

encoded by the first gene in this operon, *pstS,* are present at much higher levels than those of the

three downstream genes (5). We postulate that a potential stem-loop structure encoded just

downstream of the *pstS* gene, which we postulate serves to stabilize *pstS* mRNA from

degradation, resulting in higher steady state levels compared to the other genes of the operon. A

similar mechanism is proposed for the *pstS2* operon of *Vibrio cholerae* (31).

In complex media, PstSCAB is dispensable until late in growth, after which it is required

for cells to remain viable in the nutrient-depleted environment (**Fig.3**). As transcription of the

locus is elevated during mid-logarithmic growth (compared with stationary phase growth (5)) we

conclude that assembly of the system and Pi uptake occurs early during growth and enables

survival later when the locus is far less transcriptionally active. Accumulation of poly-P in the

pstSCAB mutant late in growth may be the result of decreased expression of the exo-

polyphosphatase genes *ppx1* and *ppx2*, which encode enzymes that metabolize poly-P.

Consistent with a need to balance poly-P production and metabolism for wild type fitness in *C.*

jejuni others, have demonstrated that *phoX* and *ppk1/ppk2* mutants also have survival defects in

nutrient limiting condition (18). Increased poly-P was also observed in phosphate transport

mutants of *M. tuberculosis* (32).

The failure of the *pstSCAB* mutant to thrive - and its loss of viability - in 10 mM lactate,

on which wild type grows to high levels, is worth further analysis. The mutant strain exhibits

reduced expression of the *pta-ackA* genes in the acetogenesis pathway, which could be partly

responsible for the decreased ATP levels in the mutant compared to the wild type. Lactate at

high concentrations causes membrane damage to *C. jejuni* and is the basis for the suggestion that

Lactobacillus strains that produce high lactic acid levels might be useful probiotics to reduce *C.*

jejuni loads in poultry (24). Concentrations required for killing *C. jejuni* are much higher than 10

mM, but in the context of loss of *pstSCAB* perhaps the cell is sensitized to lactate toxicity. A

mutant of avian pathogenic *E. coli* lacking *pst* was demonstrated to be more sensitive to

membrane-targeting polymyxin B, but we did not observe that with *pstSCAB* mutant *C. jejuni*.

How *pstSCAB* contributes to lactate-dependent growth will be the subject of future research.

417   **Methods and Materials**

Bacterial Strains and Media All strains and plasmids used in the study are listed in Table 1.

The wild type strain used in all experiments is DRH212, a streptomycin resistant derivative of *C.*

jejuni 81-176 (33). All *C. jejuni* strains were routinely grown under microaerobic conditions

(85% N2,10%CO2, 5% O2) on Muller-Hilton (MH) agar or in broth under necessary antibiotics

at the following concentrations: chloramphenicol, 15 μg ml^{-1}; kanamycin, 50 μg ml^{-1}; TMP, 10

μg ml^{-1}; streptomycin, 2 mg ml^{-1}. *C. jejuni* strains were stored in MH broth with 20% glycerol at

−80°C.

Construction of a *pstS::kan* insertion mutant and complementation *C. jejuni pstS*::*kan*

mutant was constructed by insertion of kanamycin resistance cassette into the gene as previously

described (34). Briefly, the 500bp of both upstream and downstream regions of *pstS* locus were

amplified by PCR with flanking region containing SmaI restriction site. The amplified upstream

and downstream region were joined by Soeing PCR. The 1000bp amplified PCR products was

cloned into pGEM-T easy. The *Campylobacter* kanamycin resistance cassette was excised from

PILL600 using SmaI and ligated between upstream and downstream cloned region of pGEM-T

construct. The plasmid was electroporated into DRH212 strain and grown on MH agar overnight

at 37°C under microaerobic conditions. Then cells were plated and grown on MH agar

containing kanamycin for 2-3 days, and successful integration of *pstS::kan* into the chromosome

was confirmed by PCR. As we demonstrate in Results, this insertion is polar on the *pst* genes

downstream of *pstS*. Therefore, for complementation, the coding sequences of whole *pstSCAB*

operon - from the second codon of *pstS* to the stop codon of *pstB* - were amplified by PCR and

cloned into the BamHI site of pECO102 plasmid (35), which was introduced into *E.*

*coli*DH5α/pRK212.1 strain (35). Conjugation to *C. jejuni* was performed as previously described

by Guerry et al (36).

Growth Curve analysis in MH broth and Minimal media *C. jejuni* strains WT, *pstS::kan*,

pstS::kan/C were grown from freezer stocks on MH agar containing desired antibiotics for 24h

under microaerobic condition. The strains again re-streaked and grown for 16h. The bacteria

were suspended in MH broth and inoculated in 25ml of MH broth containing desired antibiotics.

The initial O.D was approximately 0.05 and bacterial numbers were measured at different time

intervals up to 48h. To determine growth of *C. jejuni* strains in different concentrations of

inorganic phosphate, strains were grown in MCLMAN minimal media containing different

carbon sources including sodium L-pyruvate, D-lactate and L-lactate (37) as noted. Strains were

washed twice with minimal media and inoculated with 25ml of minimal containing different

concentration of Pi. Then, the growth was measured via absorbance at 600nm (OD_{600}) up to 48h

with the initial O.D was 0.05.

ATP measurements ATP levels in strains were determined by BacTitre-Glo^{TM} Assay Kit

(Promega) as previous described. Briefly, strains were grown for 24h in minimal media (L-

lactate) with high (1.6mM) and lower concentration (0.08mM) of Pi and normalized OD at 0.3.

Then ATP level were measured according to the manufacture's protocol and ATP concentrations

were measure by standard curve (26).

Measurement of intracellular metabolites Wild type strain DRH212, *pstS::kan* were grown in

presence of L-lactate with higher and lower concentration of phosphate for 18h. The cells were

collected by centrifugation and washed twice with normal saline. Then all samples were

normalized approximately with OD. 1. Then the samples were extracted for determination of

metabolites. Samples on dry ice were combined with 800 microliters of ice-cold

acetonitrile:methanol:acetone (8:1:1, v:v:v) and suitable stable isotope-labeled or synthetic

internal standards including D4-glycochenodeoxycholic acid and myristoyl phosphatidylcholine

for estimation of metabolite recovery and for relative quantitation across experimental groups.

Samples were homogenized with zirconium oxide beads for 2 minutes in a Bullet Blender (Next

Advance) at 4 degrees Celsius, incubated for 30 minutes at 4 degrees Celsius, then centrifuged

for ten minutes at 10,000g at 4 degrees Celsius. Supernatants were filtered through 0.2 micron

syringe filters (Fisher Scientific), and evaporated to dryness in a SpeedVac. Samples were

resuspended in 95% acetonitrile for use in LC-MS analysis.

Liquid chromatography-mass spectrometry Untargeted metabolite identification utilized a

Thermo model LTQ-Orbitrap Velos mass spectrometer operating in full scan negative ionization

MS mode at 60,000 resolution with data-dependent HCD-MS/MS over a mass range of 50-1000

m/z. The mass spectrometer was coupled to a Shimadzu Prominence HPLC system through an

electrospray ionization source. Ten microliters of each metabolite extract was injected and

separated on a Phenomenex HILIC LC column, 2.0 mm x 100 mm, 3.0 micron particle size, 100

Angstrom pore size, using a gradient of (A) water containing 50 mM ammonium formate, and

(B) acetonitrile from 95% to 50% B over 14 minutes as previously descrived. (38). A guard

cartridge of matching chemistry was fitted to the analytical column.

Data analysis Compound identifications, isotope correction, chromatographic peak alignment,

and peak area quantification were performed with MAVEN software (39). Due to the need to

simultaneously analyze numerous disparate classes of metabolites in untargeted experiments,

only relative quantitation of analytes against a selected internal standard were performed for

comparison of values across experimental treatment groups. Multivariate statistical analysis was

performed using MetaboAnalyst version 4.0 software (https://www.metaboanalyst.ca/).

Nutrient downshift survival *C. jejuni* were grown overnight at 37°C in MH broth under

microaerobic condition. Then strains were collected by centrifugation at 6,000 rpm for 10 min

and washed twice with MCLMAN minimum media where no carbon and no phosphate present.

Strains were diluted to an OD600 of 0.05. After different time intervals, CFU were measured by

plating on MH agar plates.

Osmotic stress survival *C. jejuni* strains were grown overnight at 37°C in MH broth under

microaerobic condition. Then strains were centrifuged and resuspended in MH media containing

1% and 1.5% NaCl. The initial OD of the culture was 0.05 and survival was measured by CFU

count at different time intervals (22).

Real Time PCR. Wild type strain DRH212, *pstS::kan* strains were grown in different condition

in vitro and total RNA was isolated by the Trizol method and RNeasy Mini Kit (Qiagen). After

digestion with TURBO Dnase treatment, cDNA was made by SuperScript III First-Strand

Synthesis Supermix (Invitrogen). Then real time PCR was performed by Syber Green method.

The gene expression of different targeted genes was measured by $2^{\text{delta deltaCT}}$ method.

Measurement of *in vivo* poly-P levels. Intracellular poly-P levels were measured as described

(40) Briefly, strains were grown in MH broth for 24h and 48h. Strains were collected in each

time point by centrifugation and dissolved in 250 µL of GITC lysis buffer (4 M guanidine

isothiocyanate, 50 mM Tris-HCl, pH 7) and lysed by incubating at 95 °C for 10 min. Lysates

were stored at -80 °C. The protein concentration was determined by Bradford assay (Bio-Rad) of

a 5-µl aliquot. Poly-P was isolated using EconoSpin silica spin columns (Epoch Life Science)

and digested with 1 µg of PPX1 exopolyphosphatase from *Saccharomyces cerevisiae*. The

resulting free phosphate was measured using a malachite green colorimetric assay and

normalized to the total protein (40).

Chicken colonization assay and *in vivo* phosphate determination   Day-of-hatch white

Leghorn chickens were inoculated orally with two doses (approximately $1\text{x}10^3$ and $1\text{x}10^6$

CFU/ml) of indicated *C. jejuni* strains (wild type strain 81-176, *pstS::kan* and *pstS::kan/C*) (8).

Birds were euthanized at day 3 and day 7 after infection, and cecal contents were collected as

described (41). Cecal contents were homogenized in sterile PBS and serial dilutions were plated

on Mueller-Hinton (MH) agar containing 10% sheep's blood, cefoperazone (40 µg/ml),

cycloheximide (100 µg/ml), trimethoprim (10 µg/ml), and vancomycin (100 µg/ml), selective for

Campylobacter. Plates were grown for 48 h under microaerobic conditions and colonies were

enumerated. For the competition experiment, wild type and *pstS::kan* strain were mixed equally

(approximately $1\text{x}10^6$ CFU/ml of each strain) and inoculated the day-of-hatch chicks by oral

gavage. At Day 7, cecal contents were collected and after serial dilution plated on

Campylobacter selective media with and without kanamycin (50 µg/ml). Competitive index

values were calculated as previously described (42).

Inorganic free phosphate was measured in chicken cecal content (100mg) at Day 3 and

Day 7 by inorganic free phosphate measurement ELISA Kit (Bioassay) as per manufacturer

protocol. All chicken experiments and protocols were approved by the Institutional Animal Care

and Use Committee in the Animal Care Program at Michigan State University.

Acknowledgements

We thank Dr. Shannon Manning, Department of Microbiology and Molecular Genetics Michigan

State University for providing human stool samples used in this study. This work was supported

in part by NIH award AI 111192 (to V.J.D.) and by the Rudolph Hugh Endowment of Michigan

State University.

Table 1:
533

Strain or plasmid	Relevant characteristics	Source of Reference
C. jejuni DRH212	Streptomycin-resistant wild type *C. jejuni* 81-176 (Sm^r)	34
DRH212 *pstS::kan*	*pstS* insertion mutant of *C. jejuni* DRH212 (Kn^r Sm^r)	In this study
DRH212 *pstS::kan/C*	*pstS* insertion mutant of *C. jejuni* DRH212 content *pstSCAB* cloned PECO212 ((Kn^r Sm^r Cm^r)	In this study
E.coli DH5a	*E. coli* strain used for cloning	
E.coli DH5a/pRK212.1	Contains conjugative plasmid for the conjugation of plasmid DNA into *Campylobacter*	36
Plasmids		
pGEMT	Subcloning vector (Ap^r)	
pECO102	pRY112 derivative with *cat* promoter in XhoI-BamHI	36
pILL600	contains *Campylobacter* kanamycin cassette	36

References

1. Burnham PM, Hendrixson DR. 2018. Campylobacter jejuni: collective components promoting a successful enteric lifestyle. Nat Rev Microbiol 16:551–565.
2. Skarp CPA, Hänninen M-L, Rautelin HIK. 2016. Campylobacteriosis: the role of poultry meat. Clin Microbiol Infect 22:103–109.
3. Marotta F, Garofolo G, Di Donato G, Aprea G, Platone I, Cianciavicchia S, Alessiani A, Di Giannatale E. 2015. Population Diversity of Campylobacter jejuni in Poultry and Its Dynamic of Contamination in Chicken Meat. Biomed Res Int 2015:859845.
4. Zeng L, Ruan M, Liu J, Wilde P, Naumova EN, Mozaffarian D, Zhang FF. 2019. Trends in Processed Meat, Unprocessed Red Meat, Poultry, and Fish Consumption in the United States, 1999-2016. J Acad Nutr Diet 119:1085-1098.e12.
5. Taveirne ME, Theriot CM, Livny J, DiRita VJ. 2013. The complete Campylobacter jejuni transcriptome during colonization of a natural host determined by RNAseq. PLoS One 8:e73586.
6. Lamarche MG, Wanner BL, Crépin S, Harel J. 2008. The phosphate regulon and bacterial virulence: a regulatory network connecting phosphate homeostasis and pathogenesis. FEMS Microbiol Rev 32:461–73.
7. Wösten MMSM, Parker CT, van Mourik A, Guilhabert MR, van Dijk L, van Putten JPM. 2006. The Campylobacter jejuni PhosS/PhosR operon represents a non-classical phosphate-sensitive two-component system. Mol Microbiol 62:278–91.
8. Vuppada RK, Hansen CR, Strickland KAP, Kelly KM, McCleary WR. 2018. Phosphate signaling through alternate conformations of the PstSCAB phosphate transporter. BMC Microbiol 18:8.
9. Chekabab SM, Harel J, Dozois CM. 2014. Interplay between genetic regulation of phosphate homeostasis and bacterial virulence. Virulence 5:786–93.
10. Lamarche MG, Dozois CM, Daigle F, Caza M, Curtiss R, Dubreuil JD, Harel J. 2005. Inactivation of the pst system reduces the virulence of an avian pathogenic Escherichia coli O78 strain. Infect Immun 73:4138–45.
11. Chekabab SM, Jubelin G, Dozois CM, Harel J. 2014. PhoB activates Escherichia coli O157:H7 virulence factors in response to inorganic phosphate limitation. PLoS One 9:e94285.
12. Tischler AD, Leistikow RL, Kirksey MA, Voskuil MI, McKinney JD. 2013. Mycobacterium tuberculosis requires phosphate-responsive gene regulation to resist host immunity. Infect Immun 81:317–28.
13. Rengarajan J, Bloom BR, Rubin EJ. 2005. Genome-wide requirements for Mycobacterium tuberculosis adaptation and survival in macrophages. Proc Natl Acad Sci U S A 102:8327–32.
14. Dar D, Sorek R. 2018. Extensive reshaping of bacterial operons by programmed mRNA decay. PLoS Genet 14:e1007354.
15. Kumar A, Gangaiah D, Torrelles JB, Rajashekara G. 2016. Polyphosphate and associated enzymes as global regulators of stress response and virulence in Campylobacter jejuni. World J Gastroenterol 22:7402–14.
16. Candon HL, Allan BJ, Fraley CD, Gaynor EC. 2007. Polyphosphate kinase 1 is a pathogenesis determinant in Campylobacter jejuni. J Bacteriol 189:8099–108.
17. Gangaiah D, Liu Z, Arcos J, Kassem II, Sanad Y, Torrelles JB, Rajashekara G. 2010.

581   Polyphosphate kinase 2: a novel determinant of stress responses and pathogenesis in
582   Campylobacter jejuni. PLoS One 5:e12142.

18.   Malde A, Gangaiah D, Chandrashekhar K, Pina-Mimbela R, Torrelles JB, Rajashekara G.
2014. Functional characterization of exopolyphosphatase/guanosine pentaphosphate
phosphohydrolase (PPX/GPPA) of Campylobacter jejuni. Virulence 5:521–33.

19.   Gaynor EC, Wells DH, MacKichan JK, Falkow S. 2005. The Campylobacter jejuni
stringent response controls specific stress survival and virulence-associated phenotypes.
Mol Microbiol 56:8–27.

20.   Peng Y-C, Lu C, Li G, Eichenbaum Z, Lu C-D. 2017. Induction of the pho regulon and
polyphosphate synthesis against spermine stress in Pseudomonas aeruginosa. Mol
Microbiol 104:1037–1051.

21.   Drozd M, Gangaiah D, Liu Z, Rajashekara G. 2011. Contribution of TAT system
translocated PhoX to Campylobacter jejuni phosphate metabolism and resilience to
environmental stresses. PLoS One 6:e26336.

22.   Cameron A, Frirdich E, Huynh S, Parker CT, Gaynor EC. 2012. Hyperosmotic stress
response of Campylobacter jejuni. J Bacteriol 194:6116–30.

23.   Luethy PM, Huynh S, Ribardo DA, Winter SE, Parker CT, Hendrixson DR. 2017.
Microbiota-Derived Short-Chain Fatty Acids Modulate Expression of Campylobacter
jejuni Determinants Required for Commensalism and Virulence. MBio 8.

24.   Neal-McKinney JM, Lu X, Duong T, Larson CL, Call DR, Shah DH, Konkel ME. 2012.
Production of organic acids by probiotic lactobacilli can be used to reduce pathogen load
in poultry. PLoS One 7:e43928.

25.   Thomas MT, Shepherd M, Poole RK, van Vliet AHM, Kelly DJ, Pearson BM. 2011. Two
respiratory enzyme systems in Campylobacter jejuni NCTC 11168 contribute to growth
on L-lactate. Environ Microbiol 13:48–61.

26.   Cameron A, Huynh S, Scott NE, Frirdich E, Apel D, Foster LJ, Parker CT, Gaynor EC.
2015. High-Frequency Variation of Purine Biosynthesis Genes Is a Mechanism of Success
in Campylobacter jejuni. MBio 6:e00612-15.

27.   Crofts AA, Poly FM, Ewing CP, Kuroiwa JM, Rimmer JE, Harro C, Sack D, Talaat KR,
Porter CK, Gutierrez RL, DeNearing B, Brubaker J, Laird RM, Maue AC, Jaep K, Alcala
A, Tribble DR, Riddle MS, Ramakrishnan A, McCoy AJ, Davies BW, Guerry P, Trent
MS. 2018. Campylobacter jejuni transcriptional and genetic adaptation during human
infection. Nat Microbiol 3:494–502.

28.   Woodall CA, Jones MA, Barrow PA, Hinds J, Marsden GL, Kelly DJ, Dorrell N, Wren
BW, Maskell DJ. 2005. Campylobacter jejuni gene expression in the chick cecum:
evidence for adaptation to a low-oxygen environment. Infect Immun 73:5278–85.

29.   Walton J, Gray TK. 1979. Absorption of inorganic phosphate in the human small
intestine. Clin Sci (Lond) 56:407–12.

30.   Gielda LM, DiRita VJ. 2012. Zinc competition among the intestinal microbiota. MBio
3:e00171-12.

31.   da C Leite DM, Barbosa LC, Mantuano N, Goulart CL, Veríssimo da Costa GC, Bisch
PM, von Krüger WMA. 2017. Transcriptional and post-transcriptional regulation of pst2
operon expression in Vibrio cholerae O1. Infect Genet Evol 51:10–16.

32.   Namugenyi SB, Aagesen AM, Elliott SR, Tischler AD. 2017. Mycobacterium tuberculosis
PhoY Proteins Promote Persister Formation by Mediating Pst/SenX3-RegX3 Phosphate
Sensing. MBio 8.

33. Hendrixson DR, Akerley BJ, DiRita VJ. 2001. Transposon mutagenesis of Campylobacter
jejuni identifies a bipartite energy taxis system required for motility. Mol Microbiol
40:214–24.
34. Johnson JG, Livny J, Dirita VJ. 2014. High-throughput sequencing of Campylobacter
jejuni insertion mutant libraries reveals mapA as a fitness factor for chicken colonization.
J Bacteriol 196:1958–67.
35. Wiesner RS, Hendrixson DR, DiRita VJ. 2003. Natural transformation of Campylobacter
jejuni requires components of a type II secretion system. J Bacteriol 185:5408–18.
36. Guerry P, Yao R, Alm RA, Burr DH, Trust TJ. 1994. Systems of experimental genetics
for Campylobacter species. Methods Enzymol 235:474–81.
37. Alazzam B, Bonnassie-Rouxin S, Dufour V, Ermel G. MCLMAN, a new minimal
medium for Campylobacter jejuni NCTC 11168. Res Microbiol 162:173–9.
38. Cai X, Li R. 2016. Concurrent profiling of polar metabolites and lipids in human plasma
using HILIC-FTMS. Sci Rep 6:36490.
39. Clasquin MF, Melamud E, Rabinowitz JD. 2012. LC-MS Data Processing with MAVEN:
A Metabolomic Analysis and Visualization EngineCurrent Protocols in Bioinformatics.
John Wiley & Sons, Inc., Hoboken, NJ, USA.
40. Pokhrel A, Lingo JC, Wolschendorf F, Gray MJ. 2019. Assaying for Inorganic
Polyphosphate in Bacteria. J Vis Exp.
41. Hendrixson DR, DiRita VJ. 2004. Identification of Campylobacter jejuni genes involved
in commensal colonization of the chick gastrointestinal tract. Mol Microbiol 52:471–84.
42. Johnson JG, Gaddy JA, DiRita VJ. 2016. The PAS Domain-Containing Protein HeuR
Regulates Heme Uptake in Campylobacter jejuni. MBio 7.
650
651

Fig 1:

Fig 1: Characterization of *pstSCAB* operon: A) RT-PCR was performed to determine whether

the four genes (*pstS, pstC, pstA* and *pstB*) are cotranscribed. Three intergenic primer sets (P1, P2

and P3) were designed to amplify transcripts crossing the gene boundaries. The product sizes in

bp were correct as predicted in all cases. RT PCR with total RNAs were extracted from both

DRH212 (left panel) and *pstS::kan* mutant (right panel) using the designed three primer sets. RT

- negative control; 16S - as a positive control.

677
678
Fig 2: **Comparative analysis of Poly-P metabolism regulatory genes and Poly-P level between DRH212 and *pstS::kan* mutant: A)** and **B)** Transcription analysis of different stress regulatory genes *ppk1, ppk2, spoT* and *ppA* at 24h and 48h after growth of DRH212 and *pstS::kan* by Quantitative RT PCR. Changes of gene expression in the *pstS::kan* mutant compared with wild type DRH212 were determined by the $2^{-\Delta\Delta CT}$ method. Data are represented as mean value of three independent experiments with standard deviation (±). **C)** and **D)** Levels of Intracellular polyphosphate (Poly-P) of three strains (DRH212, *pstS::kan, pstS::kan/C*) were measured at two different time intervals 24h and 48h. Statistical analysis were done using Unpaired t-test and One way ANOVA with Holm-Sideak's multiple test; (n=8,±SD), ∗ ,P<0.05; ∗∗,$P < 0.005$.

689
690

Fig 3:

Fig 3: Characterization of *C. jejuni pstS::kan* strain in different *in vitro* conditions:

A) Alkaline phosphatase activity was determined in pyruvate-minimal media with 1.6 mM or 0.08 mM Pi (n=3). Statistical analysis was done using two way ANOVA with Sidak's multiple comparisons test. $*$,P<0.05; $**$, $P < 0.005$, **B)** *C. jejuni* strains were grown in Mueller-Hilton (MH) broth in microaerobic condition for 48 h and after different time intervals CFU/ml were determined. **C)** *C. jejuni* DRH212 was grown in MH broth for 48h. After 48h, sterile spent media

were collected by filter sterilization. Mid-log phase cultures of DRH212, *pstS::kan, pstS::kan/C* strains were inoculated (10^8 CFU/ml) in spent media and survival was measured at different time intervals by CFU count. **D)** In nutrient down shift assay, mid-log phase of DRH212, *pstS::kan, pstS::kan/C* strains were inoculated in minimal media where no carbon and phosphate source were present. The survival assay was measured by CFU count at different time intervals. **E)** Osmotic stress tolerance: strains were inoculated in MH broth containing 1% NaCl (**F**) or 1.5% NaCl (**G**) and growth was determined by measuring O.D at 600nm and cfu counts. Each experiment represents mean value of three independent experiments. Statistical significant was done student t test. $*$,P<0.05; $**, P < 0.005$.

Fig.4:

Fig 4: Colonization ability of *pstS::kan* mutant in chickens: Day-of-hatch white Leghorn chicks were infected with either DRH212, *pstS::kan* or *pstS::kan/C* strains of *C. jejuni* (two different doses 10^3 and 10^6 CFU/ml) and cecal loads were enumerated at day three and day seven. Comparative colonization ability between DRH212 and *pstS::kan* strains at day three and day seven after oral inoculation with 10^3 (**A, B**) and 10^6 CFU/ml (**C, D).** Statistical analysis was one way ANOVA with Tukey's multiple comparison test (n=5,±SD). **E)** In competition analysis,

DRH212 versus *pstS::kan* mutant, day-of-hatch chicks were infected with equal ratio of

DRH212 and *pstS::kan* mutant. At day seven the ratio of *pstS::kan* mutant to DRH212 was

determined and is presented as a competitive index. Statistical analysis was performed using a

one-simple t test against a hypothetical value of 1. (n=5 of each group). $*$,P<0.05; $**$,$P <$

0.005.

Fig 5.

Fig 5: PstSCAB transporter of *C. jejuni* is essential for growth in lactate: *C. jejuni* strains

were grown in minimal media containing both D+L lactate (each 5mM) with two concentrations

of inorganic phosphate. Growth of DRH212 and *pstS::kan* was measured by CFU count in

minimal media (D+L lactate) with **A)** 1.6mM Pi and **B)** 0.08mM Pi at 24 and 48h. N=3; error bar

indicated SD. Statistical analysis was done by two-way analysis of variation (ANOVAs) with

Sidak's multiple comparisions test. $*$,P<0.05; $**$,$P < 0.005$**.**

Fig 6:

Fig 6: Comparative growth analysis of *C. jejuni* strains in minimal media with D- or L-

lactate. *C. jejuni* strains were grown in minimal media with D-lactate or L-lactate as a carbon

source and high or low concentration of inorganic phosphate. Growth of *C. jejuni* strains was

measured in D-lactate minimal medium with **A)** 1.6mM and **B)** 0.08mM phosphate by CFU

counts at 24h and 48h. Growth of these three strains was also determined in L-lactate minimal

medium with 1.6mM **C)** and 0.08mM **D)** phosphate. N=3; error bar indicated SD. Statistical

analysis was done by two-way analysis of variation (ANOVAs) with Sidak's multiple

comparisons test. $*$,P<0.05; $**$,$P < 0.005$.

Fig:7

Fig 7: Transcript analysis of acetogenesis regulatory and acetogenesis dependent gene

expression in wild type, ***pstS::kan*****, and complemented strains**: Bacterial strains were grown in

minimal media containing L-lactate with 1.6mM Pi up to 18h, after which cells were harvested

for RNA isolation. Acetogenesis regulatory genes (*ackA* and *pta*) and acetogenesis dependent

genes (*ggt,peb1C* and *CJJ0683*) were measured in *pstS::kan* mutant and *pstS::kan/C* strains

relative to DRH212 by real time PCR. N=3; error bar indicated SD.

 Fig 8:

Fig 8: Intracellular ATP and metabolites in DRH212 and *pstS::kan* mutant grown in L-lactate minimal media. DRH212 and *pstS::kan* strains were grown in L-lactate minimal media with 1.6mM and 0.08mM Pi at 37^0 C. After 18h, **A)** ATP levels in DRH212 , *pstS::kan* were detetermined. N=3; error bar indicated SD. The intracellular concentration of **B)** adenine and **C)** hypoxanthine were measured by liquid chromatography mass spectroscopy analysis. Data represent mean value of two independent experiments. The statistical analysis was done by one way ANOVA. $*$,P<0.05; $**$, $P < 0.005$**.**

784
785
786