**Mosaic metabolic ageing: Basal and standard metabolic rate age in opposite directions**

**and independent of environmental quality, sex or lifespan in a passerine**

Michael Briga[1,2] & Simon Verhulst[1]

[1] Groningen Institute for Evolutionary Life Sciences, University of Groningen, 9747 AG

Groningen, The Netherlands.

[2] Present address: Department of Biology, University of Turku, Turku, Finland.

**Abstract**

1. Crucial to our understanding of the ageing process is identifying how traits change with

age, which variables alter their ageing process and whether these traits associate with

lifespan.

2. We here investigated metabolic ageing in zebra finches. We longitudinally monitored 407

individuals during six years and collected 3213 measurements of two independent mass-

adjusted metabolic traits: basal metabolic rate ($BMR_m$) at thermoneutral temperatures

and standard metabolic rate ($SMR_m$), which is the same as $BMR_m$ but at ambient

temperatures below thermoneutrality.

3. $BMR_m$ *decreased* linearly with age, consistent with earlier reports. In contrast, $SMR_m$

*increased* linearly with age. To the best of our knowledge, this is the first quantification of

$SMR_m$ ageing, and thereby of the contrast between $SMR_m$ and $BMR_m$ ageing.

4. Neither metabolic rate nor metabolic ageing rate were associated with individual lifespan.

Moreover, experimental manipulations of environmental quality that decreased $BMR_m$

and $SMR_m$ and shortened lifespan with 6 months (12%) did not affect the ageing of either

metabolic trait. Females lived 2 months (4%) shorter than males, but none of the

metabolic traits showed sex-specific differences at any age.

5. Our finding that ageing patterns of metabolic rate vary depending on the ambient

temperature illustrates the importance of studying ageing in an ecologically realistic

setting.

6. Our results add to the mounting evidence that within an organism ageing is an

asynchronous process.

## 1. Introduction

31
Senescence, the decline in fitness due to declining organismal functioning with age, is common in
humans, model organisms and in the wild (Nussey et al. 2013; Belsky et al. 2015; Fontana and
Partridge 2015). Ageing is the change in traits with age, irrespective of their association with
residual reproductive value; for example, human hair may turn grey with age but this by itself
does not affect remaining lifespan. Major challenges in ageing research are to identify how
organisms change with age, the factors affecting those changes and in particular how those
changes affect fitness and remaining lifespan (Christensen et al. 2010; Nussey et al. 2013; Gaillard
and Lemaitre 2017; Partridge and Deelen 2018).

Energy metabolism is an essential resource underlying organismal functioning, reproduction and
survival and is therefore a key variable in physiology and ecology, including in the study of ageing
(Drent and Daan 1980; Burton et al. 2011; Norin and Metcalfe 2019). For example, consistent
individual differences in energy turnover can be a key determinant of behavior, reproduction,
survival (Biro and Stamps 2010; Burton et al. 2011). Energy metabolism can be quantified in
several ways. A tractable component of energy turnover that is often quantified is basal metabolic
rate (BMR), i.e. the minimum energy expenditure of a post-absorptive adult animal measured
during the rest phase at thermoneutral temperature (IUPS Thermal Commission, 2001; McNab,
1997). How BMR changes with age within indiviuals has been quantified in a variety of species
and environments, including in humans, and all studies consistently reported a decline with age
(Elliott et al. 2015). Thus BMR, an individual's minimum energy expenditure at thermoneutrality,
declines with age.

While BMR is measured in the thermoneutral zone, most endotherms spend much of their lives at
ambient temperatures that are below thermoneutrality. Standard metabolic rate (SMR), which is
the same as BMR, except that Ta is below thermoneutrality, may therefore be a metabolic trait
with higher ecological relevance, in the sense that it better reflects metabolic rate in natural
conditions. It is often implicitly assumed that BMR and SMR vary in parallel across individuals,
but we previously showed that BMR and SMR in the zebra finch *Taeniopygia guttata* correlated
poorly both between and within-individuals, despite substantial repeatabilities (Briga and
Verhulst 2017). How SMR changes with age in endotherms has to our best knowledge not
previously been investigated, but our earlier findings show that we cannot assume that it changes
with age in a similar way as BMR.

We monitored the ageing of BMR and SMR in a captive population of a small passerine, the zebra
finch. The birds used in this study lived in captivity, which may alter metabolism and possibly its
association with age compared with to that of free-living animals (Wiersma and Verhulst 2005;
Auer et al. 2016; Briga and Verhulst 2017). Indeed, an essential difference between captive and

free-living individuals is that free-living animals generally face foraging costs, which can be a key

parameter in ageing (Briga and Verhulst 2015$a$; Speakman et al. 2015). Foraging costs could drive

or reinforce many age associated declines in organismal performance in the wild, although this

was rarely studied (reviewed in: Clay et al. 2018). Hence, to broaden the range of environments

and increase the ecological relevance of our study, we housed the birds in outdoor aviaries, and

permanently exposed half of our population to high foraging costs using a manipulation of flight

cost per food reward (Koetsier and Verhulst, 2011). We applied a 2x2 design, independently

manipulating foraging costs during development and in adulthood. This manipulation shortened

lifespan and decreased body mass, $BMR_m$ and $SMR_m$ in our experimental population (Briga et al.,

2017; Briga and Verhulst, 2017; Briga et al. 2019). In our experiment, individuals experiencing

high foraging costs during development and in adulthood (HH group) lived on average six months

(12%) shorter than all other groups. Females also lived two months shorter than males (4%).

Given these findings, it was of interest to investigate whether lifespan variation between and

within experimental treatment groups and sexes was associated with ageing rate. If we assume

that factors that shorten lifespan also accelerate ageing, a common assumption in ageing research

(Williams 1999), we predict that the shortest living experimental group shows the highest rate of

metabolic ageing.

## 2. Material & methods

### 2.1 Experimental setup

Birds were reared in either experimentally small broods (with 2 or 3 chicks) or large broods (between 5 and 8 chicks). These brood sizes are within the range observed in wild (Zann 1996). Growing up in large broods increases costly begging beg more, receive less food and have impaired growth (Briga, 2016; Briga et al., 2017). After nutritional independence and before the start of the foraging cost manipulation, i.e. between 35 days till approximately 120 days, young were housed in larger indoor cages with up to 40 other young of the same sex and two male and two female adults. Once adult, birds were subject to a long-term foraging experiment, (Koetsier and Verhulst 2011). Briefly, birds were housed in eight single sex outdoor aviaries (LxHxW 310x210x150 cm) located in Groningen, the Netherlands (53° 13' 0" N / 6° 33' 0" E). Food (tropical seed mixture) water, grit and cuttlebone were provided *ad libitum* and the birds received fortified canary food (''egg food'', by Bogena, Hedel, the Netherlands) in weighed portions, which is consistent with the conditions in other facilities (Griffith et al. 2017). Each aviary contained an approximately equal number of birds and to keep densities within aviaries within a limited range, new birds were added regularly to replace those that died. The first batch was 3-24 months old when the experiment started and birds added later were 3 to 4 months old.

### 2.2 Data collection

Between December 2007 and April 2013 we collected a total of 3213 respirometry measurements on 407 individuals. Of those, 1233 measurements on 386 individuals are BMR measurements (Fig. S1). In the thermoneutral zone an organism does not increase its metabolic rate in order to maintain body temperature and for the zebra finch this zone occurs at ambient temperatures between 32°C and 39°C (Briga and Verhulst, 2017; Calder, 1964). The other 1980 measurements on 372 individuals are of standard metabolic rate (Fig. S1), which we measured in the same way as BMR, except that the ambient temperature was below thermoneutrality, between 5°C and 32°C. The vast majority of these measurements were at temperatures between 23 and 29 (Fig. S2). BMR and SMR data were collected over an age range from 0.4 months till 7.2 years (Fig. S2), in the same seasons, i.e. mostly in spring and autumn (Fig. S3) and measurements were randomized across sex and experimental treatments.

Metabolic rate was measured overnight using an open flow respirometer situated in a dark acclimatized room. Metabolic rate measurements started close to sunset (mean = 18:10 h; SD 01:17). Up to sixteen individuals were taken from the aviaries and randomly transferred to one of sixteen 1.5 L metabolic chambers. Neither food nor water was available for birds during the metabolic measurements. Birds were weighed before and after each measurement and we used

the mean value to capture mass corrected BMR or SMR. Technical details about the equipment

can be found in Bouwhuis et al. (2011). In brief, the air-flow through the metabolic chambers was

controlled at 25 l h$^{-1}$ by mass-flow controllers (5850S; Brooks, Rijswijk, the Netherlands)

calibrated with a bubble flow meter. Air was dried using a molecular sieve (3 Å ; Merck, Darmstadt,

Germany) and analyzed by a paramagnetic oxygen analyzer (Servomex Xentra 4100,

Crowborough, UK). During measurements each metabolic chamber or reference outdoor air was

sampled every 8 min for 60s to stabilize measurement levels (Bouwhuis et al. 2011). In each

sampling, we measured $O_2$ and $CO_2$ concentration and oxygen consumption was calculated using

Eq. (6) of Hill (1972). An energy equivalent of 19.7 kJ l$^{-1}$ oxygen consumed was used to calculate

energy expenditure in watt (W). Metabolic rate was taken to be the minimum value of a 30-min

running average, which included 3–6 measurement per individual. The first measurement hour

was excluded to minimize potential effects of handling stress and incomplete mixture of air in the

metabolic chamber. Body mass for the metabolic rate measurements was calculated as the

average of the before and after measurement values.

*2.3 Statistical analyses*

*2.3.1 General approach*

All analyses were done using a general linear mixed modeling approach with the function 'lmer'

of the package 'lme4' (Bates et al. 2015) in R version 3.5.3 (R Core Team 2019). All analyses

included individual as a random intercept and their age term (see below) nested within

individuals as a random slope. The random slope quantifies the variation within individual in

ageing and is required for the correct estimation of confidence intervals when investigating within

individual changes (Schielzeth and Forstmeier 2009). Such models require considerable sample

sizes to accurately estimate fixed and random effects (van de Pol 2012) and our data fulfilled those

requirements (Fig. S1). We did not have sufficient power to add other random terms to the model

(e.g. nesting individual identity in aviary). In all analyses, we found the model best supported by

the data, and hence the 'significant' predictors, using Burnham and Anderson's model selection

approach (Burnham and Anderson, 2002; Burnham et al., 2011) based on second order Akaike

Information Criterion (AICc) with the function 'dredge' of the package 'MuMIn' (Barton 2019). In

brief, this is a hypothesis-based approach that generates, given a global model, subset models that

best fit the data. Model fitting should be considered as a continuum for which alternative models

within 4 ΔAICc are plausible and become increasingly equivocal up to 14 ΔAICc, after which they

become implausible (Burnham and Anderson, 2002; Burnham et al., 2011). For models within 4

ΔAICc, we report coefficients which are the result of model averaging from the function

'model.avg' of the package 'MuMIn'. Confidence intervals of model parameters were estimated

with the Wald approximation in the function 'confint'. Residuals of all final models were normal

distributed and without influential data points or outliers.

*2.3.2 Daily and seasonal variation*

Data for all traits were collected throughout the year. To avoid confounding age patterns with

seasonal effects, we corrected for daily and seasonal variation in metabolic rates. To this end, we

first investigated for each trait which variables best characterized the effect of seasonal variation

in trait values following Briga and Verhulst (2017). We quantified the effects of daily and seasonal

variation with and tested the effects of: (i) daylength, (ii) photoperiod, quantified as a

dichotomous variable for increasing vs. decreasing daylength, and (iii) the minimum ambient

temperature (MinT). Temperature data were collected at the weather station of Eelde,

approximately 7 km from the aviaries (http://www.knmi.nl/klimatologie/), where temperature

was recorded 1.5 m above ground, every hour with accuracy of 0.1 °C. Temperature data at the

weather station reflect well the climate at the aviaries as shown from aviary temperature data

collected over the course of the eight years of observation (Briga, 2016). For $BMR_m$, model

selection approach identified that the photoperiod was the most important covariate (ΔAICc=-54,

Table S1A). The next best fitting covariate was daylength, should be included as it worsened the

model fit within the reasonable range of 4 AICc (ΔAICc=+3.1, Table S1A; Burnham et al. 2011).

There was no support for the other tested covariates (ΔAICc≥+8.9, Table S1A). For $SMR_m$, we

found that both photoperiod and MinT were important (ΔAICc=-32 and ΔAICc=-8.7 respectively,

Table S1B). The effect of MinT on trait values can last over a range of timescales (van de Pol and

Cockburn 2011; van de Pol et al. 2016). We therefore weighed MinT over a time window as we

had found earlier to affect lifespan in our study population (Briga and Verhulst, 2015a). This

weighing function included for 77% the MinT within 24 hours before measurement and reached

100% within 5 days. A weighted approach provided a slightly better model fit than using MinT of

the day before measurement (ΔAICc=-1.5). Overall, our results show that both $BMR_m$ and $SMR_m$

increased with shorter and colder days and used the variables that best captured this effect to

correct for daily and seasonal variation in both metabolic traits.

*2.3.3 Metabolic ageing*

Population level associations between trait values and age can be composed of two processes: (i)

a within individual change in trait value with age and (ii) a between individual change due to

selective mortality of individuals with certain trait values. We distinguished the contributions of

these two processes using a within subjects centering approach (van de Pol and Verhulst, 2006;

van de Pol and Wright, 2009). In this approach the within individual changes are captured in a

Δage term, which is the age at measurement mean centered per individual. Models with and

without mean-centering age gave similar conclusions (results not shown), but mean-centering age

always improved the model fit (BMR$_m$: ΔAICc=9.8; SMR$_m$: ΔAICc=13.5; for the best fitting models

in Table S4 A and B respectively). The between individual change is captured by the term lifespan,

mean centered across our population. For censored birds (N=71, see below), lifespan is unknown

and hence these received the mean-centered lifespan of zero. These birds thus contribute only to

within and not to between individual trait change. In this formulation, selective disappearance

occurs when the coefficients differ within and between individuals, which can be statistically

tested with the lifespan term in a model with age and lifespan (van de Pol and Verhulst, 2006; van

de Pol and Wright, 2009), or more directly with survival analyses, which provides straightforward

ways to test whether this disappearance varies between the experimental groups or sexes (see

below).

Within individual changes can follow different age trajectories, including no change with age or a

change that was linear, quadratic, terminal (i.e. before death) or the combination of linear and

terminal. We tested terminal changes by adding a terminal term, coded as a binomial factor for

whether or not an individual died within the year following the measurement. Age trajectories

can also be more complex following threshold models (Douhard et al. 2017; Briga et al. 2019). We

here tested these models, but we lacked the power to include random slopes, to distinguish

between one and two threshold models or to identify threshold ages within a biologically relevant

window. We therefore limited the description of these models in the supplementary materials to

show that threshold models supported the conclusions made by the linear models (Tables S10-

S12).

We tested whether (i) sex and (ii) the environmental manipulations affected metabolic ageing by

including the interaction between the age terms and sex or experimental manipulations. Sex was

always coded as a two-level factor. Effects of the environmental manipulations were tested using

(i) three-way interactions (e.g. Δage*development*adult) and (ii) by grouping experimental group

in one factor with four levels. Thirdly, because the HH group is the only group that showed

reduced lifespan relative to all other groups, we also coded experimental group as a two level

factor, i.e. with HH as one level relative to the three other groups (BB, BH, HB). These three

approaches to test for an effect of the environmental manipulations gave consistent conclusions.

Interactions between sex and the environmental manipulations and higher order interactions

between sex and the environmental manipulation age terms were never significant and we do not

show them here.

*2.3.4 Associations with lifespan*

To investigate the association between mass and survival we used Cox Proportional Hazard

analyses (CPH) using the function 'coxme' of the package 'coxme' (Therneau 2019). To avoid

pseudo-replication by repeated measurements, we used the first measurement of each individual.

These BMR and SMR values were corrected for mass, temporal and seasonal covariates. We then

used the residuals of these values for the CPH analyses. In the CPH models we included sex and

the four experimental groups as covariates, as these are known to affect lifespan (Briga et al. 2017,

Briga et al. 2019). In all these analyses, aviary was included as a random intercept to correct for

the joint housing of birds. Of the 407 individuals for which we have a metabolic rate measurement,

we monitored 336 till natural death and right censored (i) 48 individuals that were still alive after

8 years of monitoring and (ii) 23 individuals that died by accident or were euthanized for welfare

considerations. These 8 years of monitoring are consistent with the study on body mass ageing in

Briga et al. (2019). CPH analyses require predictors to be proportional which was the case as

indicated by the 'cox.zph' function ($X^2$=2.49, p>0.11).

## 3. Results

### 231

*3.1 Basal metabolic rate*

We first investigated the age trajectory of $BMR_m$ within individuals. We tested for changes with age that were either linear, quadratic terminal or a combination of linear and terminal. The best fitting shape was a linear decline with age ($\Delta AICc<-15.7$; Fig. 1A; Table S4A). Individuals lost on average 0.0024 W/Yr (-0.0034<95CI<-0.0015). The variation between individuals in within-individual change with age (i.e. random slopes) was small compared to the variation between individuals in $BMR_m$ (i.e. random intercept), with respectively a variance explained of 0.7% vs. 26%. Thus, $BMR_m$ decreased linearly with age.

We then investigated whether the environmental manipulations that shortened lifespan, also accelerated $BMR_m$ decline. High foraging costs during adulthood, but not during development decreased $BMR_m$ (development: $\Delta AICc=+9.7$; adulthood: $\Delta AICc=-12.6$; Table S2). There was no evidence for different age trajectories (Table S4A) or for a difference in the magnitude of the linear decline between experimental groups (Fig. 3A-D; $\Delta$age * manipulation: $\Delta AICc>+13.8$; terminal year * manipulation: $\Delta AICc>+21.4$; Table S5). Testing the HH group against all other experimental groups gave supported this conclusion ($\Delta AICc>+13.5$). Females lived on average two months shorter than males (4%; Briga et al., 2019). However, sexes did not differ in their $BMR_m$ ($\Delta AICc=+4.7$; Table S2), in their $BMR_m$ response to the environmental manipulations ($\Delta AICc>+11.6$; Table S2) or in their $BMR_m$ ageing (Fig. 3 E&F; $\Delta AICc>+19.2$; Table S7). Thus, $BMR_m$ decreased linearly with age and this decrease was independent of factors that affected lifespan.

We then investigated the association between $BMR_m$ and lifespan. In the best fitting general linear model, the gradual decline in $BMR_m$ with age was indistinguishable between and within individuals (slopes: -0.0018 W/Yr vs. -0.0024 W/Yr respectively), indicating no selective disappearance in any of the experimental groups (Table S5). This is confirmed with CPH models showing neither a linear nor quadratic association with lifespan (e.g. when individuals with intermediate $BMR_m$ live longest; Fig. 2; AICc>+1.3; Table S9A). Thus, there was no association between $BMR_m$ and lifespan.

*3.2 Standard metabolic rate*

We first investigated the shape of the age trajectory, testing for linear, quadratic and/or terminal changes. In contrast to $BMR_m$, which *decreased* linearly with age, $SMR_m$ *increased* linearly with age ($\Delta AICc<-4.0$; Fig. 1B; Table S4B). Individuals gained on average 0.0042 W/Yr (0.0019<95CI<-0.0063). This coefficient seemed to increase with lower ambient temperature Ta at measurement, but this interaction did not improve the model (AICc=+11). The variation between individuals in

within-individual change with age was small compared to the variation between individuals in

$SMR_m$ with respectively a variance explained of 1.5% vs. 20%.

We then investigated whether the manipulations that affected lifespan also affected $SMR_m$ ageing.

High foraging costs during adulthood, but not during development decreased $SMR_m$

(development: $\Delta AICc=+10.5$; adulthood: $\Delta AICc=-67.6$; Table S3). There was no evidence for

different age trajectories (Table S4B) or in the magnitude of the increase between experimental

groups (Fig. 3 G-J; $\Delta$age * manipulation: $\Delta AICc>+8.1$; terminal year * manipulation: $\Delta AICc>+16.1$;

Table S6). Testing the HH group against all other experimental groups supported this conclusion

($\Delta AICc>+11.8$). Just as for $BMR_m$, sexes did not differ in their $SMR_m$ ($\Delta AICc=+11.8$; Table S3), in

their $SMR_m$ response to the environmental manipulations ($\Delta AICc>+21.9$; Table S3), in their age

trajectories or in their rate of $SMR_m$ ageing ($\Delta AICc>+22.1$; Table S8). Thus, $SMR_m$ increased

linearly with age and this increase was independent of factors that affected lifespan.

Lastly, we investigated the association between $SMR_m$ and individual variation in lifespan. In a

GLM, the gradual increase in $SMR_m$ with age differed somewhat between and within individuals

(0.0022 W/Yr vs. 0.0042 W/Yr respectively; Table S6), which suggests there could be some

selective disappearance with respect to $SMR_m$. We found however no statistical support for

selective disappearance in a linear model with age and lifespan ($\Delta AICc=+12.1$) or for linear or

quadratic association between $SMR_m$ and lifespan in a CPH analyses (Fig. 2; $\Delta AICc>+1.7$; Table

S9B). Thus, we found no evidence for an association between $SMR_m$ and lifespan.

### 4. Discussion

Metabolic rate is a key physiological trait in ecology, life history and the biology of ageing. As such it is common to quantify the minimum energy expenditure at thermoneutral temperatures or $BMR_m$. We found that $BMR_m$ *decreased* with age, a result consistent with that in many other endotherms, including in zebra finches (Moe et al. 2009; Rønning et al. 2014; Elliott et al. 2015). However, in contrast to $BMR_m$, $SMR_m$ *increased* with age, at a rate twice the absolute value of $BMR_m$ decline. To the best of our knowledge, this is a new result and the discrepancy in ageing trajectory between these two energetic traits is important because it indicates that these change with age independently. Indeed, the correlations between $BMR_m$ and $SMR_m$ within individual were low ($0.04 < r < 0.22$; Briga and Verhulst, 2017) and these traits' independent ageing will have contributed to this correlation being low. An experimental manipulation of increased foraging costs that decreased lifespan also decreased $BMR_m$ and $SMR_m$, a result consistent with that of earlier studies in birds and laboratory rodents (Wiersma and Verhulst 2005; Vaanholt et al. 2007; Schubert et al. 2008, 2009). However metabolic rate or metabolic ageing were not associated with lifespan in our study. We here discuss these results in the light of other studies on metabolism and ageing.

*4.1 Metabolic ageing*

The mechanism underlying the change in $BMR_m$ with age is likely reductions in the size of metabolic expensive organs, such as heart, liver, kidneys and muscle, as was shown for humans and laboratory rodents (Roberts and Rosenberg 2006). There is also evidence for a small age associated decline in metabolism per unit of tissue (Roberts and Rosenberg 2006) and for reductions in body temperature (Florez-Duquet and McDonald 1998; Weinert 2010; Blatteis 2012). The difference in age trajectory between $BMR_m$ and $SMR_m$ indicates that there is also ageing of thermoregulation. We suggest two explanations for the increase in $SMR_m$ with age. One possibility is that birds need more energy to maintain body temperature as they age, e.g. due to poorer insulation or reduced metabolic efficiency. Secondly, birds in our study have lower body temperatures after nights at sub-thermoneutral temperatures (Briga & Verhulst, 2017), and birds might have developed lower tolerance for low body temperature as they age. This might occur because the ability to warm up decreases with age, at least in laboratory rodents (Florez-Duquet and McDonald 1998). Note however that birds differ from laboratory rodents in that thermoregulatory responses are regulated by skeletal muscles while mammals use brown adipose tissue (Dawson and O'Connor 1996; Mezentseva et al. 2008). Thus, we propose that the mechanisms underlying the different age trajectories between $BMR_m$ and $SMR_m$ reflect changes in insulation and efficiency of heat production, which is key in the light of the energetic and

thermoregulatory challenges associated with winter, predation or with high workload (Brodin

2007; Newton 2007; Speakman and Krol 2010; Nord and Nilsson 2018; Andreassen et al. 2019).

*4.2 Mosaic ageing and the association with lifespan*

Williams (1957) and Maynard-Smith (1962), suggested that traits would evolve to age in

synchrony, because "*natural selection will always be in greatest opposition to the decline of the most*

*senescence-prone system*" (Williams 1957). Our findings run counter to this prediction because

mass, $BMR_m$ and $SMR_m$ showed different age trajectories. Moreover, we previously showed mass

(Briga et al. 2019) and bill colour (Simons et al. 2012; Simons et al. 2016) in the same population

to follow ageing trajectories that differed from both metabolic traits. The age trajectory of mass

was quadratic, but this varied with foraging costs in females, but not in males, and bill color, a

sexual signal, remained constant through adult life until a decline in the terminal year. Our

combined results thus show that traits age asynchronously in zebra finches, a phenomenon coined

*´mosaic ageing'* (Cevenini et al. 2008; Walker and Herndon 2010) which has also been observed

in other systems (Herndon et al. 2002; Bansal et al. 2015; Belsky et al. 2015; Hayward et al. 2015),

but remains to be explained by evolutionary theory. The variation between traits in the net fitness

costs of changes in trait value (accounting for costs of maintaining trait level) has been proposed

as explanation for mosaic ageing, with traits with larger fitness effects predicted to be better

'canalised', i.e. change less with age (Boonekamp et al. 2018), but this hypothesis remains to be

tested in the context of ageing.

Metabolic rate (BMR, SMR) is considered a key trait in the study of life-history, ecology and

evolution (Drent and Daan 1980; Burton et al. 2011), but at the same time studies in endotherms

have not revealed consistent relationships between BMR and fitness (components). In humans,

high $BMR_m$ is associated with increased mortality risk (Ruggiero et al. 2008). In our study, we

found no consistent associations between lifespan and either BMR or SMR, and neither was the

rate of ageing in these metabolic traits associated with lifespan. On the one hand, this is reason to

question whether variation in BMR and SMR between and within individuals can be considered of

functional importance. On the other hand, because these associations are correlational, we cannot

rule out that experimental manipulations of metabolic rates (e.g. through pharmaceutic

treatment) would result in clear fitness effects. Nevertheless, it seems likely that other

physiological pathways are more important in mediating, for example, the effect of environmental

conditions on lifespan. Some promising results indicate that glucose metabolism, the insulin

pathway and the associated glucocorticoid levels are good candidates, although none of these

provides a complete picture yet (Jimeno et al. 2017, 2018; Montoya et al. 2018; Regan et al. 2019).

Hence other physiological mechanisms such as oxidative stress and the immune system also

warrant further study (De Coster et al. 2011; Simons et al. 2014; Speakman et al. 2015).

It is often implicitly assumed that factors changing lifespan will inextricably also alter ageing rate.

To what extent and for what kind of traits this is the case however remains to be identified (Bansal

et al., 2015; Christensen et al., 2009; Williams, 1999). Indeed, in our study system we found little

evidence for such association: the environmental factors affecting lifespan did not affect the

ageing of metabolic traits or body mass (Briga et al. 2019), and neither were rates of ageing higher

in shorter lived individuals. Apparently, (environmental) factors affecting lifespan can be distinct

from factors affecting ageing rate, which was also suggested for genetic factors (Burger and

Promislow 2006). Predicting when or whether an environmental variable that alters lifespan will

also affect the ageing of some traits is therefore a current challenge.

**References**

Andreasson, F., A., Nord, and J. Å. Nilsson. 2019. Age‑dependent effects of predation risk on night‑time hypothermia in two wintering passerine species. Oecologia 189:329–337.

Auer, S. K., R. D. Bassar, K. Salin, and N. B. Metcalfe. 2016. Repeatability of metabolic rate is lower for animals living under field versus laboratory conditions. Journal of Experimental Biology 219:631–634.

Bansal, A., L. J. Zhu, K. Yen, and H. A. Tissenbaum. 2015. Uncoupling lifespan and healthspan in *Caenorhabditis elegans* longevity mutants. Proceedings of the National Academy of Sciences 112:E277–E286.

Barton, K. 2019. Package MuMIn: Multi-Model Inference. Available at: https://CRAN.R-project.org/package=MuMIn.

Bates, D., M. Maechler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models using lme4. Journal of Statistical Software 67:1–48.

Belsky, D. W., A. Caspi, R. Houts, H. J. Cohen, D. L. Corcoran, A. Danese, H. Harrington, et al. 2015. Quantification of biological aging in young adults. Proceedings of the National Academy of Sciences 112:E4104–E4110.

Biro, P. A., and J.A. Stamps, 2010. Do consistent individual differences in metabolic rate promote consistent individual differences in behavior? Trends in Ecology and Evolution 25: 653–659.

Blatteis, C. M. 2012. Age-dependent changes in temperature regulation - A mini review. Gerontology 58:289–295.

Bouwhuis, S., B. C. Sheldon, and S. Verhulst. 2011. Basal metabolic rate and the rate of senescence in the great tit. Functional Ecology 25:829–838.

Boonekamp, J. J., E. Mulder, and S. Verhulst. 2018. Canalisation in the wild: effects of developmental conditions on physiological traits are inversely linked to their association with fitness. Ecology Letters 21:857–864.

Briga, M. 2016. *Growing up and growing old: A longitudinal study on aging in zebra finches*. University of Groningen, Groningen, The Netherlands.

Briga, M., B. Jimeno, and S. Verhulst. 2019. Coupling lifespan and aging? The age at onset of body mass decline associates positively with sex-specific lifespan but negatively with environment-specific lifespan. Experimental Gerontology 119:111–119.

Briga, M., E. Koetsier, J. Boonekamp, B. Jimeno, and S. Verhulst. 2017. Food availability affects adult survival trajectories depending on early developmental conditions. Proceedings of the

Royal Society B Biological Sciences 284:20162287.

Briga, M., and S. Verhulst. 2015*a*. What can long-lived mutants tell us about mechanisms causing
aging and lifespan variation in natural environments? Experimental Gerontology 71:21-26.

Briga, M., and S. Verhulst. 2015*b*. Large diurnal temperature range increases bird sensitivity to
climate change. Scientific Reports 5:16600.

Briga, M., and S. Verhulst. 2017. Individual variation in metabolic reaction norms over ambient
temperature causes low correlation between basal and standard metabolic rate. Journal of
Experimental Biology 220:3280–3289.

Brodin, A. 2007. Theoretical models of adaptive energy management in small wintering birds.
Philosophical Transactions of the Royal Societu of London B 362:1857–1871.

Burger, J. M. S., and D. E. L. Promislow. 2006. Are functional and demographic senescence
genetically independent? Experimental Gerontology 41:1108–1116.

Burnham, K., and D. Anderson. 2002. Model selection and multimodel inference: A practical
information-theoretic approach (2nd ed.). Springer-Verlag, New York.

Burnham, K. P., D. R. Anderson, and K. P. Huyvaert. 2011. AIC model selection and multimodel
inference in behavioral ecology: Some background, observations, and comparisons. Behavioral
Ecology and Sociobiology 65:23–35.

Burton, T., S. S. Killen, J. D. Armstrong, and N. B. Metcalfe. 2011. What causes intraspecific
variation in resting metabolic rate and what are its ecological consequences? Proceedings of the
Royal Society B Biological Sciences 278:3465–3473.

Calder, W. 1964. Gaseous metabolism and water relations of the zebra finch T*aeniopygia*
*castanotis*. Physiological Zoology 37:400–413.

Cevenini, E., L. Invidia, F. Lescai, S. Salvioli, P. Tieri, G. Castellani, and C. Franceschi. 2008. Human
models of aging and longevity. Expert Opinion on Biological Therapy 8:1393–1405.

Christensen, K., G. Doblhammer, R. Rau, and J. W. Vaupel. 2010. Ageing populations : the
challenges ahead. The Lancet 374:1196–1208.

Clay, T. A., E. J. Pearmain, R. A. R. McGill, A. Manica, and R. A. Phillips. 2018. Age-related variation
in non- breeding foraging behaviour and over effects on fitness in an extremely long-lived bird.
Functional Ecology 32:1832–1846.

Dawson, W., and T. O'Connor. 1996. Energetic features of avian thermoregulatory responses.
Pages 85–124 *in* C. Carey, ed. Avian energetics and nutritional ecology. Chapman & Hall, New
York, NY.

De Coster, G., S. Verhulst, E. Koetsier, L. De Neve, M. Briga, and L. Lens. 2011. Effects of early
developmental conditions on innate immunity are only evident under favourable adult
conditions in zebra finches. Naturwissenschaften 98:1049–1056.

Douhard, F., Gaillard, J.M., Pellerin, M. ands L. Jacob. 2017. The cost of growing large: costs of
post-weaning growth on body mass senescence in a wild mammal. Oikos 126:1329–1338.

Drent, R. H., and S.  Daan. 1980. The prudent parent: Energetic adjustments in avian breeding.
Ardea, 68, 225– 252.

Elliott, K. H., J. F. Hare, M. Le Vaillant, A. J. Gaston, Y. Ropert-Coudert, and W. G. Anderson. 2015.
Ageing gracefully: Physiology but not behaviour declines with age in a diving seabird. Functional
Ecology 29:219–228.

Florez-Duquet, M., and R. B. McDonald. 1998. Cold-induced thermoregulation and biological
aging. Physiological Reviews 78:339–358.

Fontana, L., and L. Partridge. 2015. Promoting health and longevity through diet: From model
organisms to humans. Cell 161:106–118.

Gaillard, J.-M., and J.-F. Lemaitre. 2017. The Williams' legacy: A critical reappraisal of his nine
predictions about the evolution of senescence. Evolution 71:2768–2785.

Griffith, S. C., O. L. Crino, S. C. Andrew, F. Y. Nomano, E. Adkins-Regan, C. Alonso-Alvarez, I. E.
Bailey, et al. 2017. Variation in reproductive success across captive populations: methodological
differences, potential biases and opportunities. Ethology 123:1–29.

Hayward, A. D., J. Moorad, C. E. Regan, C. Berenos, J. G. Pilkington, J. M. Pemberton, and D. H.
Nussey. 2015. Asynchrony of senescence among phenotypic traits in a wild mammal population.
Experimental Gerontology 71:56–68.

Herndon, L. A., P. J. Schmeissner, J. M. Dudaronek, P. A. Brown, K. M. Listner, Y. Sakano, M. C.
Paupard, et al. 2002. Stochastic and genetic factors influence tissue-specific decline in ageing C .
elegans. Nature 419:808–814.

Hill, R. W. 1972. Determination of oxygen consumption by use of the paramagnetic oxygen
analyzer. Journal of Applied Physiology 33:261–263.

Jimeno, B., M. Briga, M. Hau, and S. Verhulst. 2018. Male but not female zebra finches with high
stress-induced corticosterone have lower survival. Functional Ecology 32:713–721.

Jimeno, B., M. Briga, S. Verhulst, and M. Hau. 2017. Effects of developmental conditions on
glucocorticoid concentrations in adulthood depend on sex and foraging conditions. Hormones
and Behavior 93:175–183.Koetsier, E., and S. Verhulst. 2011. A simple technique to manipulate

foraging costs in seed-eating birds. Journal of Experimental Biology 214:1225–1229.

Maynard-Smith, J. 1962. Review lectures on senescence. I. The causes of ageing. Proceedings of
the Royal Society of London, Series B Biological Sciences B157:115–127.

Mezentseva, N. V, J. S. Kumaratilake, and S. A. Newman. 2008. The brown adipocyte
differentiation pathway in birds: an evolutionary road not taken. BMC Biology 6:17.

Moe, B., B. Rønning, S. Verhulst, and C. Bech. 2009. Metabolic ageing in individual zebra finches.
Biology Letters 5:86–89.

Montoya, B., M. Briga, B. Jimeno, S. Moonen, and S. Verhulst. 2018. Baseline glucose level is an
individual trait that is negatively associated with lifespan and increases due to adverse
environmental conditions during development and adulthood. Journal of Comparative
Physiology B: Biochemical, Systemic, and Environmental Physiology 188:517–526.

Newton, I. 2007. Weather-related mass-mortality events in migrants. Ibis 149: 453–467.

Nord, A. and J. Å. Nilsson. 2018. Heat dissipation rate constrains reproductive investment in a
wild bird. Functional Ecology 33: 250–259.

Norin, T., and N. B. Metcalfe. 2019. Ecological and evolutionary consequences of metabolic rate
plasticity in response to environmental change. Philosophical Transactions of the Royal Society
B 374:20180180.

Nussey, D. H., H. Froy, J. F. Lemaitre, J. M. Gaillard, and S. N. Austad. 2013. Senescence in natural
populations of animals: Widespread evidence and its implications for bio-gerontology. Ageing
Research Reviews 12:214–225.

Partridge, L., and J. Deelen. 2018. Facing up to the global challenges of ageing. Nature 561:45–56.

R Core Team. 2019. R: A language and environment for statistical computing. Available at:
https://CRAN.R-project.org/package=secrdesign.

Regan, J. C., H. Froy, C. A. Walling, J. P. Moatt, and D. H. Nussey. 2019. Dietary restriction and
insulin-like signalling pathways as adaptive plasticity: A synthesis and re-evaluation. Functional
Ecology, In Press.

Roberts, S. B., and I. Rosenberg. 2006. Nutrition and aging: changes in the regulation of energy
metabolism with aging. Physiological Reviews 86:651–667.

Ruggiero, C., E. J. Metter, V. Melenovsky, A. Cherubini, S. S. Najjar, A. Ble, U. Senin, D. L. Longo, and
L. Ferrucci. 2008. High basal metabolic rate is a risk factor for mortality: the Baltimore
Longitudinal Study of Aging. Journal of Gerontology A 63:698–706.

Rønning, B., B. Moe, H. H. Berntsen, E. Noreen and C. Bech 2014. Is the rate of metabolic ageing

and survival determined by basal metabolic rate in the zebra finch? Plos One 9:e108675.

Schielzeth, H., and W. Forstmeier. 2009. Conclusions beyond support: overconfident estimates in
mixed models. Behavioral Ecology 20:416–420.

Schubert, K. A., G. de Vries, L. M. Vaanholt, H. A. J. Meijer, S. Daan, and S. Verhulst. 2009. Maternal
energy allocation to offspring increases with environmental quality in house mice. American
Naturalist 173:831–840.

Schubert, K. A., L. M. Vaanholt, F. Stavasius, G. E. Demas, S. Daan, and G. H. Visser. 2008. Female
mice respond differently to costly foraging versus food restriction. Journal of Experimental
Biology 211:2214–2223.

Simons M.J.P., M. Briga, E. Koetsier, R. Folkertsma, M. D. Wubs, C. Dijkstra, and S. Verhulst. 2012.
Bill redness is positively associated with reproduction and survival in male and female zebra
finches. PLoS One 7: e40721.

Simons, M. J. P., M. Briga, B. Leenknegt, and S. Verhulst. 2014. Context-dependent effects of
carotenoid supplementation on reproduction in zebra finches. Behavioral Ecology 25:945–950.

Simons, M. J. P., M. Briga, and S. Verhulst. 2016. Stabilizing survival selection on presenescent
expression of a sexual ornament followed by a terminal decline. Journal of Evolutionary Biology
29:1368–1378.

Speakman, J.R., and E. Krol. 2010. Maximal heat dissipation capacity and hyperthermia risk:
neglected key factors in the ecology of endotherms. Journal of Animal Ecology 79: 726–746.

Speakman, J. R., J. D. Blount, A. M. Bronikowski, R. Buffenstein, C. Isaksson, T. B. L. Kirkwood, P.
Monaghan, et al. 2015. Oxidative stress and life histories: Unresolved issues and current needs.
Ecology and Evolution 5:5745–5757.

Therneau, T. 2019. coxme: Mixed Effects Cox Models. Available at: https://CRAN.R-
project.org/package=coxme.

Vaanholt, L. M., B. De Jong, T. Garland, S. Daan, and G. H. Visser. 2007. Behavioural and
physiological responses to increased foraging effort in male mice. Journal of Experimental
Biology 210:2013–2024.

van de Pol, M. 2012. Quantifying individual variation in reaction norms: how study design affects
the accuracy, precision and power of random regression models. Methods in Ecology and
Evolution 3:268–280.

van de Pol, M., L. D. Bailey, N. McLean, L. Rijsdijk, and C. R. Lawson. 2016. Identifying the best
climatic predictors in ecology and evolution. Methods in Ecology and Evolution 7:1246–1257.

van de Pol, M., and A. Cockburn. 2011. Identifying the critical climatic time window that affects
trait expression. American Naturalist 177:698–707.

van de Pol, M., and S. Verhulst. 2006. Age-dependent traits: a new statistical model to separate
within- and between-individual effects. American Naturalist 167:766–773.

van de Pol, M., and J. Wright. 2009. A simple method for distinguishing within- versus between-
subject effects using mixed models. Animal Behaviour 77:753–758.

Walker, L. C., and J. G. Herndon. 2010. Mosaic aging. Medical hypotheses 74:1048–1051.

Weinert, D. 2010. Circadian temperature variation and ageing. Ageing Research Reviews 9:51–
60.

Wiersma, P., and S. Verhulst. 2005. Effects of intake rate on energy expenditure, somatic repair
and reproduction of zebra finches. Journal of Experimental Biology 208:4091–4098.

Williams, G. C. 1957. Pleiotropy, natural-selection and the evolution of senescence. Evolution
11:398–411.

Williams, G. C. 1999. The tithonus error in modern gerontology. The Quarterly Review of Biology
74:405–415.

Zann, R. A. 1996. The Zebra Finch: a synthesis of field and laboratory studies. Oxford University
Press, Oxford.

Fig. 1. Within individuals (A) BMR$_m$ decreased with age, while (B) SMR$_m$ increased. Values are

residuals corrected for mass, seasonal variation, experimental groups, sex and for SMR$_m$ also the

ambient temperature at measurement. Boxplots show medians, quartiles and 95% CI per year.

Horizontal lines illustrate model fit. Note the Y-axes on different scales between (A) and (B).

Fig. 2. BMR$_m$ (A) or SMR$_m$ (B) were not associated with lifespan. To avoid pseudo-replication by

repeated measurements, we used the first measurement of each individual. BMR$_m$ or SMR$_m$ values

are residuals corrected for mass, seasonal variation, ambient temperature at measurement

(SMR$_m$), experimental groups and sex, as in the survival analyses in Table S9.

Fig. 3. Within-individual changes in $BMR_m$ and $SMR_m$ are independent of environmental quality ($BMR_m$: A-D; $SMR_m$: G-J) and of sex  ($BMR_m$: E, F; $SMR_m$:

K, L). Experimental group combinations are abbreviated, in chronological order, such that the first letter stands for developmental environment (B for

benign or small broods, H for harsh or large broods) and the second letter for adult environment (B for benign or low foraging costs, H for harsh or

high foraging costs). $BMR_m$ or $SMR_m$ values are residuals corrected for mass, seasonal variation, ambient temperature at measurement ($SMR_m$), sex (A-

D; G-J), experimental groups (E, F, K, L). Boxplots show medians, quartiles and 95% CI per year. Horizontal lines are group specific model fits (± 95%CI).

554