1

Characterization of *Mollivirus kamchatka*, the first modern representative

of the proposed *Molliviridae* family of giant viruses.

Running title: characterization of the first modern mollivirus

Eugene Christo-Foroux[#a], Jean-Marie Alempic[a], Audrey Lartigue[a], Sebastien Santini[a],

Karine Labadie[b], Matthieu Legendre[a] , Chantal Abergel[a] , Jean-Michel Claverie[#a]

a Aix Marseille Univ., CNRS, IGS, Information Génomique & Structurale (UMR7256),

Institut de Microbiologie de la Méditerranée (FR 3489), Marseille, France

b Genoscope, Institut François Jacob, CEA, Université Paris-Saclay, Évry, France

Correspondance to: Eugene.christo-foroux@igs.cnrs-mrs.fr, jean-michel.claverie@univ-amu.fr

Keywords: Paleovirology; Kamchatka; Comparative Genomics; Nucleo-cytoplasmic Large DNA
viruses; Genome Structure.

Abstract

Microbes trapped in permanently frozen paleosoils (permafrost) are the focus of increasing

researches in the context of global warming. Our previous investigations led to the discovery and

reactivation of two Acanthamoeba-infecting giant viruses, *Mollivirus sibericum* and *Pithovirus*

sibericum from a 30,000-year old permafrost layer. While several modern pithovirus strains have

since been isolated, no contemporary mollivirus relative was found. We now describe *Mollivirus*

kamchatka, a close relative to *M. sibericum*, isolated from surface soil sampled on the bank of the

Kronotsky river in Kamchatka. This discovery confirms that molliviruses have not gone extinct and

are at least present in a distant subarctic continental location. This modern isolate exhibits a

nucleo-cytoplasmic replication cycle identical to that of *M. sibericum*. Its spherical particle (0.6-µm

in diameter) encloses a 648-kb GC-rich double stranded DNA genome coding for 480 proteins of

which 61 % are unique to these two molliviruses. The 461 homologous proteins are highly

conserved (92 % identical residues in average) despite the presumed stasis of *M. sibericum* for the

last 30,000 years. Selection pressure analyses show that most of these proteins contribute to the

virus fitness. The comparison of these first two molliviruses clarify their evolutionary relationship

with the pandoraviruses, supporting their provisional classification in a distinct family, the

Molliviridae, pending the eventual discovery of intermediary missing links better demonstrating

their common ancestry.

Importance

Virology has long been viewed through the prism of human, cattle or plant diseases leading to a

largely incomplete picture of the viral world. The serendipitous discovery of the first giant virus

visible under light microscopy (i.e., >0.3μm in diameter), mimivirus, opened a new era of

environmental virology, now incorporating protozoan-infecting viruses. Planet-wide isolation

studies and metagenomes analyses have shown the presence of giant viruses in most terrestrial

and aquatic environments including upper Pleistocene frozen soils. Those systematic surveys have

led authors to propose several new distinct families, including the *Mimivirida*e, *Marseilleviridae*,

Faustoviridae, *Pandoraviridae*, and *Pithoviridae*. We now propose to introduce one additional

family, the *Molliviridae*, following the description of *M. kamchatka*, the first modern relative of *M.*

sibericum, previously isolated from 30,000-year old arctic permafrost.

Introduction

Studies started about 30 years ago, have provided multiple evidence that soils frozen since the late Pleistocene and predominantly located in arctic and subarctic Siberia, do contain a wide diversity of microbes that can be revived upon thawing (1-3) after tens of thousands of years. These studies culminated by the regeneration of a plant from 30,000 year-old fruit tissue (4). Inspired by those studies, we then isolated from a similar sample two different Acanthamoeba-infecting large DNA viruses, named *Pithovirus sibericum* and *Mollivirus sibericum*, demonstrating the ability for these viruses, and maybe many others, to remain infectious after similarly long periods of stasis in permafrost (5-6). Several modern relatives to the prototype Pithovirus have since been isolated and characterized, leading to the emergence of the new proposed *Pithoviridae* family (7-10). In contrast, and despite the increasing sampling efforts deployed by several laboratories, no other relative of *M. sibericum* was found. Without additional isolates, its classification as a prototype of a new family, or as a distant relative of the pandoraviruses (with which it shared several morphological features and 16% of its gene content) (6) remained an open question. Here we report the discovery and detailed characterization of the first modern *M. sibericum* relative, named *Mollivirus kamchatka*, after the location of the Kronotsky river bank where it was retrieved. The comparative analysis of these first two molliviruses highlights their evolutionary processes and suggests their provisional classification into their own family, the *Molliviridae*, distinct from the *Pandoraviridae*, pending the eventual discovery of intermediary missing links clearly establishing their common ancestry.

Results

Virus isolation

The original sample consisted of about 50 ml of vegetation-free superficial soil scooped (in sterile

tubes) from the bank of the Kronotsky river (coordinates: N 54°32'59'' E 160°34'55'') on july 6[th],

2017. Before being stored at 4°C, the sample was transported in a backpack for a week at ambient

temperature (5° C up to 24° C). This area corresponds to a continental subarctic climate: very cold

winters, and short, cool to mild summers, low humidity and little precipitation. Back in the

laboratory, few grams of the sample were used in the Acanthamoeba co-cultivation procedure

previously described (6). After a succession of passages and enrichment on *Acanthamoeba*

castellanii cultures, viral particles were produced in sufficient quantity to be recovered and

purified.

Virion morphology and ultrastructure

As for *M. sibericum*, light microscopy of infected cultures showed the multiplication of particles.

Using transmission electron microscopy (TEM), these particles - undistinguishable from that of *M.*

sibericum - appear approximately spherical, 600 nm in diameter, lined by an internal lipid

membrane and enclosed in a 20 nm-thick electron-dense thick tegument covered with a mesh of

fibers (Fig. 1).

Analysis of the replication cycle

The replication cycle in *A. castellanii* cells was monitored using TEM and light microscopy of DAPI-

stained infected cultures as previously described (6). The suite of events previously described for

host cells infected by *M. sibericum*, was similarly observed upon *M. kamchatka* replication (6, 11).

After entering the amoeba cell through phagocytosis, *M. Kamchatka* virions are found gathered in

large vacuoles individually or in groups of 2-6 particles. Multiple nuclear events occur during the

infection starting with the drift of the host cell nucleolus to the periphery of the nucleus 4-5h post

infection (p.i.). 7h p.i., the nucleus appears filled with numerous fibrils that may correspond to viral

genomes tightly packed in DNA-protein complexes (Fig. 2A). About 30% of the nuclei observed at

that time exhibit a ruptured nuclear membrane (Fig. 2B). Besides those internal nuclear events, we

observed a loss of vacuolization within the host cell from 4h p.i. to the end of the cycle, in average

9h p.i.. Large viral factories are formed in the cytoplasm at the periphery of the disorganized

nucleus (Fig. 2). These viral factories displays the same characteristics as those formed during *M.*

sibericum infections, involving an active recycling of membrane fragments (11).

Comparative genomics

DNA prepared from purified *M. kamchatka* particles was sequenced using both Illumina and

Oxford Nanopore platforms. *M. kamchatka* genome, a linear double stranded DNA molecule

(dsDNA), was readily assembled as a unique sequence of 648,864 bp. The read coverage was

uniform throughout the entire genome except for a 10 kb terminal segment presumably repeated

at both ends and exhibiting twice the average value. The *M. kamchatka* genome is thus

topologically identical to that of a *M. sibericum*, slightly larger in size (when including both terminal

repeats), and with the same global nucleotide composition (G+C= 60%). This similarity was

confirmed by the detailed comparison of their genome sequences exhibiting a global collinearity

solely interrupted by a few insertions and deletions.

Prior to the comparison of their gene contents, *M. sibericum* and *M. kamchatka* were both

annotated using the same stringent procedure that we previously developed to correct for gene

overpredictions suspected to occur in G+C rich sequences such as those of pandoraviruses (12-14).

A total of 495 and 480 genes were predicted for *M. sibericum* and *M. kamchatka*, with the

encoded proteins ranging from 51 to 2171 residues and from 57 to 2176 residues, respectively.

M. kamchatka predicted protein sequences were used in similarity search against the non-

redundant protein sequence database (15) and the re-annotated *M. sibericum* predicted

proteome. Out of the 480 proteins predicted to be encoded by *M. kamchatka* 463 had their closest

homologs in *M. sibericum*, with 92% identical residues in average. After clustering the paralogs,

these proteins corresponded to 434 distinct genes clusters delineating a first estimate of the

mollivirus core gene set. Four hundred and eleven of these clusters contained a single copy

(singletons) gene for each strain. Remarkably, 290 of the 480 (60.4 %) *M. kamchatka*-encoded

proteins did not exhibit a detectable homolog among cellular organisms or previously sequenced

viruses (excluding *M. sibericum*). Those will be referred to as "ORFans". Among the 190 proteins

exhibiting significant ($E<10^{-5}$) matches in addition to their *M. sibericum* counterparts, 78 (16% of

the total gene content) were most similar to Pandoravirus predicted proteins, 18 (3.7%) to proteins

of other virus families, 51 (10.6%) to *A. castellanii* proteins, 24 (5%) to proteins of other

eukaryotes, 17 (3.5%) to bacterial proteins and 2 (0.4%) to proteins of *Archaea* (Fig. 3).

The interpretation of these statistics are ambiguous as, on one hand, the large proportion of

"ORFans" (>60%) is characteristic of what is usually found for the prototypes of novel giant virus

families (7). On the other hand, the closest viral homologs are not scattered in diverse previously

defined virus families, but mostly belongs to the Pandoraviridae (78/96=81%) (Fig. 3). The two

molliviruses thus constitute a new group of viruses with their own specificity but with a

phylogenetic affinity with the pandoraviruses, as previously noticed (7). The proportion of *M.*

kamchatka proteins with best matching counterparts in Acanthamoeba confirms the high gene

exchange propensity with the host, already noticed for *M. sibericum* (6).

Recent evolutionary events since the *M. sibericum*/*M. kamchatka* divergence

We investigated the evolutionary events specific of each of the molliviruses by focusing on proteins

lacking reciprocal best matches between the two strains. We found 63 such cases of which 10

corresponded to unilateral strain-specific duplications of genes, and 53 were unique to a given

strain. These unique genes (Table 1 & 2) result from gains or losses in either of mollivirus strains

(20 in *M. Kamchatka*, 33 in *M. sibericum*). The likely origins of these strain-specific genes

(horizontal acquisition, *de novo* creation (13, 14), or differential loss) are listed in Table 1 and Table

2.

Six *M. kamchatka* proteins, absent from *M. sibericum*, have homologs in pandoraviruses

suggesting common gene ancestors (and loss in *M. sibericum*) or horizontal acquisitions. According

to its embedded position within the pandoravirus phylogenetic tree, only one anonymous protein

(mk_165) could be interpreted as a probable horizontal transfer from pandoraviruses (Fig. 4A).

Another candidate, mk_92, shares 75% of identical residues with a pandoravirus DNA

methyltransferase (pqer_cds_559). However the very long branch associated to the P. dulcis

homolog (eventually due to a non-orthologous replacement) raises some doubt as for the origin of

the *M. kamchatka* gene (Fig. 4B).

Two *M. kamchatka*-specific proteins, encoded by adjacent genes (mk_466, mk_467), have

homologs in Acanthamoeba, suggesting potential host-virus exchanges. Phylogenetic

reconstruction did not suggest a direction for the transfer of mk_466. However, since the unique

homolog of mk_467 is found in Acanthamoeba (and not in other eukaryotes), the corresponding

gene probably originated from a close relative of *M. kamchatka* and was recently transferred to its

host.

Three proteins unique to *M. sibericum* have homologs in pandoraviruses (Table 2), suggesting

common gene ancestors (and loss in *M. kamchatka*) or horizontal acquisitions. One protein

(ms_14) has a unique homolog in Acanthamoeba, suggesting a virus to host exchange. The

homolog of ms_312 in *Cavenderia fasciculata* (16) might be the testimony of past interactions

between molliviruses and deeply rooted ancestor of the Amoebozoa clade.

The above analyses of the genes unique to each mollivirus indicate that if horizontal transfer

may contribute to their presence, it is not the predominant mechanisms for their acquisition. We

then further investigated the 12 genes unique to *M. kamchatka* and 26 genes unique to *M.*

sibericum (i.e. "strain ORFans" w/o homolog in the databases) by computing three independent

sequence properties (Fig. 5): the codon adaptation usage index (CAI), the G+C composition, and

the ORF length.

With an average CAI value of 0.26, the strain-specific ORFans appear significantly different

from the rest of the mollivirus genes (mean = 0.40, Wilcoxon test p< 2 10^{-7}). These genes also

exhibit a significantly lower G+C content (56% for *M. kamchatka* and 57% for *M. sibericum*) than

the rest of the genes (60.5% for both viruses), also closer to the value computed for intergenic

regions (54% in average for both viruses). Moreover, the strain-specific ORFans are smaller in

average compared to the rest of the genes (115bp/378bp for *M. kamchatka* and 122bp/369bp for

M. sibericum). Altogether, those results suggest that *de novo* gene creation might occur in the

intergenic regions of molliviruses as already postulated for pandoraviruses (13, 14).

New predicted protein functions in *M. kamchatka*

Sixty four of the *M. kamchatka* predicted proteins exhibit sequence motifs associated to known

functions. Fifty nine of them are orthologous to previously annotated genes in *M. sibericum* (6).

This common subset confirms the limited complement of DNA processing and repair enzymes

found in molliviruses: mainly a DNA polymerase: mk_287, a primase: mk_236, and 3 helicases:

mk_291, mk_293, mk_351. *M. kamchatka* confirms the absence of key deoxynucleotide synthesis

pathways (such as thymidylate synthase, thymidine kinase and thymidylate kinase), and of a

ribonucleoside-diphosphate reductase (present in pandoraviruses). The five *M. kamchatka* specific

proteins (Table 1) associated to functional motifs or domain signature correspond to:

- two proteins (mk_93 and mk_104) containing a type of zinc finger (Ring domain) mediating

protein interactions,

- one protein (mk_469) with similarity to the (BI)-1 like family of small transmembrane proteins,

- one predicted LexA-related signal peptidase (mk_166),

- one DNA methyltransferase (mk_92).

Evaluation of the selection pressure exerted on mollivirus genes

The availability of two distinct strains of mollivirus allows the first estimation of the selection

pressure exerted on their shared genes during their evolution. This was done by computing the

ratio ω=dN/dS of the rate of non-synonymous mutations (dN) over the rate of synonymous

mutations (dS) for pairs of orthologous genes. ω values much lesser than one are associated to

genes the mutation of which have the strongest negative impact on the virus fitness. The high

sequence similarity of proteins shared by *M. sibericum* and *M. kamchatka* allowed the generation

of flawless pairwise alignments and the computation of highly reliable ω values for most (*i.e.*

397/411) of their orthologous singletons.

Fourteen singleton pairs were not taken into account in the selection pressure analysis

because of their either identical or quasi-identical sequences (11 of them), or unreliable pairwise

alignments (3 of them). For the 397 gene pairs retained in the analysis, the mean ω value was 0.24

± 0.14 (Fig. 6). This result corresponds to a strong negative selection pressure indicating that most

of the encoded proteins greatly contribute to the molliviruses' fitness. Together with the high level

of pairwise similarity (92%) of their proteins, this also indicates that *M. kamchatka* evolved very

little during the last 30,000 years and that the *M. sibericum* genome was not prominently damaged

during its cryostasis in permafrost.

The analysis restricted to the 244 pairs of ORFan-coding genes resulted in a very similar ω

value of 0.29± 0.15 (Fig. 6). This indicates that although homologs of these proteins are only found

in molliviruses, they have the same impact on the virus fitness than more ubiquitous proteins. This

confirms that they do encode actual proteins, albeit with unknown functions. In contrast, four

orthologous pairs (ms_160/mk_141; ms_280/mk_262; ms_171/mk_151; ms_430/mk_411;

ms_60/mk_48) exhibit ω value larger than one. Those ORFans either are under positive selection

for maintaining their functions, or newly created gene products undergoing refinement or

pseudogenization.

We further examined the selection pressure of proteins-coding genes with homologs in

pandoraviruses. We used their 10 sequenced genomes to generate the corresponding gene

clusters (Fig. 7). The 90 clusters shared by both virus groups included 64 singletons (single copy

gene present in all viruses), among which 55 were suitable for dN/dS computations. The mean ω

value (0.17 ± 0.1) was very low, indicating that these genes, forming a "super core" gene set

common to the molliviruses and pandoraviruses, are under an even stronger negative selection

pressure than those constituting the provisional (most likely overestimated) mollivirus core gene

set .

Genomic inhomogeneity

The original genome analysis of Lausannevirus (a member of the Marseilleviridae family) (17)

revealed an unexpected non-uniform distribution of genes according to their annotation.

"Hypothetical" genes (i.e. mostly ORFans) were segregated from "annotated" (i.e. mostly non-

ORFans) in two different halves of the genome. In a more recent work, we noticed a similar bias in

the distribution of Pandoravirus core genes (13). The availability of a second mollivirus isolate gave

us the opportunity to investigate this puzzling feature for yet another group of Acantamoeba-

infecting virus. In Fig. 8, we plotted the distribution of three types of genes: 1- those with

homologs in *A. castellanii* (n= 55 for *M. sibericum* and n= 51 for *M. kamchatka*), 2- those belonging

to the super core set shared by both molliviruses and pandoraviruses (n=64), 3- those unique to

either mollivirus strains (n=26 for *M. sibericum*, n=12 for *M. kamchatka*). These plots reveal a

strong bias in the distribution of the super core vs. ORFan genes (Fig. 8). The first half of the *M.*

sibericum genome exhibits 90% of its ORFans while the second half contains most of the members

of the super core gene set. In contrast, genes eventually exchanged with the host display a more

uniform distribution. The lack of an apparent segregation in the distribution of ORFans in the *M.*

kamchatka genome might be due to their underprediction as no transcriptome information is

available for this strain. Fig. 9 shows that there is also a strong bias in the distribution of single-

copy genes *vs.* those with paralogs in either *M. sibericum* and/or *M. kamchatka*. Altogether, these

analyses suggest that the two genome halves follow different evolutionary scenario, the first half

concentrating the genomic plasticity (*de novo* gene creation, gene duplication), the other half

concentrating the most conserved, eventually essential, gene content.

Discussion

Following the discovery of their first representatives, each families of giant (e.g. Mimiviridae,

Pithoviridae, Pandoraviridae) and large (e.g. Marseilleviridae) viruses infecting acanthamoeba have

expanded steadily, suggesting they were relatively abundant and present in a large variety of

environments. One noticeable exception has been the molliviruses, the prototype of which

remained unique after its isolation from 30,000-year old permafrost. The absence of *M. sibericum*

relatives from the large number of samples processed by others and us since 2014, raised the

possibility that they might have gone extinct, or might be restricted to the Siberian arctic. Our

isolation of a second representative of the proposed Molliviridae family, *M. kamtchatka*, at a

location more than 1,500 km from the first isolate and enjoying a milder climate, is now refuting

these hypotheses. Yet, the planet-wide ubiquity of these viruses remains to be established, in

contrast to other acanthamoeba-infecting giant viruses (7). Even when present, mollivirus-like

viruses appear to be in very low abundance, as judged from the very small fraction of

metagenomics reads they represent in total sample DNA for *M. kamchatka* (about 0.02 part per

million) as well as for *M. sibericum* (about one part per million)(6). Another possibility would be

that the actual environmental host is not an acanthamoeba, the model host used in our laboratory.

However, evidences of specific gene exchanges with acanthamoeba (including a highly conserved

homolog major capsid protein) (6, 18, 19) make this explanation unlikely. We conclude that

members of the proposed Molliviridae family are simply less abundant than other acanthamoeba-

infecting viruses, a conclusion further supported by the paucity of Mollivirus-related sequences in

the publicly available metagenomics data (data not shown).

As always the case, the characterization of a second representative of a new virus

representative opened new opportunities of analysis. Unfortunately, the closeness of *M.*

kamchatka with *M. sibericum* limited the amount of information that could be drawn from their

comparison. For instance, the number of genes shared by the two isolates is probably a large

overestimate of the "core" gene set characterizing the whole family. On the other hand, the

closeness of the two isolates allowed an accurate determination of the selection pressure

(ω=dN/dS) exerted on many genes, showing that most of them, including mollivirus ORFans,

encode actual proteins the sequence of which are under strong negative selection and thus

contribute to the virus fitness. Given the partial phylogenetic affinity (i.e. 90 shared gene clusters)

of the mollivirus with the pandoraviruses, we also assessed the selection pressure exerted on 55 of

these "super core" genes, and found them under even stronger negative selection (Fig. 7). This

suggests that this super core gene set might have been present in a common ancestor to both

proposed families.

If we postulate that *M. sibericum* underwent into a complete stasis when it became frozen in

permafrost while *M. kamchatka* remained in contact with living acanthamoeba, we could consider

the two viral genomes to be separated by at least 30,000 years of evolution (eventually more if

they are not in a direct ancestry relationship)(20). The high percentage of identical residues (92%)

in their proteins corresponds to a low substitution rate of 1.7×10^{-6} amino acid change/position

/year. This is an overestimate since the two viruses probably started to diverge from each other

longer than 30,000-year ago. This value is nevertheless comparable with estimates computed for

poxviruses (21) given the uncertainty on the number of replicative cycles occurring per year. The

high level of sequence similarity of *M. kamchatka* with *M. sibericum* also indicates that the later

did not suffer much DNA damage during its frozen stasis, even in absence of detectable virus-

encoded DNA repair functions.

Horizontal gene transfers with the host were suggested by the fact that 51 proteins shared by

the two mollivirus strains exhibited a second best match in acanthamoeba. Because no homolog is

detected in other eukaryotes for most of them, these transfers probably occurred in the mollivirus-

to-host direction. The clearest case is that of a major capsid protein homolog (mk_314, ml_347)

sharing 64% identical residues with a predicted acanthamoeba protein (locus: XP_004333827).

Two other genes encoding proteins that have also homologs in molliviruses flank the

corresponding host gene. However, the corresponding viral genes are not collinear in *M. sibericum*

or *M. kamchatka* and were probably transferred from a different, yet unknown mollivirus strain.

The presence of a 100% conserved major capsid protein homolog in the genome of *M. kamchatka*

and *M. sibericum* is itself puzzling. Such protein (with a double-jelly roll fold) is central to the

structure of icosahedral particles (22). Consistent with its detection in *M. sibericum* virions (6), its

conservation in *M. kamchatka*, suggests that it still plays a role in the formation of the spherical

mollivirus particles, while it has no homolog in the pandoraviruses. Inspired by previous

observations made on the unrelated Lausannevirus genome (17), we unveiled a marked

asymmetry in the distribution of different types of protein-coding genes in the Mollivirus genomes.

As shown in Fig. 8 the left half of the genome concentrates most of the genes coding for strain-

specific ORFans while the right half concentrates most of super core genes shared with

pandoraviruses. This asymmetry is even stronger for the multiple copy genes while single-copy

genes are uniformly distributed along the genome (Fig. 9). The molliviruses thus appear to confine

their genomic "creativity" (*de novo* creation and gene duplication) in one-half of their genome,

leaving the other half more stable. An asymmetry in the distribution of the core genes was

previously noticed in the pandoravirus genomes (13). Such features might be linked to the

mechanism of replication that is probably similar for the two virus families. Further studies are

needed to investigate this process. The asymmetrical genomic distribution of pandoravirus core

genes and mollivirus super core genes might be a testimony of their past common ancestry.

Despite their differences in morphology, as well as in virion and genome sizes, the comparative

analysis of the prototype *M. sibericum* and of the new isolate *M. kamchatka* confirms their

phylogenetic affinity with the Pandoraviruses (Fig. 3, Fig. 10). However, it remains unclear whether

this is due to a truly ancestral relationship between them, or if it is only the consequence of

numerous past gene exchanges favored by the use of the same cellular host. From the perspective

of the sole DNA polymerase sequence, the two known molliviruses do cluster with the

pandoraviruses, albeit at a larger evolutionary distance than usually observed between members

of the same virus family (Fig. 10). In absence of an objective threshold, and pending the

characterization of eventual "missing links", we thus propose to classify *M. sibericum* and *M.*

kamchatka as members of the proposed Molliviridae family, distinct from the Pandoraviridae.

Materials and Methods

Virus isolation

We isolated *M. kamchatka* from muddy grit collected near Kronotski Lake, Kamchatka (Russian

Federation N :54 32 59, E :160 34 55). The sample was stored for twenty days in pure rice medium

(23) at room temperature. An aliquot of the pelleted sample triggered an infected phenotype on a

culture of *Acanthamoeba castellanii Neff* (ATCC30010TM) cells adapted to 2,5µg/mL of

Amphotericin B (Fungizone), Ampicillin (100µg/ml), Chloramphrenicol (30µg/ml) and Kanamycin

(25µg/ml) in protease-peptone–yeast-extract–glucose (PPYG) medium after two days of incubation

at 32°C. A final volume of 6 mL of supernatant from two T25 flasks exhibiting infectious

phenotypes was centrifuged for 1 hour at 16,000xg at room temperature. Two T75 flasks were

seeded with 60,000 cells/cm² and infected with the resuspended viral pellet. Infected cells were

cultured in the same conditions as described below. We confirmed the presence of viral particles

by light microscopy.

Validation of the presence of *M. kamchatka* in the original sample

To confirm the origin of the *M. kamchatka* isolate from the soil of the Kronotsky river bank,

DNA was extracted from the sample and sequenced on an Illumina platform, leading to

340,320,265 pair-ended reads (mean length 150bp). These metagenomics reads were then

mapped onto the genome sequence of *M. kamchatka*. Seven matching (100 % Identity) pair-ended

reads (hence 14 distinct reads) were detected, indicating the presence of virus particles in the

original sample, although at very low concentration. However, the very low probability of such

matches by chance ($p<10^{-63}$) together with the scattered distribution of these matches along the

viral genome, further demonstrate the presence of *M. kamchatka* in the original sample.

Virus Cloning

Fresh *A. castellanii* cells were seeded on a 12-well culture plate at a final concentration of

70,000cells/cm². Cell adherence was controlled under light microscopy after 45 minute and viral

particles were added to at a multiplicity of infection (MOI) around 50. After 1 h, the well was

washed 15 times with 3 mL of PPYG to remove any viral particle in suspension. Cells were then

recovered by gently scrapping the well, and a serial dilution was performed in the next three wells

by mixing 200µL of the previous well with 500µL of fresh medium. Drops of 0.5µL of the last

dilution were recovered and observed by light microscopy to confirm the presence of a unique *A.*

castellanii cell. The 0.5µL droplets were then distributed in each well of three 24-well culture plate.

Thousand uninfected *A. castellanii* cells in 500µL of PPYG were added to the wells seeded with a

single cell and incubated at 32°C until witnessing the evidence of a viral production from the

unique clone. The corresponding viral clones were recovered and amplified prior purification, DNA

extraction and cell cycle characterization by electron microscopy.

Virus mass production and purification

A total number of 40 T75 flasks were seeded with fresh *A. castellanii* cells at a final

concentration of 60,000 cells/cm². We controlled cell adherence using light microscopy after 45

minute and flasks were infected with a single clone of *M. kamchatka* at MOI=1. After 48h hours of

incubation at 32°C, we recovered cells exhibiting infectious phenotypes by gently scrapping the

flasks. We centrifuged for 10 min at 500×g to remove any cellular debris and viruses were pelleted

by a 1-hour centrifugation at 6,800xg. The viral pellet was then layered on a discontinuous cesium

chloride gradient (1,2g/cm² / 1,3g/cm² / 1,4g/cm² / 1,5g/cm²) and centrifuged for 20h at 103,000xg.

The viral fraction produced a white disk, which was recovered and washed twice in PBS and stored

at 4 °C or −80 °C with 7,5% DMSO.

Infectious cycle observations using TEM

Twelve T25 flasks were seeded with a final concentration of 80,000 cells/cm² in PPYG medium

containing antibiotics. In order to get a synchronous infectious cycle eleven flasks were infected by

freshly produced *M. kamchatka* at substantial MOI=40. The *A. castellanii* infected flasks were fixed

by adding an equal volume of PBS buffer with 5 % glutaraldehyde at different time points after the

infection : 1h pi, 2h pi, 3h pi, 4h pi, 5h pi, 6h pi, 7h pi, 8h pi, 9h pi, 10h pi and 25h pi. After 45min of

fixation at room temperature, cells were scrapped and pelleted for 5min at 500xg. Then cells were

resuspended in 1ml of PBS buffer with 2,5 % glutaraldehyde and stored at 4°C. Each sample was

coated in 1mm³ of 2 % low melting agarose and embedded in Epon-812 resin. Optimized osmium-

thiocarbohydrazide-osmium (OTO) protocol was used for staining the samples: 1h fixation in PBS

with 2 % osmium tetroxide and 1.5 % potassium ferrocyanide, 20 min in water with 1 %

thiocarbohydrazide, 30min in water with 2 % osmium tetroxide, overnight incubation in water with

1 % uranyl acetate and finally 30min in lead aspartate. Dehydration was made using an increasing

concentration of ethanol (50 %, 75 %, 85 %, 95 %, 100 %) and cold dry acetone. Samples were

progressively impregnated with an increasing mix of acetone and Epon-812 resin mixed with DDSA

0,34v/v and NMA 0,68v/v (33 %, 50 %, 75 % and 100%). Final molding was made using a hard

Epon-812 mix with DDSA 0,34v/v NMA 0,68v/v and 0,031v/v of DMP30 accelerator and hardened

in the oven at 60°C for 5 days. Ultrathin sections (90nm thick) were observed using a FEI Tecnai G2

operating at 200kV.

DNA Extraction

M. kamchatka genomic DNA was extracted from approximately 5 x 10^9 purified virus particles

using Purelink Genomic extraction mini kit according to the manufacturer's recommendation. Lysis

was performed with in a buffer provided with the kit and extra DTT at a final concentration of

1mM.

Genome sequencing and assembly

M. kamchatka genome was assembled using Spades (24) with a stringent K-mer parameter

using both various iteration steps (k = 21,41,61,81,99,127), the "-careful" option to minimize

number of mismatches in the final contigs, and the "-nanopore" option to use long reads.

Annotation of *Mollivirus sibericum* and *Mollivirus kamchatka*

A stringent gene annotation of *M. sibericum* was performed as previously described (13) using

RNA-seq transcriptomic data (6). Stranded RNA-seq reads were used to accurately annotate

protein-coding genes. Stringent gene annotation of *M. kamchatka* was performed w/o RNA seq

data but taking into account protein similarity with *M. sibericum*. Gene predictions were manually

curated using the web-based genomic annotation editing platform Web Apollo (25). Functional

annotations of protein-coding genes of both genomes were performed using a two-sided approach

as already previously described (13). Briefly, protein domains were searched with the CD-search

tool (26) and protein sequence searching based on the pairwise alignment of hidden Markov

models (HMM) was performed against the Uniclust30 database using HHblits tool (27). Gene

clustering was done using Orthofinder's default parameters (28) adding the "-M msa -oa" option.

Strict orthology between pairs of proteins was confirmed using best reciprocal blastp matches.

Selection Pressure Analysis

Ratios of non-synonymous (dN) over synonymous (dS) mutation rates for pairs of orthologous genes were computed from MAFFT global alignment (29) using the PAML package and codeml with the « model = 2 » (30). A strict filter was applyied to the dN/dS ratio: dN > 0, dS > 0, dS ≤ 2 and dN/dS ≤ 10. The computation of the Codon Adaptation Index (CAI) of both Mollivirus was performed using the cai tool from the Emboss package (31).

Metagenome sequencing, assembly and annotation

All metagenomic sample were sequenced with DNA-seq paired-end protocol on Illumina HiSeq platform at Genoscope producing 16 datasets of 2x150bp read length. Raw reads quality was evaluated with FASTQC (32). Identified contaminants were removed and remaining reads were trimmed on the right end using 30 as quality threshold with BBTools (33). Assemblies of filtered data sets were performed using MEGAHIT (34) with the following options: "--k-list 33,55,77,99,127 --min-contig-len 1000". All filtered reads were then mapped to the generated contigs using Bowtie2 (35) with the "--very-sensitive" option.

Availability of data

The *M. kamtchatka* annotated genome sequence is freely available from the public through the Genbank repository (URL://www.ncbi.nlm.nih.gov/genbank/) under accession number XXXXX.

Acknowledgements

We are deeply indebted to our volunteer collaborator Alexander Morawitz for collecting the Kamchatka soil sample. We thank N. Brouilly, F. Richard and A. Aouane (imagery platform, Institut de Biologie du Développement de Marseille Luminy) for their expert assistance. E Christo-Foroux is the recipient of a DGA-MRIS scholarship (201760003). This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No 832601) and from the FRM prize "Lucien Tartois" to C.

Abergel. The funding bodies had no role in the design of the study, analysis, and interpretation of

data and in writing the manuscript.

Competing interests

The authors declare that they have no competing interests

Figures

Fig 1. Ultrathin section TEM image of a neo-synthetized *M. kamchatka* particle in the cell

cytoplasm 7h post infection. The structure of the mature particles appear identical to that of *M.*

sibericum.

Fig 2. Ultrathin section TEM image of *A. castellanii* cell 7 to 10 hours post infection by *M.*

kamchatka. (A) Viral factory exhibiting fibrils (F), a nascent viral particle (V), and surrounding

mitochondria (M). Fragments of the ruptured nuclear membrane are visible as dark bead strings.

(B) Details of a nuclear membrane rupture through which fibrils synthetized in the nucleus (N) are

shed into the cytoplasm (C).

472 **Fig 3.** Distribution of the best-matching NR homologs of *M. kamchatka* predicted proteins. Best-
473 matching homologous proteins were identified using BLASTP (E value <10^{-5}) against the non-
474 redundant (NR) database (15) (after excluding *M. sibericum*). Green shades are used for
475 eukaryotes, red shades for viruses.

Fig 4. Eventual gene transfers from a pandoravirus to *M. kamchatka*. Both phylogenetic trees were

computed from the global alignments of orthologous protein sequences using MAFFT (29). IQtree

(36) was used to determine the optimal substitution model (options: « -m TEST » and « –bb

1000 »). (A) Protein mk_165 (no predicted function). (B) Predicted methyltransferase mk_92. Both

M. kamchatka protein sequences appear embedded within the pandoravirus trees. In (A), the long

branch leading to the *M. kamchatka* homolog suggests its accelerated divergence after an ancient

acquisition from a pandoravirus. In (B), the long branch leading to the *P. dulcis* homolog might

alternatively be interpreted as a non-orthologous replacement of the ancestral pandoravirus

version of the gene.

Fig 5. Genomic features of strain-specific ORFans. (A) Codon adaption index (CAI). (B) G+C content.

(C) Protein length. Box plots show the median, the 25th and 75th percentiles. P-value are

calculated using the Wilcoxon test.

Fig 6. Selection pressure among different classes of genes. Values of ω (i.e. dN/dS) were computed

from the alignments of homologous coding regions in *M. kamchatka* and *M. sibericum*. (A)

Distribution of calculated ω values (n=397). (B) Box plots of the ω ratio among ORFan genes

(n=243) and non ORFan genes (n=154). Box plots show the median, the 25th, and 75th percentiles.

All p-value are calculated using the Wilcoxon test.

505   **Fig 7.** Comparison of the mollivirus and pandoravirus core gene contents. (A) The distribution of

506   the protein clusters shared by all pandoraviruses (black), the two Molliviruses (pink), and by both

507   virus groups (super core genes) (blue). (B) Box plot of ω values calculated from the alignment of

508   molliviruses core genes (pink), and super core genes (blue). Box plots show the median, the 25th,

509   and 75th percentiles.

Fig 8. Distribution of different classes of genes along mollivirus genomes. (A) Variation of the gene

density as computed by the ggpplot2 "geom_density" function (37). Genes with best-matching

homologs in *A. castellanii* (in the NR database excluding mollivirus) are uniformly distributed (in

pink) in contrast to super core genes (in green) and strain-specific ORFans (in blue). (B) Cumulative

distribution of the above classes of genes using the same color code.

Fig 9. Distribution of single-copy *vs.* multiple copy genes along mollivirus genomes. Single copy

genes (in blue) in both strains are uniformly distributed in contrast to genes with paralogs (pink) in

at least one strain that cluster in the left half of the genomes. (A) *M. sibericum* (n=48). (B) *M.*

kamchatka (n= 46).

Fig 10. Phylogeny of DNA polymerase B of large and giant dsDNA viruses. This neighbor-joining tree

was computed (JTT substitution model, 100 resampling) on 397 amino acid positions from an

alignment of 42 sequences computed by MAFFT (29). Branches with boostrap values <60% were

collapsed.

Tables

Table 1. Status of the protein-coding genes unique to *M. kamchatka*

ORF ID	Predicted function	Putative evolutionary scenario
mk_25	None (ORFan)	*de novo* creation or loss in *M. sibericum*
mk_92	DNA methyltransferase	loss in *M. sibericum* (present in some pandoraviruses)
mk_93	Ring domain	loss in *M. sibericum* (present in some pandoraviruses)
mk_104	Ring domain	loss in *M. sibericum* (present in some pandoraviruses)
mk_127	None (ORFan)	*de novo* creation or loss in *M. sibericum*
mk_159	None	loss in *M. sibericum* (present in some pandoraviruses)
mk_165	None	HGT from Pandoravirus
mk_166	Peptidase	loss in *M. sibericum* (present in some pandoraviruses)
mk_172	None (ORFan)	*de novo* creation or loss in *M. sibericum*
mk_182	None (ORFan)	*de novo* creation or loss in *M. sibericum*
mk_231	None (ORFan)	*de novo* creation or loss in *M. sibericum*
mk_313	None (ORFan)	*de novo* creation or loss in *M. sibericum*
mk_369	None (ORFan)	*de novo* creation or loss in *M. sibericum*
mk_415	None (ORFan)	*de novo* creation or loss in *M. sibericum*
mk_441	None (ORFan)	*de novo* creation or loss in *M. sibericum*
mk_466	None (ORFan)	*de novo* creation or loss in *M. sibericum*
mk_467	None	loss in *M. sibericum* (HGT to Acanthamoeba)
mk_469	B1-1 like	loss in *M. sibericum* (present in Acanthamoeba)
mk_476	None (ORFan)	*de novo* creation or loss in *M. sibericum*
mk_478	None (ORFan)	*de novo* creation or loss in *M. sibericum*

535 **Table 2.** Status of the protein-coding genes unique to *M. sibericum*

ORF ID	Predicted function	**Putative evolutionary scenario**
ms_1	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_3	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_5	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_7	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_8	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_13	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_14	None	HGT to Acanthamoeba
ms_38	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_42	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_53	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_64	None	loss in *M. Kamchatka* (present in Noumeavirus)
ms_109	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_120	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_136	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_138	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_139	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_144	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_157	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_159	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_166	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_172	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_190	None	loss in *M. kamchatka* (present in some pandoraviruses)
ms_193	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_246	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_258	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_311	Zinc-finger domain	loss in *M. kamchatka (*present in *Gossypium hirsutum)*
ms_312	Zinc-finger domain	loss in *M. kamchatka* (present in *Cavenderia fasciculata)*
ms_313	None	loss in *M. kamchatka* (present in some pandoraviruses)
ms_464	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_465	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_479	DNA methyltransferase	loss in *M. kamchatka* (present in some pandoraviruses)
ms_494	None (ORFan)	*De novo* creation or loss in *M. kamchatka*
ms_495	None (ORFan)	*De novo* creation or loss in *M. kamchatka*

References

1. Shi T, Reeves RH, Gilichinsky DA, Friedmann EI, 1997. Characterization of Viable Bacteria from Siberian Permafrost by 16S rDNA Sequencing. Microb Ecol 33:169-179. https://doi.org/10.1007/s002489900019.

2. Vishnivetskaya T, Kathariou S, McGrath J, Gilichinsky D, Tiedje JM 2000. Low-temperature recovery strategies for the isolation of bacteria from ancient permafrost sediments. Extremophiles 4:165-173. https://doi.org/10.1007/s007920070031.

3. Graham DE, Wallenstein MD, Vishnivetskaya TA, Waldrop MP, Phelps TJ, Pfiffner SM, Onstott TC, Whyte LG, Rivkina EM, Gilichinsky DA, Elias DA, Mackelprang R, VerBerkmoes NC, Hettich RL, Wagner D, Wullschleger SD, Jansson JK. 2012. Microbes in thawing permafrost: the unknown variable in the climate change equation. ISME J 6:709-712. https://doi.org/10.1038/ismej.2011.163.

4. Yashina S, Gubin S, Maksimovich S, Yashina A, Gakhova E, Gilichinsky D. 2012. Regeneration of whole fertile plants from 30,000-y-old fruit tissue buried in Siberian permafrost. Proc Natl Acad Sci U S A 109:4008-4013. https://doi.org/10.1073/pnas.1118386109.

5. Legendre M, Bartoli J, Shmakova L, Jeudy S, Labadie K, Adrait A, Lescot M, Poirot O, Bertaux L, Bruley C, Couté Y, Rivkina E, Abergel C, Claverie JM. 2014. Thirty-thousand-year-old distant relative of giant icosahedral DNA viruses with a pandoravirus morphology. Proc Natl Acad Sci U S A 111:4274-4279. https://doi.org/ 10.1073/pnas.1320670111.

6. Legendre M, Lartigue A, Bertaux L, Jeudy S, Bartoli J, Lescot, M, Alempic JM, Ramus C, Bruley C, Labadie K, Shmakova L, Rivkina E, Couté Y, Abergel C, Claverie JM. 2015. In-depth study of Mollivirus sibericum, a new 30,000-y-old giant virus infecting Acanthamoeba. Proc Natl Acad Sci U S A 112:E5327-E5335. https://doi.org/10.1073/pnas.1510795112.

7. Abergel C, Legendre M, Claverie JM. The rapidly expanding universe of giant viruses: Mimivirus,

Pandoravirus, Pithovirus and Mollivirus. 2015. FEMS Microbiol Rev 39:779-796.

https://doi.org/10.1093/femsre/fuv037.

8. Levasseur A, Andreani J, Delerce J, Bou Khalil J, Robert C, La Scola B, Raoult D. 2016. Comparison

of a Modern and Fossil Pithovirus Reveals Its Genetic Conservation and Evolution. Genome Biol

Evol 8:2333-2339. https://doi.org/10.1093/gbe/evw153.

9. Andreani J, Aherfi S, Bou Khalil JY, Di Pinto F, Bitam I, Raoult D, Colson P, La Scola B. 2016.

Cedratvirus, a Double-Cork Structured Giant Virus, is a Distant Relative of Pithoviruses. Viruses

8:300. https://doi.org/10.3390/v8110300.

10. Bertelli C, Mueller L, Thomas V, Pillonel T, Jacquier N, Greub G. 2017. Cedratvirus lausannensis -

digging into Pithoviridae diversity. Environ Microbiol 19:4022-4034.

https://doi.org/10.1111/1462-2920.13813.

11. Quemin ER, Corroyer-Dulmont S, Baskaran A, Penard E, Gazi AD, Christo-Foroux E, Walther P,

Abergel C, Krijnse-Locker J. 2019. Complex membrane remodeling during virion assembly of the

30,000-year-old Mollivirus sibericum. J Virol 93(13). pii: e00388-19.

https://doi.org/10.1128/JVI.00388-19.

12. Philippe N, Legendre M, Doutre G, Couté Y, Poirot O, Lescot M, Arslan D, Seltzer V, Bertaux L,

Bruley C, Garin J, Claverie JM, Abergel C. 2013. Pandoraviruses: amoeba viruses with genomes up

to 2.5 Mb reaching that of parasitic eukaryotes. Science 341:281-286.

https://doi.org/10.1126/science.1239181.

13. Legendre M, Fabre E, Poirot O, Jeudy S, Lartigue A, Alempic JM, Beucher L, Philippe N, Bertaux

L, Christo-Foroux E, Labadie K, Couté Y, Abergel C, Claverie JM. 2018. Diversity and evolution of the

emerging Pandoraviridae family. Nat Commun 9:2285.

https://doi.org/10.1038/s41467-018-04698-4.

14. Legendre M, Alempic JM, Philippe N, Lartigue A, Jeudy S, Poirot O, Ta NT, Nin S, Couté Y,

Abergel C, Claverie JM. 2019. Pandoravirus celtis illustrates the microevolution processes at work

in the giant Pandoraviridae genomes. Front Microbiol 10:430. https://doi.org/

10.3389/fmicb.2019.00430.

15. NCBI Resource Coordinators. 2018. Database resources of the National Center for

Biotechnology Information. Nucleic Acids Res 46:D8-D13. https://doi.org/10.1093/nar/gkx1095.

16. Heidel AJ, Lawal HM, Felder M, Schilde C, Helps NR, Tunggal B, Rivero F, John U, Schleicher M,

Eichinger L, Platzer M, Noegel AA, Schaap P, Glöckner G. 2011. Phylogeny-wide analysis of social

amoeba genomes highlights ancient origins for complex intercellular communication. Genome Res

21:1882-1891. https://doi.org/10.1101/gr.121137.111.

17. Thomas V, Bertelli C, Collyn F, Casson N, Telenti A, Goesmann A, Croxatto A, Greub G. 2011.

Lausannevirus, a giant amoebal virus encoding histone doublets. Environ Microbiol 13:1454-1466.

https://doi.org/10.1111/j.1462-2920.2011.02446.x.

18. Maumus F, Blanc G. 2016. Study of gene trafficking between acanthamoeba and giant viruses

suggests an undiscovered fof amoeba-infecting viruses. Genome Biol

Evol 8:3351-3363. https://doi.org/10.1093/gbe/evw260.

19. Chelkha N, Levasseur A, Pontarotti P, Raoult D, Scola B, Colson P. 2018. A phylogenomic study of

Acanthamoeba polyphaga draft genome sequences suggests genetic exchanges with giant viruses.

Front Microbiol 9:2098. https://doi.org/10.3389/fmicb.2018.02098.

20. Duchêne S, Holmes EC. 2018. Estimating evolutionary rates in giant viruses using

ancient genomes. Virus Evol 4:vey006. https://doi.org/10.1093/ve/vey006.

21. Hughes AL, Irausquin S, Friedman R. 2010. The evolutionary biology of poxviruses. Infect Genet

Evol 10:50-59. https://doi.org/10.1016/j.meegid.2009.10.001.

22. San Martín C, van Raaij MJ. 2018. The so far farthest reaches of the double jelly roll capsid

protein fold. Virol J 15:181. https://doi.org/10.1186/s12985-018-1097-1.

23. Arslan D, Legendre M, Seltzer V, Abergel C, Claverie JM. 2011. Distant Mimivirus relative with a

larger genome highlights the fundamental features of Megaviridae. Proc Natl Acad Sci U S A

108:17486-17491. https://doi.org/10.1073/pnas.1110889108.

24. Bankevich A, Nurk S, Antipov D, Gurevich A, Dvorkin M, Kulikov AS, Lesin V, Nikolenko S, Pham

S, Prjibelski A, Pyshkin A, Sirotkin A, Vyahhi N, Tesler G, Alekseyev MA, Pevzner P. 2012. SPAdes: A

New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. J Comput Biol

19:455-477. https://doi.org/10.1089/cmb.2012.0021.

25. Dunn NA, Unni DR, Diesh C, Munoz-Torres M, Harris NL, Yao E, Rasche H, Holmes I, Elsik C,

Lewis S. 2019. Apollo: Democratizing genome annotation. PLoS Comput Biol 15: e1006790.

https://doi.org/10.1371/journal.pcbi.1006790.

26. Marchler-Bauer A, Derbyshire MK, Gonzales NR, Lu S, Chitsaz F, Geer LY, Geer RC, He J, Gwadz

M, Hurwitz DI, Lanczycki CJ, Lu F, Marchler GH, Song JS, Thanki N, Wang Z, Yamashita RA, Zhang D,

Zheng C, Bryant SH. 2015. CDD: NCBI's conserved domain database. Nucleic Acids Res 43:D222-

226. https://doi.org/10.1093/nar/gku1221.

27. Remmert M, Biegert A, Hauser A, Söding J. 2011. HHblits: lightning-fast iterative

protein sequence searching by HMM-HMM alignment. Nat Methods. 9:173-175.

https://doi.org/10.1038/nmeth.1818.

28. Emms DM, Kelly S. 2015. OrthoFinder: solving fundamental biases in whole genome

comparisons dramatically improves orthogroup inference accuracy. Genome Biol 16:157.

https://doi.org/10.1186/s13059-015-0721-2.

29. Katoh K, Misawa K, Kuma K-I, Miyata T. 2002. MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. Nucleic Acids Res 30:3059-3066.

https://doi.org/10.1093/nar/gkf436.

30. Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol 24:1586-

1591. https://doi.org/10.1093/molbev/msm088.

31 . Rice P, Longden I, Bleasby A. 2000. EMBOSS: the European molecular biology open software

suite. Trends Genet 16:276-277. https://doi.org/10.1016/s0168-9525(00)02024-2.

32. Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. Available

online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc .

33. Bushnell B, Rood J, Singer E. 2017. BBMerge – Accurate paired shotgun read merging via

overlap. PLoS ONE 12: e0185056. https://doi.org/10.1371/journal.pone.0185056.

34. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. 2015. MEGAHIT: An ultra-fast single-node solution

for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics

31:1674-1676. https://doi.org/10.1093/bioinformatics/btv033.

35. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods

9:357-359. https://doi.org/10.1038/nmeth.1923.

36. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A fast and effective stochastic

algorithm for estimating maximum likelihood phylogenies. Mol Biol Evol 32:268-274.

https://doi.org/10.1093/molbev/msu300.

37. Wickham H. ggplot2: Elegant Graphics for Data Analysis, p 33-74. 2016. Springer-Verlag New

York.