ESI mutagenesis: A one-step method for introducing point mutations into bacterial artificial chromosome transgenes.

Arnaud Rondelet[1]*, Andrei Pozniakovsky[2]*, Marit Leuschner[2], Ina Poser[2], Andrea Ssykor[2], Julian Berlitz[1], Nadine Schmidt[1], Anthony A Hyman[2], Alexander W Bird[1]

[1]Max Planck Institute of Molecular Physiology, Dortmund, Germany
[2]Max Planck Institute of Molecular Cell Biology and Genetics, Dresden, Germany

*authors contributed equally

Abstract

Bacterial artificial chromosome (BAC)-based transgenes have emerged as a powerful tool for controlled and conditional interrogation of protein function in higher eukaryotes. While homologous recombination-based recombineering methods have streamlined the efficient integration of protein tags onto BAC transgenes, generating precise point mutations has remained less efficient and time-consuming. Here we present a simplified method for inserting point mutations into BAC transgenes requiring a single recombineering step followed by antibiotic selection. This technique, which we call ESI (**E**xogenous/**S**ynthetic **I**ntronization) mutagenesis, relies on co-integration of a mutation of interest along with a selectable marker gene, the latter of which is harboured in an artificial intron adjacent to the mutation site. Cell lines generated from ESI-mutated BACs express the transgenes equivalently to the endogenous gene, and all cells efficiently splice out the synthetic intron. Thus, ESI-mutagenesis provides a robust

and effective single-step method with high precision and high efficiency for mutating BAC transgenes.

Introduction

The ability to precisely query functional hypotheses of protein function in

cells requires the capacity to express rationally mutated proteins from genes under

their native physiological regulation. Traditionally, transgenes expressed in higher

eukaryotes are derived from cDNAs, and thus lack native cis-regulatory elements

or alternative splicing isoforms, often resulting in overexpression and

deregulation artefacts. This may hinder proper phenotypic functional

characterization of mutated genes, as well as the determination of the precise

localization and interaction partners of the protein products. While the

development of sequence-specific nucleases has enabled mutation of specific

genes at their endogenous loci(Gaj *et al.*, 2013; Ran *et al.*, 2013), mutations in

essential genes that are lethal will prevent growth and recovery of viable cells.

Additionally, deleterious mutations are prone to accumulate suppressive changes

in chromosome integrity or gene expression during the procedure of selecting and

expanding cells for analysis, particularly in the genomically instable cancer cell

lines frequently employed.

The use of bacterial artificial chromosomes (BACs) as transgenes largely

overcomes these limitations. BACs are bacterial vectors containing fragments of

an eukaryotic genome that are large enough to encode an entire gene in its

genomic context, including *cis*-regulatory elements. Transgenes expressed from

BACs thus deliver near-physiological expression in eukaryotic cells(Kittler *et al.*,

2005; Bird and Hyman, 2008; Poser *et al.*, 2008; Bird *et al.*, 2011). BAC transgenes

modified to be RNAi-resistant allow for conditional exposure of recessive

mutations through selective depletion of endogenous protein(Kittler *et al.*, 2005;

Bird and Hyman, 2008; Ding *et al.*, 2009; Bird *et al.*, 2011; Scolz *et al.*, 2012; Zheng

et al., 2014).

Because of their size, BACs do not lend themselves to restriction/ligation-based

modification techniques. Instead, homologous recombination-based

recombineering techniques are used to engineer BACs in *Escherichia coli* (Murphy,

1998; Zhang *et al.*, 1998; Yu *et al.*, 2000). Recombineering utilizes phage proteins

(Redα, β and γ from phage λ; or RecE and RecT from the Rac prophage) to promote

homologous recombination, facilitating a wide variety of DNA modifications. The

high efficiency of the single-step recombineering procedure to insert a protein tag

attached to a selectable marker gene on a DNA vector has allowed its adaptation to

genome-scale high-throughput pipelines(Sarov *et al.*, 2006; Poser *et al.*, 2008;

Sarov *et al.*, 2012; Hasse *et al.*, 2016; Sarov *et al.*, 2016). The subsequent generation

of transgenic cell lines based on these libraries has further been used to precisely

determine the cellular localisation and the quantitative interactome of more than

a thousand proteins (Hubner *et al.*, 2010; Hutchins *et al.*, 2010; Hein *et al.*, 2015).

On the contrary, current techniques to introduce point mutations in BACs still

require either a counterselection-based two-step procedure(Bird *et al.*, 2011;

Wang *et al.*, 2013) or a lower efficiency one-step procedure requiring extensive

PCR-screening (Lyozin *et al.*, 2014). Here we present a simple and efficient one-

step procedure to introduce point mutations in BAC transgenes, harnessing

introns to carry selectable markers, which reduces the time and cost of generating

mutagenized constructs.

Modifying intronic sequences is an attractive approach to modifying or

gaining gene functionality with minimal perturbation of the host gene protein

product, and has met use in a variety of applications such as conditional knock-out

and mutant constructs, or miRNA expression(Gu *et al.*, 1994; Kaulich *et al.*, 2015;

Wassef *et al.*, 2017). Synthetic introns have been designed based on the four core

splicing signals necessary for spliceosome processing: two splice sites (SS) located

at the 5' and 3' intron boundaries (5'SS and 3'SS, respectively), a branchpoint

sequence (BP) located approximately 25 nucleotides upstream of the 3'SS, and a

polypirimidine tract (PPT) directly upstream of the 3'SS (Figure 1A) (Lin *et al.*,

2003; Wang and Burge, 2008; Mercer *et al.*, 2015). Most of the sequence

determinants for splicing are thus located within the intron. In most eukaryotes,

the consensus sequences of the core splicing elements located within exons are

limited to a (C/A) A G sequence directly upstream of the intron and a G

downstream of the intron (Figure 1 A) (Zhang, 1998). Synthetic introns designed

from these minimal core splicing signals are indeed efficiently spliced out of an

exon, and have been used in multiple applications, such as making conditional

knockouts, or expressing shRNAs or miRNAs(Lin *et al.*, 2003; Greber and

Fussenegger, 2007; Lin and Ying, 2013; Seyhan, 2016; Guzzardo *et al.*, 2017).

Here we present a one-step BAC recombineering procedure in which point

mutations are introduced into a BAC along with a synthetic intron. An antibiotic-

resistance gene located within the intron allows the selection of positive clones

after the single recombineering step in bacteria, and is seamlessly spliced out of

mRNA sequences of the harbouring transgene in eukaryotic cells. This procedure

significantly increases the efficiency and decreases the screening time of BAC

mutagenesis.

RESULTS

ESI-mutagenesis: a one-step BAC recombineering strategy for introduction

of point mutations.

The ESI-mutagenesis strategy is outlined in Figure 1. First, a synthetic

intron "cassette" is amplified by PCR for use in the recombineering reaction (Fig

1B). This synthetic intron cassette contains the core intron-based splicing signal

sequences, as well as an antibiotic resistance gene. We have created multiple

cassettes with varying bacterial and eukaryotic selectable markers, presented in

Figure S1. In order to become a functional intron, the synthetic intron cassette

must be recombined into a site in the exon where it is directly bordered by an

upstream (C/A) A G sequence and a downstream G (Figure 1A, and C). While this

(C/A) A G | G sequence would occur every 128 nucleotides assuming random

distribution, similar sequences can be mutated to this sequence at amino acid

wobble positions during cassette amplification (see Figure S2 for an example),

significantly increasing the frequency of useable integration sites. Furthermore,

this consensus is not a strict requirement, and a variety of sequences are found in

cells and predicted to function(Zhang, 1998; Nguyen *et al.*, 2018; Ohno *et al.*, 2018).

Therefore, it is unlikely that a potential synthetic intron integration site will not be

found within the useable vicinity of a mutation site of interest. In addition, if the

desired mutation is close enough to an existing intron, the antibiotic resistance

cassette can be directly targeted to the endogenous intron by generating only one

new splice site, with the opposite homology arms targeting intronic sequences (see

Figure 1C, right side). This is preferable when possible, because not only does it

maintain more closely the native intron/exon structure, but it also avoids the

creation of unusually small exons (less than ~50bp), which may be prone to exon

skipping (Dominski and Kole, 1991).

An example of how to design ESI-mutagenesis is presented in supplementary

Figure S2. Once a target site for synthetic intron insertion near the mutation site is

identified, primers are designed to amplify the synthetic intron cassette and

containing 5'-extended 50 bp arms homologous to sequences directly surrounding

the insertion site. One or both homology arms contain the mutation(s) to be

integrated. The cassette is amplified by PCR and the resulting product is inserted

into the target BAC via recombineering (Figure 1 C). Correct recombination events

are selected using the antibiotic resistance encoded within the synthetic intron. A

protocol for the ESI mutagenesis recombineering procedure as well as generating

mammalian transgenic cell lines is provided in Figure S3.

Upon transfection of the modified BAC into eukaryotic cells of interest, cells that

have integrated the BAC in their genome may also be selected using the antibiotic

resistance encoded within the synthetic intron. If desired, the antibiotic resistance

gene may be removed by application of Cre recombinase via loxP sites located

within the cassette (Figure S1). Once the BAC transgene is transcribed the

synthetic intron is spliced out leading to the formation of a messenger RNA

carrying the desired mutation.

ESI mutated BAC transgenes yield proteins with the expected size and

localization

To confirm that BAC transgenes containing synthetic introns yielded proteins of the correct size and localisation when expressed in human cells, we started with a panel of 10 GFP-tagged BAC transgenes, and modified each using our ESI-mutagenesis strategy to introduce mutations to render them resistant to specific siRNA targeting sequences (Table S1). We then generated Hela cell lines stably expressing either the "parental" GFP-tagged genes, or the "ESI-mutated" (i.e. RNAi-resistant) GFP-tagged genes (Figure 2A). Because these mutations should not change the corresponding protein sequence, this allowed us to directly compare the expression and localization of the mutated transgenes to that of the parental, while offering an assay for the presence and function of the introduced mutation (RNAi resistance). Depending on the location of the mutation with respect to an existing intron, we either targeted the synthetic intron along with the desired mutation into an exon (4/10 BACs) (Figure 1 C left panel, Figure 2B), or made use of an existing exon-intron border to introduce the desired mutation and the antibiotic resistance (6/10 BACs) (Figure 1 C right panel, Figure 2B). All stably integrated "ESI-mutated" transgenes led to the expression of proteins of sizes identical to the one expressed from the corresponding parental transgenes (Figure 2 C). Moreover, we could not observe any differences in the cellular localisation of proteins expressed from "ESI-mutated" transgenes as compared to those expressed from parental transgenes (Figure 2 D). We next performed siRNA experiments on six of the parental / "ESI-mutated" BAC lines pairs to confirm functionality of the mutations. When the parental BAC-GFP lines were treated with the siRNA against the tagged transgene, we observed a reduction in the corresponding mRNA levels comparable to that observed in Hela wild type cells

(Figure 3 A). In contrast, ESI-mutated BAC lines did not show any reduction in the

corresponding mRNA levels upon siRNA treatment. We then analysed RNAi-

resistance on the protein level for two of the ESI-mutated BAC lines for which we

had the corresponding specific antibody. Both CEP135 and AURKB ESI-mutated

BAC clonal lines showed expression of their respective GFP-tagged transgenes at a

level similar to the corresponding endogenous proteins (Figure 3 B), and

treatment of both ESI-mutated lines with the corresponding siRNA resulted in

depletion of the endogenous protein while the transgene could still be detected

(Figure 3 B).

To further demonstrate that ESI-mutagenesis allows the introduction of functional

mutations in a BAC transgene, we used the procedure to mutate two EB1-

interacting motifs (SxIP) of the microtubule-associated protein GTSE1 to abolish

its interaction with the plus end tracking protein EB1(Honnappa *et al.*, 2009; Scolz

et al., 2012). In this example, we started with a GFP-tagged GTSE1 BAC, integrated

the synthetic intron between the two sites to be mutated, and included the two

mutations (SxNN) within the 5' and 3' homology arms (Figure S2). U2OS cells were

generated stably expressing the BAC transgene and analysed for GTSE1 expression

and interaction with EB1. The ESI-mutated BAC yielded a protein of the same size

as GTSE1 endogenously tagged with GFP, as expected(Figure 3 C, left panel)

(Bendre *et al.*, 2016). However, the mutated protein was no longer efficiently

pulled-down by GST-EB1 when compared to the endogenous GTSE1 or GTSE1-GFP

(Figure 3 C), verifying that the mutation was present.

A functional ESI-mutated BAC transgene is expressed in all cells of a clonal line

We showed above that ESI-mutated BACs yield proteins of the expected size and the correct cellular localisation, and that mutation may efficiently confer RNA-resistance or loss of interaction partner when analysed at the cell-population level. We next further assessed whether an ESI-mutated transgene was functionally expressed, i.e. able to sustain the cellular functions of its endogenous counterpart, uniformly in all the cells of a population. This is critical for functional and phenotypic analysis of mutants on the single cell level. To address this question, we ESI-mutated the gene encoding Aurora A kinase (AURKA) on a GFP-tagged BAC transgene to render it resistant to RNAi, transfected it into U2OS cells, and selected a cell clone (U2OS AURKA-GFP siRES (ESI)) that expressed Aurora A-GFP at the same level as the endogenous Aurora A (Figure 4 A, sixth lane). Treatment of this line with siRNA against AURKA led to the depletion of the endogenous Aurora A, while the GFP-tagged Aurora A remained expressed (Figure 4 A, fifth and sixth lanes, lowest and uppermost band respectively) and correctly localised to the centrosomes and the spindle (Figure 4 B and C). U2OS cells depleted of Aurora A accumulate in prometaphase and show smaller bipolar spindles (Marumoto et al JBC 2003, Bird and Hyman 2008) (Figure 4 D, E, and F). In contrast, cells containing the RNAi-resistant Aurora A-GFP ESI maintained a normal distribution of mitotic phases, similar to control-treated cells (Figure 4 D). Furthermore, we could not observe any differences in the length of the metaphase spindle between U2OS treated with control-siRNA and the U2OS AURKA-GFP siRES (ESI) clone treated with the siRNA against AURKA (Figure 4 E and F). This result indicates that all cells depleted for the endogenous Aurora A express a functional ESI mutated Aurora A-

GFP ESI transgene. Taken together, these data show that all cells of the U2OS

AURKA-GFP siRES (ESI) clone uniformly express a functional Aurora A-GFP

protein.

DISCUSSION

We have shown that by co-integrating a selectable marker located within an intron,

site-directed mutations can be quickly and efficiently introduced into BAC

transgenes using a one-step recombineering procedure. This recombineering

method is thus analogous to that for integrating protein tags: a single PCR reaction

to generate a cassette with homology arms followed by recombineering and

selection. The high efficiency (>90%) of the latter has facilitated high-throughput

genome-scale application of protein tagging(Poser *et al.*, 2008; Sarov *et al.*, 2012;

Hein *et al.*, 2015; Hasse *et al.*, 2016; Sarov *et al.*, 2016). While we have focused on

applications in human cell lines, this method would also be applicable to BAC or

fosmids commonly used as transgenes in other model organisms.

Critically, BAC transgenes ESI-mutated to be RNAi resistant yield proteins of the

correct size and localization that can functionally rescue the phenotypes observed

upon depletion of the endogenous counterpart. Our investigation of transgenic

lines at the single-cell level suggest that the synthetic intron is correctly spliced out

in all the cells of a population. Furthermore, although we formally describe the ESI-

mutagenesis technique here for the first time, we have previously successfully

employed it to generate mutations in BAC transgenes, for example, to functionally

analyze a microcephaly-associated mutation in the centriolar protein CPAP at the

single cell level in HeLa cells and neural progenitor cells(Zheng *et al.*, 2014; Gabriel

et al., 2016).

We cannot exclude that a small portion of synthetic introns within individual cells

are misspliced, yet without detectable consequences in our functional assays.

Indeed, although the core essential sequences for defining introns and splice sites

are located primarily within the intron and a few nucleotides into the exons, the

precise regulation of efficient splicing is a more complex process and impacted by

additional sequence information, including that within exons (Chasin, 2007).

Nevertheless, mis-spliced mRNAs are probably minimal and degraded by

nonsense-mediated decay.

One of our original concerns in this study was that the new synthetic intron or the

gene within may impact expression of the transgene to an unuseful extent. While

introns have been well characterized to generally increase gene expression(Nott

et al., 2003; Shaul, 2017), nested intronic genes found natively in eukaryotic

genomes generally impart a negative effect on the "host" gene expression(Yu *et al.*,

2005; Kumar, 2009). Thus it is possible that expression of a transgene could be

impacted by synthetic intron-mutagenesis. While we did not observe gross

changes in gene expression in the cell lines generated here, this will likely be highly

context-dependent (gene, location of intron, organism, tissue, etc) and we have

observed a case where a nested antibiotic resistance gene in a synthetic intron

directly integrated into the genome via CAS9 has decreased host gene expression

(unpublished results). We have thus also designed a version of the synthetic intron

cassette for BAC recombineering in which the antibiotic resistance only carries a

bacterial promoter, eliminating the necessity to recombine out with Cre. In the end,

because BACs integrate randomly into the genome by standard transfection

procedures, clonal variations in expression (+/- 2-fold) in transfected cell lines,

likely dependent on position effect and copy number, are anyway observed. Such

variations could compensate for synthetic intron effects, and it is thus good

practice, prior to phenotypic functional analysis, to select for clonal cell lines

expressing BAC transgenes at endogenous levels.

The general strategy outlined here of harbouring a selectable marker gene in an

artificial intron adjacent a mutation could in principle be used to increase the

efficiency of isolating point mutations via Cas9-induced homologous

recombination directly into the genome. This would particularly be useful in hard-

to-transfect cell lines where extensive screening may be necessary without a

selection step. In our preliminary studies applying this ESI-mutagenesis strategy

to direct Cas9-mediated human genome modification, we are indeed able to

recover modified cell lines at high frequency. We also found, however, more readily

observable impacts on gene expression from the synthetic intron when located in

the native gene, which could be alleviated by removing the antibiotic resistance

gene after Cre application. While further development of the synthetic intron

cassette design/strategy towards direct genome modification holds promise for

high efficiency selection of Cas9-mediated site-directed mutation, sensitive

applications (i.e. anything to be used in a clinical setting) would not be advisable

due to potential changes induced by the synthetic intron.

The use of BAC transgenes remains a powerful genetic tool for exploration of

protein function. The ESI mutagenesis method described here allows for

straightforward and rapid site-directed mutagenesis of BAC transgenes to further

improve their ease of use.

Methods and Materials

Bacterial Artificial Chromosomes and recombineering

All BACs used and generated, including modified sequences, are presented in

Table S1. LAP- and NLAP-tagged "parental" BACs were generated by

recombineering as described in Poser et al 2008. "ESI-mutated" BACs were

generated from "parental" BACs following the procedure outlined in Figure S3.

Antibodies

Mouse anti-GFP (11 814 460 001, Roche), rabbit anti-GTSE1 (custom generated;

described in Scolz *et al.*, 2012), and rabbit anti-Aurora kinase B (ab2254, Abcam)

were used for Western blots. Mouse anti-α-tubulin (DM1α, Sigma-Aldrich), rabbit

anti-Cep135 (custom generated, described in Bird and Hyman, 2008), goat anti-

GFP (MPI Dresden; described in Poser *et al.*, 2008), and goat anti-Aurora A (sc-

27883, Santa Cruz) were used for both Western Blot and immunofluorescence.

Donkey anti-goat Alexa488 (Jackson Immunoresearch; 705 545 147), donkey

anti-mouse Alexa594 (Bethyl; A90-337D4), and donkey anti-rabbit Alexa650

(Bethyl; A120-208D5) were used in immunofluorescence. Donkey anti-goat HRP

(Santa Cruz; SC-2020), sheep anti-mouse HRP (Amersham; NXA931-1ml), and

donkey anti-rabbit HRP (Amersham; NXA934-1ml) were used for Western Blots.

Cell Lines and Cell Culture

U2OS, HeLa cells, and derivatives were grown at 5% CO2 and 37 °C in DMEM (PAN

Biotech) supplemented with 10% filtered Fetal Bovine Serum (FBS, Gibco), 2 mM

L-glutamine (PAN Biotech), 0.1 mg/mL streptomycin and 100 U/ml penicillin

(PAN Biotech, Pen/Strep mix). BACs were transfected into cells using the

Effectene kit (Qiagen) following the manufacturer's instructions, and stable cell

lines were selected on antibiotics.

RNAi

siRNA sequences used are presented in Table S1. siRNAs were purchased from

Thermofisher/Ambion. Unless specified otherwise, control siRNA was Silencer

negative control No. 2 (Thermofisher). A reverse transfection approach was used

to deplete AurkA and CEP135, using 50 nM and 40 nM siRNA, respectively. The

transfection procedure for a 24-well plates with a final volume of 500 µL is

described. If required, the procedure was scaled up. The siRNA and 2.5 µL

Oligofectamine (Invitrogen) were prepared separately in a final volume of 50 and

15 µL OptiMEM (Invitrogen), respectively. After 5 min incubation at RT, the siRNA

and transfection reagent were mixed and further incubated 20 min at RT.

Meanwhile, HeLa cells, U2OS cells or their derivatives were seeded at the desired

confluency into a 24 well plates containing or not a coverslip. The transfection mix

was finally added onto the cells and the volume adjusted to 500 µL with pre-

warmed medium. The medium was changed 7 to 8 hours post-transfection. For

CEP135 depletion, cells were harvested for Western Blot 48 hours post-

transfection. For AurkA depletion, 12 hours post-transfection cells were

synchronized with 3.3 µM Nocodazole for 18 hours, and harvested for Western

Blot. Alternatively, cells seeded on coverslips were fixed and processed for

microscopy 30 hours post-transfection. A forward transfection approach was

used to deplete AurkB using 50nM siRNA. Day before transfection, cells were

seeded at ~40 % confluency in a 3.5 cm dish. The siRNA and 5 µL of Oligofectamine

(Invitrogen) were prepared separately into 200 µL OPTIMEM. After 5 min

incubation at RT siRNA and Oligofectamine solutions were mixed and further

incubated 20 min at RT. Meanwhile, cells were washed twice with PBS and 1.6 mL

OPTIMEM were added on top of them. The transfection mix was then added onto

the cells. Cells were harvested for Western Blot 48 hours post-transfection. In

experiments measuring mRNA levels in HeLa cells, a control scrambled RNA

(customed, Ambion, sense: 5'-UUCUCCGAACGUGUCACGUtt-3') was used, and

siRNA was transfected using Interferin (Polyplus Transfection) and 100nM siRNA

per well of a 6-well plate in forward or reverse transfection (140,000 cells).

Duration of the depletion is indicated for each gene in Table S1.

Western Blot

Cells were lysed on ice for 10 to 15 min in RIPA buffer or cell lysis buffer (50 mM

Na2HPO4, 150 mM NaCl, 10% glycerol, 1% Triton X-100, 1 mM EGTA, 1.5 mM

MgCl2, 50 mM HEPES pH 7.2, 1 mM DTE) both supplemented with 2X protease

inhibitor mix (SERVA). Debris were eliminated by centrifugation at 4 °C at max-

speed in a benchtop centrifuge. Samples were prepared in Laemmli buffer and run

on SDS-PAGE before transfer onto a nitrocellulose membrane. 5% milk (Carl Roth)

in PBS Tween20 0.1% (SERVA electrophoresis) was used to block the membrane

before incubation with the primary and secondary antibodies diluted in the same

blocking solution. Membranes were incubated with ECL reagent (GE Healthcare)

before development onto Hyperfilm (Amersham) or imaging on the ChemiDoc™

MP imaging system (BioRad). Digital data were obtained by scanning Hyperfilms

or using the ImageLab software (BioRad). Fiji was used to adjust levels, generate

8-bit tiffs, and crop images.

qPCR analysis

cDNA was isolated from 250,000 cells per condition using the Cells-to-cDNA II Kit

(Ambion AM1722) following the manufacturer's two-step RT-PCR protocol.

qPCR was performed with SYBR Green (ThermoFisher) on a Mx3000 qPCR

Machine (Stratagene). Primers are indicated in Table S1.

Pull-downs

Cells from a confluent 15 cm dish were lysed for 15 min on ice in 2mL cell lysis

buffer (50 mM Na$_2$HPO$_4$, 150 mM NaCl, 10% glycerol, 1% Triton X-100, 1 mM

EGTA, 1.5 mM MgCl$_2$, 50 mM HEPES pH 7.2, 1 mM DTE) supplemented with 2X

protease inhibitor mix (SERVA) and 1mM DTE (15min on ice). Debris were

removed by centrifugation at max-speed at 4 °C in a benchtop centrifuge. Total

protein concentration was measured by Bradford. Equivalent amounts of cell

lysate were incubated 20 hours at 4 °C on GST or GST-EB1 beads. Beads were

washed thrice with 1 mL cell lysis buffer supplemented with 2X Protease inhibitor.

Proteins were eluted with 45 μL Laemmli 2x.

Immunofluorescence

For most experiments, cells were seeded onto coverslips and fixed in ice-cold

methanol for 12 min at -20 °C. Cells were blocked at RT with 5% BSA (Sigma)

dissolved into PBS. For images of Aurora B-GFP and CHMP4B-GFP cell lines, cells

were fixed in 3% PFA supplemented with 5 mM EGTA, 1mM $MgCl_2$ and 2%

sucrose, they were washed, then permeabilized using PBS/0.1% Triton X-100 and

eventually blocked with PBS/0.2% Fish Skin Gelatin (Sigma, Germany). Primary

antibodies were diluted into 5% BSA in PBS and incubated onto coverslips for 1 h

at 37 °C in a wet chamber. The same procedure was applied for secondary antibodies.

Coverslips were mounted using Prolong Gold antifade mounting with DAPI (Molecular

Probes and Thermo Fisher Scientific). Cells were washed thrice with PBS between

every step.

Microscopy

Immunofluoresence images for the AurkA experiments were acquired using a

spinning-disk confocal system (Marianas, Intelligent Imaging Innovation, 3i) built

on an Axio Observer Z1 microscope (Zeiss) with an ORCA-Flash 4.0 camera

(Hamamatsu Photonics). Images were taken with a 63× 1.4 NA Plan-Apochromat

oil objective (ZEISS). The spindle size was measured using the CEP135 signal in

the Slidebook software (Marianas, Intelligent Imaging Innovation, 3i, V5.5 or

later).

Experiments assessing the localization of the GFP transgenes, were all performed

on live cells at the exception of the AurkB and CHMP4B transgenes. Live cells were

imaged in CO2-independent visualization media (Gibco). Images were acquired

using a DeltaVision imaging system (GE Healthcare) with an sCMOS camera (PCO

Edge 5.5) and a 60x 1.42NA Plan Apo N UIS2 objective (Olympus).

BIBLIOGRAPHY

Bendre, S., Rondelet, A., Hall, C., Schmidt, N., Lin, Y.-C., Brouhard, G. J., and Bird, A.
W. (2016). GTSE1 tunes microtubule stability for chromosome alignment and
segregation by inhibiting the microtubule depolymerase MCAK. The Journal of
Cell Biology *215*, 631–647.

Bird, A. W., and Hyman, A. A. (2008). Building a spindle of the correct length in
human cells requires the interaction between TPX2 and Aurora A. The Journal of
Cell Biology *182*, 289–300.

Bird, A. W., Erler, A., Fu, J., Hériché, J.-K., Maresca, M., Zhang, Y., Hyman, A. A., and
Stewart, A. F. (2011). High-efficiency counterselection recombineering for site-
directed mutagenesis in bacterial artificial chromosomes. Nat Meth *9*, 103–109.

Chasin, L. A. (2007). Searching for splicing motifs. Adv. Exp. Med. Biol. *623*, 85–
106.

Ding, L. *et al.* (2009). A genome-scale RNAi screen for Oct4 modulators defines a
role of the Paf1 complex for embryonic stem cell identity. Cell Stem Cell *4*, 403–
415.

Dominski, Z., and Kole, R. (1991). Selection of splice sites in pre-mRNAs with
short internal exons. Molecular and Cellular Biology *11*, 6075–6083.

Gabriel, E. *et al.* (2016). CPAP promotes timely cilium disassembly to maintain
neural progenitor pool. Embo J *35*, 803–819.

Gaj, T., Gersbach, C. A., and Barbas, C. F., III (2013). ZFN, TALEN, and CRISPR/Cas-
based methods for genome engineering. Trends in Biotechnology, 1–9.

Greber, D., and Fussenegger, M. (2007). Multi-gene engineering: Simultaneous
expression and knockdown of six genes off a single platform. Biotechnol. Bioeng.
96, 821–834.

Gu, H., Marth, J. D., Orban, P. C., Mossmann, H., and Rajewsky, K. (1994). Deletion
of a DNA polymerase beta gene segment in T cells using cell type-specific gene
targeting. Science *265*, 103–106.

Guzzardo, P. M., Rashkova, C., Santos, dos, R. L., Tehrani, R., Collin, P., and
Bürckstümmer, T. (2017). A small cassette enables conditional gene inactivation
by CRISPR/Cas9. Nature Publishing Group *7*, 16770.

Hasse, S., Hyman, A. A., and Sarov, M. (2016). TransgeneOmics â€" A transgenic
platform for protein localization based function exploration. Methods *96*, 69–74.

Hein, M. Y. *et al.* (2015). A Human Interactome in Three Quantitative Dimensions
Organized by Stoichiometries and Abundances. Cell *163*, 712–723.

Honnappa, S. *et al.* (2009). An EB1-Binding Motif Acts as a Microtubule Tip
Localization Signal. Cell *138*, 366–376.

Hubner, N. C., Bird, A. W., Cox, J., Splettstoesser, B., Bandilla, P., Poser, I., Hyman,
A., and Mann, M. (2010). Quantitative proteomics combined with BAC
TransgeneOmics reveals in vivo protein interactions. The Journal of Cell Biology
189, 739–754.

Hutchins, J. R. A. *et al.* (2010). Systematic Analysis of Human Protein Complexes
Identifies Chromosome Segregation Proteins. Science *328*, 593–599.

Kaulich, M., Lee, Y. J., Lonn, P., Springer, A. D., Meade, B. R., and Dowdy, S. F.
(2015). Efficient CRISPR-rAAV engineering of endogenous genes to study protein
function by allele-specific RNAi. Nucleic Acids Research.

Kittler, R., Pelletier, L., Ma, C., Poser, I., Fischer, S., Hyman, A. A., and Buchholz, F.
(2005). RNA interference rescue by bacterial artificial chromosome transgenesis
in mammalian tissue culture cells. Proceedings of the National Academy of
Sciences *102*, 2396–2401.

Kumar, A. (2009). An overview of nested genes in eukaryotic genomes.
Eukaryotic Cell *8*, 1321–1329.

Lin, S.-L., and Ying, S.-Y. (2013). Gene silencing in vitro and in vivo using intronic
microRNAs. Methods Mol Biol *936*, 209–229.

Lin, S.-L., Chang, D., Wu, D.-Y., and Ying, S.-Y. (2003). A novel RNA splicing-
mediated gene silencing mechanism potential for genome evolution. Biochem
Biophys Res Commun *310*, 754–760.

Lyozin, G. T., Bressloff, P. C., Kumar, A., Kosaka, Y., Demarest, B. L., Yost, H. J.,
Kuehn, M. R., and Brunelli, L. (2014). Isolation of rare recombinants without
using selectable markers for one-step seamless BAC mutagenesis. Nat Meth *11*,
966–970.

Mercer, T. R., Clark, M. B., Andersen, S. B., Brunck, M. E., Haerty, W., Crawford, J.,
Taft, R. J., Nielsen, L. K., Dinger, M. E., and Mattick, J. S. (2015). Genome-wide
discovery of human splicing branchpoints. Genome Research *25*, 290–303.

Murphy, K. C. (1998). Use of bacteriophage lambda recombination functions to
promote gene replacement in Escherichia coli. Journal of Bacteriology *180*,
2063–2071.

Nguyen, H., Das, U., Wang, B., and Xie, J. (2018). The matrices and constraints of
GT/AG splice sites of more than 1000 species/lineages. Gene *660*, 92–101.

Nott, A., Meislin, S. H., and Moore, M. J. (2003). A quantitative analysis of intron
effects on mammalian gene expression. Rna *9*, 607–617.

Ohno, K., Takeda, J.-I., and Masuda, A. (2018). Rules and tools to predict the
splicing effects of exonic and intronic mutations. Wiley Interdiscip Rev RNA *9*.

Poser, I. *et al.* (2008). BAC TransgeneOmics: a high-throughput method for
exploration of protein function in mammals. Nat Meth *5*, 409–415.

Ran, F. A., Hsu, P. D., Wright, J., Agarwala, V., Scott, D. A., and Zhang, F. (2013).
Genome engineering using the CRISPR-Cas9 system. Nat Protoc *8*, 2281–2308.

Sarov, M. *et al.* (2012). A genome-scale resource for in vivo tag-based protein
function exploration in C. elegans. Cell *150*, 855–866.

Sarov, M. *et al.* (2016). A genome-wide resource for the analysis of protein
localisation in Drosophila. eLife *5*, e12068.

Sarov, M., Schneider, S., Pozniakovski, A., Roguev, A., Ernst, S., Zhang, Y., Hyman,
A. A., and Stewart, A. F. (2006). A recombineering pipeline for functional
genomics applied to Caenorhabditis elegans. Nat Meth *3*, 839–844.

Scolz, M. *et al.* (2012). GTSE1 is a microtubule plus-end tracking protein that
regulates EB1-dependent cell migration. PLoS ONE *7*, e51259.

Seyhan, A. A. (2016). A multiplexed miRNA and transgene expression platform
for simultaneous repression and expression of protein coding sequences. Mol
Biosyst *12*, 295–312.

Shaul, O. (2017). How introns enhance gene expression. Int. J. Biochem. Cell Biol.
91, 145–155.

Wang, H., Bian, X., Xia, L., Ding, X., Muller, R., Zhang, Y., Fu, J., and Stewart, A. F.
(2013). Improved seamless mutagenesis by recombineering using ccdB for
counterselection. Nucleic Acids Research.

Wang, Z., and Burge, C. B. (2008). Splicing regulation: from a parts list of
regulatory elements to an integrated splicing code. Rna *14*, 802–813.

Wassef, M., Luscan, A., Battistella, A., Le Corre, S., Li, H., Wallace, M. R., Vidaud, M.,
and Margueron, R. (2017). Versatile and precise gene-targeting strategies for
functional studies in mammalian cell lines. Methods, 1–10.

Yu, D., Ellis, H. M., Lee, E. C., Jenkins, N. A., Copeland, N. G., and Court, D. L. (2000).
An efficient recombination system for chromosome engineering in Escherichia
coli. Proceedings of the National Academy of Sciences *97*, 5978–5983.

Yu, P., Ma, D., and Xu, M. (2005). Nested genes in the human genome. Genomics
86, 414–422.

Zhang, M. Q. (1998). Statistical features of human exons and their flanking
regions. Human Molecular Genetics *7*, 919–932.

Zhang, Y., Buchholz, F., Muyrers, J. P., and Stewart, A. F. (1998). A new logic for
DNA engineering using recombination in Escherichia coli. Nat Genet *20*, 123–
128.

Zheng, X. *et al.* (2014). Conserved TCP domain of Sas-4/CPAP is essential for
pericentriolar material tethering during centrosome biogenesis. Proc. Natl. Acad.
Sci. U.S.a. *111*, E354–E363.

FIGURE LEGENDS

Figure 1: ESI-mutagenesis: a one step recombineering procedure to introduce

point mutations in BAC

(A) Scheme showing the organisation of an eukaryotic gene with exons (in blue)

and their flanking introns. Introns are spliced out from pre-messenger RNAs to

produce mature messenger RNAs (mRNAs) containing only exons. The core

splicing signals constitute the minimal information required for the splicing of an

intron, and include a 5' and 3' splicing signal sequences (5'SS and 3'SS,

respectively), a branch point (B.P.), and a poly-pyrimidine tract (P.P.T). The

position of these signals relative to the exon / intron border is indicated with the

corresponding nucleotides shown in pink (in introns) or light orange (in exons).

ESI mutagenesis is a one step recombineering procedure that relies on the

introduction into a BAC of a synthetic intron coding a selectable marker along with

the mutation of interest, thereby allowing for the easy selection of correct

recombinants. After transcription of the transgene in eukaryotic cells, the

synthetic intron is spliced out to produce a mutated mRNA. **(B)** The synthetic

intron constitutes of the intronic core splicing signal (in pink), an antibiotic

resistance cassette (Atb.^{R}) under the control of a dual eukaryotic/prokaryotic

promoter (in green), and two loxP sites flanking the antibiotic resistance cassette.

In order to serve as a template for BAC recombineering, the synthetic intron is

amplified by PCR with primers containing ~50 bp long homology arms to the

targeted sequence (in blue), the mutation(s) to introduce (in orange) and the

exonic part of the 5'SS and 3'SS (in light orange). **(C)** Scheme showing the

introduction of a point mutation into a BAC by ESI-mutagenesis. Depending on the

position of the sequence to mutate, the point mutation can be inserted along with

the synthetic intron into an exon (Left panel), or, if the mutation is located in the

proximity of a pre-existing intron, the part of the synthetic intron encoding the

antibiotic resistance can be targeted into this intron (Right panel). The synthetic

intron is spliced out during RNA maturation and only the desired mutation is

present in the mRNA.

Figure 2: ESI-mutated BAC transgenes yield proteins of the right size and

localisation

(A) GFP tagged BAC transgenes (Parental BAC, P) were mutated by ESI

mutagenesis to be RNAi resistant (ESI mutated BAC, ESI) and transfected into

HeLa cells. **(B)** Depending on the position of the sequence to mutate, the synthetic

intron was either targeted into an exon (left panel, light blue label in B, C and D)

or into the neighbouring intron by making use of pre-existing 5'SS, and / or 3'SS

(right panel, light brown label in B, C and D). Names of the mutated transgene are

indicated in the panel corresponding to their mutation strategy. The ESI mutated

AURKA transgene is analysed in Figure 4. The GTSE1 transgene was ESI mutated

at its SxIP motifs and is analysed in Figure 3. **(C)** GFP-tagged BAC transgenes ESI

mutated to be RNAi resistant yield proteins of the expected size. Immunoblotting

on cell-lysate of pools of HeLa cells transfected with either parental BACs (P) or

ESI-mutated BACs (ESI). GFP antibody was used as a probe. Transgenes names are

indicated at the top. **(D)** GFP-tagged transgenes ESI mutated to be RNAi resistant

show the same cellular localization as their parental GFP-tagged transgenes. Still

images of live cell imaging on HeLa stably transfected with the parental or the ESI-

mutated BAC. CHMP4B and AurkB -GFP transgenes were detected by

immunofluorescence with anti-GFP antibody. Because of differences in expression

level within cell pool, each picture was acquired and scaled independently of the

others. Scale Bar 10 μm.

Figure 3: ESI-mutated BAC transgenes show the expected phenotype

(A) BAC transgenes ESI mutated to be RNAi resistant show RNAi resistance at the

mRNA level. HeLa wild type, parental BAC lines, and ESI-mutated BAC lines were

treated with control-siRNA or transgene-specific siRNAs. Levels of mRNA

corresponding to the transgenes were determined by qPCR and normalised to

GAPDH mRNA levels. For each cell lines, the % of mRNA remaining in the

transgene-siRNA treated cells as compared to the control-siRNA treated cells are

presented (N=2 exp.). A panel of 3 transgenes mutated by targeting the synthetic

intron into an exon (light blue label) and 3 transgenes mutated by targeting the

synthetic intron into a pre-existing intron (light brown label) were analysed. **(B)**

BAC transgenes ESI mutated to be RNAi resistant show RNAi resistance at the

protein level. HeLa and clonal HeLa lines expressing ESI-mutated CEP135 or

AurkB GFP-tagged BAC transgenes were treated with control, CEP135 or AurkB -

siRNAs. Levels of AurkB, CEP135 and Tubulin were monitored by immunoblot

using specific antibodies. **(C)** Mutation of GTES1 SxIP motifs into SxNN by ESI

mutagenesis disrupts the interaction of a GTSE1-GFP BAC transgene with EB1. Cell

lysates from U2OS, U2OS expressing an ESI-mutated GTSE1-GFP SxNN BAC

transgene, and U2OS expressing an endogenously GFP-tagged GTSE1 (GTSE1-GFP

Cas9), were used in pull-downs with GST or GST-EB1 as baits. Pull-downs inputs

and outputs were probed by immunoblot using GTSE1 antibody. GST and GST-EB1

fusion were visualized by Coomassie Blue.

Figure 4: An AURKA-GFP BAC transgene ESI-mutated to carry RNAi resistance

rescues endogenous AURKA depletion. **(A)** An AURKA-GFP transgene ESI-

mutated to be RNAi resistant is expressed at endogenous level and is RNAi

resistant. Immunoblotting on U2OS and a U2OS clone expressing the ESI-mutated

AURKA-GFP BAC transgene (U2OS AURKA-GFP siRES (ESI)), both treated with

control- or AURKA- siRNA and blocked in mitosis. Antibodies against AURKA and

Tubulin were used as probe. **(B,C)** The ESI-mutated AURKA-GFP transgene shows

the correct cellular localization. Immunofluorescence on U2OS and U2OS AURKA-

GFP siRES (ESI) treated with control- and/or AURKA- siRNA. AURKA antibody (B)

or AURKA and Tubulin antibodies (C) were used as a probe. The ESI mutated

AURKA-GFP transgene rescues the mitotic arrest **(D)** and spindle collapse **(E,F)**

observed after endogenous AURKA depletion. The percentage of cells in each

mitotic phase is presented (D) for U2OS and U2OS AURKA-GFP siRES (ESI) clone

treated with control- or AURKA- siRNA (N= 3 exp.). U2OS and U2OS AURKA-GFP

siRES (ESI) clone were treated with control- or AURKA- siRNA and stained by

immunofluorescence using DAPI and an antibody against CEP135 (E). The pole-

to-pole distance (µm) in metaphase cells is presented (F). In AURKA-depleted

U2OS, spindle length was measured in prometaphase cells with nearly aligned

chromosomes. The number of mitotic cells used in each condition (n) and

statistically significant differences to U2OS treated with control-siRNA are

indicated (Kruskal Wallis, followed by paired Wilcoxon test). All scale bar 5 µm.

Figure S1: Available synthetic intron cassettes.

The synthetic intron cassettes generated in this study are depicted (A to D). The

resistance genes and the promoters (bacterial and eukaryotic) controlling their

expression are shown. When present, the loxP sites allow the removal of the

antibiotic resistances by the CRE recombinase. The position of the core intron-

based splicing signal sequences (intronic splicing donor dite, branch point (B.P.),

polypirimidine tract (PPT), intronic splicing acceptor site) is shown. The plasmid

from which each cassette can be amplified is indicated. Numbers indicate the

nucleotide positions within the cassette.

Figure S2: ESI mutagenesis of the GTSE1 SxIP motifs

(A) Scheme representing the sequence of exon 9 of GTSE1 containing the two SxIP

motifs (in orange) and a potential insertion site for the synthetic intron (in pink).

Distance (in bp) of the potential insertion site for the synthetic intron to the

border with introns 8-9 and 9-10 are presented. **(B)** Primers used to amplify the

synthetic intron are presented. They are composed of (i) homology arms to GTSE1,

(ii) the SxIP mutations (in orange) and (iii) a synonymous mutation (in orange)

allowing the potential insertion site (in pink) to fit the CAGG consensus. **(C)** The

amplified synthetic intron is recombined onto the GTSE1 BAC. The synthetic

intron is spliced out from the pre-mRNA to yield a mRNA coding GTSE1 mutated

at both SxIP motifs.

Figure S3: The ESI mutagenesis procedure at a glance.

(A) Scheme presenting the BAC recombineering protocol. Important parameters

to pay attention to are indicated. (B) Scheme showing the procedure to obtain the

ESI mutagenesis BAC recombineering template. Three PCRs (see Figure 1B and S1

for primer design) are performed to amplify the synthetic intron and pooled for

subsequent PCR purification. The resulting product is used as recombineering

template. (C) The ESI mutated BAC is transfected into eukaryotic cells. Stably

transfected cells are selected using the antibiotic resistance coded by the synthetic

intron. If the BAC carries a fluorescent protein tagged transgene, the antibiotic

resistance associated to the tag should be use for selection.

Figure 1

A Core intron splicing signals

DNA

Exon 1   Exon 2   Exon 3   Exon 4

5'   5' Splicing signal (5' SS)   B.P.   P.P.T.   3' Splicing signal (3' SS)   3'
C(A)AG GTA(G)AGT   A   YYY(n)YY   CAG G

mRNA C(A)AG G
Ex. 2   Ex. 3

B Synthetic Intron Cassette

5'
Homology arm to targeted sequence
Mutation
Exonic 5'SS
3'

Dual prokaryotic / eukaryotic promoter

5'SS loxP   Atb.R   loxP   B.P. / P.P.T. / 3'SS
3'
Exonic 3'SS
Homology arm to targeted sequence
5'

PCR

C

Host : *Escherichi coli*

Targeting of an exon   Targeting of an existing exon/intron border

Exon   Intron   Bacterial Artificial Chromosome (BAC)   Exon   Intron

synthetic intron   Synthetic intron with   5'SS loxP   loxP
selectable marker   homology arms   selectable marker
                               and mutation(s)

                               *BAC recombineering
                               (homologous recombination)*

                               Modified BAC

BAC stably integrated in eukaryotioc cell

                               BAC DNA

                               Transcription and splicing

                               Mutated mRNA

Figure S1

A

Synthetic intron *(plasmid TH0496)* **with :**
Blasticidin resistance under eukaryotic promoter
ClonNat resistance under bacterial promoter

B

Synthetic intron *(plasmid TH0823)* **with :**
Blasticidin resistance under eukaryotic promoter
ClonNat resistance under bacterial promoter
loxP sites for removal of antibiotic resistances by the CRE recombianse

C

Synthetic intron *(plasmid TH0480)* **with :**
Neocyn / Kanamycin resistance under dual eukaryotic / bacterial promoter
loxP sites for removal of antibiotic resistances by the CRE recombianse

D

Synthetic intron *(plasmid TH0821)* **with :**
Zeocin resistance under dual eukaryotic / bacterial promoter
loxP sites for removal of antibiotic resistances by the CRE recombianse

Figure S2

Figure S3

A

E.coli strains available from BAC collections

Eukaryotic gene of interest

Atb.R #1   BAC backbone

E.coli chromosome

*NB: the procedure can be applied on BAC carrying a tagged transgene.
In this case Antibiotic #1 corresponds to the antibiotic resistance associated to the tag.*

Electroporation of plasmid coding Rec α,β,γ

Day 0: Pre-culture of BAC clone (LB, Atb. #1)
Day 1: 1) Liquid culture from pre-culture (LB, Atb. #1)
      2) Electroporation of Rec α,β,γ plasmid

Rec   Tet.R   ⚠ Replicates only at 30°C

Selection at **30°C** on LB plate:
- Antibiotic #1 (BAC)
- Tetracycline

Day 0 to day 2

Recombineering of the synthetic intron

Day 2: Pre-culture transformed clone (LB, Atb. #1, Tet)
Day 3: 1) Liquid culture from pre-culture (LB, Atb. #1, Tet.)
      2) Induction of Rec α,β,γ for 70 min. (at OD 0.3-0.4)
      3) Electroporation of synthetic intron **(see B)**
      4) Incubation 50 min at 37°C

⚠ Pre-culture and culture at 30°C. Keep bacteria on ice while preparing them for induction and electroporation.

Selection at **37°C** on LB plate:
- Antibiotic #1 (BAC)
- Antibiotic #2 (Synthetic intron)

Day 2 to day 4

Colony PCR on Antibiotic #2 resistant clones

1) Use colonies for PCR and restreak on (LB, Atb. #2) plates
2) Sequencing of positive PCR product

Synthetic intron integration results in a shift in the PCR product size shift

Day 4

BAC purification

Day 5: Pre-culture of PCR positive clones (LB, Atb. #2)
Day 6: 1) Large scale culture from pre-culture (LB, Atb. #2)
      2) BAC purification using a BAC prep. Kit

⚠ Do not freeze BAC DNA (keep at 4°C)

Day 5 and 6

NB : The integrity of BACs that have integrated the synthetic intron can be further checked by comparing their restriction profile with the one of the template BAC

B

Synthetic Intron Cassette

Homology arm to targeted sequence
Exonic 5'SS
Dual prokaryotic / eukaryotic promoter
5'SS loxP Atb.R loxP B.P. / P.P.T. / 3'SS
Homology arm to targeted sequence
Mutation
Exonic 3'SS

3 x PCR
⚠ Order oligos HPLC purified

Pool for PCR purification

Synthetic intron with mutation and homology arms

*For primer design, see **Figure S2***

C

BAC transfection into host cell line

Day 5: Seed cells for transfection (60-70% confluent on day 6)
Day 6: Transfect BAC DNA into cells
Day 7: Change media to fresh media

Day 5 to day 7

Selection of transfected clones expressing the ESI mutated transgene

Day 8: Change media to selection media (Antibiotic #2)
Day 21-28: Regularly change media to fresh selection media
Pick and expand resistant clones

Antibiotic # 2 selection

Colony picking and expansion

Clone characterization

⚠ If the BAC harbours a tagged transgene it is advised to select clones using the antibiotic corresponding to the resistance attached to the Tag.

from Day 8