1 Correlated structure of neuronal firing in macaque visual cortex limits information for
2 binocular depth discrimination.
3

4 Jackson E. T. Smith* and Andrew J. Parker
5
Department of Physiology, Anatomy and Genetics
University of Oxford
Oxford
Oxfordshire
OX1 3PT
United Kingdom
12
13 *Correspondence: jackson.smith@dpag.ox.ac.uk

14
Word count:
Title: 119/120 characters (with space)
Abstract: 150/150
Introduction: 1,481
Results: 8,125
Discussion: 2,196
Main figure legends: 1,600
Total (Intro, Results, Discussion, Fig legends): 13,402
23

24 ABSTRACT
25
Variability in cortical neural activity potentially limits sensory discriminations. Theoretical
work shows that information required to discriminate two similar stimuli is limited by the
correlation structure of cortical variability. We investigated these information-limiting
correlations by recording simultaneously from visual cortical areas V1 and V4 in macaque
monkeys, performing a binocular, stereo-depth discrimination task. Within both areas, noise
correlations on a rapid temporal scale (20-30ms) were stronger for neuron-pairs with similar
selectivity for binocular depth, meaning that these correlations potentially limit information
for making the discrimination. Between-area correlations (V1 to V4) were different, being
weaker for neuron pairs with similar tuning, and having a slower temporal scale (100+ms).
Fluctuations in these information-limiting correlations just prior to the detection event were
associated with changes in behavioural accuracy. Although these correlations limit the
recovery of information about sensory targets, their impact may be curtailed by integrative
processing of signals across multiple brain areas.
39
40 Key words: V1, V4, Cortex, Vision, Rhesus, Noise correlation, Stereo depth, Encoding,
41 Behaviour, Utah array

42
43
44 INTRODUCTION
45
Processing in the sensory areas of cerebral cortex is thought to limit the detection and
discrimination of weak sensory signals (A. J. Parker & Newsome, 1998). Initial work
examined how the changes in the firing of action potentials induced by near-threshold
changes in the sensory stimulus could be detected against the variability of the base-level of
cortical activity (Barlow, Kaushal, Hawken, & Parker, 1987; Newsome, Britten, & Movshon,
1989; A. J. Parker & Hawken, 1985; Tolhurst, Movshon, & Dean, 1983). This work elucidated
how detection and discrimination could be related to the statistics of the firing of single
neurons.
54
These initial studies indicated that it was not unusual for the detection performance of
single neurons to match the behavioural performance of the entire visual system. However,
since the firing of all neurons shows some variability, combination of the firing from a pool
of neurons to form a multi-neuron sensory signal ought to improve the detection rate for
weak stimuli. The puzzle is why the nervous system apparently cannot provide a better
sensitivity for the organism by making a statistical combination of the signals from multiple
neurons.
62
One answer from early work on the somatosensory periphery is that the correlation of
residual noise within a pool of sensory neurons may also be an important factor (Johnson,
Darian-Smith, & LaMotte, 1973). In the absence of correlations, additive pooling of neuronal
signals results in a steady improvement of sensory thresholds with increasing pool size.
However, the presence of correlations limits the potential gain in performance from
increases in the numbers of neurons. As more and more neurons are added, the portion of
noise that is uncorrelated does indeed get averaged away, but the portion of noise that is
correlated, and therefore shared by every neuron, cannot be reduced this way and comes to

dominate the limit on performance (A. J. Parker & Newsome, 1998; Shadlen, Britten,
Newsome, & Movshon, 1996).
73
74
75 *Correlation structure in the pool of sensory neurons*
76
Recent work expands that insight, showing how the nature of the pooling computation and
the structure of correlations are both critical in setting limits on the encoding and decoding
of information within sensory cortex (Abbott & Dayan, 1999; Averbeck, Latham, & Pouget,
2006; Kanitscheider, Coen-Cagli, Kohn, & Pouget, 2015b; Moreno-Bote et al., 2014; Pouget,
Dayan, & Zemel, 2003). The correlation structure of a pool of neurons in relation to sensory
discriminations is captured by making all pair-wise comparisons between members of the
pool, across a number of repeated presentations of the sensory stimuli. Critically for our
study, this theoretical work demonstrates that the signals from a neuron-pair limit the
discrimination between two stimuli in two ways (Nogueira et al., 2020). The first is fairly
straightforward, which is that the change in the stimulus should alter the mean level of
firing from both neurons; bigger differences in the mean response should improve the
discrimination ability. The second concerns the variability of neuronal firing and its
correlation structure across the population. The theoretical work demonstrates that noise
or variability is the most detrimental when its correlated structure causes the conjoint firing
of the neuron-pair to mimic a response to some change in the stimulus that did not, in fact,
occur. Such correlations have been termed "information-limiting correlations" (Kohn, Coen-
Cagli, Kanitscheider, & Pouget, 2016; Moreno-Bote et al., 2014).
94
Assessment of the correlation structure ideally requires the simultaneous recording of
multiple pairs of neurons from within the sensory cortex during the performance of a
behavioural task with repeated stimulus presentations (Kanitscheider et al., 2015b; Moreno-
Bote et al., 2014). In our work, we addressed this by training monkeys to discriminate
binocular stereoscopic depth, while we tested a range of neuronal encoding by using
chronically implanted Utah arrays to record multiple neurons simultaneously in two cortical
sites: the primary visual cortex (V1) and the extrastriate area V4. Although the fact that we
studied binocular vision is not especially critical for the questions that we investigate here,
the processing of stereoscopic information proves to be a good model system for isolating
the contribution of cortical processing, as stereo involves the direct combination of
information from the two eyes at a cortical level.
106
107 *Neuronal recording during sensory task performance*
108
Our task was designed so that we could thoroughly test all simultaneously recorded neurons
over a range of stereoscopic depths, to build up a complete tuning curve for each neuron.
Many studies have employed tasks that involve a binary decision, potentially limiting which
cortical neurons are relevant to the task, with the result that many of the recorded neurons
may never be tested under the optimal (or near-optimal) conditions for assessing their
encoding of information. Measurement over the whole range of sensory encoding also
increases the yield of relevant neuronal responses in the case of using fixed recording
electrodes, as with the Utah array.
117

The paradigm used here allowed us to estimate the extent to which the two neurons
respond in the same way to a range of stimulus values, which is termed the signal
correlation (Bair, Zohary, & Newsome, 2001) for the neuron pair. As our task involved the
discrimination between two closely similar binocular depths, we also assessed the
differential response of the sensory neurons by taking the slope of the tuning curve at each
point. By making repeated measures of the response to each binocular depth, we could also
estimate the noise correlation of all recorded neuron-pairs at each tested value of binocular
depth, and apply recently-developed methods to search for information-limiting
correlations (Kanitscheider, Coen-Cagli, Kohn, & Pouget, 2015a). We couple this approach
with a time-dependent measure of correlations (Bair et al., 2001) to estimate the temporal
scale of information encoding.
129
In earlier comparisons of the correlation structure and sensory tuning of cortical neurons,
the data were either from anaesthetized animals (M. A. Smith & Kohn, 2008) or from awake
animals that fixated without performing any other task (M. A. Smith & Sommer, 2013). In
other cases, the behavioural measures were taken on one task (judgment of heading
direction) but the measures of correlation structure were taken during the performance of a
different task (visual fixation) (Pitkow, Liu, Angelaki, DeAngelis, & Pouget, 2015). In an
alternative scenario, the measurements of signal correlation were obtained whilst the
animal was passively fixating rather than actively discriminating the stimuli (Bondy, Haefner,
& Cumming, 2018).
139
An important advance made with the recordings undertaken here is that the macaque
monkeys actively performed a behavioural task, making a stereoscopic depth judgment on
the presented stimuli. We obtained simultaneous neuronal and behavioural data across the
complete tuning curve of the neurons. A number of recent studies have demonstrated that
the correlation structure of neuronal firing in sensory areas is influenced by the
manipulation of cognitive factors such as attention or perceptual learning (Bondy et al.,
2018; Ni, Ruff, Alberts, Symmonds, & Cohen, 2018; D. A. Ruff & Cohen, 2014; Douglas A.
Ruff & Cohen, 2016; Verhoef & Maunsell, 2017). We aimed to maintain these factors as
constant as possible to focus on the discrimination performance of the neuronal population,
by ensuring that the design of the experimental task arranged for spatial and featural
attention to be consistently directed towards the depth signalled by the visual stimuli.
151
Finally, the relationship between perceptual decisions and the correlation structure of
neuronal firing has been a topic of intensive research (Krause & Ghose, 2018; Krug, Curnow,
& Parker, 2016; Pitkow et al., 2015) ever since it was suggested that the two may be linked
(Shadlen et al., 1996). To assess this link, it is critical to measure the correlation structure in
the most relevant neuronal population as the animal is performing a sensory discrimination
task. In this paper, we are primarily concerned with the state of cortical firing as the animal
waits attentively for the arrival of a change in binocular depth that must be detected and
reported. Our results show for the first time that the state of putative differential
correlations within the neuronal population has a direct association with the accuracy of the
upcoming behavioural decision.
162
163   *Correlations within and between cortical areas.*
164

A further open question is whether the noise correlation structure limits the passage of signals through each of the stages of cortical processing (Zylberberg, Pouget, Latham, & Shea-Brown, 2017). We tackle this issue by examining the correlation structure and information encoding of neuronal activity within and between cortical areas V1 and V4. Specifically, we test how information may pass between cortical areas, with respect to neuronal correlations during task performance. We examine the time-course of information-bearing signals and assess them in relation to information-limiting correlations and behavioural choice. Our results suggest a simple model, which demonstrates that the potentially deleterious effects of information-limiting correlations do not build up as sensory signals pass from one cortical area to another. We suggest that this is an important and previously unrecognized principle of neural encoding across multiple areas of the cerebral neocortex.

RESULTS

We trained two Rhesus macaques (*Macaca mulatta*) to observe 4 patches of dynamic, random dot stereograms (RDS) and perform an odd-one-out task (Figure 1), in which they detected which one of the four patches displayed a change in binocular depth. Each RDS was configured as a central circular patch, whose binocular depth varied from trial to trial, and a surround region of dots located in the fixation plane of the monkeys' eyes. In the 'present RDS' phase (Fig.1Aiii), all 4 stimuli had the same depth for the central patch, which was chosen from one out of a set of 13 different values. The 'popout' phase (iv) presented the change in depth, which was applied to the centre of one of the 4 RDS stimuli. That patch became the target for a saccade to indicate a correct choice and earn a reward; a saccade to any of the other three RDS was incorrect and earned no reward.

The monkeys' behaviour indicated attention to the RDS stimuli in the 'Present RDS' phase, as they were better at reporting the location of the disparity step change as it grew in magnitude (Fig.1B, % correct), while their reaction times (RT) decreased. On catch trials with no change in the RDS, the animals had to guess, so their accuracy was at chance levels (Fig.1B, dotted line, left panels) and their RTs were longest (Fig.1B, right panels). Fig.1C shows the relationship between percent correct as a function of RT using a 50ms moving window, grouped by magnitude of the popout disparity change. Correct decisions are broadly within the window of 100-400ms after pop-out.

Neural activity in the dorsal, gyral portions of V1 and V4 was recorded, using two 64 channel (8x8; 0.4mm spacing) Utah array electrodes (one array per area) implanted chronically in the left hemisphere. The electrode placement meant that the recorded neurons had receptive fields (RF) that were stimulated by the lower-right RDS (Figure 2A&S1). An automatic spike sorting method was used to identify action potentials followed by manual checking (see Methods for details.) As with other work with Utah arrays, analyses were based on both single units and multiunit clusters and the term 'unit' refers interchangeably to either.

Variations between neuronal RF with respect to spatial location and preference for binocular depth, dot size, dot contrast and other dimensions of the RDS stimulus mean it is impossible to simultaneously achieve optimal stimulation of every unit recorded by the arrays. We therefore focused on varying these parameters to ensure that there was significant co-activation of the V1 and V4 units (Fig.2D&E, S2D&E). To validate the co-activations of neuron pairs, we applied entry criteria, such that each unit in the pair had to show statistically significant tuning to disparity (one-way ANOVA at the 1% level) and fire at least 5 spikes per second to its preferred disparity. For all the recordings reported here, the animals were performing the stereo depth task, which requires allocation of attention to the four RDS patches.

We made separate measurements of RF location and stimulus preferences in preliminary experiments in each monkey. These measurements demonstrated that there was scatter in the positions of the RF centres of the units, both within and between cortical areas (Fig.2A&S1). We later address how this separation of RF centres may have influenced our results. For now, we confine ourselves to noting that the combination of electrode

224   placements and adjustment of the configuration of the RDS stimulus resulted in common
225   activation of pairs of V1 and V4 units by the centre portion of the same RDS. We focus on
226   the activations of units from the 'Hold fixation' and 'Present RDS' phases of the trial
227   (Fig.1Aii&iii, 2D&E), before the arrival of the discrimination stimulus itself.
228
229   *V1 and V4 units were selective for binocular disparity*
230
The V1 and V4 neural signals that we recorded are able to support task performance.
Disparity tuning curves (Fig.2B) are shown for an example V1 (circles, gold line) and V4 (plus
sign, blue line) unit. The solid lines are the best-fitting Gabor functions (equation 21)(Prince,
Pointon, Cumming, & Parker, 2002), which describe the response profiles as a function of
disparity. To quantify a unit's contribution to the estimation of stimulus disparity, we
computed how much information the firing of the neuron could contribute to the
identification of the currently displayed disparity of the RDS, called the disparity mutual
information (DMI, equation 8; Fig.2C&E, S2A-C&E). This measure correlated well with a
previously-established measure of selectivity (Fig.S2A&B) – the disparity discrimination
index (DDI; Prince et al., 2002) – for the V1 (Spearman correlation, Initial, $r = 0.976$, $p =$
9.559e−130) and V4 ($r = 0.953$, $p = 4.705e−111$) neurons.
242
243   The neurons in V1 and V4 that responded most to the onset of the stimulus were also the
244   most selective for disparity, as the peak firing rate and DMI (in the 50 to 200ms window)
245   were well correlated (Fig.2C&S2C: V1 in gold, Spearman $r = 0.609$, $p \rightarrow 0$; V4 blue, $r =$
246   0.248, $p = 2.663e−4$). Low-firing neurons were relatively poor at encoding disparity,
247   compared to highly active neurons.
248
We searched for the time point at which the units became selective for disparity by
convolving the spike trains with a causal exponential kernel (20ms time constant) to obtain
the firing rate and DMI time series for each unit. The average firing rates (Fig.2D&S2D; V1,
gold; V4, blue) had a transient burst of activity in response to the onset (black dotted) of the
RDS, which then settled down to a steady rate. As expected, V1 began responding to the
RDS before V4 and V1 neurons showed selectivity for the binocular disparity slightly earlier
than V4 neurons.
256
Firing rates in both areas were selective for binocular disparity almost as soon as the units
began responding to the RDS, with the average DMI showing a similar time course to the
changes in the firing rate (Fig.2E&S2E). We verified the link in timing between the DMI and
the peak firing for each unit by finding the Spearman correlation of firing rate and DMI,
matched by time bin, in an 800ms window starting 50ms after the RDS onset *i.e.* we took
the zero-lag cross-correlation of the firing rate and DMI time series. The result was
significantly positive for V1 (mean $r = 0.192$, right-tailed t-test $p = 3.339e−22$) and V4 ($r =$
0.245, $p = 7.819e−31$), showing that individual units were more selective for disparity at
moments in the trial when their firing rate was higher.
266
We partitioned each trial into three consecutive windows for further analysis (Fig.2D&E).
The 'Spontaneous' window (leftward arrow) captured neural responses during the 'Hold
fixation' phase (Fig.1Aii) of the trial, in which the monkeys held their gaze on the fixation
target prior to the arrival of RDS signals in V1. The response to the onset of the RDS is

captured by the 'Initial' window (thick bar and black solid lines, marking 50-200ms after
stimulus onset), which shows a transient burst of firing and rise in the DMI. The responses
to the RDS for the remainder of the 'Present RDS' phase of the trial are captured by the
'Steady state' window (rightward arrow, Fig.1Aiii). Within-unit comparisons showed that
disparity selectivity (DMI) was stronger in the 'Initial' than in the 'Steady state' window, for
V1 (mean paired difference = −0.577 bits/s, left-tailed t-test p = 1.294e−12) and V4 (−0.220
bits/s, p = 6.827e−11); in part, these differences reflect the greater firing rates in the Initial
window.
279
280 *Information-limiting correlations*
281
If the sensory discrimination task performed by the monkeys is supported by a linear
computation across pools of active neurons, then a special class of noise correlation –
differential correlation (Moreno-Bote et al., 2014) – is potentially a fundamental limit on the
information that is encoded by the sensory pool. Figure 3A–C show how differential
correlations could limit information that is retrievable from a pair of neurons, whose
activities were simultaneously recorded in V4 during task performance. The pair had similar
tuning for binocular disparity (*i.e.* $0 < r_{signal}$, Fig.3A). Both units had a common negative slope
in their tuning curves around 0.1° (black dashed), where they were sensitive to the same
small changes in disparity. Adding the signals from these two neurons potentially improves
sensitivity but this depends on the noise distributions.
292
Joint responses to multiple presentations of the 0.1° disparity stimulus were obtained by
counting spikes in the Initial window (Fig.3B, grey). At 0.1°, the pair had a positive noise
correlation (Fig.3C). Critically, the principal component (arrow) of the covariance was nearly
parallel to the tangent line (grey) of the tuning curves at 0.1°. To see the consequence of
this, consider a linear discriminator that classifies the joint responses as one of two
disparities on either side of 0.1°. If the decision boundary is perpendicular to the tangent
line (grey) then noise correlation (arrow) will make the joint responses jump across the
decision boundary, causing classification errors (Averbeck et al., 2006). Put simply,
correlated noise would mimic true changes in the stimulus, limiting the sensitivity of this
pair.
303
This outcome contrasts with the responses of a second pair (Fig.3D–F), recorded
simultaneously from V1 and V4. As before, both units had steep, overlapping slopes in their
tuning curves (Fig.3D). Unlike the V4/V4 pair, the V1 and V4 units were selective for opposite
disparities ($r_{signal} < 0$). Spike counts from both units in response to a disparity of 0° (Fig.3D,
black dashed) were obtained as before (Fig.3E), and the joint responses were examined
(Fig.3F). Although this pair has a negative signal correlation, they have a positive noise
correlation (r_{sc} = 0.203; p = 0.029): the noise correlation does not align to the tangent (noise
correlation first principal component PCA1 at 72.8° [95%CI: 58.5°, 85.3°]). In this case,
correlated noise would be less likely to leak across the decision boundary. Further examples
of neuron-pairs with opposite signs of relationship between signal and noise correlation are
shown in Supplementary Figure S3. For those examples, one pair also had opposite selectivity
and positive correlation, but two others had the inverse pattern, with similar disparity
selectivity and negative noise correlations.
317

r_{CCG} measures noise correlation at different time scales

Noise correlations (r_{SC}, equation 10&11) are captured experimentally by setting up a
window of time, in which to count the number of spikes in response to repeated
presentations of the same stimulus. In terms of timing, this encompasses the synchronous
firing of spikes with millisecond precision up to long time-scale, co-fluctuations in spike
numbers (Marlene R. Cohen & Kohn, 2011; M. A. Smith & Kohn, 2008; M. A. Smith &
Sommer, 2013). Measurement of spike count correlation confounds these slow and rapid
correlations, depending on the size of the counting window that is used. Another, time-
based measure – r_{CCG} (equation 17) – is used to reveal noise correlations at different
temporal scales (Bair et al., 2001). This measure, r_{CCG}, is the time integral of the spike-train
cross-correlation and reveals the timing relationships between correlated spikes, as well as
the strength of the correlation. The r_{CCG} measure converges on r_{SC} as the width (τ) of the
integration window reaches that of the counting window used to calculate r_{SC}. We asked
whether information-limiting correlations occupy a limited set of temporal scales by using
r_{CCG} to examine noise correlations within cortical areas V1 and V4 and between those two
cortical areas (Figures 4 & S4).

Within V1 (Fig.4A), the time scale of noise correlations changed as the trial progressed.
Before the RDS appeared, correlations occupied a slow time-scale, being strongest on the
order of hundreds of ms (dashed). The time window for correlations became tighter after
RDS onset, peaking at τ = 20ms; r_{CCG}(τ=20ms) was greater in the Initial window (teal) than
the Spontaneous window (mean paired difference = 2.141e−2, paired t-test p =
1.592e−124). Even in steady-state responses to the RDS (magenta), the slower correlations
were weaker, with a drop in r_{CCG}(τ=100ms) from the Spontaneous window to the Steady
state (−1.092e−2, p = 1.005e−21).

These changes in r_{CCG} over time scale are reflected in the shapes of the V1-V1 cross-
correlograms (CCG, equation 19; Fig.4B). In the period of Spontaneous activity, the CCG was
positive and had a broad peak, coinciding with the steady increase of Spontaneous r_{CCG} as τ
increased. In contrast, the Initial and Steady state CCGs had narrow positive peaks flanked
by negative side lobes, showing that the majority of correlated spikes occurred within tens
of ms of each other. This matched the peaks in Initial and Steady state r_{CCG} at τ = 20ms.

Unlike V1, correlations within V4 (Figs.4D & S4A) began saturating at τ = 20ms in all three
analysis windows. V4 correlations were also stronger in the Spontaneous window, with
r_{CCG}(τ=20ms) dropping in the Initial window (−2.188e−2, p = 9.053e−105) and then changing
little from the Initial to Steady state windows (9.464e−4, p = 0.041). These differences
appeared in the V4 CCGs (Figs.4E & S4B) in which the Initial and Steady state CCGs were
mainly attenuated versions of the Spontaneous CCG.

In marked contrast, the correlations between V1 and V4 (Fig.4G) saturated at τ = 100ms.
Although these between-area correlations are weaker, it is important to understand that
they were significantly positive during the stimulus presentation (mean r_{CCG}(100ms); Initial =
8.075e−3, right-tailed t-test p = 1.512e−43; Steady state = 7.398e−3, p = 4.088e−61;
Spontaneous = −1.286e−3, p = 1). There was no measurable change in the V1/V4
correlations from the Initial to the Steady state window (mean difference = −6.767e−4,

paired t-test p = 0.159). The V1/V4 CCGs (Fig.4H) reveal a consistent delay, whereby V1
spikes tended to occur up to 50ms earlier than the associated V4 spikes.
367
Bair and colleagues (Bair et al., 2001) established an important property of r_{CCG} , which is
that it provides a lower-variance estimate of short-term correlations, compared to r_{SC} . The
parameter τ can be adjusted so that $r_{CCG}(\tau)$ integrates only the central peak of the CCG and
discards the longer time-course components. These longer time-course components are
captured by r_{SC} but often add variance to that estimate. This principle is illustrated for the
"Steady state" condition by a direct comparison of r_{CCG}(100ms) and r_{SC} in V1 (Fig.4C), V4
(Fig.4F&S4C), and between areas (Fig.4I). There was a strong, positive Spearman correlation
of r_{CCG} and r_{SC} in all cases (V1 r = 0.750, V4 r = 0.772, V1/V4 r = 0.807, all $p \to 0$), showing
that they measured a common source of correlation. However, the marginal distributions
plotted at the top and right-hand sides show that the variation (standard deviation) of r_{SC}
(σ_{SC}) was greater than that of r_{CCG}(100ms) (σ_{CCG}) for all three cases (V1, V4, V1/V4, see Table
1). Hence, the Pitman's r_{SD} test for comparing variances of correlated variables (Lee, 1992)
shows that r_{CCG} is a more accurate measure of short-term correlations than r_{sc}.
381
382 *Table 1: Comparison of scatter in estimates of r_{sc} and r_{ccg}(100ms)*

Cortical Area	σ_{SC}	σ_{CCG} (τ=100ms)	Pitman's r_{SD}
V1	0.135	0.044	0.830
V4	0.112	0.056	0.766
V1/V4	0.112	0.030	0.886

383 Each r_{SD} has a p-value approaching 0
384
385 *Noise correlations varied with disparity tuning, temporal scale, and time*
386
387 We asked initially whether the noise correlations of these neurons had any consistent
388 relationship to disparity selectivity (Figure 5 & S5). We plotted r_{CCG} as a function of the signal
389 correlation between pairs (r_{signal}, equation 9), grouped by temporal scale (τ,
390 Fig.5A,C,E&S5A), or as a function of τ, grouped by r_{signal} (Fig.5B,D,F&S5B).
391
For pairs in V4, there was a clear positive relationship of r_{CCG} and r_{signal} (Fig.5C&S5A) at both
short (τ = 20ms, dashed) and long (τ = 150ms, solid) time scales in the Initial (black) and
Steady state (magenta) windows (see Table 2). This coincided with an increase in r_{CCG} across
a range of values of τ for pairs with similar disparity preferences (Fig.5D&S5B, r_{signal} > 0,
dashed), compared to those with opposite preferences (r_{signal} < 0, solid). For pairs in V1, a
similar but weaker positive relationship was observed (Fig.5A&B) that was significant in the
Initial window and Steady state window for long time scales (τ = 150ms) but not short (τ =
20ms) (see Table 2). In summary, neurons in the same cortical area that have similar
preferences for the binocular depth of the stimulus tend also to show stronger, more
positive noise correlations, suggestive of a local network of neurons that process binocular
depth.
403
404 By comparison, the pattern of correlations between V1 and V4 (Fig.5E&F) were therefore
405 different from within-area correlations in two important ways. First, there was no
406 measurable relationship between r_{CCG}(20ms) and r_{signal}. Second, at longer time-scales, a

weak relationship emerged, but it was negative (see Table 2). That is, V1/V4 pairs tended to have opposite noise and signal correlations (like in Fig.3D–F), unlike the pairs from within the same area (*e.g.* Fig.3A–C). The pattern of correlations between V1 and V4 suggests that a different principle governs their relationships. These between-area correlations are also more evident at longer time-scales.

Table 2: relationship between r_{signal} and r_{ccg}, accompanies Figure 5

Cortical areas	τ (ms)	Initial		Steady state	
		Spearman r	P-value	Spearman r	P-value
V1/V1	20	0.042	0.038 *	0.030	0.144 ns
	150	0.050	0.013 *	0.054	0.009 **
V4/V4	20	0.172	4.133e−16 ***	0.170	5.674e−16 ***
	150	0.141	2.448e−11 ***	0.131	5.906e−10 ***
V1/V4	20	−0.013	0.392 ns	−0.004	0.779 ns
	150	−0.032	0.032 *	−0.025	0.091 ●

A concern might be that some of these relationships could be driven by the fact that a dynamic RDS stimulus with the same properties will vary subtly from trial to trial in the exact positioning and sequencing of the dots. Such variation could conceivably induce a local correlation in firing due to similar responses to the details of the dot sequence rather than the same binocular depth. One way to address this is to construct pseudo-random sequences of dot presentation that are in fact identical. This is achieved by creating so-called 'frozen noise', in which the same seed is given to the pseudo-random number generator on every trial, thereby creating identical sequences of dots on repeated trials. We performed some control experiments using frozen noise, comparing neuronal responses with and without resetting of the number seed. These experiments were performed with 100% binocular correlation, as this condition may be expected to give the strongest form of false association between binocular depth sensitivity and dot sequence. As with previous uses of this approach (Britten, Newsome, Shadlen, Celebrini, & Movshon, 1996), we found that the details of the dot sequence did not matter substantially. Specifically, the information about binocular depth was not artificially inflated when frozen noise, rather than new random sequences, was used to generate the RDS stimuli.

Empirical evidence of differential noise correlations

A property of information-limiting correlations is that they limit discrimination thresholds in the neighbourhood of a specific stimulus value (Moreno-Bote et al., 2014). For this reason, they are also called differential correlations, since the concern is with the component of noise that lies along the tuning curve contour corresponding to the stimulus change. Theory therefore predicts that measured noise correlations are the sum of two sources, one that limits the sensory threshold due to differential correlations and one that does not (Moreno-Bote et al., 2014).

The differential correlations that limit information about the sensory discrimination scale
with the product of the first derivatives of the two tuning curves from the pair of neurons
under consideration (Fig.3C). It is here that the specifics of our experimental design present
an advantage, because every pair of neurons was tested across a range of binocular
disparities, as the psychophysical task was being actively performed. Hence, the relationship
between the size of the noise correlation and the value of the derivatives of the tuning
curves could be measured at multiple points along the tuning curve for every neuron pair.
Moreover, all of these measurements could be directly related to the behavioural
performance of the task.
453
The other component of noise correlations does not limit information because, in principle,
this source of noise could be eliminated by averaging or pooling across many neurons.
Nonetheless, this source of noise does present a practical limitation because it may mask
the differential correlations that need to be measured and characterized. We reasoned that
the low variance measure of noise correlation, r_{CCG}, would be more likely to reveal weak and
rapid differential correlations, rather than r_{SC}, which is noisier. Thus, we examined the
relationship of r_{CCG} at different integration times (τ) and the product of derivatives over a
range of time scales.
462
For pairs of neurons from the same area (V1 or V4), noise correlations had a positive
relationship with the product of derivatives ($f'{\times}f'$, see equation 23), during both the Initial
and Steady state phases of neuronal activity (Fig.6&S6). This relationship is shown for some
specific values of time scale (τ), plotting r_{CCG} against $f'{\times}f'$ (V1 Fig6A&B; V4 Fig.6D&E,S6A&B;
regression lines in orange). A continuous plot of the slope (Spearman correlation) between
r_{CCG} and $f'{\times}f'$ across time scales (τ) is shown in Fig.6C for V1 and Fig.6F&S6C for V4; these
continuous plots were used to select time scales for the illustrative plots
(Fig.6A,B,D,E,S6A&B). The vertical arrows on the time-scale axes show the τ of the maximum
regression slopes for V1 (Fig.6C) and V4 (Fig.6F&S6C). Statistical summaries for the plots in
Fig.6 (A,B,D,E) are shown in Table 3. It is evident that the relationship between r_{CCG} and $f'{\times}f'$
is most prominent in a narrow range of time scales (10 $\lessgtr \tau \lessgtr$ 100ms).
474
475 *Table 3: summary of relationship between r_{CCG} and $f'{\times}f'$, accompanies Fig.6*

Cortical areas	Initial			Steady state		
	τ (ms)	Spearman r	P-value	τ (ms)	Spearman r	P-value
V1/V1	41	0.033	1.137e−6 ***	95	0.049	3.423e−13***
V4/V4	21	0.072	1.437e−24***	32	0.079	2.726e−29***
V1/V4	88	−0.026	1.023e−7***	109	−0.019	9.298e−5 ***

476
For pairs of neurons recorded simultaneously between V1 and V4, the result was
unexpected (Fig.6G–I). Instead of a positive relationship, we observed a consistent negative
relationship of r_{CCG} and $f'{\times}f'$ in the Initial and Steady state time periods, which peaked for a
narrow range of τ close to 100ms (Fig6G,H,I). This negative relationship of r_{CCG} and $f'{\times}f'$ does
not mean that the values of r_{CCG} are themselves negative. On average, the values of r_{CCG} and
indeed of r_{SC} are positive (see Fig.4). It is how these positive values of r_{CCG} vary with the size
and sign of $f'{\times}f'$ that is of concern here, similar to the between-area relationship of r_{signal} and
r_{CCG}, mentioned above (Fig 5 and Table 2).

Differential noise correlations are attenuated at long time scales

The amount of information that can be encoded by a neuronal pool depends fundamentally on the number of different firing states that the pool can adopt. Thus, correlation among members of the pool reduces the capacity for information coding. For a sensory discrimination task, the impact of these correlations depends on the discrimination task that is to be performed. Differential correlations that mimic the change in neuronal response to the stimulus change that must be identified will result in a saturation of the information in a neuronal pool of a certain size. This is, differential correlations will follow the same principle identified by earlier studies, which concluded that threshold performance as a function of pool size is limited by correlation in the neuronal spike counts (Johnson et al., 1973; Shadlen et al., 1996).

Therefore, we measured the information about binocular depth that was available within neuronal pools of increasing size (Figure 7A&S7A). Information was assessed as linear Fisher information (equation 24)(Kanitscheider et al., 2015b), which measures how accurately the responses of the neurons in a pool can determine which binocular depth was presented. Fisher information is inversely related to the width of the likelihood function for estimating binocular depth. It therefore increases when the population response can better distinguish between different RDS depth values, and an increase of Fisher information is brought about from the addition of informative neurons or from changes in the pool's correlations (Kohn et al., 2016). Empirically, information-limiting correlations can be removed from measured samples of neuronal population activity by shuffling the relationship between neuronal activity across multiple presentations. Shuffling decorrelates the population activity and is therefore is predicted to increase the Fisher information, if information-limiting correlations are present in the experimental measurements.

Using responses from the Initial phase of firing, we calculated the normalized and averaged bias-corrected linear Fisher information (equation 32) for pools of units from V1-only (gold, left), V4-only (blue, middle), and V1 mixed with V4 (green, left). Units were added to each pool from the most selective for disparity down to the least selective (Sanayei et al., 2018); thus, part of the apparent saturation is due to exhaustion of informative units rather than the effect of correlations. The range of pool sizes was limited to those observed in at least half of the experimental sessions. The Fisher information available from experimental recordings was compared against the information within an equivalent pool of decorrelated units (solid black curve), for the cases of rapid (τ = 20ms, dotted) and slow (τ = 150ms, solid coloured) noise correlations. For all pool types, we found that the empirical information at τ = 20ms began to diverge from the decorrelated information for pools of 10 units. In pools with only V1 or V4 units, there was a similar but weaker divergence when τ = 150ms. However, for the mixed V1 and V4 pool, the empirical information at τ = 150ms was closely aligned with the decorrelated information (Fig.7A, right panel).

To get more detail about the effect of time scale, we plotted the average, raw (not normalised) Fisher information of the Initial responses as a function of τ (Fig.7B&S7B), using the maximum available pool size for each pool type in each experimental session. The empirical information (solid coloured line) was compared against the decorrelated

532   information (dashed, black horizontal line). For all pool types, the empirical information
533   tended to increase with τ. But the empirical information converged on the decorrelated
534   information only in the mixed V1 and V4 pool, at around τ = 100ms.
535
We confirmed this result by plotting the Fisher information after decorrelation against the
empirical information for τ = 150ms (Fig.7C&S7C). The information after decorrelation was
greater than empirical for the single area pools (V1: mean paired difference 'PD' = 66.117,
paired-sample t-test p = 4.700e−6; V4: PD = 32.316, p = 1.745e−7), but there was no
significant difference for the mixed pool (V1/V4: PD= −20.558, p = 0.183). Similar results
were observed using Steady state responses (Fig.S7D-F), except that the empirical
information never entirely converged on the decorrelated information over the range of
time scales that we analysed.
544
We tested whether the increase in the mixed V1/V4 information might be due to its larger
pool size, compared to the V1- or V4-only pools, rather than any differences in the structure
of the noise correlations. This was tested by plotting the information of a mixed pool with
12 units against the sum of information from separate V1- and V4-only pools of size 6, each
(Fig.S7G); as in Fig.7A, the most informative units were used. The average paired difference
in information (Fig.S7H) revealed that the mixed pool had extra information, over a range of
values for τ. This difference increased with τ; hence information increased in the mixed pool
as its covariance structure changed.
553
554   *Differential correlations and receptive field separation*
555
A proper analysis of this data set requires consideration of the relative positions of the
receptive fields (RFs) recorded from V1 and V4. Due to the topographic placement of the
chronic Utah arrays, the majority of V4 RFs were more eccentric than the V1 RFs.
Furthermore, the average distance between V1 and V4 RFs was greater in one monkey
(M135) than the other (M138). During our main experiments, we presented areas V1 and V4
with a single visual stimulus that extended across the RFs of both visual areas. Both set of
RFs were therefore co-activated by a single RDS stimulus (Fig.2A).
563
Nonetheless, it could be argued that the presence of a correlation structure that is the
opposite of normal information-limiting correlations is simply due to the RF separation
rather than the fact that the recorded neurons were in different cortical areas. Further, the
apparent cancellation of information-limiting noise might have been caused by the
misalignment of V1 and V4 RFs rather than any genuine corrective mechanism. To address
these concerns, we can formulate two, testable null hypotheses. First, when compared over
a range of similar values of RF separation, there will be no difference in the structure of
within-area and between-area correlations. Second, any measurable recovery of
information by a population of V1 and V4 neurons should increase as the separation of V1
and V4 RFs increases.
574
To measure RF separation, we performed an independent quantification of the positions
and sizes of the V1 and V4 RFs. RF position was mapped in separate experiments by placing
a small random dot stereogram (RDS) at randomly selected positions on a grid of possible
locations, while the animal maintained fixation; the RDS position changed between trials.

The size of the RDS was fixed (2.5° diameter) and the binocular disparity was always 0°.
Thus, we mapped the average multi-unit firing rate for each electrode across RDS positions.
We quantified the RFs by least-squares fits of two-dimensional isotropic Gaussians to each
RF map, with centre and standard deviation (SD) of the Gaussian taken as measures of the
RF centre and size.

The Gaussian fits were reasonably accurate (median coefficient of determination *i.e.* r^2;
M135 V1 = 0.875, 95% boot CI [0.788, 0.881]; M135 V4 = 0.650, [0.517, 0.813]; M138 V1 =
0.966, [0.962,0.968]; M138 V4 = 0.862, [0.739, 0.926]), exhibiting the expected,
fundamental relationship between RF eccentricity and size (all RFs: Spearman correlation
M135 = 0.829, p → 0 ; M138 = 0.634, p = 3.351e–9).

The RF separation of pairs of neurons was defined by computing the Euclidean distance
between RF centers, in degrees of visual field. For neurons i and j (i ≠ j), RF separation was

$$RF_{separation} = \sqrt{\left(x_j - x_i\right)^2 + \left(y_j - y_i\right)^2}$$ where (x_i, y_i) was the center of the Gaussian fit to
the RF map of the i^{th} neuron, similarly for the j^{th} neuron.

Information-limiting noise correlations produce a positive relationship between measured
noise correlations and the product of tuning curve derivatives (*f'×f'*); for brevity, we shall
refer to this relationship as R_D. Our main analysis revealed a positive R_D between pairs of
neurons from the same cortical area, providing direct evidence of information-limiting noise
correlations within both V1 and V4 for encoding binocular disparity. However, pairs of
neurons from different cortical areas had, on average, a negative R_D. Plotting R_D as a
function of RF separation in Figure 8 shows that within-area correlations are generally
positive, on average, whilst correlations between areas are predominantly negative – across
the spatial separations in our sample. Neuron pairs were first binned (1 circle/bin) by their
RF separation, with at least 100 pairs/bin (N bins; M135 V1/V1 = 3, V4/V4 = 9, V1/V4 = 10
(with outlier 11); M138 V1/V1 = 19 (20), V4/V4 = 11 (12), V1/V4 = 33); binning was done
after discarding the 2% of pairs with the lowest DMI *i.e.* the least disparity selectivity. The R_D
of each bin was quantified as the Spearman correlation of measured noise correlations
(r_{CCG}(150ms)) and *f'×f'* of the bin's pairs. Likewise, the average RF separation was found for
each bin. There was no provable relationship between RF separation and differential
correlations. If anything, our data indicate that all relationships, positive and negative,
declined as the spatial separation of RF centres increased.

We then asked how much linear Fisher information loss (equation 31) there was due to
potential differential correlations (Fig.S8); this was taken to be the difference between
empirical and decorrelated information levels, normalised by the expected root of the
variance (equation 31). The amount of loss for population compositions was sensitive to the
temporal scale (τ) of the measured noise correlations (Fig.S8A&B). The greatest information
loss was seen at rapid (τ < 100ms) time scales. However, this information loss disappeared
at longer time scales (τ > 100ms), suggesting an effective cancellation of information-
limiting noise. To test the effect of RF separation on information loss, we plotted loss
(Fig.S8C) across values of τ for two pools of neurons: 1) all V1 and V4 pairs with RF
separations up to a criterion value of 7.99° (V1&Close-V4, green solid), and 2) the pool of V1
and V4 pairs with separations exceeding that criterion (V1&Distant-V4, dashed black). As

before, we found strong evidence of information loss at small values of τ in both
populations, and in both subjects. However, the V1&Close-V4 population was always the
first to reach an average loss of zero. This effect was more prominent in M135, possibly due
to the greater separation of the V1 and Distant-V4 RFs recorded from this cortex.
629
To better illustrate this point, we fixed τ at 120ms. At this temporal scale, the information
loss of the V1&Close-V4 neurons was no different from zero in either M135 (average =
+0.031, 95% Boot CI [−0.012, 0.075]; right-tail t-test p = 0.083) or M138 (−0.037 [−0.111,
0.043], p = 0.820). However, the loss of the V1&Distant-V4 neurons was significantly
positive in M135 (0.125 [0.070, 0.182], p = 7.827e−6) and M138 (0.130 [0.036, 0.231], p =
0.005). A direct comparison between the two populations was possible, with the
information loss of the V1&Distant-V4 population being significantly greater (right-tail
paired-sample T-test p, M135 = 0.002, M138 = 2.731e−5). Our results suggest that the level
of information loss increased with RF separation. They further suggest that information-
limiting noise correlations are cancelled when the V1 and V4 RFs are close together.
640
641 In summary, our data support two conclusions. First, there is a categorical difference
642 between within-area correlations and between-area correlations, regardless of RF
643 separation. Second, information recovery between V1 and V4 is stronger when RF
644 separation is smaller.
645
646 *Differential correlations and behavioural choice.*
647
In the final phase of the trial (Fig.1Aiv), the binocular depth instantaneously changed in one
of the four RDS stimuli. This popout event revealed one out of the four RDS stimuli to be an
oddball target. The monkey could get a reward by correctly choosing the target RDS with a
saccade onto that stimulus. Rewards were given after the gaze fell reliably onto the target
RDS (correct choice). Given that we recorded neuronal responses to just one of these four
RDS locations, we divided the trials into four categories of outcome, specified by
combinations of two factors: Target IN or OUT of the RF, and CORRECT or INCORRECT
choice.
656
The association between behavioural choice and the neuronal state is plotted in Figure 9.
The x-axis shows time relative to the pop-out event. As before, the state of the noise
correlation structure of the recorded neuronal responses, R_D, is summarized by the
Spearman correlation between $r_{CCG}(\tau)$ and the product of differentials $f'\times f'$. Values of τ were
30ms for the V1/V1 and V4/V4 pairs, and 110ms for the V1/V4 pairs. Owing to differences
between animals that emerged in early training, the two monkeys were exposed to either 1
second (M135) or 2 seconds (M138) of indistinguishable RDS stimuli during the Presentation
phase (Fig.1Aiii), prior to the pop-out event. In a fraction of catch trials, no depth change
occurred, to measure guess rates; these trials were discarded from this analysis.
666
The data in Fig.9 show an association between the correctness of choice and the state of
correlations, R_D, prior to the behavioural decision. The effect is strongest for the within-area
correlations of V4 (Fig.9A, middle panels, blue/black data points, fill shows two-tailed
significance at 5% level). For both animals, correct choices onto the target over the RF are
associated with stronger values of R_D in V4. This is not due to an alteration of correlations by

the target stimulus, because the target is still unknown. Second, analysis of the opposite
contingency (Fig 9B), when the target occurs outside of the RF, shows that these
correlations are also stronger for correct choices. In this case, a correct choice means
selecting a target outside of the RF and choosing the target over the RF is incorrect. For
cases, where both the target and incorrect choices were out of the RF, the R_D measure of
differential correlation was similar to when incorrect choices went into the RF (not shown).
Therefore, this association in V4 is between correctness of choice and the measured state of
differential correlations, regardless of target location. For the within-area correlations of V1
(Fig.9, left panels, gold/black data points), the same general trend is present, but the effects
are much weaker and less consistent, especially in monkey M138.

For the between-area correlations of V1/V4 (Fig.9, right panels, green/black data points), an
interesting pattern emerges. Previously in Figures 6&8, we demonstrated a negative
Spearman correlation for the V1/V4 R_D. This negative relationship is even stronger for
correct choices as opposed to incorrect choices, most notably when the target is outside the
RF (but note that target IN trial count is ≈25% of target OUT).

To quantify the significance of these changes, an all-ways ANOVA was performed on the
normalised ratio of r_{CCG} over $f'×f'$ (equation 35), separately per subject, testing the effects of
time (window position), cortical area (V1 only, V4 only, V1/V4), and choice (correct or
incorrect). The normalisation permitted the pooling of target IN and OUT trials, because
both sets show a similar dependency on correctness. The effect of choice was significant in
all but one condition (M135, V1 only, −500 to 0ms window, $p = 0.979$) when controlling for
both time and cortical area (Tukey-Kramer multiple comparisons test, $p < 0.001$).

By themselves, these results do not necessarily confirm that differential correlations are
stronger in magnitude when correct choices are made (that is, more positive for within-area
correlations or more negative for between-area correlations). Another interpretation is that
correct performance is associated with a suppression of general, non-differential
correlations within the neuronal pool, and that this suppression allows a stronger coupling
between information-limiting correlations and task performance.

More direct evidence of the potential importance of differential correlations is presented in
Figure 10. Here, the goal is to obtain a unidimensional measure of the differential
correlation in a neural population on each trial, so that its trial-to-trial relationship with
perceptual choice could be quantified. The focus is upon the change in the V1 and V4
population response that is caused by the popout stimulus. By projecting the population
response onto the optimal axis for discriminating baseline vs. popout disparity (equation
36), we reduce the N-dimensional firing rate down to a 1D estimate of the popout disparity.
This is done by taking a weighted, linear sum of the firing rates, using weights given by
$w \sim \Sigma^{-1} \Delta F$, for population covariance Σ and expected size and sign of response to the
popout, ΔF; weights were optimised according to the baseline and popout disparity on each
trial. Owing to the Σ^{-1} term, the optimal axis can mitigate non-differential noise
correlations, but any differential correlations in the population response are necessarily
projected onto the optimal axis.

The values arising from this projection were normalised so that the expected response to
the baseline disparity was 0, and the expected response to the popout disparity was always
+1, regardless of whether it is actually an increase or a decrease. If pre-popout values for
the projected response are near zero, this suggests that the population contained very little
differential correlation and accurately encoded the baseline disparity. Values between 0 and
+1 suggest the presence of differential noise that tilted the population closer to the firing
pattern expected in response to the popout disparity, while values between −1 and 0
suggest the opposite. Hence the relative direction and magnitude of perturbations in
disparity encoding could be estimated on each trial.
727
Fig.10A shows the distributions of projected responses from each animal from the
combined population of all V1 and V4 units with $RF_{separation} \leq 7.99°$ (see Fig.S8). When the
analysis window is −100ms to 0ms (grey) before the popout event, then the distributions
peak around 0, regardless of the upcoming target location. After the start of the popout
event (+35ms to +135ms window, teal), the distributions shift towards +1, for the case when
the popout target was IN the RF (top). For the other cases, when the target was OUT of the
RF, then the pre- and post-popout projection distributions were indistinguishable (bottom).
Thus, the normalised projection succeeded in representing the combination of signal and
noise in encoding the RDS disparity.
737
We asked whether perturbations in this projected response that occur just prior to the
arrival of the popout stimulus might affect the ability of the animals to detect the upcoming
change in the visual stimulus. We quantified this by computing a version of Detect
Probability (DP) (Cook & Maunsell, 2002) to measure the covariation of the normalised
projection with the perceptual choice. Projected responses from the time period −100ms-
0ms were now grouped according to behavioural choice (Fig.10B top, Choice IN (open) and
OUT (filled) of the RF). These distributions, categorized by future choice, allow the
computation of receiver operating characteristic (ROC) curves (Fig.10B bottom, green),
where the choice IN rate is plotted on the vertical axis of the ROC curve and choice OUT is
on the horizontal axis. The DP is the area under this ROC curve (equation 40), such that a DP
> 0.5 indicates that larger, more positive perturbations preceded choices IN to the RF, while
DP < 0.5 suggests the opposite and DP = 0.5 indicates no effect on detection. Fig 10B shows
that both ROC curves dip below the chance line (black). This means that positive
perturbations that mimic the arrival of the target stimulus and occur just beforehand have a
negative effect on detection rates for the target.
753
It is interesting to note that the portions of the probability distributions of the projected
response that are most relevant for this effect on behaviour lie exactly within the bounds
that are also relevant for the neural detection of the target stimulus. The relevant positions
are marked as the baseline (empty cross) and popout (filled cross) disparities and are simply
transposed from the vertical lines in Fig 10A. The perturbations in the pre-popout period
have their greatest impact on behaviour, producing a DP < 0.5, when they occur within
these boundaries.
761
In Fig.10C, we computed DP for populations of units, with $RF_{separation} \leq 7.99°$, for V1 only
(gold), V4 only (blue), or mixed V1&V4 units (green), using responses measured from a
range of pre-popout windows (−400 to −200, −200 to 0, −100 to 0ms), with weights

optimised using either fast ($\tau = 20$ms) or slow ($\tau = 150$ms) noise correlations; values were
computed separately for trials with the target IN the RF (top) or OUT (bottom). We found
that the only difference in DP that was consistent across subjects was in the combined V1
and V4 responses just prior (−100 to 0ms) to the appearance of the popout target IN the RF
(arrows). Notably, the relevant temporal scale for these noise correlations was fast, at
20ms. This suggests that rapid, pre-popout differential correlations that spanned both V1
and V4 are detrimental to detecting the change in disparity.

This is a key prediction of the role for differential correlations in limiting the detection of the
stimulus. When the state of the network before the arrival of popout is such that it more
closely resembles the state of the network after the popout stimulus arrives, then this is
detrimental to perception of the disparity change. On trials when the fluctuations in
network state place it further away from the neural response to the popout, then detection
of the popout target is less impaired. In line with Figs.6,7,&S8, our results suggest that it is
the rapid differential correlations that do indeed degrade behavioural detection. Slower
correlations prior to the popout event do not appear to impair detection, suggesting that
integrative processes in the visual cortex may have enough time to attenuate rapid
differential fluctuations.

A model V4 output neuron can attenuate differential noise

Here we consider how the joint actions of V1 and V4 could preserve information
transmission in the presence of information-limiting correlations within each individual
cortical area. The finding of pairs of V1 and V4 units that had opposite disparity preferences
and positive noise correlations (*e.g.* Fig.3D–F, Fig.S3 left) provides evidence for one route to
cancelling the correlated noise without reducing the sensory signal (Abbott & Dayan, 1999;
Averbeck et al., 2006). This phenomenon has been observed experimentally in the second
order areas of the somatosensory system (Romo, Hernandez, Zainos, & Salinas, 2003).

To investigate this further, we built a simple model of a V4 output neuron (Figure 11,
equations 37-39), of a type that might project to downstream areas. The model had two
afferent neurons (Fig.11A), a neuron that projects from V1 onto the V4 output neuron and
an intrinsic, local-circuit neuron from within V4 that also connects to the V4 output neuron.
The details of this could be varied without a loss of generality, but importantly this
represents a simple feedforward system that can explain many of our results.

Correlated noise was removed by subtracting the weighted V1 responses from the input V4
responses, each input integrated across a Δt ms window, to produce the V4 output
responses. Each V1 and V4 pair in our dataset was used to simulate the responses of a
model V4 output neuron, whose disparity selectivity was then measured. The prediction is
that the V1/V4 input pairs with positive noise correlation ($r_{noise} > 0$) and negative signal
correlation ($r_{signal} < 0$) should yield V4 output neurons with better disparity selectivity
(Fig.11B, V4 input variance, dashed, output V4 variance, grey).

We test this circuitry directly with empirical data by measuring the disparity selectivity of
each simulated V4 unit, using the area under the receiver operating characteristic curve

(AROC, equation 40) as a measure of neuronal sensitivity. Focusing on the Initial analysis
window, the difference in AROC (equation 41) between the V4 model output and input
(Fig.11C left, thick solid, Not shuffled) confirmed the prediction above, where positive
differences indicate model V4 output neurons with better disparity sensitivity.

As stated, a pair could improve its stimulus sensitivity either by changing its correlation
structure, or by producing a larger modulation of firing in response to stimulus changes.
Hence, it is possible that the superior performance of the model V4 output neuron was
trivially due to the combined signalling of the V1 and V4 input neurons, rather than any
cancellation in correlated noise. We tested this by recomputing the model V4 outputs after
shuffling the afferent V1 responses between trials such that the V1 unit maintained its
stimulus sensitivity, but the noise correlations were destroyed (thin solid, V1 tuning kept). In
this scenario, the model V4 neuron can only improve its sensitivity through combined input
signals, but not through cancelling out the correlated noise of its afferents.

Model V4 neurons had better stimulus sensitivity than the afferent V4 neuron, whether
there was shuffling (thin solid) or not (thick solid). Yet, the model V4 neurons with no
shuffling achieved a further increase in stimulus sensitivity in comparison to those with
shuffling. This improvement in sensitivity occurred specifically when the afferent neurons
had opposite disparity selectivity ($r_{signal} < -0.5$, mean difference AROC = 2.961e−3, paired t-
test p = 4.974e−112); in contrast, the model without shuffling suffered a reduction in
sensitivity when the afferent neurons had similar disparity selectivity ($r_{signal} > +0.5$,
−7.894e−4, p = 3.269e−10). This is precisely the pattern that should be observed when the
afferent V1 and V4 neurons have positive noise correlations (Abbott & Dayan, 1999;
Averbeck et al., 2006). Additional stimulus sensitivity is available by cancelling out the noise
correlations of afferent neurons with opposite stimulus selectivity.

As differential correlations seemed to be reduced by slower, V1 to V4 correlations
(Fig.7&S8), we asked if the simulated V4 output neuron would improve its performance with
a wider integration window. Thus, we evaluated the Initial V4 output responses over a set of
Δt values. For each Δt, the un-shuffled model V4 sensitivity (Fig.11D, left) was subtracted by
the shuffled model V4 sensitivity; again, this tests for changes in sensitivity due to the
cancellation of afferent noise correlations. When the input V1/V4 pair had $r_{signal} < -0.5$,
there was a significant tendency for sensitivity to increase as Δt increases (Δt = 20ms vs
160ms, mean paired difference = 1.540e−3 , p = 9.490e−25).

For the Steady state responses, the AROC measure detected a weak improvement (Fig.11C,
right) when the input pair had opposite disparity preferences (Not shuffled vs V1 tuning
kept, mean paired difference = 4.9181e−4, paired t-test p = 3.1995e-26). No improvement in
selectivity was found with increasing Δt (Fig.D right, Δt = 100ms vs 800ms, mean paired
difference = −3.902e−4, p = 7.986e−16).

DISCUSSION

Cortical areas V1 and V4 contain noise correlations that could limit the information that those areas convey about stereo depth. This noise limits information chiefly at short time scales, as the animal is discriminating depth. Unexpectedly, noise correlations between the two areas were found to have a different structure at longer time scales, which could reflect the brain's capacity to limit correlations that might become harmful to perceptual performance. We show that the state of short-time course noise correlations has a direct association with the animal's performance, demonstrating a critical piece of evidence that the structure of correlations has a direct impact on behavioural decisions and sensory performance.

Detecting and interpreting differential correlations

Empirical evidence of differential correlations has rarely been reported, due in part to the prediction that they are so small (Moreno-Bote et al., 2014) that thousands of neurons and thousands of trials are needed to detect them (Kanitscheider et al., 2015b; Kohn et al., 2016).

But the magnitude of noise correlations is dynamic, and changes with a number of factors (Marlene R. Cohen & Kohn, 2011), including state of consciousness (Ecker et al., 2014), stimulus presentation (Churchland et al., 2010), perceptual grouping (Bondy et al., 2018; M. R. Cohen & Newsome, 2008; Poort & Roelfsema, 2009; Wasmuht, Parker, & Krug, 2019), attention (M. R. Cohen & Newsome, 2009; Mitchell, Sundberg, & Reynolds, 2009; D. A. Ruff & Cohen, 2014; Verhoef & Maunsell, 2017), and perceptual learning (Ni et al., 2018; Sanayei et al., 2018). Our experiment brought many of these global factors under control by using awake subjects that performed a demanding perceptual task. Indeed, we presented evidence (Figs.9&10) that the animal's fluctuations in perceptual performance may reflect the impact of differential correlations on vision perception.

Another advance we have made in detecting differential correlations is in how they are measured. Spike count correlation (r_{SC}) is often used despite the fact that correlations vary over temporal scales (Bair et al., 2001; Denfield, Ecker, Shinn, Bethge, & Tolias, 2018; Kohn & Smith, 2005; M. A. Smith & Kohn, 2008; M. A. Smith & Sommer, 2013). If differential correlations are rapid, then they could be masked by the slower correlations that are captured by r_{SC}. By using the r_{CCG} measure introduced by Bair et al. (2001), we found evidence of differential correlations operating at tens of milliseconds. This is much shorter than the width of the analysis windows that have typically been used to measure r_{SC} (Marlene R. Cohen & Kohn, 2011).

A complication is that the time scale of noise correlation is dynamic, and can change within an area due to stimulation or behavioural state. This can be seen by comparing our results with previous studies of V1 (Kohn & Smith, 2005; M. A. Smith & Kohn, 2008) and V4 (M. A. Smith & Sommer, 2013) that used r_{CCG}. They found noise correlations that saturated around 100ms during stimulation, in both areas; but the subjects were either anaesthetized (Kohn & Smith, 2005; M. A. Smith & Kohn, 2008) or passively fixating (M. A. Smith & Sommer, 2013).

We saw a similar r_{CCG} profile in the spontaneous responses of V1; but with visual stimulation

presented under a task-relevant contingency, r_{CCG} saturated at 20ms. In V4, saturation was

also near 20 ms – before and during stimulation. Our results here are more closely

compatible with other cases in which r_{CCG} also saturates at tens of ms in the dorsal area

V5/MT when monkeys perform a perceptual task (Bair et al., 2001; Wasmuht et al., 2019).

Thus, the rapid, differential correlations that we observed here may have only been readily

detectable during the performance of a perceptual task.

Recording depth and topography

We used 1mm electrodes, with the result that our recordings were likely from the

supragranular layers near the layer 3-4 border, from a mixture of neurons that projected

nearby or to other cortical areas (Harris & Mrsic-Flogel, 2013). The correlations may have

carried a stronger differential component if many neurons projected locally. This stems from

the theory that differential correlations build up between neurons that share some of the

same noisy afferents (Kanitscheider et al., 2015b). Given the functional heterogeneity across

layers in V1 and V4 (Buffalo, Fries, Landman, Buschman, & Desimone, 2011; Denfield et al.,

2018; Nandy, Nassi, & Reynolds, 2017; Pettine, Steinmetz, & Moore, 2019), cross-laminar

recordings will be required to resolve this issue.

Some of the signal from V1 will have been relayed to V4 by other areas; but there are direct

axon projections from V1 to V4 and from V4 to V1 (Ungerleider, Galkin, Desimone, &

Gattass, 2008). Although we did co-activate the V1 and V4 neurons with a single stimulus,

the measured V1 RFs were not perfectly coincident with those of V4. Hence, some of the

excitatory drive for the V4 neurons presumably came from other, unmeasured V1 neurons.

Importantly, we saw qualitatively similar V4 correlations in a set of control experiments with

the RDS centred on the aggregate V4 RF (see Figs.S1&S2, S4-S7).

In V1 and V4, noise correlations span a range of cortical (*i.e.* retinotopic) distances when

neurons respond to the same stimulus (M. R. Cohen & Newsome, 2009; Poort & Roelfsema,

2009; M. A. Smith & Kohn, 2008; M. A. Smith & Sommer, 2013) – which we observed (mean

r_{CCG}(20ms); V1 pairs < 1.44mm apart = 4.987e−2; V1 > 1.44mm = 1.805e−2; V4 < 1.44mm =

3.416e−2; V4 > 1.44mm = 2.279e−2; all t-test $p \to 0$; median distance = 1.44mm). Moreover,

noise correlations in both areas increase when neurons respond to a common stimulus

amongst an array of stimuli (Poort & Roelfsema, 2009; D. A. Ruff & Cohen, 2014). Thus, it is

likely that the correlations of perceptually grouped neurons vary systematically across the

cortex. We suggest that by stimulating V1 and V4 with the same object that we increased

the likelihood of the visual system having grouped them into a perceptual pool.

Mechanisms of noise removal

In a sense, our results are puzzling. Neuronal correlations in V1 and V4 show positive

relationships between the neuron pairs with similar stimulus tuning and tuning curve slope

(differential correlations). This kind of pattern is reported for motion in V5/MT and

orientation in V1. With simple feedforward connections, one ought to observe the same

relationship between V1 and V4. However, the general pattern of V1/V4 correlations is
inverted (Fig.5E&F,6I,8).
948
Noise correlations like that can be removed linearly without losing information (Abbott &
Dayan, 1999; Averbeck et al., 2006). Thus, our simple model (Fig.11) suggests that the
combined action of V1 and V4 attenuates the local correlations through linear operations.
However, differential correlations cannot be linearly removed by later processing
(Kanitscheider et al., 2015b; Moreno-Bote et al., 2014; Pitkow et al., 2015). One way of
resolving this discrepancy rests on our result that the covariance structure between areas is
not static, but evolves across temporal scales. Notably, the V1/V4 pattern emerged at
slower time scales than the rapid differential correlations within each area.
957
Related results come from MT (M. R. Cohen & Newsome, 2008) and V4 (D. A. Ruff & Cohen,
2014) in which positive noise correlations strengthened between neurons with opposite
tuning (but see Bondy et al., 2018). In V1, positive correlations that increase the information
in linearly-pooled responses can appear between neurons that respond to different stimuli
(Poort & Roelfsema, 2009). Lastly, to change the V1 to V4 correlation structure in a trial
would require the kind of fast and highly specific adaptations that are observed between
them in attention tasks (Grothe, Neitzel, Mandon, & Kreiter, 2012; Rohenkohl, Bosman, &
Fries, 2018).
966
Adaptable covariance may imply an underlying, non-linear computation. But what could be
biologically plausible? A spike-count variance code could overcome correlations if correlated
inputs were high-pass filtered in a spatial domain (*e.g.* across neurons) followed by a
quadratic non-linearity to compute variance (Shamir & Sompolinsky, 2004). Another
potential choice is divisive normalisation (Carandini & Heeger, 2012), which can model the
structure of correlations between areas V1 and MT (Douglas A. Ruff & Cohen, 2016).
Divisive normalisation might decorrelate responses in two ways (see Eq. 3.1 in Tripp, 2012).
First, spike count variance can be saturated by the correlated variance of other neurons,
attenuating correlations in the normalised response. Second, the normalised responses of
correlated units can move in opposite directions to their raw values, decorrelating them.
One mechanism need not exclude others, and the brain might profit from the parallel action
of several computations.
979
980 *Sources of differential correlations*
981
Present analysis suggests both bottom-up (Kanitscheider et al., 2015b) and top-down (Ecker,
Denfield, Bethge, & Tolias, 2016) origins of differential correlations. In the former, stimulus
noise is sent through diverging projections to downstream areas; differential correlations
then build up between neurons with common noisy inputs (Kanitscheider et al., 2015b),
especially if sub-optimal computations are used (Beck, Ma, Pitkow, Latham, & Pouget,
2012). In the latter, noise in the selection of an attended feature value can lead to a gain
profile that shifts position relative to the stimulus preferences of a population of neurons
– inducing differential correlations (Ecker et al., 2016).
990
991 Our results invite a compromise. Both sources are present, but take effect at different
992 moments. The initial burst of firing is a bottom-up event in which information is most

reduced by rapid noise correlations, which are unlikely to have a top-down origin (Denfield et al., 2018). In our data, only the Initial responses could reach the decorrelated information level, at a temporal scale of 100ms. The Steady state information did increase at wider temporal scales, yet failed to hit the decorrelated level within an 800ms time scale. One interpretation is that rapid differential correlations in the Initial response had a bottom-up source, followed by slow differential correlations with a top-down source that could not be locally attenuated. This fits well with the common reports of top-down or contextual effects that follow the initial response of visual neurons to a stimulus (Roelfsema, Lamme, & Spekreijse, 1998) (Pack & Born, 2001) (M. R. Cohen & Newsome, 2009; Herrero, Gieselmann, Sanayei, & Thiele, 2013; Nienborg & Cumming, 2009; Poort et al., 2012; J. E. T. Smith et al., 2015).

Neural pooling and perception

The statistical relationship between a pool of sensory neurons and perceptual behaviour is connected to the noise correlations within the pool (Haefner, Gerwinn, Macke, & Bethge, 2013; Nienborg, Cohen, & Cumming, 2012; Andrew J Parker, 2013; Pitkow et al., 2015; Shadlen et al., 1996). That relationship may evolve over time, with a sensory pool driving a brief, bottom-up effect on behavioural choice, followed afterwards by a top-down reflection of choice bias (J.E.T. Smith, Masse, Zhan, & Cook, 2012). These effects are difficult to separate in many experiments, which employ long stimulus viewing times (Britten et al., 1996; Dodd, Krug, Cumming, & Parker, 2001; Nienborg & Cumming, 2009; Pitkow et al., 2015; Poort et al., 2012; Yu & Gu, 2018).

However, a subject may base perceptual decisions on only a few hundred ms of viewing in natural conditions (M. R. Cohen & Maunsell, 2009). Hence, studies using mixtures of reaction time tasks and short-duration stimuli often observe that the transient response is accounted for by a small pool of neurons (Ghose & Harrison, 2009; Harrison, Weiner, & Ghose, 2013; Krause & Ghose, 2018; K. F. Weiner & Ghose, 2015) with a bottom-up link to behaviour (Price & Born, 2010; Jackson E. T. Smith, Zhan, an, & Cook, 2011; Katherine F. Weiner & Ghose, 2014). Our results suggest that the transient response is a high-fidelity signal that could be decoded from a small pool of neurons. Later, top-down signals may arrive that impose slow and widespread differential correlations in a larger pool, thus injecting choice correlations into neurons that are never read out (M. R. Cohen & Newsome, 2009; Nienborg & Cumming, 2010; Nienborg & Cumming, 2009; Pitkow et al., 2015; Yu & Gu, 2018).

Although the associations between behavioural decisions and network state were stronger in one monkey than the other, the timing of these changes of state, in terms of differential correlations, supports this interpretation. Rapid fluctuations in the state of the network that mimic the change of state that occurs upon the presentation of the discrimination stimulus are detrimental to stimulus detection. This is a fundamental prediction of the theoretical role of differential correlations (Pouget et al., 2003) that we have been able to validate with these experimental measures.

Correlations and the organizational structure of the neocortex

Our results lead us to propose a general framework for thinking about the effect of noise
correlations on neuronal processing. First, within-area noise correlations that potentially
limit sensory discrimination are a by-product of local information-processing (Harris &
Mrsic-Flogel, 2013; Kanitscheider et al., 2015b). Second, we suggest that the signalling
between areas acts to remove the harmful consequences of these local, intrinsic
correlations. The removal process may use at least two strategies: one is the connectivity
between cortical areas (*e.g.* Figure 8) and the other is dynamic processing that adapts to the
ongoing statistics of the neural signals. An implication of this framework is that one function
of keeping diverse neocortical areas may be to contain the impact of differential
correlations that arise from local neuronal processing within the circuitry of individual areas.
1050

1051   ACKNOWLEDGMENTS
1052
The authors wish to thank Andy Emberton and the staff at The University of Oxford's
Biomedical Services, Daniel Heaton and the DPAG workshop for expert technical support,
plus James Bennett for help with the lab setup. Thanks go to Ruben Coen-Cagli for help in
understanding the computation of bias-corrected linear Fisher information, and also to
Alexandre Pouget, Bruce Cumming and Robert Dayan for their insightful conversations. This
work was funded by MRC grant MR/K014382/1.
1059
1060   AUTHOR CONTRIBUTIONS
1061
1062   AP was the principal investigator and designed the study. JS collected and analysed the data.
1063   JS and AP wrote the manuscript.
1064
1065   DECLARATION OF INTERESTS
1066
1067   The authors declare no competing interests.
1068

METHODS

Animals

We recorded from two adult male Rhesus macaque monkeys (*Macacca mulatta*) acquired from the UK Public Health England breeding colony at Porton Down through the Centre for Macaques (M135 and M138). Each animal was implanted with a titanium headpost (Gray Matter Research, Bozeman, MT) and a CerePort pedestal (Blackrock Microsystems, Salt Lake City, UT) in separate, aseptic surgical procedures under general anaesthesia. Each CerePort was connected to a pair of 64-channel Utah arrays that were implanted subdurally in the upper layers of the exposed cerebral cortex. The exact locations for surgical implantation were determined by a preliminary magnetic resonance imaging (MRI) scan of the animal's head to reveal bone and soft tissue structures.

Animals were trained to fixate on a target and then to perform an odd-one-out task using saccadic eye movement responses. All procedures were done in compliance with Home Office (UK) regulations. There was an ample purveyance of victuals. The key results were qualitatively similar in both animals, and so we have pooled the data across animals, unlsess otherwise stated.

Visual stimulus

Visual stimuli were presented with a Wheatstone stereoscope comprising of a pair of CRT monitors (Flexscan F78, Eizo, Hakusan, Japan; or P225f, ViewSonic, Brea, CA) at 84cm from the subject's eyes. Cold mirrors were used to reflect images into the eyes so that the infrared light from an eye tracking camera (Hi Speed Primate, SensorMotoric Instruments GmBH, Teltow, Germany), set behind the mirrors, could pass through. Monitors had a resolution of 1600 x 1200 pixels, a refresh rate of 85Hz, and subtended 26.7° × 20.1° of the subject's visual field. Each pixel was 1.6e−2° wide. The VGA signal was split and duplicated, so that the left-eye monitor received three copies of the red channel for its RGB input, while the right-eye monitor received three copies of the green channel. When viewed on a conventional single colour monitor by human observers, the stereo images were visible as red/green anaglyph images; but viewed through the stereoscope, the images appeared as grayscale images presented to each eye of the subjects. Thus, the Wheatstone stereoscope produced no stereoscopic cross-talk.

Quadro K2200 (NVIDIA, Santa Clara, CA) video cards in multi-core Intel (Santa Clara, CA) processor computers running Ubuntu 14.04 (Linux kernel 4.4.0 lowlatency) were used to drive the monitors. Stimuli were programmed using PsychToolbox (3.0.14) (Brainard, 1997) running in Matlab R2015b (The Mathworks, Natick, MA). All stimuli were presented on a mid-gray background, using appropriate OpenGL alpha-blending (GL_SRC_ALPHA + GL_ONE_MINUS_SRC_ALPHA) to obtain sub-pixel resolution. A 4 × 4cm square was shown in the upper-left corner of each monitor during trials. This was recorded with a photodiode (PIN-25DP, OSI Optoelectronics, Hawthorne, CA) by the electrophysiology system, in order to synchronise high-precision frame time stamps from PsychToolbox with the neural activity. The square was blocked from the subject's view. A full white, 0.3° diameter circular dot was used as the gaze fixation target.

1116
Random dot stereograms (RDS) were used to present binocular disparities for the
behavioural, odd-one-out task and to stimulate recorded neurons. Each RDS comprised of a
circular patch of 0.16° diameter circular dots. Dot positions were randomly sampled on each
frame; the frame rate matched the refresh rate of 85Hz. Four RDS were presented, but dot
positions were sampled independently for each RDS. Half of the dots were black, and half
were white. Dots occluded each other in a random order. Each RDS had two regions, a 6°
diameter circular centre and a 1° wide annular surround. The dot density in both regions
was 25% (area occupied by dots / total area), assuming no dot overlap; in other words, the
number of dots per RDS was 25% of the RDS area divided by the area of a single dot,
rounded up. The RDS centre varied in binocular disparity such that a flat circular plane of
dots was shifted convergently (negative disparities, toward subject) or divergently (positive,
away). Surround dots were always on the fixation plane at zero disparity *i.e.* the surface of
the monitors' screens. The surround functioned to mask monocular cues when non-zero
disparities were presented in the centre, and to help the subject maintain vergence. When
the RDS centre was at a non-zero disparity, each of its monocular images had a horizontal
shift that overlapped the RDS surround on one side while leaving a gap on the other.
Surround dots in the overlap region were discarded, while uncorrelated dots were used to
fill the gaps. Thus, the resulting monocular images carried no trace of the disparity of the
RDS centre, so that the subjects could only perform the task using the stereoscopic
information.
1137
1138 **Task**
1139
Subjects performed an odd-one-out detection task (Figure 1A) that was divided into four
epochs. At the start of each trial, a fixation point was presented (Fig.1Ai). The subject then
had to maintain its gaze at the fixation point for 0.5 second to initiate the trial (Fig.1Aii).
Following this, four RDS were presented for 1 second (M135) or 2 seconds (M138), one in
each quadrant of the visual field (Fig.1Aiii). Subjects had to maintain their binocular gaze
within a 0.75° radius of the centre of the fixation dot to both initiate the trial and progress
through to the end of the presentation phase. During the presentation phase, all four RDS
had the same baseline disparity in their central region. Baseline disparities were randomly
selected on each trial with equal probability from a set of 13 values: 0°, ±0.01°, ±0.02°,
±0.05°, ±0.10°, ±0.20°, ±0.50° (similar to Shiozaki, Tanabe, Doi, & Fujita, 2012). At the end of
the presentation period, one of the four RDS had a disparity step change in its central
region, gaining an additional 0°, ±0.05°, or ±0.10° (Fig.1Aiv). Both the location and value of
the step change were randomly selected on each trial with equal probability. Thus, the RDS
that changed became the odd-one-out (*i.e.* the popout target or oddball), which the subject
was required to detect by making a saccade onto its location. With four possible target
locations, the probability of correctly selecting the target by chance was 0.25 (Figure 1B). A
juice reward was given for correct performance. No reward was given for incorrect
saccades, or if the subject responded prior to the step change. After the training or testing
session, animals had ad libitum access to water in the home enclosure for a limited period
of time every day. Animals were weighed each day and assessed.
1160
1161
1162

Recording

The animals' binocular gaze positions were sampled at 500Hz and recorded alongside the neural responses. In cortex, each subject was implanted with two Utah arrays (Blackrock Microsystems). Both arrays had 64 platinum electrodes, each 1mm in length, arranged in an 8 × 8 grid with 400μm spacing. It is unclear exactly which layer of the cortex that each electrode tip had entered, due to the curvature of the cortical surface relative to the flat array, and from slight deviations of the arrays from the plane tangential to the cortex. Nonetheless, given the electrode lengths, the presumption is that the recordings reported here are primarily from the upper, supragranular layers of the cortex. One array was implanted in cortical area V1 and the other in V4. A CerePlex E headstage (Blackrock Microsystems) was attached to the CerePort implant. The headstage applied 0.3Hz 1^{st} order high-pass and 7.5kHz 3^{rd} order low-pass anti-aliasing Butterworth filters to the analogue signals before 16-bit digitizing all 128 channels at 30kHz. Digitized signals were transmitted to the Cerebus (128 channel electrophysiology system, Blackrock Microsystems) using fibre optics, thus minimizing noise at the analogue stage. All computers, monitors, and the reward pump were located in a separate control room while the subject, headstage, stimulus monitors, digital hub, and CRTV camera were within a test room that had Faraday shielding. A 2^{nd} order 250 to 5000Hz bandpass digital Butterworth filter was applied online to each data channel before applying a threshold of −6.5 times the root-mean-squared (RMS) of the estimated noise level to detect action potentials (Rizk & Wolf, 2009); the thresholds were newly estimated on each experiment for each electrode. The −6.5×RMS threshold was the default setting for the Cerebus, which worked well in practice to detect multi-unit activity without saturating the acquisition system and losing data.

The simultaneous V1/V4 data in this study came from 6 recording sessions with M135 and 4 sessions with M138, yielding 8279 trials from M135 (1380 per-session mean [60.7 standard deviation]) and 6073 from M138 (1518 [58.0]). Approximately similar numbers of trials with each baseline disparity were obtained. We made no assumptions about the identity of each unit, treating each day's recording separately. Here, the term 'unit' simply means a cluster of spike waveforms from one electrode (see below). This yielded a total of 63 V1 (10.5 [3.2]) and 111 V4 (18.5 [3.5]) units from M135, and 132 V1 (33.0 [3.5]) and 102 V4 (25.5 [3.7]) units from M138. Thus, we retrieved 315 V1 (52.5 [32.6]), 975 V4 (162.5 [63.5]), and 1206 inter-cortical (201.0 [91.6]) neuron pairs from M135 and 2104 V1 (526.0 [108.4]), 1251 V4 (312.8 [90.6]), and 3392 inter-cortical (848.0 [183.7]) pairs from M138. All pairwise analyses compared units from separate electrodes (Figures 3 – 6 & 8).

In order to stimulate both V1 and V4 receptive fields (RF) with the central region of the lower-right RDS, the array of all four RDS stimuli was sometimes shifted upwards and to the left, in which case they had an asymmetrical arrangement relative to the fixation point (Figure 1C). A second set of V4 control data was obtained in which the lower-right RDS was centred on part of the aggregate RF of the V4 Utah array (Figure S1); in this case, the four RDS locations were symmetrically placed around the fixation point. In these sessions, 11 baseline disparity values were used, including: 0°, ±0.05°, ±0.1°, ±0.2°, ±0.3°, ±0.4°. The task was slightly different in that the odd-one-out did not have a disparity step change. Rather, it reversed the binocular correlation of its central dots from +100% correlated to -100%. That is, dots had the same contrast in the left and right-eye images during the presentation

period. After the presentation period, odd-one-out dots in the left image were paired with
dots of opposite contrast in the right image. However, we only analysed responses to the
+100% correlated dots, prior to the popout phase of the trial. Thus, despite this change of
task, the control data were obtained under similar sensory stimulation conditions to the test
data, and the animals had to attend to the stimuli to do the task. For M135, the diameter of
each RDS centre was 4°. For M138, it was 1°. 7 control sessions were recorded from M135,
yielding a total of 5257 trials (751.0 [135.7]), 413 V4 control units (59.0 [8.4]) and 1536 pairs
(219.4 [88.4]). 3 control sessions were recorded from M138, yielding 2082 trials (694.0
[240.8]), 266 V4 control units (88. 7 [15.3]) and 1101 pairs (367.0 [181.7]).
1219
1220 **Analysis**
1221
1222 The data were analysed using programs that were written and executed in Matlab R2015b &
1223 R2019a.
1224
1225 *Pre-processing*
1226
In both animals, we found incidences of cross-talk between a sub-set of electrodes. This was
apparent as a fraction of tightly synchronous waveform times between certain pairs of
electrodes, defined as waveforms occurring within ±1.3e−4 second *i.e.* ±4 samples at 30KHz
of each other. Synchronous waveforms often had highly similar, stereotypical shapes.
Testing with a digital neural signal simulator (Blackrock Microsystems) indicated that all
components downstream of the CerePort functioned properly. Subsequent testing by the
manufacturer suggests that there was a larger tolerance of fit between some pairs of
CerePort and CerePlex E than had been specified, causing small amounts of misalignment on
some recording days that could explain the emergence of cross-talk on some groups of
channels. Thus, individual electrodes in the Utah arrays were discarded from the analysis if
they had at least 6% synchronous spikes with any other electrode and the median Pearson
correlation of the synchronous waveforms was 0.5 or more. An electrode was also discarded
from analysis, if there was no significant change in the multiunit spiking rate following onset
of the RDS stimulus. This was defined by counting spikes in two 500ms wide analysis
windows, one from −500ms to the appearance of the RDS and the other from 50ms to
550ms. A paired-sample t-test was used to compare the firing rates before and after RDS
appearance at a significance level of 5%. When a significant difference was found then the
electrode's spikes were sorted for further analysis. On average, 9 V1 electrodes (minimum
5, maximum 14) and 14 V4 (min 12, max 17) electrodes from M135 were used, while 27 V1
(min 22, max 31) and 20 V4 (min 19, max 23) electrodes were used from M138 per session.
1247
Spikes detected online with the Cerebus were sorted using our implementation of the
clustering algorithm by Fee, Mitra, and Kleinfeld (1996) and the heuristic choices of
UltraMegaSort2000 (Hill, Mehta, & Kleinfeld, 2011; https://neurophysics.ucsd.edu). An
additional step was taken after aligning the spike waveform peaks but prior to the principal
component analysis. Each waveform was weighted by a Gaussian shaped window with a
standard deviation of 1ms that was centred on the waveform peak; area under the Gaussian
that overlapped the waveform was normalized to preserve the overall magnitude of the
spike. This emphasised the peak region of the waveforms during sorting. Automated spike
sorting was then verified and corrected manually. We based our analyses on both single

units and multiunit clusters and use the term 'unit' to refer to either, as in other work with
Utah array electrodes (D. A. Ruff & Cohen, 2014).

1259

1260 Pre-screening disparity tuning curves were found for each unit, using spike rates from an
1261 analysis window of 50 to 1050ms (M135) or 50 to 2050ms (M138) following onset of the
1262 RDS stimuli. Only units with significant disparity tuning (one-way ANOVA, $p < 0.01$) were
1263 kept for further analysis. Additionally, any unit with a tuning curve peak of less than 5
1264 spikes/second was also discarded. These tuning curves were not used in the main analysis.

1265

1266 *Single unit analysis*

1267

1268 The selectivity of a unit for binocular disparity was quantified in two ways. Disparity
1269 discrimination index (DDI; Prince et al., 2002) was used to validate disparity mutual
1270 information (DMI). DDI was defined as:

1271

1272
$$\frac{N_{max} - N_{min}}{(N_{max} - N_{min}) + 2R_{error}} \tag{1}$$

1273

1274 where N_{max} and N_{min} were the largest and smallest responses of the unit's disparity tuning
1275 curve, and R_{error} was the square root of the residual variance around the means over the
1276 whole tuning curve. DDI was computed using the square root of the firing rates (as in Prince
1277 et al., 2002).

1278

1279 DMI was computed non-parametrically (Wallisch et al., 2014) by first estimating the joint
1280 probability distribution of the disparity and spike count using 13 evenly spaced bins in both
1281 marginal dimensions to get a two-dimensional histogram (one marginal bin per disparity
1282 value, 11 marginal bins used for V4 control data). Disparities were ranked ordinally before
1283 binning. Joint probabilities were empirically estimated by:

1284

1285
$$P_{xy}(i,j) = \frac{1}{M} \sum_{m=1}^{M} J_m(i,j) \tag{2}$$

1286

1287 where M was the total number of trials and $J_m(i,j)$ had a binary value:

1288

1289
$$J_m(i,j) = \begin{cases} 1, \text{ordinal disparity was in bin } i \text{ and spike count was in bin } j \text{ on trial } m \\ 0, \text{otherwise} \end{cases} \tag{3}$$

1290

1291 The marginal probabilities were computed in a similar manner. For ordinal disparity values
1292 (P_x) and spike counts (P_y), they were:

1293

1294
$$P_x(i) = \frac{1}{M} \sum_{m=1}^{M} x_m(i) \tag{4}$$

1295

1296
$$P_y(j) = \frac{1}{M} \sum_{m=1}^{M} y_m(j) \tag{5}$$

1297

1298 where $x_m(i)$ and $y_m(j)$ had a value of 1 if the data fell into the respective bin on trial m,
1299 and 0 otherwise. The raw empirical mutual information was estimated by:

1300

1301
$$I_{raw} = \sum_{i=1}^{B}\sum_{j=1}^{B} P_{xy}(i,j)log_2 \frac{P_{xy}(i,j)}{P_x(i)P_y(j)}$$
(6)

1302

1303 where B was the number of bins along the marginal. However, this information estimate
1304 will be biased – without further correction. The correction was found by applying equation
1305 (6) to randomly shuffled copies of the data. Using 30 repeats (Hatsopoulos, Ojakangas,
1306 Paninski, & Donoghue, 1998), a shuffle-correction term was estimated as:

1307

1308
$$I_{shuffle} = \frac{1}{30}\sum_{s=1}^{30} i_{shuffle}(s)$$
(7)

1309

1310 where $i_{shuffle}(s)$ was the output of equation (6) following the s^{th} random shuffling of the
1311 data. The final, corrected mutual information between the disparity and spike count was
1312 taken as:

1313

1314
$$I = \frac{I_{raw} - I_{shuffle}}{\Delta t}$$
(8)

1315

1316 where Δt was the width of the spike-counting analysis window, in seconds, providing a
1317 value in bits/second.

1318

A time series analysis of firing rates and DMI was conducted on spike trains that were
convolved with a causal exponential kernel (20ms time constant). DMI was measured
independently for each time point using Δt = 0.02s (Eq.8) to estimate the instantaneous rate
of information. The firing rate and DMI time series were used to define three analysis
windows for the pairwise and population analyses. The Spontaneous window started 500ms
before RDS onset, ended 50ms after RDS onset, and covered the period of gaze fixation
prior to the units' first responses to the RDS. The Initial window was located to cover the
first transient response to the RDS, which was measured from 50ms to 200ms after the RDS
appeared. A final Steady state window ran from 200ms to 1050ms for M135 and from
200ms to 2050ms in M138, covering responses to the RDS up to but not including those to
the disparity step-change.

1330

1331 *Paired signal and noise correlations*

1332

1333 In the analysis of pairs of neurons, we focused on three key metrics, as defined by Bair et al.
1334 (2001). First was the degree to which the firing rate of two units was driven by the same
1335 stimulus values – the pair's signal correlation. Signal correlation was defined as the Pearson
1336 correlation of the pair's disparity tuning curve values:

1337

(9)

$$r_{signal} = \frac{E[N_j^d N_k^d] - E[N_j^d]E[N_k^d]}{\sigma_{N_j^d}\sigma_{N_k^d}}$$

where N_j^d and N_k^d were the average spike counts of units j and k in response to a disparity d, E was the expected value of the average spike counts across disparity values, and σ was the standard deviation of the average spike counts across disparity values.

The second metric was the degree to which the residual variances in the tuning curves were shared by the two units, the pair's noise correlation. If the stimulus value is fixed over a set of trials, then noise correlation can be quantified as spike-count correlation (r_{SC}) using:

$$r_{SC} = \frac{E[n_j n_k] - E[n_j]E[n_k]}{\sigma_{n_j}\sigma_{n_k}} \tag{10}$$

where n_j and n_k were the spike counts on a single trial from units j and k, E was the expected value across trials, and σ was the standard deviation across trials. In order to pool across trials with different disparity values we first z-score normalized the spike counts of each unit for each sub-set of trials, grouped by disparity; r_{SC} was then computed as:

$$r_{SC} = E[z_j z_k] \tag{11}$$

where z_j and z_k were the z-scored firing rates on a single trial for units j and k.

A limitation of r_{SC} is that it only measures noise correlation at the time scale of the spike-counting window, which is typically on the order of seconds. However, noise correlations may occur at a range of time scales, from near synchrony at the scale of milliseconds to slower modulations over seconds. Noise correlation can be estimated at different time scales in the same analysis window (Spontaneous, Initial, or Steady state) using r_{CCG}, which is the normalized integral of the cross-correlation between paired spike-trains. By integrating from 0 to $\pm\Delta t$ lags under the cross-correlation, r_{CCG} measures rapid noise correlations when Δt is small and includes slower noise correlations when Δt is large. r_{CCG} converges upon r_{SC} when Δt approaches the width of the analysis window. To compute r_{CCG}, the spike train of unit k on trial m was first binned into a series of ms-wide bins; the spike train become a binary series of zeros and ones such that:

$$x_k^m(t) = \begin{cases} 1, \text{if unit } k \text{ fired an action potential during ms } t \text{ on trial } m \\ 0, \text{otherwise} \end{cases} \tag{12}$$

If there were M trials and the analysis window was T ms long, then $1 \leq m \leq M$ and $1 \leq t \leq T$. The post-stimulus time histogram (PSTH) of unit k during time t was:

$$P_k(t) = \frac{1}{M}\sum_{m=1}^{M} x_k^m(t) \tag{13}$$

The spike-train correlation function at τ ms of lag is:

1379

$$C_{jk}(\tau) = \frac{1}{M} \sum_{m=1}^{M} \sum_{t=1}^{T} x_j^m(t) x_k^m(t+\tau) \tag{14}$$

1381

1382 If $j = k$ then $C_{jk}(\tau)$ is the auto-correlation, otherwise if $j \neq k$ then it is the cross-
1383 correlation between units j and k. Likewise, the auto- or cross-correlation of the PSTH was:

1384

$$S_{jk}(\tau) = \sum_{t=1}^{T} P_j(t) P_k(t+\tau) \tag{15}$$

1386

1387 In fact, $S_{jk}(\tau)$ serves as a shift-predictor for $C_{jk}(\tau)$. Thus, the integral of the shift-corrected
1388 spike-train auto- or cross-correlation was:

1389

$$A_{jk}(\theta) = \sum_{\tau=-\theta}^{+\theta} \left[C_{jk}(\tau) - S_{jk}(\tau) \right] \tag{16}$$

1391

1392 where the area $A_{jk}(\theta)$ was summed symmetrically from $-\theta$ to $+\theta$ ms of lag.

1393

1394 Following this, r_{CCG} was defined as:

1395

$$r_{CCG}(\tau) = \frac{A_{jk}(\tau)}{\sqrt{A_{jj}(\tau) A_{kk}(\tau)}} \tag{17}$$

1397

1398 Using (17), equation (10) for r_{SC} can be recast as:

1399

$$r_{SC} = \frac{A_{jk}(T)}{\sqrt{A_{jj}(T) A_{kk}(T)}} \tag{18}$$

1401

1402 where $1 \leq \tau \leq T$. When τ is small, then noise correlations on a short time scale are
1403 measured; whereas, when τ is large, slower noise correlations are estimated. We found r_{CCG}
1404 separately for each disparity value. The mean r_{CCG} across disparities was found by taking the
1405 weighted average, weighing by the number of trials per disparity condition.

1406

1407 A third metric of paired activity is the normalized spike-train cross-correlogram (CCG), which
1408 shows the relative timing of correlated spikes between the pair of units, defined here as:

1409

$$CCG(\tau) = \frac{C_{jk}(\tau)}{\Theta(\tau)\sqrt{\lambda_j \lambda_k}} \tag{19}$$

1411

$$\Theta(\tau) = T - |\tau| \tag{20}$$

1413

1414 where λ_k was the average firing rate (spikes per second) of unit k, and $-T \leq \tau \leq +T$. CCG
1415 was also found separately for each disparity value, and the mean across disparities was

1416 weighted by the number of trials. For presentation purposes, *CCGs* were convolved with the
1417 symmetrical 5ms kernel [0.05, 0.25, 0.40, 0.25, 0.05] (Kohn & Smith, 2005).
1418
1419 *First derivatives of the tuning curves*
1420
1421 In order to estimate the first derivative of the disparity tuning curves, we found the best-
1422 fitting, one-dimensional Gabor function for the tuning curves of each unit, in both the Initial
1423 and Steady state analysis windows (Prince et al., 2002; Tanabe, Umeda, & Fujita, 2004). The
1424 Gabor was expressed as:
1425

$$G(d) = c_0 + c_1 e^{-\left(\frac{(d-c_2)^2}{2c_3^2}\right)} cos(2\pi c_4 (d - c_2) + c_5) \tag{21}$$

1427
where $G(d)$ was the value of the Gabor function at a disparity value of d, and c_i were the
parameters that were optimized for each tuning curve. The baseline offset (c_0) was
constrained between zero and the maximum observed response on any trial. The amplitude
of the Gaussian envelope (c_1) was constrained between zero and twice the difference
between the maximum observed response on any trial and the minimum. The horizontal
offset of the Gaussian envelope (c_2) was constrained between the minimum and maximum
disparity values that were tested. The width of the Gaussian envelope (c_3) was constrained
between 0.1 and the total range of tested disparity values. The disparity frequency of the
cosine (c_4) was estimated from the power spectrum of the tuning curve by first linearly
interpolating (function *interp1*, MATLAB) the z-scored tuning curve from the minimum to
maximum tested disparity, at 0.001° steps. The discrete Fourier transform at frequency f
($y(f)$; function *fft*, MATLAB) was estimated from the interpolated tuning curve, and the
power spectrum at f was computed using:
1441

$$p(f) = \frac{|y(f)|^2}{D} \tag{22}$$

1443
where D was the number of interpolated points. We found the frequency between 0 and 50
cycles per degree that maximized the power spectrum (f_{max}) and then constrained the
disparity frequency to be $c_4 = f_{max} \pm 0.1 f_{max}$. The phase of the cosine (c_5) was
constrained to be within $\pm\pi$. The parameters were fit using the trust-region-reflective
algorithm to minimize the square of the residual error (function *lsqcurvefit*, MATLAB). After
optimizing the parameters for each tuning curve, we estimated the first derivative of the
tuning curves at each tested disparity with:
1451

$$G' = \frac{G(d + \Delta d) - G(d - \Delta d)}{2\Delta d} \tag{23}$$

1453
1454 using a value of 0.5e−3° for Δd. The Gabors better captured the responses to smaller
1455 disparities near zero, hence the 9 smallest disparities were used to estimate $f'\times f'$.
1456
1457 The results in Figure 6 were verified using a second method to estimate the first derivative
1458 of the tuning curves, in which the raw tuning curves were linearly interpolated at $d \pm \Delta d/2$
1459 and the difference was divided by Δd, for each tested disparity value of d.

1460

1461   *Population response analysis*

1462

In order to estimate the ability of simultaneous neuronal responses to distinguish one
disparity value from another, we calculated the bias-corrected linear Fisher information as
defined by Kanitscheider et al. (2015b); their code formed the basis of our own Matlab
implementation (Kanitscheider et al., 2015a). Linear Fisher information quantifies the ability
of a linear decoder to distinguish two sets of simultaneous population spike counts that
arose in response to two different stimulus values. Hence, this was calculated by:

1469

1470
$$\hat{I}_{bc}(\tau) = \frac{\Delta N^{tr}}{\Delta d} \Sigma(\tau)^{-1} \frac{\Delta N}{\Delta d} \left(\frac{2M_{min} - U - 3}{2M_{min} - 2} \right) - \frac{2U}{M_{min}\Delta d^2} \tag{24}$$

1471

1472
$$\Delta d = d_a - d_b \tag{25}$$

1473

1474
$$\Delta N = N^{d_a} - N^{d_b} \tag{27}$$

1475

1476
$$\Sigma(\tau) = \left(\Sigma^{d_a}(\tau) + \Sigma^{d_b}(\tau) \right)/2 \tag{26}$$

1477

where d_a and d_b were two of the tested disparity values such that $d_a \neq d_b$, N^d was a
column vector of the average spike counts for U units in response to disparity value d, $\frac{\Delta N}{\Delta d}$
was also a column vector in which each element of the column vector ΔN was divided by
Δd, tr was the transposition operator that turned $\frac{\Delta N}{\Delta d}$ into a row vector, $\Sigma^d(\tau)$ was the
$U \times U$ covariance matrix for disparity value d at a timescale of τ ms, X^{-1} was the inverse of
matrix X (Moore-Penrose psudoinverse, function *pinv*, Matlab), and $\hat{I}_{bc}(\tau)$, bias-corrected
Fisher information, was a scalar value with a unit of degrees⁻². We randomly discarded trials
so that each disparity value had the same number of trials as that which had the minimum
(M_{min}), as assumed in the derivation by Kanitscheider et al. (2015b). In practice, few trials
were discarded, with the remainder being well above the minimum required for the
covariance matrix to be invertible, which is $(U + 2)/2$. For units j and k, we defined
elements of $\Sigma^d(\tau)$ to be:

1490

1491
$$\Sigma^d_{j,k}(\tau) = r^d_{CCG}(\tau)\sigma^d_j\sigma^d_k \tag{28}$$

1492

1493   where $r^d_{CCG}(\tau)$, σ^d_j and σ^d_k were respectively the r_{CCG} and spike-count standard deviations of
1494   units j and k on trials with disparity d. Thus, we could estimate the effect of noise
1495   correlations at different time scales on the linear Fisher information of the population
1496   response. It was possible to estimate the variance of the linear Fisher information for
1497   different time scales of noise correlation as:

1498

1499
$$\mathrm{Var}\hat{I}_{bc}(\tau) = \frac{2\hat{I}_{bc}(\tau)^2}{2M_{min} - U - 5}\left(1 + \frac{4(2M_{min} - 3)}{M_{min}\hat{I}_{bc}(\tau)\Delta d^2} + \frac{4U(2M_{min} - 3)}{\left(M_{min}\hat{I}_{bc}(\tau)\Delta d^2\right)^2} \right) \tag{29}$$

1500

1501   A theoretical measure of linear Fisher information is obtained when all units in the
1502   population are decorrelated with each other such that the variance of their spike counts is

statistically independent. This is often achieved numerically by shuffling the data. However, it is possible to estimate this value analytically without shuffling – using only the diagonal of the covariance matrix (called Ibc,shuffle in Kanitscheider et al., 2015b):

$$\hat{I}_{bc,decorr} = \hat{I}_{bc,decorr}(\tau) = \frac{2U}{M_{min}\Delta d^2} \sum_{i=1}^{U} \frac{(\Delta N_i / \Delta d)^2}{\Sigma_{i,i}(\tau)} - \frac{M_{min} - 2}{M_{min} - 1} \qquad (30)$$

Eq. 30 emphasizes the fact that with this calculation the decorrelated linear Fisher information is the same for all time scales of noise correlation, because the auto-correlated value of r_{CCG} is always 1 (see eq. 17 when $j = k$) and σ_j^d does not vary with τ (eqs. 27 & 28).

Any difference between the empirical and decorrelated information could signal the presence of information-limiting noise correlations in the population. In order to make a direct comparison between all combinations of disparity values, and for different compositions of neuronal population, we computed the normalized difference *i.e.* information loss as:

$$\Delta \hat{I}_{bc}(\tau) = \frac{\hat{I}_{bc,decorr}(\tau) - \hat{I}_{bc}(\tau)}{\sqrt{Var\hat{I}_{bc}(\tau)}} \qquad (31)$$

in a unit of standard deviations of $\hat{I}_{bc}(\tau)$ (see Fig.S8). Outliers were discarded when the estimated variance resulted in a normalized difference of 40 or greater. With 13 tested disparities, there were 78 unique pairs of disparity values and therefore 78 values of $\hat{I}_{bc}(\tau)$, $Var\hat{I}_{bc}(\tau)$, $\hat{I}_{bc,decorr}$, and $\Delta \hat{I}_{bc}(\tau)$ were each computed for all neuronal pool sizes, compositions, and correlation time scales. Neuronal pools comprised of V1 units only, V4 units only, or a mixture of V1 and V4 units. Over 10 sessions, this yielded 780 information values in each case.

To see if linear Fisher information saturated with increasing population size, we first ordered a given set of units by the DMI of their Initial responses, from largest to smallest. Units were added to the neuronal pool one at a time in this order, and the Fisher information was computed at each step; that is, the most disparity-selective unit was added first, and the least selective unit was added last. The result was an information curve that was a function of pool size. Initial DMI was always used to order the units because their Initial responses were the most selective for disparity (see Figs.2&S2). This ordering imposed a saturating shape on even the decorrelated information curves. However, we could then examine whether the saturation worsened as a result of noise correlations. Information curves were found for the Initial and Steady state responses at time scales of 20 and 150ms (Initial) or 20 and 800ms (Steady state) and for decorrelated units (both). In order to average across experimental sessions and combinations of disparity values, it was necessary to normalize the information values in a way that would preserve the relative shape of information curves at different time scales. To do this, the curves were first grouped by experiment and disparity values. Normalization was then achieved within a group by:

$$\qquad (32)$$

1546
$$I_{norm}(\tau, s_{pop}) = \frac{I_{raw}(\tau, s_{pop}) - \beta_{grp}}{\alpha_{grp}}$$

1547

where $I_{raw}(\tau, s_{pop})$ was the value of $\hat{I}_{bc}(\tau)$ or $\hat{I}_{bc,decorr}$ at population size s_{pop} units. β_{grp}
was the minimum value of $I_{raw}(\tau, 1)$ across values of τ. α_{grp} was then taken to be the
maximum value of $I_{raw}(\tau, s_{pop}) - \beta_{grp}$ across values of τ and s_{pop}. Results obtained using
the unit order above were confirmed in a control analysis that randomly permuted the unit
order before computing the information curves; the control was repeated 100 times for a
total N = 78,000.

1554

1555 *Testing effect of choice on noise correlation and $f' \times f'$ relationship*

1556

We wanted to test if the relationship between noise correlation (r_{CCG}) and the product of
tuning curve derivatives ($f'{\times}f'$) was dependent upon the correctness of the choice. Values
were computed using a 500ms wide analysis window that was tiled across the Present RDS
phase (Fig.1Aiii) at 250ms steps, such that windows at adjacent positions were overlapping
each other by 50%. At each window position, a unique pair of r_{CCG} and $f'{\times}f'$ values were
computed for each pair of neurons, each baseline disparity, and each combination of
popout target location (IN or OUT of RF) and choice location (IN or OUT of RF). To visualise
the dependency on choice, the Spearman correlation of r_{CCG} and $f'{\times}f'$ was computed across
neuron pairs and baseline disparities at each window position, for each combination of
target and choice location. The results are shown in Figure 9. Each Spearman correlation
value was classified as significantly different from 0 at a 5% level (filled vs empty circles,
two-tailed test).

1569

To quantify the statistical significance of our results, r_{CCG} and $f'{\times}f'$ values from each subject
were grouped by window position, popout target location (IN or OUT), and type of neuron
pair (within V1, within V4, between V1 and V4). For the g^{th} grouping (one unique
combination of subject, window position, target location, and pair type), the r_{CCG} values
were ranked (adjusted for ties; function *tiedrank*, Matlab), and so were the $f'{\times}f'$ values. This
transformed the two sets of values from group g into uniform distributions. Within group g,
the i^{th} combination of neuron pair and baseline disparity produced $q^i_{r_{CCG}}$ and $q^i_{f'\times f'}$, the
ranked r_{CCG} and $f'{\times}f'$ values, respectively. Each ranked value was then normalised:

1578

1579
$$nq^i_{r_{CCG}} = \frac{q^i_{r_{CCG}}}{q^{max}_{r_{CCG}}} \tag{34}$$

1580
$$nq^i_{f'\times f'} = \frac{q^i_{f'\times f'}}{q^{max}_{f'\times f'}} \tag{33}$$

1581

1582 Where $q^{max}_{r_{CCG}}$ and $q^{max}_{f'\times f'}$ were the maximum ranks of each value, observed in group g. This
1583 resulted in sets of values that were uniformly distributed between 0 and 1. The log ratio of
1584 normalised ranks was taken:

1585

1586
$$\log_{10} \frac{nq^i_{r_{CCG}}}{nq^i_{f'\times f'}} \tag{35}$$

1587

Critically, the log ratio preserved the relative position between correct (Target IN & Choice
IN, or Target OUT & Choice OUT) and incorrect (Target OUT & Choice IN, or Target IN &
Choice OUT) data; note, incorrect Target OUT & Choice OUT trials were not used. The
output of equation 35 was used in an all-ways ANOVA (one per subject) testing the effect of
window position, type of neuron pair, and correctness of choice that was further examined
using Tukey-Kramer's multiple-comparisons test.

1594

1595 *Detect probability using optimal projection of population responses*

1596

To test the possible effect of differential correlations on perceptual choice, we needed a
way to obtain a trial-by-trial estimate of differential correlation. This was done following the
technique of (Nogueira et al., 2020). Let n be a column vector of firing rates measured from
the simultaneous responses of N neurons on a single trial. The expected value of the firing
rates in response to baseline disparity d was $\mu_d = E[n|d]$. For any $d \in B$, where B is the set
of baseline disparity values, μ_d was estimated using the sample mean calculated from an
analysis window −200 to 0ms before the start of the popout event. Following the start of
the popout event, the binocular disparity of the target RDS was $d_p = d_b + s$, where $d_b \in B$
and s was one of several disparity steps (0°, ±0.05°, ±0.1°). For any $d_p \notin B$, μ_{d_p} was
estimated from the measured disparity tuning curves using linear interpolation. The signal
vector was defined as $\Delta f_{d_b d_p} = \mu_{d_p} - \mu_{d_b}$, describing the expected change in firing rate
between responses to the baseline and popout disparities. Equation 28 was used to define
the $N \times N$ covariance matrix $\Sigma_d(\tau)$ for each $d \in B$, using the same −200 to 0ms window.
Note that while μ_d and $\Sigma_d(\tau)$ were always measured from the −200 to 0ms window, the
population responses n were measured from a set of different analysis window positions
using different τ values, so that the analysis could examine the potential interaction of
differential correlation and choice at different points in time, for different temporal scales
of noise correlation.

1615

In order to discriminate the baseline population response from the popout response, the
optimal weighted sum of firing rates will use weights $w_{opt} \propto \Sigma^{-1} \Delta f$, similar to the solution
for linear discriminant analysis. Hence, the weights used on a given trial with baseline

disparity d_b and popout disparity d_p were defined as $w_{d_b d_p}(\tau) = \left(\Sigma_{d_b}^{-1}(\tau) \Delta f_{d_b d_p} \right)^{tr}$. Thus,
the optimal set of weights were tailored to discriminate the baseline and popout disparity
on each trial. The raw projection, $y_{raw} = w_{d_b d_p} n$ was normalised:

1622

$$\mathrm{y}(\tau) = \frac{w_{d_b d_p}(\tau)\left(n - \mu_{d_b}\right)}{w_{d_b d_p}(\tau)\Delta f_{d_b d_p}} \tag{36}$$

1624

such that $E[y|d_b] = 0$ and $E[y|d_p] = 1$; that is, the expected value of the projection was 0
for responses to the baseline disparity, and 1 for responses to the popout disparity. Hence,
the projected responses were in units of relative degrees, in comparison to the disparity
step s. Because the optimal linear weights were used, the linear combination of population
responses could mitigate any non-differential noise correlations. Thus, any remaining
variance in y was possibly due to information-limiting, differential correlations.

1631

The optimal projection of population responses, y, was measured on each trial for each
composition of neural population (V1 units only, V4 units only, all V1 and V4 units). Hence,
we could use y to predict the behavioural choice using signal detection theory to compute a
measure analogous to Choice Probability (Britten et al., 1996). After grouping trials by target
location (IN or OUT of the RF), we labelled choices IN to the RF as 'true positives' and
choices OUT of the RF (correct choice when target OUT) as 'false positives' before
computing the area under the receiver operating characteristic curve (AROC, see Equation
40 below). We pooled y across experimental days, per subject, because this was a
normalised value and because the same cortical sites were recorded each day thanks to the
chronic Utah arrays; such normalisation and pooling was required to overcome the plurality
of stimulus × choice conditions. Thus, we pooled 1168 choice IN (197 choice OUT) trials for
M135 and 819 (285) for M138 when the popout target RDS was IN the RF, and 982 (2427)
for M135 and 791 (1918) for M138 when the target was OUT; this was an ample number of
trials to effectively compute AROC (Britten et al., 1996). The significance of AROC values was
estimated by computing bootstrap 95% confidence intervals, obtained from 2e4 bootstrap
samples using the percentile method. We called this value 'Detect Probability' ($DP(\tau)$ using
$y(\tau)$) because the animals had to detect the location of any disparity change, without
reporting the direction of the change. However, this DP is not strictly the same as the
original DP of Cook and Maunsell (Cook & Maunsell, 2002), whose task used temporal
rather than spatial uncertainty.

1652

Here, DP is the probability that an ideal observer could predict the subject's choice (IN or
OUT of RF) using the optimal projection of the population response, y. If DP > 0.5 then y
tended to be more positive (*i.e.* closer to popout disparity) prior to choices IN to the RF. DP
< 0.5 shows that y was more negative (*i.e.* further from popout disparity) prior to choices IN.
DP = 0.5 is the chance level, showing no discernible differences in the distributions of y for
choice IN and choice OUT.

1659

1660 *A model V4 output neuron can attenuate differential noise*

1661

We modelled a theoretical V4 output neuron in order to test whether a simple mechanism
could attenuate differential noise that was present upstream in V1 and V4. Thus, the model
unit had only two afferent inputs, a V1 unit and a V4 unit (see Fig.11A). There were two
chief parameters of the model, the signal correlation of the V1/V4 input pair and the width
of a window used to integrate their responses. V4 output responses were predicted by
subtracting a weighted version of the V1 input from the V4 input. V4 output neurons were
modelled for all recorded V1/V4 pairs as afferent inputs, and for window widths of 20, 40,
80, and 160ms for Initial responses or 100, 200, 400, and 800ms for Steady state responses.
The V1 input weight (w in Fig.11A) should reflect the magnitude of noise shared by the
afferent pair that was not explained by the random sampling of disparity values from trial to
trial. To measure it, we used a simple generalized linear model (GLM) to predict the spike
count of the V4 input from its disparity tuning curve and the spiking of the V1 input, using
the ln link function:

1675

1676
$$\ln n_{V4}^m = c_0^{glm} + c_1^{glm} G_{V4}(d_m) + c_2^{glm} n_{V1}^m \qquad (37)$$

1677

n_{V4}^m was the spike count of the V4 input unit on trial m, G_{V4} was the Gabor function that
best fit the V4 input's tuning curve, G_{V4} was evaluated at the disparity value d_m used on
trial m, and n_{V1}^m was the spike count of the V1 input unit on trial m. The GLM was used to fit
(function *glmfit*, MATLAB) three parameters: the baseline spike count (c_0^{glm}), the amount of
disparity signal (c_1^{glm}), and the amount of shared variance between the afferent pair (c_2^{glm}).
The Gabor term was important, as it helped to push c_2^{glm} towards accounting for shared
residual variance rather than shared signal.
1685
1686 Once obtained, we predicted the V4 output responses with:
1687
1688 $$n_{V4,OUT}^m = n_{V4}^m - exp\left(c_2^{glm} n_{V1}^m\right) + \beta \tag{38}$$
1689
where $n_{V4,OUT}^m$ was the V4 output spike count on trial m. Thus, the V4 output was the V4
input minus the weighted V1 input. β was a constant term that restored the baseline value
of the modelled V4 output disparity tuning curve to match that of the V4 input. Any
negative values were capped at zero. This produced V4 output spike counts with a similar
dynamic range to the V4 input. The exponential function was used to transform the scaled
V1 response because the logarithmic link function was used in the GLM.
1696
To avoid over-fitting, we applied equations (37) and (38) to different halves of the trials in a
two-fold, cross-validation procedure. Trials were divided into even and odd-numbered
groups. Thus, the training and test sets had a similar size and composition, balancing the
need to train the GLM against the need to accurately compute AROC (see below). One
group of trials was used to train the model with Eq. (37) and then the other group was used
to evaluate it with Eq. (38). The process was repeated a second time after reversing the
roles of each group. AROC was used to evaluate the disparity selectivity of the V4 output
responses for each test set, yielding two estimates per condition that were averaged
together.
1706
The training, testing and evaluation procedure was applied to the empirical data, in which
the inter-cortical noise correlations were intact (Fig.11C thick solid, 'Not shuffled').
However, it was necessary to determine the degree to which any change in disparity
selectivity of the V4 output responses was due to the combined disparity information
signalled by the afferent pair. Therefore, the procedure was applied a second time to data in
which the V1 responses were shuffled between trials. Shuffling was done by swapping
responses between trials with the same disparity, so that the disparity selectivity of both
the V1 and V4 input units was preserved, but the noise correlations were destroyed (thin
solid, 'V1 tuning kept'). A third evaluation was done using shuffled data in which responses
were shuffled across all trials, regardless of disparity, so that the noise correlations and V1
disparity selectivity were abolished (dashed, 'All trials shuffled').
1718
We evaluated summation of the V1/V4 inputs with integration windows of different widths
(Figure 11D). This was accomplished by sub-dividing each analysis window (Initial or Steady
state) into a set of smaller, contiguous windows that were each Δt ms wide. Spike counts
were obtained and equations (37) and (38) were applied separately for each sub-division;
that is, an independent set of GLM parameters was trained and tested for each sub-division

of the analysis window. The V4 Gabor function was scaled to produce the spike count
expected in a window of size Δt ms. Before computing AROC, the V4 output spike counts
were summed back together across sub-divisions:

1727

$$1728 \qquad n_{V4,OUT}^{m} = \sum_{\omega=1}^{W} n_{V4,OUT}^{m}(\omega) \qquad\qquad (39)$$

1729

1730 where $n_{V4,OUT}^{m}(\omega)$ was the V4 output spike count obtained in the ω^{th} sub-division of the
1731 analysis window, out of W sub-divisions with a width of Δt ms.

1732

AROC was used to evaluate the model instead of DMI. In computing DMI non-
parametrically, it is possible that the true information was underestimated as a result of
discretizing the data through binning, analogous to the pixelation of a low-resolution
photograph. Therefore, we used a continuous method based on signal detection theory
(Green & Swets, 1966) that has previously served to quantify the sensitivity of single
neurons to different stimulus values (Newsome et al., 1989). We used a similar approach to
estimate disparity sensitivity, in which a receiver operating characteristic curve (ROC) is built
by labelling one set of spike counts as true positives and another as false positives. If the
true-positive spike counts were from trials with one disparity value and the false positives
were from trials with another, then the area under the ROC (AROC) becomes the probability
that an ideal observer could correctly classify the true-positive spike counts. This leads
directly to a measure of how well the neuron can discriminate the two tested disparities. A
chance value of AROC = 0.5 means that the two spike-count distributions were
indistinguishable. Values of 0 or 1, however, indicate perfect separation of the distributions,
with intermediate values of AROC indicating different amounts of overlap. We calculated
AROC using trapezoidal integration of the true- and false-positive cumulative distribution
functions (F_T and F_F, respectively):

1750

$$1751 \qquad AROC = \frac{1}{2} \sum_{i=1}^{V} [F_F(x_i) - F_F(x_{i-1})][F_T(x_i) + F_T(x_{i-1})] \qquad\qquad (40)$$

1752

1753 where $x_i \in \{x_1, x_2, x_3, \dots x_V\}$ is the ordered (ascending) set of V unique spike counts from
1754 either the true- or false-positive distribution, and $F_T(x_0) = F_F(x_0) = 0$. The absolute
1755 difference of AROC from chance is a measure of separation between two spike count
1756 distributions regardless of which was labelled as the true- or false-positives:

1757

$$1758 \qquad |AROC| = |AROC - 0.5| \qquad\qquad (41)$$

1759

1760 $|AROC|$ was computed for each of the 78 unique pairs of disparity values. With 4598 V1/V4
1761 afferent pairs, we obtained a total of 358,644 AROC predicted values for the model V4
1762 output neurons. The difference in V4 output and V4 input disparity selectivity is shown in
1763 Figures 11C.

1764

1765 *Bootstrap confidence intervals*

1766

1767   Unless otherwise stated, 95% bootstrap confidence intervals were found using the bias
1768   corrected and accelerated percentile method (function *bootci*, MATLAB) with at least 1000
1769   bootstrap samples.
1770

1772

Figure 1. *Task and performance.* A) (i) Trials started with flashing fixation point (FP, white
circle). (ii) Once fixated (dashed diamond), FP flash stops. (iii) 500ms later, four random dot
stereograms (RDS, white dots) with identical properties appeared. (iv) After fixed duration
(1s M135, 2s M138), central dots of one RDS change disparity (grey/black dots) and FP
disappears. Animals made saccade onto the oddball target RDS (arrow) for reward. B)
Percent correct trials (chance=25%, dashed line) and reaction time (RT) over popout
disparity. Using trials with $0.117 \leq RT \leq 0.344s$ for M135 (N = 6670 of 8279), or $0.143 \leq RT \leq$
$0.463s$ for M138 (N = 5478 of 6073). RT shown separately for correct (thick) and incorrect
choices *i.e.* failed trials (thin). Error bars 95% binomial confidence intervals (CI) or standard
error of the mean (SEM). C) Percent correct across RT using 50ms moving window; with 95%
bootstrap CI (2000 samples, 2.5th & 97.5th percentiles). Data grouped by absolute popout
disparity (see legend).

1785

Figure 2. *V1 and V4 units were selective for binocular disparity.* A) Receptive field (RF)
location and size for each V1 (gold) and V4 (blue) neurons; circle radius = 25% standard
deviation of 2D Gaussians fit to RF location maps. RDS centre (short dash) and surround
(long dash) shown. B) Example disparity tuning curves from V1 (circles, yellow) and V4
(crosses, blue); symbols show average firing rates; lines show best-fitting Gabors: SEM
below symbol size. C) Disparity mutual information (DMI) for V1 (gold, N = 195) and V4
(blue, N = 213) units compared with average firing rate to preferred disparities. D) Average
firing rate (mean line, SEM ribbon) across all V1 (gold) and V4 (blue) units, over trial
sequence: analysis windows cover spontaneous responses (left arrow), Initial phasic
response (solid bar and lines), later Steady state responses (right arrow); dashed line marks
the onset of RDS. E) As in C but for DMI.

1797

Figure 3. *Alignments of noise correlations and disparity tuning.* A) Disparity tuning curves of
pair V4/1 (black) & V4/2 (grey; M138.336.94/147), signal correlation (r_{signal} = +0.851);
crosses and SEM show average firing, lines show best-fitting Gabors (V4/1 r^2 = 89%, V4/2 r^2
= 93%). Vertical dashed line at +0.1° shows trials (N = 108) examined in panels B&C. B)
Raster plots and average firing rates for V4/1 (top) and V4/2 (bottom); RDS onset at t=0; for
108 trials at baseline disparity +0.1°; shading shows time window for firing rates in C. C)
Dots show joint firing rates; slight jitter applied for visualization; noise correlation, r_{SC} =
0.241, *p* = 0.012. Tangent line (grey) shows joint tuning curves at +0.1°; arrow shows first
PCA component (×40, for visibility) of joint responses. D – F) the same as A – C but for an
example V1/V4 pair (M138.336.57/87; V1 black in D, r^2 = 98%; V4 grey in D, r^2 = 94%) with
negative signal correlation (r_{signal} = −0.982). Trials with baseline disparity of 0° (D, black
dashed, N = 116) yield significant positive noise correlation (F, r_{SC} = 0.203, *p* = 0.029).

1810

Figure 4. *r_{CCG} measures noise correlation at different time scales.* A) Average r_{CCG} as function
of temporal scale τ for all V1/V1 pairs' (N = 2419) spontaneous (dashed black), Initial (teal),
and Steady state (magenta) responses (see Fig.2D); ribbons show SEM. B) As A, but has
average V1/V1 cross-correlograms (CCG). C) Scatter plot of Steady state r_{CCG} at τ =100ms
versus spike-count correlation (r_{SC}). Marginal distributions show each metric separately. D –
F) the same as A – C but for V4/V4 pairs (N = 2226). G – I) the same as A – C but for V1/V4
pairs (N = 4598).

1818

Figure 5. *Noise correlations varied with disparity tuning, time scale, and time.* A) Noise correlation (r_{CCG}) for V1/V1 as function of signal correlation (r_{signal}), grouped by time scale (τ) of r_{CCG} (20ms, dashed; 150ms, solid) and analysis window (Initial, black; Steady state, magenta); each bin has 10% of the data; error bars are SEM. B) Average V1/V1 r_{CCG} over time scale τ, grouped by signal correlation (positive r_{signal}, dashed; negative r_{signal}, solid) and analysis window, as A; ribbon shows SEM. C&D) same as A&B but for V4/V4 pairs. E&F) same as A&B but for V1/V4 pairs.

Figure 6. *Empirical evidence of differential noise correlations.* A) $r_{CCG}(\tau)$ from V1 at τ = 41ms plotted over product of first derivatives of disparity tuning curves ($f'{\times}f'$), for Initial response phase, N = 21,771 cases (2419 V1/V1 pairs at 9 disparities −0.1° to +0.1°). Best-fitting, least-squares regression line (orange) $r_{CCG}(\tau) = k + m(f' \times f')$ had positive slope (m): k = 3.452e−2; m = 1.244e−7. B) As A but for Steady state: τ = 95ms; k = 1.594e−2; m = 2.812e−7. C) Spearman correlation coefficient of r_{CCG} and $f'{\times}f'$ for Initial (black) and Steady state (magenta) responses, at all $\tau \leq$ 400ms. Filled circles show p < 0.05 (two-tailed). Arrows show τ used in A (black) and B (magenta). D − F) Same as A − C but for V4/V4 pairs and using τ = 21ms for D and τ = 32ms for E, with N = 20,034. Positive regression slopes in D: k = 3.020e−2; m = 3.039e−6, and E: k = 3.330e−2; m = 3.891e−6. G − I) Same as A − C but for V1/V4 pairs and using τ = 88ms for G and τ = 109ms for H, with N = 41,382. Negative regression slope in G: k = 7.782e−3; m = −4.711e−7, and H: k = 6.354e−3; m = −2.804e−7.

Figure 7. *Differential noise correlations are attenuated at long time scales.* A) Average ±SEM normalized bias-corrected linear Fisher information, over neuronal pool size. Information shown for rapid (τ =20ms, dotted), slow (τ =150ms, solid), and no (black) noise correlations, N = 780 (78 disparity pairs × 10 experiments). Units added to pools by decreasing order of Initial DMI.. B) Average Fisher information across r_{CCG} τ. Information is empirical (thick solid) or decorrelated (dashed). 95% bootstrap CI. C) Decorrelated vs. empirical information at τ = 150ms. Max. pool sizes in B & C.

Figure 8. *Differential correlations and receptive field separation.* For each type of neuron pair, pairs were ordered by RF separation then grouped; at least 100 pairs to each group (N bins = N pairs / 100, rounded down). Spearman correlation of r_{CCG}(150ms) versus $f'{\times}f'$ for the pairs in each group, as a function of average log_{10} RF separation. Marginal histograms show distribution of Spearman correlation, counting groups.

Figure 9. *Effect of choice on direct evidence of differential correlation.* Spearman correlation of $r_{CCG}(\tau)$ versus $f'{\times}f'$ (see Fig.6) in 500ms window (grey) at 250ms steps (circles, centred in each window) prior to popout event (Fig.1Aiv). For V1/V1 (left, τ = 30ms), V4/V4 (middle, τ = 30ms), and V1/V4 (right, τ = 110ms) pairs. RDS onset (Fig.Aiii) at −1 for M135, −2 for M138. Circle fill shows significance (two-tailed, p ≤ 0.05). Coloured circles show choices IN to RF, black show choices OUT of RF. A) Popout target IN the RF, choices IN are correct. B) Popout appears OUT of RF, choices IN are incorrect; incorrect choices out of RF not shown.

Figure 10. *Differential correlation predicts choice.* N-dimensional firing rates of simultaneous population responses were linearly projected onto the axis that optimally discriminated baseline from popout responses. A) Example projected responses for combined V1 and V4

populations. Expected value of 0 for response to baseline disparity (dashed), and +1 for popout (solid). Target RDS appeared IN (top panels) or OUT (bottom) of the RF. Responses prior to (−100 to 0ms, grey) or after start (+35 to +135ms, teal) of popout event. B) Example ROC curves (bottom, green) using pre-popout projections from A (top, grey). Black line: chance association. Crosses show probability values equal to expected baseline (open) or popout (filled) response. Top shows distributions grouped by choice IN (empty) or OUT (filled) of RF, black vertical lines as in A, green vertical lines show averages. C) Detect Probability for populations with V1 units only (gold), V4 units only (blue), or mixed V1 and V4 units (green), relative to chance (0.5, black horizontal) with target RDS IN (top) or OUT (bottom) of the RF. Population responses taken from analysis windows at different positions (left edge = −400, −200, or −100ms before popout) and width (200 or 100ms). Projection weights optimised for noise correlations at r_{CCG} time scales τ = 20 or 150ms. * 95% CI different from chance. Arrows show DP for curves in B. All Error bars are 95% bootstrap CI (2e4 boot samples, percentile method). Neuron pairs $RF_{separation} \leq 7.99°$.

Figure 11. *A model V4 output neuron attenuates differential noise.* A) Diagram of model V4 output neuron. Diamonds, afferent neurons. Rasters, afferent spikes in Δt ms window. Weighted V1 count ($w\Sigma_{V1,IN}$) subtracted from V4 count ($\Sigma_{V4,IN}$) gets V4 output count ($\Sigma_{V4,OUT}$). B) Disparity tuning prediction, afferent V1/V4 r_{signal} < 0 and r_{noise} > 0. V4 input variance ($\Sigma_{V4,IN}$, dotted). Model V4 output variance ($\Sigma_{V4,OUT}$, grey). C) Average change in AROC across r_{signal} of afferent V1/V4 pairs, Δt = 160ms. V1 responses shuffled across trials to break V1/V4 correlations but not disparity selectivity (V1 tuning kept, thin), to break correlations and V1 tuning (All trials, dashed), or not shuffled (solid). N = 358,644 (78 baseline disparity pairs × V1/V4 pairs). Each bin has 10% of data. D) Average difference in AROC over Δt, V4 model output AROC with no shuffling minus shuffled AROC with preserved V1 tuning. Afferent V1/V4 r_{signal} < −0.5 (black) and r_{signal} > +0.5 (grey). N = 4598 model V4 output neurons, one per recorded V1/V4 pair. C&D) Initial data on right, Steady state on left. SEM error bars.

Figure S1. *RF locations.* A) RF centers for V4 control data (blue circles) relative to the center of gaze. The lower-right RDS stimulus (dashed) was placed to maximize V4 responses without stimulating the V1 sites. Accompanies Figure 1.

Figure S2. *Disparity tuning in V4 control.* Similarity of DMI and DDI shown for main V1 (gold) and V4 (blue) (A) and control V4 (B). C, D, & E replicate Fig.2C, D, & E for V4 control data (N = 679 units). Accompanies Figure 2.

Figure S3. *Alternative example V1/V4 pairs.* A, B, and C replicate Fig.3D, E, & F for three other example pairs of V1 and V4 units; one pair per panel column. Accompanies Figure 3.

Figure S4. r_{CCG} *for V4 control data.* A, B, & C replicate Fig.4D, E, & F for the V4 control data (N = 2637 pairs). Accompanies Figure 4.

Figure S5. *Noise correlations vs signal correlation and time scale for V4 control data.* A & B replicates Fig.5C&D for the V4 control data. Accompanies Figure 5.

Figure S6. *Empirical evidence of differential noise correlations in V4 control data.* A, B, & C replicates Fig.6D, E, & F for the V4 control data. Accompanies Figure 6.

1912

1913 **Figure S7.** *Differential correlation attenuation controls.* A, B, & C replicate Fig.7A, B, & C
1914 (middle) for V4 control data. D, E, & F replicate Fig.7A, B, & C using Steady state analysis
1915 window. G) Information of mixed V1/V4 pool (size 12) vs. summed information across V1-
1916 only and V4-only pools (size 6) for rapid (left) and slow (right) correlations. H) Average
1917 paired differences of data in G (left) over τ. Right, same but for Steady state. * significant
1918 paired t-tests for adjacent data (Initial α = 0.05, Steady state α = 0.05/3).

1919

1920 **Figure S8.** *Effect of RF separation on information encoding.* A) Average information loss due
1921 to differential correlation, across noise-correlation temporal scale (τ), for populations of V1
1922 units only (gold), V4 only (blue), or combined V1 & V4 units (green). B) same as A, but for V4
1923 control. C) Information loss in two populations of V1 and V4 units. Same V1 units in both
1924 populations. Sets of V4 units were non-overlapping. Each V4 unit classed by its average RF
1925 separation to all V1 units; V4 RFs were close to V1's (Close-V4, solid green) or distant
1926 (Distant-V4, black dashed). Inset panels show spline-interpolated average RF maps for each
1927 V4 group. The criterion (7.99°) for classifying 'close' and 'distant' made the number of close
1928 V1/V4 pairs in M135 the same as the number of distant V1/V4 pairs in M138. Thus, the
1929 smaller group from each animal had comparable statistical power. All) 95% bootstrap CI.

1930

1931   <u>REFERENCES</u>

1932

1933

Abbott, L. F., & Dayan, P. (1999). The effect of correlated variability on the accuracy of a
population code. *Neural Computation, 11*(1), 91-101.
Averbeck, B. B., Latham, P. E., & Pouget, A. (2006). Neural correlations, population coding
and computation. *Nat Rev Neurosci, 7*(5), 358-366. doi:nrn1888 [pii]
10.1038/nrn1888
Bair, W., Zohary, E., & Newsome, W. T. (2001). Correlated firing in macaque visual area MT:
Time scales and relationship to behavior. *Journal of Neuroscience, 21*(5), 1676-1697.
Barlow, H. B., Kaushal, T. P., Hawken, M., & Parker, A. J. (1987). Human Contrast
Discrimination and the Threshold of Cortical-Neurons. *Journal of the Optical Society*
of America a-Optics Image Science and Vision, 4(12), 2366-2371. Retrieved from <Go
to ISI>://A1987L543100024
Beck, Jeffrey M., Ma, Wei J., Pitkow, X., Latham, Peter E., & Pouget, A. (2012). Not Noisy,
Just Wrong: The Role of Suboptimal Inference in Behavioral Variability. *Neuron,*
74(1), 30-39. doi:https://doi.org/10.1016/j.neuron.2012.03.016
Bondy, A. G., Haefner, R. M., & Cumming, B. G. (2018). Feedback determines the structure
of correlated variability in primary visual cortex. *Nat Neurosci, 21*(4), 598-606.
doi:10.1038/s41593-018-0089-1
Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision, 10*(4), 433-436.
Britten, K. H., Newsome, W. T., Shadlen, M. N., Celebrini, S., & Movshon, J. A. (1996). A
relationship between behavioral choice and the visual responses of neurons in
macaque MT. *Visual Neuroscience, 13*(1), 87-100.
Buffalo, E. A., Fries, P., Landman, R., Buschman, T. J., & Desimone, R. (2011). Laminar
differences in gamma and alpha coherence in the ventral stream. *Proceedings of the*
National Academy of Sciences, 108(27), 11262. doi:10.1073/pnas.1011284108
Carandini, M., & Heeger, D. J. (2012). Normalization as a canonical neural computation. *Nat*
Rev Neurosci, 13(1), 51-62. Retrieved from http://dx.doi.org/10.1038/nrn3136
Churchland, M. M., Yu, B. M., Cunningham, J. P., Sugrue, L. P., Cohen, M. R., Corrado, G. S., .
. . Shenoy, K. V. (2010). Stimulus onset quenches neural variability: a widespread
cortical phenomenon. *Nature Neuroscience, 13*(3), 369-U325. doi:Doi
10.1038/Nn.2501
Cohen, M. R., & Kohn, A. (2011). Measuring and interpreting neuronal correlations. *Nat*
Neurosci, 14(7), 811-819.
doi:http://www.nature.com/neuro/journal/v14/n7/abs/nn.2842.html#supplementar
y-information
Cohen, M. R., & Maunsell, J. H. (2009). Attention improves performance primarily by
reducing interneuronal correlations. *Nat Neurosci, 12*(12), 1594-1600. doi:nn.2439
[pii]
10.1038/nn.2439
Cohen, M. R., & Newsome, W. T. (2008). Context-Dependent Changes in Functional Circuitry
in Visual Area MT. *Neuron, 60*(1), 162-173. Retrieved from <Go to
ISI>://000260237300016
Cohen, M. R., & Newsome, W. T. (2009). Estimates of the Contribution of Single Neurons to
Perception Depend on Timescale and Noise Correlation. *Journal of Neuroscience,*
29(20), 6635-6648. Retrieved from <Go to ISI>://000266252300026

Cook, E. P., & Maunsell, J. H. R. (2002). Dynamics of neuronal responses in macaque MT and VIP during motion detection. *Nature Neuroscience, 5*(10), 985-994. Retrieved from <Go to ISI>://000178242200018

Denfield, G. H., Ecker, A. S., Shinn, T. J., Bethge, M., & Tolias, A. S. (2018). Attentional fluctuations induce shared variability in macaque primary visual cortex. *Nat Commun, 9*(1), 2654. doi:10.1038/s41467-018-05123-6

Dodd, J. V., Krug, K., Cumming, B. G., & Parker, A. J. (2001). Perceptually bistable three-dimensional figures evoke high choice probabilities in cortical area. *Journal of Neuroscience, 21*(13), 4809-4821. Retrieved from <Go to ISI>://000169454600029

http://www.jneurosci.org/content/jneuro/21/13/4809.full.pdf

http://www.jneurosci.org/content/21/13/4809.long

Ecker, A. S., Berens, P., Cotton, R. J., Subramaniyan, M., Denfield, G. H., Cadwell, C. R., . . . Tolias, A. S. (2014). State dependence of noise correlations in macaque primary visual cortex. *Neuron, 82*(1), 235-248. doi:10.1016/j.neuron.2014.02.006

Ecker, A. S., Denfield, G. H., Bethge, M., & Tolias, A. S. (2016). On the Structure of Neuronal Population Activity under Fluctuations in Attentional State. *Journal of Neuroscience, 36*(5), 1775-1789. doi:10.1523/JNEUROSCI.2044-15.2016

Fee, M. S., Mitra, P. P., & Kleinfeld, D. (1996). Automatic sorting of multiple unit neuronal signals in the presence of anisotropic and non-Gaussian variability. *J Neurosci Methods, 69*(2), 175-188. doi:https://doi.org/10.1016/S0165-0270(96)00050-7

Ghose, G. M., & Harrison, I. T. (2009). Temporal Precision of Neuronal Information in a Rapid Perceptual Judgment. *J Neurophysiol, 101*(3), 1480-1493. doi:Doi 10.1152/Jn.90980.2008

Green, D. M., & Swets, J. A. (1966). *Signal Detection Theory and Psychophysics.* New York: Wiley.

Grothe, I., Neitzel, S. D., Mandon, S., & Kreiter, A. K. (2012). Switching Neuronal Inputs by Differential Modulations of Gamma-Band Phase-Coherence. *The Journal of Neuroscience, 32*(46), 16172. doi:10.1523/JNEUROSCI.0890-12.2012

Haefner, R., Gerwinn, S., Macke, J., & Bethge, M. (2013). Inferring decoding strategy from choice probabilities in the presence of noise correlations. *Nature Neuroscience, 16*, 235-242. Retrieved from http://hdl.handle.net/10101/npre.2012.7014.1

http://www.nature.com/neuro/journal/v16/n2/pdf/nn.3309.pdf

Harris, K. D., & Mrsic-Flogel, T. D. (2013). Cortical connectivity and sensory coding. *Nature, 503*, 51. doi:10.1038/nature12654

Harrison, I. T., Weiner, K. F., & Ghose, G. M. (2013). Inattention blindness to motion in middle temporal area. *Journal of Neuroscience, 33*(19), 8396-8410. doi:10.1523/JNEUROSCI.3409-12.2013

Hatsopoulos, N. G., Ojakangas, C. L., Paninski, L., & Donoghue, J. P. (1998). Information about movement direction obtained from synchronous activity of motor cortical neurons. *Proceedings of the National Academy of Sciences, 95*(26), 15706. doi:10.1073/pnas.95.26.15706

Herrero, Jose L., Gieselmann, Marc A., Sanayei, M., & Thiele, A. (2013). Attention-Induced Variance and Noise Correlation Reduction in Macaque V1 Is Mediated by NMDA Receptors. *Neuron, 78*(4), 729-739. doi:https://doi.org/10.1016/j.neuron.2013.03.029

Hill, D. N., Mehta, S. B., & Kleinfeld, D. (2011). Quality Metrics to Accompany Spike Sorting of Extracellular Signals. *The Journal of Neuroscience, 31*(24), 8699. doi:10.1523/JNEUROSCI.0971-11.2011

Johnson, K. O., Darian-Smith, I., & LaMotte, C. (1973). Peripheral neural determinants of temperature discrimination in man: a correlative study of responses to cooling skin. *J Neurophysiol, 36*, 347-370.

Kanitscheider, I., Coen-Cagli, R., Kohn, A., & Pouget, A. (2015a). MatLab tools for estimating linear Fisher information from population data along with synthetic data and recorded spike count responses from neurons in macaque primary visual cortex to grating images with different orientations and white noise. *CRCNS.org.* *http://dx.doi.org/10.6080/K0PK0D3B*.

Kanitscheider, I., Coen-Cagli, R., Kohn, A., & Pouget, A. (2015b). Measuring Fisher Information Accurately in Correlated Neural Populations. *Plos Computational Biology, 11*(6), e1004218. doi:10.1371/journal.pcbi.1004218

Kohn, A., Coen-Cagli, R., Kanitscheider, I., & Pouget, A. (2016). Correlations and Neuronal Population Information. *Annual Review of Neuroscience, 39*(1), 237-256. doi:10.1146/annurev-neuro-070815-013851

Kohn, A., & Smith, M. A. (2005). Stimulus dependence of neuronal correlation in primary visual cortex of the macaque. *Journal of Neuroscience, 25*(14), 3661-3673. doi:10.1523/JNEUROSCI.5106-04.2005

Krause, B. M., & Ghose, G. M. (2018). Micropools of reliable area MT neurons explain rapid motion detection. *J Neurophysiol, 120*(5), 2396-2409. doi:10.1152/jn.00845.2017

Krug, K., Curnow, T. L., & Parker, A. J. (2016). Defining the V5/MT neuronal pool for perceptual decisions in a visual stereo-motion task. *Philosophical Transactions of the Royal Society B: Biological Sciences*.

Lee, J. C. (1992). Comparison of variance between correlated samples. *Computer Applications in the Biological Sciences, 8*(4), 405-406.

Mitchell, J. F., Sundberg, K. A., & Reynolds, J. H. (2009). Spatial attention decorrelates intrinsic activity fluctuations in macaque area V4. *Neuron, 63*(6), 879-888. doi:S0896-6273(09)00695-3 [pii]
10.1016/j.neuron.2009.09.013

Moreno-Bote, R., Beck, J., Kanitscheider, I., Pitkow, X., Latham, P., & Pouget, A. (2014). Information-limiting correlations. *Nat Neurosci, 17*(10), 1410-1417. doi:10.1038/nn.3807
http://www.nature.com/neuro/journal/v17/n10/abs/nn.3807.html#supplementary-information

Nandy, A. S., Nassi, J. J., & Reynolds, J. H. (2017). Laminar Organization of Attentional Modulation in Macaque Visual Area V4. *Neuron, 93*(1), 235-246. doi:10.1016/j.neuron.2016.11.029

Newsome, W. T., Britten, K. H., & Movshon, J. A. (1989). Neuronal Correlates of a Perceptual Decision. *Nature, 341*(6237), 52-54. Retrieved from <Go to ISI>://A1989AN95800053

Ni, A. M., Ruff, D. A., Alberts, J. J., Symmonds, J., & Cohen, M. R. (2018). Learning and attention reveal a general relationship between population activity and behavior. *Science, 359*(6374), 463. doi:10.1126/science.aao0284

Nienborg, H., Cohen, M. R., & Cumming, B. G. (2012). Decision-Related Activity in Sensory Neurons: Correlations Among Neurons and with Behavior. *Annual Review of Neuroscience, 35*(1), 463-483. doi:doi:10.1146/annurev-neuro-062111-150403

Nienborg, H., & Cumming, B. (2010). Correlations between the activity of sensory neurons and behavior: how much do they tell us about a neuron's causality? *Current Opinion in Neurobiology, 20*(3), 376-381. doi:https://doi.org/10.1016/j.conb.2010.05.002

Nienborg, H., & Cumming, B. G. (2009). Decision-related activity in sensory neurons reflects more than a neuron's causal effect. *Nature, 459*, 89-92. Retrieved from http://dx.doi.org/10.1038/nature07821

http://www.nature.com/nature/journal/vaop/ncurrent/suppinfo/nature07821_S1.html

Nogueira, R., Peltier, N. E., Anzai, A., DeAngelis, G. C., Martínez-Trujillo, J., & Moreno-Bote, R. (2020). The Effects of Population Tuning and Trial-by-Trial Variability on Information Encoding and Behavior. *The Journal of Neuroscience, 40*(5), 1066. doi:10.1523/JNEUROSCI.0859-19.2019

Pack, C. C., & Born, R. T. (2001). Temporal dynamics of a neural solution to the aperture problem in visual area MT of macaque brain. *Nature, 409*(6823), 1040-1042. doi:10.1038/35059085

Parker, A. J. (2013). A micro-pool model for decision-related signals in visual cortical areas. *Frontiers in Computational Neuroscience, 7*. doi:10.3389/fncom.2013.00115

Parker, A. J., & Hawken, M. J. (1985). Capabilities of monkey cortical cells in spatial resolution tasks. *Journal of the Optical Society of America A-Optics Image Science and Vision, A2*, 1101-1114.

Parker, A. J., & Newsome, W. T. (1998). Sense and the single neuron: Probing the physiology of perception. *Annual Review of Neuroscience, 21*, 227-277. Retrieved from <Go to ISI>://000072446400009

https://www.annualreviews.org/doi/pdf/10.1146/annurev.neuro.21.1.227

Pettine, W. W., Steinmetz, N. A., & Moore, T. (2019). Laminar segregation of sensory coding and behavioral readout in macaque V4. *Proceedings of the National Academy of Sciences of the United States of America, 116*(29), 14749-14754. doi:10.1073/pnas.1819398116

Pitkow, X., Liu, S., Angelaki, D. E., DeAngelis, G. C., & Pouget, A. (2015). How Can Single Sensory Neurons Predict Behavior? *Neuron, 87*(2), 411-423. Retrieved from <Go to ISI>://WOS:000361144200015

Poort, J., Raudies, F., Wannig, A., Lamme, Victor A. F., Neumann, H., & Roelfsema, Pieter R. (2012). The Role of Attention in Figure-Ground Segregation in Areas V1 and V4 of the Visual Cortex. *Neuron, 75*(1), 143-156. doi:https://doi.org/10.1016/j.neuron.2012.04.032

Poort, J., & Roelfsema, P. R. (2009). Noise Correlations Have Little Influence on the Coding of Selective Attention in Area V1. *Cerebral Cortex, 19*(3), 543-553. doi:10.1093/cercor/bhn103

Pouget, A., Dayan, P., & Zemel, R. S. (2003). Inference and computation with population codes. *Annual Review of Neuroscience, 26*, 381-410.

Price, N. S. C., & Born, R. T. (2010). Timescales of Sensory- and Decision-Related Activity in the Middle Temporal and Medial Superior Temporal Areas. *The Journal of Neuroscience, 30*(42), 14036. doi:10.1523/JNEUROSCI.2336-10.2010

Prince, S. J. D., Pointon, A. D., Cumming, B. G., & Parker, A. J. (2002). Quantitative analysis of the responses of V1 neurons to horizontal disparity in dynamic random-dot stereograms. *J Neurophysiol, 87*(1), 191-208. Retrieved from <Go to ISI>://000173155700018

Rizk, M., & Wolf, P. D. (2009). Optimizing the automatic selection of spike detection
thresholds using a multiple of the noise level. *Medical & Biological Engineering &*
Computing, 47(9), 955-966. doi:10.1007/s11517-009-0451-2
Roelfsema, P. R., Lamme, V. A. F., & Spekreijse, H. (1998). Object-based attention in the
primary visual cortex of the macaque monkey. *Nature, 395*(6700), 376-381.
doi:10.1038/26475
Rohenkohl, G., Bosman, C. A., & Fries, P. (2018). Gamma Synchronization between V1 and
V4 Improves Behavioral Performance. *Neuron, 100*(4), 953-963 e953.
doi:10.1016/j.neuron.2018.09.019
Romo, R., Hernandez, A., Zainos, A., & Salinas, E. (2003). Correlated neuronal discharges
that increase coding efficiency during perceptual discrimination. *Neuron, 38*(4), 649-
657. Retrieved from <Go to ISI>://000183104000014
Ruff, D. A., & Cohen, M. R. (2014). Attention can either increase or decrease spike count
correlations in visual cortex. *Nature Neuroscience, 17*(11), 1591-1597.
doi:10.1038/nn.3835
Ruff, D. A., & Cohen, M. R. (2016). Stimulus Dependence of Correlated Variability across
Cortical Areas. *The Journal of Neuroscience, 36*(28), 7546.
doi:10.1523/JNEUROSCI.0504-16.2016
Sanayei, M., Chen, X., Chicharro, D., Distler, C., Panzeri, S., & Thiele, A. (2018). Perceptual
learning of fine contrast discrimination changes neuronal tuning and population
coding in macaque V4. *Nat Commun, 9*(1), 4238. doi:10.1038/s41467-018-06698-w
Shadlen, M. N., Britten, K. H., Newsome, W. T., & Movshon, J. A. (1996). A computational
analysis of the relationship between neuronal and behavioral responses to visual
motion. *Journal of Neuroscience, 16*(4), 1486-1510.
Shamir, M., & Sompolinsky, H. (2004). Nonlinear Population Codes. *Neural Computation,*
16(6), 1105-1136. doi:10.1162/089976604773717559
Shiozaki, H. M., Tanabe, S., Doi, T., & Fujita, I. (2012). Neural Activity in Cortical Area V4
Underlies Fine Disparity Discrimination. *The Journal of Neuroscience, 32*(11), 3830-
3841. doi:10.1523/jneurosci.5083-11.2012
Smith, J. E. T., Beliveau, V., Schoen, A., Remz, J., Zhan, C. A. A., & Cook, E. P. (2015).
Dynamics of the functional link between area MT LFPs and motion detection. *J*
Neurophysiol, 114(1), 80-98. doi:10.1152/jn.00058.2015
Smith, J. E. T., Masse, N. Y., Zhan, C. A., & Cook, E. (2012). Linking Neural Activity to Visual
Perception: Separating Sensory and Attentional Contributions. In S. Molotchnikoff &
J. Rouat (Eds.), *Visual Cortex: Current Status and Perspectives*: IntechOpen.
Smith, J. E. T., Zhan, C., an, A., & Cook, E. P. (2011). The Functional Link between Area MT
Neural Fluctuations and Detection of a Brief Motion Stimulus. *The Journal of*
Neuroscience, 31(38), 13458. doi:10.1523/JNEUROSCI.1347-11.2011
Smith, M. A., & Kohn, A. (2008). Spatial and temporal scales of neuronal correlation in
primary visual cortex. *Journal of Neuroscience, 28*(48), 12591-12603.
doi:10.1523/JNEUROSCI.2929-08.2008
Smith, M. A., & Sommer, M. A. (2013). Spatial and temporal scales of neuronal correlation in
visual area V4. *Journal of Neuroscience, 33*(12), 5422-5432.
doi:10.1523/JNEUROSCI.4782-12.2013
Tanabe, S., Umeda, K., & Fujita, I. (2004). Rejection of false matches for binocular
correspondence in macaque visual cortical area V4. *Journal of Neuroscience, 24*(37),
8170-8180. Retrieved from <Go to ISI>://000223884400020

Tolhurst, D. J., Movshon, J. A., & Dean, A. F. (1983). The Statistical Reliability of Signals in
Single Neurons in Cat and Monkey Visual Cortex. *Vision Res, 23*(8), 775-785.
Tripp, B. P. (2012). Decorrelation of Spiking Variability and Improved Information Transfer
Through Feedforward Divisive Normalization. *Neural Computation, 24*(4), 867-894.
doi:10.1162/NECO_a_00255
Ungerleider, L. G., Galkin, T. W., Desimone, R., & Gattass, R. (2008). Cortical connections of
area V4 in the macaque. *Cerebral Cortex, 18*(3), 477-499. Retrieved from <Go to
ISI>://000253490000001
Verhoef, B.-E., & Maunsell, J. H. R. (2017). Attention-related changes in correlated neuronal
activity arise from normalization mechanisms. *Nature Neuroscience, 20*, 969.
doi:10.1038/nn.4572
https://www.nature.com/articles/nn.4572#supplementary-information
Wallisch, P., Lusignan, M. E., Benayoun, M. D., Baker, T. I., Seth Dickey, A., & Hatsopoulos, N.
G. (2014). Information Theory. In *MATLAB for Neuroscientists : An Introduction to*
Scientific Computing in MATLAB (2 ed.). London: Elsevier Science & Technology.
Wasmuht, D. F., Parker, A. J., & Krug, K. (2019). Interneuronal correlations at longer time
scales predict decision signals for bistable structure-from-motion perception. *Nature*
Scientific Reports, in press.
Weiner, K. F., & Ghose, G. M. (2014). Rapid shape detection signals in area V4. *Frontiers in*
Neuroscience, 8, 294. Retrieved from
https://www.frontiersin.org/article/10.3389/fnins.2014.00294
Weiner, K. F., & Ghose, G. M. (2015). Population coding in area V4 during rapid shape
detections. *J Neurophysiol, 113*(7), 3021-3034. doi:10.1152/jn.01044.2014
Yu, X., & Gu, Y. (2018). Probing Sensory Readout via Combined Choice-Correlation Measures
and Microstimulation Perturbation. *Neuron, 100*(3), 715-727.e715.
doi:https://doi.org/10.1016/j.neuron.2018.08.034
Zylberberg, J., Pouget, A., Latham, P. E., & Shea-Brown, E. (2017). Robust information
propagation through noisy neural circuits. *Plos Computational Biology, 13*(4),
e1005497. doi:10.1371/journal.pcbi.1005497
2192

A
i) Start
ii) Hold
fixation
iii) Present
RDS
iv) Popout
& choice

*fixation
point*

◇ *Gaze*   ↘ *Choice saccade*

B

M135   M138   M135   M138

% correct

RT (s)

Popout change in disparity (°)

Fail
Corr.

C

M135   – –   0.00°
M138   ——   ±0.05°
        ——   ±0.10°

% correct

RT (s from popout)

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Figure 11

Figure S1

Accompanies Figure 1

Figure S2

Accompanies Figure 2

Figure S3

Accompanies Figure 3

Figure S4

Accompanies Figure 4

Figure S5

Accompanies Figure 5

Figure S6

Accompanies Figure 6

Figure S7

Accompanies Figure 7

Figure S8

Accompanies Figure 8