Herpes Simplex Virus Infection, Acyclovir and IVIG Treatment All Independently Cause Gut

Dysbiosis.

Chandran Ramakrishna[1], Stacee Mendonca[1,] Paul M. Ruegger[2], Jane Hannah Kim[2], James

Borneman[2] and Edouard Cantin[1*].

Department of Molecular Immunology[1], Beckman Research Institute of City of Hope, Duarte, CA

91010 and Department of Microbiology and Plant Pathology[2], University of California, Riverside,

CA 92521.

*Corresponding Authors: Edouard M. Cantin

Beckman Research Institute of City of Hope

Department of Molecular Immunology

Fox Plaza North, Room 100B

1500 E. Duarte Rd, Duarte CA 91010-3012

Phone: +1 (626) 301-8480

Email: ecantin@coh.org

James Borneman

University of California

Department of Microbiology and Plant Pathology

3401 Watkins Drive

Multidisciplinary Research Building Room 4130

Riverside, CA 92521

Phone: +1 (951) 827-3584

Email: borneman@ucr.edu

Abstract.

Herpes simplex virus 1 (HSV) is a ubiquitous human virus resident in a majority of the global population as a latent infection. Acyclovir (ACV), is the standard of care drug used to treat primary and recurrent infections, supplemented in some patients with intravenous immunoglobulin (IVIG) treatment to suppress deleterious inflammatory responses. We found that HSV, ACV and IVIG can all independently disrupt the gut bacterial community in a sex biased manner when given to uninfected mice. Treatment of HSV infected mice with ACV or IVIG alone or together revealed complex interactions between these drugs and infection that caused pronounced sex biased dysbiosis. ACV reduced *Bacteroidetes* levels in male but not female mice, while levels of the Anti-inflammatory Clostridia (AIC) were reduced in female but not male mice, which is significant as these taxa are associated with protection against the development of GVHD in hematopoietic stem cell transplant (HSCT) patients. Gut barrier dysfunction is associated with GVHD in HSCT patients and ACV also decreased *Akkermansia muciniphila,* which is important for maintaining gut barrier functionality. Cumulatively, our data suggest that long-term prophylactic ACV treatment of HSCT patients may contribute to GVHD and potentially impact immune reconstitution. These data have important implications for other clinical settings, including HSV eye disease and genital infections, where ACV is given long-term.

Author Summary.

        Primary and reactivated HSV and VZV infections are treated with Acyclovir (ACV), an antiviral drug that blocks viral DNA synthesis. In some patients IVIG is used as adjunctive therapy to block deleterious inflammation. Long term preventative treatment of patients who receive stem transplants for various blood cancers has been successful in preventing life threatening reactivated HSV and VZV infections, but GVHD remains a major factor limiting transplant

success. Studies reported here reveal that HSV infection, ACV and IVIG given alone can all

disrupt the gut microbiota and that complex interactions between these drugs and infection results

in even more pronounced sex biased changes in the gut bacteria community structure.

Importantly, ACV treatment decreased the levels of specific bacterial taxa, including the anti-

inflammatory *Clostriodia* and *Bacteroidetes* that have been shown to protect against development

of GVHD in stem cell transplant patients. These data suggest that long term preventative

treatment of patients with ACV may contribute to GVHD in transplant patients and have negative

consequences in other HSV induced diseases treated long term with ACV. The health effects of

long term ACV and IVIG treatments warrant further clinical studies.

Introduction.

Herpes Simplex Virus type 1 (HSV), a ubiquitous human virus is the major cause of HSV

encephalitis (HSE), the most prevalent sporadic encephalitis resulting from either primary

infection or reactivation of latent virus. However, despite improved diagnostic procedures and

effective antiviral therapies, most HSE survivors have persistent neurological impairments,

including memory and behavior disturbances, dysphasia and seizures, and only 50-65% of these

survivors return to independent living [1, 2]. A delay in initiating Acyclovir (ACV) treatment past

the second hospital day is associated with poor neurological outcomes [3, 4]. Recent clinical trials

evaluating prolonged oral ACV/valaciclovir (VACV) treatment following standard 14-day

intravenous ACV treatment reported improved neurocognitive outcomes in neonates but not

adults for reasons that are obscure [5, 6]. Although, it is generally accepted that replication

induced pathology underlies HSV related neurological dysfunction, supporting experimental or

clinical evidence is lacking. Overwhelming evidence has linked inflammation to the development

of various neurological disorders and neuropsychiatric diseases, including Alzheimer's disease

(AD), schizophrenia, autism spectrum disorder (ASD), multiple sclerosis (MS), Parkinson's

disease (PD), depression and anxiety [7-9].

Having unequivocally established that HSE arises from exaggerated CNS inflammatory

responses and that the immunomodulatory activities of intravenous immunoglobulins (IVIG) can

prevent HSE in a mouse model [10], we tested the hypothesis that persistent inflammation, which

is documented in humans and mice after HSE [11-14], causes neurobehavioral impairments in

survivors, that should be impeded by IVIG's anti-inflammatory activity [10]. Compared to treatment

of HSV infected mice with ACV or PBS alone, treatment with ACV+IVIG from day 4 pi reduced

CNS inflammation and anxiety, consistent with our hypothesis. Strikingly, development of learning

and memory (LM) deficits that were evident only in female PBS treated mice, were inhibited by

ACV treatment and counterintuitively, aggravated by ACV+IVIG treatment. Treatment of infected

male mice with ACV+IVIG also impaired LM compared to ACV or PBS alone, revealing that IVIG

antagonized the beneficial effects of ACV [15]. Intriguingly, the differential antagonistic effects of

ACV+IVIG on cognitive behavior in HSV infected mice, compared to ACV and PBS treatment

alone, were reflected in differential serum proteomic profiles [15]. These reported antagonistic

effects of ACV and IVIG on LM present a conundrum, since they are at odds with the known

mechanisms of action of these drugs.

Rapidly accumulating evidence is revealing the critical role of the microbiome in regulating

brain homeostasis and function such that perturbation of the gut bacteria community structure

and function is increasingly being implicated in a variety of neurodegenerative and

neuropsychiatric diseases. In an effort to gain insight into how HSV induces LM impairment and

the paradoxical effects of ACV and IVIG, we investigated a role for the gut microbiota. HSV

infection, ACV and IVIG were all associated with significant disruption of the gut bacterial

community structure that was sex biased. Furthermore, treating HSV infected mice with either

ACV or IVIG alone or both drugs together resulted in more pronounced sex-biased shifts in the

gut bacterial community structure compared to uninfected mice. These results have significant

clinical implications, particularly when patients receive prolonged ACV or IVIG treatment.

Results.

Equal numbers (n=8) of female and male C57BL/6 mice were bilaterally inoculated with

virulent HSV1 strain 17+ (1×10^5 PFU/eye) by corneal scarification as previously described [15].

At day 4 post infection (pi), ACV was administered at 1.25 mg / mouse by intraperitoneal injection

(ip) daily for 3 days, while IVIG was given as single dose of 25 mg/mouse by ip injection on day

4pi [15]. Fresh fecal pellets (n=1-2/ mouse) were collected on day 7 pi and stored at -80°C until

processed for Illumina 16S rRNA gene sequencing to determine the effects of infection and drug

treatment on the gut microbiome. Normal male and female mice differed in gut bacteria

composition and unexpectedly, HSV ocular infection caused further shifts in the gut bacteria

community and amplified this sex difference, as shown in a PCoA plot of Hellinger beta diversity

distance values for infected compared to uninfected male and female mice (**Figure 1A**; P<0.05,

Adonis Tests). In addition, HSV infection had a greater effect on gut bacterial communities in

males (P=0.003) compared to females (P=0.011) (**Figure 1A**). Significant differences were

observed at the phyla level, particularly for firmicutes (**Figure 1B**) with more marked differences

evident at the species level for *Clostridium aerotolerans* and other clostridial species, for example

Clostridium XIVa that ferment carbohydrates in the gut resulting in production of short chain fatty

acids (SFCs) that contribute to barrier integrity and also exhibit anti-inflammatory properties

(**Figure 1C**). A notable difference was also observed for *Akkermansia muciniphila* that has many

health promoting activities, including maintaining gut barrier health (**Figure 1C**).

Treating HSV infected mice with ACV from day 4 pi for three days resulted in even more

drastic shifts in the gut bacteria composition and exaggerated sex differences (**Figure 2A**), than

for infection alone. Considerable abundance changes were evident at the Phyla level for

Bacteroidetes, *Firmicutes* and *Verrucomicrobia* (**Figure 2B**) and at the species level (**Figure 2C**).

Notably, whereas HSV infection reduced the abundance of *Firmicutes* significantly in male but

not female mice (**Figure 1B**), ACV reversed this effect restoring the abundance to the level in

uninfected male mice, while also increasing the abundance in female mice (**Figure 2B** and **Figure**

1B). Notable abundance changes at the species level included drastic suppression of *Clostridium*

aerotolerans in infected male mice compared to increased abundance in females (**Figure 1C**),

while ACV treatment further increased this abundance only in females (**Figure 2C**). *Akkermansia*

muciniphila abundance was increased by infection in male mice but reduced in females (**Figure**

1C), while ACV treatment resulted in total suppression of this species in female mice compared

to a marked reduction in male mice (**Figure 2C**). There are many other similar changes in species

abundance that are differentially impacted by ACV treatment in a sex-biased manner, indicative

of complex interactions between infection, ACV effects on infected host cells, and bacteria, as

well as metabolites produced by bacterial metabolism of ACV.

Treatment of uninfected mice with IVIG alone also shifted the gut bacteria community

composition with a notable marked sex effect as determined by a beta diversity analysis (**Figure**

3). Males and females showed a major reduction in *A. muciniphila,* and a lesser reduction of

Verrucomicrobia in males, compared to females that showed increased abundance of this phylum

in response to IVIG treatment (**Figure 4**). The abundance of many other bacterial species was

differentially altered by IVIG treatment of males and females, for example, *Clostridium*

aerotolerans, *Bacteroides acidifaciens* and *Porphyromonadaceae* (**Figure 4B**). The response to

IVIG was distinct in HSV infected mice, and the complex interactions between infection, ACV and

IVIG were also evident at the phyla and species levels and were strongly sex biased as well

(**Figure 4A and 4B**). IVIG treatment decreased *A. muciniphila* abundance markedly in infected

males and females as did ACV, whereas in contrast, treatment with ACV+IVIG caused a notable

increase in its abundance, indicative of antagonistic effects of these two drugs in the context of

infection (**Figure 4B**) In a similar vein, *C. aerotolerans* abundance increased markedly in males,

but was unchanged in females treated with IVIG, while in contrast, it was strongly decreased in

males but slightly increased in females treated with ACV alone. In contrast, treatment with

ACV+IVIG suppressed an IVIG-induced increase in males and an ACV-induced increase in

females, revealing antagonism between ACV and IVIG in the context of HSV infection (**Figure**

4B).

Patients with hematologic and other malignancies have benefited immensely from

allogeneic hematopoietic stem cell transplantation (allo-HSCT or HSCT), which can be a potent

curative immunotherapy. However, life threatening complications such as graft-versus-host

disease (GVHD), relapse, and infections that include reactivated HSV and VZV limit its application

[16]. HSV and varicella zoster (VZV) reactivation has been successfully suppressed by

prophylactic ACV treatment, though ACV-resistant (ACVr) HSV is an emerging problem [17, 18].

Long term ACV prophylactic treatment is now routine for HSCT patients, because it was found to

correlate with reduced HSV and ACVr HSV disease in those treated for longer than 1 year [19].

Given this routine clinical practice, we evaluated the effects of ACV on fecal bacteria,

because gut microbes have been implicated in GVHD pathophysiology and because we posit that

ACV contributes to the development of GVHD by changing the gut microbiota. First, we identified

gut bacterial changes in humans with GVHD [20-30]. Next, we determined whether the ACV-

induced changes that we detected in this mouse study matched those GVHD-associated

changes. Whenever we identified taxa that were altered in both types of studies, the direction of

the change was the same, and it was consistent with our hypothesis that ACV contributes to the

development of human GVHD by changing the gut microbiota. In the following, we describe these

results, and we note that these ACV-induced changes were only observed in the HSV-infected

mice and not in the uninfected mice.

Reduced levels of several taxa belonging to the phylum *Bacteroidetes* have been shown

to be associated with GVHD, indicating that these gut bacteria may play a protective role. In a

pediatric study, GVHD patients had lower levels of the family *Bacteroidaceae* and the genus

Parabacteroides [30]. In a longitudinal study, pediatric patients that had lower levels of

Bacteroidetes prior to HSCT were more likely to develop GVHD [24]. In our study, all three if these

taxa were reduced by ACV treatment in male but not female mice (**Figure 5A**).

Reduced levels of Anti-Inflammatory Clostridia (AIC) have also been detected in human

GVHD patients [20, 23-25, 27-30], indicating that these gut bacteria may play a protective role.

This terminology was first introduced by Piper et al. [31] in the context of short bowel syndrome,

and then introduced to the GVHD literature by Simms-Waldrip et al. [30]. AIC taxa include

members of the families *Clostridiaceae*, *Erysipelotrichaceae*, *Eubacteriaceae*, *Lachnospiraceae*

and *Ruminococcaceae*. In a pediatric study, decreases in *Blautia* and *Clostridium bolteae* were

associated with the development of GVHD [30]. In an adult study, lower levels of *Blautia*, *Blautia*

hansenii, and *Blautia stercoris* were associated with the development of GVHD [28]. In a

longitudinal study, reduced levels of the *Blautia* before HSCT was shown to be a predictive marker

for the development of GVHD [27]. In our study, all of these taxa were reduced by ACV treatment

in female but not male mice (**Figure 5B**).

In a more detailed analysis of AIC bacteria, we observed that while HSV infection

increased the abundance of *Blautia hansenii* only in males, ACV treatment reduced its abundance

in females but had no effect on its abundance in males (**Supplemental Figure 1**). Remarkably,

a dramatic increase in *B. hansenii* in uninfected females was observed after IVIG treatment, and

this increase was abrogated by ACV (compare NoHSV_F, NoHSV_IVIG_F and

NoHSV_ACVplusIVIG_F) (**Supplemental Figure 1**), a result that supports sex-based differential

effects of these drugs. However, during HSV infection, both IVIG and ACV reduced *B. hansenii*

in females, whereas only IVIG reduced abundance in males. Interestingly, HSV infection

significantly increased the abundance of the AIC genera *Blautia, Allobaculum,* and *Clostridium*

XVIII but not *Turicibacter* in both males and females (**Supplemental Figure 2**). ACV treatment of

HSV infected female mice resulted in significant decreases in the abundances of 4 AIC genera:

Blautia, *Allobaculum*, *Clostridium* XVIII and *Turicibacter,* whereas in infected males, ACV

decreased the abundance of *Marvinbryantia* and *Oscillibacter* (**Supplemental Figure 2**). In

addition, ACV increased the abundance of *Turicibacter* in uninfected females but not males.

Finally, the two most abundant operational taxonomic units (OTUs), which exhibited a

change in their relative abundances due to ACV treatment, were assigned to the family

Porphyromonadaceae and the species *A. muciniphila* (**Figure 5C**). While we did not find these

taxa associated with GVHD in prior human studies, GVHD has been associated with intestinal

barrier dysfunction [32-36]. Supporting our hypothesis that ACV contributes to the development

of GVHD by changing the gut microbiota, members of the *Porphyromonadaceae* have been

shown to cause gut barrier dysfunction [37, 38], and our *Porphyromonadaceae* OTU was

increased in its abundance by ACV. In addition, *A. muciniphila* was decreased by ACV treatment

in our study, and it has been shown to strengthen gut barrier functioning [39-41].

Discussion.

Our intention in this brief report is to alert the scientific community and especially clinicians

to the fact that HSV infection, the antiviral drug ACV, and the immunomodulatory biological, IVIG,

can all independently result in significant perturbations of the gut bacterial communities. Our data

reveal complex interactions between HSV infection and ACV or/and IVIG treatment that result in

marked alterations to gut bacterial communities. Although the clinical consequences of these

changes have not yet been elucidated, they could have profound implications in several settings

including HSCT-associated GVHD.

Though the mechanisms by which ocular HSV infection causes gut dysbiosis are unclear,

neuroinflammatory mechanisms and effects on the enteric nervous system via connected

brainstem neuronal circuits can be envisaged [15, 42]. Indeed, recent paradigm-shifting reports

reveal that peripheral neurons, including nociceptive and sensory neurons, can directly sense and

respond to environmental alarms by releasing neuropeptides that can regulate immune responses

in target organs including the gut [43, 44]. Persistence of gut dysbiosis was not evaluated here,

but results from a behavioral study alluded to earlier suggest long-term effects of infection and

drug treatment on gut bacterial ecology should be investigated [15]. Sex biased effects on HSV

induced dysbiosis merit further study, as these may involve microglial responses to HSV infection

and the microglial compartment is known to be regulated by the microbiota in a sex biased manner

[45-47].

The mechanism by which ACV, the standard antiviral for HSV infections, changes the gut

microbiota likely involves its uptake into bacteria. ACV is preferentially phosphorylated by the viral

encoded thymidine kinase (Tk) resulting in cell retention and eventual incorporation into viral DNA

resulting in inhibition of viral replication via DNA chain termination. Because Tk is conserved in

numerous bacterial species, ACV can be taken up and incorporated into DNA, resulting in

bactericidal effects [48-51]. Indeed, early studies on DNA replication mechanisms relied on

labeling bacterial DNA with tritiated thymidine and many bacterial taxa can be imaged using

nucleoside analogues such as 1-(2_-deoxy-2_-fluoro-_-D-arabinofuranosyl)-5-[125I] iodouracil

([125I]FIAU) that are substrates for HSV Tk [52-55]. Incorporation of [*methyl*-^{3}H]thymidine into

DNA has been unequivocally demonstrated for members of the *Clostridium* genus [56] and our

data show ACV reduced the abundance of the *Blautia* genus (order *Clostridiales;* [57]) *Blautia*

hansenii, *Blautia stercoris*, and *Clostridium bolteae* in females but not males. Additionally,

interrogating the NCBI reference genome sequence for *Blautia hansenii* confirmed the presence

of a thymidine kinase enzyme. Our data are therefore consistent with ACV causing dysbiosis by,

at least in part, inhibiting the growth of various bacteria taxa via the Tk mechanism, though other

mechanisms involving bacterial metabolism of ACV cannot be excluded. Clearly, the mechanisms

by which ACV affects gut bacterial ecology are complex, which is further supported by the sex-

biased effects.

We also explored the effects of IVIG treatment alone and in combination with ACV in HSV-

infected and uninfected mice, because IVIG has been used to treat HSV encephalitis (HSE) and

is also a frontline therapy for autoimmune encephalitis, which is triggered by HSE and other insults

[58-60]. Moreover, IVIG is being evaluated in a randomized control trial for children with all-cause

encephalitis to determine whether neurological outcomes are improved compared to standard

antiviral therapy alone, which is similar to our behavioral study that generated paradoxical results

[15, 61]. Reports that IVIG's antigenic repertoire includes reactivities to a variety of gut commensal

antigens and metabolites have increased recently [62-64], which is consistent with a report that

gut commensals can somehow trigger systemic IgG responses under homeostatic conditions that

protect against systemic infection [65, 66]. We speculate that by neutralizing bacterial/host

antigens/metabolites, IVIG is able to influence host immunity, the nervous system, and other

physiological processes, resulting in perturbation of gut bacteria ecology. We speculate that the

disparate and complex effects of ACV and IVIG alone and in combination on the gut bacteria

ecology likely account for their antagonistic effects on cognitive behavior in mice latently infected

with HSV that we alluded to earlier [15].

This study has several limitations. Being exploratory in nature, analyses of the gut bacteria

were done at a single time point immediately after infection or drug treatment, rather than as a

longitudinal study that would have provided information on the persistence of the dysbiotic state

as well as mechanistic insights as to how HSV, ACV and IVIG provoke dysbiosis. Ideally, the

effects of ACV should be tested in latently infected mice, since virtually all HSCT patients harbor

latent HSV. However, because HSV infection alone disrupts the gut bacterial community,

assessing the effects of ACV on the gut bacteria community structure in the latently infected mice

would likely be difficult. Because ACV was given ip to mice but usually orally to HSCT patients

[67], its effects on the gut bacteria community maybe underestimated in our study.

Notwithstanding these caveats, our finding that ACV treatment of HSV infected mice

decreased the relative abundances of several bacterial taxa is important because these bacteria

have been negatively correlated with the induction of and mortality from GVHD in HSCT patients

[24, 27, 28, 30]. These results are also consistent with our hypothesis that ACV contributes to the

development of GVHD by changing the gut microbiota. In the context of allo-HSCT, GVHD occurs

when donor immune cells recognize recipient tissues as foreign, leading to immune-mediated

damage to several organs and tissues including the gastrointestinal tract. This has led

researchers to posit that the reduction of anti-inflammatory bacteria such as AIC contribute to

GVHD pathology [30]. The results from our study extend this hypothesis to include ACV treatment

as a putative contributor to GVHD, because ACV reduced AIC bacteria in the gut. ACV treatment

also decreased the relative abundances of several members of the *Bacteroidetes*, some of which

have been shown to exhibit anti-inflammatory properties [68-71]. More relevantly, the capsular

polysaccharide A (PSA) from *Bacteroides fragilis* reduced HSV-associated mortality in mice by

dramatically reducing immune-mediated inflammation [72]. In addition, the two most abundant

OTUs identified in our study, whose relative abundances were positively (*Porphyromonadaceae*)

and negatively (*A. muciniphila*) correlated with ACV treatment, have been shown to weaken [37,

38] and strengthen [39-41] gut barrier function, respectively. These results provide an additional

link between ACV treatment and GVHD, because barrier dysfunction, which can cause systemic

inflammation, is a hallmark of GVHD [32-36]. Finally, long-term ACV prophylaxis initiated early

after HSCT might also impair immune reconstitution based on results from a study of antibiotic

depletion of gut bacteria in a murine model of syngeneic bone marrow transplantation [73]. These

tantalizing results warrant independent validation and further detailed studies using a murine

autologous BMT model to more rigorously evaluate the impact of long-term ACV prophylaxis on

GVHD and engraftment, because results from such studies might eventually lead to improved

outcomes for HSCT patients. Ideally, such future studies should be performed with mice harboring

wild microbiota, because several recent reports show that immune responses in mice with wild

microbiomes model human immune responses more closely than conventional mice with SPF

microbiota [74-76].

Materials and Methods.

Ethics Statement

All animal procedures were performed with prior approval of the City of Hope Institutional Animal

Care and Use Committee (IACUC) under protocol # 07043 and within the framework of the Guide

for the Care and Use of Laboratory Animals. C57BL6/J (B6) were bred in the vivarium at City of

Hope.

Mouse Studies

Master stocks of HSV1 strain 17 composed of only of cell-released virus were prepared in

and their titers determined on mycoplasma-free CV-1 cell monolayers. Single use aliquots of virus

in Hanks balanced salt solution supplemented with 2% fetal bovine serum were stored at -80°C.

Male and female mice, 6–8 weeks of age, were infected with HSV1 17$^+$, a virulent strain. Mice

were sedated with ketamine (60 mg/kg) and xylazine (5 mg/kg) prior to HSV inoculation by corneal

scarification. B6 mice were bilaterally inoculated with 1x 10^5 PFU per eye and monitored daily as

previously described [15, 77].

Administration of Acyclovir and Intravenous Immunoglobulins.

ACV obtained from (APP Pharmaceuticals, Schaumburg, IL) was given at 50 mg/kg of

body weight by intraperitoneal (ip) injection daily for 3 days starting on day 4 pi and PBS was

given according to the same schedule to control mice. IVIG (Carimune, NF) obtained from CSL

Behring (King of Prussia, PA, USA) was given ip as a single 0.5 ml dose (25 mg/mouse) on day

4 pi or it was given in combination with a 3 day course of ACV.

Illumina Bacterial 16S rRNA gene sequencing.

Illumina bacterial 16S rRNA gene libraries were constructed as follows. PCRs were

performed in an MJ Research PTC-200 thermal cycler (Bio-Rad Inc., Hercules, CA, USA) as 25

µl reactions containing: 50 mM Tris (pH 8.3), 500 µg/ml bovine serum albumin (BSA), 2.5 mM

MgCl$_2$, 250 µM of each deoxynucleotide triphosphate (dNTP), 400 nM of the forward PCR primer,

200 nM of each reverse PCR primer, 1 µl of DNA template, and 0.25 units JumpStart Taq DNA

polymerase (Sigma-Aldrich, St. Louis, MO, USA). PCR primers 515F

(GTGCCAGCMGCCGCGGTAA) and 806R (GGACTACHVGGGTWTCTAAT) were used to

targeted the 16S rRNA gene containing portions of the hypervariable regions V4 and V5, with the

reverse primers including a 12-bp barcode [78]. Thermal cycling parameters were 94°C for 5 min;

35 cycles of 94°C for 20 s, 50°C for 20 s, and 72°C for 30 s, and followed by 72°C for 5 min. PCR

products were purified using the MinElute 96 UF PCR Purification Kit (Qiagen, Valencia, CA,

USA).

16S rRNA gene data processing.

We used the UPARSE pipeline for de-multiplexing, length trimming, quality filtering and

operational taxonomic units (OTU) picking using default parameters or recommended guidelines

that were initially described in [79] and which have been updated at

https://www.drive5.com/usearch/manual/uparse_pipeline.html. Briefly, after demultiplexing,

sequences were trimmed to a uniform length of 249 bp, then filtered at the recommended 1.0

expected error threshold. Sequences were then dereplicated and clustered into zero-radius OTUs

using the UNOISE3 algorithm [80], which also detects and removes chimeric sequences; this

method is based on making OTUs at 100% identity. An OTU table was then generated using the

otutab command. OTUs having non-bacterial DNA were identified by performing a local BLAST

search [81] of their seed sequences against the nt database. OTUs were removed if any of their

highest-scoring BLAST hits contained taxonomic IDs within Rodentia, Viridiplantae, Fungi, or

PhiX. Taxonomic assignments to the OTUs were performed with SINTAX [82] using RDP

Classifier 16S training set number 16 [83] as the reference database.

16S rRNA gene data analyses.

Beta diversity was measured using QIIME 1.9.1 [84] to calculate a Hellinger beta diversity

distance matrix, which was depicted using principle coordinates analysis (PCoA), and statistically

assessed by performing Adonis tests. Statistical differences among the taxa were determined

using edgeR [85, 86]. Taxa relative abundance figures were made using Prism (GraphPad, La

Jolla, CA). Comparative analyses of the bacterial taxa between human GVHD studies and our

mouse study excluded sequence-selective qPCR, because the selectivity of such assays is

questionable given the conserved nature of the 16S rRNA gene, and because the results of such

studies are not typically validated by sequence analyses. The bacterial sequences have been

deposited in the National Center for Biotechnology Information (NCBI)'s Sequence Read Archive

(SRA) under the BioProject Accession Number PRJNA549765.

References.

1.   McGrath N, Anderson NE, Croxson MC, Powell KF. Herpes simplex encephalitis treated
with acyclovir: diagnosis and long term outcome. Journal of Neurology, Neurosurgery and
Psychiatry. 1997;63(3):321-6. doi: 10.1136/jnnp.63.3.321.
2.   Raschilas F, Wolff M, Delatour Fdr, Chaffaut C, De Broucker T, Chevret S, et al. Outcome
of and Prognostic Factors for Herpes Simplex Encephalitis in Adult Patients: Results of a
Multicenter Study. Clin Infect Dis. 2002;35(3):254-60. doi: 10.1086/341405.
3.   Kennedy PGE, Steiner I. Recent issues in herpes simplex encephalitis. Journal of
NeuroVirology. 2013;19(4):346-50. doi: 10.1007/s13365-013-0178-6.
4.   Bradshaw MJ, Venkatesan A. Herpes Simplex Virus-1 Encephalitis in Adults:
Pathophysiology, Diagnosis, and Management. Neurotherapeutics. 2016;13(3):493-508. doi:
10.1007/s13311-016-0433-7.
5.   Gnann JW, Sköldenberg B, Hart J, Aurelius E, Schliamser S, Studahl M, et al. Herpes
Simplex Encephalitis: Lack of Clinical Benefit of Long-term Valacyclovir Therapy. Clin Infect Dis.
2015;61(5):683-91. doi: 10.1093/cid/civ369.
6.   Kimberlin DW, Whitley RJ, Wan W, Powell DA, Storch G, Ahmed A, et al. Oral Acyclovir
Suppression and Neurodevelopment after Neonatal Herpes. New England Journal of Medicine.
2011;365(14):1284-92. doi: doi:10.1056/NEJMoa1003509.
7.   Chiara G, Marcocci M, Sgarbanti R, Civitelli L, Ripoli C, Piacentini R, et al. Infectious
Agents and Neurodegeneration. Molecular Neurobiology. 2012;46(3):614-38. doi:
10.1007/s12035-012-8320-7.
8.   Hope S, Hoseth E, Dieset I, Mørch RH, Aas M, Aukrust P, et al. Inflammatory markers are
associated with general cognitive abilities in schizophrenia and bipolar disorder patients and
healthy controls. Schizophrenia Research. 2015;165(2-3):188-94. doi:
http://dx.doi.org/10.1016/j.schres.20151654.04.004.
9.   Kuntz T, Gilbert J. Does the brain listen to the gut? eLife. 2016;5:e17052. doi:
10.7554/eLife.17052.
10.   Ramakrishna C, Newo ANS, Shen Y-W, Cantin E. Passively Administered Pooled Human
Immunoglobulins Exert IL-10 Dependent Anti-Inflammatory Effects that Protect against Fatal HSV
Encephalitis. PLoS Pathog. 2011;7(6):e1002071. doi: 10.1371/journal.ppat.1002071.
11.   Cantin EM, Hinton DR, Chen J, Openshaw H. Gamma interferon expression during acute
and latent nervous system infection by herpes simplex virus type 1. J Virol. 1995;69(8):4898-905.
12.   Liu T, Tang Q, Hendricks RL. Inflammatory infiltration of the trigeminal ganglion after
herpes simplex virus type 1 corneal infection. J Virol. 1996;70(1):264-71.
13.   Khanna KM, Lepisto AJ, Decman V, Hendricks RL. Immune control of herpes simplex
virus during latency. Current Opinion in Immunology. 2004;16(4):463-9.
14.   Aurelius E, Forsgren M, Skoldenberg B, Strannegard O. Persistent intrathecal immune
activation in patients with herpes simplex encephalitis. J Infect Dis. 1993;168(5):1248-52.
15.   Ramakrishna C, Golub MS, Chiang A, Hong T, Kalkum M, Cantin EM. Effects of Acyclovir
and IVIG on Behavioral Outcomes after HSV1 CNS Infection. Behavioural Neurology.
2017;2017:14. doi: 10.1155/2017/5238402.
16.   Shono Y, van den Brink MRM. Gut microbiota injury in allogeneic haematopoietic stem
cell transplantation. Nature Reviews Cancer. 2018. doi: 10.1038/nrc.2018.10.
17.   Frobert E, Burrel S, Ducastelle-Lepretre S, Billaud G, Ader F, Casalegno J-S, et al.
Resistance of herpes simplex viruses to acyclovir: An update from a ten-year survey in France.
Antiviral Research. 2014;111(0):36-41. doi: http://dx.doi.org/10.1016/j.antiviral.2014.08.013.
18.   Baumrin E, Cheng MP, Kanjilal S, Ho VT, Issa NC, Baden LR. Severe Herpes Zoster
Requiring Intravenous Antiviral Treatment in Allogeneic Hematopoietic Cell Transplantation

Recipients on Standard Acyclovir Prophylaxis. Biology of Blood and Marrow Transplantation. 2019. doi: https://doi.org/10.1016/j.bbmt.2019.04.015.

19. Dadwal SS. Herpes Virus Infections Other than Cytomegalovirus in the Recipients of Hematopoietic Stem Cell Transplantation. Infect Dis Clin North Am. 2019;33(2):467-84. doi: https://doi.org/10.1016/j.idc.2019.02.012.

20. Taur Y, Jenq RR, Perales MA, Littmann ER, Morjaria S, Ling L, et al. The effects of intestinal tract bacterial diversity on mortality following allogeneic hematopoietic stem cell transplantation. Blood. 2014;124(7):1174-82. Epub 2014/06/19. doi: 10.1182/blood-2014-02-554725. PubMed PMID: 24939656; PubMed Central PMCID: PMCPMC4133489.

21. Weber D, Oefner PJ, Dettmer K, Hiergeist A, Koestler J, Gessner A, et al. Rifaximin preserves intestinal microbiota balance in patients undergoing allogeneic stem cell transplantation. Bone Marrow Transplant. 2016;51(8):1087-92. Epub 2016/03/22. doi: 10.1038/bmt.2016.66. PubMed PMID: 26999466.

22. Shono Y, Docampo MD, Peled JU, Perobelli SM, Velardi E, Tsai JJ, et al. Increased GVHD-related mortality with broad-spectrum antibiotic use after allogeneic hematopoietic stem cell transplantation in human patients and mice. Sci Transl Med. 2016;8(339):339ra71. Epub 2016/05/20. doi: 10.1126/scitranslmed.aaf2311. PubMed PMID: 27194729; PubMed Central PMCID: PMCPMC4991773.

23. Peled JU, Devlin SM, Staffas A, Lumish M, Khanin R, Littmann ER, et al. Intestinal Microbiota and Relapse After Hematopoietic-Cell Transplantation. J Clin Oncol. 2017;35(15):1650-9. Epub 2017/03/16. doi: 10.1200/jco.2016.70.3348. PubMed PMID: 28296584; PubMed Central PMCID: PMCPMC5455763.

24. Biagi E, Zama D, Nastasi C, Consolandi C, Fiori J, Rampelli S, et al. Gut microbiota trajectory in pediatric patients undergoing hematopoietic SCT. Bone Marrow Transplant. 2015;50(7):992-8. doi: 10.1038/bmt.2015.16.

25. Jenq RR, Ubeda C, Taur Y, Menezes CC, Khanin R, Dudakov JA, et al. Regulation of intestinal inflammation by microbiota following allogeneic bone marrow transplantation. J Exp Med. 2012;209(5):903-11. Epub 2012/05/02. doi: 10.1084/jem.20112408. PubMed PMID: 22547653; PubMed Central PMCID: PMCPMC3348096.

26. Holler E, Butzhammer P, Schmid K, Hundsrucker C, Koestler J, Peter K, et al. Metagenomic analysis of the stool microbiome in patients receiving allogeneic stem cell transplantation: loss of diversity is associated with use of systemic antibiotics and more pronounced in gastrointestinal graft-versus-host disease. Biol Blood Marrow Transplant. 2014;20(5):640-5. Epub 2014/02/05. doi: 10.1016/j.bbmt.2014.01.030. PubMed PMID: 24492144; PubMed Central PMCID: PMCPMC4973578.

27. Biagi E, Zama D, Rampelli S, Turroni S, Brigidi P, Consolandi C, et al. Early gut microbiota signature of aGvHD in children given allogeneic hematopoietic cell transplantation for hematological disorders. BMC Med Genomics. 2019;12(1):49. Epub 2019/03/09. doi: 10.1186/s12920-019-0494-7. PubMed PMID: 30845942; PubMed Central PMCID: PMCPMC6404274.

28. Jenq RR, Taur Y, Devlin SM, Ponce DM, Goldberg JD, Ahr KF, et al. Intestinal Blautia Is Associated with Reduced Death from Graft-versus-Host Disease. Biol Blood Marrow Transplant. 2015;21(8):1373-83. Epub 2015/05/16. doi: 10.1016/j.bbmt.2015.04.016. PubMed PMID: 25977230; PubMed Central PMCID: PMCPMC4516127.

29. Weber D, Jenq RR, Peled JU, Taur Y, Hiergeist A, Koestler J, et al. Microbiota Disruption Induced by Early Use of Broad-Spectrum Antibiotics Is an Independent Risk Factor of Outcome after Allogeneic Stem Cell Transplantation. Biol Blood Marrow Transplant. 2017;23(5):845-52. Epub 2017/02/25. doi: 10.1016/j.bbmt.2017.02.006. PubMed PMID: 28232086; PubMed Central PMCID: PMCPMC5546237.

30. Simms-Waldrip TR, Sunkersett G, Coughlin LA, Savani MR, Arana C, Kim J, et al. Antibiotic-Induced Depletion of Anti-inflammatory Clostridia Is Associated with the Development

of Graft-versus-Host Disease in Pediatric Stem Cell Transplantation Patients. Biol Blood Marrow
Transplant. 2017;23(5):820-9. Epub 2017/02/14. doi: 10.1016/j.bbmt.2017.02.004. PubMed
PMID: 28192251.

31. Piper HG, Fan D, Coughlin LA, Ho EX, McDaniel MM, Channabasappa N, et al. Severe
Gut Microbiota Dysbiosis Is Associated With Poor Growth in Patients With Short Bowel
Syndrome. JPEN J Parenter Enteral Nutr. 2017;41(7):1202-12. Epub 2016/07/14. doi:
10.1177/0148607116658762. PubMed PMID: 27406942.

32. Melson J, Jakate S, Fung H, Arai S, Keshavarzian A. Crypt loss is a marker of clinical
severity of acute gastrointestinal graft-versus-host disease. Am J Hematol. 2007;82(10):881-6.
Epub 2007/06/16. doi: 10.1002/ajh.20976. PubMed PMID: 17570511.

33. Spencer GD, Shulman HM, Myerson D, Thomas ED, McDonald GB. Diffuse intestinal
ulceration after marrow transplantation: a clinicopathologic study of 13 patients. Hum Pathol.
1986;17(6):621-33. Epub 1986/06/01. doi: 10.1016/s0046-8177(86)80135-6. PubMed PMID:
3011641.

34. Glucksberg H, Storb R, Fefer A, Buckner CD, Neiman PE, Clift RA, et al. Clinical
manifestations of graft-versus-host disease in human recipients of marrow from HL-A-matched
sibling donors. Transplantation. 1974;18(4):295-304. Epub 1974/10/01. doi: 10.1097/00007890-
197410000-00001. PubMed PMID: 4153799.

35. Ponec RJ, Hackman RC, McDonald GB. Endoscopic and histologic diagnosis of intestinal
graft-versus-host disease after marrow transplantation. Gastrointest Endosc. 1999;49(5):612-21.
Epub 1999/05/06. doi: 10.1016/s0016-5107(99)70390-1. PubMed PMID: 10228260.

36. Sale GE, Shulman HM, McDonald GB, Thomas ED. Gastrointestinal graft-versus-host
disease in man. A clinicopathologic study of the rectal biopsy. Am J Surg Pathol. 1979;3(4):291-
9. Epub 1979/08/01. PubMed PMID: 44107.

37. Nakajima M, Arimatsu K, Kato T, Matsuda Y, Minagawa T, Takahashi N, et al. Oral
Administration of P. gingivalis Induces Dysbiosis of Gut Microbiota and Impaired Barrier Function
Leading to Dissemination of Enterobacteria to the Liver. PLoS One. 2015;10(7):e0134234. Epub
2015/07/29. doi: 10.1371/journal.pone.0134234. PubMed PMID: 26218067; PubMed Central
PMCID: PMCPMC4517782.

38. Flak MB, Colas RA, Munoz-Atienza E, Curtis MA, Dalli J, Pitzalis C. Inflammatory arthritis
disrupts gut resolution mechanisms, promoting barrier breakdown by Porphyromonas gingivalis.
JCI Insight. 2019;4(13). Epub 2019/07/12. doi: 10.1172/jci.insight.125191. PubMed PMID:
31292292; PubMed Central PMCID: PMCPMC6629160.

39. van der Lugt B, van Beek AA, Aalvink S, Meijer B, Sovran B, Vermeij WP, et al.
Akkermansia muciniphila ameliorates the age-related decline in colonic mucus thickness and
attenuates immune activation in accelerated aging Ercc1 (-/Delta7) mice. Immun Ageing.
2019;16:6. Epub 2019/03/23. doi: 10.1186/s12979-019-0145-z. PubMed PMID: 30899315;
PubMed Central PMCID: PMCPMC6408808.

40. Grander C, Adolph TE, Wieser V, Lowe P, Wrzosek L, Gyongyosi B, et al. Recovery of
ethanol-induced Akkermansia muciniphila depletion ameliorates alcoholic liver disease. Gut.
2018;67(5):891-901. Epub 2017/05/28. doi: 10.1136/gutjnl-2016-313432. PubMed PMID:
28550049.

41. Wu W, Lv L, Shi D, Ye J, Fang D, Guo F, et al. Protective Effect of Akkermansia
muciniphila against Immune-Mediated Liver Injury in a Mouse Model. Front Microbiol.
2017;8:1804. Epub 2017/10/17. doi: 10.3389/fmicb.2017.01804. PubMed PMID: 29033903;
PubMed Central PMCID: PMCPMC5626943.

42. Gesser RM, Koo SC. Oral inoculation with herpes simplex virus type 1 infects enteric
neuron and mucosal nerve fibers within the gastrointestinal tract in mice. Journal of virology.
1996;70(6):4097-102. PubMed PMID: 8648749.

43.   Basso L, Serhan N, Tauber M, Gaudenzio N. Peripheral neurons: Master regulators of
skin and mucosal immune response. European Journal of Immunology. 2019;0(0). doi:
10.1002/eji.201848027.
44.   Pavlov VA, Tracey KJ. Neural regulation of immunity: molecular mechanisms and clinical
translation. Nat Neurosci. 2017;20(2):156-66. doi: 10.1038/nn.4477.
45.   Lokensgard JR, Cheeran MC, Hu S, Gekker G, Peterson PK. Glial cell responses to
herpesvirus infections: role in defense and immunopathogenesis. J Infect Dis. 2002;186 Suppl
2:S171-9. doi: 10.1086/344272. PubMed PMID: 22311297.
46.   Marques CP, Cheeran MCJ, Palmquist JM, Hu S, Urban SL, Lokensgard JR. Prolonged
Microglial Cell Activation and Lymphocyte Infiltration following Experimental Herpes Encephalitis.
J Immunol. 2008;181(9):6417-26.
47.   Thion MS, Low D, Silvin A, Chen J, Grisel P, Schulte-Schrepping J, et al. Microbiome
Influences Prenatal and Adult Microglia in a Sex-Specific Manner. Cell. 2018;172:1-17. doi:
10.1016/j.cell.2017.11.042.
48.   Black ME, Hruby DE. Nucleotide sequence of the Escherichia coli thymidine kinase gene
provides evidence for conservation of functional domains and quaternary structure. Molecular
Microbiology. 1991;5(2):373-9. doi: 10.1111/j.1365-2958.1991.tb02119.x.
49.   Jeffrey WH, Paul JH. Thymidine uptake, thymidine incorporation, and thymidine kinase
activity in marine bacterium isolates. Applied and Environmental Microbiology. 1990;56(5):1367-
72.
50.   Konrad A, Yarunova E, Tinta T, Piškur J, Liberles DA. The global distribution and evolution
of deoxyribonucleoside kinases in bacteria. Gene. 2012;492(1):117-20. doi:
https://doi.org/10.1016/j.gene.2011.10.039.
51.   Lönnqvist B, Palmblad J, Ljungman P, Grimfors G, Järnmark M, Lerner R, et al. Oral
acyclovir as prophylaxis for bacterial infections during induction therapy for acute leukaemia in
adults. Support Care Cancer. 1993;1(3):139-44. doi: 10.1007/bf00366060.
52.   Brewin N, Cairns J. State of the DNA replication fork during thymine deprivation of
Escherichia coli, as observed by pulse-labelling with [3H]thymidine. Journal of Molecular Biology.
1977;111(3):353-63. doi: https://doi.org/10.1016/S0022-2836(77)80057-0.
53.   Bettegowda C, Foss CA, Cheong I, Wang Y, Diaz L, Agrawal N, et al. Imaging bacterial
infections with radiolabeled 1-(2′-deoxy-2′-fluoro-β-D-arabinofuranosyl)-5-iodouracil. Proceedings
of the National Academy of Sciences of the United States of America. 2005;102(4):1145-50. doi:
10.1073/pnas.0408861102.
54.   Davis SL, Be NA, Lamichhane G, Nimmagadda S, Pomper MG, Bishai WR, et al. Bacterial
Thymidine Kinase as a Non-Invasive Imaging Reporter for <italic>Mycobacterium
tuberculosis</italic> in Live Animals. PLoS ONE. 2009;4(7):e6297. doi:
10.1371/journal.pone.0006297.
55.   Peterson KL, Reid WC, Freeman AF, Holland SM, Pettigrew RI, Gharib AM, et al. The use
of 14C-FIAU to predict bacterial thymidine kinase presence: Implications for radiolabeled FIAU
bacterial imaging. Nuclear Medicine and Biology. 2013;40(5):638-42. doi:
http://dx.doi.org/10.1016/j.nucmedbio.2013.01.005.
56.   Wellsbury P, Herbert RA, John Parkes R. Incorporation of [methyl-3H]thymidine by
obligate and facultative anaerobic bacteria when grown under defined culture conditions. FEMS
Microbiology Ecology. 1993;12(2):87-95. doi: 10.1111/j.1574-6941.1993.tb00020.x.
57.   Jenq RR, Taur Y, Devlin SM, Ponce DM, Goldberg JD, Ahr KF, et al. Intestinal *Blautia* Is
Associated with Reduced Death from Graft-versus-Host Disease. Biology of Blood and Marrow
Transplantation. 2015;21(8):1373-83. doi: 10.1016/j.bbmt.2015.04.016.
58.   Stingl C, Cardinale K, Van Mater H. An Update on the Treatment of Pediatric Autoimmune
Encephalitis. Current Treatment Options in Rheumatology. 2018;4(1):14-28. doi:
10.1007/s40674-018-0089-z.

59. Armangue T, Spatola M, Vlagea A, Mattozzi S, Cárceles-Cordon M, Martinez-Heras E, et
al. Frequency, symptoms, risk factors, and outcomes of autoimmune encephalitis after herpes
simplex encephalitis: a prospective observational study and retrospective analysis. The Lancet
Neurology. 2018;17(9):760-72. doi: https://doi.org/10.1016/S1474-4422(18)30244-8.
60. Wekerle H. Brain Autoimmunity and Intestinal Microbiota: 100 Trillion Game Changers.
Trends in Immunology. 2017;38(7):483-97. doi: https://doi.org/10.1016/j.it.2017.03.008.
61. Iro MA, Sadarangani M, Absoud M, Chong WK, Clark CA, Easton A, et al. /mmuno*g*lobuli*N*
*i*n the *T*reatment of *E*ncephalitis (IgNiTE): protocol for a multicentre randomised controlled trial.
BMJ Open. 2016;6(11). doi: 10.1136/bmjopen-2016-012356.
62. Ye SL, Lei M, Jiang P, Liu FJ, Wang ZK, Cao HJ, et al. Demonstration of the IgG antibody
repertoire against the bacteria Escherichia coli in Chinese intravenous immunoglobulins. Journal
of Pharmaceutical and Biomedical Analysis. 2017;133:8-14. doi: 10.1016/j.jpba.2016.10.018.
PubMed PMID: WOS:000392894900002.
63. Uchimura Y, Fuhrer T, Li H, Lawson MA, Zimmermann M, Yilmaz B, et al. Antibodies Set
Boundaries Limiting Microbial Metabolite Penetration and the Resultant Mammalian Host
Response. Immunity. 2018;49(3):545-59.e5. doi: https://doi.org/10.1016/j.immuni.2018.08.004.
64. Schneider C, Smith DF, Cummings RD, Boligan KF, Hamilton RG, Bochner BS, et al. The
human IgG anti-carbohydrate repertoire exhibits a universal architecture and contains specificity
for microbial attachment sites. Science Translational Medicine. 2015;7(269):269ra1-ra1. doi:
10.1126/scitranslmed.3010524.
65. Zeng Melody Y, Cisalpino D, Varadarajan S, Hellman J, Warren HS, Cascalho M, et al.
Gut Microbiota-Induced Immunoglobulin G Controls Systemic Infection by Symbiotic Bacteria and
Pathogens. Immunity. 2016;44(1-12). doi: http://dx.doi.org/10.1016/j.immuni.2016.02.006.
66. Negm OH, MacKenzie B, Hamed MR, Ahmad OAJ, Shone CC, Humphreys DP, et al.
Protective antibodies against Clostridium difficile are present in intravenous immunoglobulin and
are retained in humans following its administration. Clinical & Experimental Immunology.
2017:n/a-n/a. doi: 10.1111/cei.12946.
67. Kakiuchi S, Tsuji M, Nishimura H, Yoshikawa T, Wang L, Takayama-Ito M, et al.
Association of the Emergence of Acyclovir-Resistant Herpes Simplex Virus Type 1 With
Prognosis in Hematopoietic Stem Cell Transplantation Patients. The Journal of Infectious
Diseases. 2017;215(6):865-73. doi: 10.1093/infdis/jix042.
68. Shen Y, Giardino Torchia ML, Lawson GW, Karp CL, Ashwell JD, Mazmanian SK. Outer
membrane vesicles of a human commensal mediate immune regulation and disease protection.
Cell Host Microbe. 2012;12(4):509-20. Epub 2012/09/25. doi: 10.1016/j.chom.2012.08.004.
PubMed PMID: 22999859; PubMed Central PMCID: PMCPMC3895402.
69. Round JL, Mazmanian SK. Inducible Foxp3+ regulatory T-cell development by a
commensal bacterium of the intestinal microbiota. Proc Natl Acad Sci U S A. 2010;107(27):12204-
9. Epub 2010/06/23. doi: 10.1073/pnas.0909122107. PubMed PMID: 20566854; PubMed Central
PMCID: PMCPMC2901479.
70. Mazmanian SK, Round JL, Kasper DL. A microbial symbiosis factor prevents intestinal
inflammatory disease. Nature. 2008;453(7195):620-5. Epub 2008/05/30. doi:
10.1038/nature07008. PubMed PMID: 18509436.
71. Mazmanian SK, Liu CH, Tzianabos AO, Kasper DL. An immunomodulatory molecule of
symbiotic bacteria directs maturation of the host immune system. Cell. 2005;122(1):107-18. Epub
2005/07/13. doi: 10.1016/j.cell.2005.05.007. PubMed PMID: 16009137.
72. Ramakrishna C, Kujawski M, Chu H, Li L, Mazmanian SK, Cantin EM. Bacteroides fragilis
polysaccharide A induces IL-10 secreting B and T cells that prevent viral encephalitis. Nat
Commun. 2019;10(1):2153. Epub 2019/05/16. doi: 10.1038/s41467-019-09884-6. PubMed
PMID: 31089128; PubMed Central PMCID: PMCPMC6517419.
73. Staffas A, Burgos da Silva M, Slingerland AE, Lazrak A, Bare CJ, Holman CD, et al.
Nutritional Support from the Intestinal Microbiota Improves Hematopoietic Reconstitution after

Bone Marrow Transplantation in Mice. Cell Host & Microbe. 2018;23(4):447-57. doi: 10.1016/j.chom.2018.03.002.

74.   Rosshart SP, Herz J, Vassallo BG, Hunter A, Wall MK, Badger JH, et al. Laboratory mice born to wild mice have natural microbiota and model human immune responses. Science. 2019;365(6452):eaaw4361. doi: 10.1126/science.aaw4361.

75.   Rosshart SP, Vassallo BG, Angeletti D, Hutchinson DS, Morgan AP, Takeda K, et al. Wild Mouse Gut Microbiota Promotes Host Fitness and Improves Disease Resistance. Cell. 2017;171(5):1015-28.e13
. doi: 10.1016/j.cell.2017.09.016.

76.   Viney M, Riley EM. The Immunology of Wild Rodents: Current Status and Future Prospects. Frontiers in Immunology. 2017;8(1481). doi: 10.3389/fimmu.2017.01481.

77.   Lundberg P, Ramakrishna C, Brown J, Tyszka JM, Hamamura M, Hinton DR, et al. The immune response to herpes simplex virus type 1 infection in susceptible mice is a major cause of central nervous system pathology resulting in fatal encephalitis. J Virol. 2008;82(14):7078-88. doi: 10.1128/JVI.00619-08. PubMed PMID: 18480436; PubMed Central PMCID: PMCPMC2446972.

78.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. Proc Natl Acad Sci U S A. 2011;108. doi: 10.1073/pnas.1000080107.

79.   Edgar RC. UPARSE: highly accurate OTU sequences from microbial amplicon reads. Nat Methods. 2013;10. doi: 10.1038/nmeth.2604.

80.   Edgar RC. UNOISE2: Improved error-correction for Illumina 16S and ITS amplicon reads. bioRxiv. 2016. doi: http://dx.doi.org/10.1101/081257doi:http://dx.doi.org/10.1101/081257.

81.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol. 1990;215(3):403-10. Epub 1990/10/05. doi: 10.1016/s0022-2836(05)80360-2. PubMed PMID: 2231712.

82.   Edgar RC. Edgar RC. 2016. SINTAX, a simple non-Bayesian taxonomy classifier for 16S and ITS sequences. bioRxiv. 2016. doi: :http://dx.doi.org/10.1101/074161doi:http://dx.doi.org/10.1101/074161.

83.   Cole JR, Wang Q, Fish JA, Chai B, McGarrell DM, Sun Y, et al. Ribosomal Database Project: data and tools for high throughput rRNA analysis. Nucleic Acids Res. 2014;42(Database issue):D633-42. Epub 2013/11/30. doi: 10.1093/nar/gkt1244. PubMed PMID: 24288368; PubMed Central PMCID: PMCPMC3965039.

84.   Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al. QIIME allows analysis of high-throughput community sequencing data. Nature methods. 2010;7(5):335-6.

85.   McCarthy DJ, Chen Y, Smyth GK. Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. Nucleic Acids Res. 2012;40(10):4288-97. Epub 2012/01/31. doi: 10.1093/nar/gks042. PubMed PMID: 22287627; PubMed Central PMCID: PMCPMC3378882.

86.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics. 2010;26(1):139-40. doi: 10.1093/bioinformatics/btp616.