Real-time measurement of E2:ERα transcriptional activity in live-cells.

Manuela Cipolletti[1], Stefano Leone[1], Stefania Bartoloni[1], Claudia Busonero[1], and Filippo Acconcia[1].

Department of Sciences, Section Biomedical Sciences and Technology, University Roma Tre,

Rome, Italy.

* Correspondence should be addressed to Prof Filippo Acconcia, filippo.acconcia@uniroma3.it at

the Department of Sciences, University Roma Tre, Viale Guglielmo Marconi, 446, I-00146, Rome,

Italy. Tel.: +39 0657336320; Fax: +39 0657336321.

Running head: Real-time live-measurement ERα activity.

Keywords:

- **17β-estradiol**

- **Estrogen Receptor**

- **Transcriptional Activity**

- **Real-time Biology**

- **Breast Cancer.**

Total Number of Figures: 7

Data availability statement: Research data not shared. However, the data sets used and/or

analyzed during the current study are available from the corresponding author on reasonable

request.

Contract grant sponsor: Associazione Italiana Ricerca sul Cancro AIRC, Fondo di finanziamento

per le attività base di ricerca (FFABR) to FA. The Grant of Excellence Departments, MIUR

(ARTICOLO 1, COMMI 314 – 337 LEGGE 232/2016) to Department of Science.

Abstract.

Kinetic analyses of diverse physiological processes have the potential to unveil new aspects of the molecular regulation of cell biology at temporal levels. 17β-estradiol (E2) regulates diverse physiological effects by binding to the estrogen receptor α (ERα), which primarily works as a transcription factor. Although many molecular details of the modulation of ERα transcriptional activity have been discovered including the impact of receptor plasma membrane localization and its relative E2-evoked signalling, the knowledge of real-time ERα transcriptional dynamics in living cells is lacking. Here, we report the generation of MCF-7 and HeLa cells stably expressing a modified luciferase under the control of an E2-sensitive promoter, which activity can be continuously monitored in living cells and show that E2 induces a linear increase in ERα transcriptional activity. Ligand-independent (*e.g.,* epidermal growth factor) receptor activation was also detected in a time-dependent manner. Kinetic profiles of ERα transcriptional activity measured in the presence of both receptor antagonists and inhibitors of ERα plasma membrane localization reveals a biphasic dynamic of receptor behaviour underlying novel aspects of receptor-regulated transcriptional effects. Finally, analysis of the rate of the dose-dependent E2 induction of ERα transcriptional activity demonstrates that low doses of E2 induce an effect identical to that determined by high concentrations of E2 as a function of the duration of hormone administration. Overall, we present the characterization of sensitive stable cell lines where to study the kinetic of E2 transcriptional signaling and to identify new aspects of ERα function in different physiological or pathophysiological conditions.

Introduction.

The sex hormone 17β-estradiol (E2) is a critical regulator of cell physiology as it controls the homeostasis of female and male reproductive and non-reproductive tissues and organs. The pleiotropic E2 actions depend on the activation of E2 signaling. E2 signaling is triggered by the activation the estrogen receptor α (ERα), which works as a ligand-induced transcription factor (Busonero et al., 2019).

Indeed, in the nucleus, the E2:ERα complex activates the transcription of the genes containing the estrogen response element (ERE) sequence in their promoters. Moreover, E2 can regulate the expression of non-ERE sequence containing genes by inducing the ERα interaction with other transcription factors (Ascenzi et al., 2006). In addition, the regulation of ERα transcriptional activity can be further triggered by other hormones (*e.g.,* epidermal growth factor – EGF) in the absence of E2 (Ascenzi et al., 2006). Activation of E2:ERα complex gene transcription requires receptor phosphorylation on the serine (S) residue 118 (Lannigan, 2003; Le Romancer et al., 2011). The S118 ERα phosphorylation is a result of the E2-dependent activation of kinase cascades (*e.g.,* PI3K/AKT; ERK/MAPK), which are triggered by the plasma membrane localized ERα (Acconcia et al., 2005a; La Rosa et al., 2012; Pedram et al., 2007). ERα plasma membrane localization occurs because the receptor is palmitoylated on the cysteine (C) residue 447 by specific palmitoyl-acyl-transferases (PAT) (Acconcia et al., 2005a; Adlanmerini et al., 2014; La Rosa et al., 2012; Pedram et al., 2012; Pedram et al., 2007; Sosa et al., 2019). The ERα plasma membrane localization is a pre-requisite for E2-induced ERα transcriptional activity (La Rosa et al., 2012).

In the last years, diverse kinetic analyses have been performed in live-cells to monitor real-time cellular responses induced by different extracellular stimuli. Measurement of growth rates of breast cancer cells treated with different hormones (*e.g.,* E2, progestins, androgens and corticosteroids) or chemicals (*e.g.,* environmental pollutants) revealed complex kinetic proliferation profiles that suggested novel mechanisms of action for each compound (Rotroff et al., 2013). Another method, which has been developed to study protein turnover, allowed to identify in a

quantitative time-dependent manner new kinetic mechanistic events required for protein

degradation induced by proteolysis targeting chimeras (PROTACs) (Riching et al., 2018).

Therefore, real-time live-cell assays for measuring different parameters of cellular biology

holds the potential for the identification of unrecognized biological phenomena underlying the

physiological effects of extracellular stimuli (*e.g.,* hormones).

Remarkably, to our knowledge, the kinetic evaluation of E2-induced ERα transcriptional

activity in real-time has never been reported. In turn, we decided to generate a cell line-based model

system to detect the ERα transcriptional activity in living cells. To this purpose, we took advantage

of the available technology for which the activity of a modified luciferase (nanoluciferase-PEST -

NLuc) can be continuously detected in cells loaded with a non-toxic cell-permeable substrate (Hall

et al., 2012) and developed MCF-7 and HeLa cell lines stably expressing an E2-responsive NLuc

reporter gene construct.

Here, we report the characterization of ERα transcriptional activity in live-cells by different

hormones (*i.e.,* E2 and EGF) as well as the real-time kinetic analysis of the impact of ERα plasma

membrane localization on the E2-induced ERα transcriptional activity.

Materials and Methods.

Cell culture and reagents.

17β-estradiol (E2), epidermal growth factor (EGF), 4OH-tamoxifen (Tam), DMEM (with and

without phenol red), fetal calf serum, charcoal stripped fetal calf serum (DCC) and the palmitoyl-

acyl-transferase (PAT) inhibitor 2-bromohexadecanoic acid (2-bromo-palmitate; 2-Br) [IC_{50} of ~4

μM] (Varner et al., 2003) were purchased from Sigma-Aldrich (St. Louis, MO). Bradford protein

assay kit as well as anti-mouse and anti-rabbit secondary antibodies were obtained from Bio-Rad

(Hercules, CA). Antibodies against ERα (F-10 mouse – for WB), pS2 (FL-84 rabbit) and cathepsin

D (H75 rabbit) were obtained from Santa Cruz Biotechnology (Santa Cruz, CA); anti-vinculin

.

antibody was from Sigma-Aldrich (St. Louis, MO). All other antibodies were purchased by Cell Signalling Technology Inc. (Beverly, MA, USA). Chemiluminescence reagent for Western blotting was obtained from BioRad Laboratories (Hercules, CA, USA). Nano-Glo® Endurazine™ was purchased from Promega (Promega, Madison, MA, USA). All the other products were from Sigma-Aldrich. Analytical- or reagent-grade products were used without further purification. The identities of all the used cell lines [*i.e.,* human breast carcinoma cells (MCF-7 and T47D-1) and human cervix carcinoma cells (HeLa)] were verified by STR analysis (BMR Genomics, Italy).

Plasmids and Cloning

The reporter plasmid 3xERE TATA, the pcDNA flag 3.1 C, the pcDNA flag-ERα, the pcDNA flag-ERα C447A, and the pcDNA flag-ERα S118A were previously described (La Rosa et al., 2012). In order to generate the reporter plasmid pGL2Basic Neo_NLucPest_3xERE TATA containing the Nanoluciferase-PEST (NLuc) gene under the control of the 3xERE TATA promoter, the NLuc gene was first excised by the pNL2.2 (Promega, Madison, MA, USA) using KpnI/BamHI sites and cloned into the pGL2Basic Neo 3xERE TATA (a generous gift of Dr Wilson) (Wilson et al., 2004) using KpnI/BamHI sites. The resulting plasmid was the pGL2Basic Neo_NLucPest. Next, the fragment containing the 3xERE TATA was excised by the pGL2Basic Neo 3xERE TATA cut KpnI/HindIII and cloned into the pGL2Basic Neo_NLucPest plasmid cut KpnI/HindIII to finally generate the pGL2Basic Neo_NLucPest_3xERE TATA (Fig. 1).

Generation of stable MCF-7 and Hela ERE-NLuc cell lines.

MCF-7 and HeLa cells were transfected with pGL2Basic Neo_NLucPest_3xERE TATA using lipofectamine 2000 (Thermofisher) reagent according to the manufacturer's instructions. Twenty-four hours after transfection medium was changed and the selection antibiotic was added. In particular, MCF-7 ERE-NLuc cells were generated by using G418 (500 μg/ml) while HeLa ERE-

NLuc cells were generated by using G418 (750 μg/ml). Pooled cloned were used for the

experiments. Selection antibiotic was left in growing medium while each experiment was

performed in the absence of antibiotic.

Growth Curves.

The xCELLigence DP system ACEA Biosciences, Inc. (San Diego, CA) Multi-E-Plate station

was used to measure the time-dependent response to E2 by real-time cell analysis (RTCA). Each

experimental condition was tested in quadruplicate. A detailed description of the instrument and the

relative software has been previously published (Rotroff et al., 2013). MCF-7, T47D-1 and MCF-7

ERE-NLuc cells were seeded in E-Plates 96 in growing medium. After overnight monitoring of

growth once every 15 min, medium was changed, and cells were grown in 1% DCC medium in the

presence or in the absence of E2 (1 nM) and remained in the medium until the end of the

experiment. Cellular responses were then recorded once every 15 min for a total time of 72 hours.

For comparison of the effect of E2 in each cell line, the ratio between the normalized cell

index (NCI) (obtained by the ACEA Biosciences software) of the mean value for the E2-treated

samples and the NCI of the mean value for control sample was calculated and shown as a function

of time.

Real-time measurement of NanoLucPest expression.

MCF-7 ERE-NLuc cells were seeded in 96 well plates (5000 cells/well). Twenty-four hours

after plating, medium was changed, and cells were grown in 1% DCC medium for 24 hours and

then stimulated with E2. Each experimental condition was plated in triplicate and 3 wells were

always treated with fulvestrant (ICI182,240) (Sigma Aldrich) in order to measure the basal ERα

transcriptional activity. HeLa ERE-NLuc cells were transfected with the indicated ERα encoding

plasmids or with vector control. Twenty-four hours after transfection cells were seeded in 96 well

plates (5000 cells/well) and subsequently treated as described for MCF-7 ERE-NLuc cells. Nano-

Glo® Endurazine^{TM} was added according to manufacturer's instruction in 50µl as final

experimental volume together with ligand and/or inhibitor administration. Plates were then

transferred into a Tecan Spark microplate reader (Switzerland) set to 37°C and 5% CO_2. Light

emission (released light units - RLU) was measured for 24 hours every other 5 minutes. For

calculations, each data point was subtracted of the RLU mean value of the 3 fulvestrant treated

samples at each time point. Next each value was subtracted with the value of the corresponding

sample at time 0. The mean value of each experimental condition was calculated and subsequently

the ratio between each experimental condition and the control condition was obtained. Each

experiment was done twice in duplicate.

Western Blotting Assays.

Before any cellular and biochemical assay, cells were grown in 1% DCC medium for 24

hours and then stimulated with E2 at the indicated time points and doses. Cells were lysed in YY

buffer (50 mM HEPES at pH 7.5, 10% glycerol, 150 mM NaCl, 1% Triton X-100, 1 mM EDTA, 1

mM EGTA) plus protease and phosphatase inhibitors. Western blotting analyses were performed by

loading 20-30 µg of protein on SDS-gels. Gels were run and transferred to nitrocellulose

membranes with Biorad Turbo-Blot semidry transfer apparatus. Immunoblotting was carried out by

incubating membranes with 5% milk (60 min), followed by incubation o.n. with the indicated

antibodies. Secondary antibody incubation was continued for an additional 60 min. Bands were

detected using a Biorad Chemidoc apparatus.

BrdU Incorporation.

Bromodeoxiuridine (BrdU) was added in the last 30 minutes to the medium and then cells

were fixed, permeabilized, and the histones were dissociated with 2 M HCl as previously described

(Darzynkiewicz and Juan, 2001). BrdU positive cells were detected with an anti-BrdU primary

antibody diluted 1:100 (DAKO Cytomatation) and with an anti-mouse-Alexa488 conjugated diluted

1:100 (Thermofisher). Both antibodies were incubated for 1 hour at R.T. in the dark. BrdU

fluorescence was measured using a CytoFlex flow cytometer and the cell cycle analysis was

performed by CytExpert v1.2 software (Beckman Coulter). All samples were counterstained with

propidium iodide (PI) for DNA/BrdU biparametric analysis.

Cell cycle analysis

After treatments, cells were harvested with trypsin, and counted to obtain 10^6 cells per

condition. Then, the cells were centrifuged at 1500 rpm for 5 min at 4°C, fixed with 1 ml ice-cold

70% ethanol and subsequently stained with PI buffer (500 μg/ml Propidium Iodide, 320μg/ml

RNaseA, in 0.1% Triton X in PBS). DNA fluorescence was measured using a CytoFlex flow

cytometer and the cell cycle analysis was performed by CytExpert v1.2 software (Beckman

Coulter).

Statistical analysis.

A statistical analysis was performed using the ANOVA (One-way analysis of variance and

Tukey's as post-test) test with the InStat version 3 software system (GraphPad Software Inc., San

Diego, CA). Densitometric analyses were performed using the freeware software Image J by

quantifying the band intensity of the protein of interest respect to the relative loading control band

(*i.e.,* vinculin) intensity. In all analyses, *p* values < 0.01 were considered significant but for Western

blotting experiments for which *p* values < 0.05 were considered significant.

Results.

Characterization of MCF-7 NLuc cells.

Characterization of the generated MCF-7 ERE-NLuc cells was performed by evaluating the

ability of E2 to induce cell proliferation (Castoria et al., 2001), S118 ERα phosphorylation (Ali et

al., 1993), ERα degradation (Leclercq et al., 2006) as well as the accumulation of two well-known

ERE-containing genes (*i.e.,* presenilin 2 – pS2/TFF and cathepsin D – CatD) (Sun et al., 2005).

Growth curves analyses indicated that E2 induced a persistent time-dependent increase in cell

number in MCF-7 ERE-NLuc cells (Fig. 2A-green line). Interestingly, E2 was also able to increase

the number of parental MCF-7 cells with the same kinetics (Fig. 2A-red line). As control, we also

measured the ability of E2 to increase the number of T47D-1, another ERα expressing breast cancer

cell line (Wilson et al., 2004). Figure 2A (purple line) shows that in T47D-1 cells E2 increased the

cell number in a time-dependent manner, although with a different kinetics. MCF-7 ERE-NLuc

cells were treated with E2 for 24 hours and cell cycle analysis was further performed. E2

augmented the percentage of the cells that incorporated BrdU (Fig. 2B) and increased the number of

cells in the S phase of the cell cycle (Fig. 2C) with respect to the control untreated cells. These data

demonstrate that E2 induces MCF-7 ERE-NLuc DNA synthesis, cell cycle progression and cell

proliferation.

E2-induced S118 phosphorylation of the ERα is required for receptor transcriptional activity

(Ali et al., 1993). Therefore, we tested if E2 could trigger this receptor post-translational

modification in MCF-7 ERE-NLuc cells. Time-course analyses revealed that E2 rapidly (5 min) and

persistently (2 hours) increased the fraction of the S118 phosphorylated ERα in MCF-7 ERE-NLuc

cells, which peaked after 30 min of E2 administration (Fig. 2D and D').

Because E2-induced ERα degradation is intrinsically connected with receptor transcriptional

activity (Metivier et al., 2003; Reid et al., 2003) and we noted that E2 also determined a rapid and

persistent reduction in ERα intracellular levels (Fig. 2D), we next evaluated the ability of E2 to

trigger ERα degradation. Figure 2E and 2E' show that in 24 hours E2 reduced ERα content in MCF-

7 ERE-NLuc cells in a dose-dependent manner.

Next, we finally tested the ability of E2 to modulate the expression of the ERE-containing

genes pS2/TFF and CatD. As shown in figure 3B and 3C, the levels of both pS2/TFF and CatD

were increased by 24 hours of E2 treatment in MCF-7 ERE-NLuc cells in a dose-dependent manner.

Overall, these data indicate that the MCF-7 ERE-NLuc cells respond to E2 as expected for an

E2 sensitive cell line derived from the parental MCF-7 cells.

Real-time measurement of E2-induced ERα transcriptional activity in living cells.

To measure the E2-induced ERα transcriptional activity in real-time and in live-cells, MCF-7

ERE-NLuc cells were loaded with the live-cell substrate Nano-Glo® Endurazine[TM] (Hall et al.,

2012) in the presence or in the absence of different doses of E2 (from 10^{-12} to 10^{-8}M) and ERE-

NLuc-dependent activity (*i.e.,* released light units - RLU) was measured for 24 hours every other 5

minutes in a 37°C and 5% CO_2 controlled atmosphere (for details, please see material and methods

section).

As shown in figure 3A, E2 induced a time-dependent increase in ERE-NLuc activity at all the

tested doses with a maximal effect occurring already at 10^{-10}M. Notably, cell treatment with 10^{-12}M

E2 was ineffective in inducing ERE-NLuc activity (Fig. 3A – purple line). On the contrary, E2-

induced activation of ERE-NLuc activity occurred with the same kinetic profile but it reached a

lower level of induction when 10^{-11}M E2 was administered to MCF-7 ERE-NLuc cells (Fig. 3A –

green line). To evaluate if such differences could be ascribed to a different rate of E2-induced

effect, we calculated the slope of the curves (*i.e.,* linear regression) obtained by cells treated with E2

in our time courses real-time analyses. Figure 3A' shows a dose-dependent trend in the slope of the

curves relative to E2 administration. In particular, no differences were observed among the slopes

of the curves that refers to the cells treated with doses of E2 ranging from 10^{-10} to 10^{-8}M while the

slope extracted from the 10^{-11}M E2-treated cells was significantly lower (Fig. 3A'). Notably, the

slope of the cells treated with 10^{-12}M E2 was identical to that of the control samples (Fig. 3A').

Next, in order to directly understand if the E2 ERE-NLuc activity correlates with the E2-

induced ERE-containing gene expression, we compared the E2 dose-dependent effect on the

measured ERE-NLuc activity at the single time point 24 hours (Fig. 3C – green line) with the dose-

dependent increase in pS2/TFF and CatD detected in MCF-7 ERE-NLuc cells after 24 hours of E2

administration (Fig. 3B and 3C). As shown in figure 3C, the E2-dependent increase in both ERE-NLuc activity, pS2/TFF and CatD expression was described by a sigmoidal curve typical of the E2 effect. Remarkably, the excitatory dose 50 (ED_{50}) calculated for ERE-NLuc activity was lower (ED_{50}= 5.0 10^{-12}M – 5 pM) than the one calculated for Western blotting analysis of pS2/TFF and CatD cellular levels (ED_{50}= 2.5 10^{-11}M – 250 pM) (Fig. 3C). Therefore, the ERE-NLuc activity assay is sensitive to E2 administration.

Next, we evaluated the involvement of ERα in the E2-induced ERE-NLuc activity by administering the ERα antagonist 4OH-tamoxifen (Tam). As shown in figure 4A, the time-dependent linear E2 induction of ERE-NLuc activity was strongly prevented by Tam administration. Interestingly, the kinetic profile of Tam-treated MCF-7 ERE-NLuc cells was biphasic. In a first phase, we obtained values below the control up to 650 min (*i.e.,* about 10.8 hours) for Tam+E2 treated samples and up to 700 min (*i.e.,* about 11.7 hours) for Tam-alone treated samples. In a second phase, instead, Tam administration triggered an increase in ERE-NLuc activity up to 1440 min (*i.e.,* 24 hours) both in the presence and in the absence of E2. Accordingly, the slope of the Tam curves obtained in MCF-7 ERE-NLuc cells were significantly lower than the E2 one but also significantly increased with respect to the control samples (Fig. 4A').

To understand if this Tam-dependent behavior was due to an artifactual measurement of ERE-NLuc activity in our stable MCF-7 cell lines, we again compared the E2 effect on the measured ERE-NLuc activity at the single time point 24 hours (Fig. 4C – green bars) with the E2-induced increase in pS2/TFF and CatD detected in MCF-7 ERE-NLuc cells after 24 hours of E2 administration (Fig. 4B and 4C). As shown in figure 4B and 4C, the E2-dependent increase in both ERE-NLuc activity, pS2/TFF and CatD expression was prevented by Tam administration. However, Tam alone increased the basal levels of CatD (Fig. 4B). Notably, a similar stimulatory effect of Tam on ERE-containing genes was already scored in different cell lines (Arao et al., 2011). As expected (Busonero et al., 2019; Leclercq et al., 2006), the E2-induced degradation of ERα was

prevented by Tam (Fig. 4B). These results show that the E2-induced ERE-NLuc activation depends

on ERα transcriptional activity.

Overall, these data demonstrate that E2 induces a rapid and persistent liner increase in ERα

transcriptional activity which can be detected at low doses of E2 (*i.e.,* between 10^{-12} to 10^{-11}M) and

can be prevented by antiestrogens (*e.g.,* Tam) and further suggest that the differences in the effects

elicited by E2 at different doses could depend on the rate at which E2 induces ERα transcriptional

activation.

Real-time measurement of E2-independent ERα transcriptional activity.

Next, we evaluated the ligand-independent ERα transcriptional activation in MCF-7 ERE-

NLuc cells. To this purpose we treated cells with epidermal growth factor (EGF), which is known to

induce receptor gene transcription in the absence of E2 (Ascenzi et al., 2006).

Because EGF-dependent ERα activation is weak and could be detected only under ERα

overexpression conditions (Berno et al., 2008; El-Tanani and Green, 1997), we included a negative

control in our experiments by also administering E2 at 10^{-13}M. As expected, E2 induced a time-

dependent linear increase in ERE-NLuc activity when cells were treated at hormone dose of 10^{-8}M

(black line) but not when 10^{-13}M E2 (orange line) was administered (Fig. 5A). Interestingly, EGF

treatment also induced a time-dependent stimulation of ERE-NLuc activity (Fig. 5A). Analysis of

both hormone effects on the measured ERE-NLuc activity at the time point 24 hours (Fig. 5B) and

the slope of hormone-triggered ERE-NLuc activity curves (Fig. 5C) further confirmed such

observations.

Therefore, in MCF-7 ERE-NLuc cells EGF triggers the activation of the ERα transcriptional

function.

Real-time measurement of the impact of ERα plasma membrane localization on E2-induced

ERα transcriptional activity.

Plasma membrane localization of the ERα occurs because the receptor is palmitoylated by specific palmityl-acyl-transferases (PATs) on the C residue 447. Inhibition of ERα palmitoylation by either the PAT inhibitor 2-bromo-palmitate (2-Br) or the ERα C447 to alanine (A) mutant (C447A) prevents receptor plasma membrane localization and E2 signaling including ERα transcriptional functions (Acconcia et al., 2005a; Adlanmerini et al., 2014; La Rosa et al., 2012; Pedram et al., 2012; Pedram et al., 2007; Sosa et al., 2019). Therefore, we analyzed the effect of E2 in MCF-7 ERE-NLuc cells in the presence of the PAT inhibitor 2-Br and further evaluated the E2-triggered ERE-NLuc activity of the C447A mutant in transfected HeLa ERE-NLuc cells. Because E2-induced ERα S118 phosphorylation is also required for full receptor transcriptional activity (Ali et al., 1993), we also tested the effects of E2 in ERα S118A mutant transfected HeLa ERE-NLuc cells.

As shown in figure 6A and 6B, 2-Br treatment of MCF-7 ERE-NLuc cells prevented the E2-induced effect on the ERE-NLuc activity and strongly reduced the slope of the curve derived by E2 administration. Similar results were also obtained in wild type (wt) and C447A ERα mutant transfected HeLa ERE-NLuc cells treated with E2 (Fig. 6D and 6E). Notably, as expected for a receptor defective in S118 phosphorylation (Ali et al., 1993), the E2 the kinetic profile as well as its relative slope was reduced in S118A ERα mutant transfected HeLa ERE-NLuc cells with respect to HeLa ERE-NLuc cells expressing the wt receptor (Fig. 6D and 6E).

Interestingly, we noticed a complex kinetic profile of 2Br-treated MCF-7 ERE-NLuc cells. 2-Br treatment reduced the ERE-NLuc activity below the control. Indeed, within the first 5 hours of treatment E2 was not able to trigger ERE-NLuc activation while from 5 to 24 hours of E2 administration, the hormone stimulated the ERE-NLuc reporter activity (Fig. 6C). On the contrary, the measured E2-dependent ERE-NLuc activation of the C447A ERα mutant in transfected HeLa

329   ERE-NLuc cells was constantly lower than the one detected in the presence of the wt receptor for

330   the entire duration of the analysis (Fig. 6F).

331   These results confirm that ERα palmitoylation is required for E2-induced ERα transcriptional

332   activity and further suggest that the E2-dpendent activation of ERα plasma membrane localized

333   receptor is necessary for both rapid and persistent activation of ERα transcriptional functions.

335   Prediction of dose- and time-dependent E2 effect on ERα transcriptional activity.

Finally, based on the slope of the transcriptional profile extracted by our E2 dose-dependent

analyses in MCF-7 ERE-NLuc cells (Fig. 2A and 2B), we reasoned that the results obtained from

our stable cell lines could allow to predict the time point at which different doses of E2 elicit the

same ERE-NLuc activity. In turn, we first calculated the time at which each dose of E2 determines

a specific amount of E2 effect (Fig. 7A). On this basis, we hypothesized that 24 hours E2

administration at 10^{-11}M would determine a transcriptional effect equal to those elicited by 18 hours

E2 administration at concentration ranging from 10^{-10}M at 10^{-8}M, as detected in ERE-NLuc assays

(Fig. 7B).

To verify this prediction, we treated both parental MCF-7 and MCF-7 ERE-NLuc cells with

E2 at 10^{-11}M for 24 hours and with E2 at both 10^{-10}M, 10^{-9}M and 10^{-8}M for 18 hours and measured

the pS2/TFF expression. As shown in figure 7C and 7C', E2 induced the same accumulation of

pS2/TFF intracellular levels at all the tested doses in both parental and artificial MCF-7 cell lines.

Remarkably, under these conditions E2 lost its dose-dependent effect (Fig. 3C) on the induction of

pS2/TFF expression (Fig. 7D).

These data demonstrate that prolonged treatments of MCF-7 cells with low doses of E2

induce an increase in ERα transcriptional activity identical to that elicited by E2 administered to

cells at high concentrations, thus suggesting a critical role for time-dependent effects rather than

concentration-dependent effects in the regulation of E2:ERα transcriptional activity.

Discussion.

The main aim of the present work was to identify a method allowing the measurement of ERα

transcriptional activity both in real-time and in living cells. Here, we constructed a novel plasmid

(*i.e.,* pGL2Basic Neo_NLucPest_3xERE TATA) containing neomycin resistance cassette and a

nanoluciferase-PEST (NLuc) gene under the control of 3 repetition of the classical estrogen

response element (ERE) (Fig. 1). The NLuc gene encodes for a modified luciferase gene engineered

to be fused in frame with a PEST sequence. This fusion protein returns a high brightness than the

classic luciferases and has a shorter protein half-life because of the PEST sequence. Additionally,

the substrate for NLuc protein is cell permeable and non-toxic. These biochemical features render

such reporter protein particularly suitable for live-cell transcriptional studies (Hall et al., 2012).

In turn, we generated MCF-7 and HeLa cells in which our novel plasmid was stably

transfected. MCF-7 cells are ERα expressing cells and are considered standard ''workhorses'' to

study E2-dependent ERα effects (Kao et al., 2009). On the other hand, HeLa cells do not express

ERα and therefore do not respond to E2 unless the receptor is exogenously introduced by transient

transfection (Acconcia et al., 2005b). The resulting MCF-7-ERE NLuc and HeLa-ERE NLuc stable

cell lines are responsive tools to study live-cell ERα transcriptional dynamics in real-time under

different experimental conditions.

Initial characterization of the stable MCF-7-ERE NLuc cells revealed that E2 induces the

proliferation of MCF-7-ERE NLuc cells with a kinetic identical to the one elicited by E2 in the

parental MCF-7 cells. Moreover, either MCF-7 cell lines are more sensitive to E2 than the T47D-1

cells, possibly because an higher level of ERα is present in MCF-7 cells than in T47D-1 cells (Kao

et al., 2009). Furthermore, T47D-1 cells but not MCF-7 cells also express the ERβ (Wilson et al.,

2004) which is known to mediate the antiproliferative effects of E2 (Acconcia et al., 2005b). As

expected for an MCF-7-derived cell lines (Dutertre and Smith, 2003; La Rosa et al., 2012;

Lannigan, 2003; Leone et al., 2018; Weitsman et al., 2006), we found that the MCF-7-ERE NLuc

cells respond to E2 by inducing DNA synthesis, cell cycle progression, the rapid and persistent

phosphorylation of the ERα on the S118 residue, the degradation of the ERα as well as the increase

in the levels of both pS2/TFF and CatD, two classic ERE-containing E2-target genes (Sun et al.,

2005). Therefore, we conclude that this novel cell line responds to E2 as the parental MCF-7 cells

both in terms of cell proliferation and in terms of intracellular molecular mechanisms of the E2:ERα

complex action.

The performance of the stable MCF-7-ERE NLuc cells was evaluated by measuring different

parameters of the ERα transcriptional activity. Real-time evaluation of E2 dose-dependent effect

indicated that E2 linearly induces ERα transcriptional activity as a function of time. Moreover, we

observed that the rate of E2-dependent induction of ERE-based transcription increase in a dose-

dependent manner. The excitatory dose 50 (ED_{50}) at a single time point (24 hours) following E2

exposure was approximately 5 pM. Notably, these results are similar to those obtained in T47D-1

cells stably transfected with different standard luciferase reporter genes, where the measured ED_{50}

for E2 at 24 hours was 6 pM (Legler et al., 1999) or 3 pM (Wilson et al., 2004). Interestingly, the

fact that the ED_{50} for E2 at 24 hours for pS2/TFF and CatD measured by means of Western blotting

analysis is 25 pM demonstrates that ERE-based assays are more sensitive to variations in ERα

transcriptional activity. Accordingly, although the ligand-independent transcriptional activation of

the ERα by EGF is weak and could be detected only in cells overexpressing the receptor (Berno et

al., 2008; El-Tanani and Green, 1997), we were able to profile the temporal dynamics of the EGF-

induced ERα transcriptional activity also in endogenously expressing ERα cells.

Time-dependent E2-induced ERα transcriptional activity was prevented by Tam in MCF-7-

ERE NLuc cells. Indeed, the rate of E2-transcriptional induction was strongly reduced in the

presence of Tam. However, analysis of the time-dependent transcriptional profile of Tam-treated

MCF-7-ERE NLuc cells identified a biphasic kinetic both in the presence and in the absence of E2.

Tam is the prototype selective estrogen receptor modulator (SERM). It works by binding to the ERα

and inhibiting its transcriptional activity through receptor structural modifications (Brzozowski et

al., 1997). In our real-time live-cell analyses we observed a first phase in which the ERα

transcriptional activity was reduced below the control. However, at later time points, ERα

transcriptional activity was stimulated by Tam although the E2 effect was completely abolished.

Accordingly, while pS2/TFF expression was completely prevented by Tam, CatD basal levels were

increased by this SERM, although E2 lost its ability to induce the expression of both genes.

Interestingly, the observed stimulatory activity of Tam on ERα transcriptional activity is consistent

with already reported observations (Arao et al., 2011). However, it is additionally possible to

speculate that basal Tam-dependent ERα activation could be ascribed to the ability of Tam to

induce an accumulation in ERα intracellular levels (Busonero et al., 2019; Leclercq et al., 2006).

The increased number of ERα molecules would compensate for the inhibitory effect of Tam but

only at later time points. On the contrary, the inhibition of receptor transcriptional activity is

immediate and consistent with the ability of Tam to bind to ERα and inducing an antagonist

receptor conformation in a time-frame compatible with those of ligand:receptor association

(Brzozowski et al., 1997).

Real-time analysis of the impact of ERα plasma membrane localization on E2-induced ERα

transcriptional activity showed that the rate of this process was strongly dampened in cells in which

ERα palmitoylation was prevented. More interestingly, the ERα transcriptional activity showed a

biphasic kinetic profile in MCF-7-ERE NLuc cells treated with the PAT inhibitor. Inhibition of ERα

palmitoylation results in an inhibition of the E2-induced extra-nuclear signaling which are required

for the activation of receptor transcriptional activity and for the completion of many different E2-

dependent physiological functions both in cell lines and *in vivo* (Acconcia et al., 2005a;

Adlanmerini et al., 2014; La Rosa et al., 2012; Pedram et al., 2012; Pedram et al., 2007; Sosa et al.,

2019). Our time-dependent live-cell analyses confirm that the contribution of the rapid signaling

pathways originating from the activation of the ERα at the plasma membrane is critical for full

receptor transcriptional activity. However, data presented here further demonstrate that besides

being a pre-requisite for the rapid initial stages of E2-induced ERα transcriptional activation (La

Rosa et al., 2012), the E2:ERα plasma membrane signaling is also absolutely required for the

prolonged effect of ERα as a ligand-induced transcription factor. In support to this notion, the time-

dependent kinetic profile of E2-induced transcriptional activity of the palmitoylation defective ERα

C447A mutant measured in HeLa-ERE NLuc cells was completely flat. Therefore, we conclude that

the plasma membrane localization of the ERα is necessary and sufficient for the induction of the

receptor transcriptional activity both for rapid and prolonged times of E2 administration.

Finally, we predicted the time-dependent transcriptional behavior of different doses of an ERα

ligand. Indeed, we have calculated the time required for different E2 concentrations to reach the

same transcriptional effects and effectively reported that treatments with different E2 concentrations

at different time points of both MCF-7-ERE NLuc and parental MCF-7 cells produced the same

increase in pS2/TFF expression. Physiological blood concentration of E2 fluctuates in healthy pre-

menopausal woman between picomolar and nanomolar concentrations. However, the peak of E2

concentration in blood lasts for a maximum time of 48-72 hours. This peak is however sufficient to

induce all the physiological uterine modifications (Zittermann et al., 2000). Our results suggest that

rather than the chronic exposure to E2, the time of E2 administration is critical to achieve the

maximal ERα transcriptional activity. In this respect, the pulsatile nature of E2 action under

physiological conditions would support such concept. However, future investigations are required

to establish if short E2 treatments could lead to fully and prolonged activation of ERα action.

In conclusion, we report a method to study the real-time kinetics of E2:ERα transcriptional

activity in live-cells using the generated MCF-7-ERE NLuc and HeLa-ERE NLuc stable cell lines.

Here, we only measured the classic parameters of the E2 signaling (*i.e.,* ligand-dependent and

independent functions; the impact of ERα antagonists and of receptor plasma membrane

localization) on ERα transcriptional activity in MCF-7-ERE NLuc or in HeLa-ERE NLuc cells.

Nonetheless, our stable cell lines are suitable for diverse kind of analyses. MCF-7-ERE NLuc cells

can be easily used for drug discovery by setting up high-throughput screenings for the identification

of novel ERα antagonists with specific kinetic profiles. In addition, these cell lines could facilitate

the analysis of the transcriptional effects elicited by known or unknown endocrine disruptors or

environmental contaminants that bind to the ERα and have a weak estrogenic activity. In this

respect, the use of a live-cell real-time assay could be an advantage in understanding complex dose-

dependent effects of such receptor ligands (*e.g.,* U-shaped curves) (Acconcia et al., 2016; Acconcia

et al., 2015; Marino et al., 2012). MCF-7-ERE NLuc cells are also in principle suitable for the

analysis of the physiological involvement of a specific protein on the regulation of E2:ERα

transcriptional activity (*e.g.,* single or high-throughput screening with siRNA oligonucleotides). On

the other hand, HeLa-ERE NLuc cells are transfectable with different kinds of ERα (or ERβ)

deletion or point mutants and can be used to connect ERα structural biochemical determinants with

receptor transcriptional activity.

Therefore, these stable cell lines represent new models for the real-time live-cell analysis of

the kinetic aspects of E2 signaling under physiological or pathophysiological conditions.

Author contribution statement.

M.C. performed most of the work (generation of stable cell lines; real-time live cell

experiments; most of Western blotting analyses); S.L. performed all cell cycle analyses. S.B. and

C.B. generated the expression vectors required for transfection and performed some of Western

blotting analyses. M.C., S.L., S.B. and C.B. to figure preparation and manuscript proof-reading.

F.A. designed the research and wrote the paper.

Acknowledgments.

The research leading to these results has received funding from AIRC under IG 2018 - ID.

21325 project – P.I. Acconcia Filippo. This study was also supported by grants from Ateneo Roma

Tre and Fondo di finanziamento per le attività base di ricerca (FFABR) to FA. The Grant of

Excellence Departments, MIUR (ARTICOLO 1, COMMI 314 – 337 LEGGE 232/2016) to

Department of Science, University Roma TRE is also gratefully acknowledged. The Authors

declare no conflict of interest.

Figure Captions.

Figure 1. The pGL2Basic Neo_NLucPest_3xERE TATA plasmid.

Schematic of the produced plasmid map used to generate the MCF-7 and HeLa ERE-NLuc stable

cell lines. Important genetic determinants and restriction enzymes are indicated.

Figure 2. E2 sensitivity of the MCF-7 ERE-NLuc stable cell lines.

(A) Growth curves analysis of MCF-7 (red line), MCF-7 ERE-NLuc (green line) and T47D-1

(purple line) cells treated with E2 (10^{-9} M) for 72 hours. Measurement of cell impedence has been

detected every 15 minutes with the xCelligence DP device. Graph shows the E2 effect calculated

for each cell line at each time point with respect to its relative control cells (grey line). The data are

the means of two different experiments in which each sample was measured in quadruplicate (for

details please see the material and method section). Bromodeoxyurdine (BrdU) incorporation (B)

and cell cycle profile (C) of MCF-7 ERE-NLuc cells treated with E2 (10^{-9} M) for 24 hours. Western

blotting (D) and relative densitometric analyses (D') of S118 phosphorylated (pS118) ERα and ERα

expression levels in MCF-7 ERE-NLuc cells treated with E2 at the indicated time points (10^{-9} M).

Western blotting (E) and relative densitometric analyses (E') of ERα expression levels in MCF-7

ERE-NLuc cells treated with E2 for 24 hours at the indicated doses. The loading control was done

by evaluating vinculin expression in the same filter. * indicates significant differences with respect

to the – sample. All experiments were performed in triplicates. Data are the mean ± standard

deviations with a p value < 0.05.

Figure 3. Kinetic analysis of E2 effect in MCF-7 ERE-NLuc stable cell lines.

(A) Profile and (B) relative linear regression (Slope) of ERE-NLuc activity detected in MCF-7

ERE-NLuc cells treated with the indicated doses of E2 in the presence of the live-cell substrate

Nano-Glo® Endurazine™. Released light units (RLU) was measured for 24 hours every other 5

minutes in a 37°C and 5% CO_2 controlled atmosphere. Graph shows the E2 effect calculated at each

time point with respect to its relative control sample. The data are the means of two different

experiments in which each sample was measured in triplicate (for details please see the material and

method section). (C, green line) ERE-NLuc activity detected in MCF-7 ERE-NLuc cells treated

with the indicated doses of E2 for 24 hours. Western blotting (B) and relative densitometric

analyses (C) of pS2/TFF (red line in C) and cathepsin D (CatD) (orange line in C) expression levels

in MCF-7 ERE-NLuc cells treated with E2 for 24 hours at the indicated doses. The loading control

was done by evaluating vinculin expression in the same filter. * indicates significant differences

with respect to the – sample; ° indicates significant differences with respect to the $10^{-10} – 10^{-8}$ M

E2-treated samples. Data are the mean ± standard deviations with a p value < 0.01.

Figure 4. The role of 4OH-tamoxifen on the kinetic analysis of E2 effect in MCF-7 ERE-NLuc

stable cell lines.

(A) Profile and (A') relative linear regression (Slope) of ERE-NLuc activity detected in MCF-7

ERE-NLuc cells treated with E2 (10^{-8} M) and the live-cell substrate Nano-Glo® Endurazine™ in

the presence or in the absence of 4OH-tamoxifen (Tam - 10^{-7} M). Released light units (RLU) was

measured for 24 hours every other 5 minutes in a 37°C and 5% CO_2 controlled atmosphere. Graph

shows the E2 effect calculated at each time point with respect to its relative control sample. The

data are the means of two different experiments in which each sample was measured in triplicate

(for details please see the material and method section). (B, green bars) ERE-NLuc activity detected

in MCF-7 ERE-NLuc cells treated with E2 (10^{-8} M) in the presence or in the absence of 4OH-

tamoxifen (Tam - 10^{-7} M) for 24 hours. Western blotting (B) and relative densitometric analyses (C)

of pS2/TFF (red bars in C), cathepsin D (CatD) (orange bars in C) and ERα expression levels in

MCF-7 ERE-NLuc cells treated with E2 (10^{-8} M) in the presence or in the absence of 4OH-

tamoxifen (Tam - 10^{-7} M) for 24 hours. The loading control was done by evaluating vinculin

expression in the same filter. * indicates significant differences with respect to the – sample; °

indicates significant differences with respect to the E2-treated samples. # indicates significant

differences with respect to the Tam-treated samples. Data are the mean ± standard deviations with a

p value < 0.01 for panel (A') and $p < 0.05$ for panel (C).

Figure 5. The kinetic analysis of EGF effect in MCF-7 ERE-NLuc stable cell lines.

(A) Profile and (C) relative linear regression (Slope) of ERE-NLuc activity detected in MCF-7

ERE-NLuc cells treated with EGF (10 ng/ml) or E2 at the indicated doses and the live-cell substrate

Nano-Glo® Endurazine[TM]. Released light units (RLU) was measured for 24 hours every other 5

minutes in a 37°C and 5% CO_2 controlled atmosphere. Graph shows the E2 effect calculated at each

time point with respect to its relative control sample. The data are the means of two different

experiments in which each sample was measured in triplicate (for details please see the material and

method section). (B) ERE-NLuc activity detected in MCF-7 ERE-NLuc cells treated with EGF (10

ng/ml) or E2 at the indicated doses for 24 hours. * indicates significant differences with respect to

the – sample. Data are the mean ± standard deviations with a p value < 0.01.

Figure 6. The role of ERα palmitoylation on the kinetic analysis of E2 effect in MCF-7 and

HeLa ERE-NLuc stable cell lines.

Profile and relative linear regression (Slope) of ERE-NLuc activity detected in the presence of the

live-cell substrate Nano-Glo® Endurazine[TM] either MCF-7 ERE-NLuc cells treated with E2 (10^{-8}

M) in the presence or in the absence of 2-bromopalmitate (2-Br - 10^{-5} M) (A, B) or in HeLa ERE-

NLuc cells transfected with the pcDNA flag-ERα wild type (wt), the pcDNA flag-ERα C447A and

the pcDNA flag-ERα S118A and treated with E2 (10^{-8} M) (D, E). Released light units (RLU) was

measured for 24 hours (for MCF-7 ERE-NLuc) and for 12 hours (for HeLa ERE-NLuc) every other

5 minutes in a 37°C and 5% CO_2 controlled atmosphere. Graph shows the E2 effect calculated at

each time point with respect to its relative control sample. The data are the means of two different

experiments in which each sample was measured in triplicate (for details please see the material and

method section). (C) ERE-NLuc activity detected in MCF-7 ERE-NLuc cells treated with E2 (10^{-8}

M) in the presence or in the absence of 2-bromo-palmitate (2-Br - 10^{-5} M) or (F) in HeLa ERE-

NLuc cells transfected with the pcDNA flag-ERα wild type (wt), the pcDNA flag-ERα C447A and

the pcDNA flag-ERα S118A and treated with E2 (10^{-8} M) at the indicated time points. * indicates

significant differences with respect to the – sample; ° indicates significant differences with respect

to the E2-treated samples (for MCF-7 ERE-NLuc cells) or to the E2-treated samples in wt ERα (for

HeLa ERE-NLuc cells). # indicates significant differences with respect to the 2Br-treated samples

(for MCF-7 ERE-NLuc cells) or to the – samples in C447A ERα mutant samples (for HeLa ERE-

NLuc cells). Data are the mean ± standard deviations with a p value < 0.01.

Figure 7. The role of E2 doses and time of E2 administration on ERα transcriptional activity

in MCF-7 cells.

(A) Time required for E2 dose-dependent induction of ERE-NLuc activity in MCF-7 ERE-NLuc

cells. (B) ERE-NLuc activity detected in MCF-7 ERE-NLuc cells treated with E2 at the indicated

doses for the indicated times. Western blotting (C, C') and relative densitometric analyses (D) of

pS2/TFF expression levels in MCF-7 (C, red line in D) and in MCF-7 ERE-NLuc cells (C', green

line in D) treated with E2 at the indicated doses for the indicated times. The loading control was

done by evaluating vinculin expression in the same filter. * indicates significant differences with

respect to the – sample. All experiments were performed in triplicates. Data are the mean ± standard

deviations with a p value < 0.01 for panel (B) and p < 0.05 for panel (D).

References.

Acconcia F, Ascenzi P, Bocedi A, Spisni E, Tomasi V, Trentalance A, Visca P, Marino M. 2005a.

Palmitoylation-dependent estrogen receptor alpha membrane localization: Regulation by 17

beta-estradiol. Molecular Biology of the Cell 16(1):231-237.

Acconcia F, Fiocchetti M, Marino M. 2016. Xenoestrogen regulation of ERalpha/ERbeta balance in

hormone-associated cancers. Mol Cell Endocrinol.

Acconcia F, Pallottini V, Marino M. 2015. Molecular Mechanisms of Action of BPA. Dose-

response : a publication of International Hormesis Society 13(4):1559325815610582.

Acconcia F, Totta P, Ogawa S, Cardillo I, Inoue S, Leone S, Trentalance A, Muramatsu M, Marino

M. 2005b. Survival versus apoptotic 17beta-estradiol effect: role of ER alpha and ER beta

activated non-genomic signaling. J Cell Physiol 203(1):193-201.

Adlanmerini M, Solinhac R, Abot A, Fabre A, Raymond-Letron I, Guihot AL, Boudou F, Sautier L,

Vessieres E, Kim SH, Liere P, Fontaine C, Krust A, Chambon P, Katzenellenbogen JA,

Gourdy P, Shaul PW, Henrion D, Arnal JF, Lenfant F. 2014. Mutation of the palmitoylation

site of estrogen receptor alpha in vivo reveals tissue-specific roles for membrane versus

nuclear actions. Proc Natl Acad Sci U S A 111(2):E283-290.

Ali S, Metzger D, Bornert JM, Chambon P. 1993. Modulation of transcriptional activation by

ligand-dependent phosphorylation of the human oestrogen receptor A/B region. EMBO J

12(3):1153-1160.

Arao Y, Hamilton KJ, Ray MK, Scott G, Mishina Y, Korach KS. 2011. Estrogen receptor alpha

AF-2 mutation results in antagonist reversal and reveals tissue selective function of estrogen

receptor modulators. Proc Natl Acad Sci U S A 108(36):14986-14991.

Ascenzi P, Bocedi A, Marino M. 2006. Structure-function relationship of estrogen receptor alpha

and beta: impact on human health. Mol Aspects Med 27(4):299-402.

Berno V, Amazit L, Hinojos C, Zhong J, Mancini MG, Sharp ZD, Mancini MA. 2008. Activation

of estrogen receptor-alpha by E2 or EGF induces temporally distinct patterns of large-scale

chromatin modification and mRNA transcription. PLoS One 3(5):e2286.

Brzozowski AM, Pike AC, Dauter Z, Hubbard RE, Bonn T, Engstrom O, Ohman L, Greene GL,

Gustafsson JA, Carlquist M. 1997. Molecular basis of agonism and antagonism in the

oestrogen receptor. Nature 389(6652):753-758.

Busonero C, Leone S, Bartoloni S, Acconcia F. 2019. Strategies to degrade estrogen receptor alpha

in primary and ESR1 mutant-expressing metastatic breast cancer. Mol Cell Endocrinol

480:107-121.

Castoria G, Migliaccio A, Bilancio A, Di Domenico M, de Falco A, Lombardi M, Fiorentino R,

Varricchio L, Barone MV, Auricchio F. 2001. PI3-kinase in concert with Src promotes the

S-phase entry of oestradiol-stimulated MCF-7 cells. EMBO J 20(21):6050-6059.

Darzynkiewicz Z, Juan G. 2001. Analysis of DNA content and BrdU incorporation. Current

protocols in cytometry Chapter 7:Unit 7 7.

Dutertre M, Smith CL. 2003. Ligand-independent interactions of p160/steroid receptor coactivators

and CREB-binding protein (CBP) with estrogen receptor-alpha: regulation by

phosphorylation sites in the A/B region depends on other receptor domains. Mol Endocrinol

17(7):1296-1314.

El-Tanani MK, Green CD. 1997. Two separate mechanisms for ligand-independent activation of the

estrogen receptor. Mol Endocrinol 11(7):928-937.

Hall MP, Unch J, Binkowski BF, Valley MP, Butler BL, Wood MG, Otto P, Zimmerman K,

Vidugiris G, Machleidt T, Robers MB, Benink HA, Eggers CT, Slater MR, Meisenheimer

PL, Klaubert DH, Fan F, Encell LP, Wood KV. 2012. Engineered luciferase reporter from a

deep sea shrimp utilizing a novel imidazopyrazinone substrate. ACS chemical biology

7(11):1848-1857.

Kao J, Salari K, Bocanegra M, Choi YL, Girard L, Gandhi J, Kwei KA, Hernandez-Boussard T,

Wang P, Gazdar AF, Minna JD, Pollack JR. 2009. Molecular profiling of breast cancer cell

lines defines relevant tumor models and provides a resource for cancer gene discovery.

PLoS One 4(7):e6146.

La Rosa P, Pesiri V, Leclercq G, Marino M, Acconcia F. 2012. Palmitoylation Regulates 17beta-
Estradiol-Induced Estrogen Receptor-alpha Degradation and Transcriptional Activity. Mol
Endocrinol 26(5):762-774.

Lannigan DA. 2003. Estrogen receptor phosphorylation. Steroids 68(1):1-9.

Le Romancer M, Poulard C, Cohen P, Sentis S, Renoir JM, Corbo L. 2011. Cracking the Estrogen
Receptor's Posttranslational Code in Breast Tumors. Endocr Rev 32(5):597-622.

Leclercq G, Lacroix M, Laios I, Laurent G. 2006. Estrogen receptor alpha: impact of ligands on
intracellular shuttling and turnover rate in breast cancer cells. Curr Cancer Drug Targets
6(1):39-64.

Legler J, van den Brink CE, Brouwer A, Murk AJ, van der Saag PT, Vethaak AD, van der Burg B.
1999. Development of a stably transfected estrogen receptor-mediated luciferase reporter
gene assay in the human T47D breast cancer cell line. Toxicol Sci 48(1):55-66.

Leone S, Busonero C, Acconcia F. 2018. A high throughput method to study the physiology of
E2:ERalpha signaling in breast cancer cells. J Cell Physiol 233(5):3713-3722.

Marino M, Pellegrini M, La Rosa P, Acconcia F. 2012. Susceptibility of estrogen receptor rapid
responses to xenoestrogens: Physiological outcomes. Steroids 77(10):910-917.

Metivier R, Penot G, Hubner MR, Reid G, Brand H, Kos M, Gannon F. 2003. Estrogen receptor-
alpha directs ordered, cyclical, and combinatorial recruitment of cofactors on a natural target
promoter. Cell 115(6):751-763.

Pedram A, Razandi M, Deschenes RJ, Levin ER. 2012. DHHC-7 and -21 are
palmitoylacyltransferases for sex steroid receptors. Mol Biol Cell 23(1):188-199.

Pedram A, Razandi M, Sainson RC, Kim JK, Hughes CC, Levin ER. 2007. A conserved
mechanism for steroid receptor translocation to the plasma membrane. J Biol Chem
282(31):22278-22288.

Reid G, Hubner MR, Metivier R, Brand H, Denger S, Manu D, Beaudouin J, Ellenberg J, Gannon F.

2003. Cyclic, proteasome-mediated turnover of unliganded and liganded ERalpha on

responsive promoters is an integral feature of estrogen signaling. Mol Cell 11(3):695-707.

Riching KM, Mahan S, Corona CR, McDougall M, Vasta JD, Robers MB, Urh M, Daniels DL.

2018. Quantitative Live-Cell Kinetic Degradation and Mechanistic Profiling of PROTAC

Mode of Action. ACS chemical biology 13(9):2758-2770.

Rotroff DM, Dix DJ, Houck KA, Kavlock RJ, Knudsen TB, Martin MT, Reif DM, Richard AM,

Sipes NS, Abassi YA, Jin C, Stampfl M, Judson RS. 2013. Real-time growth kinetics

measuring hormone mimicry for ToxCast chemicals in T-47D human ductal carcinoma cells.

Chemical research in toxicology 26(7):1097-1107.

Sosa LDV, Petiti JP, Picech F, Chumpen S, Nicola JP, Perez P, De Paul A, Valdez-Taubas J,

Gutierrez S, Torres AI. 2019. The ERalpha membrane pool modulates the proliferation of

pituitary tumours. J Endocrinol 240(2):229-241.

Sun JM, Spencer VA, Li L, Yu Chen H, Yu J, Davie JR. 2005. Estrogen regulation of trefoil factor

1 expression by estrogen receptor alpha and Sp proteins. Exp Cell Res 302(1):96-107.

Varner AS, Ducker CE, Xia Z, Zhuang Y, De Vos ML, Smith CD. 2003. Characterization of human

palmitoyl-acyl transferase activity using peptides that mimic distinct palmitoylation motifs.

Biochem J 373(Pt 1):91-99.

Weitsman GE, Li L, Skliris GP, Davie JR, Ung K, Niu Y, Curtis-Snell L, Tomes L, Watson PH,

Murphy LC. 2006. Estrogen receptor-alpha phosphorylated at Ser118 is present at the

promoters of estrogen-regulated genes and is not altered due to HER-2 overexpression.

Cancer Res 66(20):10162-10170.

Wilson VS, Bobseine K, Gray LE, Jr. 2004. Development and characterization of a cell line that

stably expresses an estrogen-responsive luciferase reporter for the detection of estrogen

receptor agonist and antagonists. Toxicol Sci 81(1):69-77.

Zittermann A, Schwarz I, Scheld K, Sudhop T, Berthold HK, von Bergmann K, van der Ven H,

Stehle P. 2000. Physiologic fluctuations of serum estradiol levels influence biochemical

markers of bone resorption in young women. J Clin Endocrinol Metab 85(1):95-101.

Figure 1

Figure 2

A

B

C

D

D'

E

E'

Figure 3

A

A'

B

C

Figure 4

A

A'

B

C

Figure 5

A

B

C

Figure 6

A

D

B

E

C

F

Figure 7

A

B

C

MCF-7

C'

MCF-7 NLuc

D