Unmasking functional redundancy of deubiquitylases enables specificity profiling and discovery of novel proteasome substrates

Valentina Rossio[1], Joao A. Paulo[1], Joel Chick[1], Bradley Brasher[2], Steven P. Gygi[1], Randall W. King[1*]

[1] Department of Cell Biology, Blavatnik Institute at Harvard Medical School, Boston, MA 02115, USA.

[2] Boston Biochem, a Bio-Techne Brand 840 Memorial Drive, Cambridge, MA 02139, USA.

*Correspondence to Randall W. King: randy_king@hms.harvard.edu

Abstract

Deubiquitylating enzymes (DUBs) counteract ubiquitylation to control the stability or activity of

their substrates. Identifying DUB substrates is challenging and genetic approaches can be

thwarted by redundant action of DUBs. Here, we circumvented redundancy by broadly inhibiting

DUBs in *Xenopus* egg extract and used quantitative mass spectrometry to identify over thirty

proteins that undergo proteasomal degradation, the majority of which have not been reported as

DUB substrates. These results were confirmed with recombinant human proteins, demonstrating

the conservation of their DUB-dependent stability. We used these substrates to profile the ability

of a panel of DUBs to rescue degradation. This approach revealed that USP7, uniquely among

the 14 DUBs tested, has a broad ability to rescue degradation. USP21, which is used widely to

nonspecifically deubiquitylate proteins in vitro, was unable to rescue degradation, highlighting

the importance of profiling enzyme activity in a physiological system. Together, we identify new

DUB substrates and present a system to characterize physiological DUB specificity, overcoming

the challenges posed by DUB redundancy.

Introduction

The covalent addition of ubiquitin to proteins is a widespread, highly regulated post-translational modification. The Ubiquitin-Proteasome System (UPS) is best known for its ability to target specific proteins for degradation, but ubiquitylation also regulates protein localization, activity and function independently of degradation[1,2,3]. Ubiquitin contains seven lysine residues, each of which can be conjugated further with additional ubiquitin moieties to form polyubiquitin chains of different topologies that may result in different fates[4]. For example, lysine-63 linked chains promote non-degradative functions such as DNA repair or endocytosis[5,6], while lysine-48 linked chains and branched chains such as K11/K48 or K29/K48 promote degradation by the proteasome[7], a vital process that contributes to cellular homeostasis.

Ubiquitin is covalently attached to substrates via a cascade of E1-E2-E3 enzymes[8]. In contrast, deubiquitylating enzymes (DUBs) catalyze ubiquitin removal[9]. As such, a substrate's ubiquitylation status is a product of a carefully tuned balance between ubiquitylation and deubiquitylation activities. By counteracting substrate ubiquitylation, DUBs can rescue proteins targeted to proteasomal destruction. The human genome encodes approximately 100 DUBs, divided into two families: the zinc metalloprotease class (10 DUBs), and the cysteine-protease class that contains most other DUBs (~90)[10,11]. A small number of DUBs have been well-studied, with many characterized substrates, but most DUBs still do not have any known substrates, limiting our understanding of how DUB substrate specificity is achieved. DUB specificity is complex, as it can arise through binding specificity mediated by unique protein-protein interaction domains[12], or through recognition of specific topologies of ubiquitin chains[13,14,15]. Most DUB substrates have been discovered by first identifying proteins that interact with

specific DUBs. This approach has been used globally[16], but it may miss substrates that interact

weakly with DUBs or identify regulators or scaffold proteins rather than true substrates.

Identifying physiological UPS substrates, such as DUB targets, is challenging due to the

prominence of protein-quality control pathways that ubiquitylate newly synthesized proteins that

do not fold properly in growing cells[17]. Therefore, it can be difficult to discern which

ubiquitylation or degradation events are physiologically relevant, and which are simply a result

of misfolding during protein biogenesis. In addition, because DUBs may function

redundantly[18,19], inactivating a single DUB may not destabilize its substrates or affect a particular

biological process. Consistent with the presence of functional redundancy between DUBs, only

the metallo-protease RPN11 is essential for viability in yeast[20]. In the case of proteome stability,

it is not known which DUBs have the broadest impact in rescuing proteins from proteasome

degradation or to what extent DUBs act redundantly in this process. Identifying the DUBs that

have a major role in counteracting proteasomal degradation is of therapeutic interest, because the

UPS targets for destruction misfolded, damaged proteins and proteins involved in a wide range

of diseases[21]. Inhibiting DUBs that target these proteins may accelerate their elimination to

provide a therapeutic benefit.

Here, we first took an unbiased quantitative proteomic approach and discovered new

substrates protected from proteasomal degradation by DUBs, using the *Xenopus* egg extract

model system. By broadly inhibiting cysteine-protease DUBs, we circumvented the possible

effects of their redundancy on proteome stability. Because transcription and translation are not

active in this system, protein instability caused by DUB inhibition is not a consequence of

protein quality control ubiquitination activity. We recapitulated protein destabilization with

expression of recombinant human orthologs in extract, demonstrating that DUB-dependent

protein stability is evolutionarily conserved. Next, we took advantage of these newly identified

DUB substrates and tested the ability of a set of DUBs to rescue their degradation, uncovering a

broad ability of USP7 to rescue protein instability in extract. However, specific inhibition of

USP7 with a small molecule inhibitor was not sufficient to promote degradation of most of the

substrates we identified, suggesting that USP7 functions redundantly with other DUBs in this

system. Our work highlights the impact of DUB redundancy on proteome stability and reveals

the specificity and activity of DUBs whose function would otherwise be masked by redundancy.

Results

UbVS treatment induces rapid depletion of available ubiquitin and labels a broad set of

cysteine-protease DUBs in *Xenopus* extract

Our laboratory previously showed that the broad cysteine-protease DUB inhibitor UbVS

induces depletion of available ubiquitin in *Xenopus* extract[22]. In particular, UbVS-treatment

inhibited proteasome degradation of cyclin B, which could be efficiently rescued by addition of

50 μM exogenous ubiquitin[22]. We confirmed that 10 μM UbVS was sufficient to rapidly deplete

free ubiquitin, (Fig. 1a, 1b) which was associated with discharge of the Ube2C thioester (E2)

within 5 minutes (Fig. 1a, bottom). Addition of exogenous ubiquitin to UbVS-treated extract

fully recharged the Ube2C thioester, confirming that the E2's discharge was due to the depletion

of free ubiquitin (Fig.1a, bottom). To determine the spectrum of cysteine-containing DUBs

targeted by this concentration of UbVS, extract was treated with 10 μM HA-tagged UbVS (HA-

UbVS) and sensitive DUBs were visualized as discrete bands by anti-HA Western blotting

analysis (Fig. 1b, right). To identify these DUBs, we performed mass spectrometry analysis of

immunopurified HA-UbVS from extract and identified 88 proteins (Table S2), 35 of which

were cysteine-protease DUBs (Table S1, S2) that are likely direct targets of UbVS. All known

classes of cysteine-protease DUBs were found (Fig. 1c, Table S1), whereas metallo-protease

DUBs (JAMM) were not identified, as expected[23]. Prior proteomic analysis of *Xenopus* extract

detected 54 DUBs[24] (Table S1), 51 of which were cysteine-proteases. Thus, 10 μM HA-UbVS

labeled 35 of 51 (69%) of cysteine-protease DUBs present in extract, consistent with broad

specificity of UbVS for this class of DUBs[23]. The remaining DUBs were likely not either

because they were present in low abundance or because they do not react rapidly with UbVS.

Our HA-UbVS pull-down also isolated proteins that are likely to associate very tightly with

DUBs (Table S2), due to us carrying out the pull-down in presence of high salt (500 mM). A

large number of these proteins (31) are components of the proteasome, consistent with the

known interaction of UbVS-sensitive DUBs UCHL5 and USP14[25] with the proteasome. Other

known interactors of DUBs that we isolated included GBP1/GBP2, which binds USP10[26], and

the SPATA proteins (SPATA2/SPATA2L), which have been identified as CYLD interactors[16] in

human cells (Table S2), suggesting evolutionary conservation of these interactions.

UbVS treatment induces degradation of a limited set of proteins in *Xenopus* extract when

available ubiquitin is restored

We first wanted to investigate how inhibition of multiple DUBs influenced stability of

the proteome in *Xenopus* extract. We hypothesized that simultaneous inhibition of 35 DUBs with

UbVS might lead to destabilization of proteins that require ongoing deubiquitylation to maintain

their stability. By simultaneously inhibiting a large number of DUBs, we predicted that we might

be able to identify new substrates of DUBs, including substrates whose stability is conferred by

the action of redundant DUBs. We used a multiplexed Tandem Mass Tag (TMT)-based

quantitative proteomics approach[27] (Fig. 1d) and we measured protein abundance over time in

untreated extract, as well as extract to which ubiquitin, UbVS or the combination

(UbVS/ubiquitin) were added (Fig. 1e). Proteins decreasing specifically in the presence of UbVS

and/or UbVS/ubiquitin may normally be protected from degradation by UbVS-sensitive DUBs.

We performed 2 independent experiments using *Xenopus* extract prepared from 2 sets of eggs

(collected from different animals and processed separately), to ensure reproducibility. In each

experiment we quantified around 8000 proteins, with an overlap of 94.8% between the 2

experiments.

First, we observed that in the absence of any perturbation, the proteome was stable in

Xenopus extract over the course of 60 minutes (Fig. S1a). Few proteins changed in abundance

over time and none reproducibly in both the experiments. This degree of proteome stability is

consistent with the idea that bulk rates of translation are relatively quiescent in the *Xenopus*

system[28] and that the extract was treated with cycloheximide to prevent protein synthesis and

mitotic entry. Proteome stability in untreated extract was not a consequence of limiting ubiquitin

availability because addition of exogenous ubiquitin alone did not stimulate protein degradation

(Fig. S1a). With the exception of ubiquitin, which was added to the extract, we did not observe

any proteins that increased in abundance (Fig. S1a) in both the experiments. Addition of UbVS

alone also had no measurable effect on protein stability, possibly due to depletion of available

ubiquitin (Fig. S1a). However, addition of UbVS together with ubiquitin (UbVS/ubiquitin)

induced degradation of 34 proteins, as defined by a reduction in their abundance by at least 1.5-

fold in both experiments (Fig. 1f, g). Choosing a more relaxed threshold based on the 5% of

proteins whose abundance decreased most in UbVS/ubiquitin (FDR 2%) revealed additional

proteins that decreased in both experiments (Fig. S3a). Because transcription and translation

were inactive in extract, the decrease in protein abundance of this specific set of proteins was

likely a consequence of protein degradation that occurred as a direct consequence of DUB

inhibition. Thus, these proteins represent putative DUB substrates that are protected from

degradation by UbVS-sensitive DUBs.

Because only a limited number of proteins were destabilized by the addition of 10 μM

UbVS/ubiquitin, we tested whether stronger inhibition of UbVS-sensitive DUBs, by addition of a

higher concentration of UbVS, led to destabilization of a larger number of proteins. We

compared protein stability in untreated extract and in extract co-treated with ubiquitin and 10 μM

(as in the previous experiments) or 30 μM UbVS (Fig. S2a) using TMT-based quantitative

proteomics as before. We confirmed the degradation of the proteins observed previously using

10 μM UbVS, but we did not observe an increase in the number of proteins degraded in the

presence of 30 μM UbVS (Fig. S2b). On the contrary, the proteins decreasing in 10 μM UbVS

were more stable at the higher UbVS concentration (Fig. S2b, S2c). This finding suggested that

addition of a higher concentration of UbVS might lead to faster ubiquitin depletion, thereby

hampering the ability of UbVS to stimulate protein degradation. To test this hypothesis, we

monitored the discharge rate of the E2-Ub thioester after ubiquitin addition to extract pre-treated

with increasing concentrations of UbVS. Whereas addition of 50 μM ubiquitin was sufficient to

completely sustain Ube2C charging for 30 minutes in extract treated with 10 μM UbVS, this

concentration of ubiquitin was insufficient to maintain the E2 charged in extract treated with

higher concentration of UbVS (Fig. S2d). Thus, we cannot drive broader protein instability by

stronger DUB inhibition because ubiquitin becomes depleted too rapidly.

Confirmation of UbVS-dependent proteasomal degradation of the newly identified

substrates with human orthologs

Most of the 34 substrates that we identified have reported physiological functions, but

only 9 are known UPS substrates (Fig. 2a, Fig. S3a, Table S3). Furthermore, for only four of

them have specific DUBs been identified that control their stability. Two of these proteins are

ubiquitin ligases, MARCH7 and BIRC3, which are known to be protected from degradation by

cysteine-protease DUBs in human cells (by USP7 or USP9[29] and USP19 respectively[30]). In

addition, we identified Stam and NFX1, which are known to be deubiquitylated by the cysteine-

protease DUBs USP8[31] and USP9 respectively[32]. The ability of our screen to identify known

DUB substrates suggests that our approach is capable of identifying physiologically relevant

proteins whose stability requires DUB activity.

For the majority of the proteins we identified, DUBs have not been implicated in

regulation of their stability. We identified four additional ubiquitin ligases (MKRN1, RNF138,

ZNF598, UBOX5), whose stability is not known to be dependent on DUB activity. This

enrichment for ubiquitin ligases is again consistent with the fact that DUBs are known to protect

them from auto-ubiquitination and consequent degradation[17,33]. Other functional classes of

proteins that we identified included transcriptional regulators (GTSF1, SOX3, SOX15, TGIF2),

signaling proteins (GYGIf1, PLK3, WEE2, CSNK1E, BORA, PIM3), cytoskeletal regulators

(CRIPT, WASL, KIAA1430, HN1, ARHGEF19), proteins involved in RNA processing

(FAM32, RNASEH1, RBM18, SYF2, ZMAT22, ZFP36L11, EIF1), DNA damage components

(ETAA1, SPRTN), a lipid storage protein (PLIN2), and an uncharacterized protein (C6ORF132).

Together these findings suggest that ongoing deubiquitylation in the *Xenopus* system is

important to maintain the stability of proteins that regulate a wide variety of biological processes.

We looked for possible characteristics sequence characteristics of these substrates (Table

S3) that could contribute to their recognition by the UPS. Consistent with the PEST sequence[34]

being a signal that promotes ubiquitin and proteasome-dependent degradation, we found one or

multiple PEST sequences in 13 of the 34 hits (Table S3). Because efficient protein degradation

by the proteasome requires unstructured regions in its substrates[35], we analyzed their sequences

using ProViz[36]. We found that 8 substrates were predicted to be completely or highly disordered

whereas 25 had significantly disordered regions consistent with the possibility that they are

proteasome substrates (Table S3).

To directly test whether degradation of the identified DUB substrates was proteasome-

dependent, we quantified protein stability in extract treated with UbVS/ubiquitin in the presence

of the proteasome inhibitor MG262 or DMSO (as a control) with a TMT-based quantitative

proteomics experiment (Fig. S4a, S4b). We found that 19 proteins became unstable after

UbVS/ubiquitin addition (Fig. S4b, S4c), 14 of which were among the 34 substrates identified

(Fig. 1g). All of these proteins were stabilized by proteasome inhibition (Fig. S4b, c). Thus,

UbVS-sensitive DUBs antagonize proteasome-mediated degradation of these proteins.

To verify the findings of the proteomic experiments and directly test whether these

proteins become destabilized by DUB inhibition, we expressed 13 candidate substrates in rabbit

reticulocyte lysate and labeled them with [35]S-methionine. We chose ten proteins from our list of

34 candidates, as well as three proteins that were destabilized in at least one of the two

experiments of Fig.1f. We translated *in vitro* human orthologs, with the exception of PIM3,

where we tested the *Xenopus* protein. Subsequently, we added the translated proteins to untreated

extract or to extract pre-treated with ubiquitin, UbVS or UbVS/ubiquitin and monitored their

stability over time. We found that 12/13 proteins were degraded in the presence of

UbVS/ubiquitin but were stable in the other conditions (Fig. 2b, 2c and Fig. S3b, 3c). STAM was

the only protein that was not destabilized in the presence of UbVS/ubiquitin, but instead showed

strong poly-ubiquitination (Fig. 2d). As a control, two proteins that were stable in the proteomic

experiments (KPNA2 and HSPA8), were also tested and found to be stable when assayed by this

approach (Fig. 2b, 2c) Thus, because we could recapitulate the behaviour of the endogenous

Xenopus proteins using human orthologs, the DUB-dependent regulation of these proteins is

likely conserved, as is known already for MARCH7, BIRC3, NFX1 and STAM.

Identification of DUBs sufficient to rescue proteins from degradation in UbVS/ubiquitin-

treated extract

Next, we used these substrates to investigate the role of cysteine-protease DUBs in

counteracting their proteasomal degradation. If the proteins degraded in UbVS/ubiquitin-treated

extract were true DUB substrates, addition of recombinant DUBs to the extract should be able to

rescue their degradation. Furthermore, by testing the sufficiency of each DUB to rescue

degradation of these substrates, we can evaluate the activity and specificity of each DUB, even if

they normally function redundantly. We reasoned that the DUBs efficiently targeted by UbVS

were most likely to rescue the stability of the proteins degraded. Thus, we measured the fraction

of each DUB labeled by UbVS. After we treated the extract with 10 μM HA-UbVS, we

subsequently depleted all the HA-UbVS (and associated proteins) from the extract by

immunodepletion with anti-HA antibodies coupled to beads. Using TMT-based quantitative

proteomics, we compared the proteins remaining in extract after immunodepletion of HA-UbVS

to the proteins present in extract after incubation with empty beads. As expected, the proteins

depleted were mostly DUBs (Fig. S5a). We detected 32 cysteine-protease DUBs in undepleted

extract and found that 25 of them were depleted by HA-UbVS (Fig. 3a), with 19 DUBs being

depleted by at least 30%. There was no clear correlation between the reported abundance[24] of the

DUBs in extract (Table S1) and their fractional depletion by HA-UbVS (Fig. 3a). We selected 12

DUBs that were depleted at least 30% to test if they could rescue substrate degradation in

UbVS/ubiquitin treated extract (Fig.3b, in bold). In addition, we tested USP14 because it can

rescue several substrates from proteasomal degradation[37], as well as USP21 because it is a highly

active and non-specific DUB widely used in in vitro assays.

First, we confirmed the activity of each recombinant DUB with a UbVS-reactivity assay

(Fig. S5b, c). All DUBs tested showed almost complete labelling with UbVS, with the exception

of USP14 that requires activation by the proteasome[23,37], which was absent from these in vitro

assays. We then proceeded to add each recombinant DUB to extract pre-treated with

UbVS/ubiquitin and monitored the stability of ^{35}S-methionine-labeled ING2, CSNK1E and

CRIPT (Fig.3b). We chose these proteins as model substrates because they have not been

reported to be regulated by DUBs and they have unrelated functions. Extract was treated with

UbVS for 30 minutes, instead of 10 minutes, to ensure that the UbVS would react completely

with the endogenous DUBs and not inhibit the exogenous DUBs added to the extract (data not

shown). We began by assessing all DUBs at the same concentration (800 nM), which is

substantially greater than the endogenous concentration for ten of the fourteen DUBs tested (Fig.

S6a). We compared the ability of each DUB to rescue degradation of the selected substrates,

creating a "DUB profile" that reports the percentage of substrate rescued by each DUB (Fig. 3b).

We observed that only USP7, USP15, USP25 and USP28 could rescue degradation of these

substrates (Fig. 3b, Fig. S6b). Whereas USP28 and USP15 are present at low concentration in

extract (4 and 20 nM respectively), USP25 was not detected in the previous study indicating that

its concentration is very low, so its activity was not further evaluated. USP7 is much more

abundant in extract (150 nM), suggesting that it might be able to rescue degradation of these

substrates at its physiological concentration. In fact, when tested at physiological concentrations,

USP7 remained capable of strongly rescuing CRIPT stability, with partial rescue of ING2

stability (Fig.3c, S6c). In contrast, USP15 and USP28 did not rescue substrate degradation when

used at lower concentrations, even though these exceeded their endogenous concentrations by 5-

to 15-fold, respectively (Fig. 3c, S6c). Therefore, among all the DUBs screened, USP7 seemed

unique in its ability to rescue degradation of the substrates when tested at its physiological

concentration. In contrast, USP4, USP9X, OTUD3, USP8, USP10 and USP21 did not rescue

degradation of the three substrates, even though they were tested well above their endogenous

concentrations. We were puzzled by the fact that the two known proteasome associated DUBs,

UCHL5 and USP14, did not affect degradation of the substrates tested. Because they are very

abundant in extract, we retested them together with UCHL3 (another very abundant DUB in

extract) at 5 times their endogenous concentration and we compared them to USP7. None of

these DUBs had an effect comparable to USP7 in this assay (Fig.3d).

Because USP7, USP15, USP25 and USP28 was each sufficient to stabilize all three

substrates when used at 800 nM concentration, we wondered whether these DUBs were also

capable of restoring ubiquitin availability in UbVS-treated extracts. We treated extract with 10

μM UbVS, added each of the recombinant DUBs and monitored the charging status of Ube2C-

Ub thioester as a readout of ubiquitin availability. We included USP21 as a positive control, as it

broadly deubiquitylates substrates in vitro, as well as USP5, which can regenerate free ubiquitin

by acting on unanchored ubiquitin chains[38]. As expected, addition of exogenous ubiquitin fully

restored charging of the Ube2C-Ub thioester (Fig.3e), as did addition of USP21, consistent with

its known broad substrate specificity. USP5 did not have any effect, suggesting that free

unanchored ubiquitin chains are not generated at high levels in UbVS-treated extract. Of the 4

DUBs able to rescue substrate degradation, only USP15 was able to fully restore charging of

Ube2C. This finding suggests that USP7, USP25 and USP28 have a restricted substrate

specificity that is capable of rescuing substrate degradation without impacting ubiquitin

availability in the extract, whereas USP15 seems to have a broader specificity that allows it to

rescue degradation and also restore ubiquitin recycling. On the other hand, despite the ability of

USP21 to regenerate free ubiquitin, it is not able to rescue the degradation of any of the tested

substrates.

USP7 broadly rescues protein degradation in UbVS/ubiquitin treated extract

As USP7 seemed to be the most efficient DUB in rescuing degradation of the substrates

tested, we decided to investigate how broad an effect USP7 had in rescuing proteins from

degradation in extract treated with UbVS/ubiquitin. We performed a TMT-based quantitative

proteomics experiment comparing protein stability in extract treated with UbVS/ubiquitin, and in

extract in which USP7, USP9X or USP4 were added at 800 nM each (Fig. 4a). We tested USP9X

and USP4 as they were good targets of HA-UbVS (Fig. 3a) yet were not sufficient to rescue the

degradation of the panel of substrates that we tested (Fig 3c). We found that USP7 could rescue

16 of the 20 proteins degraded in UbVS/ubiquitin (Fig. 4c). However, we did not find any

protein stabilized by addition of USP4 or USP9X (Fig.4b, 4c), consistent with the results of our

earlier screen using the panel of [35]S-labeled substrates. The fact that human recombinant USP7

can stabilize endogenous *Xenopus* proteins suggests again that DUB-dependent regulation of the

stability of these proteins is likely evolutionarily conserved.

Because USP7 broadly rescued substrate degradation in UbVS/ubiquitin-treated extract

(Fig. 4b, c) and has been reported to be associated with the proteasome[39,40], USP7 might inhibit

the proteasome independent of its catalytic activity, as is known for USP14[37]. Thus, we tested if

the catalytic activity of USP7 was required for substrate rescue. Pre-incubation of USP7 with

UbVS inhibited rescue of substrate degradation (Fig.4d), confirming that deubiquitinating

activity of USP7 is required for the rescue. Second, we investigated if USP7 could inhibit

degradation of other known proteasome substrates. We tested if USP7 can rescue the APC

substrates cyclin B1 and securin from proteasome degradation. Surprisingly, we found that USP7

was not able to rescue them from degradation (Fig. 4e), indicating that USP7 does not generally

inhibit the degradation of all substrates of the UPS.

Inhibition of Usp7 is not sufficient to induce protein instability suggesting that DUBs

function redundantly to control protein stability in extract

Because USP7 could stabilize most of the proteins degraded in extracts treated with

UbVS/ubiquitin, we tested if inactivation of USP7 using the specific inhibitor XL-188[41] was

sufficient to induce degradation of the substrates that we identified. We also tested the effect of

USP14 inhibition on proteome stability using the specific inhibitor IU-47[42] alone or in

combination with XL-188, since USP14 inactivation promotes degradation of some proteasome

substrates in vitro and in human cells[43]. First, we verified these compounds were able to inhibit

the endogenous *Xenopus* DUBs. We pre-incubated extract with the active compounds (IU-47 and

XL-188) and the respective inactive derivatives (IU-C and XL-203) for 30 minutes, and then

added 10 μM HA-UbVS. As expected, only the active compounds were able to prevent HA-

UbVS labelling of the specific targeted DUB (Fig. S7a). After we verified that the inhibitors

were working in extract, we compared protein stability in extract pre-treated with 10 μM UbVS,

100 μM IU-47 (USP14i), 200 μM XL-188 (USP7i) or the combination of the latter inhibitors,

with TMT-based quantitative proteomics (Fig. S7b). In all conditions we added ubiquitin to

minimize any impact of DUB inhibition on ubiquitin recycling. In this experiment, we confirmed

again that 14 of the 34 candidates (Fig. S7b) were degraded in UbVS/ubiquitin treated extract.

Among these proteins, only HN1 was destabilized after specific USP7 inhibition. There were no

other proteins (detected with more than one peptide) that were destabilized following specific

USP7 inhibition. Inhibition of USP14 did not promote protein instability in *Xenopus* extract, as

none of the proteins degraded in UbVS/ubiquitin, or any other protein, was degraded in IU-47-

treated extract (Fig. S7c, d). Inhibition of both DUBs, as expected, caused the degradation of

Hn1 that was destabilized by the USP7 inhibitor. In addition, we observed degradation of three

other proteins: DBN1, KHLC1, and Supervillin (SVIL), a known USP7 substrate[44] (Fig.S7c,

bottom). Together these findings suggest that inhibition of USP7 alone or in combination with

USP14 is not sufficient to drive the broader pattern of protein instability that we observe in

extracts treated with UbVS/ubiquitin. Instead, our results support a model in which multiple

DUBs play redundant functions in maintaining stability of these substrates.

Discussion

In this study, we discovered novel DUB substrates that are protected from proteasome

degradation by cysteine-protease DUBs using a non-biased proteomic approach. One of the

challenges in identifying DUB substrates is that DUBs may function redundantly. Here, we

circumvented DUB redundancy by inactivating multiple DUBs simultaneously. We took

advantage of the *Xenopus* egg extract model system in which the quality control pathway is

silent, as protein synthesis is inhibited. Thus, proteins destabilized by DUB inhibition are more

likely to be direct DUB substrates as transcription and translation are suppressed. We

recapitulated DUB-dependent stability with recombinant human proteins, indicating that their

DUB-dependent regulation is evolutionary conserved. We showed how these novel physiological

DUB substrates can be a valuable tool for studying DUB activity and specificity. By testing the

ability of individual DUBs to rescue their degradation in a system in which DUBs are broadly

inhibited, we found that USP7 has a broad capacity to rescue substrate degradation. However, we

found that specific inhibition of USP7 is not sufficient to induce substrate instability, suggesting

that stability of the substrates that we identified is likely controlled by multiple DUBs. Together

our findings highlight the role of DUB redundancy and the importance of inhibiting multiple

DUBs to reveal new DUB substrates.

Our approach began with identifying proteins protected from proteasomal degradation by

cysteine-protease DUBs. Several of these proteins are known to be regulated by specific DUBs,

including MARCH7, BIRC3, STAM, NFX1,TRIP12, SVIL, and NEK2A[29,30,32,45,31,44,46].

Furthermore, PIM3 and WASL have not been connected previously to DUBs, but are similar in

sequence to well-established DUB substrates (PIM2[47] and WASH[48] respectively). Together,

these findings validate that our approach can identify true DUB substrates. However, the vast

majority of the proteins we identified have not been directly implicated as DUB substrates,

demonstrating the novelty of our approach. Some proteins that we identified have a proposed

physiological function, but have not been deeply studied otherwise, explaining why UPS-

dependent control of their stability has not yet been reported. As examples, we highlight CRIPT,

a neuronal cytoskeletal protein, as well as SYF2, ZMAT2 and RBM18 that are involved in RNA

processing. In contrast, a number of the other substrates we identified, including Inhibitor of

growth 2 (ING2), SPRTN, ETAA1, and the Casein Kinases ε and δ, have been extensively

studied, so it is surprising that DUB-dependent control of their stability has not yet been

reported. ING2 belongs to the family of ING proteins, which are tumor suppressors that regulate

DNA repair, cell proliferation, differentiation, senescence and apoptosis[49]. ING2 levels are low

in a variety of cancer cells and ING2 knockout mice develop a high incidence of cancer[49]. ING2

is targeted to the proteasome by the ubiquitin ligase Smurf1[50]. Here we confirmed that ING2 is a

UPS substrate and we discovered that cysteine-protease DUBs antagonize its degradation,

providing a novel mechanism that regulates its stability. SPRTN and ETAA1 are both involved

in DNA replication and DNA damage and are essential for maintaining genome stability[51,52]. A

specific mutation in SPRTN causes a syndrome that predisposes patients to cancer (the Ruijs-

Aalfs syndrome)[53] while ETAA1 (Ewing tumor-associated antigen 1) was discovered as a tumor-

specific antigen in Ewing's sarcoma[54]. Our results provide the first evidence that the UPS

controls the stability of ETAA1. SPRTN is a cell cycle-regulated protein that is known to be

ubiquitylated by APC/CDH1[55] and degraded upon exit from mitosis. Our data suggest that

UbVS-sensitive DUBs could ensure SPRTN stability during S-phase or during the DNA-damage

response when its activity is required. Casein Kinases ε and δ are versatile proteins that

participate in multiple processes such as cell cycle control, spindle organization and circadian

rhythm[56]. They auto-phosphorylate themselves inhibiting their catalytic activity[57]. Our study

suggests that DUBs could modulate their degradation by the UPS as another mechanism to

regulate their activity.

Our candidate substrates were enriched for ubiquitin ligases (n=12), extending our

understanding of the importance of DUBs in counteracting their auto-ubiquitination and

consequent degradation[17,33]. Makorin Ring Finger protein 1 (MKRN1) ubiquitylates p53 and p21

during cell growth, targeting them to proteasomal degradation[58]. Yet despite intensive study, a

role of DUBs in controlling MKRN1 stability has not been described. Our data suggest that

autoubiquitination of MKRN1 could be an important control mechanism, as it is for MDM2,

another p53 ubiquitin ligase[59]. Among the new substrates we identified, we found multiple

transcription factors (n=7) and proteins involved in RNA metabolism (n=12). These data further

support the idea that DUBs may regulate these processes[16,20] and suggest that changes in protein

levels observed in growing cells after DUB inhibition must be carefully interpreted as they could

be due to an indirect effect induced by changes in transcription/translation.

Having identified a set of novel physiological DUB substrates, we next tested the ability

of specific DUBs to antagonize their degradation under conditions in which the physiological

rates of substrate ubiquitination are maintained. The most interesting finding to emerge from this

analysis was the ability of USP7 to rescue a large number of substrates from degradation. USP7

has been highly studied and has attracted attention as a pharmacological target, as it regulates the

stability of the tumor suppressor p53 and its regulator MDM2[60]. Among the DUBs we tested,

USP7 has the greatest number of identified substrates, most of which are targeted to the

proteasome[61]. Among the substrates that we identified, a few such as MARCH7, SVIL, NEK2A

and TRIP12 are known targets of USP7[29,44,46,45] whereas two others, PIM3 and WASL, are likely

USP7 targets as USP7 is known to regulate related proteins (PIM2[47] and WASH[48] respectively)

in human cells.

What explains the broad ability of USP7 to antagonize proteasomal degradation? USP7

has been reported to associate with the proteasome[39,40], where it could intercept proteins that are

targeted for degradation. However, the physiological significance and mechanism of this

association has not been carefully studied. An intriguing possibility is that USP7 acts on

substrates that are insensitive to the other proteasome-associated DUBs, USP14 and UCHL5. For

example, USP14 has been shown to deubiquitylate proteins that carry more than one ubiquitin

chain[62]. As such, it follows that USP7 could deubiquitylate substrates bearing single ubiquitin

chain substrates that are insensitive to USP14. More studies are needed to fully understand the

substrate specificity of the proteasome-associated DUBs and the possible role of USP7 in this

context. USP7 may have a broad ability to rescue substrates from degradation because it directly

binds a wide range of proteins through recognition of common degenerate motifs (P/AxxS and

Kxxx/KxK) that are found in many proteins[61]. Another plausible explanation is that USP7 is

kinetically fast, and thus competent to counteract the rates of ubiquitination of ubiquitin ligases

that target substrates for degradation. This idea is supported by USP7 being the most active DUB

in Ub-AMC assays among a panel of 12 cysteine DUBs tested[63].

Despite USP7 being able to broadly rescue substrates from degradation, it was unable to

rescue the APC/C substrates Securin and CyclinB1. This result was not due to a lack of activity

of USP7 in mitosis because we verified that USP7 was still active in mitosis (data not shown).

The insensitivity of these proteins to USP7 could be due to the fact that USP7 is unable to

recognize K11-linked ubiquitin and mixed ubiquitin chain types that are synthetized by the

APC[22]. USP7 has been reported to deubiquitylate all types of ubiquitin linkages [63,64], but activity

towards mixed ubiquitin linkages (such as K48/K11) remains unclear.

Despite the broad ability of USP7 to rescue substrates from degradation, USP7 was

unable to rescue ubiquitin depletion in UbVS-treated extract. In contrast, Usp21 (a highly active

and nonspecific DUB) was able to rescue ubiquitin depletion, but was unable to rescue

degradation of the tested substrates. These finding suggest that even if USP7 broadly rescues

substrates from degradation, it must have a distinct and probably restricted substrate specificity

compared to USP21. For example, USP21 may deubiquitylate a limited number of highly

abundant proteins present in extract, whereas USP7 may target a distinct set of low abundance

proteins. Identifying non-degradative targets of USP7 and USP21 will be an important question

to address in future research to better understand the features underlying substrate specificity of

these enzymes.

Our study suggests that the large number of substrates protected from degradation by

USP7 reported in the literature is not simply due to USP7 having been deeply studied, but arises

from its broad ability to rescue substrates from degradation. Although many USP7 substrates

have been reported, our work suggests that USP7 may contribute to the stabilization of an even

larger number of substrates, as the identification of some substrates may have been masked by

DUB redundancy. In our system, despite USP7's ability to broadly rescue substrates from

degradation, we observed that specific inhibition of USP7 was not sufficient to induce

degradation of these substrates. Similar results were obtained for the inhibition of USP14, a

proteasome-associated DUB whose inhibition enhances proteasome activity[37]. Even

simultaneous inhibition of USP7 and USP14 destabilized very few proteins. Together these

findings support the notion that DUBs function redundantly to maintain the stability of the

proteins in this system. Because degradation is an irreversible step in the protein lifecycle, DUB

redundancy may help set a higher threshold for ubiquitination required for degradation of a

protein, beyond what is required for direct recognition by the proteasome.

Materials and Methods

Gene nomenclature

The human gene symbols have been used for consistency and simplicity in the manuscript.

Reagents

Commercial antibodies used for Western blotting analysis were as follow: anti-Ube2C (A-650, Boston Biochem), anti-ubiquitin (P4D1; sc-8017; Santa Cruz Biotechnology), anti-HA-peroxidase (3F10; Roche), anti-USP7 (A300-033A; Bethyl). Secondary antibodies used included anti-rabbit IgG-HRP (NA934V) and anti-mouse IgG-HRP (NA931V) from GE Healthcare. Chemicals used in this study included: Ubiquitin vinyl sulfone (U-202), MG262 (I-120) and Ubiquitin vinyl sulfone tagged with HA (U-212), purchased from Boston Biochem. For immunoprecipitation anti-HA-Agarose beads (A2095) from Sigma were used. Ubiquitin (U-100H) and recombinant DUBs were provided by Boston Biochem (E-519, USP7; E-520, USP8; E-320, USP5; E-325, UCHL3; E-325, UCHL5; E-546, USP25; E-552, USP9x; E-570, USP28; E-574, OTUD3; E-592, USP10; E-594, USP15; E-596, USP4).

Preparation of *Xenopus* egg extract

Interphase extract was prepared as described[65] but using 2 µg/ml calcium ionophore (A23187, Calbiochem) for egg activation. Entry into mitosis was induced by addition of 1 µM non-degradable cyclin B (MBP-Δ90) as previously described[66].

HA-UbVS immunoprecipitation

Following treatment with 10 μM HA-UbVS (10 minutes at 24°C), *Xenopus* extract was

diluted three times with XB buffer (10 mM potassium HEPES pH 7.7, 500 mM KCl, 0.1 mM

CaCl2, 1 mM MgCl$_2$, 0.5% NP40) and incubated with anti-HA beads or empty beads at 4°C (1h

and 30 minutes). At the end of the incubation, beads were washed three times with XB buffer,

SDS sample buffer was added and samples were subjected to SDS gel electrophoresis. Samples

were processed according to the GeLC-MS/MS strategy[67].

HA-UbVS immunodepletion

Xenopus extract was pre-treated with 10 μM HA-UbVS (10 minutes at 24°C) diluted

three times with XB buffer and incubated with anti-HA beads or empty beads overnight at 4°C.

Supernatants from the beads were collected and treated for Mass Spectrometry analysis by SL-

TMT method (described below).

Degradation of ^{35}S-methionine labeled substrates

Extract was pre-treated with UbVS (10 μM) for 10 min at 24°C before addition of

ubiquitin (50 μM) and substrates. The preincubation time with UbVS was extended to 30

minutes when recombinant DUBs were added to the extract. Pre-treatment of extract with

MG262 (200 μM) or specific DUB inhibitors was done at 24°C for 30 minutes. Substrates were

expressed and labeled using ^{35}S-methionine (Perkin Elmer, NEG709A500UC) with the TNT kit

(Promega: L1770). Each substrate was amplified with primers by PCR to allow T7-dependent

transcription of the PCR product or transcribed directly if plasmids contained a T7 promoter. The

translation reaction mix was added to the *Xenopus* extract at 8% final volume. Samples of the

reactions were collected at the indicated time, quenched with sodium dodecyl sulfate (SDS)

sample buffer, and processed for SDS gel electrophoresis and phosphor imaging (Bio-Rad PMI).

Quantification was performed using Quantity One software (Bio-Rad).

UbVS labeling of recombinant DUBs to verify their activity

Each DUB (1μM) was incubated with saturating amount of UbVS or HA-UbVS as

indicated (1 hour at 30°C). Reactions were stopped with addition of sodium dodecyl sulfate

(SDS) sample buffer and run on SDS-PAGE. After SDS-PAGE, the gel was stained with

Coomassie Blue.

Streamlined Tandem Mass Tag Protocol

Peptide extraction and digestion

The TMT labeling protocol and mass spectrometry analysis were based on the SL-TMT

sample preparation strategy[27]. *Xenopus* extract was collected, resuspended in the appropriate

buffer (1% SDS, 5mM DTT, 50mM Tris pH 8.8 and Pierce protease inhibitor tablet) and flash

frozen in liquid nitrogen. Methanol–chloroform precipitation was performed. Four volumes of

methanol were added to each sample and briefly vortexed. One volume chloroform was added to

the samples and vortexed again. Lastly, three volumes water was added and vortexed. The

samples were centrifuged (5 minutes, 14000 RPM) and subsequently washed twice with cold

methanol. Samples were resuspended in 200 mM EPPS, pH 8.5 and digested overnight at 24°C

with Lys-C protease (Wako Chemicals). Later samples were incubated for 6 hours at 37°C for

digestion by trypsin protease (Pierce Biotechnology).

Isobaric labeling and fractionation

Tandem mass tag (TMT) isobaric reagents (Thermo Fisher Scientific) were resuspended

in anhydrous acetonitrile (final concentration of 20 µg/µL). 10 µL of the labeling reagents plus

30 µl of acetonitrile was added to the peptides obtained by the previous digestions (~100 µg).

After incubation at room temperature (90 minutes), the reaction was quenched using

hydroxylamine to a final concentration of 0.3% (v/v). The TMT-labeled samples were pooled at

a 1:1 ratio across all the samples. Fractions were fractionated off-line using basic pH reversed

phase chromatography (BPRP). Fractions were pooled into 6 or 12 super-fractions which were

acidified with formic acid to a final concentration of 1%. Each fraction was desalted using

homemade StageTip, dried with vacuum centrifugation, and reconstituted in 5% acetonitrile and

5% formic acid for LC-MS/MS processing.

<u>Peptide detection, identification and quantification</u>

All samples were analyzed on an Orbitrap Fusion mass spectrometer (Thermo Fisher

Scientific) coupled to a Proxeon EASY-nLC 1000 liquid chromatography (LC) pump (Thermo

Fisher Scientific). Peptides were separated on a column packed with 35 cm of Accucore C18

resin (2.6 µm, 150 Å, Thermo Fisher Scientific). The column had a 100 µm inner diameter

microcapillary. For each experiment, 2 µg of peptides were loaded onto this column. Peptides

were separated, using a flow rate of 450 nL/min., with a 150-minute gradient of 3 to 25%

acetonitrile in 0.125% formic acid. Each analysis used an MS3-based TMT method, which it is

known to reduce ion interference if compared to MS2 quantification. The scan sequence starts

with an MS1 spectrum (Orbitrap analysis, resolution 120,000, 400−1400 Th, automatic gain

control (AGC) target 5E5, maximum injection time 100 ms). For subsequent MS2/MS3 analysis,

only the top 10 precursors were selected. MS2 analysis included: collision-induced dissociation

(CID), quadrupole ion trap analysis, automatic gain control (AGC) 2E4, NCE (normalized

collision energy) 35, q-value 0.25, maximum injection time 120 ms), and isolation window at

0.7. After we acquire each MS2 spectrum, we collected an MS3 spectrum in which multiple MS2

fragment ions were captured in the MS3 precursor population with isolation waveforms using

multiple frequency notches. MS3 precursors were fragmented by HCD and analyzed using the

Orbitrap (NCE 65, AGC 1.5E5, maximum injection time 150 ms, resolution was 50,000 at 400

Th). For MS3 analysis, we used charge state-dependent isolation windows: For charge state z=2,

the isolation window was set at 1.3 Th, for z=3 at 1 Th, for z=4 at 0.8 Th, and for z=5 at 0.7 Th.

Collected Spectra were processed using a Sequest-based software pipeline. Spectra were

converted to mzXML using MS Convert [68]. Database searching included all the entries from the

PHROG databas. This database includes many lower abundant proteins and multiple splice

isoforms (not present in other databases). In the original study, around 11,000 proteins were

identified[24]. Thus, our study (with ~8000 proteins) represents an acceptable coverage of the

Xenopus proteome. This database was concatenated with one composed of the sequences in the

reversed order. Searches were performed using a 50 Th precursor ion tolerance and the product

ion tolerance was set to 0.9 Da. Oxidation of methionine residues (+15.995 Da) and, where

indicated. Peptide-spectrum matches (PSMs) were adjusted to a 1% false discovery rate (FDR).

PSM filtering was performed using a linear discriminant analysis, as described previously and

assembled to a final protein-level FDR of 1%. Proteins were quantified by summing reporter ion

counts across all matching PSMs[69]. Reporter ion intensities were adjusted to correct for the

isotopic impurities of the different TMT reagents according to manufacturer specifications. The

peptides signal-to-noise (S/N) measurements assigned to each protein were summed and

normalized so that the sum of the signal for all proteins in each channel was equivalent, thereby

accounting for equal protein loading. Lastly, each protein was scaled such that the summed

signal-to-noise for that protein across all channels was 100, thereby generating a relative

abundance (RA) measurement.

TMT Mass-spectrometry analysis

A two-sided Student's t-test was used as a measure of statistical confidence of the

observed $\log_2$ fold change. Selected candidates met both thresholds (Fc$\leq$ -1.5 and *p*-value<0.05)

in the experiments. For Figure 1g, the candidates were included if at least one peptide was

identified in both of the experiments. For subsequent figures, where a single experiment was

performed, candidates were included only if they were detected with at least 2 different peptides.

When multiple isoforms of the same protein decreased in UbVS/ubiquitin, only the isoform with

more peptides was selected (for simplicity isoforms are not indicated in the figures).

References

1.   Swatek, K. N. & Komander, D. Ubiquitin modifications. *Cell Res.* **26**, 399–422 (2016).

2.   Hershko, A. & Ciechanover A. The ubiquitin system. *Annu. Rev. Biochem.* **67**, 425–79

(1998).

3.   Clague, M. J., Heride, C. & Urbé, S. The demographics of the ubiquitin system. *Trends*

Cell Biol. **25**, 417–426 (2015).

4.   Komander, D. & Rape, M. The ubiquitin code. *Annu. Rev. Biochem.* **81**, 203–29 (2012).

5.   Jackson, S. P. & Durocher, D. Regulation of DNA damage responses by ubiquitin and

SUMO. *Mol. Cell* **49**, 795–807 (2013).

6.   Lauwers, E., Jacob, C. & André, B. K63-linked ubiquitin chains as a specific signal for

protein sorting into the multivesicular body pathway. *J. Cell Biol.* **185**, 493–502 (2009).

7.   Chau, V. *et al.* A multiubiquitin chain is confined to specific lysine in a targeted short-

lived protein. *Science* **243**, 1576–83 (1989).

8.   Ye, Y. & Rape, M. Building ubiquitin chains: E2 enzymes at work. *Nat. Rev. Mol. Cell*

Biol. **10**, 755–64 (2009).

9.   Clague, M. J., Coulson, J. M. & Urbé, S. Cellular functions of the DUBs. *J Cell Sci* **125**,

277–286 (2012).

10.   Mevissen, T. E. T. & Komander, D. Mechanisms of deubiquitinase specificity and

regulation. **86**, 159–92 (2017).

11.   Clague, M. J. *et al.* Deubiquitylases from genes to organism. *Physiol. Rev.* **93**, 1289–1315

(2013).

12.   Ma, J. *et al.* C-terminal region of USP7/HAUSP is critical for deubiquitination activity

and contains a second mdm2/p53 binding site. *Arch. Biochem. Biophys.* **503**, 207–212

(2010).

13.   Bremm, A., Freund, S. M. V & Komander, D. Lys11-linked ubiquitin chains adopt

593   compact conformations and are preferentially hydrolyzed by the deubiquitinase Cezanne.

594   *Nat. Struct. Mol. Biol.* **17**, 939–947 (2010).

595   14.   Wang, T. *et al.* Evidence for bidentate substrate binding as the basis for the K48 linkage

596   specificity of Otubain 1. *J. Mol. Biol.* **386**, 1011–1023 (2009).

597   15.   Flierman, D. *et al.* Non-hydrolyzable diubiquitin probes reveal linkage-specific reactivity

598   of deubiquitylating enzymes mediated by S2 pockets. *Cell Chem. Biol.* **23**, 472–82 (2016).

599   16.   Sowa, M. E., Bennett, E. J., Gygi, S. P. & Harper, J. W. Defining the human

600   deubiquitinating enzyme interaction landscape. *Cell* **138**, 389–403 (2009).

601   17.   Kim, W. *et al.* Systematic and quantitative assessment of the ubiquitin-modified

602   proteome. *Mol. Cell* **44**, 325–40 (2011).

603   18.   Beckley, J. R., Chen, J.-S., Yang, Y., Peng, J. & Gould, K. L. A Degenerate cohort of

604   yeast membrane trafficking DUBs mediates cell polarity and survival. *Mol. Cell.*

605   *Proteomics* **14**, 3132–3141 (2015).

606   19.   Kwon, S.-K., Saindane, M. & Baek, K.-H. p53 stability is regulated by diverse

607   deubiquitinating enzymes. *Biochim. Biophys. Acta - Rev. Cancer* **1868**, 404–411 (2017).

608   20.   Clague, M. J., Urbé, S. & Komander, D. Breaking the chains: deubiquitylating enzyme

609   specificity begets function. *Nat. Rev. Mol. Cell Biol.* **20**, 338–352 (2019).

610   21.   Hegde, R. S. & Zavodszky, E. Recognition and Degradation of Mislocalized Proteins in

611   Health and Disease. *Cold Spring Harb. Perspect. Biol.* a033902 (2019).

612   doi:10.1101/cshperspect.a033902

613   22.   Dimova, N. V. *et al.* APC/C-mediated multiple monoubiquitylation provides an

alternative degradation signal for cyclin B1. *Nat. Cell Biol.* **14**, 168–176 (2012).

23. Borodovsky, A. *et al.* A novel active site-directed probe specific for deubiquitylating enzymes reveals proteasome association of USP14. *EMBO J.* **20**, 5187–96 (2001).

24. Wühr, M. *et al.* Deep proteomics of the Xenopus laevis egg using an mRNA-derived reference database. *Curr. Biol.* **24**, 1467–1475 (2014).

25. de Poot, S. A. H., Tian, G. & Finley, D. Meddling with fate: The proteasomal deubiquitinating enzymes. *J. Mol. Biol.* **429**, 3525–3545 (2017).

26. Soncini, C., Berdo, I. & Draetta, G. Ras–GAP SH3 domain binding protein (G3BP) is a modulator of USP10, a novel human ubiquitin specific protease. *Oncogene* **20**, 3869–3879 (2001).

27. Navarrete-Perea, J., Yu, Q., Gygi, S. P. & Paulo, J. A. Streamlined Tandem Mass Tag (SL-TMT) Protocol: an efficient strategy for quantitative (phospho)proteome profiling using Tandem Mass Tag-synchronous precursor selection-MS3. *J. Proteome Res.* **17**, 2226–2236 (2018).

28. Richter, J. D. & Smith, L. D. Reversible inhibition of translation by Xenopus oocyte-specific proteins. *Nature* **309**, 378–80

29. Nathan, J. A. *et al.* The Ubiquitin E3 Ligase MARCH7 is Differentially Regulated by the Deubiquitylating Enzymes USP7 and USP9X. *Traffic* **9**, 1130–1145 (2008).

30. Mei, Y., Hahn, A. A., Hu, S. & Yang, X. The USP19 deubiquitinase regulates the stability of c-IAP1 and c-IAP2. *J. Biol. Chem.* **286**, 35380–7 (2011).

31. Berlin, I., Schwartz, H. & Nash, P. D. Regulation of Epidermal Growth Factor Receptor Ubiquitination and Trafficking by the USP8·STAM Complex. *J. Biol. Chem.* **285**, 34909–34921 (2010).

32. Chen, X. *et al.* Identification of a USP9X Substrate NFX1-123 by SILAC-Based

Quantitative Proteomics. *J. Proteome Res.* **18**, 2654–2665 (2019).

33. Ventii, K. H. & Wilkinson, K. D. Protein partners of deubiquitinating enzymes. *Biochem.*

J. **414**, 161–75 (2008).

34. Rechsteiner, M. & Rogers, S. W. PEST sequences and regulation by proteolysis. *Trends*

Biochem. Sci. **21**, 267–71 (1996).

35. Aufderheide, A., Unverdorben, P., Baumeister, W. & Förster, F. Structural disorder and its

role in proteasomal degradation. *FEBS Lett.* **589**, 2552–2560 (2015).

36. Jehl, P., Manguy, J., Shields, D. C., Higgins, D. G. & Davey, N. E. ProViz—a web-based

visualization tool to investigate the functional and evolutionary features of protein

sequences. *Nucleic Acids Res.* **44**, W11–W15 (2016).

37. Lee, B.-H. *et al.* Enhancement of proteasome activity by a small-molecule inhibitor of

USP14. *Nature* **467**, 179–184 (2010).

38. Hadari, T., Warms, J. V, Rose, I. A. & Hershko, A. A ubiquitin C-terminal isopeptidase

that acts on polyubiquitin chains. Role in protein degradation. *J. Biol. Chem.* **267**, 719–27

(1992).

39. Bousquet-Dubouch, M.-P. *et al.* Affinity purification strategy to capture human

endogenous proteasome complexes diversity and to identify proteasome-interacting

proteins. *Mol. Cell. Proteomics* **8**, 1150–64 (2009).

40. Besche, H. C., Haas, W., Gygi, S. P. & Goldberg, A. L. Isolation of mammalian 26S

proteasomes and p97/VCP complexes using the ubiquitin-like domain from HHR23B

reveals novel proteasome-associated proteins †. *Biochemistry* **48**, 2538–2549 (2009).

41. Lamberto, I. *et al.* Structure-guided development of a potent and selective non-covalent

active-site inhibitor of USP7. *Cell Chem. Biol.* **24**, 1490-1500.e11 (2017).

42.   Boselli, M. *et al.* An inhibitor of the proteasomal deubiquitinating enzyme USP14 induces

tau elimination in cultured neurons. *J. Biol. Chem.* **292**, 19209–19225 (2017).

43.   Lee, B.-H. *et al.* Enhancement of proteasome activity by a small-molecule inhibitor of

USP14. *Nature* **467**, 179–184 (2010).

44.   Fang, Z. & Luna, E. J. Supervillin-mediated suppression of p53 protein enhances cell

survival. *J. Biol. Chem.* **288**, 7918–29 (2013).

45.   Cai, J.-B. *et al.* Ubiquitin-specific protease 7 accelerates p14 [ARF] degradation by

deubiquitinating thyroid hormone receptor-interacting protein 12 and promotes

hepatocellular carcinoma progression. *Hepatology* **61**, 1603–1614 (2015).

46.   Franqui-Machin, R. *et al.* Destabilizing NEK2 overcomes resistance to proteasome

inhibition in multiple myeloma. *J. Clin. Invest.* **128**, 2877–2893 (2018).

47.   Kategaya, L. *et al.* USP7 small-molecule inhibitors interfere with ubiquitin binding.

Nature **550**, 534–538 (2017).

48.   Hao, Y.-H. *et al.* USP7 acts as a molecular rheostat to promote WASH-dependent

endosomal protein recycling and is mutated in a human neurodevelopmental disorder.

Mol. Cell **59**, 956–969 (2015).

49.   Guérillon, C., Larrieu, D. & Pedeux, R. ING1 and ING2: multifaceted tumor suppressor

genes. *Cell. Mol. Life Sci.* **70**, 3753–3772 (2013).

50.   Nie, J. *et al.* HECT ubiquitin ligase Smurf1 targets the tumor suppressor ING2 for

ubiquitination and degradation. *FEBS Lett.* **584**, 3005–3012 (2010).

51.   Vaz, B. *et al.* Metalloprotease SPRTN/DVC1 orchestrates replication-coupled DNA-

protein crosslink repair. *Mol. Cell* **64**, 704–719 (2016).

52.   Haahr, P. *et al.* Activation of the ATR kinase by the RPA-binding protein ETAA1. *Nat.*

Cell Biol. **18**, 1196–1207 (2016).

53.   Lessel, D. *et al.* Mutations in SPRTN cause early onset hepatocellular carcinoma, genomic

instability and progeroid features. *Nat. Genet.* **46**, 1239–1244 (2014).

54.   Borowski, A. *et al.* Structure and function of ETAA16: a novel cell surface antigen in

Ewing's tumours. *Cancer Immunol. Immunother.* **55**, 363–374 (2006).

55.   Mosbech, A. *et al.* DVC1 (C1orf124) is a DNA damage–targeting p97 adaptor that

promotes ubiquitin-dependent responses to replication blocks. *Nat. Struct. Mol. Biol.* **19**,

1084–1092 (2012).

56.   Schittek, B. & Sinnberg, T. Biological functions of casein kinase 1 isoforms and putative

roles in tumorigenesis. *Mol. Cancer* **13**, 231 (2014).

57.   Budini, M. *et al.* Autophosphorylation of carboxy-terminal residues inhibits the activity of

protein kinase CK1α. *J. Cell. Biochem.* **106**, 399–408 (2009).

58.   Lee, E.-W. *et al.* Differential regulation of p53 and p21 by MKRN1 E3 ligase controls cell

cycle arrest and apoptosis. *EMBO J.* **28**, 2100–2113 (2009).

59.   Ranaweera, R. S. & Yang, X. Auto-ubiquitination of Mdm2 enhances its substrate

ubiquitin ligase activity. *J. Biol. Chem.* **288**, 18939–46 (2013).

60.   Sheng, Y. *et al.* Molecular recognition of p53 and MDM2 by USP7/HAUSP. *Nat. Struct.*

Mol. Biol. **13**, 285–291 (2006).

61.   Kim, R. Q. & Sixma, T. K. Regulation of USP7: A High Incidence of E3 Complexes. *J.*

Mol. Biol. **429**, 3395–3408 (2017).

62.   Lee, B.-H. *et al.* USP14 deubiquitinates proteasome-bound substrates that are

ubiquitinated at multiple sites. *Nature* **532**, 398–401 (2016).

63.   Faesen, A. C. *et al.* The differential modulation of USP activity by internal regulatory

domains, interactors and eight ubiquitin chain types. *Chem. Biol.* **18**, 1550–61 (2011).

64.   Ritorto, M. S. *et al.* Screening of DUB activity and specificity by MALDI-TOF mass

spectrometry. *Nat. Commun.* **5**, 4763 (2014).

65.   Murray AW. Cell cycle extracts. *Methods Cell Biol* **36**, 581–605 (1991).

66.   Zeng, X. *et al.* Pharmacologic inhibition of the Anaphase-Promoting Complex Induces a

spindle checkpoint-dependent mitotic arrest in the absence of spindle damage. *Cancer*

Cell **18**, 382–395 (2010).

67.   Paulo, J. A. Sample preparation for proteomic analysis using a GeLC-MS/MS strategy. *J.*

Biol. Methods **3**, 45 (2016).

68.   Adusumilli, R. & Mallick, P. Data conversion with proteoWizard msConvert. in *Methods*

in Molecular Biology **1550**, 339–368 (2017).

69.   McAlister, G. C. *et al.* Increasing the multiplexing capacity of TMTs using reporter ion

isotopologues with isobaric masses. *Anal. Chem.* **84**, 7469–78 (2012).

70   Hewings D. *et al*. Reactive-site-centric chemoproteomics identifies a distinct class of

deubiquitinase enzymes *Nat. Commun .***9** 1162 (2018)

Acknowledgments

We gratefully acknowledge W. Harper for the gift of plasmids from the ORFeome collection, S.

Buhrlage for the gift of XL-188 and XL-203 and D. Finley for the gift of IU-C, IU-47 and

USP21. We thank W. Harper for helpful discussions. This work was supported by NIH grants

R01 GM132129 (J.A.P.), R01 GM67945 (S.P.G) and R35 GM127032 (R.W.K.).

Author contributions

V.R. and R.W.K designed the experiments

V.R. performed all the experiments

B.B. provided advice on DUB selection and DUB assays

J.A.P., J.C. and S.P.G. performed mass spectrometric analysis

R.W.K and V.R. wrote the manuscript with input from all the authors

Competing interests

The authors declare no competing interests

Rossio et al., Fig1

Fig.1 Inhibition of cysteine-protease DUBs by UbVS causes rapid ubiquitin depletion and degradation of a specific set of proteins. (a) Interphase *Xenopus* extract was treated with the indicated concentrations of UbVS or **(b)** HA-UbVS. After 10 minutes (time 0), 50 μM ubiquitin or buffer were added (A, bottom). Aliquots were quenched with **(a)** non-reducing or **(b)** reducing sample buffer and subjected to immunoblotting (IB) for the indicated proteins. **(c)** Extract was incubated with 10 μM HA-UbVS (10 minutes) and subjected to anti-HA pull-down followed by analysis of eluates by tandem mass spectrometry. The number of DUBs identified with the anti-HA pull-down (gray) is compared to the number of DUBs detected previously in extract[24] (white). **(d)** Mass spectrometry analysis by Streamlined-Tandem Mass Tag (SL-TMT) method is summarized. **(e)** Ubiquitin or buffer were added to the extract (time 0) pre-incubated with buffer or 10 μM UbVS for 10 minutes. Samples were collected in duplicate at time 0 and 60 minutes and processed as described in (d). **(f)** Volcano plots of TMT analysis comparing the proteins detected in UbVS/ubiquitin at 60 minutes with the proteins detected at time 0 (2 independent experiments). Statistical significance ($-\log_{10}$ p-value) is plotted against ratio (average $\log_2$). Proteins significantly decreasing in both the experiments are showed in black. **(g)** $\log_2$ ratio heat map of the proteins significantly downregulated in UbVS/ubiquitin in both the experiments in (f).

a

Proteins	Function
MKRN1 UBOX5 (putative) **MARCH7 RNF138** BIRC3 NFX1 (putative) ZNF598	UPS (E3s)
RBM18 SYF2 EIF1 FAM32A **ZMAT2** RNASEH1 ZFP36L1	RNA metabolism
SOX3 SOX15 TGIF2 GTSF1	Transcription factors
CRIPT ARHGEF19 KIAA1430 WASL	Cytoskeleton
BORA WEE2 **PIM3** GIGYF1 HN1 PLK3	Proliferation/cell death
SPRTN ETAA1	DNA damage
STAM (trafficking) **CSNK1E** (circadian period) PLIN2 (lipid storage) C6ORF132 (?)	Others

b

c

d

Rossio et al.,Fig2

Fig. 2 Independent validation of the proteins protected from degradation by UbVS-

sensitive DUBs (a) Functions of the proteins down-regulated in UbVS/ubiquitin-treated extract

(Fig.1f, g). Proteins independently validated in are shown in bold. (**b**) Human orthologs (prefix

h) of the indicated proteins, with the exception of PIM3, where the *Xenopus* protein was used,

were expressed in rabbit reticulocyte lysate, labeled with ^{35}S-methionine and added to the

extract, treated as described for the TMT proteomic experiments (Fig.1e). Aliquots were

collected at the indicated times, quenched with SDS sample buffer and processed for SDS-PAGE

and phosphorimaging. UbVS/ubiquitin-dependent degradation of these proteins was confirmed

in 2 independent experiments. (**c**) Quantification of one experiment. Quantification of the second

experiment is shown in Fig. S3d (**d**) Performed as in (b) *: ubiquitinated species.

Rossio et al., Fig3

Fig 3. Identification of DUBs sufficient to rescue degradation of the selected substrates in UbVS/ubiquitin-treated extract (a) Extract was incubated with 10 μM HA-UbVS (10 minutes) overnight at 4°C with anti-HA or empty beads. Supernatants were collected in triplicate for mass spectrometry analysis by the SL-TMT method. Percentage of each DUB depleted by 10 μM HA-UbVS. Bold: DUBs tested in subsequent experiments. **(b)** Extract was pre-incubated with 10 μM UbVS (30 minutes), after which recombinant DUBs (800 nM),^{35}S-methionine labeled substrates and buffer or ubiquitin were added. After 60 minutes, aliquots were quenched with sample buffer and processed for SDS-PAGE and phosphorimaging. Quantitative profiles for the 3 model substrates tested were created. The percentage of each substrate rescued by the addition of the indicated DUBs is reported for two independent experiments. Gels for one of the experiments are shown in Fig. S6 **(c)** and **(d).** Dose response analysis, performed as in **(b)** but at varying DUB concentrations. The tested concentration is shown; the value in parentheses indicates the fold-increase of the tested concentration relative to the endogenous concentration. **(e)** Performed as in **(b)** with the exception that ubiquitin was not added to the extracts with the DUBs. Samples were collected at the indicated times (time 0), quenched with non-reducing sample buffer and subjected to immunoblotting.

Rossio et al., Fig4

Fig.4 USP7 broadly rescues substrates from degradation in UbVS/ubiquitin-treated extract

(a) Top: Extract was incubated with 10 μM UbVS (30 minutes) followed by addition of DUBs (800 nM) and ubiquitin. Samples were collected in duplicate and processed for mass spectrometry analysis by SL-TMT method. Bottom: R^2 of the technical replicates for each condition. **(b)** Volcano plot of quantitative proteomics analysis comparing proteins detected in the different conditions. Statistical significance ($-\log_{10}$ p-value) is plotted against ratio (average $\log_2$ ratio). Proteins significantly decreasing in UbVS/ubiquitin are indicated by blue dots in all the conditions tested. **(c)** $\log_2$ ratio heat map of the proteins significantly downregulated in UbVS/ubiquitin after addition of the indicated DUB. Bold: Proteins decreasing in both experiments in Fig. 1f. **(d)** Extract was incubated with 10 μM UbVS (30 minutes) followed by addition of USP7 (800 nM), ubiquitin and ^{35}S-methionine labeled substrates (time 0). USP7 was added to extract after pre-incubation (30 minutes) with buffer or UbVS (USP7-UbVS). Right: Quantification of the experiment. Error bars: SEM of three independent experiments. One representative experiment is shown. **(e)** ^{35}S-methionine labeled substrates were added to interphase extract or to mitotically arrested extract pre-incubated with UbVS (30 minutes). Concomitantly, USP7 and ubiquitin were added (time 0). Aliquots were collected (20 minutes), quenched as usual and processed for SDS-PAGE and phosphorimaging. Right: Quantification of the substrate present at 20 minutes in the experiment on the left. Error bars: SEM of three independent experiments. One representative experiment is shown.