Proteomics of broad deubiquitylase inhibition unmasks redundant enzyme function to re-

veal substrates

Valentina Rossio[1], Joao A. Paulo[1], Joel Chick[2] Bradley Brasher[3], Steven P. Gygi[1], Randall W.

King[1*]

[1] Department of Cell Biology, Blavatnik Institute at Harvard Medical School, Boston, MA 02115, USA.

[2] Present address: Vividion Therapeutics San Diego, CA 92121, USA.

[3] Boston Biochem, a Bio-Techne Brand, Cambridge, MA 02139, USA.

*Correspondence to Randall W. King: randy_king@hms.harvard.edu

Abstract

Deubiquitylating enzymes (DUBs) counteract ubiquitylation to control stability or activity of

substrates. Identification of DUB substrates is challenging because multiple DUBs act on the

same substrates, thwarting genetic approaches. Here, we circumvented redundancy by broadly

inhibiting DUBs in *Xenopus* egg extract. DUB inhibition increases ubiquitylation of hundreds of

proteins, depleting free ubiquitin without inducing widespread degradation. Restoring available

ubiquitin led to proteasomal degradation of over thirty proteins, indicating that deubiquitylation

is essential to maintain their stability. We confirmed their DUB-dependent stability with recom-

binant human proteins, demonstrating evolutionary conservation. We profiled the ability of

DUBs to rescue protein stability, and found that USP7 has a unique ability to broadly antagonize

proteasomal degradation. Together, we provide a comprehensive characterization of ubiquitin

dynamics in the *Xenopus* system, identify new DUB substrates, and present a new approach to

characterize physiological DUB specificity that overcomes challenges posed by DUB redun-

dancy

Introduction

Conjugation of ubiquitin to proteins is a widespread, highly regulated post-translational modification. The Ubiquitin-Proteasome System (UPS) is best known for its ability to target specific proteins for degradation. Still, ubiquitylation also regulates protein localization, activity and function independently of degradation(Swatek and Komander 2016)·(Hershko and Ciechanover A. 1998)·(Clague, Heride, and Urbé 2015). Ubiquitin can be incorporated in polyubiquitin chains of different topologies that may result in different fates(Komander and Rape 2012). For example, lysine-63 linked chains promote non-degradative functions (Jackson and Durocher 2013)·(Lauwers, Jacob, and André 2009), whereas lysine-48 linked chains and branched chains such as K11/K48 or K29/K48 promote proteasomal degradation(Chau et al. 1989). Ubiquitin is covalently attached to substrates via a cascade of E1-E2-E3 enzymes(Yihong Ye and Rape 2009). In contrast, deubiquitylating enzymes (DUBs) catalyze ubiquitin removal(Clague, Coulson, and Urbé 2012).

Comprehensive proteomic identification of physiological UPS substrates has been challenging due to the prominence of protein-quality control pathways that ubiquitylate newly synthesized proteins that do not fold properly in growing cells(W. Kim et al. 2011). Therefore, it can be difficult to distinguish regulatory ubiquitylation of mature proteins from more widespread ubiquitylation that targets misfolded proteins for degradation during protein biogenesis. To overcome this challenge, in our study we took advantage of the *Xenopus laevis* egg extract model system. Since the mature *Xenopus* egg has completed its growth prior to being laid, there is little ongoing translation and protein folding, thereby minimizing the contribution of quality-control ubiquitylation compared to actively growing cultured cells. Thus, ubiquitylation events observed in these extracts are more likely to reflect physiologic regulatory events rather than quality control.

Xenopus egg extract can be prepared with little dilution of the cytoplasm, therefore preserving

physiological integrity and enabling the reconstitution of complex biochemical processes such as

DNA damage repair cell cycle progression and mitotic spindle formation (Yardimci et al.

2012)ˌ(Glotzer, Murray, and Kirschner 1991)ˌ(Field and Mitchison 2018). *Xenopus* egg extract has

been routinely used for studies of protein ubiquitylation and degradation, including studies of mi-

totic cyclin degradation(Glotzer, Murray, and Kirschner 1991) as well as regulatory ubiquitylation

during DNA replication(Yardimci et al. 2012). Together these studies indicate that ubiquitylation

regulates important physiological processes in *Xenopus* egg extracts. Still, we have a limited un-

derstanding of ubiquitin homeostasis and the overall dynamics of ubiquitylation, deubiquitylation,

and protein degradation in this experimental system.

The human genome encodes approximately 100 DUBs, divided into two families: the zinc

metalloprotease class (10 DUBs), and the cysteine-protease class that contains most other DUBs

(~90)(Mevissen and Komander 2017)ˌ(Clague et al. 2013). A few DUBs have been well-studied

with many characterized substrates, but most DUBs still do not have any known substrates, limit-

ing our understanding of how DUB substrate specificity is achieved. DUB specificity is complex,

as it can arise through binding specificity mediated by unique protein-protein interaction do-

mains(Ma et al. 2010), or through recognition of specific topologies of ubiquitin chains(Bremm,

Freund, and Komander 2010; Wang et al. 2009)ˌ(Flierman et al. 2016). Most DUB substrates have

been discovered by first identifying proteins that interact with specific DUBs. This approach has

been used globally(Sowa et al. 2009), but it may miss substrates that interact weakly with DUBs

or identify regulators or scaffold proteins rather than true substrates. Identifying DUB targets is

difficult because DUBs may function redundantly(Beckley et al. 2015)ˌ(Kwon, Saindane, and

Baek 2017), and thus inactivating a single DUB may not destabilize its substrates or affect a

particular biological process. In the case of proteome stability, it is not known which DUBs have

the broadest impact in rescuing proteins from proteasome degradation or to what extent DUBs act

redundantly in this process. Therefore, new approaches are needed to identify physiological sub-

strates of DUBs and to overcome the challenges posed by redundant function of these enzymes.

In this study we applied unbiased quantitative proteomic approaches to characterize the

dynamics of ubiquitylation and proteasomal degradation in *Xenopus* egg extract. We found that

the proteome is stable, despite widespread ubiquitylation, suggesting that most ubiquitylation is

non-degradative. By broadly inhibiting cysteine-protease DUBs, we circumvented the possible

effects of redundancy on proteome stability and discovered a set of proteins whose stability de-

pends on DUB activity. We next took advantage of this panel of substrates to identify DUBs that

are sufficient to counteract proteasomal degradation of these proteins. By broadly inactivating

DUBs and adding back a panel of recombinant DUBs one by one, we unmasked DUB redundancy

and discovered that USP7 can rescue many substrates from degradation. However, specific inhi-

bition of USP7 with a small molecule inhibitor was not sufficient to promote degradation of most

of the substrates we identified, suggesting that USP7 functions redundantly with other DUBs. Our

work highlights the impact of DUB redundancy on proteome stability and reveals the specificity

and activity of DUBs whose function would otherwise be masked by redundancy.

Results

UbVS treatment induces rapid depletion of available ubiquitin and labels a broad set of cysteine-protease DUBs in *Xenopus* extract

To selectively inhibit cysteine protease DUBs in Xenopus extract, we used ubiquitin vinyl sulfone (UbVS), which covalently inhibits cysteine-protease DUBs, without inhibiting other cysteine proteases or other classes of DUBs(Borodovsky et al. 2001). Our laboratory previously showed that UbVS treatment efficiently blocks ubiquitin recycling in *Xenopus* extract(Dimova et al. 2012). We confirmed that 10 µM UbVS was sufficient to rapidly deplete free ubiquitin, (Fig. 1a, 1b) which was associated with rapid discharge of the Ube2C-ubiquitin thioester within 5 minutes (Fig. 1a, bottom). Other E2s were similarly rapidly discharged (data not shown). Addition of exogenous ubiquitin to UbVS-treated extract fully recharged the Ube2C thioester confirming that the E2's discharge was due to the depletion of free ubiquitin (Fig.1a, bottom). To determine the spectrum of cysteine-containing DUBs targeted by this concentration of UbVS, extract was treated with 10 µM HA-tagged UbVS (HA-UbVS) and sensitive DUBs were visualized as discrete bands by anti-HA Western blotting analysis (Fig. 1b, right). To identify these DUBs, we performed mass spectrometry analysis of immunopurified HA-UbVS from extract and identified 88 proteins (Table S2), 35 of which were cysteine-protease DUBs (Table S1, S2) that are likely direct targets of UbVS. All known classes of cysteine-protease DUBs were found (Fig. 1c, Table S1), whereas metallo-protease DUBs (JAMM) were not identified, as expected(Borodovsky et al. 2001). Prior proteomic analysis of *Xenopus* extract detected 54 DUBs(Wühr et al. 2014) (Table S1), 51 of which were cysteine-proteases. Thus, 10 µM HA-UbVS labeled 35 of 51 (69%) of cysteine-protease DUBs present in extract, consistent with broad specificity of UbVS for this class of DUBs(Borodovsky et al. 2001). The remaining DUBs were likely not identified either because

they were present in low abundance or because they do not react rapidly with UbVS. Our HA-

UbVS pull-down also isolated proteins that are likely to strongly associate with DUBs (Table S2),

as the pull-down was carried out in the presence of high salt (500 mM KCl). A high fraction of

these proteins (31) are components of the proteasome, consistent with the known interaction of

UbVS-sensitive DUBs UCHL5 and USP14(de Poot, Tian, and Finley 2017) with the proteasome.

Other known interactors of DUBs that we isolated included GBP1/GBP2, which binds

USP10(Soncini, Berdo, and Draetta 2001), and the SPATA proteins (SPATA2/SPATA2L), which

have been identified as CYLD interactors(Sowa et al. 2009) in human cells (Table S2), suggesting

evolutionary conservation of these interactions.

UbVS treatment induces degradation of a limited set of proteins when available ubiquitin is

restored

We first wanted to investigate how inhibition of multiple DUBs influenced proteome sta-

bility in *Xenopus* extract. We hypothesized that simultaneous inhibition of 35 DUBs with UbVS

might lead to destabilization of proteins that require ongoing deubiquitylation to maintain their

stability. By simultaneously inhibiting a large number of DUBs, we predicted that we might be

able to identify new substrates of DUBs, including substrates whose stability is conferred by the

action of redundant DUBs. We used a multiplexed Tandem Mass Tag (TMT)-based quantitative

proteomic approach(Navarrete-Perea et al. 2018) (Fig. 1d), and we measured protein abundance

over time in untreated extract, as well as extract to which ubiquitin, UbVS, or the combination

(UbVS/ubiquitin) were added (Fig. 1e). We reasoned that proteins that decreased specifically in

the presence of UbVS and/or UbVS/ubiquitin may normally be protected from degradation by

UbVS-sensitive DUBs. We performed two independent experiments using *Xenopus* extract

prepared from two sets of eggs (collected from different animals and processed separately), to

ensure reproducibility. In each experiment we quantified around 8000 proteins, with an overlap of

94.8% between the two experiments.

First, we observed that in the absence of any perturbation, the proteome was stable in

Xenopus extract over the course of 60 minutes. Few proteins changed in abundance over time, and

those that did were not shared between the experiments (Fig. S1a). This degree of proteome

stability is consistent with the idea that bulk rates of translation are relatively quiescent in the

Xenopus system(Richter and Smith n.d.) and that the extract was treated with cycloheximide to

prevent protein synthesis and mitotic entry. Proteome stability in untreated extract was not a con-

sequence of limiting ubiquitin availability because addition of exogenous ubiquitin alone did not

stimulate protein degradation (Fig. S1a). With the exception of ubiquitin, which was added to the

extract, we did not observe any proteins that increased in abundance (Fig. S1a) in both the

experiments. Addition of UbVS alone also had no measurable effect on protein stability, possibly

due to depletion of available ubiquitin (Fig. S1a). However, addition of UbVS together with ubiq-

uitin (UbVS/ubiquitin) induced degradation of 34 proteins, as defined by a reduction in their abun-

dance by at least 1.5-fold in both experiments (Fig. 1f, g). Choosing a more relaxed threshold based

on the 5% of proteins whose abundance decreased most in UbVS/ubiquitin (FDR 1%) revealed

additional proteins that decreased in both experiments (Fig. S4a). Because transcription and trans-

lation were inactive in extract, the decrease in protein abundance of this specific set of proteins

was likely a consequence of protein degradation that occurred as a direct consequence of DUB

inhibition. Thus, these proteins represent putative DUB substrates that are protected from degra-

dation by UbVS-sensitive DUBs.

Because only a limited number of proteins were destabilized by the addition of 10 μM

UbVS/ubiquitin, we tested whether stronger inhibition of UbVS-sensitive DUBs, by addition of a

higher concentration of UbVS, led to destabilization of a larger number of proteins. We compared

protein stability in untreated extract and in extract co-treated with ubiquitin and 10 µM (as in the

previous experiments) or 30 µM UbVS (Fig. S2a) using TMT-based quantitative proteomics as

before. We confirmed the degradation of the proteins observed previously using 10 µM UbVS, but

we did not observe an increase in the number of proteins degraded in the presence of 30 µM UbVS

(Fig. S2b). On the contrary, the proteins decreasing in 10 µM UbVS were more stable at the higher

UbVS concentration (Fig. S2b, S2c). This finding suggested that addition of a higher concentration

of UbVS might lead to faster ubiquitin depletion, thereby hampering the ability of UbVS to stim-

ulate protein degradation. To test this hypothesis, we monitored the discharge rate of the E2-Ub

thioester after ubiquitin addition to extract pre-treated with increasing concentrations of UbVS.

Whereas addition of 50 µM ubiquitin was sufficient to completely sustain Ube2C charging for 30

minutes in extract treated with 10 µM UbVS, this concentration of ubiquitin was insufficient to

maintain charged Ube2C in extract treated with 30 µM UbVS (Fig. S2d). Thus, we cannot drive

broader protein instability by stronger DUB inhibition because ubiquitin becomes depleted too

rapidly.

Non-degradative ubiquitylation targets a large number of proteins in *Xenopus* egg extract

Since addition of UbVS causes rapid depletion of available ubiquitin and discharge of E2

enzymes (Fig.1a, b), the ubiquitin conjugation machinery must be active in *Xenopus* extract. We

were therefore surprised by the fact that addition of ubiquitin or UbVS alone did not stimulate

protein degradation (Fig.S1a) and that UbVS/ubiquitin promoted degradation of only a limited set

of proteins (Fig.1g). These findings may be explained by the prevalence of non-degradative

ubiquitin conjugation pathways that preferentially consume ubiquitin, limiting ubiquitylation of

degradative substrates. To acquire a better picture of the spectrum of proteins that are ubiquitylated

in extract, we performed a proteomic experiment to identify proteins that become modified with

exogenously added ubiquitin. We treated the extract with 50 μM of HA-ubiquitin or buffer for 30

minutes to allow the incorporation of HA-ubiquitin into ubiquitylated substrates. Immunoblot re-

vealed incorporation of tagged ubiquitin into many proteins (Fig. S3a). We then performed mass

spectrometry analysis of immunopurified HA-ubiquitin from extract and identified 772 proteins

that were enriched for binding to HA-antibody beads relative to empty beads (log2 anti-HA beads

relative to empty beads)> 1, p-value < 0.05) (Fig.S3b). Because this pull-down was carried out in

the presence of high salt (500 mM KCl), these proteins are likely either directly ubiquitylated or

bind ubiquitin with high affinity. The top proteins enriched on the anti-HA beads were UPS com-

ponents such as E1s, E2s and ubiquitin ligases (Fig. S3B) that can form a thioester with HA-ubiq-

uitin, confirming the validity of this approach. Beyond ubiquitylation machinery, proteins involved

in translation, such as ribosomes, translation factors and RNA binding proteins, were the most

frequently identified. Other proteins isolated included protein folding factors, cytoskeletal compo-

nents, proteins involved in DNA replication/repair, mitochondrial proteins, and metabolic en-

zymes. We also identified 9 of the 34 proteins degraded in extract treated with UbVS/ubiquitin,

confirming that these proteins are directly ubiquitylated in extract. Thus, while the proteome is

stable in untreated extract, the ubiquitin conjugation machinery appears highly active in its ability

to modify a large number of proteins with ubiquitin in a manner that does not impact their stability.

Therefore, the failure of ubiquitin addition to stimulate widespread degradation (Fig.S1a) may be

a result of the preferential incorporation into non-degrative substrates.

Together our findings suggest that UbVS treatment induces rapid ubiquitin depletion

principally by promoting ubiquitylation of non-degradative substrates. To identify proteins that

become preferentially ubiquitylated in the presence of UbVS, we profiled global protein ubiq-

uitylation in extract using a multiplexed quantitative di-glycine (diGly) remnant method(Rose et

al. 2016) after addition of ubiquitin or UbVS/ubiquitin to the extract for 30 minutes. Overall, we

identified 883 ubiquitylation sites in 515 proteins, 219 of which were found in the previous HA-

ubiquitin pull down experiment (Fig. S3c), indicating substantial overlap with the two different

approaches. We identified 190 ubiquitylation sites (142 proteins) whose ubiquitylation increased

significantly after UbVS/ubiquitin addition compared to addition of ubiquitin alone (log$_2$ Fold

change>1) (Fig.S3d). As expected, we found 8 ubiquitylation sites belonging to 6 proteins of the

34 that were destabilized by UbVS/ubiquitin addition to the extract (Fig. S3e), confirming again

that these proteins are direct DUB substrates. Several proteins identified have been previously

reported to be DUB substrates, such as the ESCRT complex component CHMP1B (USP8)(Cre-

spo-Yàñez et al. 2018), the ubiquitin ligase SMURF2 (USP15)(Iyengar et al. 2015) and the repli-

cation factor PCNA (USP1)(Huang et al. 2006). However, the majority of the UbVS-sensitive sites

are novel. Because UbVS treatment increases ubiquitylation of a much larger number of proteins

than those that are destabilized, including abundant proteins such ribosome subunits and histones

, these substrates may sequester ubiquitin, limiting protein degradation in extract treated with

UbVS alone.

Confirmation of UbVS-dependent proteasomal degradation of the newly identified sub-

strates with human orthologs

After developing a clearer picture of the overall dynamics of ubiquitylation, deubiquityla-

tion, and protein degradation in extract, we next focused on the 34 substrates that require UbVS-

sensitive DUBs to maintain their stability. Most of these proteins have reported physiological

functions, but only 9 are known UPS substrates (Fig. 2a, Fig. S4a, Table S3). Furthermore, for

only four of them have specific DUBs been identified that control their stability. Two of these

proteins are ubiquitin ligases, MARCH7 and BIRC3, which are known to be protected from deg-

radation by cysteine-protease DUBs in human cells (by USP7 or USP9(Nathan et al. 2008) and

USP19 respectively(Mei et al. 2011)). In addition, we identified Stam and NFX1, which are known

to be deubiquitylated by the cysteine-protease DUBs USP8(Berlin, Schwartz, and Nash 2010) and

USP9 respectively(Chen et al. 2019). The ability of our screen to identify known DUB substrates

suggests that our approach is capable of identifying physiologically relevant proteins whose sta-

bility requires DUB activity.

For the majority of the proteins we identified, DUBs have not been implicated in regulation

of their stability. We identified four additional ubiquitin ligases (MKRN1, RNF138, ZNF598,

UBOX5), whose stability is not known to be dependent on DUB activity. This enrichment for

ubiquitin ligases is again consistent with the fact that DUBs are known to protect them from auto-

ubiquitylation and consequent degradation(W. Kim et al. 2011)·(Ventii and Wilkinson 2008).

Other functional classes of proteins that we identified included transcriptional regulators (GTSF1,

SOX3, SOX15, TGIF2), signaling proteins (GYGIf1, PLK3, WEE2, CSNK1E, BORA, PIM3),

cytoskeletal regulators (CRIPT, WASL, KIAA1430, HN1, ARHGEF19), proteins involved in

RNA processing (FAM32, RNASEH1, RBM18, SYF2, ZMAT22, ZFP36L11, EIF1), DNA dam-

age components (ETAA1, SPRTN), a lipid storage protein (PLIN2), and an uncharacterized pro-

tein (C6ORF132). Together these findings suggest that ongoing deubiquitylation in the *Xenopus*

system is important for maintaining the stability of proteins that regulate a wide variety of biolog-

ical processes.

We looked for possible characteristics sequence of these substrates (Table S3) that could

contribute to their recognition by the UPS. Consistent with the PEST sequence(Rechsteiner and

Rogers 1996) being a signal that promotes ubiquitin and proteasome-dependent degradation, we

found one or multiple PEST sequences in 13 of the 34 hits (Table S3). Because efficient protein

degradation by the proteasome requires unstructured regions in its substrates(Aufderheide et al.

2015), we analyzed their sequences using ProViz(Jehl et al. 2016). We found that 8 substrates

were predicted to be completely or highly disordered whereas 25 had significantly disordered re-

gions consistent with the possibility that they are proteasome substrates (Table S3).

To directly test whether degradation of the identified DUB substrates was proteasome-

dependent, we quantified protein stability in extract treated with UbVS/ubiquitin in the presence

of the proteasome inhibitor MG262 or DMSO (as a control) with a TMT-based quantitative pro-

teomics experiment (Fig. S5a, S5b). We found that 19 proteins became unstable after UbVS/ubiq-

uitin addition (Fig. S5b, S5c), 14 of which were among the 34 substrates identified (Fig. 1g). All

of these proteins were stabilized by proteasome inhibition (Fig. S5b, c). Thus, UbVS-sensitive

DUBs antagonize proteasome-mediated degradation of these proteins.

To verify the findings of the proteomic experiments and directly test whether these proteins

become destabilized by DUB inhibition, we expressed 13 candidate substrates in rabbit reticulo-

cyte lysate and labeled them with [35]S-methionine. We chose ten proteins from our list of 34 can-

didates, as well as three proteins that were destabilized in at least one of the two experiments shown

in Fig.1f. We translated *in vitro* human orthologs, with the exception of PIM3, where we tested

the *Xenopus* protein. Subsequently, we added the translated proteins to untreated extract or to ex-

tract pre-treated with ubiquitin, UbVS or UbVS/ubiquitin and monitored their stability over time.

We found that 12 of 13 proteins were degraded in the presence of UbVS/ubiquitin but were stable

in the other conditions (Fig. 2b, 2c, S4b-d), showing the same pattern that we observed for the

endogenous counterparts. STAM was the only protein that was not destabilized in the presence of

UbVS/ubiquitin, but instead showed strong poly-ubiquitylation (Fig. 2d). As a control, two pro-

teins that were stable in the proteomic experiments (KPNA2 and HSPA8), were also tested and

found to be stable when assayed by this approach (Fig. 2b, 2c) Thus, because we could recapitulate

the behaviour of the endogenous *Xenopus* proteins using human orthologs, the DUB-dependent

regulation of these proteins is likely conserved, as is known already for MARCH7, BIRC3, NFX1

and STAM.

Identification of DUBs sufficient to rescue proteins from degradation in UbVS/ubiquitin-

treated extract

Next, we used these substrates to investigate the role of cysteine-protease DUBs in coun-

teracting their proteasomal degradation. If the proteins degraded in UbVS/ubiquitin-treated extract

are true DUB substrates, addition of recombinant DUBs to the extract should be able to rescue

their degradation. Furthermore, by testing the sufficiency of each DUB to rescue degradation of

these substrates, we can evaluate the activity and specificity of each DUB, even if they normally

function redundantly. We reasoned that the DUBs efficiently targeted by UbVS were most likely

to rescue the stability of the proteins degraded. Thus, we measured the fraction of each DUB la-

beled by UbVS. After we treated the extract with 10 μM HA-UbVS, we subsequently depleted all

the HA-UbVS (and associated proteins) from the extract by immunodepletion with anti-HA anti-

bodies coupled to beads. Using TMT-based quantitative proteomics, we compared the proteins

remaining in extract after immunodepletion of HA-UbVS to the proteins present in extract after

incubation with empty beads. As expected, the proteins depleted were mostly DUBs (Fig. S6a).

We detected 32 cysteine-protease DUBs in undepleted extract and found that 25 of them were

depleted by HA-UbVS (Fig. 3a), with 19 DUBs being depleted by at least 30%. There was no clear

correlation between the reported abundance(Wühr et al. 2014) of the DUBs (Table S1) and their

fractional depletion by HA-UbVS (Fig. 3a). We selected 12 DUBs that were depleted at least 30%

to test if they could rescue substrate degradation in UbVS/ubiquitin treated extract (Fig.3a, in

bold). In addition, we tested USP14 because it can rescue several substrates from proteasomal

degradation(B.-H. Lee et al. 2010a), as well as USP21 because it is a highly active DUB widely

used in in vitro assays.

First, we confirmed the activity of each recombinant DUB with a UbVS-reactivity assay

(Fig. S6b, c). All DUBs tested showed almost complete labelling with UbVS, with the exception

of USP14 that requires activation by the proteasome(B.-H. Lee et al. 2010a; Borodovsky et al.

2001), which was absent from these in vitro assays. We then proceeded to add each recombinant

DUB to extract pre-treated with UbVS/ubiquitin and monitored the stability of ^{35}S-methionine-

labeled ING2, CSNK1E, and CRIPT (Fig.3b). We chose these proteins as model substrates be-

cause they have not been reported to be regulated by DUBs and they have unrelated functions. We

began by assessing all DUBs at the same concentration (800 nM), which is substantially greater

than the endogenous concentration for ten of the fourteen DUBs tested (Fig. S7a). We compared

the ability of each DUB to rescue degradation of the selected substrates, creating a "DUB profile"

that reports the percentage of substrate rescued by each DUB (Fig. 3b). We observed that only

USP7, USP15, USP25 and USP28 could rescue degradation of these substrates (Fig. 3b, Fig. S7b).

Whereas USP28 and USP15 are present at low concentration in extract (4 and 20 nM respectively),

USP25 was not detected in the previous study indicating that its concentration is very low, so its

activity was not further evaluated. USP7 is much more abundant in extract (150 nM), suggesting

that it might be able to rescue degradation of these substrates at its physiological concentration. In

fact, when tested at physiological concentrations, USP7 remained capable of strongly rescuing

CRIPT stability, with partial rescue of ING2 stability (Fig.3c, S7c). In contrast, USP15 and USP28

did not rescue substrate degradation when used at lower concentrations, even though these ex-

ceeded their endogenous concentrations by 5- to 15-fold, respectively (Fig. 3c, S7c). Therefore,

among all the DUBs screened, USP7 seemed unique in its ability to rescue degradation of the

substrates when tested at its physiological concentration. In contrast, USP4, USP9X, OTUD3,

USP8, USP10, and USP21 did not rescue degradation of the three substrates, even though they

were tested well above their endogenous concentrations. We were puzzled by the fact that the two

known proteasome associated DUBs, UCHL5 and USP14, did not affect degradation of the sub-

strates tested. Because they are very abundant in extract, we retested them and we tested UCHL3,

another very abundant DUB in extract, at 5 times their endogenous concentration and we compared

them to USP7. None of these DUBs had an effect comparable to USP7 in this assay (Fig. 3d, S7d).

Because USP7, USP15, USP25, and USP28 were each sufficient to stabilize all three sub-

strates when used at 800 nM concentration, we wondered whether these DUBs were also capable

of restoring ubiquitin availability in UbVS-treated extracts. We treated extract with 10 μM UbVS,

added each of the recombinant DUBs and monitored the charging status of Ube2C-Ub thioester as

a readout of ubiquitin availability. We included USP21 as a positive control, as it broadly deubiq-

uitylates substrates *in vitro*, as well as USP5, which can regenerate free ubiquitin by acting on

unanchored ubiquitin chains(Hadari et al. 1992). As expected, addition of exogenous ubiquitin

fully restored charging of the Ube2C-Ub thioester (Fig.3e), as did addition of USP21, consistent

with its known broad substrate specificity. USP5 did not have any effect, suggesting that free

unanchored ubiquitin chains are not generated at high levels in UbVS-treated extract. Of the 4

DUBs able to rescue substrate degradation, only USP15 was able to fully restore charging of

Ube2C. This finding suggests that USP7, USP25, and USP28 are capable of rescuing substrate

degradation without impacting ubiquitin availability in the extract, whereas USP15 seems to have

a different substrate specificity that allows it to rescue degradation and also restore ubiquitin recy-

cling. On the other hand, despite the ability of USP21 to regenerate free ubiquitin, it is unable to

rescue the degradation of any of the tested substrates.

USP7 broadly rescues protein degradation in UbVS/ubiquitin treated extract

As USP7 seemed to be the most efficient DUB in rescuing degradation of the substrates

tested, we decided to investigate how broad an effect USP7 had in rescuing proteins from degra-

dation in extract treated with UbVS/ubiquitin. We performed a TMT-based quantitative prote-

omics experiment comparing protein stability in extract treated with UbVS/ubiquitin, and in ex-

tract in which USP7, USP9X, or USP4 were added at 800 nM each (Fig. 4a). We tested USP9X

and USP4 as they were targets of HA-UbVS (Fig. 3a) yet were not sufficient to rescue the degra-

dation of the panel of substrates that we tested (Fig 3b). We found that USP7 could rescue 16 of

the 20 proteins degraded in UbVS/ubiquitin (Fig. 4c). However, we did not find any protein stabi-

lized by addition of USP4 or USP9X (Fig.4b, 4c), consistent with the results of our earlier screen

using the panel of [35]S-labeled substrates. The fact that human recombinant USP7 can stabilize

endogenous *Xenopus* proteins suggests again that DUB-dependent regulation of the stability of

these proteins is likely evolutionarily conserved.

Because USP7 broadly rescued substrate degradation in UbVS/ubiquitin-treated extract (Fig. 4b,

4c) and has been reported to be associated with the proteasome(Bousquet-Dubouch et al. 2009;

Besche et al. 2009), USP7 might inhibit the proteasome independent of its catalytic activity, as is

known for USP14(B.-H. Lee et al. 2010a). Thus, we tested if the catalytic activity of USP7 was

required for substrate rescue. Pre-incubation of USP7 with UbVS inhibited rescue of substrate

degradation (Fig. 4d), confirming that deubiquitylating activity of USP7 is required for the rescue.

Second, we investigated if USP7 could inhibit degradation of other known proteasome substrates.

We tested if USP7 can rescue the APC substrates cyclin B1 and securin from proteasome degra-

dation. Surprisingly, we found that USP7 was unable to rescue them from degradation (Fig. 4e),

indicating that USP7 does not generally inhibit the degradation of all substrates of the UPS.

Inhibition of Usp7 is not sufficient to induce protein instability suggesting that DUBs func-

tion redundantly to control protein stability in extract

Because USP7 could stabilize most of the proteins degraded in extracts treated with

UbVS/ubiquitin, we tested if inactivation of USP7 using the recently developed specific inhibitor

XL-188(Lamberto et al. 2017) was sufficient to induce degradation of the substrates that we iden-

tified. We also tested the effect of USP14 inhibition on proteome stability using the specific inhib-

itor IU-47(Boselli et al. 2017) alone or in combination with XL-188, since USP14 inactivation

promotes degradation of some proteasome substrates in vitro and in human cells(B.-H. Lee et al.

2010b). First, we verified these compounds were able to inhibit the endogenous *Xenopus* DUBs.

We pre-incubated extract with the active compounds (IU-47 and XL-188) and the respective inac-

tive derivatives (IU-C and XL-203) for 30 minutes, and then added 10 μM HA-UbVS. As ex-

pected, only the active compounds were able to prevent HA-UbVS labelling of the specific tar-

geted DUB (Fig. S8a). After we verified that the inhibitors were working in extract, we compared

protein stability in extract pre-treated with 10 μM UbVS, 100 μM IU-47 (USP14i), 200 μM XL-

188 (USP7i) or the combination of the latter inhibitors, with TMT-based quantitative proteomics

(Fig. S8b). In all conditions we added ubiquitin to minimize any impact of DUB inhibition on

ubiquitin recycling. In this experiment, we confirmed again that 14 of the 34 candidates (Fig. S8c)

were degraded in UbVS/ubiquitin treated extract. Among these proteins, only HN1 was destabi-

lized after specific USP7 inhibition. There were no other proteins (detected with more than one

peptide) that were destabilized following specific USP7 inhibition. Inhibition of USP14 did not

promote protein instability in *Xenopus* extract, as none of the proteins degraded in UbVS/ubiquitin,

or any other protein, was degraded in IU-47-treated extract (Fig. S8c, d). Inhibition of both DUBs,

as expected, caused the degradation of HN1 that was destabilized by the USP7 inhibitor. In addi-

tion, we observed degradation of three other proteins: DBN1, KHLC1, and Supervillin (SVIL), a

known USP7 substrate(Fang and Luna 2013) (Fig.S8c, bottom). Together these findings suggest

that inhibition of USP7 alone or in combination with USP14 is not sufficient to drive the broader

pattern of protein instability that we observe in extracts treated with UbVS/ubiquitin. Instead, our

results support a model in which multiple DUBs play redundant functions in maintaining stability

of these substrates.

Discussion

Our study offers the first broad picture of the dynamics of ubiquitylation, deubiquitylation, and protein degradation in *Xenopus* extract, a model system that has been used extensively to study the ubiquitylation and degradation of specific substrates. We provide important new insights into the relative roles of UbVS-sensitive DUBs in recycling ubiquitin and in protecting proteome stability. Identifying DUB substrates is challenging due to the fact that multiple DUBs can act redundantly on the same substrates. Here, by broadly inactivating DUBs we circumvented DUB redundancy and, using an integrated set of unbiased proteomic experiments, we identified both degradative and non-degrative targets of DUBs. Using these newly discovered set of physiological substrates whose stability is DUB-dependent, we investigated the ability of a panel specific DUBs to protect substrates from degradation. We revealed that USP7 was uniquely capable of rescuing most proteins from degradation in extract in which DUBs were broadly inhibited. However, since inhibition of USP7 alone was not sufficient to promote substrate degradation, our findings suggest that USP7 functions redundantly with other DUBs. In summary, our study provides the first comprehensive characterization of protein stability and ubiquitin dynamics in *Xenopus* extract, reveals novel DUB substrates and presents a new approach to characterize DUB specificity.

By analyzing protein abundance over time in interphase *Xenopus* extract, we observed that the flux of proteins targeted to the proteasome is low and remarkably resistant to UPS perturbation. Untreated extract showed little change in protein abundance in the absence of protein synthesis, indicating that the proteome is stable. This degree of proteome stability is consistent with the fact that the *Xenopus* egg has completed its growth and sits in a quiescent state until fertilization. Yet at the same time, ubiquitylation and deubiquitylation appear to be highly active. Our findings suggest that the robustness of protein stability to UPS perturbation arises because most ubiquitylation

and deubiquitylation in unperturbed extracts occurs on non-degrative substrates. Addition of ubiq-

uitin alone led to ubiquitin incorporation into hundreds of proteins in a manner that did not impact

their stability. Furthermore, addition of UbVS alone led to alterations in protein ubiquitylation but

caused no change in protein stability, instead causing a rapid depletion of free ubiquitin. This find-

ing highlights the predominant role of UbVS-sensitive DUBs in recycling ubiquitin in extracts.

Included among the non-degradative targets of UbVS-sensitive DUBs, we identified highly abun-

dant proteins such as histones and ribosomes that may serve to sequester ubiquitin to "store" it for

other purposes, such as stress resistance. Consistent with this idea, proteotoxic stresses such as

heat shock or proteasome inhibition that induce a high demand for free ubiquitin, cause a redistri-

bution of ubiquitin from histones to misfolded proteins or to proteins targeted to degradation in

actively growing cells(Dantuma et al. 2006)·(Rose et al. 2016)

Although most ubiquitylation in *Xenopus* extract appears to be non-degradative, our ex-

periments also identified a set of novel degradative substrates whose stability requires ongoing

deubiquitylation by UbVS-sensitive DUBs. DUB-dependent stability of these proteins was re-

vealed only when DUBs were broadly inhibited, and available ubiquitin was restored. We con-

firmed DUB-dependent stability of these substrates with recombinant human proteins, demonstrat-

ing evolutionary conservation. Several substrates are known to be regulated by specific DUBs,

including MARCH7, BIRC3, STAM and NFX1 (Nathan et al. 2008)·(Mei et al. 2011)·(Chen et al.

2019)·(Cai et al. 2015). Furthermore, PIM3 and WASL have not been connected previously to

DUBs, but are similar in sequence to well-known DUB substrates (PIM2(Kategaya et al. 2017)

and WASH(Hao et al. 2015) respectively). Together, these findings validate that our approach can

identify established DUB substrates. Still, the vast majority of the proteins we identified are not

known DUB substrates, demonstrating the novelty of our findings. Our candidate substrates were

enriched for ubiquitin ligases (n=12), extending our understanding of the importance of DUBs in

counteracting their auto-ubiquitylation and consequent degradation(W. Kim et al. 2011)·(Ventii

and Wilkinson 2008). Makorin Ring Finger protein 1 (MKRN1) ubiquitylates p53 and p21 target-

ing them to proteasomal degradation(E.-W. Lee et al. 2009). Yet despite intensive study, a role for

DUBs in controlling MKRN1 stability has not been described. Our data suggest that DUB-depend-

ent stability of MKRN1 could be an important control mechanism, as it is for MDM2, another p53

ubiquitin ligase(Ranaweera and Yang 2013).

Beyond ubiquitin ligases, we identified substrates with a broad range of functions. We

found multiple transcription factors (n=7) and proteins involved in RNA metabolism (n=12). A

number of substrates, such as SPRTN, ETAA1, and the Casein Kinases ε and δ, have been exten-

sively studied, so it is surprising that DUB-dependent control of their stability has not yet been

reported. SPRTN and ETAA1 are both involved in DNA replication/damage and are essential for

maintaining genome stability(Vaz et al. 2016)·(Haahr et al. 2016). Casein Kinases ε and δ are

versatile proteins that participate in multiple processes such as cell cycle control, spindle organi-

zation, and circadian rhythm(Schittek and Sinnberg 2014). Our study suggests that DUBs could

modulate their degradation by the UPS as mechanism to regulate their activity.

This study also describes a new approach for studying the specificity and activity of DUBs

in a system in which physiological rates of substrate ubiquitylation are maintained. Our approach

measures effects on degradation rather than deubiquitylation, providing a new way of assessing

the ability of DUBs to counter proteasome-mediated degradation. The most unexpected finding

to emerge from this analysis was the ability of USP7 to rescue a large number of substrates from

degradation. Despite this broad activity, USP7 was unable to rescue APC/C substrates from

degradation, suggesting that USP7 is not a general inhibitor of proteasome activity. This result was

not due to a lack of activity of USP7 in mitosis because we verified that USP7 was still fully active

in mitosis (data not shown). Furthermore, even though USP7 was able to broadly rescue substrates

from degradation, USP7 was unable to rescue ubiquitin depletion in UbVS-treated extract. To-

gether these findings indicate that USP7 has a defined substrate specificity that seems targeted

towards degradative versus non-degradative substrates. Interestingly, we also observed exactly the

opposite pattern of specificity with USP21(Yu Ye et al. 2011), which was able to rescue ubiquitin

availability but was unable to rescue degradation of our panel of substrates.

What explains the broad ability of USP7 to antagonize proteasomal degradation? USP7 has

been highly studied and has attracted attention as a pharmacological target, as it regulates the sta-

bility of the tumor suppressor p53 and its regulator MDM2(Sheng et al. 2006). Among the DUBs

we tested, USP7 has the greatest number of published substrates, most of which are targeted to the

proteasome(R. Q. Kim and Sixma 2017). Among the substrates that we identified, a few such as

MARCH7, SVIL, NEK2A, and TRIP12 are known targets of USP7(Nathan et al. 2008)·(Fang and

Luna 2013)·(Franqui-Machin et al. 2018)·(Cai et al. 2015) whereas two others, PIM3 and WASL,

are likely USP7 targets since it deubiquitylates related proteins (PIM2(Kategaya et al. 2017) and

WASH(Hao et al. 2015) respectively) in human cells. USP7 has been reported to associate with

the proteasome(Bousquet-Dubouch et al. 2009)·(Besche et al. 2009), where it could intercept pro-

teins that are targeted for degradation. However, the physiological significance and mechanism of

this association has not been carefully studied. USP7 may have a broad ability to rescue substrates

from degradation because it directly binds a wide range of proteins through recognition of common

degenerate motifs (P/AxxS and Kxxx/KxK) that are found in many proteins(R. Q. Kim and Sixma

2017). Another plausible explanation is that USP7 is kinetically sufficient to counteract the rates

of ubiquitylation of ubiquitin ligases that target substrates for degradation. This idea is supported

by USP7 being the most active DUB in Ub-AMC assays among a panel of 12 cysteine DUBs

tested(Faesen et al. 2011).

Our study also highlights the important impact and challenge of functional redundancy

when studying DUB activity and specificity. In the *Xenopus* system, we observed that inhibition

of USP7 alone was not sufficient to induce degradation of substrates, despite USP7's ability to

broadly rescue substrates from degradation. This finding supports the notion that DUBs function

redundantly to maintain the stability of the proteins in this system. Because degradation is an irre-

versible step in the protein lifecycle, DUB redundancy may help set a higher threshold for ubiq-

uitylation required for degradation of a protein, beyond what is required for direct recognition by

the proteasome. On the other hand, DUB redundancy poses a serious technical challenge to fully

understand the role of DUBs and identify their substrates, especially using genetic approaches.

Our work highlights how unmasking DUB redundancy it is key not only for discovery of novel

DUB substrates but also to fully understand functional specificity of these important enzymes.

Materials and Methods

Gene nomenclature

The human gene symbols have been used for consistency and simplicity in the manuscript.

Reagents

Commercial antibodies used for Western blotting analysis were as follow: anti-Ube2C (A-650, Boston Biochem), anti-ubiquitin (P4D1; sc-8017; Santa Cruz Biotechnology), anti-HA-peroxidase (3F10; Roche), anti-USP7 (A300-033A; Bethyl). Secondary antibodies used included anti-rabbit IgG-HRP (NA934V) and anti-mouse IgG-HRP (NA931V) from GE Healthcare. Chemicals used in this study included Ubiquitin vinyl sulfone (U-202), MG262 (I-120), Ubiquitin vinyl sulfone tagged with HA (U-212) and HA-ubiquitin (U-110) purchased from Boston Biochem. For immunoprecipitation anti-HA-Agarose beads (A2095) from Sigma were used. For the diGly experiment immunoaffinity beads from Cell Signaling were used (5562). Ubiquitin (U-100H) and recombinant DUBs were provided by Boston Biochem (E-519, USP7; E-520, USP8; E-320, USP5; E-325, UCHL3; E-325, UCHL5; E-546, USP25; E-552, USP9x; E-570, USP28; E-574, OTUD3; E-592, USP10; E-594, USP15; E-596, USP4).

Preparation of *Xenopus laevis* egg extract

Interphase extract was prepared as previously described[Murray AW 1991] but using 2 µg/ml calcium ionophore (A23187, Calbiochem) for egg activation. Entry into mitosis was induced by addition of 1 µM non-degradable cyclin B (MBP-Δ90) as previously described[Zeng et al. 2010].

HA-UbVS immunoprecipitation

Following treatment with 10 µM HA-UbVS (10 minutes at 24°C), *Xenopus* extract was diluted three times with XB buffer (10 mM potassium HEPES pH 7.7, 500 mM KCl, 0.1 mM CaCl2, 1 mM MgCl$_2$, 0.5% NP40) and incubated with anti-HA beads or empty beads at 4°C (1h and 30 minutes). At the end of the incubation, beads were washed three times with XB buffer, SDS sample buffer was added and samples were subjected to SDS gel electrophoresis. Samples were processed according to the GeLC-MS/MS strategy(Paulo 2016).

HA-UbVS immunodepletion

Xenopus extract was pre-treated with 10 µM HA-UbVS (10 minutes at 24°C) diluted three times with XB buffer and incubated with anti-HA beads or empty beads overnight at 4°C. Super-natants from the beads were collected and treated for Mass Spectrometry analysis by SL-TMT method (described below).

HA-ubiquitin immunoprecipitation

Following treatment with 50 µM HA-ubiquitin (30 minutes at 24°C), *Xenopus* extract was diluted three times with XB buffer (10 mM potassium HEPES pH 7.7, 500 mM KCl, 0.1 mM CaCl2, 1 mM MgCl$_2$, 0.5% NP40, Pierce proteases inhibitor tablet and NEM 5mM) and incubated with anti-HA beads or empty beads at 4°C (2h). At the end of the incubation, beads were washed four times with XB buffer and elute with the HA peptide (two times). Samples were processed for Mass Spectrometry analysis by SL-TMT method (described below).

Immunoprecipitation of diGly-containing peptides

Dried peptides (2 mg of proteins) were resuspended in IAP buffer [50 mM MOPS (pH 7.2), 10 mM sodium phosphate and 50 mM NaCl] and centrifuged at top speed (5 min). After that the supernatants were added to the diGly resin (Cell Signaling Technology) and incubated for 2 hr at 4°C. After that, beads were washed three times with ice-cold IAP buffer and twice with PBS. The diGly peptides were eluted twice with 0.15% TFA, desalted using homemade StageTips and dried via vacuum centrifugation. Peptides were immunoprecipitated twice. Samples were processed for Mass Spectrometry analysis by SL-TMT method (described below).

Degradation of ^{35}S-methionine labeled substrates

Extract was pre-treated with UbVS (10 μM) for 10 min at 24°C before addition of ubiquitin (50 μM) and substrates. The preincubation time with UbVS was extended to 30 minutes when recombinant DUBs were added to the extract. Pre-treatment of extract with MG262 (200 μM) or specific DUB inhibitors was done at 24°C for 30 minutes. Substrates were expressed and labeled using ^{35}S-methionine (Perkin Elmer, NEG709A500UC) with the TNT kit (Promega: L1770). Each substrate was amplified with primers by PCR to allow T7-dependent transcription of the PCR product or transcribed directly if plasmids contained a T7 promoter. The translation reaction mix was added to the *Xenopus* extract at 8% final volume. Samples of the reactions were collected at the indicated time, quenched with sodium dodecyl sulfate (SDS) sample buffer, and processed for SDS gel electrophoresis and phosphor imaging (Bio-Rad PMI). Quantification was performed using Quantity One software (Bio-Rad).

UbVS labeling of recombinant DUBs to verify their activity

Each DUB (1μM) was incubated with saturating amount of UbVS or HA-UbVS as indi-

cated (1 hour at 30°C). Reactions were stopped with addition of sodium dodecyl sulfate (SDS)

sample buffer and run on SDS-PAGE. After SDS-PAGE, the gel was stained with Coomassie

Brilliant Blue.

Streamlined Tandem Mass Tag Protocol

<u>Peptide extraction and digestion</u>

The TMT labeling protocol and mass spectrometry analysis were based on the SL-TMT

sample preparation strategy(Navarrete-Perea et al. 2018). *Xenopus* extract was collected, resus-

pended in the appropriate buffer (1% SDS, 5 mM DTT, 50 mM Tris pH 8.8 and Pierce protease

inhibitor tablet) and flash frozen in liquid nitrogen. Methanol–chloroform precipitation was per-

formed. Four volumes of methanol were added to each sample and briefly vortexed. One volume

chloroform was added to the samples and vortexed again. Lastly, three volumes water was added

and vortexed. The samples were centrifuged (5 minutes, 14000 RPM) and subsequently washed

twice with cold methanol. Samples were resuspended in 200 mM EPPS, pH 8.5 and digested over-

night at 24°C with Lys-C protease (Wako Chemicals). Later samples were incubated for 6 hours

at 37°C for digestion by trypsin protease (Pierce Biotechnology).

<u>Isobaric labeling and fractionation</u>

Tandem mass tag (TMT) isobaric reagents (Thermo Fisher Scientific) were resuspended in

anhydrous acetonitrile (final concentration of 20 μg/μL). 10 μL of the labeling reagents plus 30

μL of acetonitrile was added to the peptides obtained by the previous digestions (~100 μg). After

incubation at room temperature (90 minutes), the reaction was quenched using hydroxylamine to

a final concentration of 0.3% (v/v). The TMT-labeled samples were pooled at a 1:1 ratio across

all the samples. Fractions were fractionated off-line using basic pH reversed phase chromatog-

raphy (BPRP). Fractions were pooled into 6 or 12 super-fractions which were acidified with formic

acid to a final concentration of 1%. The pooled peptides were desalted using homemade StageTip,

dried with vacuum centrifugation, and reconstituted in 5% acetonitrile and 5% formic acid for LC-

MS/MS processing.

<u>Peptide detection, identification and quantification</u>

All samples were analyzed on an Orbitrap Fusion mass spectrometer (Thermo Fisher Sci-

entific) coupled to a Proxeon EASY-nLC 1000 liquid chromatography (LC) pump (Thermo Fisher

Scientific). Peptides were separated on a column packed with 35 cm of Accucore C18 resin (2.6

µm, 150 Å, Thermo Fisher Scientific). The column had a 100 µm inner diameter microcapillary.

For each experiment, 2 µg of peptides were loaded onto this column. Peptides were separated,

using a flow rate of 450 nL/min., with a 150-minute gradient of 3 to 25% acetonitrile in 0.125%

formic acid. Each analysis used an MS3-based TMT method, which it is known to reduce ion

interference if compared to MS2 quantification. The scan sequence starts with an MS1 spectrum

(Orbitrap analysis, resolution 120,000, 400−1400 Th, automatic gain control (AGC) target 5E5,

maximum injection time 100 ms). For subsequent MS2/MS3 analysis, only the top 10 precursors

were selected. MS2 analysis included: collision-induced dissociation (CID), quadrupole ion trap

analysis, automatic gain control (AGC) 2E4, NCE (normalized collision energy) 35, q-value 0.25,

maximum injection time 120 ms), and isolation window at 0.7. After we acquire each MS2 spec-

trum, we collected an MS3 spectrum in which multiple MS2 fragment ions were captured in the

MS3 precursor population with isolation waveforms using multiple frequency notches. MS3

precursors were fragmented by HCD and analyzed using the Orbitrap (NCE 65, AGC 1.5E5, max-

imum injection time 150 ms, resolution was 50,000 at 400 Th). For MS3 analysis, we used charge

state-dependent isolation windows: For charge state z=2, the isolation window was set at 1.3 Th,

for z=3 at 1 Th, for z=4 at 0.8 Th, and for z=5 at 0.7 Th. Collected Spectra were processed using

a Sequest-based software pipeline.   Spectra were converted to mzXML using MS Convert

(Adusumilli and Mallick 2017). Database searching included all the entries from the PHROG da-

tabas. This database includes many lower abundant proteins and multiple splice isoforms (not pre-

sent in other databases). In the original study, around 11,000 proteins were identified(Wühr et al.

2014). Thus, our study (with ~8000 proteins) represents an acceptable coverage of the *Xenopus*

proteome. This database was concatenated with one composed of the sequences in the reversed

order. Searches were performed using a 50 Th precursor ion tolerance and the product ion tolerance

was set to 0.9 Da. Oxidation of methionine residues (+15.995 Da) and, where indicated. Peptide-

spectrum matches (PSMs) were adjusted to a 1% false discovery rate (FDR). PSM filtering was

performed using a linear discriminant analysis, as described previously and assembled to a final

protein-level FDR of 1%. Proteins were quantified by summing reporter ion counts across all

matching PSMs(McAlister et al. 2012). Reporter ion intensities were adjusted to correct for the

isotopic impurities of the different TMT reagents according to manufacturer specifications. The

peptides signal-to-noise (S/N) measurements assigned to each protein were summed and normal-

ized so that the sum of the signal for all proteins in each channel was equivalent, thereby account-

ing for equal protein loading. Lastly, each protein was scaled such that the summed signal-to-noise

for that protein across all channels was 100, thereby generating a relative abundance (RA) meas-

urement.

TMT Mass-spectrometry analysis

A two-sided Student's t-test was used as a measure of statistical confidence of the observed $\log_2$ fold change. Selected candidates met both thresholds (Fold Change $\leq$ -1.5 and p-value<0.05) in the experiments. For Figure 1g, the candidates were included if at least one peptide was identified in both of the experiments. For subsequent figures, where a single experiment was performed, candidates were included only if they were detected with at least 2 different peptides. When multiple isoforms of the same protein decreased in UbVS/ubiquitin, only the isoform with more peptides was selected (for simplicity isoforms are not indicated in the figures).

References

Adusumilli, Ravali, and Parag Mallick. 2017. "Data Conversion with ProteoWizard MsConvert."
  In *Methods in Molecular Biology*, 1550:339–68. https://doi.org/10.1007/978-1-4939-6747-
  6_23.

Aufderheide, Antje, Pia Unverdorben, Wolfgang Baumeister, and Friedrich Förster. 2015.
  "Structural Disorder and Its Role in Proteasomal Degradation." *FEBS Letters* 589 (19):
  2552–60. https://doi.org/10.1016/J.FEBSLET.2015.07.034.

Beckley, Janel R., Jun-Song Chen, Yanling Yang, Junmin Peng, and Kathleen L. Gould. 2015.
  "A Degenerate Cohort of Yeast Membrane Trafficking DUBs Mediates Cell Polarity and
  Survival." *Molecular & Cellular Proteomics* 14 (12): 3132–41.
  https://doi.org/10.1074/mcp.M115.050039.

Berlin, Ilana, Heather Schwartz, and Piers D. Nash. 2010. "Regulation of Epidermal Growth
  Factor Receptor Ubiquitination and Trafficking by the USP8·STAM Complex." *Journal of
  Biological Chemistry* 285 (45): 34909–21. https://doi.org/10.1074/jbc.M109.016287.

Besche, Henrike C., Wilhelm Haas, Steven P. Gygi, and Alfred L. Goldberg. 2009. "Isolation of
  Mammalian 26S Proteasomes and P97/VCP Complexes Using the Ubiquitin-like Domain
  from HHR23B Reveals Novel Proteasome-Associated Proteins †." *Biochemistry* 48 (11):
  2538–49. https://doi.org/10.1021/bi802198q.

Borodovsky, A, B M Kessler, R Casagrande, H S Overkleeft, K D Wilkinson, and H L Ploegh.
  2001. "A Novel Active Site-Directed Probe Specific for Deubiquitylating Enzymes Reveals
  Proteasome Association of USP14." *The EMBO Journal* 20 (18): 5187–96.
  https://doi.org/10.1093/emboj/20.18.5187.

Boselli, Monica, Byung-Hoon Lee, Jessica Robert, Miguel A Prado, Sang-Won Min, Chialin

Cheng, M Catarina Silva, et al. 2017. "An Inhibitor of the Proteasomal Deubiquitinating Enzyme USP14 Induces Tau Elimination in Cultured Neurons." *The Journal of Biological Chemistry* 292 (47): 19209–25. https://doi.org/10.1074/jbc.M117.815126.

Bousquet-Dubouch, Marie-Pierre, Emilie Baudelet, Frédéric Guérin, Mariette Matondo, Sandrine Uttenweiler-Joseph, Odile Burlet-Schiltz, and Bernard Monsarrat. 2009. "Affinity Purification Strategy to Capture Human Endogenous Proteasome Complexes Diversity and to Identify Proteasome-Interacting Proteins." *Molecular & Cellular Proteomics : MCP* 8 (5): 1150–64. https://doi.org/10.1074/mcp.M800193-MCP200.

Bremm, Anja, Stefan M V Freund, and David Komander. 2010. "Lys11-Linked Ubiquitin Chains Adopt Compact Conformations and Are Preferentially Hydrolyzed by the Deubiquitinase Cezanne." *Nature Structural & Molecular Biology* 17 (8): 939–47. https://doi.org/10.1038/nsmb.1873.

Cai, Jia-Bin, Guo-Ming Shi, Zhao-Ru Dong, Ai-Wu Ke, Hong-Hui Ma, Qiang Gao, Zao-Zhuo Shen, et al. 2015. "Ubiquitin-Specific Protease 7 Accelerates P14 [ARF] Degradation by Deubiquitinating Thyroid Hormone Receptor-Interacting Protein 12 and Promotes Hepatocellular Carcinoma Progression." *Hepatology* 61 (5): 1603–14. https://doi.org/10.1002/hep.27682.

Chau, V, J W Tobias, A Bachmair, D Marriott, D J Ecker, D K Gonda, and A Varshavsky. 1989. "A Multiubiquitin Chain Is Confined to Specific Lysine in a Targeted Short-Lived Protein." *Science (New York, N.Y.)* 243 (4898): 1576–83. https://doi.org/10.1126/science.2538923.

Chen, Xiangling, Dayun Lu, Jing Gao, Hongwen Zhu, Yanting Zhou, Daming Gao, and Hu Zhou. 2019. "Identification of a USP9X Substrate NFX1-123 by SILAC-Based Quantitative Proteomics." *Journal of Proteome Research* 18 (6): 2654–65.

https://doi.org/10.1021/acs.jproteome.9b00139.

Clague, Michael J., Igor Barsukov, Judy M. Coulson, Han Liu, Daniel J. Rigden, and Sylvie

Urbé. 2013. "Deubiquitylases from Genes to Organism." *Physiological Reviews* 93 (3):

1289–1315. https://doi.org/10.1152/physrev.00002.2013.

Clague, Michael J., Judy M. Coulson, and Sylvie Urbé. 2012. "Cellular Functions of the DUBs."

J Cell Sci 125 (2): 277–86. https://doi.org/10.1242/JCS.090985.

Clague, Michael J., Claire Heride, and Sylvie Urbé. 2015. "The Demographics of the Ubiquitin

System." *Trends in Cell Biology* 25 (7): 417–26. https://doi.org/10.1016/j.tcb.2015.03.002.

Crespo-Yàñez, Xènia, Carmen Aguilar-Gurrieri, Anne Claire Jacomin, Agnès Journet, Magda

Mortier, Emmanuel Taillebourg, Emmanuelle Soleilhac, Winfried Weissenhorn, and Marie

Odile Fauvarque. 2018. "CHMP1B Is a Target of USP8/UBPY Regulated by Ubiquitin

during Endocytosis." *PLoS Genetics* 14 (6). https://doi.org/10.1371/journal.pgen.1007456.

Dantuma, Nico P., Tom A.M. Groothuis, Florian A. Salomons, and Jacques Neefjes. 2006. "A

Dynamic Ubiquitin Equilibrium Couples Proteasomal Activity to Chromatin Remodeling."

Journal of Cell Biology 173 (1): 19–26. https://doi.org/10.1083/jcb.200510071.

Dimova, Nevena V., Nathaniel A. Hathaway, Byung-Hoon Lee, Donald S. Kirkpatrick, Marie

Lea Berkowitz, Steven P. Gygi, Daniel Finley, and Randall W. King. 2012. "APC/C-

Mediated Multiple Monoubiquitylation Provides an Alternative Degradation Signal for

Cyclin B1." *Nature Cell Biology* 14 (2): 168–76. https://doi.org/10.1038/ncb2425.

Faesen, Alex C, Mark P A Luna-Vargas, Paul P Geurink, Marcello Clerici, Remco Merkx,

Willem J van Dijk, Dharjath S Hameed, Farid El Oualid, Huib Ovaa, and Titia K Sixma.

2011. "The Differential Modulation of USP Activity by Internal Regulatory Domains,

Interactors and Eight Ubiquitin Chain Types." *Chemistry & Biology* 18 (12): 1550–61.

715  https://doi.org/10.1016/j.chembiol.2011.10.017.

716 Fang, Zhiyou, and Elizabeth J Luna. 2013. "Supervillin-Mediated Suppression of P53 Protein

717  Enhances Cell Survival." *The Journal of Biological Chemistry* 288 (11): 7918–29.

718  https://doi.org/10.1074/jbc.M112.416842.

719 Field, Christine M., and Timothy J. Mitchison. 2018. "Assembly of Spindles and Asters in

720  Xenopus Egg Extracts." *Cold Spring Harbor Protocols* 2018 (6): 440–48.

721  https://doi.org/10.1101/pdb.prot099796.

722 Flierman, Dennis, Gerbrand J van der Heden van Noort, Reggy Ekkebus, Paul P Geurink, Tycho

723  E T Mevissen, Manuela K Hospenthal, David Komander, and Huib Ovaa. 2016. "Non-

724  Hydrolyzable Diubiquitin Probes Reveal Linkage-Specific Reactivity of Deubiquitylating

725  Enzymes Mediated by S2 Pockets." *Cell Chemical Biology* 23 (4): 472–82.

726  https://doi.org/10.1016/j.chembiol.2016.03.009.

727 Franqui-Machin, Reinaldo, Mu Hao, Hua Bai, Zhimin Gu, Xin Zhan, Hasem Habelhah, Yogesh

728  Jethava, et al. 2018. "Destabilizing NEK2 Overcomes Resistance to Proteasome Inhibition

729  in Multiple Myeloma." *The Journal of Clinical Investigation* 128 (7): 2877–93.

730  https://doi.org/10.1172/JCI98765.

731 Glotzer, M, A W Murray, and M W Kirschner. 1991. "Cyclin Is Degraded by the Ubiquitin

732  Pathway." *Nature* 349 (6305): 132–38. https://doi.org/10.1038/349132a0.

733 Haahr, Peter, Saskia Hoffmann, Maxim A. X. Tollenaere, Teresa Ho, Luis Ignacio Toledo,

734  Matthias Mann, Simon Bekker-Jensen, Markus Räschle, and Niels Mailand. 2016.

735  "Activation of the ATR Kinase by the RPA-Binding Protein ETAA1." *Nature Cell Biology*

736  18 (11): 1196–1207. https://doi.org/10.1038/ncb3422.

737 Hadari, T, J V Warms, I A Rose, and A Hershko. 1992. "A Ubiquitin C-Terminal Isopeptidase

That Acts on Polyubiquitin Chains. Role in Protein Degradation." *The Journal of Biological*

Chemistry 267 (2): 719–27. http://www.ncbi.nlm.nih.gov/pubmed/1309773.

Hao, Yi-Heng, Michael D. Fountain, Klementina Fon Tacer, Fan Xia, Weimin Bi, Sung-Hae L.

Kang, Ankita Patel, et al. 2015. "USP7 Acts as a Molecular Rheostat to Promote WASH-

Dependent Endosomal Protein Recycling and Is Mutated in a Human Neurodevelopmental

Disorder." *Molecular Cell* 59 (6): 956–69. https://doi.org/10.1016/j.molcel.2015.07.033.

Hershko, A, and Ciechanover A. 1998. "The Ubiquitin System." *Annu. Rev. Biochem.* 67: 425–

79.

Huang, Tony T., Sebastian M.B. Nijman, Kanchan D. Mirchandani, Paul J. Galardy, Martin A.

Cohn, Wilhelm Haas, Steven P. Gygi, Hidde L. Ploegh, René Bernards, and Alan D.

D'Andrea. 2006. "Regulation of Monoubiquitinated PCNA by DUB Autocleavage." *Nature*

Cell Biology 8 (4): 339–47. https://doi.org/10.1038/ncb1378.

Iyengar, Prasanna Vasudevan, Patrick Jaynes, Laura Rodon, Dilraj Lama, Kai Pong Law, Yoon

Pin Lim, Chandra Verma, Joan Seoane, and Pieter Johan Adam Eichhorn. 2015. "USP15

Regulates SMURF2 Kinetics through C-Lobe Mediated Deubiquitination." *Scientific*

Reports 5 (October). https://doi.org/10.1038/srep14733.

Jackson, Stephen P., and Daniel Durocher. 2013. "Regulation of DNA Damage Responses by

Ubiquitin and SUMO." *Molecular Cell* 49 (5): 795–807.

https://doi.org/10.1016/j.molcel.2013.01.017.

Jehl, Peter, Jean Manguy, Denis C. Shields, Desmond G. Higgins, and Norman E. Davey. 2016.

"ProViz—a Web-Based Visualization Tool to Investigate the Functional and Evolutionary

Features of Protein Sequences." *Nucleic Acids Research* 44 (W1): W11–15.

https://doi.org/10.1093/nar/gkw265.

Kategaya, Lorna, Paola Di Lello, Lionel Rougé, Richard Pastor, Kevin R Clark, Jason Drummond, Tracy Kleinheinz, et al. 2017. "USP7 Small-Molecule Inhibitors Interfere with Ubiquitin Binding." *Nature* 550 (7677): 534–38. https://doi.org/10.1038/nature24006.

Kim, Robbert Q., and Titia K. Sixma. 2017. "Regulation of USP7: A High Incidence of E3 Complexes." *Journal of Molecular Biology* 429 (22): 3395–3408. https://doi.org/10.1016/j.jmb.2017.05.028.

Kim, Woong, Eric J Bennett, Edward L Huttlin, Ailan Guo, Jing Li, Anthony Possemato, Mathew E Sowa, et al. 2011. "Systematic and Quantitative Assessment of the Ubiquitin-Modified Proteome." *Molecular Cell* 44 (2): 325–40. https://doi.org/10.1016/j.molcel.2011.08.025.

Komander, David, and Michael Rape. 2012. "The Ubiquitin Code." *Annual Review of Biochemistry* 81 (1): 203–29. https://doi.org/10.1146/annurev-biochem-060310-170328.

Kwon, Seul-Ki, Madhuri Saindane, and Kwang-Hyun Baek. 2017. "P53 Stability Is Regulated by Diverse Deubiquitinating Enzymes." *Biochimica et Biophysica Acta (BBA) - Reviews on Cancer* 1868 (2): 404–11. https://doi.org/10.1016/j.bbcan.2017.08.001.

Lamberto, Ilaria, Xiaoxi Liu, Hyuk-Soo Seo, Nathan J. Schauer, Roxana E. Iacob, Wanyi Hu, Deepika Das, et al. 2017. "Structure-Guided Development of a Potent and Selective Non-Covalent Active-Site Inhibitor of USP7." *Cell Chemical Biology* 24 (12): 1490-1500.e11. https://doi.org/10.1016/j.chembiol.2017.09.003.

Lauwers, Elsa, Christophe Jacob, and Bruno André. 2009. "K63-Linked Ubiquitin Chains as a Specific Signal for Protein Sorting into the Multivesicular Body Pathway." *The Journal of Cell Biology* 185 (3): 493–502. https://doi.org/10.1083/jcb.200810114.

Lee, Byung-Hoon, Min Jae Lee, Soyeon Park, Dong-Chan Oh, Suzanne Elsasser, Ping-Chung

Chen, Carlos Gartner, et al. 2010 "Enhancement of Proteasome Activity by a Small-

Molecule Inhibitor of USP14." *Nature* 467 (7312): 179-84

https://doi.org/10.1038/nature09299.

Lee, Eun-Woo, Min-Sik Lee, Suzanne Camus, Jaewang Ghim, Mi-Ran Yang, Wonkyung Oh,

Nam-Chul Ha, David P Lane, and Jaewhan Song. 2009. "Differential Regulation of P53 and

P21 by MKRN1 E3 Ligase Controls Cell Cycle Arrest and Apoptosis." *The EMBO Journal*

28 (14): 2100–2113. https://doi.org/10.1038/emboj.2009.164.

Ma, Jianhong, John D. Martin, Yu Xue, Leng A. Lor, Karen M. Kennedy-Wilson, Robert H.

Sinnamon, Thau F. Ho, et al. 2010. "C-Terminal Region of USP7/HAUSP Is Critical for

Deubiquitination Activity and Contains a Second Mdm2/P53 Binding Site." *Archives of*

Biochemistry and Biophysics 503 (2): 207–12. https://doi.org/10.1016/j.abb.2010.08.020.

McAlister, Graeme C, Edward L Huttlin, Wilhelm Haas, Lily Ting, Mark P Jedrychowski, John

C Rogers, Karsten Kuhn, et al. 2012. "Increasing the Multiplexing Capacity of TMTs Using

Reporter Ion Isotopologues with Isobaric Masses." *Analytical Chemistry* 84 (17): 7469–78.

https://doi.org/10.1021/ac301572t.

Mei, Yide, Allison Alcivar Hahn, Shimin Hu, and Xiaolu Yang. 2011. "The USP19

Deubiquitinase Regulates the Stability of C-IAP1 and c-IAP2." *The Journal of Biological*

Chemistry 286 (41): 35380–87. https://doi.org/10.1074/jbc.M111.282020.

Mevissen, Tycho E T, and David Komander. 2017. "Mechanisms of Deubiquitinase Specificity

and Regulation" 86: 159–92. https://doi.org/10.1146/annurev-biochem.

Murray AW. 1991. "Cell Cycle Extracts." *Methods Cell Biol* 36: 581–605.

https://www.ncbi.nlm.nih.gov/pubmed/?term=Murray+Cell+cycle+extracts+1991.

Nathan, James A., Soma Sengupta, Stephen A. Wood, Arie Admon, Gabriel Markson, Chris

Sanderson, and Paul J. Lehner. 2008. "The Ubiquitin E3 Ligase MARCH7 Is Differentially

Regulated by the Deubiquitylating Enzymes USP7 and USP9X." *Traffic* 9 (7): 1130–45.

https://doi.org/10.1111/j.1600-0854.2008.00747.x.

Navarrete-Perea, José, Qing Yu, Steven P. Gygi, and Joao A. Paulo. 2018. "Streamlined Tandem

Mass Tag (SL-TMT) Protocol: An Efficient Strategy for Quantitative (Phospho)Proteome

Profiling Using Tandem Mass Tag-Synchronous Precursor Selection-MS3." *Journal of*

Proteome Research 17 (6): 2226–36. https://doi.org/10.1021/acs.jproteome.8b00217.

Paulo, Joao A. 2016. "Sample Preparation for Proteomic Analysis Using a GeLC-MS/MS

Strategy." *Journal of Biological Methods* 3 (3): 45. https://doi.org/10.14440/jbm.2016.106.

Poot, Stefanie A.H. de, Geng Tian, and Daniel Finley. 2017. "Meddling with Fate: The

Proteasomal Deubiquitinating Enzymes." *Journal of Molecular Biology* 429 (22): 3525–45.

https://doi.org/10.1016/j.jmb.2017.09.015.

Ranaweera, Ruchira S, and Xiaolu Yang. 2013. "Auto-Ubiquitination of Mdm2 Enhances Its

Substrate Ubiquitin Ligase Activity." *The Journal of Biological Chemistry* 288 (26):

18939–46. https://doi.org/10.1074/jbc.M113.454470.

Rechsteiner, M, and S W Rogers. 1996. "PEST Sequences and Regulation by Proteolysis."

Trends in Biochemical Sciences 21 (7): 267–71.

http://www.ncbi.nlm.nih.gov/pubmed/8755249.

Richter, J D, and L D Smith. n.d. "Reversible Inhibition of Translation by Xenopus Oocyte-

Specific Proteins." *Nature* 309 (5966): 378–80. Accessed June 9, 2019.

http://www.ncbi.nlm.nih.gov/pubmed/6727992.

Rose, Christopher M., Marta Isasa, Alban Ordureau, Miguel A. Prado, Sean A. Beausoleil,

Mark P. Jedrychowski, Daniel J. Finley, J. Wade Harper, and Steven P. Gygi. 2016.

830 "Highly Multiplexed Quantitative Mass Spectrometry Analysis of Ubiquitylomes." *Cell*

831 *Systems* 3 (4): 395-403.e4. https://doi.org/10.1016/j.cels.2016.08.009.

832 Schittek, Birgit, and Tobias Sinnberg. 2014. "Biological Functions of Casein Kinase 1 Isoforms

833 and Putative Roles in Tumorigenesis." *Molecular Cancer* 13 (1): 231.

834 https://doi.org/10.1186/1476-4598-13-231.

835 Sheng, Yi, Vivian Saridakis, Feroz Sarkari, Shili Duan, Tianne Wu, Cheryl H Arrowsmith, and

836 Lori Frappier. 2006. "Molecular Recognition of P53 and MDM2 by USP7/HAUSP." *Nature*

837 *Structural & Molecular Biology* 13 (3): 285–91. https://doi.org/10.1038/nsmb1067.

838 Soncini, Chiara, Ingrid Berdo, and Giulio Draetta. 2001. "Ras–GAP SH3 Domain Binding

839 Protein (G3BP) Is a Modulator of USP10, a Novel Human Ubiquitin Specific Protease."

840 *Oncogene* 20 (29): 3869–79. https://doi.org/10.1038/sj.onc.1204553.

841 Sowa, Mathew E., Eric J. Bennett, Steven P. Gygi, and J. Wade Harper. 2009. "Defining the

842 Human Deubiquitinating Enzyme Interaction Landscape." *Cell* 138 (2): 389–403.

843 https://doi.org/10.1016/j.cell.2009.04.042.

844 Swatek, Kirby N, and David Komander. 2016. "Ubiquitin Modifications." *Cell Research* 26 (4):

845 399–422. https://doi.org/10.1038/cr.2016.39.

846 Vaz, Bruno, Marta Popovic, Joseph A. Newman, John Fielden, Hazel Aitkenhead, Swagata

847 Halder, Abhay Narayan Singh, et al. 2016. "Metalloprotease SPRTN/DVC1 Orchestrates

848 Replication-Coupled DNA-Protein Crosslink Repair." *Molecular Cell* 64 (4): 704–19.

849 https://doi.org/10.1016/j.molcel.2016.09.032.

850 Ventii, Karen H, and Keith D Wilkinson. 2008. "Protein Partners of Deubiquitinating Enzymes."

851 *The Biochemical Journal* 414 (2): 161–75. https://doi.org/10.1042/BJ20080798.

852 Wang, Tao, Luming Yin, Eric M. Cooper, Ming-Yih Lai, Seth Dickey, Cecile M. Pickart, David

Fushman, Keith D. Wilkinson, Robert E. Cohen, and Cynthia Wolberger. 2009. "Evidence

for Bidentate Substrate Binding as the Basis for the K48 Linkage Specificity of Otubain 1."

Journal of Molecular Biology 386 (4): 1011–23. https://doi.org/10.1016/j.jmb.2008.12.085.

Wühr, Martin, Robert M Freeman, Marc Presler, Marko E Horb, Leonid Peshkin, Steven Gygi,

Marc W Kirschner, and Marc W Kirschner. 2014. "Deep Proteomics of the Xenopus Laevis

Egg Using an MRNA-Derived Reference Database." *Current Biology : CB* 24 (13): 1467–

75. https://doi.org/10.1016/j.cub.2014.05.044.

Yardimci, Hasan, Anna B. Loveland, Antoine M. van Oijen, and Johannes C. Walter. 2012.

"Single-Molecule Analysis of DNA Replication in Xenopus Egg Extracts." *Methods* 57 (2):

179–86. https://doi.org/10.1016/j.ymeth.2012.03.033.

Ye, Yihong, and Michael Rape. 2009. "Building Ubiquitin Chains: E2 Enzymes at Work."

Nature Reviews. Molecular Cell Biology 10 (11): 755–64. https://doi.org/10.1038/nrm2780.

Ye, Yu, Masato Akutsu, Francisca Reyes-Turcu, Radoslav I Enchev, Keith D Wilkinson, and

David Komander. 2011. "Polyubiquitin Binding and Cross-reactivity in the USP Domain

Deubiquitinase USP21." *EMBO Reports* 12 (4): 350–57.

https://doi.org/10.1038/embor.2011.17.

Zeng, Xing, Frederic Sigoillot, Shantanu Gaur, Sungwoon Choi, Kathleen L. Pfaff, Dong-Chan

Oh, Nathaniel Hathaway, Nevena Dimova, Gregory D. Cuny, and Randall W. King. 2010.

"Pharmacologic Inhibition of the Anaphase-Promoting Complex Induces a Spindle

Checkpoint-Dependent Mitotic Arrest in the Absence of Spindle Damage." *Cancer Cell* 18

(4): 382–95. https://doi.org/10.1016/j.ccr.2010.08.010.

Acknowledgments

We gratefully acknowledge W. Harper for the gift of plasmids from the ORFeome collection, S.

Buhrlage for the gift of XL-188 and XL-203 and D. Finley for the gift of IU-C, IU-47 and USP21.

We thank Sandhya Manohar for the help in preparing the *Xenopus* egg extract. We thank W. Har-

per for helpful discussions. This work was supported by NIH grants R01 GM132129 (J.A.P.), R01

GM67945 (S.P.G) and R35 GM127032 (R.W.K.).

Author contributions

V.R. designed, performed, and analyzed all the experiments

V.R. prepared all samples for mass spectrometry with assistance from J.P.

J.P., J.C. and S.P.G. performed mass spectrometric analysis and provide reagents

B.B. provided advice on DUB selection and DUB assays

R.W.K. assisted with experimental design

V.R. and R.W.K. conceived of the project and wrote the manuscript with input from the other

authors.

Competing interests

The authors declare no competing interests

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the
PRIDE partner repository with the dataset identifier ….

Fig.1 Inhibition of cysteine-protease DUBs by UbVS causes rapid ubiquitin depletion and degradation of a set of proteins. (a) Extract was treated with UbVS or **(b)** HA-UbVS. After 10 minutes (time 0), 50 μM ubiquitin or buffer were added (**a**, bottom). Samples were subjected to immunoblotting (IB). **(c)** Extract was incubated with 10 μM HA-UbVS (10 minutes) and subjected to anti-HA pull-down followed by mass spectrometry analysis.The number of DUBs identified (gray) is compared to the number of DUBs detected previously in extract (Wuhr et al. 2014) (white). **(d)** Mass spectrometry analysis by SL-TMT method is summarized **(e)** Ubiquitin or buffer were added to the extract (time 0) pre-incubated with buffer or UbVS (10 minutes). Samples were collected as indicated **(f)** Volcano plots of TMT analysis comparing the proteins detected in UbVS/ubiquitin at 60 minutes with the proteins detected at time 0 (2 independent experiments). Statistical significance (-$\log_{10}$ p-value) is plotted against the log-transformed ratio of the samples. Proteins significantly decreasing in both the experiments are shown in black. **(g)** Heat map of the proteins downregulated in UbVS/ubiquitin in both the experiments in **(f)**.

a

Proteins	Function
MKRN1 UBOX5 (putative) **MARCH7 RNF138** BIRC3 NFX1 (putative) ZNF598	UPS (E3s)
RBM18 SYF2 EIF1 FAM32A **ZMAT2** RNASEH1 ZFP36L1	RNA metabolism
SOX3 SOX15 TGIF2 GTSF1	Transcription factors
CRIPT ARHGEF19 KIAA1430 WASL	Cytoskeleton
BORA WEE2 **PIM3** GIGYF1 HN1 PLK3	Proliferation/cell death
SPRTN ETAA1	DNA damage
STAM (trafficking) **CSNK1E** (circadian period) PLIN2 (lipid storage) C6ORF132 (?)	Others

b

c

d

Fig. 2 Independent validation of the proteins protected from degradation by UbVS-sensitive DUBs (a) Functions of the proteins down-regulated in UbVS/ubiquitin-treated extract (Fig.1g). Proteins independently validated are in bold. **(b)** Human orthologs (prefix h) or Xenopus protein (prefix Xe) were expressed in reticulocyte lysate, labeled with 35S-methionine and added to the extract, treated as in Fig.1e. Aliquots were collected and processed for SDS-PAGE and phosphorimaging. UbVS/ubiquitin-dependent degradation of these proteins was confirmed in 2 independent experiments. One of the expeiments is shown **(c)** Quantification of the experiment in **(b)**. Quantification of the second experiment is shown in Fig. S3d **(d)** Performed as in **(b)** *: ubiquitylated species.

Rossio et al.,Fig.2

Fig 3. Identification of DUBs sufficient to rescue degradation of the selected substrates in UbVS/ubiquitin-treated extract (a) Percentage of DUBs depleted by HA-UbVS. Extract was treated with 10μM HA-UbVS (10 minutes) and incubated overnight with anti-HA or empty beads. Supernatants were analyzed in triplicate by the SL-TMT method. Bold: DUBs tested in subsequent experiments.**(b)** Extract was incubated with 10 μM UbVS (30 minutes),after which recombinant DUBs (800 nM), labeled substrates and ubiquitin were added. After 60 minutes, aliquotes were processed for phosphorimaging. Quantitative profiles for 3 model substrates were created. The percentage of each substrate rescued by the addition of the DUBs is reported (two independent experiments). **(c)** Dose response analysis, performed as in **(b)**. The tested concentration is shown; the value in parentheses indicates the fold-increase of the tested concentration relative to the endogenous (two independent expeirments). One representative experiment is shown in Fig. S6**(c)** and **(d)** respectively. Performed as in **(b)** with the exception that ubiquitin was not added with the DUBs. Samples were processed for immunoblotting.

Rossio et al., Fig3

Fig.4 USP7 broadly rescues substrates from degradation in UbVS/ubiquitin-treated extract (a) Extract was incubated with UbVS followed by addition of DUBs and ubiquitin. Samples were analyzed by SL-TMT method. Bottom: R2 of the technical replicates **(b)** Volcano plot of quantitative proteomics analysis comparing proteins detected in the different conditions. Statistical significance is plotted against the log transformed ratio of the samples collected in duplicate. Proteins significantly decreasing in UbVS/ubiquitin are in blue **(c)** Heat map of the proteins downregulated in UbVS/ubiquitin after addition of DUBs. Bold: Proteins decreasing in both experiments in Fig. 1f. **(d)** Extract was treated such as in **(a)** 35S-methionine labeled substrates were added (time 0). USP7 was pre-incubated (30 min.) with buffer or UbVS (USP7-UbVS). Right: Quantification. Error bars: SEM of three independent experiments. One representative experiment is shown. (e) 35S-methionine labeled substrates were added to interphase and mitotic extract incubated with UbVS (30 min.). USP7 and ubiquitin were added (time 0). Aliquots were collected and processed for phosphorimaging. Right: Quantification of the substrate present at 20 minutes Error bars: SEM of three independent experiments. One representative experiment is shown.

Conditions	R² replicates exp.1	R² replicates exp.2
Time 0	0.9978	0.9942
Buffer	0.9968	0.9975
Ubiquitin	0.9917	0.9975
UbVS	0.9891	0.996
UbVS+ubiquitin	0.9831	0.999

Fig. S1 Analysis of the Xenopus proteome over time, after ubiquitin and UbVS addition.(a) Volcano plot of the 2 proteomic experiments comparing protein abundance between the indicated conditions (60 minutes) and time 0 as in Fig.1f. Statistical significance (-$\log_{10}$ *p-value*) is plotted against fold change (average log2 ratio). Proteins significantly decreasing in the indicated conditions are in black. (**b**) R2 of the technical replicates are reported.

Fig. S2 Increasing UbVS concentration does not enhance protein degradation in extract. (a) Extract was incubated with buffer, 10 or 30 μM UbVS (10 minutes) and 50 μM of ubiquitin was added (time 0). Samples were collected in triplicate and processed for SL-TMT method analysis **(b)** Volcano plot of quantitative proteomics analysis. Statistical significance (-log$_{10}$ *p-value*) is plotted against ratio (average log$_2$ ratio). Blue dots: proteins significantly decreasing in 10 μM UbVS/ubiquitin (Fig.1g) **(c)** Log2 ratio heat map compares the proteins decreasing in **(b)**. Proteins decreasing in the experiments in Fig.1f are in bold. **(d)** Extract was treated with UbVS (10 minutes), after which 50 μM of ubiquitin was added (time 0). Aliquots were subjected to immunoblotting.

Fig.S3 Identification of non-degradative substrate-ubiquitin conjugates (a) Extract was treated with 50 μM HA-ubiquitin (HA-UB) for 30 minutes and subjected to immunoblotting (IB) **(b)** After incubation with HA-ubiquitin, extract was subjected to anti-HA pull-down and analyzed by TMT-mass spectrometry. Volcano plot comparing abundance of protein bound to anti-HA beads relative to empty beads. Statistical significance (-log10 *p-value*) is plotted against ratio (average log2). Samples were collected in technical triplicates. Proteins significantly increasing are in black. Abundant UPS components are labeled **(c)** Overlap between the HA-ubiquitin pull down and the diGly enrichment **(d)**Volcano plot of the TMT diGly remnant analysis comparing changes in ubiquitination sites detected after addition UbVS/ubiquitin to the changes after addition of only ubiquitin (30 min.).Samples were collected in duplicate.Statistical significance is plotted against ratio (average log2). Proteins significantly increasing are in black. **(e)** Ubiquitination sites of the proteins destabilized in UbVS/ubiquitin (Fig.1f) are shown. The graph shows the relative amount (R.A.) of ubiquitination in the conditions tested (average between duplicates). A.U. Arbitrary unit. Error bars: standard deviation (N=2)

a

Proteins	Function
RNF10 TRIM37 TRIP12 PPIL2 NOSIP	UPS (E3s)
SNRPA CPEB1 DND1 ZC3HAV1 UTP14A	RNA metabolism
TCEANC2 ACRC (putative) POUF51	Transcription factors
CCDC120 CTTN PKP3 UPK1A SVIL SSFA2	Cytoskeleton
NEK2 BTG4	Proliferation/Cell death
SFXN1 **SSBP1** MTG1 HTRA2	Mitochondrial proteins
OVCH1 SERPINA1	Proteases
CSNK1D EEA1 (trafficking) DNAJB7 (folding) CRIP3 (?) KIAA1551 (?) WDFY1(immunity)	Others

Fig. S4 Validation of the substrates protected from degradation by the UbVS-sensitive DUBs (a) Functions of the 5% of proteins whose abundance decreased most in UbVS/ubiquitin (FDR 1%) in both the experiments (Fig.1f). The 34 proteins included in the more selective threshold are not shown. In bold, proteins validated with independent experiments. **(b)** The indicated proteins were expressed and labeled as described previously and added to extract pre-treated as indicated. Levels of the proteins were assessed by SDS-PAGE and phosphorimaging **(c)** Quantification of the substrate levels (60 minutes) of the experiment in **(b)**. Substrates have been validated with 2 independent experiments. One representative experiment is shown. **(d)** Quantification of the second set of independent experiments in **(c)** and in Fig.2.

d

Conditions	R^2 between replicates
Time 0	0.916
UbVS+Ub+ DMSO	0.99
UbVS+Ub+MG262	0.9953

Fig. S5 Degradation of proteins in UbVS/ubiquitin-treated extract is proteasome-dependent (a) Extract was incubated with 200 µM MG262 or DMSO (20 minutes), after which 10 µM UbVS was added (10 minutes), followed by addition of 50 µM of ubiquitin (time 0). Samples were collected in duplicate (time 0 and 60 minutes) and processed for mass spectrometry analysis by the SL-TMT method **(b)** Volcano plot of quantitative proteomics analysis comparing the proteins detected in UbVS/ubiquitin in presence of DMSO or MG262 at 60 minutes with the proteins detected at time 0. Statistical significance (-log10 *p-value*) is plotted against ratio (average log2 ratio). Blue dots represent the proteins significantly decreasing in UbVS/ubiquitin and ubiquitin are labeled **(c)** Log2 ratio heat map of the proteins significantly downregulated in UbVS/ubiquitin. Proteins decreasing in the experiments in Fig.1G are in bold. **(d)** R2 of the technical replicates of each condition.

Fig S6. Identification of the fraction of each DUB depleted by HA-UbVS and validation of the catalytic activity of recombinant DUBs (a) Volcano plot of quantitative proteomics analysis comparing the proteins detected in the supernatant of the empty beads with the proteins detected in the supernatant of the anti-HA beads. Statistical significance (-$\log_{10}$ *p-value*) is plotted against ratio (average $\log_2$ ratio). DUBs are labeled **(b)** Indicated DUBs were incubated with saturating amount of UbVS or buffer (30 minutes). Reactions were stopped with SDS sample buffer, run on SDS-PAGE and the gel was stained with Comassie Blue. **(c)** USP9X was incubated with HA-UbVS and its activity was examined by Immunoblotting.

Fig.S7. Experiments used to create the quantitative DUB activity profile (Fig.3).(a) Estimated endogenous concentration of the DUBs tested (Fig.3a). One of the representative experiments reported in the quantitative profile in Fig.3b is shown in **(b)**, in Fig.3c is shown in **(c)** and in Fig.3d is shown in **(d)**.The blue rectangles in **(a)** indicate the four DUBs rescuing all the three substrates tested.

Fig.S8 Usp7 or Usp14 inhibition does not promote broad protein instability (a) Extract was incubated with the indicated compounds (20 minutes), followed by HA-UbVS addition (20 minutes). Aliquots were analyzed by immunoblotting. IU-47 and IU-C: Usp14 inhibitor (Usp14i) and control compound. XL-188 and XL-203: Usp7 inhibitor (Usp7i) and control compound. **(b)** Extract was incubated with DMSO or compounds. UbVS was added for 10 minutes(time 0) to the samples with DMSO. Ubiquitin was added to all samples, samples were collected in duplicate and analyzed by the SL-TMT method. **(c)** Top: heat map compares the effect of the specific DUB inhibitors on the proteins significantly decreasing in UbVS/ubiquitin in this experiment and in the experiments in Fig. 1F. Bottom: heat map of the proteins affected by the specific DUB inhibitors. **(d)** Volcano plot of quantitative proteomicsanalysis. Statistical significance (-$\log_{10}$ *p-value*) is plotted against ratio (average log2 ratio).Blue dots: proteins decreasing in UbVS/ubiquitin; black dots:proteins decreasing in UbVS/ubiquitin and in presence of Usp7i; orange dots: proteins decreasing in presence of both the inhibitors. **(e)** R2 of the technical replicates

Table S1: List of the Xenopus DUBs

DUBs	Estimated concentration (nM)	Class	
CYLD	16	USP	**n=35: UbVS-reactive**
USP4	5.65	USP	
USP5	546.4	USP	
USP7	146.9	USP	
USP8	46.2	USP	
USP9X	73.99	USP	
USP10	209	USP	
USP11	78	USP	
USP14	425	USP	
USP15	21.48	USP	
USP19	15.7	USP	
USP24	11.1	USP	
USP28	3.44	USP	
USP37	5.84	USP	
USP38	13.08	USP	
USP40	48.32	USP	
USP47	63.6	USP	
USP48	23	USP	
USP25		USP	
OTUB1	102	OTU	
OTUB2	42	OTU	
YOD1	66	OTU	
OTUD4	31.4	OTU	
OTUD3	8.11	OTU	
OTUD6B	53.5	OTU	
VCPIP1	53.6	OTU	
OTUD5		OTU	
OTUD7B	3.57	OTU	
UCHL1	2523	UCH	
UCHL3	1465	UCH	
UCHL5	600	UCH	
FAM188A	45	MINDY	
FAM188B	2.8	MINDY	
ATXN3	91.6	MJD	
ZUFSP	10.35	ZUFSP	
USP1	5.96	USP	**n=19 Expressed but not UbVS-reactive**
USP16	38.27	USP	
USP39 *	45.07	USP	
USP3	3.04	USP	
USP6	1.15	USP	
USP12	1.82	USP	
USP13	7.96	USP	
USP20	5.71	USP	
USP30	7.21	USP	
USP32	1.81	USP	
USP33	1.29	USP	
USP34	14.22	USP	
USP36	0.91	USP	
USP13	7.96	USP	
ALG13*	29.8	OTU	
FAM63A	28.6	MINDY	
BRCC3	*23.62*	*JAMM*	
CSN5=cops	*459.82*	*JAMM*	
RPN11=PS	*749.98*	*JAMM*	
USP2		USP	**n=43 Not detected or UbVS-reactive**
USP9Y		USP	
USP17L2		USP	
USP18		USP	
USP21		USP	
USP22		USP	
USP26		USP	
USP27X		USP	
USP29		USP	
USP31		USP	
USP35		USP	
USP41		USP	
USP42		USP	
USP43		USP	
USP44		USP	
USP45		USP	
USP46		USP	
USP49		USP	
USP50 *		USP	
USP51		USP	
PAN2/USP52 *		USP	
USP53 *		USP	
USP54 *		USP	
BAP1		UCH	
ATXN3L		MJD	
JOSD1		MJD	
JOSD2		MJD	
H1N1L		OTU	
A20		OTU	
Cezanne 2		OTU	
FAM105A		OTU	
OTUD1		OTU	
OTUD5		OTU	
OTUD6A		OTU	
OTUD7A		OTU	
TNFAIP3		OTU	
OTU1		OTU	
ZRANB1=TRABID		OTU	
AMSH		*JAMM*	
AMSH-LP		*JAMM*	
MPND		*JAMM*	
MYSM1		*JAMM*	
*PRPF8**		*JAMM*	

*: probably inactive
JAMM=Metallo-protease DUBs
Concentration is based on Wuhr et al., 2014

TableS2: HA-UbVS interactors

Protein	Gene symbol	Gene Description	Empty beads rep1	Empty beads rep2	Anti-HA beads rep1	Anti-HA beads rep2	NOTES
CL492Conti	PSMC3	26S protease regulatory subunit 6A	8	12	176	190	
zeinaSSns	PSMC5	26S protease regulatory subunit 8	6	8	163	178	DUBs
CL3721Cor	PSMD2	26S proteasome non-ATPase regulatory subunit 2	3	4	148	150	Proteasome, proteasome-associated proteins, ubiquitin
CL1908Cor	PSMD11	26S proteasome non-ATPase regulatory subunit 11	5	7	139	151	
CL2523Cor	PSMD1	26S proteasome non-ATPase regulatory subunit 1	5	3	138	144	
CL2720Cor	USP5	Ubiquitin carboxyl-terminal hydrolase 5	5	2	130	146	
CL4411Cor	UCHL5	Ubiquitin carboxyl-terminal hydrolase isozyme L5	2	2	112	128	
TC449583	UBC	Polyubiquitin-C	2	3	109	84	
CL6Contig5	USP7	Ubiquitin carboxyl-terminal hydrolase 7	2	1	101	113	
CL5541Cor	PSMC2	26S protease regulatory subunit 7	6	5	97	90	
CL2791Cor	PSMC6	26S protease regulatory subunit 10B	4	5	90	105	
zeinaSSns	PSMC1	26S protease regulatory subunit 4	7	6	82	86	
CL4759Cor	PSMD13	26S proteasome non-ATPase regulatory subunit 13	0	2	80	80	
dsrrswapns	USP14	Ubiquitin carboxyl-terminal hydrolase 14	1	3	79	79	
CL18578Cc	UCHL1	Ubiquitin carboxyl-terminal hydrolase isozyme L1	5	4	78	87	
CL5907Cor	PSMD12	26S proteasome non-ATPase regulatory subunit 12	2	3	76	84	
CL2698Cor	PSMD6	26S proteasome non-ATPase regulatory subunit 6	3	4	73	71	
CL60Contig	PSMD3	26S proteasome non-ATPase regulatory subunit 3	6	5	73	70	
CL2846Cor	USP9X	Probable ubiquitin carboxyl-terminal hydrolase FAF-X	0	0	63	67	
CL1008Cor	PSMC4	26S protease regulatory subunit 6B	0	1	60	66	
zeinaSSns	USP47	Ubiquitin carboxyl-terminal hydrolase	0	0	60	55	
TC426391	UCHL3	Ubiquitin carboxyl-terminal hydrolase isozyme L3	1	1	58	75	
CL10383Cc	USP28	Ubiquitin carboxyl-terminal hydrolase 28	0	0	54	48	
CL5540Cor	PSMD14	26S proteasome non-ATPase regulatory subunit 14	3	5	53	56	
CL2142Cor	USP11	Ubiquitin carboxyl-terminal hydrolase 11	0	0	49	43	
CL2142Cor	USP4	Ubiquitin carboxyl-terminal hydrolase 4	0	0	47	38	
CL1100Cor	PSMD4	26S proteasome non-ATPase regulatory subunit 4	2	3	42	44	
CL5178Cor	VCPIP1	Deubiquitinating protein VCIP135	0	0	40	40	
dsrrswapns	USP40	Ubiquitin carboxyl-terminal hydrolase 40	0	0	32	42	
CL306Conti	PSMD8	26S proteasome non-ATPase regulatory subunit 8	0	1	31	44	
dsrrswapns	OTUD4	OTU domain-containing protein 4	0	0	30	34	
CL4416Cor	PSMB3	Proteasome subunit beta type-3	4	3	24	22	
TC424927	PSMA5	Proteasome subunit alpha type-5	0	1	23	23	
TC423038	OTUB1	Ubiquitin thioesterase OTUB1	0	0	22	22	
CL7391Cor	CYLD	Ubiquitin carboxyl-terminal hydrolase CYLD	0	0	21	18	
CL3942Cor	USP15	Ubiquitin carboxyl-terminal hydrolase	0	0	20	22	
CL150Conti	USP10	Ubiquitin carboxyl-terminal hydrolase 10	0	0	20	18	
CL3001Cor	USP38	Ubiquitin carboxyl-terminal hydrolase 38	0	0	20	18	
CL4026Cor	PSMA2	Proteasome subunit alpha type-2	2	2	19	20	
CL3430Cor	PSMA1	Proteasome subunit alpha type-1	0	0	19	15	
dsrrswapns	G3BP1	Ras GTPase-activating protein-binding protein 1	0	0	18	16	Usp10 regulator (Soncini et al., 2001)
dsrrswapns	YOD1	Ubiquitin thioesterase OTU1	0	0	17	22	
CL6350Cor	PSMD5	26S proteasome non-ATPase regulatory subunit 5	0	0	17	20	
dsrrswapns	ATXN3	Ataxin 3 variant ref	0	0	17	16	
CL6078Cor	PSMA6	Proteasome subunit alpha type-6	0	4	16	13	
CL3715Cor	USP8	Ubiquitin carboxyl-terminal hydrolase 8	0	0	15	20	
CL5195Cor	UBE3A	Ubiquitin-protein ligase E3A	0	0	15	19	
CL8846Cor	PSMB4	Proteasome subunit beta type-4	1	2	14	16	
CL6048Cor	PSMA4	Proteasome subunit alpha type-4	0	0	14	13	
CL362Conti	USP24	Ubiquitin carboxyl-terminal hydrolase 24	0	0	14	13	
CL5308Cor	PSMB7	Proteasome subunit beta type	0	0	14	8	
CL6056Cor	RNF216	E3 ubiquitin-protein ligase RNF216	0	0	12	11	
CL4431Cor	ADRM1	Proteasomal ubiquitin receptor ADRM1	0	0	12	8	
CL4421Cor	OTUD6B	OTU domain-containing protein 6B	0	0	11	8	
zeinaSSns	G3BP2	Ras GTPase-activating protein-binding protein 2	1	0	10	12	Usp10 regulator (Soncini et al., 2001)
CL6932Cor	TXNL1	Thioredoxin-like protein 1	0	0	9	6	Proteasome associated protein
TC414297	USP25	Ubiquitin carboxyl-terminal hydrolase 25	0	0	8	9	
CL7369Cor	USP48	Ubiquitin carboxyl-terminal hydrolase 48	0	0	7	10	
CL1858Cor	USP19	Ubiquitin carboxyl-terminal hydrolase (Fragment)	0	0	7	7	
dsrrswapns	OTUB2	Ubiquitin thioesterase OTUB2	0	0	7	5	
CL5807Cor	SENP8	Sentrin-specific protease 8	0	0	6	9	UbVS target
CL947Conti	FAM188A	Protein FAM188A	0	0	6	6	
CL3708Cor	RNF213	E3 ubiquitin-protein ligase RNF213	0	0	6	4	
CL2263Cor	UBE3C	Ubiquitin-protein ligase E3C	0	0	6	4	UbVS target
CL11640Cc	ULK3	Serine/threonine-protein kinase ULK3	0	0	6	4	
CL4995Cor	OTUD3	OTU domain-containing protein 3	0	0	6	3	
zeinaSSns	ACTN1	Alpha-actinin-1	0	0	5	7	
CL8177Cor	LMNA	Prelamin-A/C	0	0	5	5	
CL10519Cc	METTL18	Histidine protein methyltransferase 1 homolog	0	0	5	5	
zeinaSSns	DYNC1LI1	Cytoplasmic dynein 1 light intermediate chain 1	0	0	5	2	
TC432167	PSMB7	Proteasome subunit beta type-7	0	0	4	5	
CL12340Cc	PSMD10	26S proteasome non-ATPase regulatory subunit 10	0	0	4	5	
CL2481Cor	USP37	Ubiquitin carboxyl-terminal hydrolase 37	0	0	4	3	
zeinaSSns	ADPRHL2	Poly(ADP-ribose) glycohydrolase ARH3	0	0	4	2	
CL8288Cor	FAM188B	Protein FAM188B	0	0	3	5	
CL7134Cor	DRG2	Developmentally-regulated GTP-binding protein 2	0	0	3	4	
CL11278Cc	ZUFSP	Zinc finger with UFM1-specific peptidase domain protein	0	0	3	3	
CL391Conti	RPL18A	60S ribosomal protein L18a	0	0	3	3	
TC464126	HNRNPC	Heterogeneous nuclear ribonucleoproteins C1/C2	0	0	3	2	
dsrrswapns	ZFYVE16	Zinc finger FYVE domain-containing protein 16	0	0	3	2	
zeinaSSns	OTUD5	OTU domain containing 5, isoform CRA_c	0	0	2	2	
CL20Contig	ACTR2	Actin-related protein 2	0	0	2	2	
CL99Contig	PLEC	Plectin	0	0	2	2	
dsrrswapns	RNF14	E3 ubiquitin-protein ligase RNF14 (Fragment)	0	0	2	2	
CL752Conti	SPATA2L	Spermatogenesis-associated protein 2-like protein	0	0	2	2	CYLD interactor (Sowa et al., 2009)
CL871Conti	VIM	Vimentin	0	0	2	2	
dsrrswapns	SPATA2	Spermatogenesis-associated protein 2	0	0	1	2	CYLD interactor (Sowa et al., 2009)
CL12666Cc	OTUD7B	OTU domain-containing protein 7B	0	0	1	1	

rep:replicate
Selection of the interactors
Proteins are present in both replicates of the anti-HA beads and if present on empty beads they should be enriched at least 6 fold on the anti-HA beads
If multiple isoforms were present, only the isoform with more peptides is reported (for semplicity)
UbVS target are based on Hewings et al., 2018

TableS3: UbVS-sensitive proteolitic DUB substrates

Substrates	Function	Unstructured regions	PEST sequence (number)	UPS reported substrate	DUBs reported substrate
NFX1	E3	N-term and C-term			USP9
MARCH 7	E3	highly disordered		YES	USP7/USP9
MKRN1	E3	N-term and multiple regions			
FAM32A	Putative mRNA binding protein	highly disordered			
RNASEH1	Endonuclease	C-term			
PIM3	Kinase	C-term	YES	YES	USP7 is reported to act on the homolog PIM2
CRIPT	Cytoskeleton binding protein	C-term			
WASL	Actin binding protein	highly disordered	YES	YES	USP7 is reported to act on a member of the same family WASH
KIAA1430	Cilia and flagella associated protein	highly disordered	YES (2)		
RBM18	Probable RNA binding protein	C-term			
HN1	Microtuble associated protein	highly disordered	YES		
RNF138	E3	N-term and C-term		YES	
GTSF1	Transcriptional repressor	N-term and C-term	YES (2)		
SYF2	Pre-mrna splicing factor	highly disordered			
ZNF598	E3	big disordered region in the center	YES		
ARHGEF19	Rho guanine nucleotide exchanging	N-term and C-term			
ZMAT2	Spliceosome component	N-term and C-term			
BORA	Kinase activator	N-term and C-term		YES	
SOX3	Transcription factor	N-term and C-term	YES		
CSNK1E	Kinase	N-term and C-term			
SOX15	Transcription factor	N-term and C-term			
ZFP36L1	mRNA binding protein	highly disordered	YES		
BIRC3	E3	mostly ordered		YES	USP19
EIF1	Translation initiation factor	N-term and C-term			
UBOX5	Hypotetical E3 (for similarity)	multiple disordered regions			
ETAA1	DNA damage associated protein	c-term			
SPRTN	DNA damage associated protein	multiple disordered regions		YES	
WEE2	Kinase	N-term	YES		
PLIN2	Lipid storage	N-term and C-term		YES	
STAM	Signal transducing adapter	multiple disordered regions and C-term		YES	USP8
PLK3	Kinase	N-terminal and small C-term	YES	YES	
TGIF2	Transcription factor	multiple disordered regions	YES		
C6ORF132	Uncharachterized protein	highly disordered	YES (5)		
GIGYF1	GRB10 interactor protein	multiple disordered regions	YES (2)		