1 Title:
2 Evaluation of Dose Fractionated Polymyxin B on Acute Kidney Injury: A Translational *In Vivo* Model
3

4 Authors:
5 Jiajun Liu[1,2,3]; Gwendolyn M. Pais[1,2,3]; Sean N. Avedissian[4,5]; Annette Gilchrist[1,2,6]; Andrew Lee[6]; Nathaniel J.
6 Rhodes[1,2,3]; Alan R. Hauser[6]; Marc H. Scheetz[1,2,3]
7
8 [1]Midwestern University, Downers Grove, IL [2]Midwestern University Chicago College of Pharmacy
9 Pharmacometrics Center of Excellence, Downers Grove, IL [3]Northwestern Memorial Hospital, Chicago, IL
10 [4]Antiviral Pharmacology Laboratory, University of Nebraska Medical Center (UNMC) Center for Drug Discovery,
11 UNMC, Omaha, NE [5]University of Nebraska Medical Center, College of Pharmacy, Omaha, NE [6]Northwestern
12 University, Chicago, IL
13
Corresponding Author:
Marc H. Scheetz, PharmD, MSc; Professor of Pharmacy Practice and Director of Pharmacometrics Center of
Excellence; Midwestern University Chicago College of Pharmacy; 555 31[st] Street, Downers Grove, IL 60515, Phone:
630-515-6116; Fax: 630-515-6958; Email: mschee@midwestern.edu

Financial disclosure: None

Keywords: polymyxins, polymyxin B, colistin, CMS, pharmacokinetic, toxicodynamic, animal model, acute kidney
injury, biomarker, KIM1

Acknowledgements: None

Abstract

The polymyxins are last-line defense for highly resistant infections. Nephrotoxicity, however, is a dose-limiting

factor. Yet, approaches to mitigate nephrotoxicity are poorly defined. This study aimed to investigate the impact of

dose fractionated (once, twice and thrice daily) polymyxin B (PB) on acute kidney injury (AKI) in a pre-clinical

model. Secondarily, we aimed to describe the pharmacokinetic (PK) profile of PB. Sprague-Dawley rats were

assigned to experimental groups with different dosing intervals but constant total daily exposure (12 mg/kg/day into

single, twice daily, and thrice daily doses) and controls received normal saline subcutaneously over 3 days. Blood

and urine samples were collected, and kidneys were harvested at necropsy. A three-compartment model best

described the data and Bayesian observed vs. predicted concentration demonstrated bias, imprecision, and R^2 of

0.129 mg/L, 0.729 mg^2/L^2 and 0.652, respectively. PB exposure (i.e. AUC_{24h}) were similar across treatment groups

over time (p=0.87). As a representative, urinary KIM-1 were elevated on days 1 and 2 for experimental groups

compared to controls, and thrice daily group experienced the most KIM-1 increase [mean increase (95% CI) day 1

from day -1, 4.44 (0.89, 8.00) ng/mL; p=0.018] as compared to control [mean increase (95% CI) day 1 from day -1,

0.03 (-0.42, 0.49) ng/mL; p=0.99]. Correspondingly, significant histopathological damage was observed with the

same group (p=0.013) (controls as a referent). Our findings suggested that fractionating the PB dose thrice daily

resulted in the most injury in a rat model.

Background

The widespread use of broad-spectrum antimicrobial agents has led to an increasing rate of resistant infections, and

Gram-negative pathogens including *Pseudomonas aeruginosa*, *Acinetobacter* spp., and *Enterobacteriaceae* spp. are

particularly problematic (1, 2). The Center for Disease Control and Prevention (CDC) has published a

comprehensive report detailing the top antibiotic resistant threats in the U.S., stating that at least 2.86 million

Americans contract resistant bacterial and fungal infections and at least 35,900 die annually as a result (3). It is

estimated that by 2050, global deaths due to antimicrobial resistance will balloon to 10 million people per year and

become the leading cause of mortality (4). Multiple Gram-negative species are resistant to nearly all available

antibiotics (3), including newer combination agents (5, 6). Although multiple drugs in the antibiotic pipeline are

promising, there is a prudent need to maximize clinical efficacy and safety of currently available agents (7). As a

result of the paucity of active antibiotics for these difficult-to-treat infections, the polymyxins remain last resort

options (8, 9).

The polymyxins are a group of polypeptide antibiotics discovered more than 70 years ago, with activity and efficacy

against Gram-negative pathogens (10, 11). Polymyxin use has declined as a result of associated renal and

neurological adverse effects and newer agents with more favorable safety profiles have emerged (12, 13). Thus, the

treatment-limiting adverse effects of the polymyxins such as kidney injury have greatly limited their utility for

patient care (14-16). Contemporary studies utilizing widely accepted dosing regimens have demonstrated that

nephrotoxicity rates range from 21-48% (17-20). The mechanism for which nephrotoxicity develops appears to

involve several processes. First, the drug is selectively reabsorbed by the renal brush border membrane and

accumulates in renal cells, directly exerting cytotoxic effects to the proximal tubule cells (21). Secondly,

accumulation of drug in the kidneys leads to increased membrane permeability and cell lysis, causing acute tubular

necrosis (15, 22). Lastly, oxidative stress may also play a role in the development of nephrotoxicity associated with

polymyxin therapy (16, 23, 24).

While the various mechanisms of polymyxin toxicity are being elucidated, dosing strategies that accelerate/diminish

toxicity remain poorly defined. Toxicity thresholds for plasma 24-hour area under the curve (AUC_{24h}) of polymyxin

B and colistin have recently been highlighted, yet approaches to minimize nephrotoxicity risk resulted in mixed

outcomes (20, 25-29). More specifically, it remains unclear whether dividing the total daily dose of polymyxins into

fractions (e.g. giving twice or thrice daily) can circumvent kidney injury during treatment. In this study, we

examined the impact of dose fractionated systemic polymyxin B on acute kidney injury (AKI) in a pre-clinical,

humanized model with novel urinary biomarkers (30, 31). In addition, we aimed to describe the polymyxin B

pharmacokinetic (PK) profile.

Material and Methods

Chemical and reagents

Clinical grade Polymyxin B sulfate (PB) (Lot# CD807) for injection (USP) was purchased from X-GEN

Pharmaceuticals (Horseheads, NY, USA). Study drug was reconstituted and diluted with normal saline (NS) for

injection. Unused portions were properly discarded to minimize loss of potency in subsequent experiments (32).

Colistin sulfate (Sigma-Aldrich Chemical Company, Milwaukee, WI, USA) and creatinine-d3 (Cayman Chemicals,

Ann Arbor, MI, USA) were used as internal standards for the LC-MS/MS assay (Agilent Technologies). All

solvents for the LC-MS/MS analysis were of LC-MS/MS grade. Acetonitrile and methanol were purchased from

VWR International (Radnor, PA, USA). Formic acid was obtained from Fisher Scientific (Waltham, MA, USA).

Pooled male Sprague-Dawley rat plasma was used for sample preparation and calibration of standard curves

(BioreclamationIVT, Westbury, NY, USA).

Experimental design and animals

Allocation and number of animals for experimental (i.e. PB-treated) and control protocol groups are described in Fig.

1. The experimental arm was further divided into three groups based on dose fractionation design: once daily (QD),

twice daily (BID), and thrice daily (TID). Each experimental group received subcutaneous injections of PB, and

control groups received equal volumes of NS based on the QD protocol. In all, there were four study groups. Total

daily dose of PB was fixed at 12 mg/kg/day (allometrically scaled) for all experimental groups administered

subcutaneously for 72 hours (i.e. 3 PB doses for QD group, 6 for BID group, and 9 for TID group during the 3-day

study period) (33).

Male Sprague-Dawley rats (n=32, approximately 8 to 10 weeks old; Harlan, Indianapolis, IN, USA) were used.

Animals were housed in a light- and temperature-controlled rooms during acclimation and study periods. Animals

were maintained in plastic cages on a 12-hour light and 12-hour dark cycle. Food and water were freely accessible

at all times except during periods in which sampling catheters (one per animal) were surgically placed prior to

initiation of study protocol. Post-operative pain was monitored according to protocol. Data were analyzed for all

protocol-initiated animals unless terminated early, in which case data were treated as missing.

This study was conducted at Midwestern University, Downers Grove, IL. The study methods were reviewed and

approved by the Midwestern University Institutional Animal Care and Use Committee (IACUC; protocol 2677). All

animals were cared for and handled in concordance with animal care and use standards and ethical principles.

Blood and urine sampling

Fig. 2 provides a schematic flow of study design for each study group. Blood samples were obtained via an internal

jugular vein catheter that was surgically cannulated on day 0 after the acclimation period. Animals were under

ketamine (100 mg/kg) and xylazine (10 mg/kg) anesthesia for surgical procedure and were allowed 24-hour

recovery periods prior to initiation of the study protocol (i.e. drug dosing or blood sampling). When not in use,

catheters were locked with heparin solution (100 IU/mL). Blood samples (0.125 mL aliquots) were obtained after

the first dose (day 1) with a staggered sampling design. A maximum of 16 samples per animal were obtained during

a 4-day period (pre-euthanasia) and no more than 8 samples were drawn in a single day. As an example, for QD and

control groups blood samples were drawn at 5, 20, 60, 120, 180, 240, 360, and 480 minutes after the first dose on

day 1. Eight blood samples total were then obtained on days 2 and 3. A terminal sample was also drawn under

terminal anesthesia. Blood sampling schemes for BID and TID groups follow the same protocol (i.e. maximum

number of samples and volume per animal) while the sampling times were adjusted accordingly based on the dosing

intervals. Each sample was replaced with equal volume of NS to maintain euvolemia. Blood samples were

immediately transferred to a disodium EDTA (Sigma-Aldrich Chemical Company, Milwaukee, WI, USA) treated

microcentrifuge tube and centrifuged at 600 *xg* for 10 minutes. Plasma supernatant was collected and stored at -

80°C for batch sample analysis. Animals were placed in metabolic cages for urine collection as previously

described (34, 35). In brief, discrete entry times were recorded for initial transfer of animals to the metabolic cages

(catalogue number 650-0350; Nalgene, Rochester, NY) on day -1 (baseline), and urine collections and volume

measurements followed the 24-hour period on days 2, 3 and 4. All urine samples were collected in laboratory-

controlled ambient conditions, and urinary biomarkers were stable throughout as previously described (35). Urine

samples were centrifuged at 500 *xg* at 4°C for 10 minutes, and supernatant was stored at -80°C for batch analysis.

Determination of PB and creatinine concentrations in plasma

For quantification of plasma PB concentration, 40 µL plasma sample was combined with 4 µL of internal standard

of colistin sulfate at a concentration of 0.1 mg/mL. Protein precipitation was then performed with 456 µL of

methanol containing 0.1% formic acid. Following centrifugation for 10 minutes at 16,000 xg (Eppendorf model:

5424), 100 µL supernatant was collected for analysis. Processed samples were injected (injection volume at 2 µL)

into an Agilent 1260 infinity binary liquid chromatograph paired with Agilent 6420 triple quadrupole mass

spectrometer (MS). A Poroshell 120 EC-C18 column (100 mm x 3 mm, 2.7 µm) was used. The following

quantifier transitions (m/z) for polymyxin B1 (PB1) and colistin A were identified and utilized: 402.2 → 101.1,

390.6 → 101.3, respectively. The assay was linear between 0.5 to 40 mg/L (R^2=0.997) for PB1 after an applied

weight of 1/x. To quantify plasma creatinine, m/z transitions of 117.09 → 89.2, 114.1 → 44.3 were utilized for

creatinine-d3 and creatinine, respectively. After adjusting for endogenous creatinine in pooled blank plasma and 1/x

weighting, the linear range for creatinine assay was between 0.3 and 40 mg/dL (R^2=0.999). The coefficient of

variation (CV%) values for PB1 and creatinine assays were below 10% for intra- and inter-day measures. All

samples measuring above the upper limit of quantification underwent serial dilution for analysis.

Determination of urinary biomarkers of kidney injury

Urine samples were analyzed for creatinine content and urinary biomarkers. Urine aliquots were analyzed in

batches to determine the concentration of KIM-1, IP-10, TIMP-1, CLN and OPN. Urinary biomarkers were assayed

using a microsphere based MAGPIX kit as previously described (35). In brief, urine samples were aliquoted into

96-well black plates supplied with MILLIPLEX® MAP Rat Kidney Toxicity Magnetic Beed Panel 1 (EMD

Millipore Corporation, Charles, MO, USA), prepared and analyzed per manufacturer's recommendations.

Histopathological examination of renal cell damage

Kidney tissues were harvested following euthanasia. Each animal's kidneys were removed and briefly washed in

cold NS. The left kidney was fixed in 10% formalin for histopathology examination. Histopathological analysis of

kidneys (n=32) was conducted by IDEXX BioAnalytics (Columbia, MO, USA). A validated, ordinal scoring

system was employed to grade pathological lesions as previously described (31, 36, 37). Briefly, a scale of 0

indicates no abnormality while a scale of 5 indicates massive and extensive renal damage. The final histopathology

score for an individual animal was calculated based on the highest score from the anatomical structural segment.

PB pharmacokinetic model and exposure determination

To construct the base PK models and generate exposure estimates for each individual animal, the Nonparametric

Adaptive Grid (NPAG) algorithm (38, 39) within the Pmetrics (Version 1.5.2) package (39) for R (6) was utilized.

Multiple models were built and assessed. A three-compartmental structural model of PB disposition accounting for

absorption constant (K_a) from injection site to the central compartment (V_C) was fitted to all PK data. The PK

model was parameterized with K_a, V_C, intercompartmental transfer rates (K_{23}, K_{32}) between central and peripheral

(V_P) compartments, and total elimination rate constant (K_e). Assay error was included in the model using a

polynomial equation in the form of standard deviation (SD) as a function of each observed concentration, Y (i.e. SD

$= C0 + C1 * Y$). Observation weighting was performed using lambda (i.e. error $= SD^2 * lambda^2)^{0.5}$, an additive

variance model to account for extra process noise. Lambda was initially set at 1 with C0 and C1 equal to 0.1 and 0.1,

respectively. Comparative model performance was examined by the change in objective function value (OFV)

calculated as differences in -2 log-likelihood (-2LL), with a reduction of 3.84 in OFV corresponding to p <0.05

based on chi-square distribution with one degree of freedom. Further, the best-fit model was selected based on the

rule of parsimony and the lowest Akaike's information criterion (AIC) scores. Goodness-of-fit of the competing

models were evaluated by regression on observed vs. predicted plots, coefficients of determination, and visual plots

of individual Bayesian predicted concentration-time profiles. Bias was defined as mean weighted prediction error;

imprecision was defined as bias-adjusted mean weighted squared prediction error. Using the final model, PB

exposure indices (AUC_{24h}, C_{MAX}, and C_{MIN}) were calculated from individual Bayesian posterior-predicted

concentrations using 'makeNCA' within Pmetrics package across 24-hour intervals (39).

Statistical analysis for biomarkers, PK indices and histopathological scoring

Analysis of variance (ANOVA) with Geisser and Greenhouse epsilon hat correction method accounting for subjects,

treatment groups, repeated measures over time or a mixed-effects model (when data were missing) were utilized for

statistical analyses of urinary biomarkers, plasma creatinine, and PK indices between study groups using GraphPad

Prism (version 8.2.1 for Windows, GraphPad Software, La Jolla, CA). Ordinal logistical regressions on

histopathological scores were performed with observed nominal scores treated as dependent variables and control

group as the referent category using Stata version 13 (40). All tests were two-tailed with an α level of 0.05 for

statistical significance.

Results

Differences between animal cohorts

A total of 32 PB-treated animals contributed PK model data and completed all protocols. All animals weighed

between 291.1 to 321.1 g. Pre-surgery (i.e. day -1) mean (SD) urine volumes were 5.2 (1.7), 8.6 (2.2), 6.4 (2.3), and

5.8 (2.2) mL for QD, BID, TID and control groups, respectively. The QD group had significantly lower urine

volume compared to BID group at day -1 (p=0.01), while no statistical differences in urine output were observed

between experimental groups on days 1, 2, and 3 (Fig. 3). All experimental groups produced significantly more

urine compared to controls on study days 1, 2 and 3, except BID group on day 1 (p=0.23).

PB pharmacokinetic models and exposures

Various modeling approaches were utilized to fit the PK data. A three-compartment model was chosen as the final

model given that it was the most parsimonious with the least bias and imprecision and displayed the most significant

OFV change with the lowest AIC against competing models (Table 1). Table 2 provides a summary of the

population mean parameter values for K_a, V_C, K_{23}, and K_{32}. Model predictive performance for observed vs.

Bayesian posterior-predicted concentrations for bias, imprecision, and R^2 were: 0.129 mg/L, 0.729 mg^2/L^2 and 0.652,

respectively (Fig. 4). PK exposures were calculated based on NCA analysis on the Bayesian posterior-predicted

concentrations from the best-fit PK model and are graphically represented in Fig. 5. Mean AUC$_{24h}$ (SD) for QD,

BID, and TID groups were 171.1 (41.0), 168.6 (58.9), and 129.6 (50.7) mg*h/L, respectively. Similarly, mean (SD)

C$_{MAX}$ for QD, BID, and TID groups were 10.7 (0.80), 9.7 (0.47), and 6.7 (0.49) mg/L, respectively and mean C$_{MIN}$

(SD) were 2.3 (1.3), 3.7 (1.4), and 3.2 (1.2) mg/L, respectively. Statistical procedures were conducted to evaluate

the differences in AUC$_{24h}$, C$_{MAX}$ and C$_{MIN}$ between experimental groups and respective 24-hour intervals. Although

the QD group exhibited higher AUC$_{24h}$ than TID group (p=0.003) on day 1, no significant effects were observed for

overall exposure vs. time (p=0.87). Similarly, the QD group had an overall lower mean C$_{MIN}$ compared to the BID

group (p <0.0001) or TID group (p<0.0001) during the first 24 hours, but no significant differences were observed

between experimental groups over the entire study period (p=1.00). Compared to the TID group, both QD and BID

groups showed significantly higher C$_{MAX}$ during the first 24 hours (p=0.0083 and p=0.049, respectively). The QD

group exhibited the highest C$_{MAX}$ on days 2 and 3 (p=0.0025 and p=0.0017, respectively). On days 2 and 3, the

C_{MAX} in the BID group was numerically elevated when compared to the TID group, but the values were not

statistically significant (p= 0.051 and p=0.055, respectively).

Kidney injury biomarkers and histopathological examinations

Plasma creatinine and urinary biomarkers are graphically represented in Fig. 6. Plasma creatinine did not differ

across treatments over time (p=0.18); however, the following biomarkers showed a significant treatment effect over

the study period: KIM-1 (p<0.0001), OPN (p=0.029), IP-10 (p=0.046), and TIMP-1 (p<0.0001). As a representative

biomarker, KIM-1 rose rapidly on days 1 and 2 for all experimental groups. Notably, the TID group experienced the

largest KIM-1 increase [mean increase (95% CI) day 1 from day -1, 4.44 (0.89, 8.00) ng/mL; p=0.018] as compared

to control [mean increase (95% CI) day 1 from day -1, 0.03 (-0.42, 0.49) ng/mL; p=0.99]. Increases were also

observed with QD [mean increase (95% CI) day 1 from day -1, 2.58 (-0.12, 5.27) ng/mL; p=0.06] and BID groups

[mean difference (95% CI) day 1 from day -1, 0.84 (-0.05, 1.73) ng/mL; p=0.06]. Further, the TID group exhibited

a significant KIM-1 increase on day 2 [mean increase (95% CI), 2.44 (1.22, 3.67) ng/mL; p=0.0013] and a

nonsignificant decrease was observed on day 3 [mean decrease (95% CI), 2.39 (-0.053, 4.83) ng/mL; p=0.055).

Mean KIM-1 changes did not differ on days 2 and 3 between QD and BID groups. Similar trends and significant

treatment and time effects were also observed for OPN (p=0.029), IP-10 (p=0.046), and TIMP-1 (p<0.0001) but no

significant effects were observed with CLN (p=0.093) (Fig. 6).

Histopathological scorings are summarized in Table 3 and graphically displayed in Supplemental Figure S1.

Representative histopathology images are provided in supplemental materials (Figures S2, S3, S4 and S5).

Significant histopathological damage was observed with the TID group with a median [range] score of 2 [2, 3.5]

(p=0.013; 95% CI, 0.70, 5.88) using controls as a referent category. While damage was also observed with QD

[median score (range), 2 (1, 3)] and BID [2 (1, 3)] groups, no statistically significant difference was found when

comparing either group to the referent group [p=0.156 (95% CI, -0.61, 3.77), p=0.092 (95% CI, -0.34, 4.50),

respectively].

Discussion

A dose fractionation scheme in this study effectively maintained constant exposure and separated maximal

concentrations across experimental groups. Our data demonstrated that fractionating the PB dose into three daily

aliquots resulted in the most kidney injury as measured by the urinary biomarkers: KIM-1, OPN, IP-10, and TIMP-1.

Significant differences were detectable within 24 hours. These markers have previously demonstrated high

sensitivity for kidney injury (41-43). Further, KIM-1 and OPN were directly linked to proximal tubular toxicity, and

this is concordant with previous reports (25, 41). Histopathological findings also indicated that thrice daily dosing

of PB led to the most severe kidney insults within the study period. While the differences in urinary biomarkers and

categorical damage scales were not significant for QD and BID groups, both groups demonstrated less extensive

kidney injuries consistent with results from urinary biomarkers such as elevations in KIM-1 levels. Additionally, we

derived a best-fit PK model for PB in rats using rich PK data. Utilizing the best-fit model, we found that the PB

exposure (i.e. AUC_{24h}) derived were similar across all experimental groups (QD, BID, and TID) and separation of

peak concentrations was observed.

Contemporary dosing of intravenous PB recommends it be administered in 2 divided doses based on a weight-based

total daily dose (26, 44). It has been suggested that PB-induced kidney injury could be minimized by optimizing

dosing intervals, similar to that observed with aminoglycosides (7, 45). Wallace *et al.* utilized a preclinical rat

model to explore this possibility for colistin. The authors found that the colistin methanesulfonate (CMS, prodrug of

colistin) regimen corresponding to once daily dosing in humans led to a greater number and severity of renal lesions

when compared to the group received fractionated dosage corresponding to twice daily dosing in human. They

concluded that extended interval dosing of CMS resulted in more extensive renal damage (27). Abdelraouf *et al.*

also utilized a rat model and administered PB subcutaneously at 20 mg/kg/day or 5 mg/kg every 6 hours (25). In

contrast, they found a lower rate of nephrotoxicity associated with PB in the once daily group, while the split-dosage

group experienced a quicker onset of nephrotoxicity (defined by elevation of creatinine from baseline); however, the

rate of nephrotoxicity converged between groups towards the end of study. The authors suggested that an increased

active, saturable, carrier-mediated uptake may have been responsible for this effect when PB was given repeatedly

(as opposed to once daily) and led to the higher rate of renal injury. Most recently, Okoduwa *et al.* conducted a

retrospective, propensity-score matched clinical study on 200 patients who received once-daily or twice-daily

265 systemic PB across different medical centers (28). In contrast to the animal study by Abdelraouf et al., they found

266 that a higher proportion of nephrotoxicity (using clinical criteria and considering all stages of AKI) was observed in

267 the once-daily group than the twice-daily group (47% vs. 17%, respectively; p=0.0005). The findings from

268 Okoduwa et al. are consistent with the animal model employed by Wallace et al., though CMS (and not PB) was

269 utilized; however, it is unclear whether patients receiving once daily PB in clinical studies actually received

270 equivalent (or greater) exposures compared to those receiving twice daily dosing. Our findings agree with the

271 results from Abdelraouf et al. that dose fractionated PB strategy led to more extensive AKI. Additionally, our study

272 provided rich PK data to confirm the exposure status across experimental groups and employed highly sensitive and

273 specific urinary biomarkers for early detection of AKI in addition to plasma creatinine (46). The level of injury was

274 further confirmed by histopathological examination.

276 We acknowledge several limitations to our study. First, our study was limited to 72-hour dosing compared to the

277 relatively longer study period (up to 10 days) by Abdelraouf et al. The shorter time frame did not allow us to

278 observe levels of plasma creatinine nor urinary biomarkers beyond this time frame; however, the urinary biomarkers

279 utilized are highly sensitive in detecting early stages of AKI and our histopathology examinations confirmed the

280 injury (30, 31, 46-48). Secondly, PB exposure (i.e. AUC_{24h}) was held constant in our study and linking exposure to

281 toxicodynamic data (i.e. injury biomarkers) would not be ideal as this was not an objective of our study. Thus,

282 further studies are warranted to examine the PK/PD indices to toxicodynamic outcomes. Thirdly, this is a pre-

283 clinical model and additional translational studies defining the lower limit of PB toxicity are needed to design

284 maximally safe and effective dosing regimens.

286 To date, this is the first study that employed a rat model with a rich PK sampling design with dose fractionated

287 systemic PB that also allowed PK estimates at an individual level. We also demonstrated that TID dosing of PB

288 induces AKI as early as 24 hours. These findings may have clinical implications for PB dosing schemes in difficult-

289 to-treat infections while minimizing nephrotoxicity. Further studies are warranted to explore PB exposure linked to

290 toxicity while maximizing efficacy.

References

(1) Waterer, G.W. & Wunderink, R.G. Increasing threat of Gram-negative bacteria. *Crit Care Med* **29**, N75-81 (2001).

(2) Aslam, B. *et al.* Antibiotic resistance: a rundown of a global crisis. *Infect Drug Resist* **11**, 1645-58 (2018).

(3) CDC. Antibiotic Resistance Threats in the United States, 2019. Atlanta, GA: U.S. Department of Health and Human Services, CDC; 2019.

(4) Resistance, I.C.G.o.A. No Time to Wait: Securing the future from drug-resistant infections. Report to the Secretary-General of the United Nations. 2019.

(5) MacVane, S.H., Pandey, R., Steed, L.L., Kreiswirth, B.N. & Chen, L. Emergence of Ceftolozane-Tazobactam-Resistant Pseudomonas aeruginosa during Treatment Is Mediated by a Single AmpC Structural Mutation. *Antimicrob Agents Chemother* **61**, (2017).

(6) Shields, R.K., Nguyen, M.H., Press, E.G., Chen, L., Kreiswirth, B.N. & Clancy, C.J. Emergence of Ceftazidime-Avibactam Resistance and Restoration of Carbapenem Susceptibility in Klebsiella pneumoniae Carbapenemase-Producing K pneumoniae: A Case Report and Review of Literature. *Open Forum Infect Dis* **4**, ofx101 (2017).

(7) Trusts, T.P.C. Antibiotics Currently in Global Clinical Development. (2019).

(8) Evans, M.E., Feola, D.J. & Rapp, R.P. Polymyxin B sulfate and colistin: old antibiotics for emerging multiresistant gram-negative bacteria. *Ann Pharmacother* **33**, 960-7 (1999).

(9) Thomas, T.A. *et al.* High performance liquid chromatography-mass spectrometry assay for polymyxin B1 and B2 in human plasma. *Ther Drug Monit* **34**, 398-405 (2012).

(10) Falagas, M.E. & Kasiakou, S.K. Colistin: the revival of polymyxins for the management of multidrug-resistant gram-negative bacterial infections. *Clin Infect Dis* **40**, 1333-41 (2005).

(11) Yahav, D., Farbman, L., Leibovici, L. & Paul, M. Colistin: new lessons on an old antibiotic. *Clin Microbiol Infect* **18**, 18-29 (2012).

(12) Koch-Weser, J., Sidel, V.W., Federman, E.B., Kanarek, P., Finer, D.C. & Eaton, A.E. Adverse effects of sodium colistimethate. Manifestations and specific reaction rates during 317 courses of therapy. *Ann Intern Med* **72**, 857-68 (1970).

(13) Tamma, P.D. *et al.* The use of intravenous colistin among children in the United States: results from a multicenter, case series. *Pediatr Infect Dis J* **32**, 17-22 (2013).

(14) Falagas, M.E. & Kasiakou, S.K. Toxicity of polymyxins: a systematic review of the evidence from old and recent studies. *Crit Care* **10**, R27 (2006).

(15) Justo, J.A. & Bosso, J.A. Adverse reactions associated with systemic polymyxin therapy. *Pharmacotherapy* **35**, 28-33 (2015).

(16) Zavascki, A.P. & Nation, R.L. Nephrotoxicity of Polymyxins: Is There Any Difference between Colistimethate and Polymyxin B? *Antimicrob Agents Chemother* **61**, (2017).

(17) Crass, R.L., Rutter, W.C., Burgess, D.R., Martin, C.A. & Burgess, D.S. Nephrotoxicity in Patients with or without Cystic Fibrosis Treated with Polymyxin B Compared to Colistin. *Antimicrob Agents Chemother* **61**, (2017).

(18) Phe, K. *et al.* In vitro assessment and multicenter cohort study of comparative nephrotoxicity rates associated with colistimethate versus polymyxin B therapy. *Antimicrob Agents Chemother* **58**, 2740-6 (2014).

(19) Temocin, F., Erdinc, S., Tulek, N., Demirelli, M., Bulut, C. & Ertem, G. Incidence and Risk Factors for Colistin-Associated Nephrotoxicity. *Jpn J Infect Dis* **68**, 318-20 (2015).

(20) Tsuji, B.T. *et al.* International Consensus Guidelines for the Optimal Use of the Polymyxins: Endorsed by the American College of Clinical Pharmacy (ACCP), European Society of Clinical Microbiology and Infectious Diseases (ESCMID), Infectious Diseases Society of America (IDSA), International Society for Anti-infective Pharmacology (ISAP), Society of Critical Care Medicine (SCCM), and Society of Infectious Diseases Pharmacists (SIDP). *Pharmacotherapy* **39**, 10-39 (2019).

(21) Suzuki, T., Yamaguchi, H., Ogura, J., Kobayashi, M., Yamada, T. & Iseki, K. Megalin
contributes to kidney accumulation and nephrotoxicity of colistin. *Antimicrob Agents Chemother*
57, 6319-24 (2013).
(22) Kelesidis, T. & Falagas, M.E. The safety of polymyxin antibiotics. *Expert Opin Drug Saf* **14**,
1687-701 (2015).
(23) Yousef, J.M., Chen, G., Hill, P.A., Nation, R.L. & Li, J. Melatonin attenuates colistin-induced
nephrotoxicity in rats. *Antimicrob Agents Chemother* **55**, 4044-9 (2011).
(24) Yousef, J.M., Chen, G., Hill, P.A., Nation, R.L. & Li, J. Ascorbic acid protects against the
nephrotoxicity and apoptosis caused by colistin and affects its pharmacokinetics. *J Antimicrob*
Chemother **67**, 452-9 (2012).
(25) Abdelraouf, K., Braggs, K.H., Yin, T., Truong, L.D., Hu, M. & Tam, V.H. Characterization of
polymyxin B-induced nephrotoxicity: implications for dosing regimen design. *Antimicrob Agents*
Chemother **56**, 4625-9 (2012).
(26) Nation, R.L., Rigatto, M.H.P., Falci, D.R. & Zavascki, A.P. Polymyxin Acute Kidney Injury:
Dosing and Other Strategies to Reduce Toxicity. *Antibiotics (Basel)* **8**, (2019).
(27) Wallace, S.J. *et al.* Subacute toxicity of colistin methanesulfonate in rats: comparison of various
intravenous dosage regimens. *Antimicrob Agents Chemother* **52**, 1159-61 (2008).
(28) Okoduwa, A. *et al.* Nephrotoxicity Associated with Intravenous Polymyxin B Once- versus
Twice-Daily Dosing Regimen. *Antimicrob Agents Chemother* **62**, (2018).
(29) Zavascki, A.P. Polymyxins for the treatment of extensively-drug-resistant Gram-negative bacteria:
from pharmacokinetics to bedside. *Expert Rev Anti Infect Ther* **12**, 531-3 (2014).
(30) O'Donnell, J.N. *et al.* 24-Hour Pharmacokinetic Relationships for Vancomycin and Novel
Urinary Biomarkers of Acute Kidney Injury. *Antimicrob Agents Chemother* **61**, (2017).
(31) Rhodes, N.J. *et al.* Evaluation of Vancomycin Exposures Associated with Elevations in Novel
Urinary Biomarkers of Acute Kidney Injury in Vancomycin-Treated Rats. *Antimicrob Agents*
Chemother **60**, 5742-51 (2016).
(32) Lim, T.P. *et al.* Physicochemical Stability Study of Polymyxin B in Various Infusion Solutions
for Administration to Critically Ill Patients. *Ann Pharmacother* **50**, 790-2 (2016).
(33) Sharma, V. & McNeill, J.H. To scale or not to scale: the principles of dose extrapolation. *Br J*
Pharmacol **157**, 907-21 (2009).
(34) Prozialeck, W.C. *et al.* Kidney injury molecule-1 is an early biomarker of cadmium
nephrotoxicity. *Kidney Int* **72**, 985-93 (2007).
(35) Rhodes, N.J., Liu, J., McLaughlin, M.M., Qi, C. & Scheetz, M.H. Evaluation of clinical outcomes
in patients with Gram-negative bloodstream infections according to cefepime MIC. *Diagn*
Microbiol Infect Dis **82**, 165-71 (2015).
(36) Sistare, F.D. *et al.* Towards consensus practices to qualify safety biomarkers for use in early drug
development. *Nat Biotechnol* **28**, 446-54 (2010).
(37) Vaidya, V.S. *et al.* Kidney injury molecule-1 outperforms traditional biomarkers of kidney injury
in preclinical biomarker qualification studies. *Nat Biotechnol* **28**, 478-85 (2010).
(38) Leary, R.J., Roger & Schumitzky, Alan & Van Guilder, M. . An adaptive grid non-parametric
approach to pharmacokinetic and dynamic (PK/PD) population models. *Proceedings of the IEEE*
Symposium on Computer-Based Medical Systems, 389-94.
(39) Neely, M.N., van Guilder, M.G., Yamada, W.M., Schumitzky, A. & Jelliffe, R.W. Accurate
detection of outliers and subpopulations with Pmetrics, a nonparametric and parametric
pharmacometric modeling and simulation package for R. *Ther Drug Monit* **34**, 467-76 (2012).
(40) StataCorp. 2013. Stata Statistical Software: Release 13. College Station, TX: StataCorp LP.
(41) Vaidya, V.S., Ferguson, M.A. & Bonventre, J.V. Biomarkers of acute kidney injury. *Annu Rev*
Pharmacol Toxicol **48**, 463-93 (2008).
(42) Musial, K. & Zwolinska, D. Pleiotropic functions of TIMP-1 in patients with chronic kidney
disease. *Cell Mol Life Sci* **71**, 1547-8 (2014).

(43)   Erez, D.L. *et al.* Urinary CXCL10 and CXCL9 Are Associated with Acute Kidney Injury in
392   Children after Hematopoietic Stem Cell Transplantation: Results of a Discovery and Validation
393   Cohort. *Biology of Blood and Marrow Transplantation* **25**, S185-S6 (2019).
394   (44)   Polymyxin B for injection USP [package insert]. Oakville, ON: SteriMax Inc.; 2016.
395   (45)   Barza, M., Ioannidis, J.P., Cappelleri, J.C. & Lau, J. Single or multiple daily doses of
396   aminoglycosides: a meta-analysis. *BMJ* **312**, 338-45 (1996).
397   (46)   Avedissian, S.N. *et al.* Twenty-four hour pharmacokinetic relationships for intravenous
398   vancomycin and novel urinary biomarkers of acute kidney injury in a rat model. *J Antimicrob*
399   *Chemother* **74**, 2326-34 (2019).
400   (47)   Joshi, M.D. *et al.* Evaluation of Fetal and Maternal Vancomycin-Induced Kidney Injury during
401   Pregnancy in a Rat Model. *Antimicrob Agents Chemother* **63**, (2019).
402   (48)   Pais, G.M. *et al.* Comparative Performance of Urinary Biomarkers for Vancomycin-Induced
403   Kidney Injury According to Timeline of Injury. *Antimicrob Agents Chemother* **63**, (2019).

1   **Table 1. Model selection summary**

Compartmental Models	-2LL	OFV Change	AIC	Bias (mg/L)	Imprecision (mg^2/L^2)	Bayesian R^2
One	1709	Ref	1715	-0.48	0.95	0.10
Two	1709	0	1720	-0.05	0.90	0.20
Three	1450	259	1463	0.13	0.73	0.65

2   *One-compartment model served as the base model to derive two- and three-compartment models;*
3   *three-compartment model is the final model*

4   **Table 2. Mean population parameters from the final model**

	Mean	SD
K_a (h^{-1})	0.290	0.460
K_e (h^{-1})	0.411	0.076
V_C (L)	0.056	0.079
K_{23} (h^{-1})	6.214	11.430
K_{32} (h^{-1})	3.163	43.00

6 **Table 3. Ordinal logistic regression of histopathology scorings by groups**

	Median Score [range]	p value	95% CI
Control	1.5 [1-2]	Referent	Referent
QD	2 [1-3]	0.156	-0.61-3.77
BID	2 [1-3]	0.092	-0.34-4.50
TID	2 [2-3.5]	0.013	0.70-5.88

1   **Figure 1.**
2

3   **a)**   **Allocation of experimental groups.**

5   *QD: once daily; BID: twice daily; TID: thrice daily*
6   *Control: once daily protocol*

Figure 2. Schematic of study design for each study group.

Day 0 depicts surgical cannulation of catheter for blood sampling

*Kidney image from Open Michigan

Figure 3. Mean 24-h urine volume between experimental and control groups

15   **Figure 4. Goodness-of-fit plot for Bayesian observed vs. predicted plasma PB concentrations utilizing**
16   **the final three-compartment model.**

17
18

19   **Figure 5. Violin plots of PK indices from the final best-fit model by days**

20   **a) 24-h AUC**

22   **b) 24-h C$_{MAX}$**

24   **c) 24-h C$_{MIN}$**

 Figure 6. Plasma creatinine and urinary biomarkers

Supplemental Materials

Figure S1. Violin plot of histopathological scores between all groups

Figure S2. QD group histopathological examination images

a) **renal tubular regeneration (arrows) & vacuolation (arrowheads); score = 2**

b) **renal tubular degeneration (arrows) and regeneration (arrowheads); score = 3**

Figure S3. BID group histopathological examination images

a) **renal tubular regeneration (arrows) & dilatation (arrowheads); score = 2**

b) **renal tubular regeneration (arrows), cell sloughing (circle) & vacuolation (arrowheads); score = 3**

Figure S4. TID group histopathological examination images: renal tubular regeneration (arrows) & vacuolation (arrowheads); score = 3

Figure S5. Control group: renal tubular regeneration (arrows); score = 2