Adaptive divergence in shoot gravitropism creates intrinsic reproductive isolation in an

Australian wildflower

Melanie J. Wilkinson[1†], Federico Roda[1,2†], Greg M. Walter[1,3], Maddie E. James[1], Rick Nipper[4], Jessica

Walsh[4], Scott L. Allen[1], Diana M. Bernal[1], Henry L. North[1,5], Christine A. Beveridge[1] and Daniel Ortiz-

Barrientos[1*].

[1]The University of Queensland, School of Biological Sciences, St Lucia QLD 4072, Australia. [2]Current

address: Universidad Nacional de Colombia, Departamento de Biología, Bogotá, Colombia. [3]Current

address: Monash University, School of Biological Sciences, Clayton Vic 3800. [4]Floragenex, Inc., 4640

SW Macadam Avenue, Suite 200F, Portland, OR 97239, USA. [5]Current address: University of

Cambridge, Department of Zoology, Downing St., Cambridge CB2 3EJ, UK.

† These authors contributed equally to the paper.

* Author for correspondence: Daniel Ortiz-Barrientos

Email: d.ortizbarrientos@uq.edu.au

Abstract

Adaptation to the local environment is a major driver of speciation. Yet, it remains largely unknown whether natural selection directly causes intrinsic reproductive isolation (hybrid sterility or inviability) between locally adapted populations. Here, we show that adaptive divergence in shoot gravitropism, the ability of a plant's shoot to bend upwards in response to the downward pull of gravity, contributes to the evolution of intrinsic reproductive isolation in an Australian wildflower, *Senecio lautus*. We find that shoot gravitropism has evolved multiple times in association with plant height between adjacent populations inhabiting contrasting environments, suggesting that these traits have evolved by natural selection. We directly tested this prediction in multi-year reciprocal transplant experiments using hybrid populations. We show that shoot gravitropism and plant height respond to natural selection in the expected direction of the locally adapted population. Remarkably, we find that crossing F11 hybrid individuals with extreme differences in gravitropism significantly reduces their ability to produce seeds, providing strong evidence that this adaptive trait is genetically correlated with intrinsic reproductive isolation. Our results suggest that natural selection can drive the evolution of locally adaptive traits that incidentally create intrinsic reproductive isolation, thus increasing our understanding of the origin and maintenance of new species.

Main

Ever since Darwin's work on the origin of species by natural selection[1], researchers have sought to understand how natural selection creates reproductive barriers between populations[2]. On one hand, many studies have established that adaptation to contrasting environments often reduces migrant and hybrid fitness in the wild, a process commonly known as extrinsic reproductive isolation[3]. These extrinsic barriers to reproduction can dramatically reduce gene flow between populations[4,5]; however, they only act in the local environment of populations and therefore are susceptible to changes in environmental conditions. As a consequence, it is unclear whether the evolution of extrinsic reproductive isolation alone can complete speciation[4]. On the other hand, intrinsic reproductive barriers such as hybrid sterility or inviability, can accumulate regardless of environmental change and are therefore expected to be more stable over time and contribute more reliably to the completion of speciation[2]. One way in which natural selection can simultaneously create both extrinsic and intrinsic reproductive isolation is when the beneficial mutations that drive local adaptation in each population also fail to interact properly between populations through negative epistasis[6] (i.e., Dobzhansky-Muller genetic incompatibilities). Therefore, natural selection could in principle reduce gene flow through multiple extrinsic and intrinsic barriers between populations, and thus genetically link the beginning and completion of speciation[7].

Although several genes that impact plant hybrid sterility and inviability have been identified[8], there is limited evidence supporting local adaptation as the major process responsible for the evolution of these loci. In one notable exception, some intraspecific crosses in *Arabidopsis thaliana* produce dwarf, necrotic, and sterile hybrid individuals as a result of negative epistatic interactions from divergent selection of pathogen resistance genes[9]. In a second example in *Mimulus*, tight genetic linkage between alleles selected for copper tolerance and alleles that cause hybrid mortality has led to divergence between populations growing next to copper mines and those occupying typical *Mimulus* habitats[10]. Although plant-pathogen coevolution and tight linkage between genes performing different functions (e.g., stress tolerance and seed development) could be powerful mechanisms for driving species divergence[11], we need more studies on the genetics of adaptation and speciation to reveal the mechanisms by which natural selection can drive the origin of new species. Here, we introduce a novel example where we take advantage of the repeated evolution of divergent growth

habits – adjacent erect and prostrate populations – to study the contribution of adaptation to the

evolution of intrinsic reproductive barriers in an Australian wildflower, *Senecio lautus* (G.Forst. ex

Willd)[12,13]. This highly replicated system of evolution not only implies a role for natural selection in the

evolution of complex morphological or physiological differences[14,15], but also provides multiple

independent opportunities to test whether loci that contribute to local adaptation also create genetic

incompatibilities that could lead to speciation.

Previous population genetic studies in *S. lautus* found that different sets of genes related by similar

functions were repeatedly differentiated in populations with these contrasting growth habits[16,17]. One

of these sets contained genes belonging to the auxin pathway, where genes related to the transport

and regulation of this key plant hormone had repeatedly diverged between erect and prostrate

populations. In congruence, studies across the plant kingdom have established that the auxin

hormone plays a major role in how plants orient themselves with respect to light and gravity cues[18]

and in creating variation in height[19,20], branching[21,22] and reproduction[23]. With these three traits

differentiating erect and prostrate *S. lautus* populations, the auxin pathway is a compelling candidate

to link the molecular basis of adaptation and speciation in *S. lautus*. We reasoned that if divergence in

the auxin pathway contributed to the evolution of local adaptation and intrinsic reproductive isolation

in *S. lautus*, we would observe the following three predictions. 1) In multiple populations with distinct

growth habits, we would consistently detect divergence in auxin-dependent traits, such as differences

in the ability of seedlings to alter the direction of growth against the pull of gravity[18,24]. 2) These auxin-

dependent traits would respond to natural selection in the local environment. Finally, 3) crosses

between hybrid individuals with extreme differences in auxin-dependent traits would be genetically

incompatible, thus demonstrating a clear genetic link between local adaptation and intrinsic

reproductive isolation.

We test these predictions on multiple coastal populations of *S. lautus* (Fig. 1a, Table S1), where there

is evidence of local adaptation and intrinsic reproductive isolation between adjacent Dune and

Headland populations[25-29]. Populations inhabiting sand dunes (Dune hereafter) are erect, while

populations growing on adjacent rocky headlands (Headland hereafter) are prostrate (Fig. 1b). Erect

and prostrate growth habits can also be found in related populations from the alpine regions of

Australia, with a prostrate population inhabiting an exposed alpine meadow and an erect population

inhabiting adjacent sheltered alpine gullies (Fig. 1c). Dune populations are continually exposed to

high temperatures, high solar radiation, low salinity, and low nutrient sand substrate, whereas

Headland populations are exposed to high salinity, high nutrients and powerful winds[28]. These

ecotypes are genetically grouped into two major monophyletic clades based on geography (eastern

and south-eastern Australia), and adjacent Dune and Headland populations are often sister taxa,

suggesting that they have evolved their contrasting growth habits independently multiple

times[12,25,30,31]. These Dune and Headland ecotypes also show variable levels of intrinsic reproductive

isolation, where hybrid sterility in F1 hybrids is weak[28] (<10% crossing failure, n=65), but strong in F2

hybrids (58% crossing failure, n=163). Genetic incompatibilities are largely removed in one generation

as F3 hybrids also have weak hybrid sterility (similar to F1 hybrids)[28]. The inability to produce seed in

F2's is therefore likely due to few, yet strong recessive negative epistatic interactions between these

ecotypes[32,33]. Furthermore, demographic analyses recently revealed minimal levels of gene flow

between adjacent Dune and Headland populations[30], indicating that reproductive barriers (intrinsic or

extrinsic) can prevent hybridisation from occurring in the field. Given the presence of reproductive

isolation between multiple locally adapted erect and prostrate populations, *S. lautus* provides an

excellent system in which to understand whether local adaptation driven by morphological and

physiological traits can lead to the accumulation of intrinsic reproductive barriers.

Results

Repeated evolution of auxin-related genes in *Senecio lautus*

We first investigated if auxin-related genes are associated with variation in three growth-habit traits

(plant height, main stem angle and the number of branches; Fig. S1) that have evolved repeatedly in

S. lautus[31]. Specifically, we created four genetic mapping populations representing different origins of

erect and prostrate plant architectures by crossing individuals from each of the four pairs of Dune and

Headland populations[31] (Fig. S2). Bulk Segregant Analyses (BSA)[34] were used to compare the

genomes of pools of plants with extreme values for the three traits in the four mapping populations. In

agreement with previous population genetic studies in *S. lautus*[16], we found that highly differentiated

genomic regions (candidates) between hybrid individuals with contrasting traits were often not the

same across the different mapping populations (Data S1). However, the overlap between candidates

in different populations was higher than expected by chance (number of shared Single Nucleotide

Polymorphisms (SNPs) = 2,548; mean for permutations = 1,770; 99% upper CI = 1,897) and we

discovered 20 candidate regions shared across all four mapping populations. Out of these 20

repeatedly differentiated candidate regions, 11 contained genes regulating the auxin pathway (Table

S2), and four of these 11 candidate regions contained multiple auxin-related genes.

Auxin candidate genes identified in the BSA were also genetically differentiated between adjacent

natural populations, and had significant associations with survivorship in reciprocal transplants where

we examined shifts in allelic frequency after selection in the wild (Table S2 and Data S1). Additionally,

several of these auxin genes displayed differential expression between the shoot apical meristems of

erect and prostrate populations, which suggest that they are important for morphogenesis (Fig. S3).

The homologs of the BSA auxin candidate genes include genes involved in intracellular auxin

transport (Table S2), an essential process for creating auxin concentration gradients that drive a

directional growth response to environmental cues such as light (phototropism) or gravity

(gravitropism)[24,35]. Some homologs also have pleiotropic effects on reproduction (Table S2), which is

consistent with a previous mapping experiment between erect and prostrate populations of *S.*

lautus[36], where quantitative trait loci for plant architecture (e.g. plant height) clustered with the total

number of flowers an individual produces. Together with previous results, these results strengthen the

conjecture that the auxin pathway is important for adaptation to sand dunes and rocky headlands, and

that there might be a link between complex adaptations and reproductive barriers between genetically

and morphologically divergent populations of *S. lautus*.

Repeated height and gravitropism divergence across *Senecio lautus* populations

The regulation and transport of auxin is a well-established modulator of gravitropism in plants[37,38], and

therefore might also contribute to growth habit in *S. lautus*. To test whether plant height predicts

gravitropism across *S. lautus* populations, we first quantified plant height and gravitropism in 16 *S.*

lautus populations (Fig. 1 and Fig. S5). We defined the gravitropic response as the change in angle of

a seedling's stem 24 hours after a 90° rotation[39,40], where 0° is no gravitropic response and 90° is a

complete re-orientation. If differences in the auxin pathway have contributed to divergence in growth

habit, we expected that prostrate populations would display a smaller gravitropic angle than their

adjacent erect population. We found that plant height in natural environments (field) strongly

correlates with estimates of plant height in common garden conditions (glasshouse) ($F_{1,9}$=12.41,

P=0.0065), demonstrating a genetic basis of this trait in *S. lautus*. Across these populations, the

average height of a population predicted its average gravitropic angle (Fig. 2a; $F_{1,13}$=17.65,

P=0.0010), with the average magnitude of gravitropism differing between the two monophyletic clades

of this system ($F_{1,13}$=32.58, P<0.0001). Therefore, changes in gravitropism appear to be genetically

correlated with divergent growth habit traits such as plant height, and this relationship has evolved at

least once in each clade.

To test whether natural selection drove the evolution of differences between locally adapted

populations with differences in plant height, we first compared gravitropism between seven adjacent

Dune and Headland population pairs (Table S1). Four population pairs showed the expected

correlation, where plants from the Headland population exhibited a smaller gravitropic angle and were

shorter than plants from their adjacent Dune population (Fig. 2b, c). In two population pairs – Millicent

and Point Labatt – plant height did not differ between Dune and Headland population pairs (Fig. 2c),

possibly due to their similarity in environmental variables[31]. In the population pair at Stradbroke Island,

however, we observed differences in height but not in gravitropism (Fig. 2b), suggesting that similar

plant architectures could have evolved through other convergent developmental routes in coastal

populations of *S. lautus*. We assessed height and gravitropism in divergent populations from the

alpine region of Australia, and also discovered the expected pattern, where the exposed Alpine

population in the meadow was shorter and had a smaller gravitropic angle than the population in the

sheltered alpine gully (Fig. 2c). These results suggest that natural selection repeatedly acting on

auxin-related genes has contributed to differences in gravitropism across coastal and alpine

populations in this system.

Auxin drives gravitropic differences in *Senecio lautus*

To directly test whether synthetic auxin and auxin transport inhibitors would influence gravitropism

differently between erect and prostrate populations, we grew Dune (n=90) and Headland (n=98)

seeds collected from Lennox Head with synthetic auxin 2,4-D (2,4-Dichlorophenoxyacetic acid), and

polar auxin transport inhibitor NPA (naphthylphthalamic acid)[41,42]. Because a gravitropic response

requires an auxin concentration gradient, we reasoned that removing the gradient would reduce the

gravitropic angle in Dune individuals (Fig. S4). As expected, addition of synthetic auxin 2,4-D reduced

the gravitropic angle more in Dune individuals than in Headland individuals (LR chi-square=18.49,

P<0.0001). Similarly, addition of auxin transport inhibitor NPA reduced the gravitropic angle in our

experiments more in Dune individuals than in Headland individuals (LR chi-square=21.18, P<0.0001).

This difference in hormone response in Dune and Headland individuals supports our hypothesis that

differences in the auxin pathway contribute to the different gravitropic responses observed in *S.*

lautus.

Local adaptation drives the evolution of height and gravitropism

We next conducted two independent sets of field adaptation experiments to assess the role of natural

selection on the evolution of auxin-dependent traits, height and gravitropism. In each field experiment,

Dune and Headland parental and hybrid seeds were transplanted into replicated blocks at the sand

dune and rocky headland at Lennox Head. In all field experiments, natural selection consistently

favoured the local Dune or Headland population over the foreign population (Fig. 3), indicating that

our experiment was exposed to the divergent selection that drove local adaptation in each

environment.

For the height adaptation experiments, we tested whether differences in height could drive differences

in fitness in the rocky headlands. We hypothesised that offspring produced by short hybrid parents

would live longer than offspring produced by tall hybrid parents in the rocky headland. We focused on

prostrate growth because it is likely the derived trait, given that the majority of *S. lautus* populations

have an erect growth habit[16,28]. Our goal was to introgress Dune alleles associated with height onto a

Headland genomic background to examine their effect on fitness in the headland environment. Briefly,

we crossed Dune and Headland seeds from Lennox Head and then completed two rounds of

backcrossing followed by one round of crossing between tall hybrid offspring (tallest 10%) and

between short hybrid offspring (shortest 10%; Table S3 and Fig. S6). We transplanted 558 of these

seeds (from 28 families) into the rocky headland. As predicted, shorter hybrid parents produced

offspring that lived longer in the rocky headland relative to offspring from taller hybrid parents

($F_{1,26.23}$=4.87, P=0.0362). These results suggest that traits genetically correlated with plant height

contributed to early developmental fitness in the rocky headlands and contributed directly or indirectly

to the evolution of divergent growth habit.

For the gravitropism adaptation experiments, we tested whether rapid adaptation to contrasting

environments can lead to the evolution of gravitropic responses in the direction of the local population.

We hypothesised that exposing an advanced recombinant population to multiple rounds of viability

selection in the field would drive genetic covariation between fitness and gravitropism. To test this

hypothesis, we crossed 23 Dune and 22 Headland individuals from Lennox Head and then

maintained three independent genetic lines for eight generations (Table S4). This process

disassembled allelic combinations and constructed an F8 recombinant hybrid population[36] (Fig. 4a).

We planted 2,403 of these F8 seeds (from 89 families) into both the sand dune and rocky headland

(Fig. 4b and Table S5) and conducted family-based truncation selection for three generations. The

fittest families in each environment were selected based on the highest germination and survival (top

50%; see Methods for selection details). Full-siblings from these families were crossed (within a

genetic line) to produce the next generation. In the F10 generation, we tested whether families that

had the greatest number of survivors also produced offspring (F11) with the local gravitropic response

under controlled conditions.

In agreement with our hypothesis, F10 families with the greatest number of survivors in the sand dune

produced F11 offspring with a higher gravitropic angle (Table 1). We discovered that this relationship

between fitness and gravitropism was driven by the fitness of the F10 dam family and not the F10 sire

family (Table 1), suggesting maternal effects might contribute to the evolution of gravitropism in the

sand dunes. In contrast, in the rocky headland we did not detected an association between the fitness

of the F10 families and gravitropism in their offspring. Instead, Headland survivors had a positive

association between the number of days until death in a controlled environment (intrinsic viability) and

gravitropism, where individuals that died early in development had a smaller gravitropic angle

(agravitropic). We are left with the conjecture that there could be intrinsic fitness costs for agravitropic

alleles on a hybrid genomic background.

We then tested whether adaptation to contrasting environments can recreate trait correlations

observed in nature. As shown in the previous section, there is a strong correlation between

gravitropism and height across many natural populations of *S. lautus*. We expect the correlation

would be lost in the creation of the hybrid population if the traits are not pleiotropic, but if natural

selection for these traits was strong it could be reconstructed after three rounds of selection in the

rocky headland and sand dune. There was no genetic correlation between height and gravitropism in

the control population grown only in the glasshouse ($F_{1,114.3}=0.08$, $P=0.7801$, $r^2=0.04$), but as

predicted this genetic correlation was strong after three rounds of selection in the rocky headland

($F_{1,169.5}=7.09$, $P=0.0085$, $r^2=0.27$) and weak after selection in the sand dune ($F_{1,151.3}=3.20$, $P=0.0756$,

$r^2=0.09$). Together, these results indicate that natural selection can act on standing genetic variation

and reconstitute genetic architectures and trait combinations favoured in these coastal environments.

The genetics underlying gravitropism in *Senecio lautus*

To better understand the genetics of adaptation, we performed selective genotyping on the tails of the

gravitropic distribution of the F11 hybrid populations (Fig. 4c; Dune survivors, Headland survivors and

Control). We identified candidate gene regions containing the most extreme allelic differences

between individuals with the smallest gravitropic angle (agravitropic tail, <20°; mean of tail =

6.46±1.10°, n=68) and the largest gravitropic angle (gravitropic tail, >56°; mean of tail = 62.03±0.45°,

n=77). We found 55 sites (0.2% of all SNPs) across 49 genomic contigs (Data S2) with an allelic

difference in the 99.8% quantile (0.15<*p*<0.22). We discovered that these candidate gene regions

disproportionally contained homologous genes with predicted gene ontology categories of transport

and localisation of molecules within and between cells (Table S6). This is consistent with

expectations, as redistribution of auxin is required for a gravitropic response[18]. Five of the 55 sites

(11%) are located in gene homologs with functions related to the auxin pathway, including the second

(*ENODL1*; early nodulin-like protein 1)[43] and fourth (*ABA3*; molybdenum cofactor sulfurase)[44] most

differentiated SNPs between the agravitropic and gravitropic F11 tails.

Auxin-related genes, *ENODL1* and *ABA3*, have likely contributed to the adaptive divergence of

gravitropism in *S. lautus*. In both the *ENODL1* and *ABA3* genes, natural selection recreated the

expected allelic frequency shift in the F11 tails towards the parent with the same trait (Data S2). For

instance, the alleles favoured in the agravitropic F11 tail were at high frequencies in the Headland

natural population (*ENODL1* cytosine allele (C)=0.87 and *ABA3* guanine allele (G)=0.89), with the

alternate alleles being favoured in the gravitropic F11 tail and the Dune natural population (*ENODL1*

adenine allele (A)=0.69 and *ABA3* A=0.97). *ENODL1* and *ABA3* were in strong linkage disequilibrium

in the survivors of the gravitropism adaptation experiment in the rocky headland (Fisher's exact test,

n=57, P=0.0008) and sand dune (Fisher's exact test, n=48, P=0.0107), but not in the control

population reared in the glasshouse (Fisher's exact test, n=37, P=0.4093), suggesting natural

selection has likely reconstituted this favourable allelic combination. *ENODL1* C/C and *ABA3* G/G

genotypes were associated with a reduction in gravitropism of 25.10° relative to all other genotype

combinations (Fig. 5; $t_{34.30}$=4.86, P<0.0001), suggesting that not sensing the pull of gravity, or reacting

to it, is a recessive trait, and these two loci likely contribute to the genetic architecture responsible for

the gravitropic difference between the F11 tails, and potentially in nature. *ENODL1* and *ABA3* have

gene homologs with functions in not only gravitropism and plant height, but also in salt tolerance and

pollen tube reception[43-49], therefore suggesting that evolution of auxin-related genes might not only

explain the adaptive evolution of gravitropism, but also the evolution of other known trait differences

between Dune and Headland populations. Overall, our results suggest that divergence in auxin-

regulated molecular processes contributed to local adaptation to contrasting environments in coastal

populations of *S. lautus*.

The consequences of gravitropism divergence for the evolution of intrinsic reproductive

isolation

We are now in a position to investigate whether intrinsic reproductive isolation arises as a by-product

of local adaptation. We used the F11 hybrids to test whether divergent gravitropic forms are

reproductively isolated. If gravitropism was contributing directly to the intrinsic reproductive isolation

previously observed between *S. lautus* ecotypes[28,50], we expected to observe increased hybrid

sterility between F11 gravitropic and agravitropic individuals. Hybrid sterility was defined as fewer

than three seeds produced per flower head with at least three mating attempts for reciprocal crosses.

We created two groups of crosses among F11 families: crosses within (agravitropic × agravitropic,

n=28; gravitropic × gravitropic, n=37) and crosses between (agravitropic × gravitropic, n=67) the tails

of the gravitropic distribution. Consistent with gravitropism contributing to intrinsic reproductive

isolation in *S. lautus*, we found that hybrid sterility was more common (Odd's ratio=4.8x, P=0.0188) in crosses between F11 agravitropic and gravitropic plants (21%) than within each of these tails (5%; LR chi-square=6.86, P=0.0088). Five percent (n=3) of within-tail crosses failed, all of which were in the gravitropic tail. Overall this indicates that gravitropism alleles have linked or pleiotropic effects on intrinsic reproductive isolation in *S. lautus.*

To understand what might be driving this association between gravitropism and intrinsic reproductive isolation, we examined whether genetic line, F11 population or specific individuals were correlated with hybrid sterility. Crosses within (n=77) and between (n=55) the three genetic lines, regardless of their gravitropic response, did not have an effect on hybrid sterility (LR chi-square=1.10, P=0.2937). This indicates that the hybrid sterility found in the F11's is unlikely to be from genetic drift leading to incompatible differences between the genetic lines. Crosses within (n=80) and between (n=52) the three F11 populations (Dune survivors, Headland survivors and Control), again regardless of their gravitropic response, did not have an effect on hybrid sterility (LR chi-square=0.15, P=0.6995), increasing the likelihood that gravitropism, and not another trait driven by selection to one of these environments, caused intrinsic reproductive isolation in this experimental system. Next, we determined whether specific F11 individuals (n=129) were driving this association between gravitropism and hybrid sterility. Thirty-two individuals were a parent in a sterile cross; 26 of these individuals were crossed against a different individual and they successfully produced seeds, indicating that sterility is specific to the parental combination. Finally, we found that crosses were symmetric; only 6 crosses displayed differences in hybrid sterility in the reciprocal cross, indicating the maternal parent did not dictate sterility. Together, these results imply a genetic association between the adaptive divergence of gravitropism and intrinsic reproductive isolation.

Discussion

Whether local adaptation can directly drive the formation of intrinsic reproductive isolation is still a topic of debate[5,8]. Genetic incompatibilities are considered to evolve late in the speciation process, often after extrinsic barriers have formed (e.g. low fitness migrants)[4], and genetic drift or natural selection has led to the accumulation of genetic differences[2]. With few studies[9,10] on the genetics of local adaptation and intrinsic reproductive isolation within a species, the role of natural selection in

creating intrinsic reproductive barriers at the start of the speciation process remains unclear. Here, we

provide empirical evidence to suggest that natural selection is a major driving force behind the

intrinsic reproductive isolation found between recently derived erect and prostrate populations in *S.*

lautus. We identified a trait in natural populations that could only result from divergence in certain

pathways, thus linking the coordinated function of many genes with the evolution of a divergent trait.

Variation in this trait, shoot gravitropism, 1) evolved multiple times between adjacent populations, 2)

predicted the height of adult individuals, 3) responded to natural selection in the direction of the local

population, and 4) was controlled by genomic regions containing genes in the predicted genetic

pathway. Having established gravitropism as a locally adapted trait, we showed that it was genetically

correlated with hybrid sterility. This set of experiments demonstrates an elusive link between the

genetics of local adaptation and the evolution of intrinsic reproductive isolation, which to our

knowledge has only convincingly been shown twice – in *Arabipdopsis*[9] and *Mimulus*[10].

Our results suggest that the evolution of genes related to the auxin pathway could facilitate

colonisation of new habitats through concomitant changes in multiple developmental and architectural

traits. Transitions from erect to prostrate growth and the associated traits of height, main stem angle

and number of branches, are common in plants that colonise coastal headlands[51-54]. This indicates

that there are strong selective agents common to headland environments. For example, powerful

winds or salty substrates could impose early selective pressures on traits controlled by auxin such as

responses to mechanical cues[55] and halotropism[56-58]. As the auxin pathway is common across plant

species, local adaptation to headland environments through auxin-related genes could be leading to

intrinsic reproductive isolation in other plant species. In *Mimulus*, gibberellin (another key plant

hormone), has been recently implicated in local adaptation and also associated with reproductive

isolation between coastal and inland populations[59,60]. Similar to *S. lautus* Headland populations,

coastal *Mimulus* populations have reduced gibberellin, a prostrate growth form and a higher salt

tolerance compared to inland *Mimulus* populations[60,61]. Results from *Mimulus* and our study

presented here suggest that divergence in plant hormonal pathways that affect multiple traits during

plant development could be a powerful mechanism to link the genetic, ecological and physiological

basis of adaptation with speciation in plants.

Previous studies in *S. lautus* support a general link between local adaptation and intrinsic

reproductive isolation. Walter, et al. [50] found that additive variance responsible for local adaptation in

four ecotypes was also responsible for the loss of genetic variation in an F2 generation that displayed

elevated levels of hybrid sterility. This is consistent with the observation that adaptation and intrinsic

reproductive isolation are genetically correlated in some *S. lautus* populations. A similar occurrence of

hybrid sterility occurred in the F2's that created the F11 generation used here[28], suggesting that

genetic incompatibilities are segregating in multiple populations and ecotypes of *S. lautus*. The other

generations of the hybrid population (F3-F11), however, are highly fertile, implying that a large

proportion of the incompatible alleles between Dune and Headland populations were removed in the

F2 generation. Remarkably, we were able to recreate intrinsic reproductive isolation in the F11 hybrid

population when we sort F11 families according to the strength of their gravitropic response; crosses

between individuals with divergent gravitropism values fail to produce seed. A likely explanation is

that incompatibility alleles segregate at low frequencies in the population after the F2 generation, and

these incompatibilities are directly associated with gravitropism. Future research will investigate

whether the same genetic incompatibilities are causing this intrinsic reproductive isolation throughout

the system.

The mechanism by which local adaptation leads to intrinsic reproductive isolation could be pleiotropy

(e.g. Bomblies and Weigel [9]) or genetic linkage (e.g. Wright, et al. [10])[2]. The closer a hybrid sterility

locus is to a target of selection, the more likely reproductive isolation is to concomitantly evolve. We

speculate that divergence in adaptive gravitropism and hybrid sterility could result from pleiotropic

effects in *ENODL* genes. *ENODL1* was not only a strong gravitropism candidate gene in this hybrid

population but members of the ENODL family display functions related to pollen tube reception. For

example, *Arabidopsis* mutants of *ENODL* failed to arrest growth and rupture the entering pollen

tube[49]. This suggests a female-specific role in which *ENODL* is used to communicate with the male

pollen tube to enable reproduction. Consistent with this mechanism, the fitness of the dam family and

not the sire family explained gravitropism in our selection experiment. However, one finding that does

not support a dam-sire interaction for the source of intrinsic reproductive isolation in these hybrids is

the symmetric reciprocal crosses. Dam-sire interactions are more likely to cause differences in hybrid

sterility depending on the maternal parent[62]. Instead, symmetric reciprocal crosses are consistent with

the presence of negative nuclear genetic interactions, which is observed in the genetics of

reproductive isolation in F2 hybrids[28,50]. Future genetic studies will focus on isolating the loci causing

intrinsic reproductive isolation in *S. lautus* so we can reveal the molecular mechanisms by which

intrinsic barriers evolve as a by-product of local adaptation.

Overall, our study illustrates a powerful strategy to explore the genetic basis of complex adaptation

and speciation in natural systems that challenges the common gene-centric paradigm and moves

towards identifying sets of genes related by function. We postulate that the evolution of hormonal

pathways in plants provides a simple mechanism for the rapid formation of integrated traits that

facilitate colonisation of new habitats, and to the incidental evolution of intrinsic reproductive isolation.

Materials and Methods

Reference genome

Headland individuals from Lennox Head were used for the creation of an Illumina *S. lautus* reference

genome. Firstly, we collected seeds from the Headland at Lennox Head (Table S1) and germinated

seeds from two individuals by cutting 1mm off the micropyle side of the seed and placing in petri

dishes with dampened filter paper. The seeds were placed in darkness for two days to allow for roots

to grow and then transferred to a 12-hour light cycle in a constant temperature room at 25°C for seven

days to allow for shoots to grow. Seedlings were then transferred into large individual pots with

standard potting mix and grown in a glasshouse. These two individuals were crossed by rubbing

flower heads together and collecting the seeds produced. Siblings from the seeds produced were

grown and crossed by rubbing flower heads together to produce a family of individuals capable of

selfing (self-fertilisation is rare in *S. lautus*). One generation of selfing was completed to increase

homozygosity. We extracted DNA from leaf tissue of one individual using a modified CTAB protocol[63].

A draft genome of *S. lautus* was de novo assembled using second-generation short reads and

AllPaths-LG 6.1.2 using default settings. We utilised a series of eight read libraries (Table S7). The

reads were trimmed to remove residual adapter sequences and low-quality bases (minimum quality

15). The final assembly was ~843mb long and consisted of 96,372 scaffolds with an N50 of 21K.

Although 843mb is much shorter than the expected haploid size of 1.38GB[64] of the full genome, the

BUSCO gene content completeness of 84% (5% fragmented and 11% missing) suggests that this

assembly is primarily missing intergenic repetitive DNA sequences, which are notoriously difficult to

assemble. This Illumina assembly was used for mapping of all F11 samples.

A PacBio *S. lautus* reference genome was then created in an attempt to overcome the problem

caused by repetitive DNA. The same individual described above was put in darkness for three days

before leaf samples were collected and sent to BGI for DNA extraction and sequencing. The genome

was de novo assembled using next-generation PacBio long reads (56 SMRT cells) and CANU 1.6[65]

with default settings. A total of 11,481,638 filtered subreads (5,072,756 from RSII sequencer and

6,408,882 from Sequel sequencer) were used with a mean read length of ~8K base pairs and an

expected genome coverage of ~60x. The final assembly consisted of 60,947 contigs with an N50 of

34K. The BUSCO gene content completeness was 86% (4% fragmented and 10% missing).

Genetic mapping of convergent traits

Creation of the Bulk Segregant Analysis populations

To identify the genomic regions associated to growth habit in *S. lautus*, we created recombinant

populations by crossing individuals from four pairs of parapatric populations (Fig. S2) that represent

independent origins of the erect and prostrate phenotypes[12,16,31,66]. We crossed individuals from

contrasting populations growing in the following locations: (1) headland (prostrate) and dune (erect)

from Lennox Head (NSW), (2) headland (prostrate) and dune (erect) from Coffs Harbour (NSW), (3)

headland (prostrate) and muttonbird island (erect) from Coffs Harbour, and (4) exposed (prostrate)

and sheltered (erect) alpine sites from Mt Kosciuszko (NSW) and Falls Creek (VIC), respectively.

Using a random-mating-equal-contributions breeding design[67], we crossed multiple individuals from

pairs of adjacent populations to create F1's (2 crosses per plant on average) and conducted

intercrosses (2 crosses per plant on average) for eight generations between Lennox Head

populations (F8-LH), four generations between Alpine populations (F4-A) and three generations

between both Coffs Harbour populations (F3-CHD and F3-CHI) (Fig. S2).

We measured three phenotypic traits that distinguish erect and prostrate architectures and are

possibly regulated by auxin: height, angle of the main stem, and number of tertiary branches. Height

(vegetative) was measured as the vertical distance from the soil to the highest point of the plant that

has vegetative leaves (flowers and stems are not included). The angle of the main stem was

measured with respect to the soil. Higher values of this angle indicate a more erect stem. To evaluate

the number of tertiary branches, we collected three basal secondary branches (i.e. branches that

originate from the main stem) and counted the number of branches arising from these. Trait

distributions and population statistics are provided in Data S3.

Genotyping of the Bulk Segregant Analysis populations

We genotyped the four populations described above by pooling individuals from extremes of the three

traits. More specifically, young leaves were collected, and DNA was extracted following a modified

version of the CTAB protocol[63]. The DNA was quantified with the PicoGreen reagent (Invitrogen,

Carlsbad, CA) and pools of recombinant individuals were created as described in Roda, et al. [31]. For

each recombinant population we pooled DNA from plants from the tails of the distribution (upper and

lower 10%) of the three convergent traits (height, angle of the main stem and number of tertiary

branches). We thus produced 24 pools of DNA (3 traits x 4 populations x 2 tails; Data S4). Pooled

DNA was cleaned using the Wizard PCR-cleanup kit (Promega, Sydney, Australia). Libraries of

Restriction-site Associated DNA (RAD) tags[68] were created and sequenced at Floragenex (University

of Oregon High Throughput Sequencing Facility) following the methodology described in Roda, et al.

[31] but sequencing depth and coverage was increased to 50x depth and 2m reads / sample.

The Bulk Segregant Analysis

We identified genomic regions associated to convergent traits by conducting pairwise comparisons

between contrasting pools from the same trait and population (3 traits x 4 populations = 12

comparisons). To identify Single Nucleotide Polymorphisms (SNPs), determine allelic frequencies and

calculate genetic differentiation between pairs of pools, RADseq data was processed using the

PoPoolation2 pipeline[69]. Illumina reads were mapped to the PacBio genome draft of *S. lautus* using

BWA 0.7.15[70] with default parameters. We then used SAMtools 1.5[71] to create an mpileup file for

each pair of pools (Phred quality score of 20). Finally we used the PoPoolation2 pipeline 1201[69] to

extract variable positions (minimum count of 2, minimum coverage of 15, and maximum coverage of

800), estimate allelic frequency differences[72] between pools and calculate Fisher Exact tests (FET) of

allelic differentiation[73-75]. We thus obtained an average number of SNPs per comparison of 39,161 ±

7,244 SNPs located in 10,826 ± 1,405 genomic contigs. Candidate SNPs for each comparison were

based on two conditions; located in the upper 5 percentiles of the genome-wide distribution of allele

frequency differences and having Bonferroni-corrected p-values lower than 5% in the FET. Genomic

contigs containing at least one candidate SNP were considered candidates (Table S2 and Data S5).

To determine if candidates from the BSA of the three traits show footprints of natural selection we re-

analysed previous pool-seq samples of natural populations[16,31] and survivors from reciprocal

transplants[36] (Table S2). We used the pipeline described above to identify candidate SNPs and

contigs for pairwise comparisons between either adjacent populations[16,31] or survivors from

transplants to different environments[36]. To identify SNPs differentiated in natural populations we

performed 12 pairwise genomic comparisons between DNA pools from adjacent Dune and Headland

populations (Data S4). To identify loci putatively under natural selection in the sand dune and rocky

headland we performed comparisons between pools of survivors from reciprocal transplants using F8

populations (6 comparisons; Data S6). Outlier SNPs and contigs were defined as described

previously for both types of pairwise comparisons.

Permutation analyses

We ran a permutation analysis in R v3.6.0[76] to determine if the number of candidates from BSA

shared between populations and traits is greater than expected by chance. Specifically, we randomly

re-assigned outliers to SNP positions for each of the 12 comparisons (3 traits x 4 populations) and

calculated the number of positions shared between at least two comparisons. We repeated this

process 10,000 times and calculated 99% confidence intervals.

Differential gene expression between erect and prostrate natural populations

Plant material and RNA extraction

To obtain plant tissue to evaluate gene expression, we used mature plants derived from seeds

collected in the field from three erect natural populations: Alpine (A03), Dune (D01), and Tableland

(T01), and from one prostrate natural population, Headland (H01). Field-collected seeds were grown

(see above for details) under natural light conditions and temperature was controlled to 22°C during

511 the night (6:00pm-6:00am) and 25°C during the day (6:00am-6:00pm). Plants were randomly crossed

512 within each natural population, and 96 seeds of each of the four natural populations were germinated

513 as described above and grown in 48-cell trays in a controlled temperature room at 25°C with a 12-

514 hour light cycle. We sampled the shoot apical meristem (SAM) along with the youngest three leaf

515 primordia (LP 1, 2 and 3) when plants were 7 weeks old.

517 *cDNA library construction*

518 For each natural population, SAM-LP from each plant was dissected under a dissecting microscope

519 and conserved in RNAlater solution (Ambion, Inc., Austin, USA). Total RNA was extracted using

520 TRIzol[77]. Total RNA was quantified using a NanoDrop (Thermo Fisher Scientific, Waltham, USA) and

521 quality was checked using a 2100 Bioanalyzer (Agilent technologies, USA) at the Australian Genome

522 Research Facility. Only samples with an RNA integrity number[78] > 6.5 and a ratio of the 28S and 18S

523 ribosomal bands > 1.0 were further processed. 0.5 μg of total RNA were used to construct sequencing

524 libraries with the TruSeq RNA Sample Preparation Kits, and sequenced using 100bp pair-end

525 sequencing with Illumina HiSeq 2000 at BGI.

527 *Gene expression quantification*

528 We used the new tuxedo suite, composed of HISAT, StringTie and Ballgown, to identify differentially

529 expressed genes between the prostrate population, H01, and each of the three erect populations,

530 A03, D01 and T01. We first filtered RNAseq raw reads by removing sequencing adaptors, reads with

531 more than 10% unknown bases and more than 50% bases with quality lower than five. Clean reads of

532 each population were mapped to the PacBio *S. lautus* genome draft using HISAT2 2.0.0[79] to align

533 pair-end reads, and Samtools 1.5[71] to obtain sorted BAM files. Stringtie 1.3.4d[80] was used to 1)

534 assemble mapped reads of each population to transcripts, 2) merge the transcripts of the four

535 populations into a non-redundant set of transcripts contained in the four populations, and 3) quantify

536 transcript abundances in each of the four populations. Finally, we used the Ballgown[81] R package to

537 obtain the FPKM (fragments per kilobase of exon per million reads mapped) by gene. We calculated

538 the relative expression between erect (A03, D01 and T01) and prostrate (H01) samples as

539 Log2(FPKM erect/ FPKM prostrate) (Data S7).

Gene annotation

To identify auxin genes associated to the evolution of erect and prostrate phenotypes, we annotated

candidate loci from the BSA using databases from *Arabidopsis thaliana*, which is the plant model

where most studies of auxin signalling have been conducted, and has the best genome annotation

among plants. We first conducted a BLASTx of the *S. lautus* genes, as predicted from transcriptome

data, against the TAIR protein database (https://www.arabidopsis.org/) using standalone BLAST

2.6.0, and a minimum *e-value* of 1e-6 (Data S8). For genomic contigs showing associations to

adaptive traits in the BSA we also searched for genes not detected in the transcriptome by doing a

BLASTx of the contigs against the TAIR protein database. We then retrieved the function of significant

hits using the DAVID server[82,83] (Data S9). Auxin genes were identified by searching for the term

"auxin" in this annotation. We refined our functional description of these auxin genes by searching for

the terms "biosynthesis", "branch", "efflux", "gravitropism", "influx", "morphogenesis", "polar", "protein

modification", "reproduction", "transcription factor", "transport", and "tropism" (Table S2). We also

searched in the TAIR database for literature references associated to the function of these

Arabidopsis homologs. Given their large number, these references are not provided in the manuscript.

Phenotyping of natural populations

Height measurements

We measured height as described above in all 16 populations in the glasshouse and 12 of the

populations in their native field environment (Data S10 and Data S11). In the field, we measured

height in 32 individuals evenly across the range of each population. In the controlled conditions of the

glasshouse, we sampled an average of 14 individuals per population after plants reached maturity.

For both the glasshouse and field measurements, we used a linear model to determine whether Dune

populations were taller than their adjacent Headland pair: $y_{ijk} = P_i + E_{j(i)} + e_{k(ij)}$, where pair ($P_i$) is an

adjacent Dune and Headland population at the same locality and ecotype ($E_{j(i)}$) is Dune or Headland

and is nested within pair. All factors are fixed effects and $e_{k(ij)}$ is the residual error. Population height

(y_{ijk}) for each pair was compared using a one-tailed t-test. The Alpine populations were also included

with the prediction that the sheltered Alpine population (A03) would be taller than the wind exposed

Alpine population (A07). All statistical results reported here were produced in JMP 13 (SAS 2015).

Gravitropism measurements

Gravitropism was measured *in vitro* on agar plates in a dark growth cabinet using seeds from all 16

natural populations. For each population, 2-4 seeds per family were grown for ~40 maternal families

(1,278 seeds in total). The seeds from a population were combined in a falcon tube and were first

sterilised with a quick rinse in 70% EtOH, followed by four 10 minute inversions in a sterilising solution

of 6% sodium hypochlorite and 1% Tween 20. Seeds were then rinsed three times with distilled water

and vertically orientated on Murashiga and Skoog (MS) agar plates containing 0.15% MS, 0.05% 2-

(*N*-morpholino)ethanesulfonic acid (MES), 0.15% sucrose and 1% agar. We placed eight seeds on

each plate and incubated the plates at 21°C in the dark to avoid any light effects. After seven days, all

plates were rotated clockwise by 90° and a photograph of each plate was taken 24 hours after

rotation. The photographs were imported into ImageJ[84] to determine gravitropism by measuring the

angle to which the stem reorientated to the horizontal[85-87].

Overall there was a 63.8% germination success but we excluded seedlings that were shorter than

5mm, contacted the edge of the plate, or germinated after rotation. This left a total of 736 seedlings

across all 16 populations (57.6% of the total number of seeds planted; Data S10). To test the

hypothesis that Dune populations would have a stronger gravitropic response in their stem than their

adjacent Headland pair, we used a mixed linear model: $y_{ijkl} = P_i + E_{j(i)} + A_k + e_{l(ijk)}$, where pair ($P_i$) is an

adjacent Dune and Headland population at the same locality, ecotype ($E_{j(i)}$) is Dune or Headland and

is nested within pair, and agar plate (A_k) is the MS plate that the seeds were grown on. Agar plate was

fitted as a random effect while the rest were fixed effects, and $e_{l(ijk)}$ is the residual error. Gravitropism

(y_{ijkl}) measures were averaged for each population and compared between each population pair using

a one-tailed t-test. We then tested the correlation between height and gravitropism by performing a

linear regression with mean height against mean gravitropism for all 16 populations, where

populations were grouped into their respective clades (eastern and south-eastern). All statistical

results reported here were produced in JMP 13 (SAS 2015).

Synthetic auxin and auxin transport inhibitor experiments

We tested if gravitropism is governed by auxins in *S. lautus* by evaluating gravitropic responses in

seedlings treated with chemicals affecting auxin signalling. Specifically we used 2,4-

Dichlorophenoxyacetic acid (2,4-D), a carrier-dependent synthetic auxin[41,88] and Naphthylphthalamic

acid (NPA) an efflux inhibitor[42]. We collected seeds from Lennox Head sand dune and rocky headland

and grew them on MS plates containing the different chemicals at concentrations of 0 μM, 0.5 μM, 5

μM and 50μM. For this we used the following stock solutions: 2,4-D: 1mM in Ethanol and NPA: 10mM

in dimethyl sulfoxide (DMSO). We created 1ml dilutions of stock solutions (in ethanol or DMSO),

which were dissolved in 500 ml of media. Plates were incubated, rotated, photographed and

gravitropism was measured in ImageJ[84], as outlined above.

Approximately 40 seeds from different maternal families were planted for each population and

chemical concentration. Seeds were excluded from analyses following the same protocols as above,

leaving a total of 188 seedlings to quantify the gravitropic response. The 50μM concentration of 2,4-D

treatment was excluded, as less than six seeds germinated under this high concentration (Data S12).

For each chemical we used a mixed linear model using the lmer function of lme4 package in R

v3.1.3[76]: $y_{ijkl} = A_k + E_i + C_j + E_i \times C_j + e_{l(ijk)}$, where agar plate ($A_k$) is the MS plate that the seeds were

grown on, ecotype (E_i) is Dune or Headland and concentration (C_j) are the 4 different concentrations

of the chemical. Agar plate was fitted as a random effect, while ecotype, concentration and their

interaction ($E_i \times C_j$) were fixed effects, and $e_{l(ijk)}$ is the residual error. Gravitropism (y_{ijkl}) was compared

using a Type II Wald chi-square test.

Field experiments

All field experiments were conducted at the sand dune and rocky headland at Lennox Head (NSW) in

the same location where native *S. lautus* grow. We tracked each individual by gluing each seed to a

toothpick with nondrip superglue and placing them 1-2mm under the ground within a grid cell (Fig. 3b)

that was randomly assigned (for details see Walter, et al. [28,]Walter [89]). 50% shade cloth was

suspended 15cm above all plots to replicate the shade given by surrounding vegetation and was later

replaced with very light bird netting. Seeds were watered twice a day to keep the soil moist and

replicate ideal germination conditions to maximise the number of seeds in the experiment. Once

germination plateaued for all genotypes watering was gradually ceased.

Height adaptation experiments

We created genetic lines that aimed to isolate height differences on a common genomic background

(Fig. 1). Firstly, Lennox Head Dune and Headland seeds were grown and crossed to create an F1

generation. Secondly, we backcrossed to Headland parental plants for two generations to produce a

BC2F1 generation. We then grew and crossed the tallest BC2F1 individuals (n=16, tallest 10% of the

population) among one another, and the shortest individuals (n=18, shortest 10% of the population)

among one another. See Table S3 for the number of individuals and families that contributed to every

generation to create these BC2F2 genetic lines. The BC2F2 seeds were planted into the rocky

headland at Lennox Head in October 2016 (Australian spring). We planted five replicate plots, where

each plot (1.08x0.33m) consisted of the same 12 families with four (occasionally three) individuals per

family in each plot, totalling 1,116 BC2F2 seeds. Germination and mortality were recorded every day

for 49 days, then every 3-4 days until day 79 and then weekly for the remainder of the experiment,

until day 159 (Data S13).

We implemented a mixed linear model to test the hypothesis that individuals with short parents will

have higher fitness in the headland environment: $y_{ijkl} = H_i + F_j + B_k + e_{l(ijk)}$, where parental height (H_i)

was the average height of the parents measured in the glasshouse, family (F_j) was individuals with the

same parents, and block (B_k) is the five replicate plots across the rocky headland. Parental height is a

fixed effect, and family and block are random effects, and $e_{l(ijk)}$ was the residual error. Offspring fitness

(y_{ijk}) was the total number of days alive in the rocky headland from planting. All statistical results

reported here were produced in JMP 13 (SAS 2015).

Gravitropism adaptation experiments

We created an advanced recombinant population (F8) from 23 Dune and 22 Headland Lennox Head

individuals using a North Carolina 2 breeding design[67,90] as described in Roda, et al. [36]. Briefly, we

replicated the construction of the F8 using three independent replicate crossing lines (A, B and C), all

derived from the same base population. See Table S4 for the number of families per replicate genetic

line for every generation. F8 seeds were planted into the sand dune and rocky headlands in 2012 as

described above. The fittest families (top 50%) within an environment (and genetic line) were selected

using Aster modelling[91,92] implemented with the 'Aster' package in R[76]. The fitness components

included germination and survival success, where germination was the total number of individuals that

germinated in each family and survival success was the total number of individuals per family that

survived to day 85 in the F8 generation. One full-sibling from the 101 selected families (50 Dune

survivor and 51 Headland survivor families) and 44 families randomly selected from all F8 families for

the control population were then grown in the glasshouse. Each individual was randomly assigned as

a dam or sire and after reaching maturity crossed twice in a full-sibling, half-sibling crossing design.

This maintained ~100 full-sibling families for each population (Dune survivors, Headland survivors and

Control). The same field experiment and selection procedure was conducted on the F9 (in 2013) and

F10 generations (in 2014), for a total of three rounds of selection, where survival success was

measured at day 138 in the F9 generation, and day 232 in the F10 generation. See Table S5 for the

number of seeds and families planted per genetic line and environment for the three transplant

experiments.

F11 gravitropism measurements

We measured gravitropism, in the F11 generation described above, as the angle of the stem after a

90° rotation of a seedling. This included 39 Dune families, 37 Headland families and 25 inbred control

families with 12 individuals per family (1,212 seeds in total). These families were germinated in three

separate batches ~7 days apart. Briefly, we germinated the F11 seeds by cutting 1mm off the

micropyle side of the seed and placing in petri dishes with dampened filter paper. The seeds were

placed in darkness for two days to enable roots to grow and then transferred to light for 4 days in

order for shoots to grow. Seedlings were then transferred into small square pots with standard potting

mix in a constant temperature room at 25°C with a 12-hour light cycle. After one week of growing in

the pot, the plants were rotated by 90° and a photograph of each individual was taken 12 hours after

rotation. The photographs were imported into ImageJ[84] to measure gravitropism as described above.

Data S14 contains gravitropism measures and dam and sire fitness values.

Gravitropism tests of selection

We implemented a linear model to test the hypothesis that high fitness dune families will produce

gravitropic plants and high fitness headland families will produce agravitropic plants. Independent

models were used for the sand dune and rocky headland to test the effect of gravitropism on fitness in

each environment: $y_{ijklmn} = B_i + V_j + G_{k(i)} + D_{l(ik)} + S_{m(ik)} + e_{n(ijklm)}$, where temporal block ($B_i$) is the three

time points in which the F11 seeds were grown (~seven days apart); intrinsic viability (V_j) is the

number of days until the death of F11 plants in controlled conditions; and genetic line, which consists

of the three independent genetic lines (A, B and C), is nested within temporal block ($G_{k(i)}$). Dam fitness

was nested in genetic line and temporal block ($D_{l(ik)}$) and sire fitness was also nested in genetic line

and temporal block ($S_{m(ik)}$). Dam and sire fitness are the F10 family fitness values for the individuals

that were crossed to create the F11 offspring where gravitropism was measured. All factors were

included as fixed effects and $e_{n(ijklm)}$ was the residual error. Genetic line C was removed from analyses

as it has little variation in fitness values which means it did not converge. Shapiro-Wilk W test shows

the residuals from the model are normally distributed for both the sand dune (W=0.98, P=0.3879) and

rocky headland (W=0.98, P=0.2776). The linear model was performed in JMP 13 (SAS 2015).

Genetic association between height and gravitropism

We tested the genetic association between height and gravitropism after segregation in an advanced

recombinant population. We implemented a mixed linear model for the three F11 populations (Dune

survivors, Headland survivors and a Control population) that accounts for family variation: $y_{ijk} = H_i + F_j$

$+ e_{k(ij)}$, where gravitropism (y_{ijk}) is the angle of the growth response 12 hours after a 90° rotation,

height (H_i) is the vertical distance from the soil to the top of the vegetative leaves, measured after

maturity in the glasshouse and family (F_j) is a random effect that consists of individuals that have the

same dam and sire. The mixed linear model was performed in JMP 13 (SAS 2015).

Genotyping of F11 gravitropism tails

To isolate gravitropism candidate genes, we genotyped 77 gravitropic (>56°) and 68 agravitropic

(<20°) F11 individuals (Data S14). We extracted DNA from leaf tissue using a modified CTAB

protocol[63] and quantified the DNA using the PicoGreen reagent (Invitrogen, Carlsbad, CA). To

determine which parent the alleles were derived from, we included 39 Dune parentals (D01) and 41

Headland parentals (H01). Leaves from the Lennox Head Dune and Headland natural populations

were collected directly from the field and the same DNA extraction protocol was followed. To increase

accuracy of genotyping, each F11 individual was duplicated in independent wells and libraries of

Restriction-site Associated DNA (RAD) tags were created at Floragenex following the protocol from

Baird, et al. [68] but using the PstI restriction enzyme. We sequenced 380 samples on four lanes of an

Illumina HiSeq 4000 with 91bp single end reads at Floragenex. A total of 1.39 billion reads with a

mean of 3.62 million reads per sample were produced. Reads were aligned to the reference genome

using Bowtie 1.1.1[93] and a FASTQ quality score of above 20. We then used SAMtools 0.1.16[71] to

create an mpileup file of all samples with a minimum Phred quality score of 10, minimum sequencing

depth per sample of 6x and minimum percent of population genotyped of 75%. This produced 26.8K

variable positions (224K variants before filtering) with a minimum distance between variants of 50bp.

The gravitropism candidate gene set consisted of SNPs in the 99.9% quantile of the distribution of

differentiated SNPs between the gravitropism tails. The region of the scaffold containing the SNP was

annotated using the BLASTx NCBI database[94].

Senecio lautus gravitropism candidate genes

In the gravitropism candidate gene set, we tested for overrepresentation of gene function and linkage

disequilibrium between gravitropism loci. A statistical overrepresentation test was performed in

PANTHER (http://pantherdb.org/) using the TAIR identification for 32 unique gravitropism candidate

genes matched to a reference list of 27,502 *Arabidopsis thaliana* genes. To calculate linkage

disequilibrium between loci, a likelihood-ratio chi-square test was performed in JMP 13 (SAS 2015)

with each F11 population independently (Dune survivors, Headland survivors and Control); Data

S15).

Intrinsic reproductive isolation

We tested whether hybrid fertility was associated with gravitropism by randomly crossing within and

between the F11 gravitropic and agravitropic groups in a controlled temperature room and recording

successful and failed crosses. To maximise sample size, all three replicate genetic lines (A, B and C)

were used across all three environments (Dune survivors, Headland survivors and Control). A total of

138 crosses were completed (Data S16), 71 crosses between the tails and 67 within the tails

(agravitropic tail = 29 and gravitropic tail = 38). Crosses were completed by rubbing multiple flower

heads of two individuals together and collecting the seeds produced from both plants. To remove

genetic incompatibilities that might be caused by relatedness, crosses within the same family were

not performed. Hybrid sterility (a failed cross) was considered when, in the presence of pollen, less

than three seeds were produced per flower head from both plants (reciprocal crosses) with three

mating attempts. Six crosses displayed differences in sterility in the reciprocal cross (asymmetry), with

one successful and one failed cross when using the same parents, and so were removed from the

analysis (4 crosses between the tails and 2 within the tails).

If gravitropism was contributing to divergence in *S. lautus* then we expected to observe an increase in

hybrid sterility in crosses between gravitropic and agravitropic individuals. We performed a linear

model in JMP 14 (SAS. 2018) to determine whether there was a significant association between cross

type (within vs between gravitropism tails) and hybrid sterility, while accounting for genetic line and

F11 population: $y_{ijkl} = T_i + G_j + E_k + e_{l(ijk)}$, where tails ($T_i$) is either crosses within or between

gravitropism tails, genetic line (G_j) are the three independent genetic lines (A, B and C) and

population (P_k) are the three F11 populations (Dune survivors, Headland survivors and Control). All

factors were included as fixed effects and $e_{l(ijk)}$ was the residual error. Hybrid fertility (y_{ijkl}) was

compared using a likelihood-ratio chi-square test.

Acknowledgments

We are grateful to S. Smith, L.H. Rieseberg, M. Cooper, J. Engelstaedter, M.A.F. Noor and members

of the Ortiz-Barrientos laboratory for insightful comments on previous versions of this manuscript. S.

Karrenberg and S. Chenoweth provided very useful feedback on M.J. Wilkinson PhD dissertation. We

thank P. Brewer for his help in the design and execution of gravitropism experiments, T. Richards for

his contributions to the construction of recombinant populations for the BSA, and J. Donohoe for his

help in the phenotyping and rearing of plants for the BSA. We would also like to thank everyone that

helped with the field selection experiments: E. Johnston, E. Monley, G. Wilkinson, P. Wilkinson, A.

Mather, S. Thang, K. Giarola, G. Lebbink, E. Firkins-Barriere, P. Arnold, J. Donohoe, B. Brittain, H.

Liu, D. Bernal, M.C. Melo, T. Richards, J. Walter, L. Ambrose, B. Ayalon, S. Carrol, M. Gallo, A.

Maynard, C. Palmer and S. Edgley.

Author Contributions

MJW, FR and DO conceived the projects and experiments. MJW conducted height adaptation

experiments, reared and phenotyped various glasshouse populations and performed reproductive

isolation experiment. FR created the BSA mapping populations and performed the BSA analyses. FR

and MJW performed physiological and molecular biology experiments. GMW reared and phenotyped

the natural populations in the glasshouse with help from MJW and MEJ. GMW conducted the field

experiment for the gravitropism adaptation experiments with assistance from MJW, MEJ, FR and DO.

DMB performed expression analyses. MJW, MEJ and HLN phenotyped natural populations. MEJ and

MJW extracted the DNA from the F11's and RN and JW called the genotypes. MEJ reared the plants

and prepared the Illumina libraries for SA to assemble the *S. lautus* genomes. CB guided the

physiological experiments. MJW, FR, DO wrote the paper with input from all authors. DO is mentor

and supervisor for the research program.

References

Darwin, C. *On the origins of species by means of natural selection*. Vol. 247 (1859).

Coyne, J. A. & Orr, H. A. *Speciation*. (Sinauer Associates, 2004).

Rundle, H. D. & Whitlock, M. C. A genetic interpretation of ecologically dependent isolation. *Evolution* **55**, 198-201 (2001).

Seehausen, O. *et al.* Genomics and the origin of species. *Nat. Rev. Genet.* **15**, 176-192, doi:10.1038/nrg3644 (2014).

Baack, E., Melo, M. C., Rieseberg, L. H. & Ortiz-Barrientos, D. The origins of reproductive isolation in plants. *New Phytol.* **207**, 968-984, doi:10.1111/nph.13424 (2015).

Schluter, D. & Conte, G. L. Genetics and ecological speciation. *Proc. Natl. Acad. Sci. USA* **106**, 9955 (2009).

Lowry, D. B., Rockwood, R. C. & Willis, J. H. Ecological reproductive isolation of coast and inland races of *Mimulus guttatus*. *Evolution* **62**, 2196-2214, doi:10.1111/j.1558-5646.2008.00457.x (2008).

Fishman, L. & Sweigart, A. L. When two rights make a wrong: The evolutionary genetics of plant hybrid incompatibilities. *Annu. Rev. Plant Biol.*, doi:10.1146/annurev-arplant-042817-040113 (2018).

Bomblies, K. & Weigel, D. Hybrid necrosis: autoimmunity as a potential gene-flow barrier in plant species. *Nat. Rev. Genet.* **8**, 382, doi:10.1038/nrg2082 (2007).

Wright, K. M., Lloyd, D., Lowry, D. B., Macnair, M. R. & Willis, J. H. Indirect evolution of hybrid lethality due to linkage with selected locus in *Mimulus guttatus*. *PLoS Biol.* **11**, e1001497, doi:10.1371/journal.pbio.1001497 (2013).

Kirkpatrick, M. & Barton, N. Chromosome inversions, local adaptation and speciation. *Genetics* **173**, 419 (2006).

Radford, I. J., Cousens, R. D. & Michael, P. W. Morphological and genetic variation in the *Senecio pinnatifolius* complex: are variants worthy of taxonomic recognition? *Aust. Syst. Bot.* **17**, 29-48, doi:10.1071/sb03013 (2004).

Thompson, I. *Taxonomic studies of Australian Senecio (Asteraceae): 5. The S. lautus/S. lautus complex.*, 23-76 (Royal Botanic Gardens, 2005).

Stinchcombe, J. R. & Hoekstra, H. E. Combining population genomics and quantitative genetics: finding the genes underlying ecologically important traits. *Heredity* **100**, 158-170 (2007).

Barrett, R. D. H. & Schluter, D. Adaptation from standing genetic variation. *Trends Ecol. Evol.* **23**, 38-44, doi:10.1016/j.tree.2007.09.008 (2008).

Roda, F. *et al.* Convergence and divergence during the adaptation to similar environments by an Australian groundsel. *Evolution* **67**, 2515-2529, doi:10.1111/evo.12136 (2013).

James, M. E., Wilkinson, M. J., North, H. L., Engelstädter, J. & Ortiz-Barrientos, D. A framework to quantify phenotypic and genotypic parallel evolution. *bioRxiv*, doi:10.1101/2020.02.05.936450 (2020).

Friml, J. Auxin transport—shaping the plant. *Curr. Opin. Plant Biol.* **6**, 7-12 (2003).

Xu, M., Zhu, L., Shou, H. & Wu, P. A PIN1 family gene, OsPIN1, involved in auxin-dependent adventitious root emergence and tillering in rice. *Plant Cell Physiol.* **46**, 1674-1681, doi:10.1093/pcp/pci183 (2005).

Wallace, J. G. *et al.* Genome-wide association for plant height and flowering time across 15 tropical maize populations under managed drought stress and well-watered conditions in Sub-Saharan Africa. *Crop Sci.* **56**, 2365-2378, doi:10.2135/cropsci2015.10.0632 (2016).

Gallavotti, A. The role of auxin in shaping shoot architecture. *J. Exp. Bot.* **64**, 2593-2608, doi:10.1093/jxb/ert141 (2013).

Domagalska, M. a. & Leyser, O. Signal integration in the control of shoot branching. *Nat. Rev. Mol. Cell Biol.* **12**, 211-221, doi:10.1038/nrm3088 (2011).

840   23   Sundberg, E. & Østergaard, L. Distinct and dynamic auxin activities during reproductive
841   development. *Cold Spring Harb. Perspect. Biol.* **1**, a001628-a001628,
842   doi:10.1101/cshperspect.a001628 (2009).

843   24   Vanneste, S. & Friml, J. Auxin: a trigger for change in plant development. *Cell* **136**, 1005-
844   1016, doi:10.1016/j.cell.2009.03.001 (2009).

845   25   Melo, M. C., Grealy, A., Brittain, B., Walter, G. M. & Ortiz-Barrientos, D. Strong extrinsic
846   reproductive isolation between parapatric populations of an Australian groundsel. *New*
847   *Phytol.* **203**, 323-334, doi:10.1111/nph.12779 (2014).

848   26   Richards, T. J. & Ortiz-Barrientos, D. Immigrant inviability produces a strong barrier to gene
849   flow between parapatric ecotypes of *Senecio lautus*. *Evolution* **70**, 1239-1248,
850   doi:10.1111/evo.12936 (2016).

851   27   Richards, T. J., Walter, G. M., McGuigan, K. & Ortiz-Barrientos, D. Divergent natural selection
852   drives the evolution of reproductive isolation in an Australian wildflower. *Evolution* **70**, 1993-
853   2003, doi:10.1111/evo.12994 (2016).

854   28   Walter, G. M. *et al.* Diversification across a heterogeneous landscape. *Evolution* **70**, 1979-
855   1992, doi:10.1111/evo.13009 (2016).

856   29   Walter, G. M., Wilkinson, M. J., Aguirre, J. D., Blows, M. W. & Ortiz-Barrientos, D.
857   Environmentally induced development costs underlie fitness tradeoffs. *Ecology* **99**, 1391-
858   1401, doi:doi:10.1002/ecy.2234 (2018).

859   30   James, M. E., Arenas-Castro, H., Groh, J. S., Engelstädter, J. & Ortiz-Barrientos, D. Highly
860   replicated evolution of parapatric ecotypes. *bioRxiv*, 2020.2002.2005.936401,
861   doi:10.1101/2020.02.05.936401 (2020).

862   31   Roda, F. *et al.* Genomic evidence for the parallel evolution of coastal forms in the *Senecio*
863   *lautus* complex. *Mol. Ecol.* **22**, 2941-2952, doi:10.1111/mec.12311 (2013).

864   32   Fishman, L. & Willis, J. H. Evidence for Dobzhansky-Muller incompatibilites contributing to
865   the sterility of hybrids between *Mimulus guttatus* and *M. nasutus*. *Evolution* **55**, 1932-1942
866   (2001).

867   33   Stelkens, R. B., Schmid, C. & Seehausen, O. Hybrid breakdown in cichlid fish. *PLOS One* **10**,
868   e0127207, doi:10.1371/journal.pone.0127207 (2015).

869   34   Michelmore, R. W., Paran, I. & Kesseli, R. V. Identification of markers linked to disease-
870   resistance genes by bulked segregant analysis: a rapid method to detect markers in specific
871   genomic regions by using segregating populations. *Proc. Natl. Acad. Sci. USA* **88**, 9828-9832,
872   doi:10.1073/pnas.88.21.9828 (1991).

873   35   Petrasek, J. & Friml, J. Auxin transport routes in plant development. *Development* **136**, 2675-
874   2688, doi:10.1242/dev.030353 (2009).

875   36   Roda, F., Walter, G. M., Nipper, R. & Ortiz-Barrientos, D. Genomic clustering of adaptive loci
876   during parallel evolution of an Australian wildflower. *Mol. Ecol.* **26**, 3687–3699,
877   doi:10.1111/mec.14150 (2017).

878   37   Tan, L. *et al.* Control of a key transition from prostrate to erect growth in rice domestication.
879   *Nat. Genet.* **40**, 1360-1364, doi:10.1038/ng.197 (2008).

880   38   Li, P. *et al.* LAZY1 controls rice shoot gravitropism through regulating polar auxin transport.
881   *Cell Res.* **17**, 402-410, doi:10.1038/cr.2007.38 (2007).

882   39   Lopez, D., Tocquard, K., Venisse, J.-S., Legué, V. & Roeckel-Drevet, P. Gravity sensing, a
883   largely misunderstood trigger of plant orientated growth. *Front. Plant Sci.* **5**,
884   doi:10.3389/fpls.2014.00610 (2014).

885   40   Blancaflor, E. B. Regulation of plant gravity sensing and signaling by the actin cytoskeleton.
886   *Am. J. Bot.* **100**, 143-152, doi:10.3732/ajb.1200283 (2013).

887   41   Yamamoto, M. & Yamamoto, K. T. Differential effects of 1-naphthaleneacetic acid, indole-3-
888   acetic acid and 2,4-dichlorophenoxyacetic acid on the gravitropic response of roots in an
889   auxin-resistant mutant of *Arabidopsis*, *auxl*. *Plant Cell Physiol.* **39**, 660-664 (1998).

42   Ottenschläger, I. *et al.* Gravity-regulated differential auxin transport from columella to
lateral root cap cells. *Proc. Natl. Acad. Sci. USA* **100**, 2987-2991,
doi:10.1073/pnas.0437936100 (2003).
43   Ranocha, P. *et al.* Walls are thin 1 (WAT1), an *Arabidopsis* homolog of *Medicago truncatula*
NODULIN21, is a tonoplast-localized protein required for secondary wall formation in fibers.
The Plant Journal **63**, 469-483, doi:10.1111/j.1365-313X.2010.04256.x (2010).
44   Promchuea, S., Zhu, Y., Chen, Z., Zhang, J. & Gong, Z. ARF2 coordinates with PLETHORAs and
PINs to orchestrate ABA-mediated root meristem activity in *Arabidopsis*. *J. Integr. Plant Biol.*
59, 30-43, doi:10.1111/jipb.12506 (2017).
45   Wu, H., Shen, Y., Hu, Y., Tan, S. & Lin, Z. A phytocyanin-related early nodulin-like gene,
BcBCP1, cloned from *Boea crassifolia* enhances osmotic tolerance in transgenic tobacco. *J.*
Plant Physiol. **168**, 935-943, doi:10.1016/j.jplph.2010.09.019 (2011).
46   Ma, H., Zhao, H., Liu, Z. & Zhao, J. The phytocyanin gene family in rice (Oryza sativa L.):
Genome-wide identification, classification and transcriptional analysis. *PLOS One* **6**, e25184,
doi:10.1371/journal.pone.0025184 (2011).
47   Garcia de la Garma, J. *et al.* New insights into plant salt acclimation: the roles of vesicle
trafficking and reactive oxygen species signalling in mitochondria and the endomembrane
system. *New Phytol.* **205**, 216-239, doi:10.1111/nph.12997 (2015).
48   Mittler, R. & Blumwald, E. The roles of ROS and ABA in systemic acquired acclimation. *The*
Plant Cell **27**, 64-70 (2015).
49   Hou, Y. *et al.* Maternal ENODLs are required for pollen tube reception in *Arabidopsis*. *Curr.*
Biol. **26**, 2343-2350, doi:10.1016/j.cub.2016.06.053 (2016).
50   Walter, G. M. *et al.* Loss of ecologically important genetic variation in late generation hybrids
reveals links between adaptation and speciation. *Evolution Letters* **4**, 302-316,
doi:10.1002/evl3.187 (2020).
51   Beeftink, W., Rozema, J. & Huiskes, A. *Ecology of coastal vegetation*. Vol. 6 (Springer, 1985).
52   Auld, T. & Morrison, D. Genetic determination of erect and prostrate growth habit in five
shrubs from windswept headlands in the Sydney region. *Aust. J. Bot.* **40**, 1-11 (1992).
53   Morrison, D. & Rupp, A. Patterns of morphological variation within *Acacia suaveolens*
(Mimosaceae). *Aust. Syst. Bot.*, 1013-1027 (1995).
54   Crutsinger, G. M., Strauss, S. Y. & Rudgers, J. a. Genetic variation within a dominant shrub
species determines plant species colonization in a coastal dune ecosystem. *Ecology* **91**,
1237-1243 (2010).
55   Li, T. *et al.* Calcium signals are necessary to establish auxin transporter polarity in a plant
stem cell niche. *Nature Communications* **10**, 726, doi:10.1038/s41467-019-08575-6 (2019).
56   Naser, V. & Shani, E. Auxin response under osmotic stress. *Plant Mol. Biol.* **91**, 661-672,
doi:10.1007/s11103-016-0476-5 (2016).
57   van den Berg, T., Korver, R. A., Testerink, C. & ten Tusscher, K. H. Modeling halotropism: a
key role for root tip architecture and reflux loop remodeling in redistributing auxin.
Development **143**, 3350-3362 (2016).
58   Galvan-Ampudia, C. S. *et al.* Halotropism is a response of plant roots to avoid a saline
environment. *Curr. Biol.* **23**, 2044-2050 (2013).
59   Lowry, D. B. & Willis, J. H. A widespread chromosomal inversion polymorphism contributes
to a major life-history transition, local adaptation, and reproductive isolation. *PLoS Biol.* **8**,
e1000500, doi:10.1371/journal.pbio.1000500 (2010).
60   Lowry, D. B., Popovic, D., Brennan, D. J. & Holeski, L. M. Mechanisms of a locally adaptive
shift in allocation among growth, reproduction, and herbivore resistance in *Mimulus*
guttatus*. *Evolution* **73**, 1168-1181, doi:10.1111/evo.13699 (2019).
61   Lowry, D. B., Hall, M. C., Salt, D. E. & Willis, J. H. Genetic and physiological basis of adaptive
salt tolerance divergence between coastal and inland *Mimulus guttatus*. *New Phytol.* **183**,
776-788, doi:10.1111/j.1469-8137.2009.02901.x (2009).

62   Turelli, M. & Moyle, L. C. Asymmetric Postmating Isolation: Darwin's Corollary to Haldane's
Rule. *Genetics* **176**, 1059, doi:10.1534/genetics.106.065979 (2007).
63   Clarke, J. D. Cetyltrimethyl ammonium bromide (CTAB) DNA miniprep for plant DNA
isolation. *Cold Spring Harb. Protoc.* **2009**, pdb.prot5177, doi:10.1101/pdb.prot5177 (2009).
64   Liu, H. *Developing genomic resources for an emerging ecological model species Senecio
lautus* PhD Thesis thesis, The University of Queensland, (2015).
65   Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer weighting
and repeat separation. *Genome Res.* **27**, 722-736, doi:10.1101/gr.215087.116 (2017).
66   Ali, S. *Senecio lautus* complex in Australia. I. Taxonomic considerations and discussion of
some of the related taxa from New Zealand. *Aust. J. Bot.* **12**, 282-291,
doi:10.1071/BT9640282 (1964).
67   Rockman, M. V. & Kruglyak, L. Breeding designs for recombinant inbred advanced intercross
lines. *Genetics* **179**, 1069-1078, doi:10.1534/genetics.107.083873 (2008).
68   Baird, N. A. *et al.* Rapid SNP discovery and genetic mapping using sequenced RAD markers.
PLOS One **3**, 1-7, doi:10.1371/journal.pone.0003376 (2008).
69   Kofler, R., Pandey, R. V. & Schlotterer, C. PoPoolation2: identifying differentiation between
populations using sequencing of pooled DNA samples (Pool-Seq). *Bioinformatics* **27**, 3435-
3436, doi:10.1093/bioinformatics/btr589 (2011).
70   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics **25**, 1754-1760 (2009).
71   Li, H. *et al.* The sequence alignment/map format and SAMtools. *Bioinformatics* **25**, 2078-
2079 (2009).
72   Rosenzweig, B. K., Pease, J. B., Besansky, N. J. & Hahn, M. W. Powerful methods for
detecting introgressed regions from population genomic data. *Mol. Ecol.* **25**, 2387-2397
(2016).
73   Benfey, P. N. & Mitchell-Olds, T. From Genotype to Phenotype: Systems Biology Meets
Natural Variation. *Science* **320**, 495-497, doi:10.1126/science.1153716 (2008).
74   Futschik, A. & Schlötterer, C. The next generation of molecular markers from massively
parallel sequencing of pooled DNA samples. *Genetics* **186**, 207-218,
doi:10.1534/genetics.110.114397 (2010).
75   Magwene, P. M., Willis, J. H. & Kelly, J. K. The statistics of bulk segregant analysis using next
generation sequencing. *PLoS Comp. Biol.* **7**, e1002255, doi:10.1371/journal.pcbi.1002255
(2011).
76   R Core Team. R: A language and environment for statistical computing. *R Foundation for
Statistical Computing, Vienna, Austria* (2013).
77   Simms, D., Cizdziel, P. E. & Chomczynski, P. J. F. TRIzol: A new reagent for optimal single-step
isolation of RNA. **15**, 532-535 (1993).
78   Schroeder, A. *et al.* The RIN: an RNA integrity number for assigning integrity values to RNA
measurements. *BMC Mol. Biol.* **7**, 3 (2006).
79   Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory
requirements. *Nat. Methods* **12**, 357-360, doi:10.1038/nmeth.3317 (2015).
80   Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from RNA-seq
reads. *Nat. Biotechnol.* **33**, 290-295, doi:10.1038/nbt.3122 (2015).
81   Pertea, M., Kim, D., Pertea, G. M., Leek, J. T. & Salzberg, S. L. Transcript-level expression
analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nat. Protoc* **11**, 1650-
1667, doi:10.1038/nprot.2016.095 (2016).
82   Huang da, W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of large
gene lists using DAVID bioinformatics resources. *Nat. Protoc* **4**, 44-57,
doi:10.1038/nprot.2008.211 (2009).

83   Huang da, W., Sherman, B. T. & Lempicki, R. A. Bioinformatics enrichment tools: paths
toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* **37**, 1-13,
doi:10.1093/nar/gkn923 (2009).
84   Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image
analysis. *Nat. Methods* **9**, 671-675 (2012).
85   Sang, D. *et al.* Strigolactones regulate rice tiller angle by attenuating shoot gravitropism
through inhibiting auxin biosynthesis. *Proc. Natl. Acad. Sci. USA* **111**, 11199-11204,
doi:10.1073/pnas.1411859111 (2014).
86   Rigo, G. *et al.* Inactivation of plasma membrane-localized CDPK-RELATED KINASE5
decelerates PIN2 exocytosis and root gravitropic response in *Arabidopsis*. *The Plant Cell* **25**,
1592-1608, doi:10.1105/tpc.113.110452 (2013).
87   Rashotte, A. M., Brady, S. R., Reed, R. C., Ante, S. J. & Muday, G. K. Basipetal auxin transport
is required for gravitropism in roots of *Arabidopsis*. *Plant Physiol.* **122**, 481-490,
doi:10.1104/pp.122.2.481 (2000).
88   Yang, Y., Hammes, U. Z., Taylor, C. G., Schachtman, D. P. & Nielsen, E. High-affinity auxin
transport by the AUX1 influx carrier protein. *Curr. Biol.* **16**, 1123-1127,
doi:10.1016/j.cub.2006.04.029 (2006).
89   Walter, G. M. *The genetic and ecological basis of diversification* PhD Thesis thesis, The
University of Queensland, (2016).
90   Lynch, M. & Walsh, B. *Genetics and analysis of quantitative traits*. (Sinaeur Associates Inc,
1998).
91   Geyer, C. J., Wagenius, S. & Shaw, R. G. Aster models for life history analysis. *Biometrika* **94**,
415-426 (2007).
92   Shaw, R. G., Geyer, C. J., Wagenius, S., Hangelbroek, H. H. & Etterson, J. R. Unifying
life-history analyses for inference of fitness and population growth. *The American Naturalist*
172, E35-E47, doi:10.1086/588063 (2008).
93   Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biology* **10**, R25,
doi:10.1186/gb-2009-10-3-r25 (2009).
94   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search
tool. *J. Mol. Biol.* **215**, 403-410, doi:10.1016/s0022-2836(05)80360-2 (1990).
1021

 Figures and Tables

Fig. 1. Sample locations and growth habit differences between adjacent *Senecio lautus* populations. a, Map of Australia showing locations of the 16 populations used in this study. The seven coastal localities have a Dune (orange) and Headland (green) population occurring adjacent to each other. The populations are split into two monophyletic clades[31], the eastern clade (closed circles) and the south-eastern clade (open circles). **b,** *Senecio lautus* native to the sand dunes have an erect growth habit and *S. lautus* native to the rocky headlands have a prostrate growth habit. **c,** Alpine populations of *S. lautus* include a sheltered alpine gully and a wind exposed alpine meadow, containing individuals with an erect and prostrate growth habit, respectively.

1032

Fig. 2. Gravitropism and height variation across 16 *Senecio lautus* populations. a, The

correlation between gravitropism and height across *S. lautus* populations split into their monophyletic

clades – see Fig. 1 for details. Each point in the graph represents a population mean where height

was measured in the glasshouse and gravitropism was measured 24 hours after a 90° rotation. **b,c**,

Divergence in gravitropism, height in the glasshouse and height in the field between adjacent *S.*

lautus populations (D = Dune, H = Headland, A-S=Alpine Sheltered and A-E = Alpine Exposed). **b**,

eastern clade and **c**, south-eastern clade. Height in the field for Falls Creek, Kosciusko NP and Point

Labatt were not measured. Data are mean ± s.e.m.; one tailed Student's t-test (Table S8), *P ≤ 0.05,

**P ≤ 0.01, ***P ≤ 0.001, NS not significant.

1042

Fig. 3. Parental survival curves in the height (2016) and gravitropism (2012-2014) adaptation experiments at the sand dune and rocky headland at Lennox Head. a,**b**, Survival (Estimated S(t)) over the length of the field experiments is shown for the Lennox Head Dune population (orange) and the Lennox Head Headland population (green) for four independent field selection experiments at the **a**, sand dune and **b**, rocky headland. Asterisks indicate a significant difference in mortality risk between the Dune and Headland ecotypes (**P ≤ 0.01, ***P ≤ 0.001).

Fig. 4. The creation of the recombinant hybrid generation, the design of the gravitropism adaptation experiments and sequencing of the tails of the gravitropic distribution. a, 23 parental Dune and 22 parental Headland individuals from Lennox Head were crossed randomly and with equal contribution for eight generations. **b**, Seeds from this F8 recombinant population were glued to toothpicks and transplanted into the sand dune and rocky headland at Lennox Head. Among-family based selection occurred for three generations (F8, F9 and F10), where full-siblings from the fittest families were grown in the glasshouse and crossed amongst their respective genetic lines (A, B and C) and their environment (Dune survivors or Headland survivors). An inbred control was kept in the glasshouse and underwent the same crossing scheme but free from viability selection. **c**, Gravitropism was measured in the F11 recombinant population by re-orienting the plant by 90°. Here, agravitropic plants are define as individuals with gravitropic angles <20°, while gravitropic plants have gravitropic angles >56° as they reorient their growth and subsequently grow upright. Individuals in the tails of the gravitropism distribution were sequenced on four lanes of the Illumina HiSeq 4000 platform.

1064

Fig. 5. Association between *ENODL1* and *ABA3* alleles and gravitropism in *Senecio lautus*. The

average gravitropism angle is shown for each allelic combination, independently and when the

agravitropic alleles (dark grey) are combined. Different letters denote significant differences between

genotypes (t>1.98, P<0.05).

Table 1. General linear model for the effect of dam and sire on gravitropism (°) after a field

selection experiment on a recombinant hybrid Dune and Headland population. Field

selection experiments were performed on F8, F9 and F10 recombinant hybrid generations to

achieve three rounds of selection in the sand dune and rocky headland at Lennox Head (see Fig.

4 for the experimental design). Dam and sire fitness are the F10 family fitness values for the

individuals that were crossed to create the F11 offspring where gravitropism was measured.

Intrinsic viability is the number of days until death of the F11 generation in the controlled

temperature room. This experiment was conducted three times (temporal block) with three

independent genetic lines.

Source	Dune				Headland			
	DF	SS	F-Ratio	P-value	DF	SS	F-Ratio	P-value
Dam family fitness	6	8515.77	4.779	**<0.001**	6	1884.31	0.701	0.650
Sire family fitness	6	1806.62	1.014	0.424	5	2315.36	1.033	0.405
Intrinsic viability	1	260.85	0.878	0.352	1	5209.38	11.624	**0.001**
Genetic lines	3	1135.49	1.275	0.290	3	3357.19	2.497	0.067
Temporal block	2	193.14	0.325	0.724	2	2234.96	2.494	0.090

1078   DF, degrees of freedom; SS, sum of squares.