Divergence in hormone signaling drives local adaptation and intrinsic reproductive isolation

Melanie J. Wilkinson[1†], Federico Roda[1,2†], Greg M. Walter[1,3], Maddie E. James[1], Rick Nipper[4], Jessica Walsh[4], Scott L. Allen[1,5], Diana M. Bernal[1], Henry L. North[1,6], Christine A. Beveridge[1] and Daniel Ortiz-Barrientos[1*].

[1]The University of Queensland, School of Biological Sciences, St Lucia QLD 4072, Australia. [2]Current address: Universidad Nacional de Colombia, Departamento de Biología, Bogotá, Colombia. [3]Current address: University of Bristol, School of Biological Sciences, Senate House, Tyndall Ave, Bristol BS8 1TH, UK. [4]Floragenex, Inc., 4640 SW Macadam Avenue, Suite 200F, Portland, OR 97239, USA. [5]Current address: Institut für Populationsgenetik, Vetmeduni Vienna, Veterinärplatz 1, 1210, Wien, Austria. [6]Current address: University of Cambridge, Department of Zoology, Downing St., Cambridge CB2 3EJ, UK.

† These authors contributed equally to the paper.

* Author for correspondence: Daniel Ortiz-Barrientos

Email: d.ortizbarrientos@uq.edu.au

Abstract

Natural selection is a major driver of both adaptation and speciation. Whether or not the processes driving adaptation and speciation share a molecular basis remains largely unknown. Here, we show that divergence in hormone signaling contributed to the evolution of complex adaptations and intrinsic reproductive isolation in an Australian wildflower, *Senecio lautus*. We provide evidence that differences in the auxin pathway, a hormone signaling pathway required for plant growth and development, has led to the repeated evolution of erect and prostrate forms to contrasting coastal environments. Using multiple hybrid and natural populations, we show that adjacent erect and prostrate populations have repeatedly diverged in auxin-related genes and auxin-dependent phenotypes, such as gravitropism. Analysis of an advanced hybrid population in a multi-year field selection experiment revealed that high fitness families produce offspring with the local gravitropic response. We found that two auxin-related genes explained 45% of the variation in gravitropism in hybrid individuals. Remarkably, crossing hybrid individuals with extreme differences in gravitropism significantly reduces their ability to produce seed, showing gravitropism is genetically correlated with intrinsic reproductive isolation. Together, our results suggest that divergence in hormonal pathways can drive rapid adaptation to contrasting environments and to the origin of new species.

Introduction

Ever since Darwin's work on the origin of new species by natural selection (1), a fundamental

question in evolutionary biology remains unanswered: do adaptation and speciation share a common

genetic basis? Although several genes that impact speciation have been identified (2), we still remain

ignorant as to whether genes driving speciation have evolved as a by-product of local adaptation. One

notable exception exists in *Mimulus* populations adapted to copper mines where genetically linked loci

for local adaptation and hybrid mortality were found (3). Here, we introduce a novel system and a

powerful way to uncover the molecular link between local adaptation and intrinsic reproductive

isolation by studying these processes in populations that have repeatedly and independently adapted

to contrasting environments. These highly replicated systems imply a role for natural selection in the

evolution of complex morphological or physiological differences (4, 5), thus providing a unique

opportunity to examine if new species originate via the incidental evolution of intrinsic barriers to

reproduction between locally adapted populations.

We examine the genetics of repeated evolution of growth habit and its contribution to the formation of

intrinsic reproductive barriers between erect and prostrate populations of an Australian wildflower,

Senecio lautus (G.Forst. ex Willd) (6, 7). A previous population genetic study in *S. lautus* found that

sets of genes related by similar functions were repeatedly differentiated in populations with these

contrasting growth habits (8). One of these sets contained genes belonging to the auxin pathway,

where genes involved in the polar transport of this key plant hormone had repeatedly diverged

between erect and prostrate populations. Transport and regulation of auxin is required to create an

auxin-concentration gradient that alters the directional growth of a plant (9) and contributes to

variation in height (10, 11), branching (12, 13) and reproduction (14). These traits distinguish the erect

and prostrate growth habits of *S. lautus* populations, making the auxin pathway a natural candidate to

link the molecular basis of both adaptation and speciation. We reasoned that if divergence in the

auxin pathway contributed to the evolution of adaptation and speciation in *S. lautus*, we would make

the following three observations. First, we would detect similar patterns of genetic divergence in

auxin-related pathways across multiple erect and prostrate populations. Second, these populations

would differ in auxin-dependent phenotypes, such as their ability to alter the direction of growth in

relation to gravity (9, 15). Finally, divergence in these auxin-dependent phenotypes would contribute

to local adaptation and intrinsic reproductive isolation between populations.

We test these hypotheses primarily on coastal populations of *S. lautus* (Fig. 1a, Table S1), which

exhibit strong correlations between growth habit and the environments they occupy (6). Populations

inhabiting sand dunes (Dune hereafter) are erect, while populations growing on adjacent rocky

headlands (Headland hereafter) are prostrate (Fig. 1b). Erect and prostrate growth habits can also be

found in related populations from the alpine regions of Australia, with a prostrate population inhabiting

an exposed alpine meadow and an erect population inhabiting a sheltered alpine gully (Fig. 1c). Dune

populations are continually exposed to high temperatures and sun radiation, low salinity, and low

nutrient sand substrate, whereas Headland populations are exposed to high salinity, high nutrients

and powerful winds (16). Neighboring Dune and Headland populations are often sister taxa, group

into two major monophyletic clades (eastern and south-eastern) and have evolved their contrasting

growth habits independently multiple times (6, 17). These Dune and Headland population pairs are

locally adapted (16-19) and their F2 hybrids have low fitness (20), indicating the presence of intrinsic

reproductive isolation. Furthermore, performing genetic, physiological, and ecological experimental

studies is achievable in this system due to its short life cycle and small vegetative size. Therefore, the

evolution of growth habit in *S. lautus* provides an excellent system in which to understand the shared

molecular basis of adaptation and intrinsic reproductive isolation.

Results

Repeated evolution of auxin-related genes in *S. lautus*

To investigate whether there are similar patterns of genetic divergence in auxin-related pathways

across erect and prostrate populations, we assessed the genes that govern variation in three traits

that distinguish growth habit in *S. lautus* (plant vegetative height, main stem angle and the number of

tertiary branches; Fig. S1). We performed a Bulk Segregant Analysis (BSA) (21) in four independent

mapping populations that represent replicates for the evolution of the erect and prostrate

morphologies in *S. lautus* (22) (Fig. S2). For each mapping population, we pooled individuals from the

tails of each of the three trait distributions and created reduced representation libraries for

Genotyping-by-Sequencing. In agreement with a previous study in *S. lautus* (8), we found that highly

differentiated genomic regions (candidates) between extreme forms were often not the same across

mapping populations (Data S1). Only 20 candidate regions were shared across all four mapping

populations: 14 candidates (out of 2,648) for main stem angle, six candidates for branch number (out

of 1,782) and zero candidates for height (out of 1,198), suggesting repeated evolution at the gene

level is limited. However, we found that 11 out of these 20 candidate regions contained genes

regulating the auxin pathway (Table S2), implying evolution might have repeatedly occurred in

hormone signaling pathways. Four of these 11 candidate regions contained multiple auxin-related

genes, which could indicate the existence of pleiotropy or clusters of genes controlling variation in

multiple adaptive traits. Auxin candidate genes from the BSA also showed footprints of natural

selection such as genetic differentiation between adjacent populations, and significant associations

with survivorship in reciprocal transplants (Table S2 and Data S1). This suggests that the auxin

pathway might be important for adaptation to sand dunes and rocky headlands. Several of these

auxin genes also displayed differential expression between shoot apical meristems of erect and

prostrate populations (Fig. S3), providing further evidence for their involvement in mediating plant

architecture differences. The homologs of BSA auxin candidate genes include genes largely involved

in intracellular auxin transport (Table S2), an essential process for creating auxin concentration

gradients that drive a directional growth response to environmental cues such as light (phototropism)

or gravity (gravitropism) (15, 23). Some homologs also have pleiotropic effects on reproduction (Table

S2), which is consistent with a previous mapping experiment where quantitative trait loci were the

same for plant architecture and hybrid fecundity between erect and prostrate populations of *S. lautus*

(24). This indicates that a genetic link between complex adaptations and reproductive barriers

between genetically and morphologically divergent populations might exist in *S. lautus*.

The physiological basis of repeated evolution in *S. lautus*

Since we identified a multitude of auxin related genes associated with erect and prostrate

phenotypes*,* and that the regulation and transport of auxin is a well-established modulator of

gravitropism in plants, we predicted that the evolution of divergent growth habits in many plants may

be a direct consequence of changes in the auxin pathway (25, 26). If this is true in *S. lautus* then plant

height should predict gravitropism across populations, and differences in gravitropism would have

occurred repeatedly between locally adapted populations with contrasting morphologies. Additionally,

synthetic auxin and auxin transport inhibitors should influence gravitropism differently between these

erect and prostrate populations.

To test whether plant height predicts gravitropism across populations, we first quantified plant height

and gravitropism in 16 *S. lautus* populations (Fig. 1 and Fig. S4). We defined the gravitropic response

as the change in angle of a seedlings stem 24 hours after a 90° rotation (27, 28), where 0° is no

gravitropic response and 90° is a complete re-orientation in relation to the new gravity vector. If

differences in the auxin pathway have contributed to divergence in growth habit, we expect that

prostrate populations will display a smaller gravitropic angle than their adjacent erect population. First,

we show that height has a genetic basis in *S. lautus* by confirming that plant height in natural

environments (field) and plant height in common garden conditions (glasshouse) are strongly

correlated ($F_{1,9}=12.41$, p=0.0065). Next, we found that the average height of a population predicted its

average gravitropic angle (Fig. 2a; $F_{1,13}=17.65$, p=0.0010), where the average magnitude of

gravitropism differed between the two monophyletic clades ($F_{1,13}=32.58$, p<0.0001). These results

suggest that changes in gravitropism are genetically correlated with divergent growth habit traits such

as plant height, and that this relationship has evolved at least twice in the system.

To test whether differences in gravitropism have occurred repeatedly between locally adapted

populations with differences in plant height, we compared gravitropism between seven adjacent Dune

and Headland population pairs (Table S1). Four population pairs showed the expected correlation,

where plants from the Headland population exhibited a smaller gravitropic angle and were shorter

than plants from their adjacent Dune population (Fig. 2b, c). In two population pairs – Millicent and

Point Labatt – plant height did not differ between Dune and Headland population pairs (Fig. 2c),

possibly due to their similarity in environmental variables (22). In the population pair at Stradbroke

Island, however, we observed differences in height but not in gravitropism (Fig. 2b, c), suggesting that

similar plant architectures could have evolved through other developmental routes in coastal

populations of *S. lautus*. We also assessed height and gravitropism in divergent populations from the

alpine region of Australia, and discovered the expected pattern, where the exposed Alpine population

in the meadow was shorter and had a smaller gravitropic angle than the population in the sheltered

Alpine gully (Fig. 2c). These results suggest that natural selection repeatedly acting on auxin-

dependent genes contribute to differences in gravitropism across coastal and alpine populations in

this system.

To directly test whether synthetic auxin and auxin transport inhibitors will influence gravitropism

differently between erect and prostrate populations, we grew Lennox Head Dune and Headland seeds

with synthetic auxin 2,4-D (2,4-Dichlorophenoxyacetic acid), and polar auxin transport inhibitor NPA

(naphthylphthalamic acid) (29, 30). Because a gravitropic response requires an auxin concentration

gradient, we reasoned that removing the gradient would reduce the gravitropic angle in Dune

individuals (Fig. S5). As expected, addition of synthetic auxin 2,4-D reduced the gravitropic angle

more in Dune individuals (from 66.34±2.27° to 19.44±7.64°) than in Headland individuals (LR chi-

square χ^2_2=18.49, p<0.0001). Similarly, addition of auxin transport inhibitor NPA reduced the

gravitropic angle in our experiments more in Dune individuals (from 62.57±3.62° to 20.47±13.67°)

than in Headland individuals (LR chi-square, χ^2_3=21.18, p<0.0001). This difference in hormone

response in Dune and Headland individuals supports our hypothesis that differences in the auxin

pathway underlie the different gravitropic responses in *S. lautus*.

Together, these three results reveal that genetic differences in the auxin pathway have contributed to

the repeated evolution of growth habit in this system. Additionally, our findings are consistent with the

hypothesis that natural selection is the major evolutionary force responsible for the physiological and

morphological diversity underpinning the differences in growth habit between Dune and Headland

populations.

Adaptive evolution of auxin-dependent phenotypes in *S. lautus*

Having established that auxin-related genes have diverged between erect and prostrate forms in *S.

lautus,* we conducted two independent sets of field adaptation experiments to experimentally assess

the role of natural selection on the evolution of auxin-dependent phenotypes, height and gravitropism.

In each field experiment, Dune and Headland parental and hybrid seeds were transplanted into

replicated blocks at the sand dune and rocky headland at Lennox Head. In all field experiments,

natural selection consistently favored the local Dune or Headland population over the foreign

population (Fig. 3), indicating that experimental conditions captured local adaptation in both environments.

For the height adaptation experiments, we tested whether differences in height could drive differences in fitness in the rocky headlands. We hypothesized that offspring produced by short hybrid parents would live longer than offspring produced by tall hybrid parents in the rocky headland. We focused on prostrate growth as it is likely the derived trait because the majority of populations in the *S. lautus* species complex have an erect growth habit (8, 16). We introgressed Dune alleles associated with height onto a Headland genomic background. Briefly, we crossed Lennox Head Dune and Headland parentals and then completed two rounds of backcrossing followed by one round of crossing between tall hybrid offspring and between short hybrid offspring (Fig. S6). We transplanted 558 of these seeds (from 28 families) into the rocky headland. As predicted, shorter hybrid parents produced offspring that lived longer in the rocky headland relative to offspring from taller hybrid parents ($F_{1,26.23}=4.87$, p=0.0362). These results suggest that traits genetically correlated with plant height contributed to early developmental fitness in the rocky headlands and contributed directly or indirectly to the evolution of divergent growth habit.

For the gravitropism adaptation experiments, we tested whether rapid adaptation to contrasting environments can lead to the evolution of gravitropic responses in the direction of the local population, and to the formation of a genetic correlation with plant height. We hypothesized that exposing an advanced recombinant population to multiple rounds of viability selection in the field would drive genetic covariation between fitness and gravitropism. To test this hypothesis, we crossed the Dune and Headland populations from Lennox Head for eight generations using recombination to disassemble allelic combinations and construct a recombinant hybrid population (24) (Fig. 4a). We planted 2,403 of these F8 seeds (from 89 families) into the sand dune and rocky headland (Fig. 4b) and conducted family-based truncation selection for three generations, planting a similar number of seeds and families in the F9 and F10 generation (Table S3). We selected the top 50% of surviving families in each environment, and subsequently crossed their full-siblings in the glasshouse to produce the next generation. In the F10 generation, we tested whether families that had the greatest number of survivors also produced offspring with the local gravitropic response under laboratory

conditions. In agreement with our hypothesis, F10 families with the greatest number of survivors in

the sand dune produced offspring with a higher gravitropic angle (Table 1). We discovered that the

relationship between fitness of the F10 families in the sand dune and the gravitropic response in the

F11 generation in the glasshouse was dependent on fitness of the F10 dam family (Table 1),

suggesting maternal genotypes contribute to the evolution of gravitropism in the sand dunes. On the

other hand, all F10 families in the rocky headland produced offspring with intermediate levels of

gravitropism (Table 1). Notably, in the F11 recombinant Control population that was grown and

crossed in the glasshouse only, height and gravitropism were not genetically correlated ($F_{1,114.3}=0.08$,

p=0.7801, r^2=0.04), but this genetic correlation was strong after three rounds of selection in the rocky

headland ($F_{1,169.5}=7.09$, p=0.0085, r^2=0.27) and weak after selection in the sand dune ($F_{1,151.3}=3.20$,

p=0.0756, r^2=0.09). Together, these results suggest that natural selection can act on standing genetic

variation and reconstitute genetic architectures that favor a correlation between survivorship and

auxin-dependent phenotypes in the environments where the coastal populations are found.

Our results imply that natural selection of hormonal pathways can aid in the rapid evolution of multiple

integrated phenotypes when encountering a new selection pressure.

The genes underlying gravitropism in *S. lautus*

Having shown that natural selection drove the rapid evolution of gravitropism in *S. lautus* under

experimental conditions in the field, we identified genes underlying adaptation by detecting extreme

allelic differences between the F11 individuals with the smallest gravitropic angle (agravitropic tail,

<20°; mean of tail = 6.46±1.10°, n=68) and the largest gravitropic angle (gravitropic tail, >56°; mean of

tail = 62.03±0.45°, n=77). We performed selective genotyping on the tails of the gravitropic distribution

in the F11 recombinant populations (Fig. 4c; Dune and Headland survivors, and Control individuals)

and created a gravitropism candidate gene set with allelic differences in the 99.8% quantile

(0.15<*p*<0.22). We identified 55 sites (0.2% of all SNPs) across 49 genomic scaffolds in this quantile

(Data S2) and discovered that these genes disproportionally represented gene ontology categories of

transport and localization of molecules across and between cells (Table S4).

Five of the 55 sites (11%) are located in gene homologs with functions related to the auxin pathway,

including the second (*ENODL1*; early nodulin-like protein 1) and fourth (*ABA3*; molybdenum cofactor

sulfurase) most differentiated SNPs between the agravitropic and gravitropic F11 tails. In both the

ENODL1 and *ABA3* gene, the allele frequency of the F11 tails shifted in the direction of the most

common allele in the native population with the same phenotype (Data S2). For instance, the alleles

favored in the agravitropic F11 tail were at high frequencies in the Headland population (*ENODL1*

C=0.87 and *ABA3* G=0.89), with the alternate alleles being favored in the gravitropic F11 tail and the

Dune population (*ENODL1* A=0.69 and *ABA3* A=0.97). *ENODL1* and *ABA3* were in strong linkage

disequilibrium in the survivors of the gravitropism adaptation experiment in the rocky headland

(Fisher's exact test, n=57, p=0.0008) and sand dune (Fisher's exact test, n=48, p=0.0107), but not in

the control population reared in the glasshouse (Fisher's exact test, n=37, p=0.4093), suggesting

natural selection drove the evolution of this allelic combination. *ENODL1* C/C and *ABA3* G/G

genotypes were associated with a reduction in gravitropism of 25.10° relative to all other genotype

combinations (Fig. 5; $t_{34.30}=4.86$, p<0.0001; 45% of the difference in gravitropism between these F11

tails). Together, these results suggest that a small gravitropic angle is a recessive phenotype, and

that natural selection has targeted few genetic regions of large effect. *ENODL1* and *ABA3* have

functions in gravitropism, plant height, salt tolerance and pollen tube growth across different plant

species (31-37), and therefore are strong candidates to not only explain the adaptive evolution of

gravitropism, but other phenotypic differences between Dune and Headland populations. Moreover,

ENODL1 was also associated with field survivorship in a previous BSA study in this system (24) and

with height in one of the BSA populations we used here for genetic mapping of convergent traits.

Overall, our results suggest that divergence in auxin-regulated molecular processes contributed to

local adaptation to contrasting environments in coastal populations of *S. lautus*.

The consequences of gravitropism divergence on the evolution of intrinsic reproductive

isolation

We are now in a unique position to address whether adaptation and speciation have a common

molecular basis. Specifically, we can test whether divergent gravitropic forms are reproductively

isolated. To assess this problem, we investigated the effects of gravitropism on reproductive success

in the F11 hybrids. We created two groups of crosses among F11 families: crosses within

(agravitropic × agravitropic, n=28; gravitropic × gravitropic, n=37) and crosses between (agravitropic ×

gravitropic, n=67) the tails of the gravitropic distribution. We found that crosses failed more often

between F11 agravitropic and gravitropic plants (21%) than crosses within each of these tails (5%;

Fisher's exact test, n=132, p=0.0079). Of the 11 failed crosses with known genotypes, only one cross

had the *ENODL1* C/C genotype, none had the *ABA3* G/G genotype, and none were double

homozygous for the two genes, suggesting that evolution of a small gravitropic angle possibly relied

on the recessive × recessive interaction. These results imply a genetic association between

adaptation and intrinsic reproductive isolation among populations that occupy contrasting

environments.

Discussion

Here we have discovered that divergence in auxin-related genes contribute to the repeated evolution

of growth habit in *S. lautus*, suggesting that evolution in hormone signaling could facilitate colonization

of new habitats through concomitant changes in multiple developmental and architectural traits.

Consistent with our results, transitions from erect to prostrate growth and the associated phenotypes

of height, main stem angle and number of branches, are common in plants that colonize coastal

headlands (38-41). This indicates that there are strong selective agents common to headland

environments. For example, powerful winds or salty substrates could impose early selective pressure

in traits controlled by auxins such as responses to mechanical cues (42) and halotropism (43-45).

Further, we discovered that divergence in growth habit was genetically correlated with intrinsic

reproductive isolation in *S. lautus*, implying that hormone signaling divergence might lead to

pleiotropic or indirect evolution of genetic incompatibilities that reduce hybrid fitness or pollen-pistil

interactions (e.g., differences in pollen tube growth (46)) that prevent fertilization. Together, these

results suggest that divergence in plant hormonal pathways that affect multiple traits during plant

development and reproduction could be a powerful mechanism to link the genetic, ecological and

physiological basis of adaptation and speciation in plants.

The roles of hormone signaling in plant adaptation and intrinsic reproductive isolation have been

explored independently but only in rare cases have the two been linked. The hormonal pathways

most commonly involved in plant adaptation are the abscisic acid pathway, which is involved in stress

response (47), the jasmonate signaling pathway that contributes to the growth-defense conflict (48),

and the auxin pathway that responds to salt tolerance (43-45) and mechanical stress responses (42).

On the other hand, hormonal pathways involved in intrinsic reproductive isolation include the cytokinin

signaling pathway in hybrid sterility (49), and the auxin pathway in pollen tube growth (46). In more

recent work in *Arabidopsis* (50) and *Mimulus* (51), variation in genes involved in the gibberellin

pathway were associated with variation in both morphological differentiation and local adaptation to

different environments. The consequences of such adaptation for the evolution of intrinsic

reproductive isolation remain unclear, but at least in *Mimulus*, many of the gibberellin genes

associated with local adaptation reside within a chromosomal inversion strongly associated with

flowering time differences between coastal and inland populations (51, 52). Our study takes us a step

further by directly linking natural selection and divergence in hormone signaling with intrinsic

reproductive isolation in a highly replicated system of evolution, thus paving the way to uncover the

molecular basis of repeated evolution and its consequences on the evolution of intrinsic reproductive

isolation.

Overall, our study illustrates a powerful strategy to explore the genetic basis of adaptation in natural

systems that takes us away from a gene-centric paradigm and moves us towards identifying sets of

genes related by function. We conjecture that the evolution of hormonal pathways in plants provides a

simple mechanism for the rapid formation of integrated phenotypes that facilitate colonization of new

habitats, and to the incidental evolution of intrinsic reproductive isolation as a consequence.

Materials and Methods

Reference genome

Two Lennox Head Headland individuals were used for the creation of the Illumina *S. lautus* reference

genome. Firstly, we collected seeds from the Headland at Lennox Head (Table S1) and germinated

seeds from two individuals by cutting 1mm off the micropyle side of the seed and placing in petri

dishes with dampened filter paper. The seeds were placed in darkness for two days for roots to grow

and then transferred to 12 hours light/day in a constant temperature room at 25°C for seven days for

shoots to grow. Seedlings were then transferred into pots with standard potting mix and grown in the

glasshouse. Siblings from the seeds produced were grown and crossed by rubbing flower heads

together to produce a family of individuals capable of selfing (rare in *S. lautus*). One generation of

selfing was completed to increase homozygosity. We extracted DNA from leaf tissue using the CTAB

protocol (53) with column cleaning and overnight incubation. All sequencing was performed by Beijing

Genomics Institute (BGI).

A draft genome of *S. lautus* was de novo assembled using second-generation short reads and

AllPaths-LG 6.1.2 by utilizing a series of eight read libraries (Table S2). The reads were trimmed to

remove residual adapter sequence and low quality bases (minimum quality 15). The final assembly

was ~843MB long and consisted of 96,372 scaffolds with an N50 of 21K. Although 843MB is much

shorter than the expected haploid size of 1.38GB (54) of the full genome, the BUSCO gene content

completeness of 84% (5% fragmented and 11% missing) suggests that this assembly is primarily

missing intergenic repetitive DNA sequences, which are notoriously difficult to assemble. This Illumina

assembly was used for mapping of all F11 samples.

A Pacbio *S. lautus* reference genome was then created in an attempt to overcome the problem

caused by repetitive DNA. The genome was de novo assembled using next-generation Pacbio long

reads and CANU 1.6 (55). A total of 11,481,638 reads (5,072,756 from RSII sequencer and 6,408,882

from Sequel sequencer) were used with a mean read length of ~8K base pairs and an expected

genome coverage of ~60x. The final assembly consisted of 60, 947 contigs with an N50 of 34K. All

sequencing was performed by BGI.

Genetic mapping of convergent traits

Creation of BSA populations

To identify the genomic regions associated to growth habit in *S. lautus*, we created recombinant

populations by crossing individuals from four pairs of parapatric populations (Fig. S2) that represent

independent origins of the erect and prostrate phenotypes (6, 8, 22, 56). We crossed individuals from

contrasting populations growing in the following locations: (1) headland (prostrate) and dune (erect)

from Lennox Head (NSW), (2) headland (prostrate) and dune (erect) from Coffs Harbour (NSW), (3)

headland (prostrate) and muttonbird island (erect) from Coffs Harbour, and (4) exposed (prostrate)

and sheltered (erect) alpine sites from Mt Kosciuszko (NSW) and Falls Creek (VIC), respectively.

Using a random-mating-equal-contributions breeding design (57), we crossed multiple individuals

from pairs of adjacent populations to create F1s (2 crosses per plant on average) and conducted

intercrosses (2 crosses per plant on average) for eight generations between Lennox Head

populations (F8-LH), four generations between Alpine populations (F4-A) and three generations

between both Coffs Harbour populations (F3-CHD and F3-CHI) (Fig. S2).

We measured three phenotypic traits that distinguish erect and prostrate architectures and are

possibly regulated by auxins: height, angle of the main stem, and number of tertiary branches. Height

(vegetative) was measured as the vertical distance from the soil to the highest point of the plant that

has vegetative leaves (flowers and stems are not included). The angle of the main stem was

measured with respect to the soil. Higher values of this angle indicate a more erect stem. To evaluate

the number of tertiary branches we collected three basal secondary branches (i.e. branches that

originate from the main stem) and counted the number of branches rising from these. Trait

distributions and population statistics are provided in Data S3.

Genotyping of BSA populations

Genomic DNA of these pools described above was extracted following a modified version of the

CTAB protocol (53). Young leaves were collected and stored at -80°C and ground using a Tissuelyser

(Qiagen, Chatsworth, CA). DNA was quantified with the Picogreen reagent (Invitrogen, Carlsbad, CA)

and pools of recombinant individuals were created as described in ref. (22). For each recombinant

population we pooled DNA from plants from the 10% tails of the distribution of the three convergent

traits (height, angle of the main stem and number of tertiary branches). We thus produced 24 pools of

DNA (3 traits x 4 populations x 2 tails; Data S4). Pooled DNA was cleaned using the Wizard PCR-

cleanup kit (Promega, Sydney, Australia). Libraries of Restriction-site Associated DNA (RAD) tags

(58) were created and sequenced at Floragenex (University of Oregon High Throughput Sequencing

Facility) following the methodology described in ref. (22) but increasing sequencing depth and

coverage (50x depth and 2M reads / sample).

Bulk Segregant Analysis

We identified genomic regions associated to convergent traits by conducting pairwise comparisons

between contrasting pools from the same trait and population (3 traits x 4 populations = 12

comparisons). To identify SNPs, determine allelic frequencies and calculate genetic differentiation

between pairs of pools, RADseq data was processed using the PoPoolation2 pipeline (59). Illumina

reads were mapped to the PacBio genome draft of *S. lautus* using BWA 0.7.15 (60) with default

parameters. We then used SAMtools 1.5 (61) to create an mpileup file for each pair of pools (Phred

quality score of 20). Finally we used the Popoolation2 pipeline 1201 (59) to extract variable positions

(minimum count of 2, minimum coverage of 15, and maximum coverage of 800), estimate allelic

frequency differences (62) between pools and calculate Fisher Exact tests (FET) of allelic

differentiation (63-65). We thus obtained an average number of SNPs per comparison of 39,161 ±

7,244 SNPs located in 10,826 ± 1,405 genomic scaffolds. We called candidate SNPs for each

comparison based on two conditions; located in the upper 5 percentiles of the genome-wide

distribution of allele frequency differences and having Bonferroni-corrected p-values lower than 5% in

the FET. Genomic scaffolds containing at least one candidate SNP were considered candidates for

the comparison (Table S1 and Data S5).

To determine if candidates from BSA of the three traits show footprints of natural selection we re-

analyzed previous pool-seq samples of natural populations (8, 22) and survivors from reciprocal

transplants(24) (Table S3). We used the pipeline described above to identify candidate SNPs and

contigs for pairwise comparisons between adjacent populations (8, 22) or survivors from transplants

to different environments(24). To identify SNPs differentiated in natural populations we performed 12

pairwise genomic comparisons between DNA pools from adjacent Dune and Headland populations

(Data S4). To identify loci putatively under natural selection in the sand dune and rocky headland we

performed comparisons between pools of survivors from reciprocal transplants using F8 populations

(6 comparisons; Data S6). Outlier SNPs and scaffolds were defined as described previously for both

types of pairwise comparisons.

Differential gene expression between erect and prostrate natural populations

Plant material and RNA extraction

To obtain plant tissue to evaluate gene expression, we used mature plants derived from seeds collected in the field from three erect natural populations: Alpine (A03), Dune (D01), and Tableland (T01), and from one prostrate natural population, Headland (H01). Field-collected seeds were grown under natural light conditions and temperature was controlled to 22°C during the night (6:00pm-6:00am) and 25°C during the day (6:00am-6:00pm). Plants were randomly crossed within each natural population, and 96 seeds of each of the four natural populations were germinated as described above and grown in 48-cell flats in a controlled temperature room at 25°C with a 12-hour light/day cycle. We sampled the shoot apical meristem (SAM) along with the youngest 3 leaf primordia (LP 1, 2 and 3) when plants were 7 weeks old.

cDNA library construction

For each natural population, SAM-LP from each plant was dissected under a dissecting microscope and conserved in RNAlater solution (Ambion, Inc., Austin, USA). Total RNA was extracted using TRIzol (66). Total RNA was quantified using a NanoDrop (Thermo Fisher Scientific, Waltham, USA) and quality was checked using a bioanalyzer (Agilent technologies, USA) at the Australian Genome Research Facility. Only samples with RNA integrity number (67) > 6.5 and a ratio of the 28S and 18S ribosomal bands > 1.0 were further processed. 0.5 µg of total RNA were used to construct sequencing libraries with the TruSeq RNA Sample Preparation Kits, and sequenced using 100bp pair-end sequencing with Illumina HiSeqTM 2000 at BGI.

Gene expression quantification

We used the new tuxedo suite, composed of HISAT, StringTie and Ballgown, to identify differentially expressed genes between the prostrate population, H01, and each of the three erect populations, A03, D01 and T01. We first filtered RNAseq raw reads by removing sequencing adaptors, reads with more than 10% unknown bases and more than 50% bases with quality lower than five. Clean reads of each population were mapped to the PacBio *S. lautus* genome draft using HISAT2 2.0.0 (68) to align pair-end reads and Samtools 1.5 (61) to obtain a sorted BAM files. Stringtie 1.3.4d (69) was used to

assemble mapped reads of each population to transcripts, to merge the transcripts of the four

populations into a non-redundant set of transcripts contained in the four populations, and to quantify

transcript abundances in each of the four populations. Finally, we used the Ballgown (70) R package

to obtain the FPKM (fragments per kilobase of exon per million reads mapped) by gene. We

calculated the relative expression between erect (A03, D01 and T01) and prostrate (H01) samples as

Log2(FPKM erect/ FPKM prostrate) (Data S7).

Gene annotation

To identify auxin genes associated to the evolution of erect and prostrate phenotypes, we annotated

candidate loci from BSA using databases from *Arabidopsis thaliana*, which is the plant model where

most studies of auxin signaling have been conducted and has the best genome annotation among

plants. We first conducted a BLASTx of the *S. lautus* genes, as predicted from transcriptome data,

against the TAIR protein database (https://www.arabidopsis.org/) using standalone BLAST 2.6.0, and

a minimum *e-value* of 1e-6 (Data S8). For genomic scaffolds showing associations to adaptive traits

in BSA we also searched for genes not detected in the transcriptome by doing a BLATx of the

scaffolds against TAIR protein database. We then retrieved the function of significant hits using the

DAVID server (71, 72) (Data S9). Auxin genes were identified by searching for the term "auxin" in this

annotation. We refined our functional description of these auxin genes by searching for the terms

"biosynthesis", "branch", "efflux", "gravitropism", "influx", "morphogenesis", "polar", "protein

modification", "reproduction", "transcription factor", "transport", and "tropism" (Table S2). We also

searched in the TAIR database for literature references associated to the function of these

Arabidopsis homologs. Given their large number, these references are not provided in the manuscript.

Phenotyping of natural populations

Height measurements

We measured height as described above in all 16 populations in the glasshouse and 12 of the

populations in their native field environment (Data S10 and Data S11). In the field, we measured

height in 32 individuals evenly across the range of each population. In the controlled conditions of the

glasshouse, we sampled an average of 14 individuals per population after plants reached maturity.

For both the glasshouse and field measurements, we used a linear model to determine whether Dune

populations were taller than their adjacent Headland pair: $y_{ijk} = P_i + E_{j(i)} + e_{k(ij)}$, where pair ($P_i$) is an

adjacent Dune and Headland population at the same locality and ecotype ($E_{j(i)}$) is Dune or Headland

and is nested within pair. All factors are fixed effects and $e_{k(ij)}$ is the residual error. Population height

(y_{ijk}) for each pair was compared using a one-tailed t-test. The Alpine populations were also included

with the prediction that the sheltered Alpine population (A03) would be taller than the wind exposed

Alpine population (A07). All statistical results reported here were produced in JMP 13 (SAS 2015).

Gravitropism measurements

Gravitropism was measured *in vitro* on agar plates in a dark growth cabinet using seeds from all 16

natural populations. For each population, 2-4 seeds per family was grown for ~40 maternal families

(1,278 seeds in total). The seeds from a population were combined in a falcon tube and were first

sterilized with a quick rinse in 70% EtOH, followed by four 10 minute inversions in a sterilizing solution

of 6% sodium hypochlorite and 1% Tween 20. Seeds were then rinsed three times with distilled water

and vertically orientated on Murashiga and Skoog agar plates containing 0.15% MS, 0.05% MES,

0.15% sucrose and 1% agar. We placed eight seeds on each plate and incubated the plates at 21°C

in the dark to avoid any light effects. After seven days, all plates were rotated clockwise by 90° and a

photograph of each plate was taken 24 hours after rotation. The photographs were imported into

ImageJ (73) to determine gravitropism by measuring the angle to which the stem reorientated to the

horizontal (74-76).

Overall there was a 63.8% germination success but we excluded seedlings that were shorter than

5mm, contacted the edge of the plate, or germinated after rotation. This left a total of 736 seedlings

across all 16 populations (57.6% of the total number of seeds planted; Data S11). To test the

hypothesis that Dune populations would have a stronger gravitropic response in their stem than their

adjacent Headland pair, we used a mixed linear model: $y_{ijkl} = P_i + E_{j(i)} + A_k + e_{l(ijk)}$, where pair ($P_i$) is an

adjacent Dune and Headland population at the same locality, ecotype ($E_{j(i)}$) is Dune or Headland and

is nested within pair, and agar plate (A_k) is the MS plate that the seeds were grown on. Agar plate was

fitted as a random effect while the rest were fixed effects, and $e_{l(ijk)}$ is the residual error. Gravitropism

measures were averaged for each population and compared between each population pair using a

one-tailed t-test. We then tested the correlation between height and gravitropism by performing a

linear regression with mean height against mean gravitropism for all 16 populations, where

populations were grouped into their respective clades (eastern and south-eastern). All statistical

results reported here were produced in JMP 13 (SAS 2015).

Synthetic auxin and auxin transport inhibitor experiments

We tested if gravitropism is governed by auxins in *S. lautus* by evaluating gravitropic responses in

seedlings treated with chemicals affecting auxin signaling. Specifically we used 2,4-

Dichlorophenoxyacetic acid (2,4-D), a carrier-dependent synthetic auxin (29, 77) and

Naphthylphthalamic acid (NPA) an efflux inhibitor (30). We collected seeds from Lennox Head sand

dune and rocky headland and grew them on MS containing the different chemicals at concentrations

of 0 µM, 0.5 µM, 5 µM and 50µM. For this we used the following stock solutions: 2,4-D: 1mM in

Ethanol and NPA: 10mM in DMSO. We created 1ml dilutions of stock solutions (in ethanol or DMSO),

which were dissolved in 500 ml of media. Plates were incubated, rotated, photographed and

gravitropism was measured in ImageJ (73), as outlined above.

Approximately 40 seeds from different maternal families were planted for each population and

chemical concentration. Seeds were excluded from analyses following same procedure above,

leaving a total of 188 seedlings to calculate gravitropism. Treatments where few seeds germinated

were excluded (Data S12). For each chemical we used a mixed linear model using the lmer function

of lme4 package in R v3.1.3(78): $y_{ijkl} = A_k + E_i + C_j + E_i \times C_j + e_{l(ijk)}$, where agar plate ($A_k$) is the MS

plate that the seeds were grown on, ecotype (E_i) is Dune or Headland and concentration (C_j) are the 4

different concentrations of the chemical. Agar plate was fitted as a random effect, while ecotype,

concentration and their interaction ($E_i \times C_j$) were fixed effects, and $e_{l(ijk)}$ is the residual error.

Gravitropism (y_{ijkl}) was compared using a Type II Wald chi-square test.

Field experiments

All field experiments were conducted at the sand dune and rocky headland at Lennox Head, NSW, in

the same location where native *S. lautus* grow. We tracked each individual by gluing each seed to a

toothpick and placing 1-2mm under the ground within a grid cell (Fig. 3b) that was randomly assigned

(for details see ref. (79) and ref. (20)). 50% shade cloth was initially suspended over all plots to

replicate the shade given by surrounding vegetation. Seeds were watered twice a day to keep the soil

moist and replicate ideal germination conditions to maximize the number of seeds in the experiment.

Once germination plateaued for all genotypes watering was gradually ceased.

Height adaptation experiments

We created genetic lines that aimed to isolate height differences on a common genomic background

(Fig. 1). Firstly, Lennox Head Dune and Headland seeds were grown and crossed to create an F1

generation. Secondly, to isolate the height alleles on a common genomic background, we

introgressed the tall trait onto a headland genomic background by backcrossing the tallest F1 to a

Headland parental. This created a BC1F1 generation, where the tallest individuals (n=11, tallest 20%

of the population) were again backcrossed to the Headland parental, creating a BC2F1 generation.

We then grew and crossed the tallest BC2F1 individuals (n=18, tallest 10% of the population) among

one another and the shortest individuals (n=19, shortest 10% of the population) among one another.

These BC2F2 seeds were planted into the rocky headland at Lennox Head on the 2nd October 2016

(Australian spring). We planted five replicate plots, where each plot (1.08x0.33m) consisted of the

same 12 families with four (occasionally three) individuals per family in each plot, totalling 843 BC2F2

seeds. The entire field experiment had a total of 4,708 seeds (108 families) but the other genotypes

were not relevant to this study. Germination and mortality were recorded every day for 49 days, then

every 3-4 days until day 79 and then weekly for the remainder of the experiment (Data S13).

We implemented a mixed linear model to test the hypothesis that individuals with short parents will

have higher fitness in the headland environment: $y_{ijkl} = H_i + F_j + B_k + e_{l(ijk)}$, where parental height (H_i)

was the average height of the parents measured in the glasshouse, family (F_j) was individuals with the

same parents, and block (B_k) is the five replicate plots across the rocky headland. Parental height is a

fixed effect, and family and block are random effects, and $e_{l(ijk)}$ was the residual error. Offspring fitness

(y_{ijk}) was the total number of days alive in the rocky headland from planting. All statistical results

reported here were produced in JMP 13 (SAS 2015).

Gravitropism adaptation experiments

We created an advanced recombinant population (F8) from 30 Lennox Head Dune and Headland

individuals as described earlier. We replicated the construction of the F8 using three independent

replicate crossing lines (A, B and C), all derived from the same base population. F8 seeds were

planted into the sand dune and rocky headlands as described above and the 50% highest surviving

families within an environment (and genetic line) were identified, and full-sibling seeds from these

families were grown in the glasshouse. To produce the subsequent generations, we used an among-

family based selection scheme (80) by conducting crosses among the best surviving families for both

the sand dune and rocky headland separately. In crossing the surviving families, we also doubled the

crosses per family, ensuring we maintained ~100 full-sibling families for each selection generation

planted into the natural environments. Selection was conducted on the F8, F9 and F10 generations.

We selected the top 50% of surviving families in the F8, F9 and F10 field selection experiment using

Aster modelling (81, 82) implemented with the 'Aster' package in R (78). The fitness components

included germination and survival success, where germination was the total number of individuals that

germinated in each family and survival success was the total number of individuals per family that

survived to day 85 in the F8 generation, day 138 in the F9 generation, and day 232 in the F10

generation. Finally, to produce the F11 generation, we crossed the F10 generation such that survivors

were designated as sires and dams. We then crossed two sires each to two dams in a full-sib, half-sib

crossing design.

F11 gravitropism measurements

We measured gravitropism, in the F11 generation described above, as the angle of the stem after a

90° rotation of a seedling (39 Dune families, 37 Headland families and 25 inbred control families with

12 individuals per family (1,212 seeds in total were germinated). The families were germinated in

three separate batches ~7 days apart. Briefly, we germinated the F11 seeds by cutting 1mm off the

micropyle side of the seed and placing in petri dishes with dampened filter paper. The seeds were

placed in darkness for two days for roots to grow and then transferred to light for 4 days for shoots to

grow. Seedlings were then transferred into small square pots with standard potting mix in a constant

temperature room at 25°C with a 12 hours light/day cycle. After one week of growing in the pot, the

plants were rotated by 90° and a photograph of each individual was taken 12 hours after rotation. The

photographs were imported into ImageJ (73) to measure gravitropism as described above. Data S14

contains gravitropism measures and dam and sire fitness values.

Gravitropism tests of selection

We implemented a linear model to test the hypothesis that high fitness dune families will produce

gravitropic plants and high fitness headland families will produce agravitropic plants. Independent

models were used for the sand dune and rocky headland to test the effect of gravitropism on fitness in

each environment: $y_{ijklmn} = B_i + V_j + R_{k(i)} + D_{l(ik)} + S_{m(ik)} + e_{n(ijklm)}$, where time-block ($B_i$) is the three time

points in which the F11 seeds were grown (~seven days apart); viability (V_j) is the number of days

until the death of F11 plants in controlled conditions; and replicate, which consists of the three

independent genetic replicates, is nested within time-block ($R_{k(i)}$). Dam fitness was nested in replicate

and time-block ($D_{l(ik)}$) and sire fitness was also nested in replicate and time-block ($S_{m(ik)}$). Dam and sire

fitness are the F10 family fitness values for the individuals that were crossed to create the F11

offspring where gravitropism was measured. All factors were included as fixed effects and $e_{n(ijklm)}$ was

the residual error. Replicate C was removed from analyses as it has little variation in fitness values

which means it did not converge. Shapiro-Wilk W test shows the residuals from the model are

normally distributed for both the sand dune (W=0.98, p=0.3879) and rocky headland (W=0.98,

p=0.2776). The linear model was performed in JMP 13 (SAS 2015).

Genetic association between height and gravitropism

We tested the genetic association between height and gravitropism after segregation in an advanced

recombinant population. We implemented a mixed linear model for the three F11 populations (Dune

survivors, Headland survivors and a Control population) that accounts for family variation: $y_{ijk} = H_i + F_j$

$+ e_{k(ij)}$, where gravitropism (y_{ij}) is the angle of the growth response 12 hours after a 90° rotation, height

(H_i) is the vertical distance from the soil to the top of the vegetative leaves, measured after maturity in

the glasshouse and family (F_j) is a random effect that consists of individuals that have the same dam

and sire. The mixed linear model was performed in JMP 13 (SAS 2015).

Genotyping of F11 gravitropism tails

To isolate gravitropism candidate genes we genotyped 77 gravitropic (>56°) and 68 agravitropic

(<20°) F11 individuals (Data S14). We extracted DNA from leaf tissue using the CTAB protocol (53)

with column cleaning and overnight incubation and quantified the DNA using the Picogreen reagent

(Invitrogen, Carlsbad, CA). To determine which parent the alleles were derived from, we included 39

Dune parentals (D01) and 41 Headland parentals (H01). Leaves from the Lennox Head Dune and

Headland natural populations were collected directly from the field and the same DNA extraction

protocol was followed. To increase accuracy of genotyping, each F11 individual was duplicated in

independent wells and libraries of Restriction-site Associated DNA (RAD) tags were created at

Floragenex following the protocol from ref. (58) but using the PstI restriction enzyme. We sequenced

380 samples on four lanes of an Illumina HiSeq 4000 with 91bp single end reads at Floragenex. A

total of 1.39 billion reads with a mean of 3.62 million reads per sample were produced. Reads were

aligned to the reference genome using Bowtie 1.1.1 (83) and a FASTQ quality score of above 20. The

gravitropism candidate gene set consisted of SNPs in the 99.9% quantile of the distribution of

differentiated SNPs between the gravitropism tails. The region of the scaffold containing the SNP was

annotated using the blastx NCBI database (84).

Senecio lautus gravitropism candidate genes

In the gravitropism candidate gene set we tested for overrepresentation of gene function and linkage

disequilibrium between gravitropism loci. Statistical overrepresentation test was performed in

PANTHER (http://pantherdb.org/) using the TAIR identification for 32 unique gravitropism candidate

genes matched to a reference list of 27,502 *Arabidopsis thaliana* genes. To calculate linkage

disequilibrium between loci, a likelihood-ratio chi-square test was performed in JMP 13 (SAS 2015)

with each F11 group independently (Control population, Dune survivors and Headland survivors; Data

S15).

Intrinsic reproductive isolation

We tested whether intrinsic reproductive isolation was associated with the gravitropism phenotype by

randomly crossing within and between the F11 gravitropic and agravitropic groups in a controlled

temperature room and recording successful and failed crosses. To maximize sample size, all three

replicate genetic lines (A, B and C) were used across all three populations (Dune survivors, Headland

survivors and Control). A total of 132 crosses were completed (Data S16), 67 crosses between the tails and 65 within the tails (agravitropic tail = 28 and gravitropic tail 37). Crosses were completed by rubbing multiple flower heads of two individuals together and collecting the seeds produced from both plants. To remove genetic incompatibilities that might be caused by relatedness, crosses within the same family were not performed. A failed cross was considered when, in the presence of pollen, less than three seeds were produced per flower head from both plants with three mating attempts. We performed a two-tailed fisher's exact test in JMP 13 (SAS. 2015) to determine whether there was a significant association between cross type (within vs between gravitropism tails) and failed crosses.

Next, we tested whether intrinsic reproductive isolation was associated with ENDOL1 and *ABA3* by calculating the percentage of failed crosses with each allelic combination in the F11 population (Data S17 and Data S18).

Acknowledgments

We are grateful to S. Smith, L.H. Rieseberg, M. Cooper, J. Engelstaedter, M.A.F. Noor and members of the Ortiz-Barrientos laboratory for insightful comments on previous versions of this manuscript. S. Karrenberg and S. Chenoweth provided very useful feedback on M.J. Wilkinson PhD dissertation. We thank P. Brewer for his help in the design and execution of gravitropism experiments, T. Richards for his contributions to the construction of recombinant populations for BSA, and J. Donohoe for his help in the phenotyping and rearing of plants for BSA. We would also like to thank everyone that helped with the field selection experiments: E. Johnston, E. Monley, G. Wilkinson, P. Wilkinson, A. Mather, S. Thang, K. Giarola, G. Lebbink, E. Firkins-Barriere, P. Arnold, J. Donohoe, B. Brittain, H. Liu, D. Bernal, M.C. Melo, T. Richards, J. Walter, L. Ambrose, B. Ayalon, S. Carrol, M. Gallo, A. Maynard, C. Palmer and S. Edgley.

Author Contributions

MJW, FR and DO conceived the projects and experiments. MJW conducted height adaptation experiments, reared and phenotyped various glasshouse populations and performed reproductive isolation experiment. FR created BSA mapping populations and performed BSA analyses. FR and

MJW performed physiological and molecular biology experiments. GMW reared and phenotyped the

natural populations in the glasshouse and conducted the field experiment for the gravitropism

adaptation experiments with assistance from MJW, MEJ, FR and DO. DMB performed expression

analyses. MJW, MEJ and HLN phenotyped natural populations. RN and JW called the genotypes of

the F11 population. SA assembled the current draft genome of *S. lautus*. CB guided the physiological

experiments. MJW, FR, DO wrote the paper with input from all authors. DO is mentor and supervisor

for the research program.

References

1. Darwin C (1859) *On the origins of species by means of natural selection* (London: Murray) p 1859.
2. Fishman L & Sweigart AL (2018) When two rights make a wrong: The evolutionary genetics of plant hybrid incompatibilities. *Annu. Rev. Plant Biol.*
3. Wright KM, Lloyd D, Lowry DB, Macnair MR, & Willis JH (2013) Indirect evolution of hybrid lethality due to linkage with selected locus in *Mimulus guttatus*. *PLoS Biol.* 11(2):e1001497.
4. Stinchcombe JR & Hoekstra HE (2007) Combining population genomics and quantitative genetics: finding the genes underlying ecologically important traits. *Heredity* 100(2):158-170.
5. Barrett RDH & Schluter D (2008) Adaptation from standing genetic variation. *Trends Ecol. Evol.* 23(1):38-44.
6. Radford IJ, Cousens RD, & Michael PW (2004) Morphological and genetic variation in the *Senecio pinnatifolius* complex: are variants worthy of taxonomic recognition? *Aust. Syst. Bot.* 17(1):29-48.
7. Thompson I (2005) *Taxonomic studies of Australian Senecio (Asteraceae): 5. The S. lautus/S. lautus complex.* (Royal Botanic Gardens, Melbourne).
8. Roda F*, et al.* (2013) Convergence and divergence during the adaptation to similar environments by an Australian groundsel. *Evolution* 67(9):2515-2529.
9. Friml J (2003) Auxin transport—shaping the plant. *Curr. Opin. Plant Biol.* 6(1):7-12.
10. Xu M, Zhu L, Shou H, & Wu P (2005) A PIN1 family gene, OsPIN1, involved in auxin-dependent adventitious root emergence and tillering in rice. *Plant Cell Physiol.* 46(10):1674-1681.
11. Wallace JG*, et al.* (2016) Genome-wide association for plant height and flowering time across 15 tropical maize populations under managed drought stress and well-watered conditions in Sub-Saharan Africa. *Crop Sci.* 56(5):2365-2378.
12. Gallavotti A (2013) The role of auxin in shaping shoot architecture. *J. Exp. Bot.* 64(9):2593-2608.
13. Domagalska Ma & Leyser O (2011) Signal integration in the control of shoot branching. *Nature Reviews Molecular Cell Biology* 12:211-221.
14. Sundberg E & Østergaard L (2009) Distinct and dynamic auxin activities during reproductive development. *Cold Spring Harbor perspectives in biology* 1(6):a001628-a001628.
15. Vanneste S & Friml J (2009) Auxin: a trigger for change in plant development. *Cell* 136:1005-1016.
16. Walter GM*, et al.* (2016) Diversification across a heterogeneous landscape. *Evolution* 70(9):1979-1992.
17. Melo MC, Grealy A, Brittain B, Walter GM, & Ortiz-Barrientos D (2014) Strong extrinsic reproductive isolation between parapatric populations of an Australian groundsel. *New Phytol.* 203(1):323-334.
18. Richards TJ & Ortiz-Barrientos D (2016) Immigrant inviability produces a strong barrier to gene flow between parapatric ecotypes of *Senecio lautus*. *Evolution* 70(6):1239-1248.
19. Richards TJ, Walter GM, McGuigan K, & Ortiz-Barrientos D (2016) Divergent natural selection drives the evolution of reproductive isolation in an Australian wildflower. *Evolution* 70(9):1993-2003.
20. Walter GM, Aguirre JD, Blows MW, & Ortiz-Barrientos D (2018) Evolution of genetic variance during adaptive radiation. *The American Naturalist* 191(4):E108-E128.
21. Michelmore RW, Paran I, & Kesseli RV (1991) Identification of markers linked to disease-resistance genes by bulked segregant analysis: a rapid method to detect markers in specific genomic regions by using segregating populations. *Proc. Natl. Acad. Sci. USA* 88(21):9828-9832.

22.   Roda F, *et al.* (2013) Genomic evidence for the parallel evolution of coastal forms in the
Senecio lautus complex. *Mol. Ecol.* 22(11):2941-2952.

23.   Petrasek J & Friml J (2009) Auxin transport routes in plant development. *Development*
136(16):2675-2688.

24.   Roda F, Walter GM, Nipper R, & Ortiz-Barrientos D (2017) Genomic clustering of adaptive
loci during parallel evolution of an Australian wildflower. *Mol. Ecol.* 26(14):3687–3699.

25.   Tan L, *et al.* (2008) Control of a key transition from prostrate to erect growth in rice
domestication. *Nat. Genet.* 40:1360-1364.

26.   Li P, *et al.* (2007) LAZY1 controls rice shoot gravitropism through regulating polar auxin
transport. *Cell Res.* 17:402-410.

27.   Lopez D, Tocquard K, Venisse J-S, Legué V, & Roeckel-Drevet P (2014) Gravity sensing, a
largely misunderstood trigger of plant orientated growth. *Frontiers in Plant Science* 5(610).

28.   Blancaflor EB (2013) Regulation of plant gravity sensing and signaling by the actin
cytoskeleton. *Am. J. Bot.* 100(1):143-152.

29.   Yamamoto M & Yamamoto KT (1998) Differential effects of 1-naphthaleneacetic acid,
indole-3-acetic acid and 2,4-dichlorophenoxyacetic acid on the gravitropic response of roots
in an auxin-resistant mutant of *Arabidopsis*, *auxl*. *Plant Cell Physiol.* 39(6):660-664.

30.   Ottenschläger I, *et al.* (2003) Gravity-regulated differential auxin transport from columella to
lateral root cap cells. *Proc. Natl. Acad. Sci. USA* 100(5):2987-2991.

31.   Ranocha P, *et al.* (2010) Walls are thin 1 (WAT1), an *Arabidopsis* homolog of *Medicago*
truncatula NODULIN21, is a tonoplast-localized protein required for secondary wall
formation in fibers. *The Plant Journal* 63(3):469-483.

32.   Promchuea S, Zhu Y, Chen Z, Zhang J, & Gong Z (2017) ARF2 coordinates with PLETHORAs
and PINs to orchestrate ABA-mediated root meristem activity in *Arabidopsis*. *Journal of*
Integrative Plant Biology 59(1):30-43.

33.   Wu H, Shen Y, Hu Y, Tan S, & Lin Z (2011) A phytocyanin-related early nodulin-like gene,
BcBCP1, cloned from *Boea crassifolia* enhances osmotic tolerance in transgenic tobacco. *J.*
Plant Physiol. 168(9):935-943.

34.   Ma H, Zhao H, Liu Z, & Zhao J (2011) The phytocyanin gene family in rice (Oryza sativa L.):
Genome-wide identification, classification and transcriptional analysis. *PLOS One*
6(10):e25184.

35.   Garcia de la Garma J, *et al.* (2015) New insights into plant salt acclimation: the roles of
vesicle trafficking and reactive oxygen species signalling in mitochondria and the
endomembrane system. *New Phytol.* 205(1):216-239.

36.   Mittler R & Blumwald E (2015) The roles of ROS and ABA in systemic acquired acclimation.
The Plant Cell 27(1):64-70.

37.   Hou Y, *et al.* (2016) Maternal ENODLs are required for pollen tube reception in *Arabidopsis*.
Curr. Biol. 26(17):2343-2350.

38.   Beeftink W, Rozema J, & Huiskes A (1985) *Ecology of coastal vegetation* (Springer).

39.   Auld T & Morrison D (1992) Genetic determination of erect and prostrate growth habit in
five shrubs from windswept headlands in the Sydney region. *Aust. J. Bot.* 40(1):1-11.

40.   Morrison D & Rupp A (1995) Patterns of morphological variation within *Acacia suaveolens*
(Mimosaceae). *Aust. Syst. Bot.*:1013-1027.

41.   Crutsinger GM, Strauss SY, & Rudgers Ja (2010) Genetic variation within a dominant shrub
species determines plant species colonization in a coastal dune ecosystem. *Ecology* 91:1237-
1243.

42.   Li T, *et al.* (2019) Calcium signals are necessary to establish auxin transporter polarity in a
plant stem cell niche. *Nature Communications* 10(1):726.

43.   Naser V & Shani E (2016) Auxin response under osmotic stress. *Plant Mol. Biol.* 91(6):661-
672.

44. van den Berg T, Korver RA, Testerink C, & ten Tusscher KH (2016) Modeling halotropism: a
key role for root tip architecture and reflux loop remodeling in redistributing auxin.
Development 143(18):3350-3362.
45. Galvan-Ampudia CS*, et al.* (2013) Halotropism is a response of plant roots to avoid a saline
environment. *Curr. Biol.* 23(20):2044-2050.
46. Guan Y, Guo J, Li H, & Yang Z (2013) Signaling in pollen tube growth: Crosstalk, feedback, and
missing links. *Molecular Plant* 6(4):1053-1064.
47. Tuteja N (2007) Abscisic acid and abiotic stress signaling. *Plant Signaling & Behavior*
2(3):135-138.
48. Guo Q, Major IT, & Howe GA (2018) Resolution of growth-defense conflict: mechanistic
insights from jasmonate signaling. *Curr. Opin. Plant Biol.* 44:72-81.
49. Bomblies K & Weigel D (2007) Hybrid necrosis: autoimmunity as a potential gene-flow
barrier in plant species. *Nature Reviews Genetics* 8:382.
50. Luo Y*, et al.* (2015) A single nucleotide deletion in *gibberellin20-oxidase1* causes alpine
dwarfism in *Arabidopsis*. *Plant Physiol.* 168(3):930.
51. Gould BA, Chen Y, & Lowry DB (2017) Pooled ecotype sequencing reveals candidate genetic
mechanisms for adaptive differentiation and reproductive isolation. *Mol. Ecol.* 26(1):163-
177.
52. Lowry DB & Willis JH (2010) A widespread chromosomal inversion polymorphism contributes
to a major life-history transition, local adaptation, and reproductive isolation. *PLoS Biol.*
8(9):e1000500.
53. Clarke JD (2009) Cetyltrimethyl ammonium bromide (CTAB) DNA miniprep for plant DNA
isolation. *Cold Spring Harbor Protocols* 2009(3):pdb.prot5177.
54. Liu H (2015) Developing genomic resources for an emerging ecological model species
Senecio lautus. PhD Thesis (The University of Queensland).
55. Koren S*, et al.* (2017) Canu: scalable and accurate long-read assembly via adaptive k-mer
weighting and repeat separation. *Genome Res.* 27(5):722-736.
56. Ali S (1964) *Senecio lautus* complex in Australia. I. Taxonomic considerations and discussion
of some of the related taxa from New Zealand. *Aust. J. Bot.* 12(2):282-291.
57. Rockman MV & Kruglyak L (2008) Breeding designs for recombinant inbred advanced
intercross lines. *Genetics* 179(2):1069-1078.
58. Baird NA*, et al.* (2008) Rapid SNP discovery and genetic mapping using sequenced RAD
markers. *PLOS One* 3(10):1-7.
59. Kofler R, Pandey RV, & Schlotterer C (2011) PoPoolation2: identifying differentiation
between populations using sequencing of pooled DNA samples (Pool-Seq). *Bioinformatics*
27(24):3435-3436.
60. Li H & Durbin R (2009) Fast and accurate short read alignment with Burrows–Wheeler
transform. *Bioinformatics* 25(14):1754-1760.
61. Li H*, et al.* (2009) The sequence alignment/map format and SAMtools. *Bioinformatics*
25(16):2078-2079.
62. Rosenzweig BK, Pease JB, Besansky NJ, & Hahn MW (2016) Powerful methods for detecting
introgressed regions from population genomic data. *Mol. Ecol.* 25(11):2387-2397.
63. Benfey PN & Mitchell-Olds T (2008) From Genotype to Phenotype: Systems Biology Meets
Natural Variation. *Science* 320(5875):495-497.
64. Futschik A & Schlötterer C (2010) The next generation of molecular markers from massively
parallel sequencing of pooled DNA samples. *Genetics* 186(1):207-218.
65. Magwene PM, Willis JH, & Kelly JK (2011) The statistics of bulk segregant analysis using next
generation sequencing. *PLoS Comp. Biol.* 7(11):e1002255.
66. Simms D, Cizdziel PE, & Chomczynski PJF (1993) TRIzol: A new reagent for optimal single-step
isolation of RNA. 15(4):532-535.

67.   Schroeder A*, et al.* (2006) The RIN: an RNA integrity number for assigning integrity values to
842   RNA measurements. 7(1):3.

843   68.   Kim D, Langmead B, & Salzberg SL (2015) HISAT: a fast spliced aligner with low memory
844   requirements. *Nat. Methods* 12(4):357-360.

845   69.   Pertea M*, et al.* (2015) StringTie enables improved reconstruction of a transcriptome from
846   RNA-seq reads. *Nat. Biotechnol.* 33(3):290-295.

847   70.   Pertea M, Kim D, Pertea GM, Leek JT, & Salzberg SL (2016) Transcript-level expression
848   analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nature Protocols*
849   11(9):1650-1667.

850   71.   Huang da W, Sherman BT, & Lempicki RA (2009) Systematic and integrative analysis of large
851   gene lists using DAVID bioinformatics resources. *Nature Protocols* 4(1):44-57.

852   72.   Huang da W, Sherman BT, & Lempicki RA (2009) Bioinformatics enrichment tools: paths
853   toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* 37(1):1-
854   13.

855   73.   Schneider CA, Rasband WS, & Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image
856   analysis. *Nat. Methods* 9(7):671-675.

857   74.   Sang D*, et al.* (2014) Strigolactones regulate rice tiller angle by attenuating shoot
858   gravitropism through inhibiting auxin biosynthesis. *Proc. Natl. Acad. Sci. USA* 111(30):11199-
859   11204.

860   75.   Rigo G*, et al.* (2013) Inactivation of plasma membrane-localized CDPK-RELATED KINASE5
861   decelerates PIN2 exocytosis and root gravitropic response in *Arabidopsis*. *The Plant Cell*
862   25(5):1592-1608.

863   76.   Rashotte AM, Brady SR, Reed RC, Ante SJ, & Muday GK (2000) Basipetal auxin transport is
864   required for gravitropism in roots of *Arabidopsis*. *Plant Physiol.* 122(2):481-490.

865   77.   Yang Y, Hammes UZ, Taylor CG, Schachtman DP, & Nielsen E (2006) High-affinity auxin
866   transport by the AUX1 influx carrier protein. *Curr. Biol.* 16(11):1123-1127.

867   78.   Team RC (2013) R: A language and environment for statistical computing. *R Foundation for*
868   *Statistical Computing, Vienna, Austria*.

869   79.   Walter GM (2016) The genetic and ecological basis of diversification. PhD Thesis (The
870   University of Queensland).

871   80.   Walsh B & Lynch M (2013) *Evolution and Selection of Quantitative Traits: I. Foundations* p 1.

872   81.   Geyer CJ, Wagenius S, & Shaw RG (2007) Aster models for life history analysis. *Biometrika*
873   94(2):415-426.

874   82.   Shaw RG, Geyer CJ, Wagenius S, Hangelbroek HH, & Etterson JR (2008) Unifying life-history
875   analyses for inference of fitness and population growth. *The American Naturalist* 172(1):E35-
876   E47.

877   83.   Langmead B, Trapnell C, Pop M, & Salzberg SL (2009) Ultrafast and memory-efficient
878   alignment of short DNA sequences to the human genome. *Genome Biology* 10(3):R25.

879   84.   Altschul SF, Gish W, Miller W, Myers EW, & Lipman DJ (1990) Basic local alignment search
880   tool. *J. Mol. Biol.* 215(3):403-410.

Figures and Tables

883
Fig. 1. Sample locations and growth habit differences between adjacent *S. lautus* populations.
a, Map of Australia showing locations of the 16 populations used in this study. The seven coastal
localities have a Dune (orange) and Headland (green) population occurring adjacent to each other.
The populations are split into two monophyletic clades (22), the eastern clade (closed circles) and the
south-eastern clade (open circles). **b,** *Senecio lautus* native to the sand dunes have an erect growth
habit and *S. lautus* native to the rocky headlands have a prostrate growth habit. **c,** Alpine populations
of *S. lautus* include a sheltered alpine gully and a wind exposed alpine meadow, containing
individuals with an erect and prostrate growth habit, respectively.

Fig. 2. Gravitropism and height variation across 16 *S. lautus* populations. a, The correlation between gravitropism and height across *S. lautus* populations split into their monophyletic clades – see Fig. 1 for details. Each point in the graph represents a population mean where height was measured in the glasshouse and gravitropism was measured 24 hours after a 90° rotation. **b,c**, Divergence in gravitropism, height in the glasshouse and height in the field between adjacent *S. lautus* populations (D = Dune, H = Headland, A-S=Alpine Sheltered and A-E = Alpine Exposed). **b**, South-eastern clade and **c**, southern clade. Height in the field for Falls Creek, Kosciusko NP and Point Labatt were not measured. Data are mean ± s.e.m.; one tailed Student's t-test, *P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.001, NS not significant.

904
905
Fig. 3. Parental survival curves in the height (2016) and gravitropism (2012-2014) adaptation
experiments at the sand dune and rocky headland at Lennox Head. a,**b**, Survival over the length
of the field experiments is shown for the Lennox Head Dune population (orange) and the Lennox
Head Headland population (green) for four independent field selection experiments at the sand dune
(**a**) and rocky headland (**b**). Asterisks indicate a significant difference in mortality risk between the
Dune and Headland ecotypes (**P ≤ 0.01, ***P ≤ 0.001).

912
913
Fig. 4. The creation of the recombinant hybrid generation, the design of the gravitropism
adaptation experiments and sequencing of the tails of the gravitropic distribution. a, A total of
30 parental Dune and Headland individuals from Lennox Head were crossed randomly and with equal
contribution for eight generations. **b**, Seeds from this F8 recombinant population were glued to
toothpicks and transplanted into the sand dune and rocky headland at Lennox Head. Among-family
based selection occurred for three generations (F8, F9 and F10), where full-siblings from the fittest
families were grown in the glasshouse and crossed amongst their respective genetic lines (A, B and
C) and their environment (Dune survivors or Headland survivors). An inbred control was kept in the
glasshouse and underwent the same crossing scheme but free from viability selection. **c**,
Gravitropism was measured in the F11 recombinant population by re-orientating the plant by 90°.
Agravitropic plants have small gravitropic angles, while gravitropic plants have large gravitropic
angles as they reorient their growth and grow upright. Individuals in the tails of the gravitropism
distribution were sequenced on four lanes of the Illumina HiSeq 4000 platform.

927
928 **Fig. 5. Association between *ENODL1* and *ABA3* alleles and gravitropism in *S. lautus*.** The
929 average gravitropism angle is shown for each allelic combination, independently and when the
930 agravitropic alleles (dark grey) are combined. Different letters denote significant differences between
931 genotypes (t>1.98, p<0.05).

Table 1. General linear model for the effect of dam and sire on gravitropism (°) after a field selection experiment on a recombinant hybrid Dune and Headland population. Field selection experiments were performed on F8, F9 and F10 recombinant hybrid generations to achieve three rounds of selection in the sand dune and rocky headland at Lennox Head (see Fig. 4 for the experimental design). Dam and sire fitness are the F10 family fitness values for the individuals that were crossed to create the F11 offspring where gravitropism was measured. Intrinsic viability is the number of days until death of the F11 generation in the controlled temperature room. This experiment was conducted three times (temporal block) with three independent genetic replicates.

Source	Dune				Headland			
	DF	SS	F-Ratio	Prob > F	DF	SS	F-Ratio	Prob > F
Dam family fitness	6	8515.77	4.779	<0.001	6	1884.31	0.701	0.650
Sire family fitness	6	1806.62	1.014	0.424	5	2315.36	1.033	0.405
Intrinsic viability	1	260.85	0.878	0.352	1	5209.38	11.624	0.001
Genetic replicate	3	1135.49	1.275	0.290	3	3357.19	2.497	0.067
Temporal block	2	193.14	0.325	0.724	2	2234.96	2.494	0.090

DF, degrees of freedom; SS, sum of squares.