Comparative transcriptomics of a monocotyledonous geophyte reveals shared molecular mechanisms of underground storage organ formation.

Carrie M. Tribble[1*], Jesús Martínez-Gómez[2], Fernando Alzate-Guarin[3], Carl J. Rothfels[1], Chelsea D. Specht[2]

1. Department of Integrative Biology and the UC and Jepson Herbaria, University of California, Berkeley, Berkeley, CA 94720

2. School of Integrative Plant Sciences and L.H. Bailey Hortorium, Cornell University, Ithaca, NY 14853 USA

3. Grupo de Estudios Botánicos (GEOBOTA) and Herbario Universidad de Antioquia (HUA), Instituto de Biología, Facultad de Ciencias Exactas y Naturales, Universidad de Antioquia, Calle 67 Nº 53-108, Medellín, Colombia

Corresponding author: Carrie M Tribble, ctribble09@gmail.com, (510) 643-0633

Summary

- Many species from across the vascular plant tree-of-life have modified standard plant tissues into tubers, bulbs, corms, and other underground storage organs (USOs). *Bomarea multiflora* (Alstroemeriaceae) is a tropical climbing monocot with unique underground morphology, including tuberous roots. We take a comparative transcriptomics approach to characterizing the molecular mechanisms of tuberous root formation in *B. multiflora* and compare these mechanisms to those identified in other underground storage structures across diverse plant lineages.

- We sequenced transcriptomes from the growing tip of four tissue types (aerial shoot, rhizome, fibrous root, and root tuber) of three individuals of *B. multiflora*. We identify differentially expressed isoforms and test the expression of candidate genes that have been implicated in underground storage in other taxa.

- We identify 271 genes that are differentially expressed in root tubers versus fibrous roots, including genes implicated in cell wall modification, environmental signaling and flowering time, and starch biosynthesis. We also identify a phosphatidylethanolamine-binding protein (PEBP) that is over-expressed in tuberous roots.

- These findings demonstrate that deeply parallel processes underlie the formation of underground storage structures despite long evolutionary distances between taxa and non-homologous morphologies, illustrating that repeated co-option of similar genetic pathways can lead to convergent morphologies.

Introduction

The vast majority of scientific attention in botanical fields focuses exclusively on aboveground biomass. However, a holistic understanding of plant morphology, ecology, and evolution requires that considerable research effort go towards generating a comprehensive understanding of belowground biomass. While most studies of plant form and function focus on aboveground organs, on average 50% of an individual plant's biomass lies beneath the ground (Niklas, 2005). Often, belowground biomass is thought to consist solely of standard root tissue, but in some cases, plants modify 'ordinary' structures for specialized underground functions. Plants called "geophytes" are toward the extreme end of this belowground/aboveground allocation spectrum. These species rely on nutrients stored in belowground organs (underground storage organs or "USOs"), and their ephemeral aboveground parts resprout from buds located on belowground organs (Raunkiaer, 1934; Dafni *et al.*, 1981a,b; Al-Tardeh *et al.*, 2008; Veselý *et al.*, 2011). Geophytes are ecologically and economically important, morphologically diverse, and have evolved independently in all major groups of vascular plants except gymnosperms (Howard *et al.*, 2019a,b). These plants and their associated underground structures are a compelling example of evolutionary convergence; diverse taxa form a variety of structures, often from different tissues, that serve parallel physiological and ecological functions. However, our understanding of the molecular processes that drive this convergence, and the extent to which these processes are themselves convergent, remains remarkably limited, due in part to the lack of molecular studies in geophyte lineages. This is particularly true for monocotyledonous geophytic taxa, which comprise the majority of ecologically and economically important geophyte diversity.

Some of the world's most important crop plants are geophytes, including potato (stem tuber, *Solanum tuberosum*), sweet potato (tuberous root, *Ipomoea batatas*), yam (epicotyl- and hypocotyl-derived tubers, *Dioscorea spp.*)*,* cassava (tuberous root, *Manihot esculenta*), radish (swollen hypocotyl and taproot, *Raphanus raphanistrum*), onion (bulb, *Allium cepa*), lotus

(rhizome, *Nelumbo nucifera*), and more. While several of these crop plants are well studied and
have sequenced genomes or other genetic or genomic data that may inform the molecular
mechanisms underlying underground storage organ development, most research has focused on a
select few taxa that do not represent the diversity of geophyte morphology, taxonomy, or
ecology. In particular, most genetic research on geophytes and their associated underground
storage organs has been conducted in eudicotyledons (with the exception of important studies in
onion, such as in Navarro *et al.*, 2011), which neglects a huge proportion of geophytic diversity
in the monocots. Furthermore, geophytes form important components of many ecosystems,
particularly Mediterranean biomes and other seasonal habitats, as their underground nutrient
reserves fuel regrowth following periods of seasonal dormancy, prolonged dormancy, or short
term resource limitation (Rundel, 1996; Hoffmann *et al.*, 1998; Parsons, 2000; Proches *et al.*,
2006; Cuéllar-Martínez & Sosa, 2016; Sosa & Loera, 2017; Ott *et al.*, 2019; Howard *et al.*,
2019b). The geophytic habit has evolved multiple times across the vascular plant tree of life, and
geophyte lineages have been shown to diversify at faster rates than do related non-geophytic
taxa, particularly in taxa with bulbs, corms, and tubers (Howard *et al.*, 2019b), indicating that the
geophytic habit may be correlated with increased diversification and/or reduced extinction rates.
113
Underground storage organs originate from all major types of plant vegetative tissue: roots,
stems, leaves, and hypocotyls. Bulbs, corms, rhizomes, and tubers are some of the most common
underground storage organ morphologies (Pate & Dixon, 1982), but the full breadth of
morphological variation in USOs includes various root modifications (tuberous roots, taproots,
etc.), swollen hypocotyls that merge with swollen root tissue (e.g. *Adenia*: Hearn, 2009), and
intermediate structures such as rhizomes where the terminal end of the rhizome forms a bulb
from which aerial shoots emerge (e.g. *Iris*: Wilson, 2006). Despite this morphological
complexity, USOs all develop through the expansion of standard plant tissue, either derived from
the root or shoot, into swollen, discrete storage organs. These storage organs also serve similar
functions as belowground nutrient reserves (Veselý *et al.*, 2011), often containing starch or other
non-structural carbohydrates (NSCs), storage proteins, and water. The functional and
physiological similarities of underground storage organs may be driven by deep molecular
homology with parallel evolution in the underlying genetic architecture of storage organ
development, despite differences in organismal level morphology and anatomy.
128
The economic importance of some geophytes and the relevance of understanding the formation
of storage organs for crop improvement have motivated studies on the genetic basis for storage
organ development in select taxa. Potato has become a model system for understanding the
molecular basis of USO development, and numerous studies have demonstrated the complex and
interacting roles of plant hormones such as auxin, abscisic acid, cytokinin, and gibberellin on the
tuber induction process (reviewed in Hannapel *et al.*, 2017). These hormones have been
additionally identified in USO formation in other tuberous root crops including sweet potato
(Noh *et al.*, 2010; Dong *et al.*, 2019) and cassava (Melis & van Staden, 1985; Sojikul *et al.*,
2015), in rhizome formation in *Panax japonicus* var. *major* (Tang *et al.*, 2019) and *Nelumbo*
nucifera (Cheng *et al.*, 2013b; Yang *et al.*, 2015), and in corm formation in *Sagittaria trifolia*
(Cheng *et al.*, 2013a), suggesting that parallel processes trigger tuberization in both root- and
stem-originating USOs. Three primary mobile signaling genes have been implicated in triggering
the onset of tuberization in potato: SELF-PRUNING6A (StSP6A), a FLOWERING LOCUS T-
like gene (FT gene); StBEL5 in the BEL1-like transcription factor family; and POTH1, a KNOX
type transcription factor. FT-like genes have been additionally implicated in USO formation in
Dendrobium (Wang *et al.*, 2017), *Callerya speciosa* (Xu *et al.*, 2016), tropical lotus (*Nelumbo*

nucifera; Yang *et al.*, 2015), and onion (*Allium cepa*; Lee *et al.*, 2013), indicating either deep
homology of FT involvement in USO formation across angiosperms or multiple independent co-
option of FT orthologues in geophytic taxa.
148
The lateral expansion of roots into tuberous roots may be driven by either cellular proliferation
concurrent with primary growth (primary thickening growth; Kaplan, unpublished), by cellular
proliferation subsequent to primary growth (secondary thickening growth; Kaplan, unpublished),
by cellular expansion, where individual cells expand in size, or by a combination of these
processes. Expansion in plant cells requires the modification of the rigid cell wall to
accommodate increases in cellular volume (Dolan & Davies, 2004; Humphrey *et al.*, 2007), and
genes such as expansins have been implicated in cellular expansion during tuberous root
development in some taxa such as cassava and *Callerya speciosa* (Sojikul *et al.*, 2015; Xu *et al.*,
2016). Recent studies of the tuberous roots of sweet potato (*Ipomoea batatas*) and other
members of Convolvulaceae indicate that USO formation in these taxa involves a MADS-box
gene implicated in the vascular cambium (SRD1; Noh *et al.*, 2010) and a WUSCHEL-related
homeobox gene (WOX4; Eserman *et al.*, 2018), also involved in vascular cambium
development. Additional work on cassava (*Manihot esculenta*) also suggests that tuberous root
enlargement is due to secondary thickening growth originating in the vascular cambium
(Chaweewan & Taylor, 2015). However, geophytes are especially common in
monocotyledonous plants (Howard *et al.*, 2019a,b), which lack a vascular cambium entirely. No
previous study has addressed the molecular mechanisms of USO development in this major
clade, so the causes of root thickening are particularly enigmatic. Do monocots form tuberous
roots through genetic machinery that shares deep homology with the eudicot vascular-cambium-
related pathways, or have they evolved an entirely independent mechanism?
169
Bomarea multiflora (L. f.) Mirb. is a scandent monocotyledonous geophyte that is native to
Venezuela, Colombia, and Ecuador (Hofreiter, 2008) and is invasive in New Zealand (National
Pest Plant Accord: https://www.mpi.govt.nz/protection-and-response/long-term-pest-
management/national-pest-plant-accord). It typically grows in moist cloud forests between 1800 –
3800 meters elevation (Hofreiter, 2008). *Bomarea multiflora* is an excellent model in which to
study the molecular mechanisms underlying underground storage organ formation in the
monocots because it has two types of underground modifications: tuberous roots and rhizomes.
However, prior to this study, no genomic or transcriptomic data was available for any species of
Bomarea. Comparative transcriptomics permits comprehensive examination of the molecular
basis of development, tissue differentiation, and physiology in ecologically relevant taxa by
comparing the genes expressed in different organs, developmental stages, or ecological
conditions (Ekblom & Galindo, 2011; Oppenheim *et al.*, 2015). Because no prior genomic or
transcriptomic data is needed for comparative transcriptomic studies, this method is especially
appropriate for studies of non-model organisms and can yield novel insights into the expression
profiles of specific tissues. In this study, we investigate the molecular mechanisms underlying
the formation of tuberous roots in *Bomarea multiflora* using a comparative transcriptomics
approach and quantify the extent to which these mechanisms are shared across the taxonomic
and morphological breadth of geophytic taxa.
188
189 **Materials and Methods**
190
191 **Greenhouse and laboratory procedures** Seeds were collected from a single inflorescence of
192 *Bomarea multiflora* in Antioquia, Colombia [vouchered as Tribble 194, deposited at UC; Index

Herbariorium http://sweetgum.nybg.org/science/ih/] and germinated in greenhouse conditions at the University of California, Berkeley designed to replicate *B. multiflora*'s native conditions (70°F – 85°F and ~50% humidity). Six months after germination, three sibling individuals were harvested as biological replicates. The aerial shoot apical meristem (SAM) of a single branch, the rhizome apical meristem (RHI), root apical meristems (RAM) of several fibrous roots, and the growing tip of a tuberous root (TUB; Figure 1) were dissected from each of three individuals, for a total of 12 tissue samples. Samples were immediately frozen in liquid nitrogen and maintained at -80°C until extraction. Total RNA was extracted from all samples using the Agilent Plant RNA Isolation Mini Kit (Agilent, Santa Clara, Ca), optimized for non-standard plant tissues, especially those that may be high in starch. As this protocol did not work well for the vegetative SAM samples, RNA was extracted from two dissected SAMs using an alternative plant tissue RNA extraction protocol (Yockteng *et al.*, 2013). Quality of total RNA was measured with Qubit (ThermoFisher, Waltham MA) and Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA); if needed, a Sera-Mag bead clean-up was used to further clean extracted RNA (Yockteng *et al.*, 2013). Two SAM samples failed to extract at sufficient concentrations, so we harvested the SAMs of two additional individuals, froze, and extracted using the Yockteng et al. 2013 protocol. Samples with an RNA integrity (RIN) score > 7 proceeded directly to library prep. The KAPA Stranded mRNA-Seq Kit (Kapa Biosystems, Waltham MA) protocol was used for library prep. Half reactions were used with an input of at least 500 ng of RNA; however, most samples (all but two) had ~1 ug of RNA. RNA fragmentation time depended on RIN score (7< RIN < 8: 4 min; 8 < RIN < 9: 5 min; 9 < RIN: 6 min). Samples were split in half after the second post-ligation clean up (Step 10 in the Kapa protocol) in order to fine-tune the enrichment step. The first half of the samples were amplified with 12 PCR cycles; this proved too low and was increased to 15 cycles for the second half of the samples. Samples were combined and library quality was assessed with a Bioanalyzer 2100 using the DNA 1000 kit. A bead clean-up was performed on libraries showing significant adaptor peaks (Yockteng *et al.*, 2013). Samples were cleaned, multiplexed, and sequenced on a single lane of HiSeq4000 at the California Institute for Quantitative Biosciences (QB3) Vincent J. Coates Genomics Sequencing Lab.

Bioinformatics Data Processing Raw reads were cleaned, processed, and assembled using the Trinity RNA-Seq De novo Assembly pipeline (Grabherr *et al.*, 2011) under the default settings unless otherwise stated in associated scripts. All analyses were run using the Savio supercomputing resource from the Berkeley Research Computing program at UC Berkeley. Reads were cleaned with Trim Galore! (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/), keeping unpaired reads and using a minimum fragment length of 36 base pairs. Data from all tissue types and biological replicates were concatenated in order to generate a consensus transcriptome, assembled de novo from the concatenated data using Trinity. Each sample was compared back to the assembled consensus transcriptome, aligned using Bowtie 2 (Langmead & Salzberg, 2012), and quantified using RSEM (RNASeq by Expectation Maximization; Li & Dewey, 2011). The consensus transcriptome was annotated with a standard Trinotate pipeline (https://trinotate.github.io/), comparing assembled transcripts to SWISS-PROT (Boeckmann *et al.*, 2003), RNAmmer (Lagesen *et al.*, 2007), Pfam (Finn *et al.*, 2014), eggNOG (Powell *et al.*, 2014), KEGG (Tanabe & Kanehisa, 2012), and Gene Ontology (Gene Ontology Consortium, 2004) databases. The concordance of biological replicates was tested by looking for significant differences between the total number of fragments per replicate, by comparing the transcript quantities of all replicates to each other, and by checking the correlations between replicates. Transcripts with less than 10 total counts were discarded for all downstream analyses. Transcript counts were

241 transformed using the variance-stabilized transformation (VST) and all 12 samples were
242 compared using a principal components analysis. All scripts for these analyses are available on
243 GitHub (github.com/cmt2/bomTubers).
244
Data Analysis Differentially expressed isoforms (hereafter referred to as DEGs) between fibrous
(FR) and tuberous (TR) roots were identified with DESeq2 package in R (Love *et al.*, 2014),
extracting the results of comparing tuberous roots to fibrous roots and using a p-adjusted cut-off
(padj, uses a Benjamini-Hochberg correction for false discovery rate) of 0.01 and a log2-fold
change cut-off of 2 to specifically only statistically significant and sufficiently differentially
expressed isoforms for downstream analyses. To identify specific candidate genes, gene families,
or molecular processes that might be involved in the development of underground storage
organs, we surveyed the literature for recent publications on the molecular basis of USO
formation. For each group of genes hypothesized to be involved in USO formation (either gene
families or molecular/ physiological processes), the annotated transcriptome was queried for
transcript annotations matching the associated process or family (See Table S1 for the specific
search terms used). The resulting transcripts were 1) compared using the distribution of the
group's log2-fold change values to the distribution of values from all isoform using a non-
parametric Mann-Whitney test, and 2) tested for the presence of any statistically significant
differentially expressed isoforms. For all targeted candidate genes, the amino acid sequence of
the candidate gene was blasted to the assembled consensus transcriptome (see Table S2 for the
blasted sequences specifications) using an e-value cut-off of 0.01 to assess if the identified
homologs were differentially expressed. All associated scripts for data analysis are available on
GitHub (github.com/cmt2/bomTubers).
264
The evolutionary history of the phosphatidylethanolamine-binding protein (PEBP) gene family
was reconstructed by combining amino acid sequences from an extensive previously published
alignment (Liu *et al.*, 2016) with the addition of sequences specifically implicated in USO
formation in onion and potato or from geophytic taxa such as *Narcissus tazetta* (accession
AFS50164.1), *Tulipa gesneriana* (accessions MG121853, MG121854, and MG121855), *Crocus*
sativa (saffron, accession ACX53295.1), and *Lilium longiflorum* (accessions MG121858,
MG121857, MG121859) (Navarro *et al.*, 2011; Tsaftaris *et al.*, 2012; Lee *et al.*, 2013; Li *et al.*,
2013; Leeggangers *et al.*, 2017) and with copies identified in our transcriptome. Amino acids
were aligned with MAFFT as implemented in AliView v1.18 (Larsson, 2014), using trimAl
v1.4.rev15 (Capella-Gutierrez *et al.*, 2009) with the -gappyout option. The best evolutionary
model was selected with ModelTest-NG v0.1.5 (Darriba *et al.*, 2016) and unrooted gene trees
were reconstructed under a maximum likelihood framework as implemented in IQtree (Nguyen
et al., 2014), run on XSEDE using the CIPRES portal (Miller *et al.*, 2010).
278
279 **Results**
280
Transcriptome Data A total of 359 M paired-end 100 bp reads were recovered from the single
HiSeq 4000 lane for the multiplexed 12 samples, to be made available as SRAs in an NCBI
BioProject prior to publication. The assembled consensus transcriptome consists of 370,672
transcripts, corresponding to 224,661 unigenes, also to be made available on Dryad prior to
publication. The combined data have a GC content of 45.14%, N50 of 1191 bp, median
transcript length of 317 bp, and mean transcript length of 556.95 bp. All four tissue types showed
concordance between the three biological replicates with generally 1:1 ratios of transcript
quantities to each other (see Figures S1 – S4) so no further replicates were discarded and all

differentially expression analyses incorporated data from all three biological replicates. A principal component analysis (PCA) of the VST transcript counts (Figure 2) shows that the first PC axis (45% of the variance in samples) generally explains the variation between tissue types. The shoot tissues (SAM and RHI) cluster separately, while the root tissues (ROO and TUB) cluster together. The underground rhizome samples (RHI) fall out intermediate between the aerial shoot samples (SAM) and the underground root and tuber samples (ROO and TUB) along this axis. The co-clustering of fibrous and tuberous root samples in the PCA indicates that the overwhelming, general pattern of expression between all root samples is similar, especially in contrast to the very distinct shoot samples. This broad pattern of similarity provides an excellent opportunity to identify the particular genes that cause such obvious morphological differences between the tuberous and fibrous roots, given their similar background patterns of expression. The second PC axis (22% of variance) generally explains variance between biological replicates, with Individual A particularly distinct from other individuals.

Differential Expression A total of 271 differentially expressed isoforms (DEGs) were recovered between fibrous and tuberous roots (FR vs. TR) (Table S3). Of these, 226 correspond to regions of the annotated consensus transcriptome with functional annotations. The most common gene ontology (GO) terms associated with these DEGs are presented in Table 1. The top cellular components include the nucleus, the cell membrane and cell wall, and plastids; the top molecular functions include kinase activity and binding to ATP, nucleic acids, proteins, and various metals; and the top biological functions include cell wall organization, morphological development, environmental sensing, and protein transport. Of the 271 DEGs, 126 (46.5%) were over-expressed in tuberous roots while the remaining 145 (53.5%) were under-expressed. All top ten most differentially expressed isoforms (the ten DEGs with the highest absolute value log2-fold change values between fibrous and tuberous roots) were functionally annotated. These ten DEGs are implicated in various cellular and biological processes (Table 2). All but nine of these top ten DEGs are overexpressed in tuberous roots and are generally implicated in nucleotide and ATP binding, cell wall modification, root morphogenesis, and carbohydrate and fatty acid biosynthesis. The top most differentially expressed isoform with a 40.25 log2-fold change value, TRINITY_DN116220_c0_g1_i4, is a Zinc finger CCCH domain-containing protein 55, a possible transcription factor of unknown function. Other notable top DEGs include TRINITY_DN128685_c1_g3_i4, callose synthase 3, which regulates cell shape, TRINITY_DN121298_c2_g2_i5, a heat shock protein, TRINITY_DN127064_c0_g3_i1, an LRR receptor-like serine implicated in lateral root morphogenesis, and TRINITY_DN121430_c10_g2_i, a carbohydrate metabolism protein. The tenth most differentially expressed DEG, under-expressed in tuberous roots, is implicated in abscisic acid signaling.

Parallel Processes Across Taxa Eleven gene groups — either gene families, physiological signaling pathways, or biosynthesis pathways — were identified that have been implicated in USO formation by expression of functional analyses across various geophytic organisms: abscisic acid response genes, calcium-dependent protein kinases (CDPK), expansins, lignin biosynthesis, MADS-Box genes, starch biosynthesis, auxin response genes, cytokinin response genes, 14-3-3 genes, gibberellin response genes, and KNOX genes (See Figure 3). For each group, putative homologs were located in *Bomarea multiflora* and their expression patterns were analyzed. Of these 11 gene groups, the log2-fold change values of six are distributed significantly differently than the overall distribution of log2-fold change values for all isoforms; in two cases (starch biosynthesis and MADS-Box genes) the expression levels overall are

significantly greater than expected (generally over-expressed in TR compared to FR) and in the remaining four cases (abscisic acid, CDPK, expansins, and lignin biosynthesis) the groups' log2-fold change values are under-expressed. Fifteen individual DEGs were identified in these gene groups (Table 3); interestingly, there seems to be no generalizable relationships between the significance and directionality of a particular group's distribution with the presence and directionality of log2-fold change values for individual DEGs. For example, the expression distribution of gibberellin genes does not deviate significantly from the global pool of isoforms, but there is one significantly under-expressed DEG in the gibberellin group; similarly, the CDPK genes are under-expressed as a group, but the only CDPK-related significant DEG is over-expressed (Table 3).

Five specific candidate genes were identified from the literature: qRT9 has been implicated in root thickening in rice (Li *et al.*, 2015); IDD5 and WOX4 are implicated in starch biosynthesis and TR formation, respectively, in Convolvulaceae (Eserman *et al.*, 2018); sulfite reductase is associated with TR formation in *Manihot esculenta* (Sojikul *et al.*, 2010); and FLOWERING LOCUS T (FT) has been implicated in signaling the timing of USO formation in a variety of taxa, notably *Allium cepa* and *Solanum tuberosum* (Navarro *et al.*, 2011; Hannapel *et al.*, 2017). We recover between nine and 63 putative homologs of these candidates using a BLAST E-value cut-off of 0.01 (Table 4), but only one putative homolog is significantly differentially expressed (padj < 0.01): a putative FT homolog (TRINITY_DN129076_c1_g1_i1), further investigated in PEBP Gene Family Evolution (below). One putative qRT9 homolog is marginally significant (padj = 0.050), and the E-value from the BLAST result to this isoform was 0.09. Given these marginal significance values, it is likely the result is spurious and we do not follow up with further analysis.

PEBP Gene Family Evolution Thirty-seven *Bomarea* isoforms were identified as putative FLOWERING LOCUS T (FT) homologs. The longest isoform per gene was selected to include in an alignment of phosphatidylethanolamine-binding protein (PEBP) amino acid sequences. Coding sequences from the *Bomarea multiflora* transcriptome were translated to amino acid sequences using TransDecoder v5.5.0 (Haas *et al.*, 2013), removing isoforms that failed to align properly; ultimately, we include six sequences, including the significantly differentially expressed copy. Using the JTT+G4 amino acid substitution models (Jones *et al.*, 1992) we recover three major clusters in our unrooted gene tree, all with strong bootstrap support (Figure 4a); these correspond to the FT cluster, TERMINAL FLOWER 1 (TFL1) cluster, and MOTHER OF FT AND TFL1 (MFT) cluster recovered in previous analyses (Liu *et al.*, 2016). Three of the six *Bomarea multiflora* isoforms fall out with FT genes and three fall out with TFL1 genes. The *Bomarea* DEG homolog is highly supported in the TFL1 cluster with sequences from other monocot taxa (Figure 4b). FT homologs from *Allium cepa* and *Solanum tuberosum* that have been functionally implicated in stem tuber and bulb formation, respectively, are in the FT cluster but do not cluster together; rather all USO-implicated PEBP genes are more closely related to non-USO copies than to each other.

Discussion

How to Make a Tuberous Root The top 10 most highly represented biological processes of the 271 Differentially expressed isoforms (DEGs) include cell wall organization, responses to environmental signals, growth and development, and carbohydrate biosynthesis (Table 1). Together, these processes describe the various components of development by which the plant

385   modifies fibrous roots into tuberous roots: 1) *how* expansion occurs, 2) *when* tuberization is
386   triggered, and 3) *what* the tuberous roots store.
387
Root expansion likely occurs due to cellular expansion and primary thickening growth in *B.*
multiflora. Due to the absence of a vascular cambium, secondary growth is not likely to be
involved, despite the prevalence of this mechanism in other taxa such as sweet potato (Noh *et al.*,
2010; Eserman *et al.*, 2018) and cassava (Melis & van Staden, 1985). DEGs with cell wall
functional annotations, the most common GO biological process, likely contribute to root
enlargement through permitting cellular expansion. For example, pectinesterase
TRINITY_DN122210_c6_g1_i1 (log2-fold change = 21.91; padj = $1.22E^{-8}$) modifies pectin in
cell walls leading to cell wall softening, as demonstrated, for example, in Arabidopsis
(Braybrook & Peaucelle, 2013). Interestingly, expansins were not over-expressed in *B.*
multiflora, though this is the mechanism by which cell expansion occurs in other taxa (see
Expansins discussion below). The DEGs functionally annotated as contributing to cell division
(one of the top GO biological processes in the DEG dataset) may contribute to root enlargement
through increased cellular proliferation at the growing tip of the tuberous root (primary
thickening growth).
402
Several of the top GO biological processes are responses to environmental stimuli, including
cold, water deprivation, flowering, and defense, which may trigger tuberization. Flowering
development genes in particular may be co-opted for tuber formation, a hypothesis discussed in
more detail below (PEBP gene family evolution). Tuberization signaling may also be mediated
by callose production, influencing symplastic signaling pathways through plasmodesmata
modification. Callose synthase 3 is one of the most highly differentially expressed DEGs
(TRINITY_DN128685_c1_g3_i4, Table 2). Callose is a much less common component of cell
walls than is cellulose (Schneider *et al.*, 2016), but it is often implicated in specialized cell walls
and in root-specific expression (Vatén *et al.*, 2011; Benitez-Alfonso *et al.*, 2013). Callose
synthase has been implicated in the development of other unique root-based structures such as
root nodules (Gaudioso-Pedraza *et al.*, 2018) and mutations in callose synthase 3 affect root
morphology (Vatén *et al.*, 2011), suggesting that callose synthase 3 plays an integral role in
triggering tuberous roots development in *B. multiflora* through symplastic signaling pathways.
Callose signaling-induced USO formation has not previously been reported and be unique to *B.*
multiflora or to monocotylendous taxa.
418
Finally, starch is thought to be the primary nutrient reserve in *Bomarea* tubers (Kubitzki, 1998).
Many previous studies have found evidence of overexpression of carbohydrate and starch
synthesis molecules in USOs (for example in sweet potato; Eserman *et al.*, 2018)). Differentially
expressed isoforms implicated in the carbohydrate metabolic process support the presence of
active starch synthesis in our data. One of the most differentially expressed isoforms is a
homolog of sucrose non-fermenting 4-like protein (Table 2, TRINITY_DN121430_c10_g2_i1)
and participates in carbohydrate biosynthesis, demonstrating that *B. multiflora* tubers were
actively synthesizing starch when harvested. Additionally, genes implicated in defense response,
such as TRINITY_DN127064_c0_g3_i1 (LRR receptor-like serine/ threonine-protein kinase
HSL2, Table 2) may be differentially expressed in tuberous roots to protect starch reserves
against potential predation by belowground herbivores.
430
431   **Similarities in Molecular Mechanisms of USO Formation** We identify six molecular
432   processes, previously implicated in USO formation in other taxa, which are either over- or under-

433   expressed in the tuberous roots of *Bomarea multiflora* (Figure 3). These processes show parallel
434   function across deeply divergent evolutionary distances and in distinct plant structures.
435
Abscisic Acid (ABA) has been shown to increase initially and then decrease during USO
formation in the rhizomes of *Nelumbo nucifera* (Yang *et al.*, 2015), the tuberous roots of sweet
potato *Ipomoea batatas* (Dong *et al.*, 2019), and the stem tubers of potato *Solanum tuberosum*
(Xu *et al.*, 1998). We find that ABA signaling genes are under-expressed in tuberous roots
(Figure 3), perhaps indicating that our samples were in a later developmental stage, characterized
by lower ABA levels. Experimental manipulation of *B. multiflora* tuberous roots and
developmental time series would be needed to confirm the role of ABA in tuberous root
formation. An ABA signaling DEG (Table 3) is one of the most differentially expressed isoforms
(TRINITY_DN122787_c0_g1_i1: Protein IQ-DOMAIN 32, Table 2). This isoform and other
ABA genes may signal the cessation of continued elongation in monocot tuberous roots, similar
to their role in deeply divergent taxa and non-homologous USOs.
447
Calcium-dependent Protein Kinases (CDPKs) play an integral role in tuber formation in cassava
Manihot esculenta (Sojikul *et al.*, 2010). Sujikul et al. (2010) propose that CDPKs may signal
the initiation of tuberous root development, similarly to the process described in stem tuber
formation in potato (Raíces *et al.*, 2003) and rhizome development in *Nelumbo nucifera* (Cheng
et al., 2013b). In these studies, CDPKs are over-expressed and signal the initiation of USO
development. Our results show that on average CDPKs as a group are under-expressed in the
tuberous roots of *B. multiflora.* However, the only significant CDPK DEG
(TRINITY_DN124121_c3_g1_i11: Calcium-dependent protein kinase 2) is over-expressed,
suggesting that CDPK 2 expression plays a role in initiating tuberous root development and that
CDPK involvement in tuberous root formation is ubiquitous in all studied taxa.
458
Expansins are cell wall modifying genes known to loosen cell walls in organ formation (Dolan
& Davies, 2004; Humphrey *et al.*, 2007; Braybrook & Peaucelle, 2013). Their involvement in
USO formation has been documented in the tuberous roots of cassava (Sojikul *et al.*, 2015), and
Callerya speciosa (Xu *et al.*, 2016) the rhizomes of *Nelumbo nucifera* (Cheng *et al.*, 2013b), the
tuberous roots of various Convolvulaceae (Eserman *et al.*, 2018), and the stem tubers of potato
(Jung *et al.*). As a group expansins are under-expressed in tuberous compared to fibrous roots,
but none are statistically significant, so it seems unlikely that expansins play an important role in
tuberous root formation in *Bomarea multiflora*. It is possible that expansin involvement in USO
formation is unique to eudicots.
468
Lignin biosynthesis genes are under-expressed in several geophytic taxa with tuberous roots,
including cassava (Sojikul *et al.*, 2015), wild sweet potato (*Ipomoea trifida*; (Li *et al.*, 2019), and
Callerya speciosa (Xu *et al.*, 2016). Similarly, we find that lignin biosynthesis overall is under-
expressed in tuberous compared to fibrous roots, and one isoform in particular is significantly
under-expressed: TRINITY_DN125451_c5_g1_i3: Cinnamoyl-CoA reductase-like SNL6 (Table
3). This gene has been found to significantly decrease lignin content without otherwise affecting
development in tobacco (Chabannes *et al.*, 2001). Decreased lignin in tuberous roots may further
allow for cell expansion and permit lateral swelling of tuberous roots during development.
477
MADS-Box genes are implicated in USO formation in the tuberous roots of wild sweet potato
(*Ipomoea trifida*; (Li *et al.*, 2019), and sweet potato (*Ipomoea batatas*; (Noh *et al.*, 2010; Dong *et*
al., 2019), the rhizomes of *Nelumbo nucifera* (Cheng *et al.*, 2013b), and the corms of *Sagittaria*

trifolia (Cheng *et al.*, 2013a), indicating widespread parallel use of MADS-Box genes in the
formation of USOs. Similarly, we find that MADS-Box genes overall, and one DEG in
particular, are over-expressed in *Bomarea* tuberous roots. MADS-Box genes are implicated
widely as important transcription factors regulating plant development (Buylla *et al.*, 2000). It is
thus unsurprising that MADS-Box genes are regularly implicated in USO formation. It remains
unclear if the MADS-Box genes identified in the aforementioned studies represent independent
neofunctionalizations of MADS-Box genes from other aspects of plant development, or if they
form a clade of USO-specific copies.
489
Starch biosynthesis genes are very commonly identified in the formation of USOs, including in
cassava (Sojikul *et al.*, 2010; 2015), *Nelumbo nucifera* (Cheng *et al.*, 2013b; Yang *et al.*, 2015),
wild and domesticated sweet potatoes (Eserman *et al.*, 2018; Li *et al.*, 2019; Dong *et al.*, 2019),
and potato (Xu *et al.*, 1998). Since starch is so ubiquitous in USOs, this is unsurprising. We also
find starch isoforms overall to be over-expressed in *Bomarea* tuberous roots, and three genes in
particular are significantly over-expressed (see Table 3).
496
The other molecular processes we tested failed to show group-level differences from the global
distribution of expression levels. However, the presence of DEGs in some of these groups
indicates that the phytohormones in particular may play a role in tuberous root formation. One
gibberellin response isoform that is significantly under-expressed in tuberous roots, which aligns
with previous research suggesting that decreased gibberellin concentrations in roots can lead to
root enlargement (Tanimoto, 2012) and tuber formation (Xu *et al.*, 2016; Li *et al.*, 2019; Dong *et*
al., 2019). The lack of significant auxin-related isoforms as differentially expressed is surprising,
as auxin has been implicated in USO formation in several previous studies (Noh *et al.*, 2010;
Cheng *et al.*, 2013b; Sojikul *et al.*, 2015; Yang *et al.*, 2015; Xu *et al.*, 2016; Hannapel *et al.*,
2017; Li *et al.*, 2019; Dong *et al.*, 2019; Kolachevskaya *et al.*, 2019), but it is possible this
function is not utilized in the tuberous root formation for monocot taxa.
508
PEBP and FT-Like Gene Evolution in Geophytic Taxa Gene tree analysis of PEBPs indicates
that FT and TFL1 genes have been independently co-opted several times in USO formation in
diverse angiosperms (including monocots and eudicots) and in diverse USO morphologies
(including tuberous roots, bulbs, and stem tubers). Furthermore, the presence of TFL1 and FT
homologs in gymnosperms and other non-flowering plants (Liu *et al.*, 2016) suggests that the
origin of these genes predates the evolution of flower as a reproductive structure. Instead, it
seems likely that these genes originally evolved as environmental signaling genes with wider
involvement in triggering the seasonality of various aspects of plant development. Subsequent
specialization of these genes in the timing of shifts to reproductive development as well as USO
development occurred. Given our results, in seems likely that USO-specialized FT and TFL1
genes arose at least four times independently, indicating broadly parallel molecular evolution
underlies the convergent morphological evolution of USOs. Additional USO-specific PEBP
genes would shed more light on this pattern, but the dearth of studies on the molecular basis of
USO development impedes such analysis. With increased sampling, follow-up studies could
identify unique patterns of convergent molecular evolution on the USO-specific FT genes. Do
these copies share independently derived subsequences or motifs that could reflect or cause
shared function?
526
527   In *Bomarea multiflora*, TFL1 involvement in USO formation is surprising, as neotropical cloud
528   forests are generally considered relatively aseasonal environments. However, *B. multiflora*

shows two annual peaks in flowering corresponding to two peaks in annual rainfall (Ortiz & Idárraga Piedrahita, 2009), suggesting that important aspects of the plant's phenology are tied to seasonal fluctuations. While no previous studies have looked at the timing of tuber production in *Bomarea*, it is possible that tuber development is also tied to seasonal cues.

Conclusion We provide the first evidence of the molecular mechanisms of tuberous root formation in a monocotyledonous taxon, filling a key gap in understanding the commonalities of storage organ formation across taxa. We demonstrate that many molecular processes are shared across geophytic taxa, suggesting that deep parallel evolution at the molecular level underlies the convergent evolution of an adaptive trait. In particular, we demonstrate that PEBP genes implicated in underground storage organ formation have been recruited multiple times across the gene tree, demonstrating that repeated morphological convergence is matched by repeated molecular convergence. These findings suggest further avenues for research on the molecular mechanisms of how plants retreat underground and evolve strategies enabling adaptation to environmental stresses. More molecular studies on diverse, non-model taxa and more thorough sampling of underground morphological diversity will enhance our understanding of the full extent of these convergences and add to our general understanding of the molecular basis for adaptive, convergent traits.

Acknowledgements
We thank Lydia Smith (Evolutionary Genetics Lab, UC Berkeley) for training and sharing her expertise on RNA-Seq, NSF GRFP, SSB, ASPT, Pacific Bulb Society, UC Berkeley's Integrative Biology Department, and the Tinker Foundation for support to CMT, and UC Berkeley CNR and the University and Jepson Herbarium for supporting sequencing costs.

Author Contribution
FA provided materials for the experiment, CMT and CDS designed the experiment, CMT and JMG executed the experiment, CMT, JMG, and CJR analyzed the data, and CMT, JMG, CJR and CDS contributed to writing the manuscript.

References

Al-Tardeh S, Sawidis T, Diannelidis B-E, Stylianos Delivopoulos. **2008**. Water content and reserve allocation patterns within the bulb of the perennial geophyte red squill (Liliaceae) in relation to the Mediterranean climate. *Botany* **86**: 291–299.

Benitez-Alfonso Y, Faulkner C, Pendle A, Miyashima S, Helariutta Y, Maule A. **2013**. Symplastic Intercellular Connectivity Regulates Lateral Root Patterning. *Developmental Cell* **26**: 136–147.

Boeckmann B, Bairoch A, Apweiler R, Blatter M-C, Estreicher A, Gasteiger E, Martin MJ, Michoud K, O'Donovan C, Phan I, *et al.* **2003**. The SWISS-PROT protein knowledgebase and its supplement TrEMBL in 2003. *Nucleic acids research* **31**: 365–370.

Braybrook SA, Peaucelle A. **2013**. Mechano-chemical aspects of organ formation in Arabidopsis thaliana: the relationship between auxin and pectin. *PLOS ONE* **8**: e57813.

Buylla ERA, Liljegren SJ, Pelaz S, Gold SE, Burgeff C, Ditta GS, Silva FV, Yanofsky MF. **2000**. MADS-box gene evolution beyond flowers: expression in pollen, endosperm, guard cells, roots and trichomes. *The Plant Journal* **24**: 457–466.

Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. **2009**. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**: 1972–1973.

Chabannes M, Barakate A, Lapierre C, Marita JM, Ralph J, Pean M, Danoun S, Halpin C, Pettenati JG, Boudet AM. **2001**. Strong decrease in lignin content without significant alteration of plant development is induced by simultaneous down-regulation of cinnamoyl CoA reductase (CCR) and cinnamyl alcohol dehydrogenase (CAD) in tobacco plants. *The Plant Journal* **28**: 257–270.

Chaweewan Y, Taylor N. **2015**. Anatomical Assessment of Root Formation and Tuberization in Cassava (Manihot esculenta Crantz). *Tropical Plant Biology* **8**: 1–8.

Cheng L, Li S, Xu X, Hussain J, Yin J, Zhang Y, Li L, Chen X. **2013a**. Identification of differentially expressed genes relevant to corm formation in *Sagittaria trifolia*. *PLOS ONE* **8**: e54573.

Cheng L, Li S, Yin J, Li L, Chen X. **2013b**. Genome-wide analysis of differentially expressed genes relevant to rhizome formation in lotus root (*Nelumbo nucifera* Gaertn). *PLOS ONE* **8**: e67116.

Cuéllar-Martínez M, Sosa V. **2016**. Diversity patterns of monocotiledonous geophytes in Mexico. *Botanical Sciences* **94**: 699–13.

Dafni A, Cohen D, Noy-Mier I. **1981a**. Life-cycle variation in geophytes. *Annals of the Missouri Botanical Garden* **68**: 652–660.

Dafni A, Shmida A, Avishai M. **1981b**. Leafless autumnal-flowering geophytes in the Mediterranean region - phytogeographical, ecological and evolutionary aspects. *Plant Systematics and Evolution* **137**: 181–193.

Darriba D, Weiß M, Stamatakis A. **2016**. Prediction of missing sequences and branch lengths in phylogenomic data. *Bioinformatics* **32**: 1331–1337.

Dolan L, Davies J. **2004**. Cell expansion in roots. *Current Opinion in Plant Biology* **7**: 33–39.

Dong T, Zhu M, Yu J, Han R, Tang C, Xu T, Liu J, Li Z. **2019**. RNA-Seq and iTRAQ reveal multiple pathways involved in storage root formation and development in sweet potato (*Ipomoea batatas* L.). *BMC Plant Biology* **19**: 136–16.

Ekblom R, Galindo J. **2011**. Applications of next generation sequencing in molecular ecology of non-model organisms. *Heredity* **107**: 1–15.

Eserman LA, Jarret RL, Leebens-Mack JH. **2018**. Parallel evolution of storage roots in morning glories (Convolvulaceae). *BMC Plant Biology* **18**: 1–11.

Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, Heger A,
Hetherington K, Holm L, Mistry J, *et al.* **2014**. Pfam: the protein families database. *Nucleic*
acids research **42**: D222–30.

Gaudioso-Pedraza R, Beck M, Frances L, Kirk P, Ripodas C, Niebel A, Oldroyd GED,
Benitez-Alfonso Y, de Carvalho-Niebel F. **2018**. Callose-Regulated Symplastic
Communication Coordinates Symbiotic Root Nodule Development. *Current Biology* **28**: 3562–
3577.e6.

Gene Ontology Consortium. **2004**. The Gene Ontology (GO) database and informatics
resource. *Nucleic acids research* **32**: 258D–261.

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,
Raychowdhury R, Zeng Q, *et al.* **2011**. Full-length transcriptome assembly from RNA-Seq data
without a reference genome. *Nature Biotechnology* **29**: 644–652.

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB,
Eccles D, Li B, Lieber M, *et al.* **2013**. De novo transcript sequence reconstruction from RNA-
seq using the Trinity platform for reference generation and analysis. *Nature Protocols* **8**: 1494–
1512.

Hannapel DJ, Sharma P, Lin T, Banerjee AK. **2017**. The Multiple Signals That Control Tuber
Formation. *Plant Physiology* **174**: 845–856.

Hearn DJ. **2009**. Descriptive anatomy and evolutionary patterns of anatomical diversification in
Adenia (Passifloraceae). *Aliso* **27**: 13–38.

Hoffmann AJ, Liberona F, Hoffmann AE. **1998**. *Distribution and ecology of geophytes in*
Chile. Conservation threats to geophytes in mediterranean-type regions. Berlin, Germany:
Springer-Verlag.

Hofreiter A. **2008**. A Revision of *Bomarea* Subgenus *Bomarea* s.str. Section *Multiflorae*
(Alstroemeriaceae). *Systematic Botany* **33**: 661–684.

Howard CC, Folk RA, Beaulieu JM, Cellinese N. **2019a**. The monocotyledonous
underground: global climatic and phylogenetic patterns of geophyte diversity. *American Journal*
of Botany **106**: 850–863.

Howard CC, Landis JB, Beaulieu JM, Cellinese N. **2019b**. Geophytism in monocots leads to
higher rates of diversification. *New Phytologist*: 1-10.

Humphrey TV, Bonetta DT, Goring DR. **2007**. Sentinels at the wall: cell wall receptors and
sensors. *New Phytologist* **176**: 7–21.

Jones DT, Taylor WR, Thornton JM. **1992**. The rapid generation of mutation data matrices
from protein sequences. *Computer applications in the biosciences: CABIOS* **8**: 275–282.

Jung J, O'Donoghue EM, Dijkwel PP, Brummell DA. Expression of multiple expansin genes
is associated with cell expansion in potato organs. *Plant Science* **179**: 77–85.

Kolachevskaya OO, Lomin SN, Arkhipov DV, Romanov GA. **2019**. Auxins in potato: molecular aspects and emerging roles in tuber formation and stress resistance. *Plant cell reports* **38**: 681–698.

Kubitzki K. **1998.** Vol III Monocotyledons: Lilianae (except Orchidaceae). *The families and genera of vascular plants*. Berlin, Germany: Springer-Verlag.

Lagesen K, Hallin P, Rødland EA, Stærfeldt H-H, Rognes T, Ussery DW. **2007**. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic acids research* **35**: 3100–3108.

Langmead B, Salzberg SL. **2012**. Fast gapped-read alignment with Bowtie 2. *Nature methods* **9**: 357–359.

Larsson A. **2014**. AliView: a fast and lightweight alignment viewer and editor for large datasets. *Bioinformatics* **30**: 3276–3278.

Lee R, Baldwin S, Kenel F, McCallum J, Macknight R. **2013**. FLOWERING LOCUS T genes control onion bulb formation and flowering. *Nature Communications* **4**: 2884.

Leeggangers HACF, Nijveen H, Bigas JN, Hilhorst HWM, Immink RGH. **2017**. Molecular Regulation of Temperature-Dependent Floral Induction in Tulipa gesneriana. *Plant Physiology* **173**: 1904–1919.

Li B, Dewey CN. **2011**. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC bioinformatics* **12**: 323.

Li J, Han Y, Liu L, Chen Y, Du Y, Zhang J, Sun H, Zhao Q. **2015**. qRT9, a quantitative trait locus controlling root thickness and root length in upland rice. *Journal of experimental botany* **66**: 2723–2732.

Li M, Yang S, Xu W, Pu Z, Feng J, Wang Z, Zhang C, Peng M, Du C, Lin F, *et al.* 2019. The wild sweetpotato (*Ipomoea trifida*) genome provides insights into storage root development. *BMC Plant Biology* **19**: 119–17.

Li X-F, Jia L-Y, Xu J, Deng X-J, Wang Y, Zhang W, Zhang X-P, Fang Q, Zhang D-M, Sun Y, *et al.* 2013. FT-Like NFT1 Gene May Play a Role in Flower Transition Induced by Heat Accumulation in Narcissus tazetta var. chinensis. *Plant & cell physiology* **54**: 270–281.

Liu Y-Y, Yang K-Z, Wei X-X, Wang X-Q. **2016**. Revisiting the phosphatidylethanolamine-binding protein (PEBP) gene family reveals cryptic FLOWERING LOCUS Tgene homologs in gymnosperms and sheds new light on functional evolution. *New Phytologist* **212**: 730–744.

Love MI, Huber W, Anders S. **2014**. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**: 31–21.

Melis RJ, van Staden J. **1985**. Tuberization in Cassava (*Manihot esculenta*): Cytokinin and Abscisic Acid Activity in Tuberous Roots. *Journal of plant physiology* **118**: 357–366.

Miller MA, Pfeiffer W, Schwartz T. **2010**. Creating the CIPRES Science Gateway for
inference of large phylogenetic trees. 2010 gateway computing environments workshop (GCE).
Ieee: 1-10.

Navarro C, Abelenda JA, Cruz-Oró E, Cuéllar CA, Tamaki S, Silva J, Shimamoto K, Prat
S. **2011**. Control of flowering and storage organ formation in potato by FLOWERING LOCUS
T. *Nature* **478**: 119–122.

Nguyen L-T, Schmidt HA, Haeseler von A, Minh BQ. **2014**. IQ-TREE: A Fast and Effective
Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and*
Evolution **32**: 268–274.

Niklas KJ. **2005**. Modelling below-and above-ground biomass for non-woody and woody plants.
Annals of Botany **95**: 315–321.

Noh SA, Lee H-S, Huh EJ, Huh GH, Paek K-H, Shin JS, Bae JM. **2010**. SRD1 is involved in
the auxin-mediated initial thickening growth of storage root by enhancing proliferation of
metaxylem and cambium cells in sweetpotato (Ipomoea batatas). *Journal of experimental botany*
61: 1337–1349.

Oppenheim SJ, Baker RH, Simon S, DeSalle R. **2015**. We can't all be supermodels: the value
of comparative transcriptomics to the study of non-model insects. *Insect molecular biology* **24**:
139–154.

Ortiz RDC, Idárraga Piedrahita Á. **2009**. Catálogo de las Plantas Vasculares del
Departamento de Antioquia. http://www.tropicos.org/Project/Catalogo-de-Antioquia.

Ott JP, Klimešová J, Hartnett DC. **2019**. The ecology and significance of below-ground bud
banks in plants. *Annals of Botany* **123**: 1099–1118.

Parsons RF. **2000**. Monocotyledonous geophytes: comparison of California with Victoria,
Australia. *Australian Journal of Botany* **48**: 39–43.

Pate JS, Dixon KW. **1982**. *Tuberous, cormous and bulbous plants: biology of an adaptive*
strategy in Western Australia. Nedlands, Australia: University of Western Australia Press.

Powell S, Forslund K, Szklarczyk D, Trachana K, Roth A, Huerta-Cepas J, Gabaldón T,
Rattei T, Creevey C, Kuhn M, *et al*. **2014**. eggNOG v4.0: nested orthology inference across
3686 organisms. *Nucleic acids research* **42**: D231–9.

Proches S, Cowling RM, Goldblatt P, Manning JC, Snijman DA. **2006**. An overview of the
Cape geophytes. *Biological Journal of the Linnean Society* **87**: 27–43.

Raíces M, Gargantini PR, Chinchilla D, Crespi M, Téllez-Iñón MT, Ulloa RM. **2003**.
Regulation of CDPK isoforms during tuber development. *Plant molecular biology* **52**: 1011–
1024.

Raunkiaer C. **1934**. *The life forms of plants and statistical plant geography; being the collected*
papers of C. Raunkiaer.

Rundel PW. **1996**. Monocotylendous geophytes in the California flora. *Madroño* **43**: 355–368.

Schneider R, Hanak T, Persson S, Voigt CA. **2016**. Cellulose and callose synthesis and
organization in focus, what's new? *Current Opinion in Plant Biology* **34**: 9–16.

Sojikul P, Kongsawadworakul P, Viboonjun U, Thaiprasit J, Intawong B, Narangajavana
J, Svasti MRJ. **2010**. AFLP-based transcript profiling for cassava genome-wide expression
analysis in the onset of storage root formation. *Physiologia Plantarum* **140**: 189–198.

Sojikul P, Saithong T, Kalapanulak S, Pisuttinusart N, Limsirichaikul S, Tanaka M,
Utsumi Y, Sakurai T, Seki M, Narangajavana J. **2015**. Genome-wide analysis reveals
phytohormone action during cassava storage root initiation. *Plant molecular biology* **88**: 531–
543.

Sosa V, Loera I. **2017**. Influence of current climate, historical climate stability and topography
on species richness and endemism in Mesoamerican geophyte plants. *PeerJ* **5**: e3932–20.

Tanabe M, Kanehisa M. **2012**. Using the KEGG database resource. *Current protocols in*
bioinformatics **Chapter 1**: Unit1.12–1.12.43.

Tang JR, Lu YC, Gao ZJ, Song WL, Wei KH, Zhao Y, Tang QY, Li XJ, Chen JW, Zhang
GH, *et al.* 2019. Comparative transcriptome analysis reveals a gene expression profile that
contributes to rhizome swelling in *Panax japonicus* var. *major. Plant Biosystems*.

Tanimoto E. **2012**. Tall or short? Slender or thick? A plant strategy for regulating elongation
growth of roots by low concentrations of gibberellin. *Annals of Botany* **110**: 373–381.

Thiers B. Index Herbariorum: a global directory of public herbaria and associated staff.
http://sweetgum.nybg.org/science/ih/

Tsaftaris A, Pasentsis K, Kalivas A, Michailidou S, Madesis P, Argiriou A. **2012**. Isolation
of a CENTRORADIALIS/TERMINAL FLOWER1 homolog in saffron (Crocus sativus L.):
characterization and expression analysis. *Molecular Biology Reports* **39**: 7899–7910.

Vatén A, Dettmer J, Wu S, Stierhof Y-D, Miyashima S, Yadav SR, Roberts CJ, Campilho
A, Bulone V, Lichtenberger R, *et al.* 2011. Callose Biosynthesis Regulates Symplastic
Trafficking during Root Development. *Developmental Cell* **21**: 1144–1155.

Veselý P, Bureš P, Šmarda P, Pavlíček T. **2011**. Genome size and DNA base composition of
geophytes: the mirror of phenology and ecology? *Annals of Botany* **109**: 65–75.

Wang Y, Liu L, Song S, Li Y, Shen L, Yu H. **2017**. DOFT and DOFTIP1 affect reproductive
development in the orchid *Dendrobium* Chao Praya Smile. *Journal of experimental botany* **68**:
5759–5772.

Wilson CA. **2006**. Patterns in evolution in characters that define Iris subgenera and sections.
Aliso **22**: 425–433.

Xu L, Wang J, Lei M, Li L, Fu Y, Wang Z, Ao M, Li Z. **2016**. Transcriptome analysis of
storage roots and fibrous roots of the traditional medicinal herb *Callerya speciosa* (Champ.)
ScHot (X-Q Li, Ed.). *PLOS ONE* **11**: e0160338–20.

Xu X, van Lammeren AA, Vermeer E, Vreugdenhil D. **1998**. The role of gibberellin, abscisic
acid, and sucrose in the regulation of potato tuber formation in vitro. *Plant Physiology* **117**: 575–
584.

Yang M, Zhu L, Pan C, Xu L, Liu Y, Ke W, Yang P. **2015**. Transcriptomic analysis of the
regulation of rhizome formation in temperate and tropical lotus (*Nelumbo nucifera*). *Nature*
Publishing Group **5**: 13059.

Yockteng R, Almeida AMR, Yee S, Andre T, Hill C, Specht CD. **2013**. A Method for
Extracting High-Quality RNA from Diverse Plants for Next-Generation Sequencing and Gene
Expression Analyses. *Applications in Plant Sciences* **1**: 1300070–6.

New Zealand Ministry for Public Industries. National Pests Plant Accord.
https://www.mpi.govt.nz/protection-and-response/long-term-pest-management/national-pest-
plant-accord/. Accessed 1 October 2019.
764

765 **Figures and Tables**
766
767 **Figure 1:** Sampling scheme of tissue types. *Bomarea multiflora* has modified underground stems
768 (rhizomes) and modified roots (tuberous roots). Aerial shoots and roots develop from the
769 underground rhizome. We extracted RNA from the aerial shoot meristem, rhizome meristem,
770 fibrous root tip, and tuberous root tip of three *B. multiflora* individuals.

771
772

Figure 2: Principal component analysis of all samples. We performed a principal component analysis of variance stabling transformed (VST) transcript counts from all biological replicates of all tissue types. Point color indicates the tissue type of the sample and the letter corresponds to individual plants sampled. The first PC axis (PC1) explains 45% of the variance in the data, corresponding to variance due to tissue type. The second PC axis (PC2) explains 22% of the variance, corresponding to variance due to individual plant differences and other factors.

Figure 3: Differential expression of candidate gene groups. We identify isoforms corresponding
to specific pathways and gene families (gene groups) and categorize the log2-fold change of
expression between tuberous and fibrous roots of those groups. Boxplots correspond to the log2-
fold change value for the gene groups and grey points correspond to individual isoforms. Groups
with distributions that are significantly different from the entire dataset (All Genes) are labeled in
red with asterisks on the X-axis. Isoforms are significantly differentially expressed (padj > 0.01)
are labeled as red asterisks within the scatter plots.

Figure 4: Evolution of PEBP genes: (a) Unrooted gene tree of 540 PEBP gene copies from across land plants. Stars indicate all included copies from *Bomarea multiflora*; the red star corresponds to the significantly differentially expressed isoform TRINITY_DN129076_c1_g1_i1). Red and arrows indicate PEBP copies that have been implicated in USO formation in other taxa (*Allium cepa* and *Solanum tuberosum*). The major clusters correspond to the FLOWERING LOCUS T (FT, in purple), TERMINAL FLOWER 1 (TFL1, in yellow), and the MOTHER OF FLOWERING LOCUS T AND TERMINAL FLOWER 1 (MFT, in red) gene groups and are labeled with high bootstrap support. (b) Detailed view of the cluster indicated by a dashed circle in (a), including monocot-specific copies of TERMINAL FLOWER 1 genes. Line thickness corresponds to bootstrap support.

829 **Table 1:** Top ten GO annotations for all differentially expressed isoforms.

Cellular Components		Molecular Functions		Biological Processes	
Annotation	Frequency	Annotation	Frequency	Annotation	Frequency
nucleus	0.129	ATP binding	0.089	cell wall organization	0.020
integral component of membrane	0.098	metal ion binding	0.069	flower development	0.020
plasma membrane	0.078	DNA binding	0.044	multicellular organism development	0.016
cytoplasm	0.067	protein serine/threonine kinase activity	0.027	protein transport	0.016
cytosol	0.053	DNA-binding transcription factor activity	0.024	response to water deprivation	0.016
chloroplast	0.038	RNA binding	0.020	regulation of transcription, DNA-templated	0.013
mitochondrion	0.024	zinc ion binding	0.013	response to cold	0.013
plasmodesma	0.024	heme binding	0.011	carbohydrate metabolic process	0.011
apoplast	0.022	identical protein binding	0.011	cell division	0.011
cell wall	0.022	iron ion binding	0.011	defense response	0.011

830 **Table 2:** Top ten most differentially expressed isoforms (with padj < 0.01) and their
831 corresponding annotations.

		Gene Ontology		
Transcript ID: Annotated Name (SPROT)	Log2 Fold Change	Cellular Components	Molecular Functions	Biological Processes
TRINITY_DN116220_c0_g1_i4: Zinc finger CCCH domain-containing protein 55	40.25		DNA binding, metal ion binding	
TRINITY_DN128685_c1_g3_i4: Callose synthase 3	33.85	1,3-beta-D-glucan synthase complex, integral component of membrane, plasma membrane	1,3-beta-D-glucan synthase activity	(1-¿3)-beta-D-glucan biosynthetic process, cell wall organization, regulation of cell shape
TRINITY_DN121298_c2_g2_i5: Heat shock 70 kDa protein 15	33.52	cell wall, cytosol, nucleus, plasma membrane, plasmodesma	ATP binding	
TRINITY_DN115892_c0_g1_i3: Pre-mRNA-splicing factor ATP-dependent RNA helicase DEAH1	33.52	membrane, spliceosomal complex	ATP binding, ATP-dependent 3'-5' RNA helicase activity, RNA binding	mRNA processing, posttranscriptional gene silencing by RNA, RNA splicing
TRINITY_DN127064_c0_g3_i1: LRR receptor-like serine/threonine-protein kinase HSL2	33.32	integral component of membrane, plasma membrane	ATP binding, protein serine/threonine kinase activity	defense response to Gram-negative bacterium, lateral root morphogenesis, leaf abscission, regulation of gene expression
TRINITY_DN128839_c3_g1_i6: Palmitoyl-acyl carrier protein thioesterase, chloroplastic	33.01	chloroplast	thiolester hydrolase activity	fatty acid biosynthetic process
TRINITY_DN121430_c10_g2_i1 Sucrose nonfermenting 4-like protein	32.64	chloroplast, chloroplast starch grain, cytoplasm, nucleus	kinase binding, maltose binding, protein kinase activator activity, protein kinase regulator activity, protein serine/threonine kinase activity	carbohydrate metabolic process, cellular response to glucose starvation, mitochondrial fission, peroxisome fission, pollen hydration, protein autophosphorylation, regulation of protein kinase activity, regulation of reactive oxygen species metabolic process
TRINITY_DN124527_c1_g1_i5: Bifunctional TH2 protein, mitochondrial	32.64	cytosol, mitochondrion	thiaminase activity, thiamine phosphate phosphatase activity	thiamine biosynthetic process
TRINITY_DN120224_c4_g1_i1: Enhancer of polycomb homolog 2	32.25	Piccolo NuA4 histone acetyltransferase complex		DNA repair, histone acetylation, regulation of transcription by RNA polymerase II
TRINITY_DN122787_c0_g1_i1: Protein IQ-DOMAIN 32	-32.16	chloroplast envelope, cytosol, microtubule associated complex, nucleus, plasma membrane		response to abscisic acid

Table 3: Differentially expressed isoforms (with padj < 0.01) in specific gene groups and their corresponding annotations.

Process Group	Transcript ID: Annotated Name (SPROT)	Log2 Fold Change	Gene Ontology		
			Cellular Components	Molecular Functions	Biological Processes
Abscisic Acid Signaling	TRINITY_DN122359_c1_g1_i3: AMP deaminase	-8.67	cytosol, endoplasmic reticulum, integral component of mitochondrial outer membrane, intracellular membrane-bounded organelle, nucleus	AMP deaminase activity, ATP binding, metal ion binding, protein histidine kinase binding	embryo development ending in seed dormancy, IMP salvage, response to abscisic acid
Abscisic Acid Signaling	TRINITY_DN117636_c1_g2_i8: Dual specificity protein phosphatase PHS1	-9.32	cytoplasm	kinase activity, phosphoprotein phosphatase activity, protein tyrosine phosphatase activity, protein tyrosine/serine/threonine phosphatase activity	abscisic acid-activated signaling pathway, cortical microtubule movement, regulation of gene expression, regulation of stomatal movement, response to abscisic acid
Abscisic Acid Signaling	TRINITY_DN121543_c6_g2_i1: Probable RNA-binding protein ARP1	-10.84	nucleus, ribonucleoprotein complex	mRNA binding, RNA binding	mRNA processing, regulation of seed germination, response to abscisic acid, response to salt stress, response to water deprivation
Abscisic Acid Signaling	TRINITY_DN122705_c2_g1_i2: ACT domain-containing protein ACR8	-22.92		amino acid binding	response to abscisic acid
Abscisic Acid Signaling	TRINITY_DN122787_c1_g1_i1: Protein IQ-DOMAIN 32	-32.16	chloroplast envelope, cytosol, microtubule associated complex, nucleus, plasma membrane		response to abscisic acid
Calcium-Dependent Protein Kinases	TRINITY_DN124121_c3_g1_i11: Calcium-dependent protein kinase 2	13.05	cytoplasm, nucleus	ATP binding, calcium ion binding, calcium-dependent protein serine/threonine kinase activity, calmodulin binding, calmodulin-dependent protein kinase activity	abscisic acid-activated signaling pathway, intracellular signal transduction, peptidyl-serine phosphorylation, protein autophosphorylation
Cytokinin Signaling	TRINITY_DN128252_c1_g4_i1: Ferritin-3, chloroplastic	8.22	chloroplast, chloroplast envelope, chloroplast stroma, chloroplast thylakoid membrane, cytoplasm, membrane, mitochondrion, thylakoid	ferric iron binding, ferrous iron binding, ferroxidase activity, identical protein binding, iron ion binding	flower development, intracellular sequestering of iron ion, iron ion transport, leaf development, photosynthesis, response to bacterium, response to cold, response to cytokinin, response to hydrogen peroxide, response to iron ion, response to reactive oxygen species
Cytokinin Signaling	TRINITY_DN124688_c1_g1_i3: Temperature-induced lipocalin-1	-9.18	chloroplast envelope, chloroplast membrane, cytoplasm, cytoplasmic side of plasma membrane, endoplasmic reticulum, Golgi apparatus, mitochondrion, plasma membrane, plasmodesma, vacuolar membrane, vacuole	nutrient reservoir activity, transporter activity	cellular chloride ion homeostasis, cellular sodium ion homeostasis, heat acclimation, hyperosmotic salinity response, lipid metabolic process, positive regulation of response to oxidative stress, positive regulation of response to salt stress, response to cold, response to cytokinin, response to freezing, response to heat, response to high light intensity, response to light stimulus, response to paraquat, response to reactive oxygen species, response to water deprivation, seed maturation
Cytokinin Signaling	TRINITY_DN126720_c3_g1_i6: Two-component response regulator ARR2	-19.29	nucleus	DNA binding, DNA-binding transcription factor activity, phosphorelay response regulator activity	cellular response to cytokinin stimulus, cytokinin-activated signaling pathway, ethylene-activated signaling pathway, leaf senescence, regulation of root meristem growth, regulation of seed growth, regulation of stomatal movement, response to cytokinin, response to ethylene, root development
Lignin Biosynthesis	TRINITY_DN125451_c5_g1_i3: Cinnamoyl-CoA reductase-like SNL6	-20.52		3-beta-hydroxy-delta5-steroid dehydrogenase activity, oxidoreductase activity, oxidoreductase activity, acting on the CH-OH group of donors, NAD or NADP as acceptor	defense response to bacterium, lignin biosynthetic process, steroid biosynthetic process
MADS-Box Genes	TRINITY_DN127253_c5_g3_i1: MADS-box transcription factor 33	6.60	nucleus	DNA-binding transcription factor activity, protein dimerization activity, RNA polymerase II regulatory region sequence-specific DNA binding	positive regulation of transcription by RNA polymerase II
Response to Gibberellin	TRINITY_DN110988_c0_g1_i2: Gibberellin 3-beta-dioxygenase 1	-8.35	cytoplasm	dioxygenase activity, gibberellin 3-beta-dioxygenase activity, metal ion binding	gibberellic acid mediated signaling pathway, gibberellin biosynthetic process, response to gibberellin, response to red light, response to red or far red light
Starch Biosynthesis	TRINITY_DN119512_c6_g1_i4: Granule-bound starch synthase 1, chloroplastic/amyloplastic	23.77	amyloplast, chloroplast	alpha-1,4-glucan synthase activity, glycogen (starch) synthase activity, starch synthase activity	starch biosynthetic process
Starch Biosynthesis	TRINITY_DN118583_c0_g1_i7: 1,4-alpha-glucan-branching enzyme, chloroplastic/amyloplastic	22.46	amyloplast, chloroplast	1,4-alpha-glucan branching enzyme activity, 1,4-alpha-glucan branching enzyme activity (using a glucosylated glycogenin as primer for glycogen synthesis), cation binding, hydrolase activity, hydrolyzing O-glycosyl compounds	carbohydrate metabolic process, glycogen biosynthetic process, starch biosynthetic process, starch metabolic process
Starch Biosynthesis	TRINITY_DN114909_c0_g1_i2: Sedoheptulose-1,7-bisphosphatase, chloroplastic	9.19	apoplast, chloroplast envelope, chloroplast stroma, thylakoid	metal ion binding, sedoheptulose-bisphosphatase activity	defense response to bacterium, reductive pentose-phosphate cycle, starch biosynthetic process, sucrose biosynthetic process

Table 4: Specific candidate genes and results from blasting to assembled transcriptome. Asterisk indicates marginal significance.

Gene Names	Original Taxon	Isoforms Found	Number DE
OsbHLH120	*Oryza sativa*	22	1*
IDD5	*Ipomea batatas*	63	0
WOX4	*Ipomea batatas*	31	0
Sulfite reductase	*Manihot esculenta*	9	0
FT-like	*Solanum tuberosum, Allium cepa*	37	1