Throwing shade: Physiological responses to light explain competition and facilitation in a tree diversity experiment

Shan Kothari,[1*] Rebecca Montgomery,[2] Jeannine Cavender-Bares[1,3]

[1]Department of Plant and Microbial Biology, University of Minnesota, 140 Gortner Laboratory, 1479 Gortner Ave., St. Paul, MN, 55108
[2]Department of Forest Resources, University of Minnesota, St. Paul, MN, 55108
[3]Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, MN, 55108

*corresponding author
Email: kotha020@umn.edu
Phone: (734) 502-2817

Author contributions: SK conceived the project with JCB, conducted physiological measurements, analyzed and interpreted the data with JCB and RM, and wrote the first draft of the paper. JCB and RM designed the FAB experiment, coordinated annual biomass surveys, and contributed to subsequent revisions of the paper.

Summary

- Interspecific facilitation is often invoked in explanations of biodiversity-ecosystem function relationships in plant communities, but it is seldom clear how it occurs. Physiological experiments show that excess light causes stress that may depress long-term carbon assimilation. If shading by a plant's neighbors reduces light stress, it may facilitate that plant's growth. On the other hand, when light is a limiting factor for growth, shading will often have a net negative, competitive effect.

- In a tree diversity experiment, we measured growth rates and photosynthetic physiology of broadleaf tree species across a gradient of light availability imposed by their neighbors. At the extremes, trees experienced nearly full sun (monoculture), or were shaded by nearby fast-growing conifers (shaded biculture).

- Although most species had lower growth with larger neighbors, implying a net competitive effect, the two most shade-tolerant species (*Tilia americana* and *Acer negundo*) had positive responses to neighbor size. Compared to the others, these two species were especially susceptible to photoinhibition (reduced dark-acclimated F_v/F_m) in full sun. While most species had lower

assimilation rates in shaded bicultures, *T. americana* had carbon assimilation rates up to 25%

higher. *T. americana* also dropped its leaves 3-4 weeks later in the shaded biculture, extending its

growing season. We conclude that although large neighbors can cause light limitation in shade-

intolerant species, they can also increase growth through abiotic stress amelioration in shade-

tolerant species.

• Both positive and negative species interactions in our experiment can be explained by the

photosynthetic responses of trees to the light environment created by their neighbors. We show

that physiological measurements can deepen our understanding of the species interactions that

underlie biodiversity-ecosystem function relationships.

Keywords: Photoprotection; photoinhibition; photosynthetic light-response curves; competition;

facilitation; tree diversity; biodiversity and ecosystem function

Introduction

The sun's light powers photosynthesis, the chain of reactions through which plants turn carbon dioxide into the living matter they use to grow and reproduce. A dominant view in plant ecology is that light is often a limiting resource, and competition for scarce light shapes the fate of plant species (Braun-Blanquet 1932; Canham et al. 1994; Monsi & Saeki 2005; Dybzinski & Tilman 2007). While this is often true, physiologists have also shown that, as other factors begin to limit photosynthesis, plants often absorb more light than they are able to use. Beyond an initial linear rise with increasing light availability, photosynthesis begins to saturate, such that additional light contributes little to the plant's carbon gain. Moreover, external factors that limit photosynthesis—including water limitation, cold temperatures, or nutrient-poor conditions—can exacerbate this excess of light (Fig. 1; Long et al. 1991; Demmig-Adams & Adams 1992; Demmig-Adams & Adams 2018).

Light is thus both an essential resource and a potential stressor. A chronic excess of light can cause lasting oxidative damage to plant cells, especially Photosystem II (PSII), and this damage reduces the efficiency of photosynthesis. Plants can avoid damage through mechanisms of photoprotection. These include biochemical pathways that shunt off excess light safely as heat—most notably, the xanthophyll cycle. They also include structural means of avoiding excess light interception, such as self-shading, reflective leaves, or steep leaf angles (Lovelock et al. 1992; Pearcy et al. 2005; Kothari et al. 2018). However, once induced, many forms of biochemical photoprotection are slow to reverse. Even when light conditions are no longer stressful, these mechanisms may continue to suppress photosynthesis for some time (Demmig-Adams and Adams 2006; Kromdijk et al. 2016). Similarly, structural mechanisms like self-shading may be costly under low ambient light. In general, photoprotective ability often increases with exposure to conditions that put plants at risk of damage in the habitat of origin, including high light (Montgomery et al. 2008) and cold or dry conditions (Cavender-Bares 2007; Savage et al. 2009; Wujeska et al. 2013; Ramírez-Valiente et al. 2013). In these environments, photoprotection may be necessary to prevent even more costly damage.

Depending on a plant's level of investment in photoprotection, these mechanisms may lack the

capacity to protect against prolonged exposure to excess light, which can reduce net carbon assimilation

rates (Egerton et al. 2000; Howell et al. 2002; Murchie & Niyogi 2011). Consistent with Long et al.

(1991), we use the term "photoinhibition" to describe a potential drop in carbon assimilation caused by

either photodamage or slow-reversing biochemical photoprotection. Aside from its direct effects, light can

also alter other microclimatic factors, like temperature, soil moisture, and vapor pressure deficit (VPD),

which can compound photoinhibition (Björkman & Powles 1984). For example, high VPD can lower

carbon assimilation by causing stomatal closure, which reduces the potential to use light for

photosynthesis; this effect, in turn, may increase the risk of photoinhibition, reducing photosynthetic

capacity further. Competition may also may increase a plant's risk of photoinhibition if it reduces its

access to resources in a way that limits photosynthesis. Thus, photodamage and photoprotective

investments are likely to affect plants' fitness and distribution across environmental gradients (Külheim et

al. 2002).

Given that light can cause stress, it may be possible for one plant species to facilitate another

through shading. Whether shading by one species has a positive or negative overall effect on another

depends on the balance of two effects:

1. Potential reduction of carbon gain through declines in light available for photosynthesis at

any given instant; and,

2. Avoidance of chronic stress, which may increase photosynthetic rates at any given light level.

Monteith et al. (1977) defined light-use efficiency (LUE) as net photosynthesis divided by the amount of

photosynthetically active radiation (PAR) absorbed. In terms of this formalism, an individual plant will

have higher photosynthetic rates in the shade when the proportional benefit of (2) is greater than the

proportional loss caused by (1). Changes in instantaneous irradiance alter the position along a

photosynthetic light-response curve; changes in LUE alter the parameters of the curve such that plants

assimilate more or less carbon at a given irradiance (Fig. 1).

Here, we test whether the shade created by trees can facilitate the carbon assimilation and growth

of neighboring trees in a biodiversity experiment. Experimental and observational research alike has

shown that biodiversity often promotes ecosystem functions such as productivity (Tilman et al. 2014). In

forests, this trend has been shown through both experimental (Potvin & Gotelli 2008; Tobner et al. 2016;

Grossman et al. 2017; Williams et al. 2017; Huang et al. 2018; Zemp et al. 2019) and observational

(Gamfeldt et al. 2013; Liang et al. 2016; Oehri et al. 2017) data, across temperate (Tobner et al. 2016;

Grossman et al. 2017; Williams et al. 2017), subtropical (Huang et al. 2018), and tropical (Potvin &

Gotelli 2008; Zemp et al. 2019) forests. These positive relationships are robust to variation in climate,

successional stage, and other factors (reviewed in Grossman et al. 2018; Ammer 2018).

One of the many factors that could contribute to positive biodiversity-productivity relationships is

interspecific facilitation through reduction of abiotic stress (Wright et al. 2017). Some research has

marshaled indirect evidence that facilitative interactions promote these relationships by linking patterns of

productivity to species traits (Fichtner et al. 2017). Nevertheless, the physiological mechanisms that

underlie these effects are poorly understood, despite their potential relevance for projecting species

interactions under environmental change.

In this paper, we examined the interactions among tree species in the Forests and Biodiversity

(FAB) experiment at Cedar Creek Ecosystem Science Reserve (East Bethel, MN). We ask: Do certain

species aid the growth of others by shielding them from stresses caused by high irradiance? In particular,

given that conifer species had higher initial growth rates (Table 1), we focused on whether they could

facilitate the growth of broadleaf species through shading.

We addressed this question by measuring the physiology of carbon gain among broadleaf species

across neighborhood environments. We examined whether photosynthetic physiology might explain

patterns in aboveground woody growth. If shading by neighbors does have a net positive effect for some

species, we would expect larger neighbors to cause focal individuals to have higher growth and carbon

assimilation rates. We also posed the hypothesis that trees exposed to high light would use structural and

biochemical photoprotective mechanisms more than those in lower light environments. If true, this

finding would imply that trees growing in high light must invest in photoprotection to avoid damage,

reinforcing our proposition that light stress influences the interactions within these communities.

Methods

Experimental design

The Forests and Biodiversity (FAB) experiment began in 2013 and comprises 142 4×4 m plots,

each planted with 64 trees in a 0.5×0.5 m grid. These plots form three blocks, each with 49 plots

arranged in a 7×7 square. Neighboring plots within blocks are immediately adjacent to one another.

Each plot is planted with 1, 2, 5, or 12 species from a pool of 12 total species (Table 1). These species

include four evergreen conifers with needle-like leaves (two distinct families) and eight deciduous,

broadleaf angiosperms (four distinct families). Except for five-species plots, each distinct species

composition has a replicate in each block. All species in a plot have approximately equal frequency, and

the placement of species within a plot is random. Further details about the experimental design can be

found in Grossman et al. (2017).

The mean relative growth rate in monoculture during the first three years varied dramatically

among species, being highest in conifers and *B. papyrifera* (Table 1). During these first three years,

Grossman et al. (2017) found that diverse plots overyield in productivity—their growth is greater than a

weighted average of the constituent species' growth in monoculture. Using the partition of Loreau &

Hector (2001), most of this overyielding was due to complementarity effects, which are often interpreted

as a result of synergistic interactions among species, rather than selection effects.

Tree growth

We surveyed tree growth in late fall of each year. For all living trees, we measured diameter and

height to the tallest leader. We measured basal diameter (5 cm from the ground) for trees less than 137 cm

tall; and diameter at breast height for trees more than 137 cm tall. For a subset of individuals, we

measured both diameters. To predict basal diameter when it was not measured, we used linear

relationships predicting basal diameter from the height of the same individual in the same year and the

basal diameter in the previous year.

We estimated woody stem volume as $V = \pi r^2 h$, where h is height and r is the basal radius (half

the basal diameter). This assumes each tree stem is a cylinder, an assumption that has been used and

justified in other tree diversity studies (Tobner et al. 2016; Williams et al. 2017), in the absence of

system-specific allometric equations. We then calculated the yearly relative growth rate (RGR) between

2016 and 2018 as $RGR = (\ln V_{2018} - \ln V_{2016})/2$ (Hoffman & Poorter 2002). We chose 2016 as a

starting point because no replanting of dead trees occurred after that point.

For each tree, we also calculated the average stem volume of its eight neighbors (in cardinal and

intercardinal directions) in 2018 as a proxy for the intensity of aboveground interactions. We assessed

neighbor effects on focal individual growth using ordinary least-squares (OLS) regression. When a

neighbor was missing, either due to mortality or because the focal individual was on the edge of the

experiment, we assigned that neighbor a volume of zero.

Photosynthetic physiology

We measured dark- and light-acclimated chlorophyll fluorescence parameters and reflectance

spectra across all eight broadleaf species. We also measured photosynthetic light-response curves across

four focal species (*A. rubrum*, *B. papyrifera*, *Q. ellipsoidalis*, and *T. americana*).

For measurements of physiology, we focused on three kinds of plots: (1) monocultures of

broadleaf species ("monoculture"); (2) bicultures comprising one broadleaf and one conifer species

("shaded biculture"); and (3) twelve-species plots ("twelve-species") (see Fig. S1 for sample images). For

each species in each treatment, we measured physiological parameters on six individuals—two in each of

three plots. Neither *A. negundo* nor *Q. macrocarpa* was planted with a conifer in any biculture plots. We

used *B. papyrifera* as a shaded biculture partner for *Q. macrocarpa* because it is a fast-growing species

that draws down light availability. We omitted the shaded biculture treatment for *A. negundo*.

Photosynthetic light-response curves

We measured photosynthetic-light response curves from the four focal species during July 2018

using an LI-6400 gas exchange system (LI-COR BioSciences, Lincoln, NE) with a 6400-40 Leaf

Chamber Fluorometer head. From each tree ($n = 72$ total), we selected a fully expanded upper leaf with

no visible sign of disease or herbivory. We noted the angle of each leaf relative to horizontal to the

nearest 15˚ before beginning measurements. Each curve had nine steps in descending order of brightness:

2000, 1500, 1000, 500, 200, 100, 50, 20, and 0 µmol m^{-2} s^{-1}. We also measured relative electron transport

rate (ETR) using chlorophyll fluorescence at each light level. We maintained favorable conditions inside

the chamber during each measurement. Following each curve, we removed the leaf to measure leaf mass

per area (LMA). Further details about light-response curve procedures and ETR calculations can be found

in the Supplemental Methods. Finally, we fitted a non-rectangular hyperbolic model (Lobo et al. 2013) to

each light-response curve using *R* code written by Nick Tomeo, available at:

https://github.com/Tomeopaste/AQ_curves.

We aimed to determine how carbon assimilation rates vary with the ambient light level, which we

define as the photosynthetic photon flux density (PPFD) one might measure in an open field adjacent to

the experiment. For most leaves we measured, *in situ* light availability is lower than the ambient light

level, mainly because each tree is shaded by its neighbors. To estimate each tree's carbon assimilation as

a function of ambient light, we have to transform its measured photosynthetic light-response curve along

the *x*-axis according to its light availability.

For example, if the topmost leaves of a particular tree only receive 50% of ambient light, we

would stretch its light-response curve two-fold along the *x*-axis. For this tree, the realized assimilation

rate we estimate at an ambient (or open-field) PPFD of 2000 µmol photons m^{-2} s^{-1} would be the

assimilation rate at an leaf-level PPFD of 1000 µmol m^{-2} s^{-1}. The exact scaling factor is estimated for

each individual as $RLA \times \cos(\theta)$, where *RLA* is relative light availability (see *Light availability*) and θ is

the leaf angle. (Taking the cosine of θ approximately corrects for the reduction in horizontally projected

leaf area in steeply inclined leaves when the sun is directly overhead, as simplified from Ehleringer &

Werk [1986].) We used this procedure on each fitted curve in order to compare across treatments in a way

that accounts for the variable light intensity each individual receives. We refer to photosynthetic light-

response curves that describe a function of the leaf chamber PPFD as *chamber light-response curves*, and

the rescaled light-response curves as a function of ambient PPFD as *ambient light-response curves*.

Finally, we used ambient light-response curves to estimate carbon assimilation by top leaves

throughout July. We used an hourly averaged time series of ambient solar radiation from Carlos Avery

Wildlife Management Area in Columbus, MN, 13.6 km away from the study site. Following Carruthers et

al. (2001), we assumed a conversion of 0.217 µmol photons m^{-2} s^{-1} of PAR per Watt m^{-2} of solar radiation.

By using these data as inputs to the ambient light-response curves, we could estimate integrated carbon

assimilation across the month of July. This procedure assumes that the photosynthetic response to light

remains constant, and is not affected by factors like leaf temperature and stomatal closure. This

assumption is unrealistic, but we nevertheless consider it a useful way of estimating the consequences of

realistic fluctuations in the light environment. We compared treatments within species using ANOVA.

Throughout the paper, we express assimilation per unit of leaf dry mass because we aim to

determine the plants' return on investment in biomass. As expected, leaves growing in low light

environments tended to have lower LMA in most species (Poorter et al. 2019), and expressing data on a

mass basis accounts for these large differences in construction cost. We also express stomatal

conductance (g_s) and electron transport rate (ETR) on a mass basis for comparison with assimilation rates,

with the caveat that both of these processes typically occur near the leaf surface.

Instantaneous chlorophyll fluorescence, pigments, and spectral reflectance

Across all eight broadleaf species, we measured chlorophyll fluorescence parameters using an

FMS2 pulse-modulated fluorometer (Hansatech Instruments Ltd., Norfolk, UK) over two days in late July

($n = 138$ total). We measured both dark- and light-acclimated parameters at the same spot on the same

leaves. For dark-acclimation, we placed opaque leaf clips on the leaf the evening before measurement.

We then measured F_v/F_m within two hours after sunrise; this parameter is a general index of

photoinhibition (including sustained photoprotection).

We took light-acclimated measurements between 12:00 and 14:00 each day. The protocol

involved the following steps: actinic light at 1000 µmol m^{-2} s^{-1} for 15 seconds, a saturating pulse, and two

seconds of far-red light. We exposed all leaves to the same actinic light because we aimed to assess

photoprotective capacity under comparable light conditions, even though individual trees had different

light environments.

From these data, we estimated qN, a parameter that indicates the extent of non-photochemical

quenching under the imposed actinic light; this parameter is correlated with de-epoxidation of

xanthophyll cycle pigments (Watling et al. 1997; Cavender-Bares and Bazzaz 2004). Following Kramer

et al. (2004), we also calculated the quantum yields ϕ_{PSII}, ϕ_{NPQ}, and ϕ_{NO}, which sum to 1. These three

parameters are the proportions of light energy dissipated through photochemistry, non-photochemical

quenching, and non-regulated energy dissipation, respectively. The last is a crucial parameter because it

represents the inability of a plant to dissipate light in a safe way; this portion of the absorbed light may

contribute to photodamage. The Supplemental Methods contain formulas for all parameters and

justifications for our choices in data analysis.

On separate top leaves of the same trees, we measured reflectance spectra (350-2500 nm) using a

PSR+ 3500 field spectroradiometer (Spectral Evolution, Lawrence, MA). We used these spectra to

calculate the photochemical reflectance index (PRI), a two-band index that most strongly indicates

carotenoid : chlorophyll ratios (Gamon et al. 1992; Wong & Gamon 2015; Gitelson et al. 2017). This

index is calculated as PRI = $(R_{531} - R_{570}) / (R_{531} + R_{570})$, where R_n is reflectance at a wavelength of n nm.

Low PRI tends to indicate high carotenoid : chlorophyll ratios. We estimated species-specific OLS

regression slopes of PRI and of each chlorophyll fluorescence parameter against relative light availability

(see *Light availability*).

To understand how photoprotective allocation varies with adaptations to shade, we used shade

tolerance values compiled by Niinemets & Valladares (2006) on a five-point scale based on prior

literature. Higher values along this scale indicate that plants are able to grow in lower light conditions.

Phenology

We also monitored the timing of leaf abscission in monoculture and shaded biculture plots for *T.*

americana to assess whether shade could delay senescence and abscission, extending the period for

carbon gain (Cavender-Bares et al. 2000). In early August, we surveyed 120 *T. americana* trees (60 in

monoculture, 60 in shaded biculture) to determine the proportion of leaves that had fallen based on how

many of the top five axillary buds lacked a leaf. From each of these trees, we marked a single leaf and

returned every 1-2 weeks to monitor its senescence. We assessed each leaf on a binary basis (senesced or

not) based on whether it had at least 50% remaining green leaf area. We chose this criterion as a simple

proxy for the continuous scale used by Cavender-Bares et al. (2000). Based on the initial survey and the

repeated surveys of marked leaves, we tracked the proportion of leaves that had senesced over time. To

ensure that leaves were photosynthetically active until near the point of senescence, we collected a one-

time measurement of dark-acclimated F_v/F_m in mid-September among remaining leaves.

We also performed a one-time measurement of 60 (30 per treatment) *A. rubrum* plants on October

1, using the same protocol we used to do our initial early August survey of *T. americana*. We aimed for

this survey to help test the validity of our results across species.

Water potential

Water deficits—even moderate ones—can reduce carbon gain by causing stomata to close

(Brodribb et al. 2003). To assess how our treatments affected water status, we measured leaf water

potential using a Scholander pressure bomb. We measured pre-dawn water potential (Ψ_{PD}) in all eight

broadleaf species ($n = 96$) and midday water potential (Ψ_{MD}; between 12:00 and 14:00) in only the four

focal species ($n = 48$). In these measurements, we included monoculture and shaded biculture treatments,

except in *A. negundo*, where we used the twelve-species treatment in place of the absent shaded biculture

treatment. For all water potential measurements, we removed leaves and immediately placed them in

impermeable plastic bags with wet paper towels, which we stored in a cooler. We then transported leaves

indoors to be measured within 90 minutes.

Because we only intended to test whether there was a general tendency, across species, for water

potential to vary with treatment, we included species identity as a random effect and treatment as a fixed

effect in both models.

Leaf angles

Leaf angle is a key control on light exposure because more vertical leaves intercept less light

when the sun is directly overhead, and allow light to percolate deeper into the canopy. As a result, steep

leaf angles can be a key structural strategy for avoiding light, temperature, and water stress (Valladares &

Pugnaire 1999; Werner et al. 1999; Pastenes et al. 2004; Valiente-Banuet et al. 2010). We predicted that

trees in monoculture would have steeper leaf angles than those in other treatments because they need to

avoid excess light. From each of three species (*A. rubrum*, *Q. ellipsoidalis*, and *T. americana*), we

randomly selected five individuals from each of six plots—three from monoculture treatments, three from

shaded bicultures ($n = 15$ per species × treatment). On these individuals, we measured the leaf angle from

horizontal, to the nearest 15˚, of the top five leaves in the crown.

Because angles of leaves within individuals may be non-independent, we tested differences

among treatments using a mixed-effects model with treatment as a fixed effect and individual tree nested

within plot as a random effect. Because deviations from horizontal in either direction can reduce light

interception, we used the absolute value of leaf angle as the dependent variable in statistical analyses. We

calculated *p*-values using Satterthwaite's degrees of freedom method in the *lmerTest* package

(Kuznetsova et al. 2017), with the caveat that such values should be interpreted conservatively.

Environmental data

Light availability

On a cloudy day, under diffuse light conditions, we measured light interception above each tree

selected for this study ($n = 138$) using an AccuPAR LP-80 ceptometer (METER Group, Pullman, WA).

We took the mean of two to four PAR measurements in the open and the mean of two measurements

directly above the topmost leaf of each tree. By calculating a ratio of these values, we could calculate the

percent of light transmitted to the top of each tree. We called this value *relative light availability*. This

value is typically well-correlated with the percentage of ambient light available over much longer time-

scales (Parent & Messier 1996).

Soil water content

We collected a one-time measurement of soil volumetric water content in mid-August using a

FieldScout Time-Domain Reflectometer 350 (Spectrum Technologies, Inc., Aurora, IL). This single

measurement followed six days from the last major rain. At midday, we collected and averaged three

measurements in each of 32 plots, spanning monocultures of the four core broadleaf species, shaded

bicultures, and twelve-species plots.

Air temperature

For nine days during early September, we set out four Thermochron iButtons (model DS1921G;

Maxim Integrated Products, Sunnyvale, CA, USA) in shaded plots and two in twelve-species plots,

setting them to log air temperature hourly. Each iButton was suspended on mesh inside a 1 m-tall PVC

pipe that was capped with an elbow to shield against solar irradiance. We compared these data to air

temperature measured hourly under open, sunny conditions at the Cedar Creek LTER weather station

about 0.77 km away. We assumed that data from this station would be representative of air temperature in

non-shaded areas. We ignored nighttime and early morning (20:00-08:00 h) readings because iButtons

may have been influenced by evaporative cooling of condensed water droplets.

Results

Tree size and growth

Neighborhood-scale relationships between focal individual RGR and total neighbor stem volume

were noisy but highly significant in most species (Fig. 2). The sign and strength of the relationships

varied among species. In most species, individuals with larger neighbors had lower RGR, including in *A.*

rubrum ($R^2 = 0.033$, $p <10^{-5}$), *Q. alba* ($R^2 = 0.038$, $p <10^{-6}$), *Q. ellipsoidalis* ($R^2 = 0.030$, $p <10^{-5}$), *Q.*

macrocarpa ($R^2 = 0.027$, $p <10^{-5}$), and *Q. rubra* ($R^2 = 0.056$, $p <10^{-9}$). On the other hand, some species

grew faster with larger neighbors, including *A. negundo* ($R^2 = 0.018$, $p < 0.005$) and *T. americana* ($R^2 =$

0.019, $p <10^{-4}$). *B. papyrifera* showed no significant relationship with neighbor size. These results are

qualitatively unchanged by eliminating trees on the edge of the experiment or incorporating plot as a

random effect.

Although these individual-level effects are noisy, they show strong tendencies in average stem

growth across the full range of potential neighbor size. Imagine two *T. americana* trees that begin at the

same size in 2016: One in monoculture, where the average neighbor in 2018 is 573 cm^3, and one growing

in a shaded biculture with *Pinus strobus*, where the average neighbor is 2175 cm^3. The neighborhood-

level regression predicts that after two years, the average *T. americana* in the shaded biculture would be

39.7% larger due to the difference in RGR. Or, consider *Q. rubra*, which had the strongest negative

response to neighbor size: We predict that an average tree in the shaded biculture would be 39.1% smaller

than in monoculture. Because these figures assume that treatments show no average size differences by

2016, they may underrepresent the true differences in size that have accumulated and may continue to

accumulate with time.

We can also aggregate these observations to the plot scale—for example, by considering the RGR

of summed stem volume of each species in each plot (Fig. 2). (This approach is distinct from an average

of individual RGRs, which assigns equal weight to small and large individuals.) Although aggregation

leaves less power to detect significant relationships, the relationships we find here explain much more of

the variation in growth across plots, including negative RGR responses to average neighbor biomass in *Q.*

alba ($R^2 = 0.311$, $p < 0.001$) and *Q. macrocarpa* ($R^2 = 0.174$, $p < 0.01$) and a positive response in *T.*

americana ($R^2 = 0.282$, $p < 0.001$).

Trees that died between 2016 and 2018 dropped out of our individual-level analyses of RGR

because their stem volume was 0 in 2018, making it impossible to calculate RGR. To see whether

mortality could alter our conclusions, we also performed statistical analyses on untransformed growth in

stem volume from 2017 to 2018, in which we alternately (1) treated mortality as a decline in volume to 0,

or (2) removed trees that died from analyses. The rate of mortality was low between 2016 and 2018

(~7.5%), and accounting for mortality makes no qualitative difference when considering absolute growth

as the response variable (Fig. S2).

Photosynthetic physiology

Photosynthetic light-response curves

In three out of the four focal species (*Q. ellipsoidalis*, *A. rubrum*, and *T. americana*),

photosynthetic rates (per unit dry mass) were higher in shaded biculture and twelve-species treatments

than in monoculture at any given chamber light level (Fig. 3, left). In the lone pioneer species, *B.*

papyrifera, photosynthetic rates in shaded biculture were lower than in the other two treatments.

The rate of photosynthesis is often limited by either ETR or by RuBP carboxylation rate, which in

turn is often limited by stomatal diffusion of CO_2 (Farquhar et al. 1980). In our data, both ETR and

stomatal conductance (g_s) increase with light availability (Fig. S3). (ETR tends to decline at very high

light levels, especially in shaded treatments, perhaps due to acute photoinhibition.) Within each species,

the rank-order of treatments in photosynthetic rates, ETR, and g_s tend to be broadly congruent. One

exception is that *T. americana* in shaded biculture has much higher g_s than in twelve-species or

monoculture plots, despite having slightly lower photosynthetic rates than in twelve-species plots.

Compared to chamber light-response curves, the picture that emerges from ambient light-

response curves is more complex (Fig. 3, right). In *B. papyrifera*, the shaded biculture treatment still had

the lowest assimilation rate across light levels. In late-successional *T. americana*, the shaded biculture and

twelve-species treatments had higher assimilation rates across most light levels, by up to 25%. In the

other two species, assimilation rates were highest in monoculture at low light availability; as top-of-

canopy light increases, assimilation rates for twelve-species and (in *Q. ellipsoidalis*) shaded biculture

plots eventually intersect and surpass monoculture rates.

Using a time series of solar radiation, we estimated total assimilation rates in July 2018 based on

each ambient light-response curve (Fig. S4). We found that trees would have lower total assimilation in

shaded biculture than in monoculture in *A. rubrum* (ANOVA; $p = 0.012$) and *B. papyrifera* ($p = 0.003$); in

both species, trees in twelve-species plots did not differ from those in monoculture. In the other two focal

species, there were no significant differences among treatments. In these integrated estimates, the costs of

shade seem larger than one might suppose from looking only at instantaneous assimilation rates under

high ambient light. We attribute these high costs to the fact that light levels were frequently low; PPFD

was under 500 µmol m^{-2} s^{-1} more than 60% of the time. It is under these low-light conditions that trees in

monoculture most outperform those in shaded biculture.

Instantaneous chlorophyll fluorescence, pigments, and spectral reflectance

Among all eight species except the early-successional *B. papyrifera*, dark-acclimated F_v/F_m

declined as relative light availability increased (Fig. 4), showing that high light exposure causes

photoinhibition. Across all trees of the four focal species, dark-acclimated F_v/F_m is positively correlated

with light-saturated photosynthetic rates (A_{sat}), as estimated from the non-rectangular hyperbolic model

(OLS regression; $R^2 = 0.44$; $p < 10^{-9}$; Fig. 5). Slopes are not significantly different among species. This

effect suggests that photoinhibitory declines in PSII efficiency may contribute to variation in light-

response curves.

In most species, qN rose with *in situ* light availability, indicating that trees that grow in high light

have a higher capacity to dissipate excess light with non-photochemical quenching (Fig. 4). However, two

species were clear exceptions: In *T. americana* and *A. negundo*, qN was nearly invariant across the range

of light availability they experienced, which suggests that these species are unable to upregulate

photoprotection in full sun.

Extracting the species-specific slopes of the change in dark-acclimated F_v/F_m with relative light

availability, we find that that the species with the greatest decline in F_v/F_m are the most shade-tolerant

(Fig. 5; $R^2 = 0.601$; $p = 0.025$). Similarly, more shade-tolerant species have less (or no) increase in qN at

high light availability ($R^2 = 0.619$; $p = 0.022$). These results suggest that shade-tolerant species may fail to

protect themselves from excess light when growing in light-saturated environments, making them

susceptible to photoinhibition. This suggestion is borne out in patterns of ϕ_{NO} across light environments,

which show that *T. americana* and *A. negundo* exposed to 1000 μmol photons m^{-2} s^{-1} are just as

susceptible to non-regulated dissipation of light when growing in full sun as in the shade (Fig. 4).

On a ternary plot, we plotted ϕ_{PSII}, ϕ_{NPQ}, and ϕ_{NO}—the quantum yields of PSII photochemistry,

non-photochemical quenching, and non-regulated dissipation, respectively—to illustrate how light energy

is dissipated differently across treatments (Fig. S5). Our results broadly show that across species, when

exposed to an equivalent, high level of light per unit area, trees that develop in the sun vary in ϕ_{PSII} and

ϕ_{NPQ} but have lower ϕ_{NO}. However, because they regularly experience higher light levels, they may still be

at high risk of damage.

PRI declined with increasing light availability in all species, which supports the idea that trees in

monoculture allocate more to carotenoids (Fig. 4). Although this relationship had remarkably consistent

slopes and intercepts across nearly all species, *B. papyrifera* had higher PRI at all light levels within its

narrow domain.

Phenology

In *T. americana*, leaf abscission in monoculture began in early August, and more than half of all

leaves had senesced by September 3 (Fig. 6). In shaded bicultures, more than 90% of leaves remained by

September 3, and no survey found greater than 50% senescence until October 1. We assigned the

senescence date of each leaf as the date of the first survey by which it had senesced; using this response

variable, leaves in monoculture senesced much earlier than those in shaded cultures (t-test; $p < 10^{-11}$).

These results suggest that leaves in shaded bicultures had, on average, one additional month for carbon

gain. Here, we assume that leaves in shaded biculture did not downregulate photosynthesis long before

abscission, which was supported by mid-September measurements of dark-acclimated F_v/F_m. In shaded

biculture, dark-acclimated F_v/F_m averaged 0.745 at this point, which was higher than the mean among

remaining monoculture leaves, 0.642 (t-test; $p = 0.061$).

Our one-time measurement of *A. rubrum* leaves in early October suggested that this pattern is not

limited to *T. americana*. *A. rubrum* trees in shaded biculture plots retained about 56% of their leaves,

while those in monoculture retained only 10% (t-test; $p < 10^{-4}$).

Water potential

For Ψ_{PD}, we removed two data points—one in shaded biculture, one in monoculture—with values

more negative than -0.8 MPa on the grounds that they may have been caused by measurement error.

Across treatments, the mean Ψ_{PD} and Ψ_{MD} were -0.15 and -1.29 MPa, respectively. Ψ_{PD} in shaded

bicultures was more negative than in monocultures by about 0.07 MPa ($p = 0.004$). This result holds even

when including the outliers ($p = 0.049$), with a mean difference of 0.08 MPa between treatments (Fig.

S6). By midday, this pattern reversed; Ψ_{MD} was more negative in monocultures than shaded bicultures,

with a mean difference of 0.33 MPa ($p = 0.004$).

Leaf angles

Among each of the three species in which we measured leaf angles, leaves were more horizontal

in the shaded biculture treatment (Fig. 7). The size of this effect varied among species: 9.0˚ in *Q.*

ellipsoidalis ($p = 0.037$), 22.0˚ in *A. rubrum* ($p = 0.022$), and 25.0˚ in *T. americana* ($p = 0.002$). The

direction of this shift varied among species. Because *T. americana* leaves usually drooped downward in

monoculture, leaf angles became less negative in shade. In contrast, *Q. ellipsoidalis*, whose leaves were

often inclined upwards in monoculture, had less positive leaf angles in the shade. In *A. rubrum*, leaf

angles in monoculture were inclined both up and down, while in shade, leaves had a tendency to become

more horizontal.

Environmental factors

In our survey, soil moisture varied among treatments (ANOVA; $p < 0.01$); it was lowest (5.7%)

in shaded biculture plots and highest (7.2%) in broadleaf monoculture plots. We found that daytime air

temperature in the open was higher than in shaded bicultures or twelve-species plots at 62.5% of hourly

log-points during the nine-day logging period. The air temperature in monocultures averaged 1.03˚ C

higher over this period.

Discussion

We investigated how species interactions in a tree diversity experiment might emerge from the

physiological responses of trees to the light environment created by their neighbors. We found that two

species (*A. negundo*, *T. americana*) grew faster on average with larger neighbors, while five (*A. rubrum*,

Q. alba, *Q. ellipsoidalis*, *Q. macrocarpa*, and *Q. rubra*) grew slower. The two species with positive

responses to neighbor size were shade-tolerant and faced a high risk of photoinhibition in high light,

which could cause carbon assimilation to be lower in monoculture. In contrast, the five species with

negative responses increased non-photochemical quenching in high light and tended to have lower carbon

gain in shaded bicultures. These results suggest that the divergent responses of these two groups of

species were driven, at least in part, by their tolerance to excess light. Here, we interpret these patterns in

further detail.

Physiology of carbon gain

We chose our four focal species along a gradient from early-successional, shade-intolerant *B.*

papyrifera, to mid-successional *A. rubrum* and *Q. ellipsoidalis*, to late-successional, shade-tolerant *T.*

americana (Braun 1954). This successional gradient aligns with the varied consequences of shade for

carbon gain across species. Growing in a shaded biculture can either: (1) decrease assimilation rates

(compared to monoculture) at all ambient light levels, as in *B. papyrifera*; (2) decrease assimilation under

low ambient light, but increase it under high light, as in *A. rubrum* and *Q. ellipsoidalis*; or (3) increase

assimilation rates at nearly all ambient light levels, as in *T. americana*. Our finding that carbon gain in *T.*

americana increases under shade is reinforced by Carter and Cavaleri (2018), who find that its

photosynthetic rates may increase down vertical gradients from the upper canopy to the sub-canopy.

The untransformed light-response curves can help us identify the causes of these patterns. As a

function of leaf chamber PPFD—without penalizing for lowered light availability—three of our four

species (*A. rubrum*, *Q. ellipsoidalis*, and *T. americana*) had higher photosynthetic rates in the shaded

biculture and twelve-species treatments than in monoculture (Fig. 3, left). In all species except early-

successional *B. papyrifera*, dark-acclimated F_v/F_m declines with light availability. Because dark-

acclimated F_v/F_m correlates with A_{sat}, this result suggests a role for photoinhibition in determining

photosynthetic rates. Because photoinhibition most often occurs through damage or regulation of PSII, it

is usually thought to depress photosynthesis by lowering ETR. However, both ETR and stomatal

conductance seem to reflect the patterns in photosynthesis across treatments in chamber light-response

curves (Fig. S3), suggesting that both have some role in explaining why most species showed lower

photosynthesis in monoculture.

Given that light availability fluctuates—diurnally, seasonally, and with cloud cover—the

treatment that performs best may also vary from moment to moment. We produced a more integrated

picture by estimating total assimilation during the month of July, assuming that each tree's photosynthetic

light-response curve remains constant (Fig. S4). While trees in shaded bicultures had lower estimated

assimilation than in monoculture in two of our focal species (*B. papyrifera* and *A. rubrum*), those in

twelve-species plots had similar assimilation compared to those in monoculture in all four species. It

seems plausible that the intermediate light environment in these twelve-species plots allowed trees to

avoid photoinhibition without becoming frequently light-limited. We caution that these measurements are

only based on top leaves, such that we cannot directly evaluate whole-plant carbon gain.

These estimates of assimilation during July cannot explain certain features in our stem biomass

data—including, for instance, the clearly positive effect of neighbor size on growth in *T. americana*. We

next discuss two kinds of phenomena that could contribute to variation in growth rates, but which

analyses based on light-response curves do not account for: Delayed senescence and microclimatic

effects. Both may contribute to potential positive effects of shading by neighbors.

T. americana in shade dropped their leaves nearly a month later into the fall compared to in

monoculture (Fig. 6). We also found accelerated senescence in full sun in *A. rubrum*. Given that most

leaves remained photosynthetically active until shortly before abscission (as in Mattila et al. 2018), shade

can allow plants a longer period of carbon gain. The pattern that the light environment affects senescence

timing admits at least two explanations: (1) that plants in the sun accumulate phenological cues for

senescence related to temperature, or (2) that excess light and other stresses cause damage that accelerates

senescence (Cavender-Bares et al. 2000), especially as the temperature declines, rendering leaves more

susceptible to photodamage (Juvany et al. 2013; Renner & Zohner 2019). Contrary to the first

explanation, air temperatures in full sun remained higher than in shaded plots.

Although we emphasize the role of light stress, other microclimatic factors could contribute to

variation in photosynthetic rates among treatments. Because we aimed to control the chamber

microclimate during light-response curves, we may have excluded some effects of the ambient

microclimate from our photosynthesis estimates. For example, midday leaf water potential tended to be

somewhat more negative in monoculture—perhaps due to greater VPD—which could make stomata close

more often (Fig. S6). On the other hand, pre-dawn leaf water potential was slightly more negative in

shaded bicultures, which may owe to lower soil moisture. Our water potential measurements show that

water stress was not very severe in any treatment, but even modest differences could affect stomatal

behavior (Brodribb et al. 2003). In addition, it seems plausible that greater light exposure may make leaf

temperature higher in monoculture (Schymanski et al. 2013), which could push leaves above thermal

optima for photosynthesis.

In summary, we contend that shade has both costs and benefits. Being shaded reduces the light

available for photosynthesis, but it may also prevent chronic stress due to excess light, allowing the plant

to use the light it absorbs more efficiently. Both factors must be jointly considered to determine whether

shade benefits or harms a plant.

Interpreting patterns of stem growth

The stem growth data are consistent with the hypothesis that larger neighbors can facilitate

growth in *T. americana* and *A. negundo*, the most shade-tolerant species. In light of the physiological

data, we interpret this pattern as a result of abiotic stress amelioration. Because we lacked measures of

root allocation, we cannot rule out the hypothesis that this trend is caused by a shade avoidance response

that increases allocation to shoots compared to roots (Schall et al. 2012). Nevertheless, the idea that much

of this effect is facilitative is supported both by the physiological data and by other studies that have

found that shade can increase total (root and shoot) biomass (e.g. Ryser & Eek 2000; Semchenko et al.

2012).

In the majority of species, including all four *Quercus* species and *A. rubrum*, larger neighbor size

caused focal individual size to be smaller, which we interpret as the result of net competitive interactions,

potentially both aboveground and belowground. The occurrence of belowground competition is suggested

by the fact that soil moisture was lowest in our dense-growing shaded biculture plots. *B. papyrifera*, a

shade-intolerant early-successional species whose leaves had lower photosynthetic rates in the shaded

biculture, had a neutral response to neighbor size, perhaps because its rapid height growth allowed it to

overtop neighbors rather than be overtopped. In contrast to all the other broadleaf species, its top leaves

suffered no reduction of light availability in shaded biculture, compared to monoculture.

Mechanisms of photoprotection

Plants use multiple means to avoid damage from excess light, which can be classified broadly

into biochemical and structural strategies. Our results show that trees in full sun allocate more to pigment-

based methods of photoprotection, as shown by lower PRI, and higher qN in most species. The two most

shade-tolerant species—*A. negundo* and *T. americana*—do not show a greater capacity for non-

photochemical quenching when grown in full sun. Across a gradient of shade to sun, these species may

experience large increases in ϕ_{NO}, and may suffer from greater photodamage as a result, as supported by

patterns of dark-acclimated F_v/F_m. This result is consistent with other research that finds that shade-

adapted species often fail to develop strong photoprotective abilities in the sun (Montgomery et al. 2008).

Our survey of leaf angles is also consistent with the idea that plants may steeply incline their

leaves to avoid intercepting excess light—a form of structural photoprotection. Leaves that are steeply

inclined tend to intercept less light per unit area, particularly during midday, when solar radiation is

otherwise most intense. The dramatic trend we observe in leaf angles suggests that these species are able

to use gross structural characteristics to partially regulate their light absorption.

Plants—especially those that grow in the shade—may lack perfect photoprotective mechanisms

in part because they are costly. Building pigments and proteins for photoprotection uses nutrients that

could otherwise be allocated to chlorophyll, RuBisCO, or structural tissue components. Furthermore,

slow relaxation of NPQ may hinder photosynthesis even when periodic declines in light availability

remove the immediate threat of damage (Zhu et al. 2004; Murchie & Niyogi 2011; Kromdijk et al. 2016).

Steep leaf angles may also reduce carbon assimilation by reducing light interception even under low light.

However, plants in full sun may need these mechanisms to avoid even greater long-term costs caused by

photodamage. Our measurements of the mechanisms of photoprotection suggest that these species may

incur some risk of damage under high light, reinforcing our conclusions about the role of light stress in

shaping species interactions.

The case for facilitation

An important result of our study is that shade from large neighbors increases both net

photosynthetic rates and stem growth rates in shade-tolerant species. This finding has precedents: Ball et

al. (1991) and Egerton et al. (2000) found that shade enhanced photosynthesis and growth of evergreen

Eucalyptus seedlings during the winter, when cold temperatures impose a risk of photoinhibition. Howell

et al. (2002) also found that the leafless outer branches of New Zealand's divaricating shrubs reduce

photoinhibition by shading inner leaves.

Much of the work on facilitation in plant communities is guided by the stress-gradient hypothesis

(SGH), which proposes that facilitative interactions are more common in stressful environments (Bertness

& Callaway 1994; Maestre et al. 2009). We can understand photoinhibition in light of the SGH as

follows: Stressful conditions limit the use of light for photochemistry, causing high light exposure to

become a potential stressor. Under these conditions, shade from neighbors can ameliorate stress, causing

facilitation. Under benign conditions, where plants can use more of the available light for photochemistry,

high light exposure may not cause damage. Here, shading by neighbors can cause light limitation,

resulting in an adverse, competitive effect on growth. As we show here, species' physiological traits also

influence whether they are stressed by their environment.

Ever since the SGH was first described, the role of facilitative interactions in structuring plant

communities has received more attention. For example, Bimler et al. (2018) suggest that models of

species occurrence will be more accurate when they incorporate positive interactions. Nevertheless, the

role of facilitation in structuring plant communities is still often overlooked (Wright et al. 2017), and its

mechanisms are seldom assessed by measuring physiological rates (but see, for example, Barron-Gafford

et al. 2017).

Competition and facilitation in biodiversity-ecosystem function relationships

Overyielding—or biodiversity-enhanced growth—is a hallmark of positive biodiversity-

ecosystem function relationships. A whole community can only overyield if at least one of its constituent

species overyields. Although we show how species growth changes with neighbor size, not neighbor

diversity, our results show that shading allows some species to achieve greater biomass in shaded

bicultures or twelve-species plots. In the same experiment, Grossman et al. (2017) found that *T.*

americana has the greatest overyielding in multi-species plots of any broadleaf species, especially when

growing with conifers or *B. papyrifera*. Grossman et al. (2017) further showed that conifer partners in

these species combinations also grow faster than in monoculture, resulting in community overyielding.

In general, the species that faced the strongest negative effects from shade had physiological

tolerance to high light and experienced shading as a form of competition. On the other hand, species that

were facilitated by shade had low tolerance to excess light and likely benefited from the milder

microclimate their neighbors created. We can place these findings in the context of the three classes of

facilitative mechanisms distinguished by Wright et al. (2017):

1. Biotic facilitation, which is mediated by the activity of higher-order trophic interactions, such as

promotion of shared mutualists or dilution of natural enemy loads;

2. Abiotic facilitation through resource enrichment, in which some species increase resource

availability for others; and,

3. Abiotic facilitation through stress amelioration, in which some species create a favorable

microclimate for others.

In the FAB experiment, enemy-mediated biotic facilitation is likely to have a limited role; Grossman et al.

(2018) found that diversity had a mild influence on damage by herbivores and pathogens, and the

direction of the relationship varied among tree species and herbivore guilds. Nutrient enrichment is

unlikely to be a major mechanism for facilitation here because, although nitrogen is the main limiting

mineral nutrient at Cedar Creek (Tilman 1987), none of the planted tree species associates with nitrogen-

fixing microbes. As a result, stress amelioration is likely to be the main class of facilitation in this

experiment, perhaps acting in concert with niche partitioning and (to a limited extent) selection effects to

drive overyielding.

In much of the literature on biodiversity and ecosystem function, the attempt to parse out

explanations—such as niche partitioning and facilitation—often depends on ever-finer statistical

partitions of biomass data (Loreau & Hector 2001; Fox & Kerr 2012), whose interpretation is often

contested (Carroll et al. 2011; Pillai et al. 2019). Because the processes that drive biomass accumulation

are ultimately physiological, physiological measurements can guide a mechanistic interpretation of

biomass patterns. These techniques can help us determine what specific kinds of plant-plant and plant-

environment interactions produce biodiversity-productivity relationships, and how these interactions

emerge from basic aspects of plant function.

In this study, the survey-based biomass data and the physiological measurements generally

converge in showing that certain species respond positively to the size of their neighbors, while others

respond negatively. However, the physiological measurements allow us to reach a deeper level of

explanation, showing how these interactions emerge in part from the photosynthetic responses of trees to

the light environment created by their neighbors. Thus, we demonstrate that community ecology can

benefit by incorporating mechanistic approaches from physiology.

Acknowledgements

We acknowledge that University of Minnesota, including Cedar Creek ESR, lies on the ancestral, traditional, and contemporary Land of the Dakota. This research was inspired in large part by conversations with Jake Grossman and Laura Williams. Beth Fallon, German Vargas G., Artur Stefanski, Daniel Stanton, and Danielle Way all gave valuable advice about measuring photosynthesis. Cathleen Nguyen and Chris Buyarski provided tremendous logistical support for data collection in FAB; Troy Mielke, Kally Worm, Jim Krueger, Pam Barnes, Mark Saxhaug, Dan Bahauddin, and others supported other crucial aspects of Cedar Creek infrastructure. Many Cedar Creek interns helped in the stem volume survey. Sarah Hobbie and Peter Reich contributed to the design of the FAB experiment. The Cavender-Bares lab (especially Gerard Sapès), Daniel Stanton, German Vargas G., Artur Stefanski, Jamie Mosel, José Lopez, and David Kramer all provided feedback on the results or manuscript. The FAB experiment is maintained with support from National Science Foundation under DEB #1234162 to Cedar Creek LTER. Spectral measurements were conducted as part of NSF/NASA DEB #1342778 to J.C.B. and R.M.. S.K. has been supported by an NSF Graduate Research Fellowship (Grant No. 00039202) and a UMN Doctoral Dissertation Fellowship. R.M. is also supported by Minnesota Agricultural Experiment Station project MIN-42-060.

References

Ammer C. Diversity and forest productivity in a changing climate. New Phytologist 221, 50–66 (2019).

Ball MC, Hodges VS, Laughlin GP. Cold-Induced Photoinhibition Limits Regeneration of Snow Gum at Tree-Line. Functional Ecology 5, 663–668 (1991).

Barron-Gafford GA, Sanchez-Cañete EP, Minor RL, Hendryx SM, Lee E, Sutter LF, Tran N, Parra E, Colella T, Murphy PC et al. Impacts of hydraulic redistribution on grass–tree competition vs facilitation in a semi-arid savanna. New Phytologist 215, 1451–1461 (2017).

Bertness MD, Callaway R. Positive interactions in communities. Trends in Ecology & Evolution 9, 191–193 (1994).

Bimler MD, Stouffer DB, Lai HR, Mayfield MM. Accurate predictions of coexistence in natural systems require the inclusion of facilitative interactions and environmental dependency. Journal of Ecology 106, 1839–1852 (2018).

Björkman O, Powles SB. Inhibition of photosynthetic reactions under water stress: interaction with light level. Planta 161, 490–504 (1984).

Braun-Blanquet J. Plant Sociology (trans. Fuller GD, Conard HS). New York: McGraw-Hill (1932).

Brodribb TJ, Holbrook NM, Edwards EJ, Gutiérrez MV. Relations between stomatal closure, leaf turgor and xylem vulnerability in eight tropical dry forest trees. Plant, Cell & Environment 26, 443–450 (2003).

Canham CD, Finzi AC, Pacala SW, Burbank DH. Causes and consequences of resource heterogeneity in forests: interspecific variation in light transmission by canopy trees. Can. J. For. Res. 24, 337–349 (1994).

Carroll IT, Cardinale BJ, Nisbet RM. Niche and fitness differences relate the maintenance of diversity to ecosystem function. Ecology 92, 1157–1165 (2011).

Cavender-Bares J. Chilling and freezing stress in live oaks (Quercus section Virentes): Intra- and interspecific variation in PS II sensitivity corresponds to latitude of origin. Photosynthesis Research 94, 437–453 (2007).

Cavender-Bares J, Bazzaz FA. From Leaves to Ecosystems: Assessing Photosynthesis and Plant Function in Ecological Studies. In Papageorgiou GC and Govindjee (Eds.), Chlorophyll fluorescence: A Signature of Photosynthesis (pp. 737–755). Dordrecht, Netherlands: Kluwer Academic Publishers (2004).

Cavender-Bares J, Potts M, Zacharias E, Bazzaz FA. Consequences of CO2 and light interactions for leaf phenology, growth, and senescence in Quercus rubra. Global Change Biology 6, 877–887 (2000).

Demmig-Adams B, Adams, WW III. Photoprotection and Other Responses of Plants to High Light Stress. Annual Review of Plant Physiology and Plant Molecular Biology 43, 599–626 (1992).

Demmig-Adams B, Adams, WW III. An integrative approach to photoinhibition and photoprotection of photosynthesis. Environmental and Experimental Botany 154, 1–3 (2018).

Dybzinski R, Tilman D. Resource Use Patterns Predict Long-Term Outcomes of Plant Competition for Nutrients and Light. The American Naturalist 170, 305–318 (2007).

755 Egerton JJG, Banks JCG, Gibson A, Cunningham RB, Ball MC. Facilitation of Seedling Establishment:
756 Reduction in Irradiance Enhances Winter Growth of *Eucalyptus pauciflora*. Ecology 81, 1437–1449
757 (2000).
758

759 Ehleringer J, Werk KS. Modifications of solar radiation absorption patterns and the implications for
760 carbon gain at the leaf level. In T. Givnish (ed.), On the Economy of Plant Form and Function.
761 Cambridge Univ. Press, London. 57-82 (1986).
762

763 Farquhar GD, von Caemmerer S, Berry JA. A biochemical model of photosynthetic CO_2 assimilation in
764 leaves of C_3 species. Planta 149, 78–90 (1980).
765

766 Fichtner A, Härdtle W, Li Y, Bruelheide H, Kunz M, von Oheimb G. From competition to facilitation:
767 how tree species respond to neighbourhood diversity. Ecology Letters 20, 892–900 (2017).
768

769 Fox JW, Kerr B. Analyzing the effects of species gain and loss on ecosystem function using the extended
770 Price equation partition. Oikos 121, 290–298 (2012).
771

772 Gamfeldt L, Snäll T, Bagchi T, Jonsson M, Gustafsson L, Kjellander P, Ruiz-Jaen MC, Fröberg M,
773 Stendahl J, Philipson CD et al. Higher levels of multiple ecosystem services are found in forests with
774 more tree species. Nature Communications 4, 1340 (2013).
775

776 Gamon JA, Peñuelas J, Field CB. A narrow-waveband spectral index that tracks diurnal changes in
777 photosynthetic efficiency. Remote Sensing of Environment 41, 35–44 (1992).
778

779 Gitelson AA, Gamon JA, Solovchenko, A. Multiple drivers of seasonal change in PRI: Implications for
780 photosynthesis 1. Leaf level. Remote Sensing of Environment 191, 110–116 (2017).
781

782 Grossman JJ, Cavender-Bares J, Reich PB, Montgomery RA, Hobbie SE. Neighborhood diversity
783 simultaneously increased and decreased susceptibility to contrasting herbivores in an early stage forest
784 diversity experiment. Journal of Ecology 107, 1492–1505 (2019).
785

786 Grossman JJ, Vanhellemont M, Barsoum N, Bauhus J, Bruelheide H, Castagneyrol B, Cavender-Bares J,
787 Eisenhauer N, Ferlian O, Gravel D et al. Synthesis and future research directions linking tree diversity to
788 growth, survival, and damage in a global network of tree diversity experiments. Environmental and
789 Experimental Botany 152, 68–89 (2018).
790

791 Hoffmann WA, Poorter H. Avoiding Bias in Calculations of Relative Growth Rate. Annals of Botany 90,
792 37 (2002).
793

794 Howell CJ, Kelly D, Turnbull MH. Moa ghosts exorcised? New Zealand's divaricate shrubs avoid
795 photoinhibition. Functional Ecology 16, 232–240 (2002).
796

797 Huang Y, Chen Y, Castro-Izaguirre N, Baruffol M, Brezzi M, Lang A, Li Y, Härdtle W, von Oheimb G,
798 Yang V et al. Impacts of species richness on productivity in a large-scale subtropical forest experiment.
799 Science 362, 80–83 (2018).
800

801 Juvany M, Müller M, Munné-Bosch S. Photo-oxidative stress in emerging and senescing leaves: a mirror
802 image? J Exp Bot 64, 3087–3098 (2013).
803

804   Kothari S, Cavender-Bares J, Bitan K, Verhoeven AS, Wang R, Montgomery RA, Gamon J. Community-
805   wide consequences of variation in photoprotective physiology among prairie plants. Photosynthetica 56,
806   455–467 (2018).
807
808   Kramer DM, Johnson G, Kiirats O, Edwards GE. New Fluorescence Parameters for the Determination of
809   QA Redox State and Excitation Energy Fluxes. Photosynthesis Research 79, 209–218 (2004).
810
811   Kromdijk J, Głowacka K, Leonelli L, Gabilly ST, Iwai M, Niyogi KK, Long SP. Improving
812   photosynthesis and crop productivity by accelerating recovery from photoprotection. Science 354, 857–
813   861 (2016).
814
815   Külheim C, Ågren J, Jansson S. Rapid Regulation of Light Harvesting and Plant Fitness in the Field.
816   Science 297, 91–93 (2002).
817
818   Liang J, Crowther TW, Picard N, Wiser S, Zhou M, Alberti G, Schulze E-D, McGuire AD, Bozzato F,
819   Pretzsch H et al. Positive biodiversity-productivity relationship predominant in global forests. Science
820   354, aaf8957 (2016).
821
822   Lobo FA, de Barros MP, Dalmagro HJ, Dalmolin ÂC, Pereira WE, de Souza ÉC, Vourlitis GL,
823   Rodríguez Ortíz CE. Fitting net photosynthetic light-response curves with Microsoft Excel — a critical
824   look at the models. Photosynthetica 51, 445–456 (2013).
825
826   Long SP, Humphries S, Falkowski PG. Photoinhibition of Photosynthesis in Nature. Annual Review of
827   Plant Physiology and Plant Molecular Biology 45, 633-62 (1994).
828
829   Loreau M, Hector A. Partitioning selection and complementarity in biodiversity experiments. Nature 412,
830   72–76 (2001).
831
832   Lovelock CE, Clough BF. Influence of solar radiation and leaf angle on leaf xanthophyll concentrations
833   in mangroves. Oecologia 91, 518–525 (1992).
834
835   Maestre FT, Callaway RM, Valladares F, Lortie CJ. Refining the stress-gradient hypothesis for
836   competition and facilitation in plant communities. Journal of Ecology 97, 199–205 (2009).
837
838   Mattila H, Valev D, Havurinne V, Khorobrykh S, Virtanen O, Antinluoma M, Mishra KB, Tyystjärvi E.
839   Degradation of chlorophyll and synthesis of flavonols during autumn senescence—the story told by
840   individual leaves. AoB Plants 10, (2018).
841
842   Monsi M, Saeki T. On the Factor Light in Plant Communities and its Importance for Matter Production.
843   Ann Bot 95, 549–567 (2005).
844
845   Monteith JL, Moss CJ, Cooke GW, Pirie NW, Bell GDH. Climate and the efficiency of crop production
846   in Britain. Philosophical Transactions of the Royal Society of London B: Biological Sciences 281, 277–
847   294 (1977).
848
849   Montgomery RA, Goldstein G, Givnish TJ. Photoprotection of PSII in Hawaiian lobeliads from diverse
850   light environments. Functional Plant Biol. 35, 595–605 (2008).
851
852   Murchie EH, Niyogi KK. Manipulation of Photoprotection to Improve Plant Photosynthesis. Plant
853   Physiology 155, 86–92 (2011).
854

855 Niinemets Ü, Valladares F. Tolerance to Shade, Drought, and Waterlogging of Temperate Northern
856 Hemisphere Trees and Shrubs. Ecological Monographs 76, 521–547 (2006).
857
858 Oehri J, Schmid B, Schaepman-Strub G, Niklaus PA. Biodiversity promotes primary productivity and
859 growing season lengthening at the landscape scale. PNAS 114, 10160–10165 (2017).
860
861 Parent S, Messier C. A simple and efficient method to estimate microsite light availability under a forest
862 canopy. Can. J. For. Res. 26, 151–154 (1996).
863
864 Pastenes C, Porter V, Baginsky C, Horton P, González J. Paraheliotropism can protect water-stressed
865 bean (Phaseolus vulgaris L.) plants against photoinhibition. Journal of Plant Physiology 161, 1315–1323
866 (2004).
867
868 Pearcy RW, Muraoka H, Valladares F. Crown architecture in sun and shade environments: assessing
869 function and trade-offs with a three-dimensional simulation model. New Phytologist 166, 791–800
870 (2005).
871
872 Pillai P, Gouhier TC. Not even wrong: the spurious measurement of biodiversity's effects on ecosystem
873 functioning. Ecology 100, e02645 (2019).
874
875 Poorter H, Niinemets Ü, Ntagkas N, Siebenkäs A, Mäenpää M, Matsubara S, Pons TL. A meta-analysis
876 of plant responses to light intensity for 70 traits ranging from molecules to whole plant performance. New
877 Phytologist 223, 1073–1105 (2019).
878
879 Potvin C, Gotelli NJ. Biodiversity enhances individual performance but does not affect survivorship in
880 tropical trees. Ecology Letters 11, 217–223 (2008).
881
882 Ramírez-Valiente JA, Koehler K, Cavender-Bares J. Climatic origins predict variation in photoprotective
883 leaf pigments in response to drought and low temperatures in live oaks (Quercus series Virentes). Tree
884 Physiology 35, 521–534 (2015).
885
886 Renner SS, Zohner CM. The occurrence of red and yellow autumn leaves explained by regional
887 differences in insolation and temperature. New Phytologist 224, 1464–1471 (2019).
888
889 Ryser P, Eek L. Consequences of phenotypic plasticity vs. interspecific differences in leaf and root traits
890 for acquisition of aboveground and belowground resources. American Journal of Botany 87, 402–411
891 (2000).
892
893 Savage, J., J. Cavender-Bares, and A. Verhoeven. Habitat generalists and wetland specialists in the genus
894 Salix vary in their photoprotective responses to drought. Functional Plant Biology 36, 300–309 (2009).
895
896 Schall P, Lödige C, Beck M, Ammer C. Biomass allocation to roots and shoots is more sensitive to shade
897 and drought in European beech than in Norway spruce seedlings. Forest Ecology and Management 266,
898 246–253 (2012).
899
900 Schymanski SJ, Or D, Zwieniecki M. Stomatal Control and Leaf Thermal and Hydraulic Capacitances
901 under Rapid Environmental Fluctuations. PLOS ONE 8, e54231 (2013).
902
903 Semchenko M, Lepik M, Götzenberger L, Zobel K. Positive effect of shade on plant growth: amelioration
904 of stress or active regulation of growth rate? Journal of Ecology 100, 459–466 (2012).
905

Tilman D. Secondary Succession and the Pattern of Plant Dominance Along Experimental Nitrogen
Gradients. Ecological Monographs 57, 189–214 (1987).

Tilman D, Isbell F, Cowles JM. Biodiversity and Ecosystem Functioning. Annual Review of Ecology,
Evolution, and Systematics 45, 471–493 (2014).

Tobner CM, Paquette A, Gravel D, Reich PB, Williams LJ, Messier C. Functional identity is the main
driver of diversity effects in young tree communities. Ecol Lett 19, 638–647 (2016).

Valiente-Banuet A, Verdú M, Valladares F, García-Fayos P. Functional and evolutionary correlations of
steep leaf angles in the mexical shrubland. Oecologia 163, 25–33 (2010).

Valladares F, Pugnaire FI. Tradeoffs Between Irradiance Capture and Avoidance in Semi-arid
Environments Assessed with a Crown Architecture Model. Annals of Botany 83, 459–469 (1999).

Watling JR, Robinson SA, Woodrow IE, Osmond CB. Responses of Rainforest Understorey Plants to
Excess Light during Sunflecks. Australian Journal of Plant Physiology 24, 17–25 (1997).

Werner C, Correia O, Beyschlag W. Two different strategies of Mediterranean macchia plants to avoid
photoinhibitory damage by excessive radiation levels during summer drought. Acta Oecologica 20, 15–23
(1999).

Williams LJ, Paquette A, Cavender-Bares J, Messier C, Reich PB. Spatial complementarity in tree crowns
explains overyielding in species mixtures. Nature Ecology & Evolution 1, 0063 (2017).

Wong CYS, Gamon JA. Three causes of variation in the photochemical reflectance index (PRI) in
evergreen conifers. New Phytol 206, 187–195 (2015).

Wright AJ, Wardle DA, Callaway R, Gaxiola A. The Overlooked Role of Facilitation in Biodiversity
Experiments. Trends in Ecology & Evolution 32, 383–390 (2017).

Wujeska A, Bossinger G, Tausz M. Responses of foliar antioxidative and photoprotective defence
systems of trees to drought: a meta-analysis. Tree Physiology 33, 1018–1029 (2013).

Zemp DC, Gérard A, Hölscher D, Ammer C, Irawan B, Sundawati L, Teuscher M, Kreft H. Tree
performance in a biodiversity enrichment experiment in an oil palm landscape. Journal of Applied
Ecology 56, 2340–2352 (2019).

Zhu X-G, Ort DR, Whitmarsh J, Long SP. The slow reversibility of photosystem II thermal energy
dissipation on transfer from high to low light may cause large losses in carbon gain by crop canopies: a
theoretical analysis. J Exp Bot 55, 1167–1175 (2004).

Tables

958   Table 1: Characteristics of species in FAB. Except for *B. papyrifera*, broadleaf species tended to grow

959   slower than needleleaf conifers. Shade tolerance is drawn from Niinemets and Valladares (2006), who

960   evaluated it along a 1-5 scale where higher values indicate an ability to grow under lower light.

Scientific name	Species code	Common name	Family	Leaf form	Mean RGR in monoculture, 2013-6 (/year)	Shade tolerance	Partner in shaded biculture
Acer negundo	ACNE	Box elder	Sapindaceae	Broadleaf	0.039	3.47	None
Acer rubrum	ACRU	Red maple	Sapindaceae	Broadleaf	0.379	3.44	*Pinus banksiana*
Betula papyrifera	BEPA	Paper birch	Betulaceae	Broadleaf	1.272	1.54	*Pinus banksiana*
Juniperus virginiana	JUVI	Eastern redcedar	Cupressaceae	Needleleaf	0.975	1.28	NA
Pinus banksiana	PIBA	Jack pine	Pinaceae	Needleleaf	2.288	1.36	NA
Pinus resinosa	PIRE	Red pine	Pinaceae	Needleleaf	1.619	1.89	NA
Pinus strobus	PIST	White pine	Pinaceae	Needleleaf	1.722	3.21	NA
Quercus alba	QUAL	White oak	Fagaceae	Broadleaf	0.735	2.85	*Pinus strobus*
Quercus ellipsoidalis	QUEL	Northern pin oak	Fagaceae	Broadleaf	0.722	NA	*Pinus strobus*
Quercus macrocarpa	QUMA	Bur oak	Fagaceae	Broadleaf	0.547	2.71	*Betula papyrifera*
Quercus rubra	QURU	Northern red oak	Fagaceae	Broadleaf	0.505	2.75	*Pinus strobus*
Tilia americana	TIAM	American basswood	Tiliaceae	Broadleaf	0.724	3.98	*Pinus strobus*

Figure captions

Fig. 1: (A) Consider two hypothetical photosynthetic light-response curves of unstressed (blue) and

photoinhibited (red) plants, where the unstressed plant's neighbors reduce the amount of light it absorbs.

Given a light-response curve, an instantaneous reduction in light tends to reduce photosynthetic rates

(represented by the small black arrow along the red curve). However, if a long-term reduction in light

spares the plant from chronic photoinhibition, it can alter the parameters of the curve such that carbon

assimilation at any given instantaneous light level is greater (represented by the small black vertical

arrow). The balance (represented by the large dashed arrow) may either increase or reduce net carbon

assimilation.

(B) To allow us to compare photosynthetic rates visually, we can put the two light-response curves on a

common *x*-axis that represents light arriving at the top of the canopy. To do this, we need to scale the blue

light response curve horizontally to account for the fact that when the stressed plant receives the amount

of light corresponding to the red dashed line in (A), the unstressed plant receives a proportionally lower

amount, here represented by the blue dashed line. By stretching the blue curve proportionally such that

these two lines coincide, we can compare the true estimated photosynthetic rates at a given light

availability.

Fig. 2: Relative growth rate (RGR) of woody stem volume in individuals of each species between fall

2016 and 2018 as a function of the total stem volume of all neighbors. Gray dots represent individuals,

and the regression line is fit through the individual-level data. Large colored dots are aggregated to the

plot scale, as described in the main text, and color-coded by treatment. A shaded biculture is a plot in

which the focal broadleaf species grows with either a conifer or *B. papyrifera* (unless *B. papyrifera* is the

focal broadleaf species). An open biculture is a plot in which the focal broadleaf grows with another

broadleaf species (other than *B. papyrifera*).

Fig. 3: Mass-based photosynthetic light-response curves for four broadleaf species. The left panels

display photosynthetic rates as a function of chamber light intensity; the right panels represent an estimate

of realized photosynthetic rates as a function of light intensity at the top of the canopy, after accounting

for light interception by neighbors (as in Fig. 1B). Species are arranged from the least shade-tolerant (*B.*

papyrifera) to the most (*T. americana*). Error bars in left panels are ± 1 SE.

Fig. 4: Dark-acclimated F_v/F_m, qN, ϕ_{NO}, and PRI as a function of relative light availability across species

and treatments.

1002

1003 Fig. 5: Compared to shade-intolerant species, shade-tolerant species show greater declines in dark-

1004 acclimated F_v/F_m and smaller increases in qN as light availability increases. A_{sat} derived from light-

1005 response curves correlates positively and strongly with dark-acclimated F_v/F_m; slopes are statistically

1006 indistinguishable among species.

1007

1008 Fig. 6: Timing of fall leaf abscission in monoculture and shaded biculture treatments of *T. americana*.

1009

1010 Fig. 7: Top leaf angles of *A. rubrum* (left), *Q. ellipsoidalis* (middle), and *T. americana* (right) across

1011 treatments.

1012

1013 Fig. S1: Example images of *Tilia americana* in shaded biculture (top) and monoculture (bottom). Red

1014 scale bars represent about 1 m in the foreground.

1015

1016 Fig. S2: Growth of woody stem volume in individuals of each species between fall 2017 and 2018 as a

1017 function of the total stem volume of all neighbors. Gray dots represent individuals, while colored dots are

1018 aggregated to the plot scale and color-coded by treatment. Two regression lines are fit: One (blue) to all

1019 data, and one (red) excluding trees that died between 2017 and 2018.

1020

1021   Fig. S3: Responses of ETR and stomatal conductance to light availability for the four focal broadleaf

1022   species. Error bars are ± 1 SE.

1023

1024   Fig. S4: An estimate of total carbon assimilation across species and treatments throughout the month of

1025   July, calculated by applying a time series of solar radiation from a nearby weather station as inputs to our

1026   ambient light-response curves.

1027

Fig. S5: A ternary plot depicting variation among species and treatments in ϕ_{PSII}, ϕ_{NPQ}, and ϕ_{NO}, as

calculated using the lake model derivations of Kramer et al. (2004). These three quantities describe light

energy dissipation through photochemistry, through non-photochemical quenching, and through non-

regulated dissipation, respectively. A thick dashed line represents 0.83 the theoretical maximum

efficiency of PSII posited in Tietz et al. (2017). Deviations from this line represent increasingly high ϕ_{NO},

which can result in photodamage.

1034

1035   Fig. S6: Water potential before dawn (top) and at midday (bottom) among species and treatments.

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046   **Figures**

1047

1048   **Fig. 1**

 Fig. 2

1087   **Fig. 3**

1088

1089 **Fig. 4**

Fig. 5

1120   **Fig. 6**

1121
1122
1123   **Fig. 7**

1130 **Supplement to: "Throwing shade: Physiological responses to light mediate competition and**

1131 **facilitation in a tree diversity experiment"**

1132

Materials and Methods

Photosynthetic light-response curve methods

We measured photosynthetic-light response curves from the four focal species (*Acer negundo*,

Betula papyrifera, *Quercus ellipsoidalis*, and *Tilia americana*) during July 2018 using an LI-6400 gas

exchange system (LI-COR BioSciences, Lincoln, NE) with a 6400-40 Leaf Chamber Fluorometer head.

From each tree ($n = 72$ total), we selected a fully expanded, sun-exposed leaf with no visible sign of

disease or herbivory. We noted the angle of each leaf relative to horizontal to the nearest 15˚ before

beginning measurements. Each curve had nine steps in descending order of brightness: 2000, 1500, 1000,

500, 200, 100, 50, 20, and 0 μmol m^{-2} s^{-1}.

During each curve, we controlled environmental conditions to reduce the immediate effects of

environmental conditions, other than light, on stomatal conductance. We maintained relative humidity at a

favorable level, typically between 65 and 75%, and kept block temperature as close as possible to 25 ˚C.

The reference CO_2 concentration remained at 400±10 ppm. We kept each leaf in the chamber for at least

five minutes before beginning each curve. We checked that outputs were stable and matched IRGAs

before each measurement.

While we sought to control environmental conditions in the chamber, any remaining differences

among treatments could, in theory, contribute to the differences we observe in photosynthetic function.

During light-response curves, we found no difference among treatments in leaf-to-air VPD within the

chamber. There were significant but slight differences in leaf temperature inside the chamber; leaves in

monoculture, shaded biculture, and twelve-species plots had average temperatures of 27.7 ˚C, 26.5 ˚C,

and 27.0 ˚C, respectively. Previous studies on some of the same species in Minnesota have typically

found thermal optima below 26 ˚C (Sendall et al. 2015). However, while cooler leaves may be closer to

thermal optima, the small differences in temperature we observed seem unlikely to explain much of the

observed variation in photosynthetic rates within the leaf chamber.

In all but five light-response curves, we also used a saturating pulse to measure light-acclimated

chlorophyll fluorescence at each light level. These data allowed us to calculate ϕ_{PSII}, a measure of the

efficiency of Photosystem II (PSII) at each actinic light level, as:

$$\phi_{PSII} = \frac{F'_m - F_s}{F'_m}$$

where F_s is the steady-state fluorescence yield under actinic light, and F_m' is the maximum fluorescence

yield of the actinic light-acclimated sample following a saturating light pulse, which temporarily closes

all PSII reaction centers. From this parameter, we could calculate electron transport rate as:

$$ETR = \phi_{PSII} \times PPFD \times 0.42 \times 0.5$$

where PPFD is photosynthetic photon flux density (the light level inside the chamber, in μmol m^{-2} s $^{-1}$),

0.42 is an estimate of leaf absorptance, and 0.5 is an estimate of the fraction of photons that are captured

by PSII rather than PSI. Electron transport rate is a key parameter because electron transport is one of the

main steps that is capable of limiting photosynthesis (Farquhar et al. 1980), especially when light

availability or PSII efficiency are low.

1170

Chlorophyll fluorescence procedure

As described in the main text, we used dark- and light-acclimated measurements of chlorophyll

fluorescence to calculate several derived parameters, from which we make inferences about

photosynthetic and photoprotective function. Most importantly, we wanted to assess: (1) capacity for non-

photochemical quenching, as measured through the parameters qN or NPQ; and (2) the partitioning of

light energy among different processes, as assessed using the quantum yields ϕ_{PSII}, ϕ_{NPQ}, and ϕ_{NO}.

These derived chlorophyll fluorescence parameters are calculated from a few basic measurements

taken during the dark- and light-acclimated measurements, including: F_o, the minimum fluorescence yield

of a dark-acclimated sample with all PSII reaction centers open; F_o', the minimum fluorescence yield of a

light-acclimated sample with all PSII reaction centers open (following far-red light exposure); and F_m, the

maximum fluorescence yield of a dark-acclimated sample following a saturating pulse. F_v, the variable

fluorescence yield of a dark-acclimated sample, is calculated as $F_m - F_o$.

1183   The parameters qN and NPQ are meant to indicate the rate constant of the thermal dissipation of

1184   energy from PSII (Bilger and Schreiber 1986; Bilger and Björkman 1990). These parameters increase

1185   with the use of NPQ to dissipate light—in this case, under exposure to 1000 µmol m^{-2} s^{-1}. They are

1186   calculated, respectively, as:

1187
$$qN = 1 - \frac{F_m' - F_o'}{F_m - F_o}; NPQ = \frac{F_m - F_m'}{F_m'}$$

1188   In our dataset, qN is highly correlated ($R^2 = 0.96$) with the natural logarithm of NPQ; we chose to report

1189   qN to easily visualize variation among leaves with low NPQ and to help satisfy the error term normality

1190   assumption of OLS regression.

1191   The quantum yields ϕ_{PSII}, ϕ_{NPQ}, and ϕ_{NO} represent the proportions of absorbed light energy that are

1192   dissipated through photochemical processes, non-photochemical quenching, and non-regulated

1193   dissipation, respectively. As proportions, these quantum yields sum to 1. We calculated these parameters

1194   following the "lake model" derivation of Kramer et al. (2004), which treats photosynthetic reaction

1195   centers as connected by shared antennae. We achieve qualitatively similar results using the simpler

1196   analogues found in Hendricksen et al. (2005).

1197   The parameter ϕ_{PSII} is as calculated above (Supplemental Methods, *Photosynthetic light-response*

1198   *curve procedure*). In the derivation from Kramer et al. (2004), ϕ_{NPQ} and ϕ_{NO} are calculated using the

1199   parameter qL, an estimate of the fraction of open PSII reaction centers. qL is calculated as:

1200
$$qL = \frac{F_m' - F_s}{F_m' - F_o'} \frac{F_o'}{F_s}$$

1201   Given this parameter, we can calculate ϕ_{NO} as:

1202
$$\phi_{NO} = \frac{1}{NPQ + 1 + qL\left(\frac{F_m}{F_o} - 1\right)}$$

and given the other two quantum yields, we can calculate ϕ_{NPQ} as:

$$\phi_{NPQ} = 1 - \phi_{PSII} - \phi_{NO}$$

Many researchers, including Murchie and Lawson (2013) and Tietz et al. (2017) treat ~0.83 as a

theoretical maximum of ϕ_{PSII} or dark-acclimated F_v/F_m after a period of dark-acclimation long enough to

reduce non-photochemical quenching to zero. Similarly, one may treat ~0.83 as a theoretical maximum of

$\phi_{PSII} + \phi_{NPQ}$, deviations from which indicate high ϕ_{NO}, and thus, a declining ability to dissipate light

energy in a regulated way. We indicate this putative maximum in our ternary plot of the quantum yields

(Fig. S5).

1211

1212   References
1213
1214   Bilger W, Björkman O. Role of the xanthophyll cycle in photoprotection elucidated by measurements of
1215   light-induced absorbance changes, fluorescence and photosynthesis in leaves of *Hedera canariensis*.
1216   Photosynthesis Research 25, 173–185 (1990).
1217
1218   Bilger W, Schreiber U. Energy-dependent quenching of dark-level chlorophyll fluorescence in intact
1219   leaves. Photosynthesis Research 10, 303–308 (1986).
1220
1221   Farquhar GD, von Caemmerer S, Berry JA. A biochemical model of photosynthetic CO2 assimilation in
1222   leaves of C 3 species. Planta 149, 78–90 (1980).
1223
1224   Hendrickson L, Förster B, Pogson BJ, Chow WS. A simple chlorophyll fluorescence parameter that
1225   correlates with the rate coefficient of photoinactivation of Photosystem II. Photosynthesis Research 84,
1226   43–49 (2005).
1227
1228   Kramer DM, Johnson G, Kiirats O, Edwards GE. New Fluorescence Parameters for the Determination of
1229   QA Redox State and Excitation Energy Fluxes. Photosynthesis Research 79, 209–218 (2004).
1230
1231   Murchie EH, Lawson T. Chlorophyll fluorescence analysis: a guide to good practice and understanding
1232   some new applications. Journal of Experimental Botany 64, 3983–3998 (2013).
1233
1234   Sendall, KM, Reich PB, Zhao C, Jihua H, Wei X, Stefanski A, Rice K, Rich RL, Montgomery RA.
1235   Acclimation of photosynthetic temperature optima of temperate and boreal tree species in response to
1236   experimental forest warming. Global Change Biology 21, 1342–1357 (2015).
1237
1238   Tietz S, Hall CC, Cruz, JA, Kramer DM. NPQ(T): a chlorophyll fluorescence parameter for rapid
1239   estimation and imaging of non-photochemical quenching of excitons in photosystem-II-associated
1240   antenna complexes. Plant, Cell & Environment 40, 1243–1255 (2017).
1241
1242
1243
1244

1245   **Fig. S1**

1259   **Fig. S2**

1260

1261   **Fig. S3**

1262

 Fig. S4

 Fig. S5

1268   **Fig. S6**

1269