Physiological responses to light explain competition and facilitation in a tree diversity experiment

Shan Kothari,[1,2]* Rebecca Montgomery,[3] Jeannine Cavender-Bares[1,4]

[1]Department of Plant and Microbial Biology, University of Minnesota, 140 Gortner Laboratory, 1479 Gortner Ave., St. Paul, MN, 55108, United States of America
[2]Institut de recherche en biologie végétale, Université de Montréal, Montréal, Québec, H1X 2B2, Canada
[3]Department of Forest Resources, University of Minnesota, St. Paul, MN, 55108, United States of America
[4]Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, MN, 55108, United States of America

*corresponding author
Email: kotha020@umn.edu

Author contributions: SK conceived the project with JCB, conducted physiological measurements, analyzed and interpreted the data with JCB and RM, and wrote the first draft of the paper. JCB and RM designed the FAB experiment, coordinated annual growth surveys, and contributed to subsequent revisions of the paper.

Abstract

- Ecologists often invoke interspecific facilitation to help explain positive biodiversity-ecosystem function relationships in plant communities, but seldom test how it occurs. One mechanism through which one species may facilitate another is by ameliorating abiotic stress. Physiological experiments show that a chronic excess of light can cause stress that depresses carbon assimilation. If shading by a plant's neighbors reduces light stress enough, it may facilitate that plant's growth. If light is instead most often a limiting factor for photosynthesis, shading may have an adverse, competitive effect.

- In a temperate tree diversity experiment, we measured stem growth rates and photosynthetic physiology in broadleaf trees across a gradient of light availability imposed by their neighbors. At the extremes, trees experienced nearly full sun (monoculture), or were shaded by nearby fast-growing conifers (shaded biculture).

- Most species had slower growth rates with larger neighbors, implying a net competitive effect. On the other hand, the two most shade-tolerant species (*Tilia americana* and *Acer negundo*) and

the most shade-intolerant one (*Betula papyrifera*) had faster stem growth rates with larger

neighbors. The two most shade-tolerant species had large increases in photoinhibition (reduced

dark-acclimated F_v/F_m) across the gradient of increasing light availability, which suggests they

are more vulnerable to chronic light stress. While most species had lower carbon assimilation

rates in the shaded biculture treatment, *T. americana* had rates up to 25% higher. *T. americana*

also dropped its leaves 3-4 weeks earlier in monocultures, curtailing its growing season. We

conclude that although large neighbors can cause light limitation in shade-intolerant species, they

can also increase growth through abiotic stress amelioration in shade-tolerant species. Finally, in

shade-intolerant *B. papyrifera*, we find a pattern of stem elongation in trees with larger neighbors,

which may suggest that a shade avoidance response accounts for the apparent positive trend in

stem volume.

• Synthesis: Both positive and negative species interactions in our experiment can be explained in

large part by the photosynthetic responses of trees to the light environment created by their

neighbors. We show that photosynthetic physiology can help explain the species interactions that

underlie biodiversity-ecosystem function relationships. The insights that ecologists gain by

searching for such physiological mechanisms may help us forecast species interactions under

environmental change.

Keywords: Biodiversity and ecosystem function; facilitation; light competition; photoinhibition;

photoprotection; photosynthetic light-response curves; shade tolerance; species interactions; tree diversity

Introduction

The sun's light powers photosynthesis—the chain of reactions through which plants turn carbon dioxide into the living matter they use to grow and reproduce. A common view in plant ecology is that light is often a limiting resource, such that competition for scarce light can govern the fate of plant species (Braun-Blanquet 1932; Canham et al. 1994; Monsi & Saeki 2005; Dybzinski & Tilman 2007; Hautier et al. 2009). But physiologists have also shown that plants often absorb light beyond their capacity to use it for photosynthesis (Long et al. 1994; Demmig-Adams & Adams 1992). Beyond an initial linear rise with increasing light, photosynthesis begins to saturate, and additional light contributes little to the plant's carbon gain. Any factor that limits photosynthesis can exacerbate the excess of light, including photosynthetic downregulation and environmental stresses like water limitation, cold temperatures, or nutrient-poor conditions (Fig. 1; Long et al. 1994; Demmig-Adams & Adams 1992).

Light is thus both an essential resource and a potential stressor. A chronic excess of light can cause plant cells lasting oxidative damage, especially to Photosystem II (PSII). Such damage reduces the efficiency of photosynthesis and can be costly to repair, but plants can avoid it using mechanisms of photoprotection (Murchie & Niyogi 2011). These mechanisms include biochemical pathways that safely dissipate excess light as heat—most notably, the xanthophyll cycle. They also include structural means to avoid absorbing too much light, such as self-shading canopies, reflective leaves, or steep leaf angles (Lovelock & Clough 1992; Streb et al. 1997; Pearcy et al. 2005; Kothari et al. 2018). In general, plants use such photoprotective mechanisms more under environmental conditions that put them at high risk of damage, including high light (Montgomery et al. 2008) and cold or dry climate (Cavender-Bares 2007; Savage et al. 2009; Wujeska et al. 2013; Ramírez-Valiente et al. 2013). Under such conditions, plants need strong photoprotection to prevent even more costly damage.

Depending on how well-protected a plant is, prolonged exposure to excess light can still cause enough damage to reduce carbon assimilation rates (Egerton et al. 2000; Howell et al. 2002; Murchie & Niyogi 2011). Consistent with Long et al. (1994), we use the term 'photoinhibition' to describe a drop in carbon assimilation caused by either photodamage or sustained biochemical photoprotection. Bright

sunlight causes photoinhibition directly by exposing leaves to more radiation, but it also acts indirectly by

altering other microclimatic conditions like temperature, soil moisture, and vapor pressure deficit (VPD;

Björkman & Powles 1984). For example, high VPD can cause stomata to close, which limits the use of

light for photosynthesis; without an energetic sink, the resulting excess of light can then cause

photoinhibition, which limits photosynthetic capacity even further. Alternately, limitation in water or

other resources may arrest tissue expansion and cause carbon sink limitation (Tardieu et al. 2014); in

response, plants may downregulate photosynthesis and initiate sustained photoprotection (Adams III et al.

2013). Such scenarios may be worsened by belowground competition for resources. Because such stresses

vary along environmental gradients, photoinhibition and photoprotective strategies could affect the

distribution and persistence of plant populations (Külheim et al. 2002).

Given that excess light can cause stress, it may be that one plant species can facilitate another

through shading. Whether a plant receives net costs or benefits from its neighbors' shade depends on the

net balance resulting from (1) the increasing frequency and severity of light limitation and (2) the

avoidance of chronic light stress. We can describe these effects in the terms of Monteith et al. (1977),

who defined light-use efficiency (LUE) as net photosynthesis divided by the amount of photosynthetically

active radiation (PAR) absorbed. Consider a typical light-response curve that plots photosynthesis (y-axis)

against PAR (x-axis; Fig. 1). Chronic photoinhibitory stress may change the shape of the curve by

lowering LUE, and thus photosynthesis, at any given amount of PAR. By relieving photoinhibition,

shading may raise LUE. But shading would also move the x-coordinate leftward along the curve at any

given moment, raising the chance the leaf will be in the curve's light-limited portion. Whether shade

helps or hurts depends on whether, integrated across time, the gain of avoiding chronic stress exceeds the

loss caused by potential light limitation.

The species interactions that result from shading could help to explain the results of experiments

that manipulate biodiversity, including tree diversity. Experimental and observational research alike has

shown that biodiversity often promotes ecosystem functions such as productivity (Tilman et al. 2014). In

forests, this trend has been shown in both experimental (Potvin & Gotelli 2008; Tobner et al. 2016;

Grossman et al. 2017; Williams et al. 2017; Huang et al. 2018; Zemp et al. 2019) and observational

(Gamfeldt et al. 2013; Liang et al. 2016; Oehri et al. 2017) data, across temperate (Tobner et al. 2016;

Grossman et al. 2017; Williams et al. 2017), subtropical (Huang et al. 2018), and tropical (Potvin &

Gotelli 2008; Zemp et al. 2019) biomes. These positive relationships are robust to variation in climate,

successional stage, and other factors (reviewed in Grossman et al. 2018; Ammer 2018).

One of the potential drivers of positive biodiversity-productivity relationships is interspecific

facilitation (Wright et al. 2017; Barry et al. 2018). In biodiversity experiments, patterns of productivity

within species or functional groups can sometimes serve as evidence for facilitation (Mulder et al. 2001;

Fichtner et al. 2017), but they cannot tell us the physiological mechanisms that cause one species to

facilitate another. Ecologists may need to understand these mechanisms to forecast species interactions

and productivity under environmental change (Wright et al. 2017; Barry et al. 2018).

Here, we tested whether the shade created by tree crowns can facilitate the carbon assimilation

and growth of neighboring trees in the Forests and Biodiversity (FAB) experiment at Cedar Creek

Ecosystem Science Reserve (East Bethel, MN). We asked: Do some species aid others by shielding them

from stress caused by intense sunlight? In particular, we sought to test whether slow-growing, shade-

tolerant broadleaf trees might be facilitated by adjacent fast-growing conifers (Table 1). We addressed

this question by measuring the physiology and woody growth of broadleaf species across plots that vary

in species composition. If certain species do benefit from the shade of their neighbors, we would expect

individuals with larger neighbors to have higher growth and carbon assimilation rates. We also posed the

hypothesis that trees exposed to high light would use structural and biochemical photoprotective

mechanisms more than trees shaded by their neighbors. If true, this finding would imply that trees

growing in high light must invest in photoprotection to avoid damage, reinforcing the claim that light

stress influences the interactions within these communities.

Methods

Experimental design

The FAB experiment was planted in 2013 at Cedar Creek Ecosystem Science Reserve in central

Minnesota, USA. Cedar Creek has a continental climate with cold winters and warm summers; the mean

annual temperature is about 7 °C and the mean annual precipitation is about 660 mm. The site lies on the

Anoka Sand Plain, where upland regions tend to have well-drained, sandy soils (Johnston et al. 1996).

FAB comprises 142 4 × 4 m plots, each planted with 64 trees in a 0.5 × 0.5 m grid. The

experiment has three blocks, each with 49 plots arranged in a 7 × 7 square. Neighboring plots within a

block share a boundary. Each plot is planted with one- or two-year-old seedlings of 1, 2, 5, or 12 species

from a pool of 12 species total, including four evergreen conifers with needle-like leaves and eight

winter-deciduous angiosperms with broad leaves (Table 1). Except for five-species plots, each distinct

species composition is replicated in each block. All species in a plot have nearly equal frequency and are

placed at random. Grossman et al. (2017) describes the experimental design in greater detail. The mean

relative growth rate in monoculture during the first three years varied dramatically among species—it was

highest in conifers and *Betula papyrifera*, and lowest in *Acer negundo* (Table 1).

Tree growth

Within each species, we sought to determine how an individual's stem growth rates were

influenced by the size of its neighbors. We assumed that neighbors' size affects the focal individual's

growth rates, but not vice versa. This assumption of exogeneity was warranted because nearly all of the

cross-species variation in growth rates is explained by species identity, so the size of a focal individual's

neighbors is explained much more by their identity than by their interactions with the focal individual.

We surveyed tree growth in late fall of each year. For all living trees, we measured diameter and

height to the tallest leader. We measured basal diameter (5 cm from the ground) for trees less than 1.37 m

tall and diameter at breast height for trees more than 1.37 m. For the first year that a tree crossed the 1.37

m threshold, we measured both diameters. To predict a tree's basal diameter when it was not measured,

we used species-specific linear relationships ($R^2 = 0.634$–0.784) predicting basal diameter additively from

height in the same year and the basal diameter in 2016, which was the last year that the basal diameter of

all trees was measured. For the conifer *Pinus banksiana*, basal diameter was measured on few enough

trees in 2018 that we could not apply this approach; we instead estimated stem volume under the

assumption that relative growth rate (RGR) was the same from 2017 to 2018 as from 2016 to 2017.

Because we did not consider *Pinus banksiana* growth as a response variable, this approach avoided

circularity and allowed us to get reasonable estimates of size in 2018.

In all other cases, we estimated woody volume as $V = \pi r^2 h$, where h is height and r is the basal

radius. This equation assumes each tree's woody volume can be approximated as a cylinder, an

assumption that has been used and justified in other tree diversity studies in the absence of system-

specific allometric equations (Tobner et al. 2016; Williams et al. 2017). We then calculated the RGR (yr$^-$

^{1}) between 2016 and 2018 as $(\ln V_{2018} - \ln V_{2016})/2$ (Hoffman & Poorter 2002). We chose 2016 as a

starting point because no replanting of dead trees occurred after that point. Volume loss could cause RGR

to be negative (Mahmoud & Grime 1974), which often occurred when stems or branches broke, or

occasionally when plants lost all aboveground tissue and resprouted the following year.

For each tree, we calculated the average woody volume of its eight neighbors (in cardinal and

intercardinal directions) in 2018 as a proxy for the intensity of aboveground interactions. Neighbors were

sometimes missing because of mortality or because the focal tree was on the experiment's edge. In such

cases, we assigned the neighbor a volume of zero. For each broadleaf species, we tested how individual

growth changed as average neighbor size increased using a mixed-effects regression model with plot as a

random intercept, implemented in *R v. 3.6.1* (R Core Team 2019) using package *lme4 v. 1.1.2* (Bates et al.

2015). We calculated the marginal R^2 ($R_m{}^2$) of the fixed effect using the method of Nakagawa et al. (2017)

as implemented in *MuMIn v. 1.43.17* (Bartoń 2020), and *p*-values using Satterthwaite's degrees of

freedom method as implemented in *lmerTest v. 3.1.2* (Kuznetsova et al. 2017), noting that such values

should be interpreted conservatively. We also tested whether any species showed non-monotonic

responses to neighbor size using the two-lines test (Simonsohn 2018), without random effects.

Some species showed a positive response in stem growth to neighbor size. Besides facilitation,

such a response could also be caused by shade avoidance, which may cause plants to increase shoot

biomass at the expense of root biomass in order to compete for light (Shipley & Meziane 2002). We

aimed to discern whether such a shade avoidance syndrome could contribute to the positive stem growth

response. Lacking root biomass sorted by species, we could not test this idea directly, but we could test

another symptom of a typical shade avoidance syndrome: stem elongation, which prioritizes vertical

growth over lateral growth (Henry & Aarssen 1999). We considered height and basal diameter in 2016,

the last year in which basal diameter was measured directly on all trees. For all species that showed a

positive stem growth response, these two variables were linearly correlated. We used mixed-effects linear

models with plot as a random intercept to predict diameter from height in each species, then tested

whether the residuals—the observed diameters' deviations from the predictions for a tree of equal

height—were associated with average neighbor size. If trees with larger neighbors show a shade

avoidance syndrome, we would expect them to have smaller diameters for a given height. We obtained

similar conclusions when considering the log-transformed diameter-to-height ratio as the response

variable rather than the residuals of the linear diameter-height relationship.

Photosynthetic physiology

We measured chlorophyll fluorescence parameters and leaf reflectance spectra among all eight

broadleaf species, and photosynthetic light-response curves among four focal species (*A. rubrum*, *B.*

papyrifera, *Q. ellipsoidalis*, and *T. americana*). For physiological measurements, we focused on plots

belonging to one of three treatments: (1) relatively open monocultures of broadleaf species

('monoculture'); (2) bicultures comprising one broadleaf and one conifer species ('shaded biculture'); and

(3) twelve-species plots (see Fig. S1 for sample images). For each species in each treatment, we measured

physiological parameters on six individuals—two in each of three plots. Neither *A. negundo* nor *Q.*

macrocarpa was planted with a conifer in any biculture. We used *B. papyrifera* as a shaded biculture

partner for *Q. macrocarpa* because it is a fast-growing species and creates shade. We omitted the shaded

biculture treatment for *A. negundo*. When possible, we chose the same trees within a plot for each kind of

physiological measurement so that we could compare these aspects of physiology within individuals.

Photosynthetic light-response curves

We measured photosynthetic light-response curves from the four focal species during July 2018

using an LI-6400 gas exchange system (LI-COR BioSciences, Lincoln, NE, USA) with a 6400-40 Leaf

Chamber Fluorometer head. From each tree ($n = 72$ total), we selected a fully expanded upper leaf with

no visible sign of disease or herbivory. We noted the angle of each leaf relative to horizontal (to the

nearest 15˚) before beginning measurements. Each curve had nine steps in descending order of brightness:

2000, 1500, 1000, 500, 200, 100, 50, 20, and 0 µmol m^{-2} s $^{-1}$. We also measured the relative electron

transport rate (ETR) using chlorophyll fluorescence at each light level. We maintained favorable

conditions inside the chamber during each measurement. Following each curve, we removed the leaf to

measure leaf mass per area (LMA) using a balance and a flatbed scanner. Further details about light-

response curve procedures and ETR calculations can be found in the Supplemental Materials (Appendix

1). Finally, we estimated parameters like the light-saturated carbon assimilation rate (A_{sat}) by fitting a

non-rectangular hyperbolic model (Johnson & Thornley 1984) to each light-response curve using *R* code

written by Nick Tomeo, available at: https://github.com/Tomeopaste/AQ_curves.

We aimed to determine how realized carbon assimilation rates vary with the photosynthetic

photon flux density (PPFD) at the top of the canopy, a measure of light availability in the absence of

shading. While all trees received the same range of light intensities inside the instrument chamber, some

trees received much less light than others *in situ* because their neighbors shaded them more. At any

moment, trees that are more shaded are farther to the left on their light-response curve than trees that are

less. For example, consider a tree crown whose topmost leaves only receive 50% of the light reaching the

top of the canopy. For this tree crown, the realized assimilation rate we estimate at a top of the canopy

PPFD of 2000 µmol photons m^{-2} s $^{-1}$ would be the assimilation rate at a leaf-level PPFD of 1000 µmol m^{-2}

s $^{-1}$. To visualize the impacts of shading by neighbors, we followed the approach of Howell et al. (2002):

we estimated how much light each leaf receives as a constant fraction of light at the top of the canopy,

and rescaled each leaf's fitted light-response curve along the *x*-axis by this fraction's reciprocal. We

estimated the constant rescaling factor for each individual as *RLA* $\times$ cos*(θ)*, where *RLA* is relative light

availability (see *Light availability*) and θ is the leaf angle from horizontal. (Taking the cosine of θ

approximately corrects for the reduction in horizontally projected leaf area in steeply inclined leaves

when the sun is directly overhead, as simplified from Ehleringer & Werk [1986].) We used this procedure

on each fitted curve in order to compare across treatments in a way that accounts for the varying fractions

of light that individuals receive due to differences in relative height and vertical light transmission.

Finally, we used the rescaled light-response curves to estimate carbon assimilation by top leaves

throughout July. We used an hourly averaged time series of solar radiation from Carlos Avery Wildlife

Management Area in Columbus, MN, 13.6 km away from the study site. Following Udo and Aro (1999),

we assumed a conversion rate of 2.08 μmol photons m^{-2} s^{-1} of PAR per Watt m^{-2} of solar radiation. By

using these data as inputs to the rescaled light-response curves, we estimated carbon assimilation over the

month of July. This procedure assumes that the photosynthetic response to light remains constant and

unaffected by factors like leaf temperature and stomatal closure. While this assumption is never quite

true, we consider it a useful way of estimating the consequences of realistic fluctuations in light.

We mainly express assimilation per unit of leaf dry mass because we aim to determine the plants'

return on the carbon invested in leaf construction. As expected, leaves growing in low light tended to

have lower LMA in most species (Fig. S2; Poorter et al. 2019; Williams et al. 2020). More carbon is spent

on constructing a leaf with high LMA, which means that the leaf has to assimilate more carbon to recoup

its construction cost. Expressing data on a mass basis accounts for these large differences in construction

cost, although we also discuss area-based assimilation rates when they provide additional physiological

insight. We also express stomatal conductance (g_s) and electron transport rate (ETR) on a mass basis to

compare them with mass-based assimilation rates.

Instantaneous chlorophyll fluorescence and spectral reflectance

In all eight broadleaf species, we measured chlorophyll fluorescence parameters using an FMS2

pulse-modulated fluorometer (Hansatech Instruments Ltd., Norfolk, UK) over two days in late July ($n =$

138 total). We measured dark- and light-acclimated parameters at the same spot on the same leaves. We

attached opaque clips to leaves in the evening before measuring dark-acclimated F_v/F_m within two hours

after sunrise. This parameter describes the maximum quantum yield of PSII and is a general index of

photoinhibition (Murchie & Lawson 2013). We then took light-acclimated measurements between 12:00

and 14:00 each day. The protocol involved the following steps: actinic light at 1000 µmol m^{-2} s^{-1} for 15

seconds, a saturating pulse, and two seconds of far-red light. We exposed all leaves to the same actinic

light because we aimed to assess photoprotective capacity under comparable light conditions, even though

each tree had a different light environment.

From these data, we estimated qN, a parameter that indicates how much a plant relies on

photoprotective dissipation (non-photochemical quenching) under the imposed actinic light. This

parameter is correlated with the de-epoxidation state of xanthophyll cycle pigments (Watling et al. 1997;

Cavender-Bares & Bazzaz 2004). Following Kramer et al. (2004), we also calculated the quantum yields

ϕ_{PSII}, ϕ_{NPQ}, and ϕ_{NO}, which sum to 1. These three parameters are the proportions of light energy dissipated

by photosynthetic chemistry, non-photochemical quenching, and non-regulated dissipation. The last is a

crucial parameter because it represents light that a plant cannot dissipate safely; this portion of the

absorbed light may contribute to photodamage. The Supplemental Materials contain formulas for all

parameters and justifications for our choices in data analysis (Appendix 1).

On separate top leaves of the same trees, we measured reflectance spectra (350-2500 nm) using a

PSR+ 3500 field spectroradiometer (Spectral Evolution, Lawrence, MA, USA). We used these spectra to

calculate the photochemical reflectance index (PRI), calculated as PRI = $(R_{531} - R_{570}) / (R_{531} + R_{570})$,

where R_n is the reflectance at a wavelength of n nm. PRI shows a negative correlation with carotenoid :

chlorophyll ratios and, over shorter time scales, also tracks the xanthophyll de-epoxidation state (Gamon

et al. 1992; Wong & Gamon 2015; Gitelson et al. 2017). We estimated species-specific linear responses

of PRI and chlorophyll fluorescence parameters to relative light availability (see *Light availability*).

To understand how photoinhibition and photoprotective strategies vary with adaptations to shade,

we used shade tolerance values that Niinemets & Valladares (2006) compiled on a five-point scale.

Higher values along this scale indicate that plants are able to grow in lower light conditions.

Phenology

We monitored the timing of leaf abscission for *T. americana* in monoculture and shaded biculture

plots to assess whether shade could delay senescence and abscission, extending the period for carbon gain

(Cavender-Bares et al. 2000). In early August, we surveyed 120 *T. americana* trees (60 in monoculture,

60 in shaded biculture) to determine the proportion of leaves that had fallen based on how many of the top

five axillary buds lacked a leaf. From each of these trees, we marked a single leaf and returned every 1-2

weeks to monitor its senescence. We assessed each leaf on a binary basis (senesced or not) based on

whether it had at least 50% remaining green leaf area. We chose this criterion as a simple proxy for the

continuous scale used by Cavender-Bares et al. (2000). Based on the initial survey and the repeated

surveys of marked leaves, we tracked the proportion of leaves that had senesced over time. To check

whether leaves were photosynthetically active until near the point of senescence, we collected a one-time

measurement of dark-acclimated F_v/F_m in mid-September among remaining leaves.

We also performed a one-time measurement of 60 (30 per treatment) *A. rubrum* plants on October

1, using the same protocol we used to do our initial early August survey of *T. americana*. We aimed for

this survey to help test whether our results hold across species.

Water potential

Water deficits—even moderate ones—can arrest tissue growth (Tardieu et al. 2014) and reduce

carbon gain by causing stomata to close (Brodribb et al. 2003). To see how our treatments affected water

status, we measured leaf water potential using a Scholander pressure bomb. We measured pre-dawn water

potential (Ψ_{PD}) in all eight broadleaf species ($n = 96$) and midday water potential (Ψ_{MD}; between 12:00

and 14:00) in only the four focal species ($n = 48$). We included monoculture and shaded biculture

treatments, except in *A. negundo*, where we used the twelve-species treatment in place of the absent

shaded biculture treatment. We removed leaves and immediately placed them in impermeable plastic bags

with wet paper towels. We carried leaves in coolers to be measured indoors within 90 minutes.

Because we only intended to test whether there was a general tendency across species for water

potential to vary with light, we included species as a random intercept and light availability as a fixed

effect in mixed-effect regression models for both Ψ_{PD} and Ψ_{MD}. We calculated R_m^2 and *p*-values as in

models for growth rate.

Leaf angles

Leaf angle is a key control on light exposure because more vertical leaves intercept less light

when the sun is directly overhead, and may allow light to penetrate deeper into the canopy (Posada et al.

2009). As a result, steep leaf angles can be a structural strategy to avoid light, temperature, and water

stress (Valladares & Pugnaire 1999; Werner et al. 1999; Pastenes et al. 2004; Valiente-Banuet et al.

2010). We predicted that trees in monoculture would have steeper leaf angles than those in other

treatments. From each of three species (*A. rubrum*, *Q. ellipsoidalis*, and *T. americana*), we randomly

selected five individuals from each of six plots—three from monoculture treatments, three from shaded

bicultures ($n = 15$ per species × treatment). On these individuals, we measured the leaf angle from

horizontal of the top five leaves in the crown to the nearest 15˚.

Because angles of leaves within individuals may be non-independent, we tested whether

treatments differ using a mixed-effects model with treatment as a fixed effect and individual tree as a

random intercept. Deviations from horizontal in either direction can reduce light interception, so we used

the absolute value of leaf angle as the dependent variable. We again calculated *p*-values using

Satterthwaite's degrees of freedom method.

Environmental data

Light availability

On a cloudy day with diffuse light conditions, we measured the available light above each tree

selected for this study ($n = 138$) using an AccuPAR LP-80 ceptometer (METER Group, Pullman, WA,

USA). We took the mean of two to four PAR measurements in the open and the mean of two

measurements directly above the topmost leaf of each tree. By calculating a ratio of these values, we

could calculate the percent of light transmitted to the top of each tree. We called this value *relative light*

availability (RLA). This value usually correlates well with the percentage of light a tree can access over

much longer time-scales (Parent & Messier 1996).

Soil water content

We collected a one-time measurement of soil volumetric water content in mid-August using a

FieldScout Time-Domain Reflectometer 350 (Spectrum Technologies, Inc., Aurora, IL, USA). This single

measurement followed six days from most recent rain. At midday, we collected and averaged three

measurements in each of 32 plots, spanning monocultures of the four core broadleaf species, shaded

bicultures, and twelve-species plots.

Air temperature

For nine days during early September, we set out four Thermochron iButtons (model DS1921G;

Maxim Integrated Products, Sunnyvale, CA, USA) in shaded plots and two in twelve-species plots,

setting them to log air temperature hourly. Each iButton was suspended on mesh inside a 1 m-tall PVC

pipe that was capped with an elbow to shield against solar irradiance. We compared these data to air

temperature measured hourly under open, sunny conditions at the Cedar Creek LTER weather station

about 0.77 km away. We assumed that data from this station would be representative of air temperature in

non-shaded areas. We ignored nighttime and early morning (20:00-08:00 h) readings because they

seemed to be influenced by water condensation and evaporation.

Results

Tree growth

Species varied in their relationship between neighbor stem volume and focal individual relative growth rate (RGR; mixed-effects ANCOVA, neighbor volume $\times$ species interaction; $p < 10^{-15}$; $F_{11,7279} = 12.440$). For individual species, relationships between RGR and neighbor volume were noisy but often highly significant (Fig. 2). In most species, individuals with larger neighbors had lower RGR, including in *A. rubrum* ($R_m^2 = 0.027$, $p < 0.005$, $t(366) = -3.257$), *Q. alba* ($R_m^2 = 0.030$, $p < 10^{-4}$, $t(486) = -3.932$), *Q. ellipsoidalis* ($R_m^2 = 0.051$, $p < 10^{-7}$, $t(341) = -5.485$), *Q. macrocarpa* ($R_m^2 = 0.006$, $p = 0.0380$, $t(697) = -2.079$), and *Q. rubra* ($R_m^2 = 0.053$, $p < 10^{-7}$, $t(444) = -5.785$). But *B. papyrifera* had a weak positive response to neighbor size ($R_m^2 = 0.010$, $p = 0.049$, $t(124) = 1.993$), and two more species grew faster with larger neighbors, but only when average neighbor size was log-transformed: *A. negundo* ($R_m^2 = 0.014$, $p = 0.019$, $t(380) = 2.361$) and *T. americana* ($R_m^2 = 0.018$, $p < 0.005$, $t(350) = 2.154$). These results are qualitatively unchanged by removing trees on a block edge or including focal plant volume in 2016 as a covariate, although the effect of neighbor size in *B. papyrifera* or *Q. macrocarpa* sometimes became statistically insignificant under these alternate model specifications (Supplemental Materials, Appendix 2). No species had a non-monotonic response to neighbor size.

Trees that died between 2016 and 2018 dropped out of our individual-level analyses of RGR because their stem volume was 0 in 2018, making it impossible to calculate RGR. To see whether mortality could alter our conclusions, we also performed statistical analyses on absolute change in stem volume from 2017 to 2018, in which we alternately (1) treated mortality as a decline in volume to 0, or (2) removed trees that died from analyses. The rate of mortality was low between 2016 and 2018 (~7.5%), and accounting for mortality makes little qualitative difference when considering absolute growth rate as the response variable (Fig. S3; Supplemental Materials, Appendix 2).

Although the individual-level effects are noisy, they reveal strong tendencies in average stem growth across the full range of neighbor size. Imagine two *T. americana* trees that begin at the same size: One in monoculture, where the average neighbor in 2018 is 370 cm^3, and one growing in a shaded

biculture with *Pinus strobus*, where the average neighbor is 2120 cm^3. The individual-level regression

(with log-transformation) predicts that after two years, the average *T. americana* in the shaded biculture

would be 26.7% larger because of the difference in mean RGR. Or consider *Q. rubra*, which had the

strongest negative response to neighbor size: We predict that an average tree in the shaded biculture

(mean neighbor size 1988 cm^3) would be 40.2% smaller than in monoculture (mean neighbor size 160

cm^3). These figures assume that individuals across treatments start at the same size at the initial point for

RGR estimates, so they may underrepresent the true differences in size that have developed. The absolute

growth rate gives some insight into these differences. For example, the median *T. americana* stem in

monoculture grew 99 cm^3 (25–75th percentile: 21–237 cm^3) between 2017 and 2018, compared to 267

cm^3 (29–633 cm^3) in shaded biculture with *P. strobus* or *J. virginiana*. In comparison, the median *Q.*

rubra stem grew 32 cm^3 (25–75th percentile: 3.9–124 cm^3) in monoculture, compared to only 12 cm^3 (-

5.2–89 cm^3) in shaded biculture with *P. strobus*. In general, trends in the absolute growth rate across the

gradient of neighbor size mirrored those in the relative growth rate (Supplemental Materials, Appendix 2).

We can also aggregate these observations to the plot scale—for example, by considering the RGR

of summed stem volume of each species in each plot from 2016 to 2018 (Fig. 2). (This approach is

distinct from an average of individual RGRs, which assigns equal weight to small and large individuals.)

Although aggregation leaves less statistical power to detect relationships, it also reduces noise, such that

the relationships we find explain much more of the variation in growth across plots. As average neighbor

size increases, RGR declines in *Q. alba* ($R^2 = 0.297$, $p < 0.005$, $t(28) = -3.642$) and *Q. macrocarpa* ($R^2 =$

0.265, $p < 0.005$, $t(31) = -3.540$), but increases in *T. americana* ($R^2 = 0.312$, $p < 0.001$, $t(32) = 3.995$).

Finally, for the three species that showed positive individual-level stem growth responses to

neighbor size (*T. americana*, *A. negundo*, and *B. papyrifera*), we performed a follow-up analysis to test

whether they showed a shade avoidance response. We found that trees with larger neighbors had smaller

diameters than expected based on their height in *B. papyrifera* ($R^2 = 0.009$, $p = 0.008$, $t(795) = -2.645$),

but not *T. americana* and *A. negundo* ($p > 0.05$).

    Photosynthetic physiology

    Photosynthetic light-response curves

As a function of the chamber light level, mass-based assimilation rates were always higher in

shaded biculture and twelve-species treatments than in monoculture in three out of the four focal species

(*Q. ellipsoidalis*, *A. rubrum*, and *T. americana*; Fig. 3, left). In the early-successional species *B.*

papyrifera, mass-based assimilation rates in shaded biculture were lower than in the other two treatments.

Area-based assimilation rates showed similar trends in *T. americana*, but varied less among treatments in

B. papyrifera and *Q. ellipsoidalis*—and in *A. rubrum*, they were lower in shaded biculture than the other

treatments (Fig. S4) as a result of large differences among treatments in LMA (Fig. S2).

In all broadleaf species except *B. papyrifera* and *T. americana*, LMA was higher in the

monoculture than the other two treatments (Fig. S2). Across the four focal species, the light-saturated

assimilation rate A_{sat} on an area-basis was positively correlated with LMA ($R^2 = 0.431$; $p < 10^{-9}$; $t(69) =$

7.349). Within species, there was no correlation in *B. papyrifera* and *A. rubrum*, but a positive correlation

in *Q. ellipsoidalis* ($R^2 = 0.192$; $p = 0.039$; $t(16) = 2.243$) and *T. americana* ($R^2 = 0.436$; $p = 0.002$; $t(16) =$

3.759). In *T. americana*, the intercept was statistically indistinguishable from zero, making the

relationship between area-based A_{sat} and LMA close to a direct proportionality.

The rate of photosynthesis is often limited either by ETR or by the RuBP carboxylation rate; the

latter may in turn be limited by stomatal diffusion of CO_2 (Farquhar et al. 1980). In our data, both ETR

and stomatal conductance (g_s) increased with light availability (Fig. S5). (ETR tended to decline at very

high light levels, especially in shaded treatments, perhaps because of acute photoinhibition.) Within each

species, the rank-order of treatments in carbon assimilation rates, ETR, and g_s were broadly congruent.

One exception is that *T. americana* in shaded biculture had much higher g_s than in twelve-species or

monoculture plots, despite having slightly lower assimilation rates than in twelve-species plots.

Compared to the unscaled chamber light-response curves, the picture that emerged from rescaled

light-response curves is more complex (Fig. 3, right). In *B. papyrifera*, the assimilation rate was still

lowest across light levels in the shaded biculture treatment. In late-successional *T. americana*, the

assimilation rate was higher in shaded biculture and twelve-species treatments by up to 25% compared to

monoculture across most light levels. In the other two species, the mass-based assimilation rate was

highest in monoculture at low light, but twelve-species and (in *Q. ellipsoidalis*) shaded biculture plots

intersected and surpassed monocultures when enough light was available. *A. rubrum* and *Q. ellipsoidalis*

grew less and had lower light availability in shaded bicultures than *T. americana* (Fig. S2), so their mass-

based assimilation rates dropped considerably more in the rescaled light-response curves. The area-based

assimilation rate in these two species was lower in shaded bicultures (and in *Q. ellipsoidalis*, twelve-

species plots) than in monocultures across the full domain of light availability (Fig. S4), but otherwise,

mass-based and area-based rates showed similar patterns across treatments within species.

454   Using a time series of solar radiation, we estimated total mass-based assimilation rates in July

2018 based on each rescaled light-response curve (Fig. S6). We found that trees would have lower total

assimilation in shaded biculture than in monoculture in *A. rubrum* (ANOVA; $p = 0.009$; $F_{2,14} = 6.759$) and

B. papyrifera ($p = 0.002$; $F_{2,15} = 10.27$). In both species, Tukey's HSD test showed that trees in twelve-

species plots did not significantly differ from those in monoculture. There were no significant differences

at all among treatments in *Q. ellipsoidalis* ($p = 0.294$; $F_{2,15} = 1.330$) or *T. americana* ($p = 0.359$; $F_{2,15} = $

1.099). Considering area-based assimilation, we see similar results, except that there are no significant

differences among treatments in *B. papyrifera* ($p = 0.0892$; $F_{2,15} = 2.852$), only in *A. rubrum*.

Instantaneous chlorophyll fluorescence and spectral reflectance

464   Among all eight species except *B. papyrifera*, dark-acclimated F_v/F_m declined as relative light

availability (RLA) increased (Fig. 4). Extracting the species-specific slopes of this relationship, we found

that species with high shade tolerance (as quantified by Niinemets & Valladares 2006) had the greatest

decline in F_v/F_m as RLA increased (Fig. 5; $R^2 = 0.601$; $p = 0.025$; $t(5) = -3.167$). In most species, non-

photochemical quenching (qN) rose with RLA, but in *T. americana* and *A. negundo*, qN was nearly

constant across light environments (Fig. 4). Consequently, we found that shade-tolerant species also had

smaller rises in qN with RLA (Fig. 5; $R^2 = 0.619$; $p = 0.022$; $t(5) = -3.276$). PRI declined as RLA

increased, with statistically indistinguishable slopes among species (Fig. 4).

To see why shade-tolerant species showed such large increases in photoinhibition with light,

consider ϕ_{PSII}, ϕ_{NPQ}, and ϕ_{NO}—the quantum yields of PSII photochemistry, non-photochemical quenching,

and non-regulated dissipation. In most species, ϕ_{NO} decreased with light in the growth environment (Fig.

4)—for trees growing in the open (RLA $\approx$ 1), the sum $\phi_{PSII} + \phi_{NPQ}$ at 1000 µmol photons m^{-2} s^{-1} was

usually above 0.8, keeping ϕ_{NO} close to the oft-cited theoretical minimum of ~0.17 (Tietz et al. 2017; Fig.

S7). But in *T. americana* and *A. negundo*, ϕ_{NO} was nearly equal across the gradient of light availability.

To illustrate how photoinhibition affected photosynthetic function, we examined the relationship

between dark-acclimated F_v/F_m and A_{sat}. Among the four focal species, dark-acclimated F_v/F_m was

positively correlated with mass-based A_{sat} (OLS regression; $R^2 = 0.493$; $p < 10^{-11}$; $t(69) = 8.304$; Fig. 5),

and slopes were not significantly different among species (ANCOVA, species $\times$ dark-acclimated F_v/F_m

interaction; $p = 0.658$; $F_{3,63} = 0.539$). The same overall relationship holds when A_{sat} is expressed on an

area basis ($R^2 = 0.447$; $p < 10^{-9}$; $t(69) = 7.588$), although within species, *A. rubrum* and *B. papyrifera* had

no significant relationship.

One way of disentangling the role of photoinhibition from leaf plasticity in influencing

photosynthetic rates is to consider the residuals of the species-specific relationships between area-based

A_{sat} and LMA. These residuals quantify how much greater or lesser A_{sat} is than predicted based on LMA

alone for a given species. Dark-acclimated F_v/F_m showed a positive correlation with these residuals in *Q.*

ellipsoidalis ($R^2 = 0.351$; $p = 0.006$; $t(16) = 3.189$) and *T. americana* ($R^2 = 0.368$; $p = 0.004$; $t(16) =$

3.304). In these two species, a rise of 0.01 in dark-acclimated F_v/F_m would be predicted to cause a rise of

1.41 and 0.85 µmol m^{-2} s^{-1} in carbon assimilation, respectively. We found no such relationship in *B.*

papyrifera or in *A. rubrum*, although in *A. rubrum* alone, dark-acclimated F_v/F_m showed a strong negative

correlation with LMA ($R^2 = 0.460$; $p = 0.002$; $t(15) = -3.826$). This multicollinearity makes it hard to

estimate the true contribution of either variable to area-based A_{sat}, and leaves open the possibility that a

potential positive effect of LMA on A_{sat} is offset by photoinhibition as light availability increases.

Phenology

In *T. americana*, leaf abscission in monoculture began in early August, and more than half of all

leaves had senesced by September 3 (Fig. 6). In shaded bicultures, more than 90% of leaves remained by

September 3, and no survey found greater than 50% senescence until October 1. We assigned the

senescence date of each leaf as the date of the first survey by which it had senesced; using this response

variable, leaves in monoculture senesced 22 days earlier than those in shaded bicultures (t-test; $p < 10^{-11}$;

$t(116) = 7.822$). In mid-September, dark-acclimated F_v/F_m averaged 0.745 among remaining leaves in

shaded biculture, but only 0.642 in monoculture (t-test; $p = 0.061$; $t(33) = 1.936$).

Our one-time measurement of *A. rubrum* leaves in early October confirmed that this pattern is not

limited to *T. americana*. *A. rubrum* trees in shaded biculture plots retained about 56% of their leaves,

while those in monoculture retained only 10% (t-test; $p < 10^{-4}$; $t(44) = -4.486$).

Water potential

We removed two data points, one in each treatment, with Ψ_{PD} more negative than -0.8 MPa on the

grounds that they may have been subject to measurement error. Across treatments, the mean Ψ_{PD} and Ψ_{MD}

were -0.15 and -1.29 MPa. Ψ_{PD} increased with light availability ($R_m^2 = 0.272$; $p < 10^{-5}$; $t(70.0) = -5.152$),

making it less negative in monocultures than shaded bicultures by an average of 0.07 MPa (Fig. S8). This

result holds even including the outliers ($R_m^2 = 0.085$; $p = 0.014$; $t(54.3) = -2.555$). By midday, the pattern

reversed; Ψ_{MD} declined with light availability ($R_m^2 = 0.346$; $p < 10^{-4}$; $t(36.6) = 4.450$), making it more

negative in monocultures than shaded bicultures by an average of 0.33 MPa.

Leaf angles

Among each of the three species whose leaf angles we measured, leaves were nearer to horizontal

in the shaded biculture treatment (Fig. 7). The size of this fixed effect varied among species: 11.0˚ in *Q.*

ellipsoidalis ($p = 0.005$; $t(34.2) = -3.010$), 21.2˚ in *A. rubrum* ($p < 10^{-9}$; $t(108.4) = -6.807$), and 14.3˚ in *T.*

americana ($p = 0.003$; $t(124.5) = -3.040$). Because *T. americana* leaves usually drooped downward in

monoculture, leaf angles became less negative in the shade. In *Q. ellipsoidalis*, whose leaves were often

inclined upward in monoculture, leaf angles became less positive in the shade. In *A. rubrum*, leaf angles

were inclined both up and down in monoculture, but varied less in either direction in the shade.

Environmental factors

In our survey, soil moisture varied among treatments (ANOVA; $p < 0.01$; $F_{2,29} = 6.115$); it was

lowest (5.7%) in shaded biculture plots and highest (7.2%) in broadleaf monoculture plots. We found that

daytime air temperature in the open was higher than in shaded bicultures or twelve-species plots at 62.5%

of hourly log-points during the nine-day logging period. The air temperature in monocultures was higher

on average by 1.03 ˚C over this period.

Discussion

We investigated how species interactions in a tree diversity experiment might emerge from the

physiological responses of trees to the light environment created by their neighbors. We found that as

neighbor size increased, five species (*A. rubrum*, *Q. alba*, *Q. ellipsoidalis*, *Q. macrocarpa*, and *Q. rubra*)

had faster stem growth on average, while the remaining three species (*A. negundo*, *T. americana*, and *B.*

papyrifera) had slower stem growth. The five species with declining stem growth responses to neighbor

size were shade-intolerant, resistant to photoinhibition in high light, and tended to have lower carbon

assimilation in shaded bicultures. Two of the three species with increasing responses were shade-tolerant

and susceptible to photoinhibition. The divergent responses of these two groups of species seem driven in

part by their tolerance to excess light. The remaining species, *B. papyrifera*, had a positive response to

neighbor size despite lower photosynthetic rates. Here, we interpret these patterns further.

Competition and shade-intolerant species

Individuals of all four *Quercus* species and *A. rubrum* had lower stem growth when surrounded

by larger neighbors (Fig. 2). This growth pattern implies that neighborhood interactions were dominated

by competition, potentially both above- and belowground. (The fact that soil moisture was lowest in

shaded biculture plots suggests an important role for belowground competition.)

The physiological data can help us make sense of these growth patterns. We measured light-

response curves of two species in this group: *A. rubrum* and *Q. ellipsoidalis*. In the untransformed

chamber light-response curves, both had lower mass-based assimilation rates in monoculture than in the

other two treatments (Fig. 3, left). But on an area basis, *A. rubrum* had its lowest assimilation rates in the

shaded biculture treatments, and *Q. ellipsoidalis* was similar across all three treatments. The discrepancy

between area- and mass-based results is explained by variation in LMA across treatments (Fig. S2).

Multiple structural and physiological factors contribute to explaining variation in photosynthesis

across treatments. *Q. ellipsoidalis* and *T. americana* had positive relationships between LMA and area-

based A_{sat}, which may imply that much of the variation in LMA is driven by the mass of photosynthetic

tissue (Osnas et al. 2018). In contrast, *B. papyrifera* and *A. rubrum* had no correlation between LMA and

area-based A_{sat}, which may imply that much of the variation in LMA could be driven by the mass of non-

photosynthetic structural tissue. But for *A. rubrum* in particular, there was a strong negative correlation

between LMA and dark-acclimated F_v/F_m, two variables jointly affected by light availability (Fig. S2; Fig.

4). As a result, a potential positive relationship between LMA and A_{sat} in *A. rubrum* could have been

masked by photoinhibition. We also found evidence that photoinhibition lowers A_{sat} in *Q. ellipsoidalis*

and *T. americana*. Since photoinhibition occurs through damage or downregulation of PSII, it is usually

thought to depress photosynthesis by lowering ETR. But both ETR and stomatal conductance mirrored

assimilation across treatments in chamber light-response curves (Fig. S5), so both may have some role in

explaining why most species had lower assimilation in monoculture at a given chamber light level.

Rescaling the light-response curves allowed us to visualize the costs of shading—the fact that

some trees receive less light and face more frequent photosynthetic light limitation than others. Compared

to monoculture, *A. rubrum* and *Q. ellipsoidalis* in shaded biculture had lower mass-based carbon

assimilation under low light at the top of the canopy, but higher mass-based carbon assimilation under

high light (Fig. 3, right). For these two species, shading often appears severe enough to cause light

limitation of growth and photosynthesis, especially under otherwise low light.

Given that light availability fluctuates—diurnally, seasonally, and with cloud cover—the

treatment that performs best may also vary from moment to moment. We used a time-series of solar

radiation to estimate total carbon assimilation throughout July. We found that *A. rubrum* and (only on a

mass basis) *B. papyrifera* had lower carbon assimilation in shaded biculture than in the other treatments

(Fig. S6). PPFD was under 500 µmol m^{-2} s^{-1} more than 60% of the time during the month; it is under such

dim conditions that trees in monoculture have the greatest photosynthetic advantage over those in shaded

biculture, no matter what benefits shading may confer at sunnier times (Fig. 3; Fig. S4). Trees in twelve-

species plots had similar assimilation rates to those in monoculture in all species. The intermediate light

environment in twelve-species plots may have allowed trees to avoid photoinhibition without becoming

frequently light-limited (Fig. S2), although we caution that we only measured top leaves, so we cannot

directly compare whole-plant carbon gain.

Trees in FAB and elsewhere often vary in size by orders of magnitude, which foregrounds the

size-asymmetry of light competition. Research on light competition in biodiversity-ecosystem function

research underscores a tension between competitive imbalance caused by size-asymmetry and

competitive relaxation caused by light-use complementarity, in which species may partition their

exploitation of the light environment (Yachi & Loreau 2007; Sapijanskas et al. 2014; Williams et al.

2017). We leave the potential for light-use complementarity unaddressed, but we show that competitive

imbalance can suppress the photosynthesis and growth of species that are poorly adapted to shade. Such

imbalances could promote selection effects (*sensu* Loreau & Hector 2001), in which the most productive

species become even more dominant in mixtures.

Facilitation and shade-tolerant species

In *T. americana*, *A. negundo*, and *B. papyrifera*, having larger neighbors increases stem growth

(Fig. 2; Fig. S3). The former two are the most shade-tolerant broadleaf species in the experiment, while

the last is the most shade-intolerant. Moreover, these three species showed divergent photosynthetic

responses to the light environment, which suggests that different mechanisms may explain the positive

stem growth response in *B. papyrifera* than in the other two species.

We first discuss the positive trend in *T. americana* and *A. negundo* before returning to *B.*

papyrifera in the following section. In these two species, we interpret this trend mainly as a result of

facilitation, reinforcing prior results that showed that shade-tolerant species may often have facilitative

responses to being shaded (Montgomery et al. 2010). In particular, these two species' growth trends are

best fit by a model where neighbor size is log-transformed (Fig. 2, Appendix 2). Increasing neighbor size

thus has positive but diminishing marginal benefits. This outcome may result if neighbor size has a non-

linear influence on light availability, or if deep shade has escalating costs that begin to offset its benefits.

Here, the physiological data may help us explain the growth patterns. *T. americana* was among

the four focal species, and the only one of the four that had the lowest carbon assimilation rates in the

monoculture treatment across most light levels in both chamber and rescaled light-response curves (Fig.

3). The steep decline in dark-acclimated F_v/F_m with increasing light (Fig. 4) suggests that the low A_{sat} in

monoculture results from photoinhibition (Fig. 5). This pattern aligns with *T. americana*'s role as a late-

successional dominant species throughout much of its range (Braun 1950). Our finding that carbon gain

in *T. americana* can increase under shade is reinforced by Carter and Cavaleri (2018), who found that its

assimilation rates may increase down vertical gradients from the upper canopy to the sub-canopy.

Nevertheless, we found no significant differences among treatments in estimated assimilation across the

whole month of July (Fig. S6), mainly because under low light, monocultures still had higher assimilation

rates in the rescaled light-response curves. The amelioration of photoinhibition is very likely to contribute

to the positive effect of neighbor size on growth in *T. americana*, but it cannot fully explain it.

We next discuss two mechanisms of facilitation that could also contribute to positive growth

responses to neighbor size, but which analyses based on light-response curves do not account for:

Delayed senescence and microclimatic effects. Both may contribute to potential positive effects of

shading by neighbors in the species that show such responses.

T. americana in monoculture dropped its leaves nearly a month earlier into the fall compared to

the shaded biculture (Fig. 6). We also found that *A. rubrum* in monoculture had earlier leaf senescence.

Given that leaves remained photosynthetically active until shortly before abscission (as suggested by

Mattila et al. 2018), shade could allow these trees a longer period of carbon gain. This pattern admits at

least two explanations: (1) that plants in the sun are earlier to accumulate phenological cues for

senescence related to temperature or photoperiod; or (2) that abiotic stress, especially light and water

stress, accelerate senescence (Cavender-Bares et al. 2000; Estiarte & Peñuelas 2015; Brelsford et al.

2019), especially as the temperature declines, rendering leaves more susceptible to photodamage (Juvany

et al. 2013; Renner & Zohner 2019). Contrary to the first explanation, air temperatures in full sun were

higher than in shaded plots, suggesting that early leaf drop in monoculture was caused by stress. Without

measuring the timing of senescence in other species, we do not yet know how general this effect is.

Although we emphasize the role of light stress, other microclimatic factors could contribute to

variation in assimilation rates among treatments. For example, an alternate perspective on photoinhibition

is that it mainly results not from photodamage but from carbon sink limitation, which leads plants to

downregulate light capture and electron transport as a photoprotective response (Adams III et al. 2013).

From this perspective, photoinhibition seldom causes carbon limitation or reduces growth rates in nature;

rather, slow growth rates cause photoinhibition. A potential cause of sink limitation that could affect

monocultures of *T. americana* and *A. negundo* more than shaded bicultures is water limitation, which

could physically constrain the expansion of tissues (Tardieu et al. 2014). We found that leaf Ψ_{PD} was

slightly more negative in shaded bicultures, while Ψ_{MD} was more negative in monocultures (Fig. S8).

(These results are consistent with the idea that soil moisture is higher in low biomass plots, as we found,

while daytime VPD is lower in high biomass plots, as Wright et al. (2018) found on hot, dry days at

Cedar Creek.) If Ψ_{MD} responds in other tissues as in leaves, highly negative values could arrest growth in

monoculture, causing carbohydrates to accumulate in leaves under high light.

But from this perspective, it is hard to explain why this sink limitation would particularly matter

in *T. americana* and *B. papyrifera* among the four focal species; for example, *A. rubrum* also has more

negative Ψ_{MD} in monoculture (Fig. S8). Moreover, trends in ϕ_{NO} suggest that *T. americana* and *A.*

negundo in high light are especially vulnerable to photodamage (Fig. 4). Photoinhibition due to source-

sink imbalance is usually accompanied by compensatory upregulation in protective non-photochemical

quenching (Adams III et al. 2013), but these two species fail to increase qN under high light (Fig. 4).

These findings reinforce the idea that damage, not just downregulation, contributes to the steep declines

in dark-acclimated F_v/F_m these species show under increasing light. Sink limitation may contribute to the

photoinhibition and the growth patterns we observe in these species, but we believe it is not the main

explanation.

The microclimate could also alter photosynthesis in ways our light-response measurements did

not capture because we controlled the chamber microclimate. Our water potential measurements show

that water stress was not very severe in any treatment, but even modest differences could change the

timing of stomatal closure (Brodribb et al. 2003). It also seems plausible that greater light exposure may

make leaf temperature higher in monoculture (Schymanski et al. 2013), which could push leaves above

thermal optima for photosynthesis. Such microclimatic factors could contribute to the physiological

benefits of shading beyond what we could measure in light-response curves.

Shade avoidance and early-successional species

In the only early-successional species, *B. papyrifera*, the stem growth response is weak and

dependent on the model specification (Supplemental Materials, Appendix 2), but it does appear to reflect

a longer-term trend that has caused trees with larger neighbors to show faster stem growth in absolute

terms. Consider the absolute growth rate from 2017 to 2018: *B. papyrifera* stems grew faster in shaded

bicultures (median: 578 cm^3; 25^{th}–75^{th} percentile: 231–1168 cm^3) than in bicultures with other broadleaf

species (median: 496 cm^3; 25^{th}–75^{th} percentile: 87–836 cm^3). But there is a striking mismatch between the

growth patterns and photosynthetic physiology: In untransformed and rescaled light-response curves

alike, *B. papyrifera* had lower photosynthetic rates in the shaded biculture compared to the other

treatments. Because it tended to overtop even its largest neighbors, the top leaves received nearly the full

amount of available light in all treatments (Fig. S2) and showed few signs of photoinhibition (Fig. 4).

These results suggest that the increase in stem volume is unrelated to photosynthetic physiology.

One possibility is that rather than (or in addition to) representing an increase in total growth due

to facilitation, the positive trend in stem volume in *B. papyrifera* is driven by a competitive shade

avoidance response, which could increase allocation to shoot biomass at the expense of root biomass.

Such responses tend to be especially strong in shade-intolerant, early-successional species like *B.*

papyrifera (Henry & Aarssen 1997; Gilbert et al. 2001). While we could not test directly for a change in

the root-to-shoot ratio, we did find evidence that *B. papyrifera*, but not *A. negundo* or *T. americana*, had

another symptom of the shade avoidance syndrome: greater allocation to elongation growth near larger

neighbors, which caused trees to have smaller diameters than expected based on their height (Henry &

Aarssen 1999). This finding may suggest that a broader shade avoidance syndrome may contribute to the

positive influence of neighbor size on stem volume in *B. papyrifera*. This finding would be in contrast to

another tree diversity experiment, where both root-to-shoot ratios and the allometric relationship between

basal area and biomass were nearly unaffected by the local neighborhood (Guillemot et al. 2020). In the

absence of direct evidence, changes in root-to-shoot ratios driven by shade avoidance remain a tentative

but plausible contributor to positive growth responses in *B. papyrifera*. Such changes in allocation are not

mutually exclusive with facilitation, but could obviate the need to invoke facilitation.

Mechanisms of photoprotection

Plants' strategies to avoid damage from the stress of excess light can be classified broadly into

biochemical and structural strategies. We show that trees in full sun allocate more to biochemical

photoprotection, as indicated by lower PRI in all species and higher qN in all species but *T. americana*

and *A. negundo* (Fig. 4). Our leaf angle survey is also consistent with the idea that plants may steeply

incline their leaves to avoid intercepting excess light—a form of structural photoprotection (Fig. 6).

Leaves that are steeply inclined intercept less light per unit area, particularly during midday, when solar

radiation is otherwise most intense. The dramatic trend in leaf angles suggests that gross structural

characteristics allow these species use to regulate their light absorption.

704   Some plants may lack perfect photoprotective mechanisms in part because they are adapted to

environments where the costs of high photoprotective investment outweigh the benefits. Delayed

relaxation of non-photochemical quenching may hinder photosynthesis even when light declines enough

to relieve the imminent threat of damage (Zhu et al. 2004; Murchie & Niyogi 2011; Kromdijk et al.

2016). Building pigments and proteins for photoprotection uses resources that plants could otherwise

allocate to chlorophyll, RuBisCO, or structural tissue components, although the concentrations of such

photoprotective compounds are often low. Steep leaf angles may also decrease photosynthesis by

reducing light interception even when light is limiting—and all plants experience light-limiting conditions

at least sometimes, as during cloudy days or nighttime. All the same, plants in more open environments

may need photoprotective mechanisms to avoid even greater costs due to photodamage under brighter

conditions. That most species upregulate photoprotection in high light, despite the costs, implies that

these species may otherwise incur a risk of damage from excess light. These findings reinforce that light

stress and its alleviation through shading can contribute to net species interactions.

Competition and facilitation in biodiversity-ecosystem function relationships

719   The finding that shade can increase growth and photosynthesis has precedents. Ball et al. (1991)

and Egerton et al. (2000) showed that shade enhances assimilation and growth of evergreen *Eucalyptus*

seedlings during the winter, when cold may cause photoinhibition. Howell et al. (2002) also found that

among evergreen divaricating shrubs of New Zealand, leafless outer branches increase photosynthesis

during the winter by reducing photoinhibition. Species can also facilitate each other from light stress in

communities of phytoplankton (Gerla et al. 2011), which nearly all face some risk of photoinhibition

(Edwards et al. 2015; Edwards et al. 2016) These studies show that facilitation through shading can occur

in many environments, and perhaps more so in otherwise stressful ones.

Much of the work on facilitation in plant communities is guided by the stress-gradient hypothesis

(SGH), which proposes that facilitative interactions are more common in stressful environments (Bertness

& Callaway 1994; Maestre et al. 2009). We can place photoinhibition in light of the SGH as follows:

Stressful conditions limit the use of light for photochemistry, turning high light exposure into an

additional stressor. Under such stressful conditions, shade from neighbors can ameliorate stress, resulting

in facilitation. Under benign conditions, where plants can use more of the available light for

photochemistry, high light exposure may not cause damage. Here, shading by neighbors is more likely to

cause light limitation, resulting in an adverse, competitive effect on growth. As we show here, whether a

species is stressed by its environment also depends on its own physiological tolerances.

The SGH brought attention to facilitation in plant ecology, including in biodiversity-ecosystem

function research. For example, Mulder et al. (2001) showed that in experimental moss communities,

biodiversity increased productivity only under drought, as species interactions became more positive. But

the mechanisms and consequences of facilitation are still overlooked in biodiversity-ecosystem function

research (Wright et al. 2017). A small but growing literature based in biodiversity experiments has

investigated various mechanisms through which plant species can facilitate each other, including

amelioration of water stress (Wright et al. 2015), dilution of natural enemies (Grossman et al. 2019),

nitrogen fixation by legume-rhizobia mutualisms (Temperton et al. 2007), and other forms of nutrient-

related belowground facilitation (Li et al. 2014). We show that shading can reduce stress induced by the

light environment—a specific form of abiotic stress amelioration, in which certain species alter the local

environment such that it is less stressful to other species (Wright et al. 2017). But shading can also cause

light limitation, resulting in a net competitive effect.

During the first three years of the FAB experiment, Grossman et al. (2017) found that diverse

plots overyield—they are more productive than a weighted average of the constituent species in

monoculture. Using the partition of Loreau & Hector (2001), most overyielding in FAB was a result of

complementarity effects, which are often interpreted as arising from synergistic interactions among

species, including facilitation. Overyielding is a hallmark of positive biodiversity-ecosystem function

relationships, and a whole community can only overyield if at least one of its constituent species

overyields. Even though we test how growth changes with neighbor size, not neighbor diversity, we show

that shading allows some species to grow faster in shaded bicultures or twelve-species plots. In the same

experiment, Grossman et al. (2017) found that *T. americana* had the greatest overyielding in multi-species

plots of any broadleaf species, especially when growing with conifers or *B. papyrifera*. Grossman et al.

(2017) further showed that conifer partners in these species combinations also grew faster than they did in

monoculture, resulting in community overyielding.

The two broadleaf species that show the strongest evidence for facilitation by shade had low

tolerance to excess light, and likely benefited from the milder microclimate their larger neighbors created.

The conifer species, which are fast-growing and less shade-tolerant (except for *P. strobus*), may also have

grown faster in bicultures with broadleaf species because they faced less competition for light than in

monoculture. Other researchers have found that heterogeneity in shade tolerance explains the effect of

tree diversity on productivity, but they attributed this result to higher stand-level light capture enabled by

crown complementarity (Zhang et al. 2012; Toïgo et al. 2017; Searle and Chen 2019). We show that

facilitation of shade-tolerant species by shade-intolerant species is an equally compelling (and perhaps

complementary) explanation.

In the literature on biodiversity and ecosystem function, the attempt to parse out mechanisms—

such as niche partitioning and facilitation—often depends on statistical partitions of biomass or

productivity data (Loreau & Hector 2001; Fox & Kerr 2012) whose interpretation is often debated

(Carroll et al. 2011; Pillai & Gouhier 2019). But because productivity has physiological determinants, we

can explain its patterns using physiological measurements. In this study, survey-based tree growth data

and physiological measurements both show that certain species respond positively and others respond

negatively to the size of their neighbors. But only the physiological measurements allow us to explain this

divergence—they show that these interactions emerge in part from the photosynthetic responses of trees

to the light environment created by their neighbors. The insights gained from such physiological

techniques may benefit plant ecology more widely as we seek to explain community patterns in terms of

basic aspects of plant function.

Acknowledgements

University of Minnesota, including Cedar Creek ESR, lies on the ancestral, traditional, and contemporary

Land of the Dakota. Conversations with Jake Grossman and Laura Williams inspired much of this

research. Beth Fallon, German Vargas G., Artur Stefanski, Daniel Stanton, and Danielle Way all gave

valuable advice about measuring photosynthesis. We are indebted to Cathleen Nguyen, Chris Buyarski,

Troy Mielke, Kally Worm, Jim Krueger, Pam Barnes, Mark Saxhaug, Susan Barrott, and Dan Bahauddin

for making research in FAB possible. Sarah Hobbie and Peter Reich helped design the FAB experiment,

and countless Cedar Creek interns have taken part in the stem growth survey. The Cavender-Bares lab

(especially Jake Grossman and Gerard Sapès), Daniel Stanton, German Vargas G., Artur Stefanski, David

Kramer, and the UMN Physiological Ecology Group all provided feedback on the results or manuscript.

FAB is maintained with support from National Science Foundation under DEB #1234162 to Cedar Creek

LTER. Spectral measurements were conducted as part of NSF/NASA DEB #1342778 to J.C.B. and R.M.

S.K. was supported by an NSF Graduate Research Fellowship (Grant No. 00039202) and a UMN

Doctoral Dissertation Fellowship. R.M. is also supported by Minnesota Agricultural Experiment Station

project MIN-42-060.

References

Adams III WW, Muller O, Cohu CM, Demmig-Adams B. May photoinhibition be a consequence, rather

than a cause, of limited plant productivity? Photosynthesis Research 117, 31–44 (2013).

Ammer C. Diversity and forest productivity in a changing climate. New Phytologist 221, 50–66 (2019).

Ball MC, Hodges VS, Laughlin GP. Cold-induced photoinhibition limits regeneration of snow gum at

tree-line. Functional Ecology 5, 663–668 (1991).

Barry KE, Mommer L, van Ruijven J, Wirth C, Wright AJ, Bai Y, Connolly J, De Deyn G, de Kroon H,

Isbell F et al. The future of complementarity: disentangling causes from consequences. Trends in Ecology

& Evolution 34: 167–180 (2019).

Bartoń K. MuMIn: Multi-model inference. *R* package version 1.43.17 (2020).

Bates D, Maechler M, Bolker B, Walker S. Fitting linear mixed-effects models using *lme4*. Journal of

Statistical Software 67: 1–48 (2015).

Bertness MD, Callaway R. Positive interactions in communities. Trends in Ecology & Evolution 9, 191–

193 (1994).

Björkman O, Powles SB. Inhibition of photosynthetic reactions under water stress: interaction with light

level. Planta 161, 490–504 (1984).

Braun EL. Deciduous Forests of Eastern North America. Philadelphia: Blakiston (1950).

Braun-Blanquet J. Plant Sociology (trans. Fuller GD, Conard HS). New York: McGraw-Hill (1932).

Brelsford CC, Trasser M, Paris T, Hartikainen SM, Robson TM. Understory light quality affects leaf

pigments and leaf phenology in different plant functional types. biorXiv (2019). DOI: 10.1101/829036.

Brodribb TJ, Holbrook NM, Edwards EJ, Gutiérrez MV. Relations between stomatal closure, leaf turgor

and xylem vulnerability in eight tropical dry forest trees. Plant, Cell & Environment 26, 443–450 (2003).

Canham CD, Finzi AC, Pacala SW, Burbank DH. Causes and consequences of resource heterogeneity in

forests: interspecific variation in light transmission by canopy trees. Canadian Journal of Forest Research

24, 337–349 (1994).

Carroll IT, Cardinale BJ, Nisbet RM. Niche and fitness differences relate the maintenance of diversity to

ecosystem function. Ecology 92, 1157–1165 (2011).

Cavender-Bares J. Chilling and freezing stress in live oaks (Quercus section Virentes): Intra- and

interspecific variation in PS II sensitivity corresponds to latitude of origin. Photosynthesis Research 94,

437–453 (2007).

Cavender-Bares J, Bazzaz FA. From leaves to ecosystems: Assessing photosynthesis and plant function in

ecological studies. In Papageorgiou GC and Govindjee (Eds.), Chlorophyll a Fluorescence: A Signature

of Photosynthesis, 737–755. Dordrecht, Netherlands: Kluwer Academic Publishers (2004).

Cavender-Bares J, Potts M, Zacharias E, Bazzaz FA. Consequences of CO2 and light interactions for leaf

phenology, growth, and senescence in *Quercus rubra*. Global Change Biology 6, 877–887 (2000).

Demmig-Adams B, Adams, WW III. Photoprotection and other responses of plants to high light stress.

Annual Review of Plant Physiology and Plant Molecular Biology 43, 599–626 (1992).

Dybzinski R, Tilman D. Resource use patterns predict long-term outcomes of plant competition for

nutrients and light. The American Naturalist 170, 305–318 (2007).

Edwards KF, Thomas MK, Klausmeier CA, Litchman E. Phytoplankton growth and the interaction of

light and temperature: A synthesis at the species and community level. Limnology and Oceanography 61,

1232–1244 (2016).

Egerton JJG, Banks JCG, Gibson A, Cunningham RB, Ball MC. Facilitation of seedling establishment:

Reduction in irradiance enhances winter growth of *Eucalyptus pauciflora*. Ecology 81, 1437–1449

(2000).

Ehleringer J, Werk KS. Modifications of solar radiation absorption patterns and the implications for

carbon gain at the leaf level. In Givnish T (ed.), On the Economy of Plant Form and Function, 57–82.

London: Cambridge University Press (1986).

Estiarte M, Peñuelas J. Alteration of the phenology of leaf senescence and fall in winter deciduous species

by climate change: Effects on nutrient proficiency. Global Change Biology 21, 1005–1017 (2015).

Farquhar GD, von Caemmerer S, Berry JA. A biochemical model of photosynthetic CO_2 assimilation in

leaves of C_3 species. Planta 149, 78–90 (1980).

Fichtner A, Härdtle W, Li Y, Bruelheide H, Kunz M, von Oheimb G. From competition to facilitation:

how tree species respond to neighbourhood diversity. Ecology Letters 20, 892–900 (2017).

Fox JW, Kerr B. Analyzing the effects of species gain and loss on ecosystem function using the extended

Price equation partition. Oikos 121, 290–298 (2012).

Gamfeldt L, Snäll T, Bagchi T, Jonsson M, Gustafsson L, Kjellander P, Ruiz-Jaen MC, Fröberg M,

Stendahl J, Philipson CD et al. Higher levels of multiple ecosystem services are found in forests with

more tree species. Nature Communications 4, 1340 (2013).

Gamon JA, Peñuelas J, Field CB. A narrow-waveband spectral index that tracks diurnal changes in

photosynthetic efficiency. Remote Sensing of Environment 41, 35–44 (1992).

Gerla DJ, Mooij WM, Huisman J. Photoinhibition and the assembly of light-limited phytoplankton

communities. Oikos 120, 359–368 (2011).

Gilbert IR, Jarvis PG, Smith H. Proximity signal and shade avoidance differences between early and late

successional trees. Nature 411, 792–795 (2001).

Gitelson AA, Gamon JA, Solovchenko, A. Multiple drivers of seasonal change in PRI: Implications for

photosynthesis 1. Leaf level. Remote Sensing of Environment 191, 110–116 (2017).

Grossman JJ, Cavender-Bares J, Hobbie SE, Reich PB, Montgomery RA. Species richness and traits

predict overyielding in stem growth in an early-successional tree diversity experiment. Ecology 98, 2601–

2614 (2017).

Grossman JJ, Cavender-Bares J, Reich PB, Montgomery RA, Hobbie SE. Neighborhood diversity

simultaneously increased and decreased susceptibility to contrasting herbivores in an early stage forest

diversity experiment. Journal of Ecology 107, 1492–1505 (2019).

Grossman JJ, Vanhellemont M, Barsoum N, Bauhus J, Bruelheide H, Castagneyrol B, Cavender-Bares J,

Eisenhauer N, Ferlian O, Gravel D et al. Synthesis and future research directions linking tree diversity to

growth, survival, and damage in a global network of tree diversity experiments. Environmental and

Experimental Botany 152, 68–89 (2018).

Guillemot J, Kunz M, Schnabel F, Fichtner A, Madsen CP, Gebauer T, Härdtle W, Oheimb G, Potvin C.

Neighbourhood-mediated shifts in tree biomass allocation drive overyielding in tropical species mixtures.

New Phytologist 228, 1256–1268 (2020).

Hautier Y, Niklaus PA, Hector A. Competition for light causes plant biodiversity loss after

eutrophication. Science 324, 636–638 (2009).

Henry HAL, Aarssen LW. On the relationship between shade tolerance and shade avoidance strategies in

woodland plants. Oikos 80, 575–582 (1997).

Henry HAL, Aarssen LW. The interpretation of stem diameter–height allometry in trees: Biomechanical

constraints, neighbour effects, or biased regressions? Ecology Letters 2, 89–97 (1999).

Hoffmann WA, Poorter H. Avoiding bias in calculations of relative growth rate. Annals of Botany 90, 37

(2002).

924 Howell CJ, Kelly D, Turnbull MH. Moa ghosts exorcised? New Zealand's divaricate shrubs avoid

925 photoinhibition. Functional Ecology 16, 232–240 (2002).

927 Huang Y, Chen Y, Castro-Izaguirre N, Baruffol M, Brezzi M, Lang A, Li Y, Härdtle W, von Oheimb G,

928 Yang V et al. Impacts of species richness on productivity in a large-scale subtropical forest experiment.

929 Science 362, 80–83 (2018).

931 Johnson IR, Thornley JHM. A model of instantaneous and daily canopy photosynthesis. Journal of

932 Theoretical Biology 107, 531–545 (1984).

934 Johnston MH, Homann PS, Engstrom JK, Grigal DF. Changes in ecosystem carbon storage over 40 years

935 on an old-field/forest landscape in east-central Minnesota. Forest Ecology and Management 83, 17–26

936 (1996).

938 Juvany M, Müller M, Munné-Bosch S. Photo-oxidative stress in emerging and senescing leaves: a mirror

939 image? Journal of Experimental Botany 64, 3087–3098 (2013).

941 Kothari S, Cavender-Bares J, Bitan K, Verhoeven AS, Wang R, Montgomery RA, Gamon J. Community-

942 wide consequences of variation in photoprotective physiology among prairie plants. Photosynthetica 56,

943 455–467 (2018).

945 Kramer DM, Johnson G, Kiirats O, Edwards GE. New fluorescence parameters for the determination of

946 QA redox state and excitation energy fluxes. Photosynthesis Research 79, 209–218 (2004).

Kromdijk J, Głowacka K, Leonelli L, Gabilly ST, Iwai M, Niyogi KK, Long SP. Improving

photosynthesis and crop productivity by accelerating recovery from photoprotection. Science 354, 857–

861 (2016).

Külheim C, Ågren J, Jansson S. Rapid regulation of light harvesting and plant fitness in the field. Science

297, 91–93 (2002).

Kuznetsova A, Brockhoff PB, Christensen RHB. *lmerTest* package: Tests in linear mixed effects models.

Journal of Statistical Software 82, 1-26 (2017).

Liang J, Crowther TW, Picard N, Wiser S, Zhou M, Alberti G, Schulze E-D, McGuire AD, Bozzato F,

Pretzsch H et al. Positive biodiversity-productivity relationship predominant in global forests. Science

354, aaf8957 (2016).

Li L, Tilman D, Lambers H, Zhang F-S. Plant diversity and overyielding: Insights from belowground

facilitation of intercropping in agriculture. New Phytologist 203, 63–69 (2014).

Long SP, Humphries S, Falkowski PG. Photoinhibition of photosynthesis in nature. Annual Review of

Plant Physiology and Plant Molecular Biology 45, 633-62 (1994).

Loreau M, Hector A. Partitioning selection and complementarity in biodiversity experiments. Nature 412,

72–76 (2001).

Lovelock CE, Clough BF. Influence of solar radiation and leaf angle on leaf xanthophyll concentrations

in mangroves. Oecologia 91, 518–525 (1992).

Maestre FT, Callaway RM, Valladares F, Lortie CJ. Refining the stress-gradient hypothesis for

competition and facilitation in plant communities. Journal of Ecology 97, 199–205 (2009).

Mahmoud A, Grime JP. A Comparison of negative relative growth rates in shaded seedlings. New

Phytologist 73, 1215–1219 (1974).

Mattila H, Valev D, Havurinne V, Khorobrykh S, Virtanen O, Antinluoma M, Mishra KB, Tyystjärvi E.

Degradation of chlorophyll and synthesis of flavonols during autumn senescence—the story told by

individual leaves. AoB Plants 10, ply028 (2018).

Monsi M, Saeki T. On the factor light in plant communities and its importance for matter production.

Annals of Botany 95, 549–567 (2005).

Monteith JL, Moss CJ, Cooke GW, Pirie NW, Bell GDH. Climate and the efficiency of crop production

in Britain. Philosophical Transactions of the Royal Society of London B: Biological Sciences 281, 277–

294 (1977).

Montgomery RA, Goldstein G, Givnish TJ. Photoprotection of PSII in Hawaiian lobeliads from diverse

light environments. Functional Plant Biology 35, 595–605 (2008).

Montgomery RA, Reich PB, Palik BJ. Untangling positive and negative biotic interactions: views from

above and below ground in a forest ecosystem. Ecology 91, 3641–3655 (2010).

Mulder CPH, Uliassi DD, Doak DF. Physical stress and diversity-productivity relationships: The role of

positive interactions. Proceedings of the National Academy of Sciences 98: 6704–8 (2001).

Murchie EH, Niyogi KK. Manipulation of photoprotection to improve plant photosynthesis. Plant

Physiology 155, 86–92 (2011).

1002

1003   Nakagawa S, Johnson PCD, Schielzeth H. The coefficient of determination R^2 and intra-class correlation

1004   coefficient from generalized linear mixed-effects models revisited and expanded. Journal of the Royal

1005   Society Interface 14, 20170213 (2017).

1006

1007   Niinemets Ü, Valladares F. Tolerance to shade, drought, and waterlogging of temperate Northern

1008   Hemisphere trees and shrubs. Ecological Monographs 76, 521–547 (2006).

1009

1010   Oehri J, Schmid B, Schaepman-Strub G, Niklaus PA. Biodiversity promotes primary productivity and

1011   growing season lengthening at the landscape scale. Proceedings of the National Academy of Sciences

1012   114, 10160–10165 (2017).

1013

1014   Parent S, Messier C. A simple and efficient method to estimate microsite light availability under a forest

1015   canopy. Canadian Journal of Forest Research. 26, 151–154 (1996).

1016

1017   Pastenes C, Porter V, Baginsky C, Horton P, González J. Paraheliotropism can protect water-stressed

1018   bean (*Phaseolus vulgaris* L.) plants against photoinhibition. Journal of Plant Physiology 161, 1315–1323

1019   (2004).

1020

1021   Pearcy RW, Muraoka H, Valladares F. Crown architecture in sun and shade environments: assessing

1022   function and trade-offs with a three-dimensional simulation model. New Phytologist 166, 791–800

1023   (2005).

1024

1025   Pillai P, Gouhier TC. Not even wrong: the spurious measurement of biodiversity's effects on ecosystem

1026   functioning. Ecology 100, e02645 (2019).

1027

1028   Poorter H, Niinemets Ü, Ntagkas N, Siebenkäs A, Mäenpää M, Matsubara S, Pons TL. A meta-analysis

1029   of plant responses to light intensity for 70 traits ranging from molecules to whole plant performance. New

1030   Phytologist 223, 1073–1105 (2019).

1031

1032   Posada JM, Lechowicz M, Kitajima K. Optimal photosynthetic use of light by tropical tree crowns

1033   achieved by adjustment of individual leaf angles and nitrogen content. Annals of Botany 103, 795–805

1034   (2009).

1035

1036   Potvin C, Gotelli NJ. Biodiversity enhances individual performance but does not affect survivorship in

1037   tropical trees. Ecology Letters 11, 217–223 (2008).

1038

1039   R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical

1040   Computing, Vienna, Austria (2019). https://www.R-project.org/.

1041

1042   Ramírez-Valiente JA, Koehler K, Cavender-Bares J. Climatic origins predict variation in photoprotective

1043   leaf pigments in response to drought and low temperatures in live oaks (*Quercus* series *Virentes*). Tree

1044   Physiology 35, 521–534 (2015).

1045

1046   Renner SS, Zohner CM. The occurrence of red and yellow autumn leaves explained by regional

1047   differences in insolation and temperature. New Phytologist 224, 1464–1471 (2019).

1048

1049   Sapijanskas J, Paquette A, Potvin C, Kunert N, Loreau M. Tropical tree diversity enhances light capture

1050   through crown plasticity and spatial and temporal niche differences. Ecology 95, 2479–2492 (2014).

1051

1052 Savage J, Cavender-Bares J, Verhoeven A. Habitat generalists and wetland specialists in the genus *Salix*

1053 vary in their photoprotective responses to drought. Functional Plant Biology 36, 300–309 (2009).

1054

1055 Schymanski SJ, Or D, Zwieniecki M. Stomatal control and leaf thermal and hydraulic capacitances under

1056 rapid environmental fluctuations. PLOS ONE 8, e54231 (2013).

1057

1058 Searle EB, Chen HYH. Complementarity effects are strengthened by competition intensity and global

1059 environmental change in the central boreal forests of Canada. Ecology Letters 23, 79–87 (2020).

1060

1061 Shipley B, Meziane D. The balanced-growth hypothesis and the allometry of leaf and root biomass

1062 allocation. Functional Ecology 16, 326–331 (2002).

1063

1064 Simonsohn U. Two Lines: A valid alternative to the invalid testing of U-shaped relationships with

1065 quadratic regressions. Advances in Methods and Practices in Psychological Science 1, 538–555 (2018).

1066

1067 Streb P, Tel-Or E, Feierabend J. Light stress effects and antioxidative protection in two desert plants.

1068 Functional Ecology 11, 416-424 (1997).

1069

1070 Tardieu F, Parent B, Caldeira CF, Welcker C. (2014). Genetic and physiological controls of growth under

1071 water deficit. Plant Physiology 164, 1628–1635.

1072

1073 Temperton VM, Mwangi PN, Scherer-Lorenzen M, Schmid B, Buchmann N. Positive interactions

1074 between nitrogen-fixing legumes and four different neighbouring species in a biodiversity experiment.

1075 Oecologia 151, 190–205 (2007).

1076

1077   Tietz S, Hall CC, Cruz, JA, Kramer DM. NPQ(T): a chlorophyll fluorescence parameter for rapid

1078   estimation and imaging of non-photochemical quenching of excitons in photosystem-II-associated

1079   antenna complexes. Plant, Cell & Environment 40, 1243–1255 (2017).

1080

1081   Tilman D, Isbell F, Cowles JM. Biodiversity and ecosystem functioning. Annual Review of Ecology,

1082   Evolution, and Systematics 45, 471–493 (2014).

1083

1084   Tobner CM, Paquette A, Gravel D, Reich PB, Williams LJ, Messier C. Functional identity is the main

1085   driver of diversity effects in young tree communities. Ecology Letters 19, 638–647 (2016).

1086

1087   Toïgo M, Perot T, Courbaud B, Castagneyrol B, Gégout J-C, Longuetaud F, Jactel H, Vallet P. Difference

1088   in shade tolerance drives the mixture effect on oak productivity. Journal of Ecology 106: 1073–1082

1089   (2017).

1090

1091   Udo SO, Aro TO. Global PAR related to global solar radiation for central Nigeria. Agricultural and Forest

1092   Meteorology 97: 21–31 (1999).

1093

1094   Valiente-Banuet A, Verdú M, Valladares F, García-Fayos P. Functional and evolutionary correlations of

1095   steep leaf angles in the mexical shrubland. Oecologia 163, 25–33 (2010).

1096

1097   Valladares F, Pugnaire FI. Tradeoffs between irradiance capture and avoidance in semi-arid environments

1098   assessed with a crown architecture model. Annals of Botany 83, 459–469 (1999).

1099

1100   Watling JR, Robinson SA, Woodrow IE, Osmond CB. Responses of rainforest understorey plants to

1101   excess light during sunflecks. Australian Journal of Plant Physiology 24, 17–25 (1997).

1102

1103   Werner C, Correia O, Beyschlag W. Two different strategies of Mediterranean macchia plants to avoid

1104   photoinhibitory damage by excessive radiation levels during summer drought. Acta Oecologica 20, 15–23

1105   (1999).

1106

1107   Williams LJ, Paquette A, Cavender-Bares J, Messier C, Reich PB. Spatial complementarity in tree crowns

1108   explains overyielding in species mixtures. Nature Ecology & Evolution 1, 0063 (2017).

1109

1110   Williams LJ, Cavender-Bares J, Paquette A, Messier C, Reich PB. Light mediates the relationship

1111   between community diversity and trait plasticity in functionally and phylogenetically diverse tree

1112   mixtures. Journal of Ecology 108, 1617–1634 (2020).

1113

1114   Wong CYS, Gamon JA. Three causes of variation in the photochemical reflectance index (PRI) in

1115   evergreen conifers. New Phytologist 206, 187–195 (2015).

1116

1117   Wright AJ, Schnitzer SA, Reich PB. Daily environmental conditions determine the competition–

1118   facilitation balance for plant water status. Journal of Ecology 103, 648–656 (2015).

1119

1120   Wright AJ, Wardle DA, Callaway R, Gaxiola A. The overlooked role of facilitation in biodiversity

1121   experiments. Trends in Ecology & Evolution 32, 383–390 (2017).

1122

1123   Wujeska A, Bossinger G, Tausz M. Responses of foliar antioxidative and photoprotective defence

1124   systems of trees to drought: a meta-analysis. Tree Physiology 33, 1018–1029 (2013).

1125

1126   Yachi S, Loreau M. Does complementary resource use enhance ecosystem functioning? A model of light

1127   competition in plant communities. Ecology Letters 10, 54–62 (2007).

1128

1129   Zemp DC, Gérard A, Hölscher D, Ammer C, Irawan B, Sundawati L, Teuscher M, Kreft H. Tree

1130   performance in a biodiversity enrichment experiment in an oil palm landscape. Journal of Applied

1131   Ecology 56, 2340–2352 (2019).

1132

1133   Zhang Y, Chen HYH, Reich PB. Forest productivity increases with evenness, species richness and trait

1134   variation: a global meta-analysis. Journal of Ecology 100, 742–749 (2012).

1135

1136   Zhu X-G, Ort DR, Whitmarsh J, Long SP. The slow reversibility of photosystem II thermal energy

1137   dissipation on transfer from high to low light may cause large losses in carbon gain by crop canopies: a

1138   theoretical analysis. Journal of Experimental Botany 55, 1167–1175 (2004).

1139 **Tables**

1140

1141 Table 1: Characteristics of species in FAB. Except for *Betula papyrifera*, broadleaf species tended to

1142 grow slower than needleleaf conifers. Shade tolerance is drawn from Niinemets and Valladares (2006),

1143 who evaluated it along a 1-5 scale where higher values indicate an ability to grow under lower light.

1144

Scientific name	Species code	Common name	Family	Leaf form	Mean RGR in monoculture, 2013-6 (/year)	Shade tolerance	Partner in shaded biculture
Acer negundo	ACNE	Box elder	Sapindaceae	Broadleaf	0.039	3.47	None
Acer rubrum	ACRU	Red maple	Sapindaceae	Broadleaf	0.379	3.44	*Pinus banksiana*
Betula papyrifera	BEPA	Paper birch	Betulaceae	Broadleaf	1.272	1.54	*Pinus banksiana*
Juniperus virginiana	JUVI	Eastern redcedar	Cupressaceae	Needleleaf	0.975	1.28	NA
Pinus banksiana	PIBA	Jack pine	Pinaceae	Needleleaf	2.288	1.36	NA
Pinus resinosa	PIRE	Red pine	Pinaceae	Needleleaf	1.619	1.89	NA
Pinus strobus	PIST	White pine	Pinaceae	Needleleaf	1.722	3.21	NA
Quercus alba	QUAL	White oak	Fagaceae	Broadleaf	0.735	2.85	*Pinus strobus*
Quercus ellipsoidalis	QUEL	Northern pin oak	Fagaceae	Broadleaf	0.722	NA	*Pinus strobus*
Quercus macrocarpa	QUMA	Bur oak	Fagaceae	Broadleaf	0.547	2.71	*Betula papyrifera*
Quercus rubra	QURU	Northern red oak	Fagaceae	Broadleaf	0.505	2.75	*Pinus strobus*
Tilia americana	TIAM	American basswood	Tiliaceae	Broadleaf	0.724	3.98	*Pinus strobus*

1145

1146

1147

1148

1149

1150

   Figure captions

Fig. 1, top: Imagine the same slow-growing broadleaf species (round) found in two communities: a

monoculture (left), and a biculture (right) with a faster growing conifer (triangular). In the absence of

shading by neighbors, all trees would receive the amount of light available at the top of the canopy

(yellow line). We define the relative light availability (RLA) for a tree as the fraction of this light that the

tree's top leaves receive. In monoculture, most broadleaf trees have an RLA close to 1. In shaded

biculture, they are shaded by their larger conifer neighbors, so their RLA is much less than 1. These trees

may be more likely to suffer from light limitation, but they are also at less risk for photoinhibition.

Middle: For a single leaf, carbon assimilation tends to increase along a saturating curve as light intensity

at the leaf surface increases. Consider two hypothetical photosynthetic light-response curves (left) of

leaves from shaded (blue) and unshaded (red) plants. Given a light-response curve, a reduction in light

reduces photosynthesis (represented by the small black arrow along the red curve). But if a long-term

reduction in light spares the plant from photoinhibition, it can change the shape of the curve such that

carbon assimilation at any given light level is greater (represented by the small black vertical arrow). The

balance (represented by the large dashed arrow) may either increase or reduce carbon assimilation. In this

framework, trees benefit most from shading when light is otherwise abundant: the risk of photoinhibition

is greater, and the proportional cost of a given percentage decline in light is lower, since light-response

curves are usually concave-down. Whatever the amount of light at the top of the canopy, the top leaves of

the shaded plant only receive a corresponding fraction (blue dashed line), while the unshaded plant

receives the full amount (red dashed line). To make it easy to compare assimilation rates as a function of

light at the top of the canopy, we can rescale the blue light-response curve along the x-axis by the inverse

of its RLA, making the two dashed lines coincide (middle). In this case, after rescaling, the shaded plant

has higher rates of photosynthesis except when the amount of light at the top of the canopy is low. When

shaded plants have higher rates of photosynthesis summed over time, they may have faster growth, so

growth rates increase with larger neighbors (right).

Bottom: We can imagine another tree species (left) in which photoinhibition has a smaller effect (the

shaded and unshaded curves are more similar) and the RLA in the shade is lower (the blue dashed line is

at a lower light level), such that shading has greater costs and smaller benefits. Here, rescaling makes it

clear that the shaded plant has lower photosynthesis at any given level of light at the top of the canopy

(middle), so plants shaded by large neighbors may have lower growth rates (right).

1182

Fig. 2: Relative growth rate (RGR) of woody stem volume in individuals of each species between fall

2016 and 2018 as a function of the average stem volume of all neighbors. Gray dots represent individuals,

and the regression is fit based on a mixed-effects model with plot as a random intercept. For *Tilia*

americana and *Acer negundo*, the relationship with neighbor volume is modeled as logarithmic rather

than linear. The large colored dots aggregate data to the plot scale, as described in the main text, and are

color-coded by treatment. A shaded biculture is a plot where the focal broadleaf species grows with either

a conifer or *Betula papyrifera* (unless *B. papyrifera* is the focal broadleaf species). An open biculture is a

plot where the focal broadleaf grows with a broadleaf species other than *B. papyrifera*. About 1.1% of

trees fall outside the plot bounds.

1192

Fig. 3: Mass-based photosynthetic light-response curves for four broadleaf species. The left panels depict

carbon assimilation rates as a function of chamber light intensity; the right panels present estimates of

realized carbon assimilation rates as a function of light intensity at the top of the canopy. These rates are

lower because of shading by neighbors (as in Fig. 1, bottom right). Species are arranged from the least

shade-tolerant (*Betula papyrifera*) to the most (*Tilia americana*). Error bars in left panels and gray

ribbons in right panels are ± 1 SE.

1199

Fig. 4: Chlorophyll fluorescence parameters and the photochemical reflectance index (PRI) as a function

of relative light availability across species and treatments. Best-fit lines come from species-specific OLS

regressions. Dark-acclimated F_v/F_m is the maximal quantum efficiency of PSII photochemistry and

declines as a plant becomes more photoinhibited. qN describes a plant's capacity to use non-

photochemical quenching to dissipate energy. ϕ_{NO} represents the proportion of light dissipated in an

unregulated way, which may contribute to photodamage. PRI correlates negatively with carotenoid :

chlorophyll ratios.

1207

1208   Fig. 5: Compared to shade-intolerant species, shade-tolerant species show greater declines in dark-

1209   acclimated F_v/F_m (top) and smaller increases in qN (middle) as light availability increases. A_{sat} derived

1210   from light-response curves correlates positively and strongly with dark-acclimated F_v/F_m; slopes are

1211   statistically indistinguishable among species (bottom). Species codes are found in Table 1.

1212

1213   Fig. 6: In *Tilia americana*, fall leaf abscission occurs later in monoculture than in shaded biculture.

1214

1215   Fig. 7: The angles of top leaves of three species are more horizontal in shaded biculture plots than in

1216   monoculture.

 Figures

 Fig. 1

Fig. 2

1245   **Fig. 3**

Fig. 4

Fig. 5

1266   **Fig. 6**

1267
1268
1269   **Fig. 7**

1270

1271   **Supplement to: "Physiological responses to light explain competition and facilitation in a tree**

1272   **diversity experiment"**

1273

Appendix 1

Photosynthetic light-response curve methods

We measured photosynthetic light-response curves from the four focal species (*Acer negundo*,

Betula papyrifera, *Quercus ellipsoidalis*, and *Tilia americana*) during July 2018 using an LI-6400 gas

exchange system (LI-COR BioSciences, Lincoln, NE, USA) with a 6400-40 Leaf Chamber Fluorometer

head. From each tree ($n = 72$ total), we selected a fully expanded, sun-exposed leaf with no visible disease

or herbivory. We noted the angle of each leaf relative to horizontal to the nearest 15˚ before beginning

measurements. Each curve had nine steps in descending order of brightness: 2000, 1500, 1000, 500, 200,

100, 50, 20, and 0 µmol m^{-2} s^{-1}.

During each curve, we controlled environmental conditions to reduce the immediate effects of

environmental conditions other than light on stomatal conductance. We kept relative humidity at a

favorable level, typically between 65 and 75%, and set the block temperature as close as possible to 25

˚C. The reference CO_2 concentration remained at 400±10 ppm. We kept each leaf in the chamber for at

least five minutes before beginning each curve. We checked that outputs were stable, and matched IRGAs

before each measurement.

Since it is hard to control chamber conditions perfectly, any differences that remain among

treatments could contribute to the variation we observe in photosynthetic function. During light-response

curves, we found no difference among treatments in leaf-to-air VPD within the chamber. There were

significant but slight differences in leaf temperature inside the chamber; leaves in monoculture, shaded

biculture, and twelve-species plots had average temperatures of 27.7 ˚C, 26.5 ˚C, and 27.0 ˚C,

respectively. Previous studies on the temperature response of some of the same species in Minnesota have

typically found thermal optima below 26 ˚C (Sendall et al. 2015). But while cooler leaves may be closer

to thermal optima, the small differences we find in leaf temperature cannot explain much of the variation

in assimilation rates.

In all but five light-response curves, we also used a saturating pulse to measure light-acclimated

chlorophyll fluorescence at each light level. These data allowed us to calculate ϕ_{PSII}, a measure of the

efficiency of Photosystem II (PSII) at each actinic (or photosynthetic) light level, as:

1301
$$\phi_{PSII} = \frac{F'_m - F_s}{F'_m}$$

1302 where F_s is the steady-state fluorescence yield under actinic light, and F_m' is the maximum fluorescence

1303 yield of the actinic light-acclimated sample following a saturating light pulse, which temporarily closes

1304 all PSII reaction centers. From this parameter, we calculated electron transport rate as:

1305
$$ETR = \phi_{PSII} \times PPFD \times 0.42 \times 0.5$$

1306 where PPFD is photosynthetic photon flux density (the light level inside the chamber, in μmol m^{-2} s^{-1}),

1307 0.42 is an estimate of leaf absorptance, and 0.5 is an estimate of the fraction of photons that are captured

1308 by PSII rather than PSI. Electron transport rate is a key parameter because electron transport is one of the

1309 main steps that is capable of limiting photosynthesis (Farquhar et al. 1980), especially when light

1310 availability or PSII efficiency are low.

1311

Chlorophyll fluorescence parameters

As described in the main text, we used dark- and light-acclimated measurements of chlorophyll

fluorescence to calculate several derived parameters, from which we make inferences about

photosynthetic and photoprotective function. Most importantly, we wanted to assess: (1) capacity for non-

photochemical quenching, as measured through the parameters qN or NPQ; and (2) the partitioning of

light energy dissipation among pathways, as assessed using the quantum yields ϕ_{PSII}, ϕ_{NPQ}, and ϕ_{NO}.

These derived chlorophyll fluorescence parameters are calculated from a few basic measurements

taken during the dark- and light-acclimated measurements, including: F_o, the minimum fluorescence yield

of a dark-acclimated sample with all PSII reaction centers open; F_o', the minimum fluorescence yield of a

light-acclimated sample with all PSII reaction centers open (following far-red light exposure); and F_m, the

maximum fluorescence yield of a dark-acclimated sample following a saturating pulse. F_v, the variable

fluorescence yield of a dark-acclimated sample, is calculated as $F_m - F_o$.

The parameters qN and NPQ are meant to indicate the rate constant of the thermal dissipation of

energy from PSII (Bilger & Schreiber 1986; Bilger & Björkman 1990). These parameters increase with

the use of NPQ to dissipate light—in this case, under exposure to 1000 µmol m^{-2} s $^{-1}$. They are calculated,

respectively, as:

1328
$$qN = 1 - \frac{F'_m - F'_o}{F_m - F_o}; NPQ = \frac{F_m - F'_m}{F'_m}$$

In our dataset, qN is highly correlated ($R^2 = 0.96$) with the natural logarithm of NPQ; we chose to report

qN to easily visualize variation among leaves with low NPQ and to help satisfy the error term normality

assumption of OLS regression.

The quantum yields ϕ_{PSII}, ϕ_{NPQ}, and ϕ_{NO} represent the proportions of absorbed light energy that are

dissipated through photochemistry, non-photochemical quenching, and non-regulated dissipation. Since

these pathways exhaust the options, their quantum yields sum to 1. We calculated them following the

'lake model' derivation of Kramer et al. (2004), which treats photosynthetic reaction centers as connected

by shared antennae. We achieve qualitatively similar results using the simpler analogues found in

Hendricksen et al. (2005).

The parameter ϕ_{PSII} is calculated as shown above (Supplemental Methods, *Photosynthetic light-*

response curve procedure). In the derivation from Kramer et al. (2004), ϕ_{NPQ} and ϕ_{NO} are calculated using

the parameter qL, an estimate of the fraction of open PSII reaction centers. qL is calculated as:

1341
$$qL = \frac{F'_m - F_s}{F'_m - F'_o} \frac{F'_o}{F_s}$$

1342   Given this parameter, we can calculate ϕ_{NO} as:

1343
$$\phi_{NO} = \frac{1}{NPQ + 1 + qL \left(\frac{F_m}{F_o} - 1\right)}$$

1344 and given the other two quantum yields, we can calculate ϕ_{NPQ} as:

1345
$$\phi_{NPQ} = 1 - \phi_{PSII} - \phi_{NO}$$

Many researchers, including Murchie and Lawson (2013) and Tietz et al. (2017) treat ~0.83 as a

theoretical maximum of ϕ_{PSII} (or dark-acclimated F_v/F_m) after a period of dark-acclimation long enough to

reduce non-photochemical quenching to zero. Similarly, one may treat ~0.83 as a theoretical maximum of

$\phi_{PSII} + \phi_{NPQ}$, deviations from which indicate high ϕ_{NO}, and thus, a declining ability to regulate energy

dissipation. We show this putative maximum in our ternary plot of the quantum yields (Fig. S7).

Appendix 2

Here, we compare statistical models testing the influence of average neighbor volume on relative growth

rate (RGR) of focal individuals (or whole plots). The t-statistics, degrees of freedom (df), and p-values

provided are only for the parameter of interest (β_1, the slope term associated with average neighbor

volume), even for multivariate models. For the mixed-effect models with plot incorporated as a random

intercept, we calculated marginal and conditional R^2 using the method of Nakagawa et al. (2017) as

implemented in the R package *MuMIn v. 1.43.17* (Bartoń 2020), and estimated p-values using

Satterthwaite's degrees of freedom method, as implemented in the R package *lmerTest v.3.1.2*

(Kuznetsova et al. 2017). For the plot-aggregated fixed-effect model, we report a standard adjusted R^2. As

usual, the R^2 values are calculated for the full model, including when models have multiple fixed effects.

1361

In the main text, we present summary statistics from models including only neighbor size as a fixed

effect, either untransformed or log-transformed (A or B), based on which fits better for a given species.

The results, including slope estimates, are mostly robust to the model specification (C-E). We highlight in

blue the results presented in the main text, and in yellow, cases where our inferences about how species

respond to neighbor volume would be different under an alternate model specification (based on

comparing p-values to an $\alpha = 0.05$ threshold). For models that include focal individual size in 2016, we

note that these values are not usually correlated with RGR between 2016 and 2018 ($R^2 < 0.01$ except in

Acer negundo), so there is no problem of multicollinearity.

1370

The final three models have independent variables other than focal individual RGR. In F, we predict plot-

level RGR from the plot-scale average of each individual's average neighbor biovolume. In G and H, we

predict the absolute growth rate of focal individuals, which allow us to compare model parameters

including (G) and excluding (H) trees that died between 2016 and 2018. Because of extreme

heteroskedasticity and non-normality of errors, we implemented these models using robust regression

with a smoothed Huber loss function in R package *robustlmm 2.3* (Koller et al. 2016).

A. FocalRGR$_{2016\text{-}2018}$ = α + β_1 × NeighborVolume + (1|Plot)

1378

Species	$\beta_1 \times 10^4$ (SE $\times 10^4$)	R_m^2	R_c^2	t	df	p
Acer negundo	0.394 (0.324)	0.003	0.374	1.217	380	0.224
A. rubrum	-0.775 (0.238)	0.027	0.191	-3.257	366	0.00123
Betula papyrifera	0.279 (0.140)	0.010	0.056	1.993	124	0.049
Quercus alba	-1.062 (0.270)	0.030	0.164	-3.932	486	9.65×10^{-5}
Q. ellipsoidalis	-1.270 (0.232)	0.051	0.108	-5.485	341	8.05×10^{-8}
Q. macrocarpa	-0.423 (0.203)	0.006	0.218	-2.079	697	0.0380
Q. rubra	-1.405 (0.243)	0.053	0.087	-5.785	444	1.37×10^{-8}
Tilia americana	0.348 (0.191)	0.006	0.138	1.818	481	0.0697

1379

1380 B. FocalRGR$_{2016\text{-}2018}$ = α + β_1 × log(NeighborVolume) + (1|Plot)

Species	$\beta_1 \times 10^2$ (SE $\times 10^2$)	R_m^2	R_c^2	t	df	p
Acer negundo	4.005 (1.697)	0.014	0.364	2.361	380	0.0187
A. rubrum	-0.963 (2.293)	0.001	0.182	-0.420	337	0.675
Betula papyrifera	2.498 (1.958)	0.004	0.046	1.276	167	0.204
Quercus alba	-6.403 (2.372)	0.018	0.164	-2.700	367	0.00725
Q. ellipsoidalis	-6.292 (1.928)	0.021	0.093	-3.264	282	0.00124
Q. macrocarpa	-1.742 (1.768)	0.002	0.227	-0.985	661	0.325
Q. rubra	-11.119 (2.083)	0.049	0.076	-5.339	224	2.29×10^{-7}
Tilia americana	6.680 (2.174)	0.018	0.143	3.073	405	0.00226

1381

1382 C. FocalRGR$_{2016\text{-}2018}$ = α + β_1 × NeighborVolume + (1|Plot) (excluding individuals on a block edge)

Species	$\beta_1 \times 10^4$ (SE $\times 10^4$)	R_m^2	R_c^2	t	df	p
Acer negundo	0.481 (0.332)	0.005	0.356	1.447	344	0.149
A. rubrum	-0.830 (0.249)	0.030	0.204	-3.338	359	9.32×10^{-4}
Betula papyrifera	0.264 (0.145)	0.009	0.069	1.823	136	0.0705
Quercus alba	-0.985 (0.281)	0.026	0.180	-3.502	448	5.09×10^{-4}
Q. ellipsoidalis	-1.217 (0.237)	0.048	0.126	-5.135	372	4.56×10^{-7}
Q. macrocarpa	-0.305 (0.206)	0.003	0.241	-1.480	653	0.139
Q. rubra	-1.381 (0.248)	0.052	0.081	-5.580	407	4.40×10^{-8}
Tilia americana	0.321 (0.195)	0.005	0.132	1.651	430	0.0996

1383

1384 D. FocalRGR$_{2016\text{-}2018}$ = α + β_1 × NeighborVolume + β_2 × Focal_size$_{2016}$

1385

Species	$\beta_1 \times 10^4$ (SE $\times 10^4$)	R_m^2	R_c^2	t	df	p
Acer negundo	0.393 (0.322)	0.013	0.344	1.219	376	0.223
A. rubrum	-0.803 (0.242)	0.027	0.187	-3.322	389	9.79×10^{-4}
Betula papyrifera	0.268 (0.140)	0.011	0.056	1.912	130	0.058
Quercus alba	-1.045 (0.272)	0.031	0.165	-3.845	484	1.37×10^{-4}
Q. ellipsoidalis	-1.270 (0.232)	0.051	0.108	-5.480	344	8.26×10^{-8}
Q. macrocarpa	-0.423 (0.203)	0.008	0.225	-2.080	699	0.038
Q. rubra	-1.398 (0.242)	0.057	0.089	-5.769	435	1.52×10^{-8}
Tilia americana	0.352 (0.192)	0.007	0.143	1.839	492	0.0666

1386

1387

1388 E. $FocalRGR_{2016-2018} = \alpha + \beta_1 \times \log(NeighborVolume) + \beta_2 \times Focal\_size_{2016}$

1389

Species	$\beta_1 \times 10^2$ (SE $\times 10^2$)	R_m^2	R_c^2	t	df	p
Acer negundo	3.851 (1.692)	0.025	0.339	2.277	377	0.0234
A. rubrum	-0.983 (2.315)	0.001	0.181	-0.424	358	0.672
Betula papyrifera	2.324 (1.970)	0.005	0.046	1.179	181	0.240
Quercus alba	-6.246 (2.382)	0.020	0.167	-2.622	372	0.00910
Q. ellipsoidalis	-6.312 (1.934)	0.021	0.094	-3.264	290	0.00123
Q. macrocarpa	-1.702 (1.769)	0.003	0.233	-0.962	665	0.337
Q. rubra	-10.930 (2.081)	0.052	0.078	-5.250	220	3.59×10^{-7}
Tilia americana	6.752 (2.178)	0.018	0.148	3.100	416	0.00206

1390

1391 F. $PlotRGR_{2016-2018} = \alpha + \beta_1 \times MeanPlotNeighborVolume$

Species	$\beta_1 \times 10^4$ (SE $\times 10^4$)	R_{adj}^2	t	df	p
Acer negundo	-0.615 (1.339)	-0.037	-0.459	21	0.651
A. rubrum	-0.589 (0.455)	0.022	-1.296	29	0.205
Betula papyrifera	0.126 (0.302)	-0.030	0.417	27	0.680
Quercus alba	-2.258 (0.620)	0.297	-3.642	28	0.00109
Q. ellipsoidalis	-0.870 (0.722)	0.014	-1.205	31	0.237
Q. macrocarpa	-2.225 (0.629)	0.265	-3.540	31	0.00129
Q. rubra	-1.237 (0.695)	0.070	-1.781	28	0.085
Tilia americana	1.026 (0.257)	0.312	3.995	32	3.55×10^{-4}

1392

1393 G. $FocalAbsoluteGrowth_{2017-2018} = \alpha + \beta_1 \times NeighborVolume + (1|Plot)$

1394

Species	$\beta_1 \times 10^2$ (SE $\times 10^2$)	R_m^2	R_c^2	t
Acer negundo	0.301 (0.141)	0.009	0.395	2.133
A. rubrum	-0.482 (0.159)	0.024	0.200	-3.040
Betula papyrifera	8.023 (2.023)	0.044	0.215	3.966
Quercus alba	-0.652 (0.240)	0.014	0.057	-2.715
Q. ellipsoidalis	-0.674 (0.201)	0.018	0.041	-3.359
Q. macrocarpa	-0.582 (0.208)	0.012	0.079	-2.802
Q. rubra	-1.807 (0.345)	0.039	0.039	-5.236
Tilia americana	5.357 (0.918)	0.057	0.130	5.838

1395

1396 H. $FocalAbsoluteGrowth_{2017-2018} = \alpha + \beta_1 \times NeighborVolume + (1|Plot)$ (excluding individuals that died
1397 by 2018)

1398

Species	$\beta_1 \times 10^2$ (SE $\times 10^2$)	R_m^2	R_c^2	t
Acer negundo	0.315 (0.176)	0.007	0.449	1.794
A. rubrum	-0.237 (0.180)	0.005	0.212	-1.319
Betula papyrifera	9.208 (2.027)	0.059	0.254	4.543
Quercus alba	-0.618 (0.280)	0.010	0.054	-2.207
Q. ellipsoidalis	-0.674 (0.221)	0.016	0.045	-3.051
Q. macrocarpa	-0.610 (0.213)	0.013	0.073	-2.866
Q. rubra	-1.804 (0.384)	0.033	0.033	-4.699
Tilia americana	6.212 (0.954)	0.072	0.175	6.511

1399

Supplementary references

Bartoń K. MuMIn: Multi-model inference. *R* package version 1.43.17 (2020).

Bilger W, Björkman O. Role of the xanthophyll cycle in photoprotection elucidated by measurements of light-induced absorbance changes, fluorescence and photosynthesis in leaves of *Hedera canariensis*. Photosynthesis Research 25, 173–185 (1990).

Bilger W, Schreiber U. Energy-dependent quenching of dark-level chlorophyll fluorescence in intact leaves. Photosynthesis Research 10, 303–308 (1986).

Farquhar GD, von Caemmerer S, Berry JA. A biochemical model of photosynthetic CO2 assimilation in leaves of C 3 species. Planta 149, 78–90 (1980).

Hendrickson L, Förster B, Pogson BJ, Chow WS. A simple chlorophyll fluorescence parameter that correlates with the rate coefficient of photoinactivation of Photosystem II. Photosynthesis Research 84, 43–49 (2005).

Kramer DM, Johnson G, Kiirats O, Edwards GE. New fluorescence parameters for the determination of QA redox state and excitation energy fluxes. Photosynthesis Research 79, 209–218 (2004).

Kuznetsova A, Brockhoff PB, Christensen RHB. *lmerTest* package: Tests in linear mixed effects models. Journal of Statistical Software 82, 1-26 (2017).

Koller M. *robustlmm*: An R package for robust estimation of linear mixed-effects models. Journal of Statistical Software 75, 1–24 (2016).

1426

1427 Murchie EH, Lawson T. Chlorophyll fluorescence analysis: a guide to good practice and understanding

1428 some new applications. Journal of Experimental Botany 64, 3983–3998 (2013).

1429

1430 Nakagawa S, Johnson PCD, Schielzeth H. The coefficient of determination R^2 and intra-class correlation

1431 coefficient from generalized linear mixed-effects models revisited and expanded. Journal of the Royal

1432 Society Interface 14, 20170213 (2017).

1433

1434 Sendall, KM, Reich PB, Zhao C, Jihua H, Wei X, Stefanski A, Rice K, Rich RL, Montgomery RA.

1435 Acclimation of photosynthetic temperature optima of temperate and boreal tree species in response to

1436 experimental forest warming. Global Change Biology 21, 1342–1357 (2015).

1437

1438 Tietz S, Hall CC, Cruz, JA, Kramer DM. NPQ(T): a chlorophyll fluorescence parameter for rapid

1439 estimation and imaging of non-photochemical quenching of excitons in photosystem-II-associated

1440 antenna complexes. Plant, Cell & Environment 40, 1243–1255 (2017).

1441

1442   **Supplemental figure captions**

1443

1444   Fig. S1: Example images of *Tilia americana* in shaded biculture (top) and monoculture (bottom). Red

1445   scale bars represent about 1 m in the foreground.

1446

1447   Fig. S2: Relative light availability (top) and leaf mass per area (bottom) of each species across treatments.

1448   We show LMA values from all leaves used in carbon assimilation and chlorophyll fluorescence

1449   measurements. Error bars in left panels are ± 1 SE.

1450

1451   Fig. S3: Growth of woody stem volume in individuals of each species between fall 2017 and 2018 as a

1452   function of the average stem volume of all neighbors. Gray dots represent individuals, while colored dots

1453   are aggregated to the plot scale and color-coded by treatment. Based on mixed-effects robust regression

1454   models with plot as a random intercept, two lines are fit for each species: One (blue) to all data, and one

1455   (red) excluding trees that died between 2017 and 2018.

1456

1457   Fig. S4: Area-based photosynthetic light-response curves for four broadleaf species. The left panels

1458   display carbon assimilation rates as a function of chamber light intensity; the right panels represent an

1459   estimate of realized assimilation rates as a function of light intensity at the top of the canopy. Error bars in

1460   left panels and gray ribbons in right panels are ± 1 SE.

1461

1462   Fig. S5: Responses of ETR, stomatal conductance, and C_i (internal CO_2 concentration) to light availability

1463   for the four focal broadleaf species. Error bars are ± 1 SE.

1464

1465   Fig. S6: Estimates of total mass-based (top) and area-based (middle) carbon assimilation across species

1466   and treatments throughout the month of July, calculated by applying a time series of solar radiation from a

1467   nearby weather station as inputs to our rescaled light-response curves. Error bars are ± 1 SE. (bottom)

1468   Area-based A_{sat} is correlated with LMA across species; within species, there is only a positive relationship

1469   for *Quercus ellipsoidalis* and *Tilia americana*.

1470

Fig. S7: A ternary plot depicting variation among species and treatments in ϕ_{PSII}, ϕ_{NPQ}, and ϕ_{NO}, as

calculated using the lake model derivations of Kramer et al. (2004). These three quantities describe light

energy dissipation through photochemistry, through non-photochemical quenching, and through non-

regulated dissipation, respectively. A thick dashed line represents 0.83, the theoretical maximum

efficiency of PSII posited in Tietz et al. (2017). Deviations to the lower right of this line represent

increasingly high ϕ_{NO}, which can result in photodamage.

1477

1478   Fig. S8: Water potential before dawn (Ψ_{PD}; top) and at midday (Ψ_{MD}; bottom) as a function of light

1479   availability. Data are from monoculture and shaded biculture treatments in all eight species (Ψ_{PD}) or only

1480   the four focal species (Ψ_{MD}). Best-fit lines come from species-specific OLS regressions. Two outliers in

1481 Ψ_{PD} are not included.

1482   **Supplemental figures**
1483
1484   **Fig. S1**
1485

1496   **Fig. S2**
1497
1498

1518   **Fig. S3**

1519

1520   **Fig. S4**

1521

Fig. S5

1532   **Fig. S6**

1533

1534
1535   **Fig. S7**

1554   **Fig. S8**

1555

1556