Neighbor GWAS: incorporating neighbor genotypic identity into genome-

wide association studies of field herbivory on *Arabidopsis thaliana*

Yasuhiro Sato[1,2], Eiji Yamamoto[1,3], Kentaro K. Shimizu[4,5*], Atsushi J. Nagano[6*]

[1]PRESTO, Japan Science and Technology Agency, Kawaguchi 332-0012, Japan

[2]Research Institute for Food and Agriculture, Ryukoku University, Yokotani 1-5, Seta Oe-

cho, Otsu, Shiga 520-2194, Japan

[3]Kazusa DNA Research Institute, Kazusa-kamatari 2-6-7, Kisarazu, Chiba 292-0818, Japan

[4]Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Winterthurerstrasse 190, 8057 Zurich, Switzerland

[5]Kihara Institute for Biological Research, Yokohama City University, 641-12 Maioka, 244-

0813 Totsuka-ward, Yokohama, Japan

[6]Department of Plant Life Sciences, Faculty of Agriculture, Ryukoku University, Yokotani 1-

5, Seta Oe-cho, Otsu, Shiga 520-2194, Japan

*Co-correspondence:

Atsushi J. Nagano

anagano@agr.ryukoku.ac.jp

Kentaro K. Shimizu

kentaro.shimizu@ieu.uzh.ch

ABSTRACT

An increasing number of field studies show that the phenotype of an individual plant depends not only on its genotype but also on those of neighboring plants; however, this fact is not taken into consideration in genome-wide association studies (GWAS). Based on the Ising model of ferromagnetism, we incorporated neighbor genotypic identity into a regression model in this study. The proposed method, named "neighbor GWAS", was applied to simulated and real phenotypes using *Arabidopsis thaliana* accessions. Our simulations showed that phenotypic variation explained by neighbor effects approached a plateau when an effective spatial scale became narrow. Thus, the effective scale of neighbor effects could be estimated by patterns of the phenotypic variation explained. The power to detect causal variants of neighbor effects was moderate to strong when a trait was governed by tens of variants. In contrast, there was a reasonable power down when hundreds of variants underlay a single trait. We applied the neighbor GWAS to field herbivory data on 200 accessions of *A. thaliana*, and found that the neighbor effects more largely contributed to the observed damage variation than self-genotype effects. Interestingly, several defensin family genes were associated with neighbor effects on the herbivory, while self-genotype effects were related to flavin-monooxygenase glucosinolate S-oxygenase 2 (FMO GS-OX2). Overall, the neighbor GWAS highlights the overlooked but significant role of plant neighborhood effects in shaping phenotypic variation, thereby providing a novel and powerful tool to dissect complex traits in spatially structured environments.

Keywords: *Arabidopsis thaliana*, GWAS, Ising model, Neighbor effects, Plant-insect interaction

p. 2

INTRODUCTION

Plants are immobile and thus cannot escape their neighbors. In natural and agricultural fields, individual phenotypes depend not only on the plants' own genotype but also on those of other plants in a neighborhood (Tahvanainen and Root 1972; Barbosa et al. 2009; Underwood et al. 2014). This phenomenon has been termed neighbor effects or associational effects in plant ecology (Barbosa et al. 2009; Underwood et al. 2014; Sato 2018). Such neighbor effects were initially reported as a form of interspecific interaction among different plant species (Tahvanainen and Root 1972), but many studies have thus far illustrated that neighbor effects occur among different genotypes within a plant species in herbivory (Schuman et al. 2015; Sato 2018; Ida et al. 2018), pathogen infections (Mundt 2002; Zeller et al. 2012), and pollinator visitations (Genung et al. 2012). Although neighbor effects are of considerable interest in plant science (Dicke and Baldwin 2010; Erb 2018) and its potential application to agriculture (Zeller et al. 2012; Dahlin et al. 2018), these effects are not often considered in quantitative genetics of field-grown plants.

Complex mechanisms underlie neighbor effects through direct competition (Weiner 1990), herbivore and pollinator movement (Bergvall et al. 2006; Genung et al. 2012; Verschut et al. 2016), and volatile communication among plants (Schuman et al. 2015; Dahlin et al. 2018). For example, lipoxygenase (*LOX*) genes govern jasmonate-mediated volatile emissions that induce defenses of neighboring plants in *Nicotiana* (Schuman et al. 2015). Even if direct plant-plant communications are absent, herbivores can mediate indirect interactions between plant genotypes (Sato and Kudoh 2017; Ida et al. 2018): *GLABRA1* (*GL1*) polymorphism determines hairy or glabrous phenotypes in *Arabidopsis* plants (Hauser et al. 2001) and allow a flightless leaf beetle to avoid hairy plants when encountered at a low frequency (Sato and Kudoh 2017; Sato et al. 2017). Yet, there are few hypothesis-free

p. 3

approaches to seek key genetic variants responsible for plant neighborhood effects.

Genome-wide association studies (GWAS) have been increasingly adopted to

resolve the genetic architecture of complex traits in the model plant, *Arabidopsis thaliana*

(Atwell et al. 2010; Togninalli 2018) and crop species (Hamblin et al. 2011). Plant

interactions with herbivores (Brachi et al. 2015; Nallu et al. 2018), microbes (Horton et al.

2014; Wang et al. 2018), and other plant species (Frachon et al. 2019), are one of such

complex traits dissected through the lens of GWAS. To distinguish causal variants from the

genome structure, GWAS often employs a linear mixed model with kinship considered in a

random effect (Kang et al. 2008; Korte and Farlow 2013). However, it is generally impossible

to test all the gene-by-gene interactions due to combinatorial explosion (Gondro et al. 2013);

thus, some feasible and reasonable approach should be invented for GWAS of neighbor

effects.

To incorporate neighbor effects into GWAS, we focused on a theoretical model of

neighbor effects in a magnetic field, known as the Ising model (Ising 1925; McCoy and

Maillard 2012), which has been applied to forest gap dynamics and community assembly in

plant ecology (Kizaki and Katori 1999; Schlicht and Iwasa 2004; Azaele et al. 2010).

Assuming that an individual plant is a magnet, two alleles at each locus correspond to the

north or south dipole, whereby genome-wide multiple testing across all loci is analogous to a

number of parallel two-dimensional layers. The Ising model has a clear advantage in its

interpretability, such that (i) the optimization problem for a population sum of trait values can

be regarded as an inverse problem of a simple linear model, (ii) the sign of neighbor effects

determines a model trend to generate a clustered or checkered spatial pattern of the two

states, and (iii) the self-genotypic effect determines the general tendency to favor one allele

over another (Fig. 1).

p. 4

In this study, we proposed a new methodology integrating GWAS and the Ising

model, named "neighbor GWAS". The method was applied to simulated phenotypes and real

data of field herbivory on *A. thaliana*. We addressed two specific questions: (i) what spatial

and genetic factors influenced the power to detect causal variants? (ii) were neighbor effects

significant sources of leaf damage variation in field-grown *A. thaliana*? Based on the

simulation and application, we determined the feasibility of our approach to detect neighbor

effects in field-grown plants.

MATERIALS & METHODS

Neighbor GWAS

Basic model. We analyzed neighbor effects in GWAS as an inverse problem of the two-

dimensional Ising model (Fig. 1). We consider a situation where a plant accession has either

of two alleles at each locus, and a number of accessions occupy a finite set of field sites in a

two-dimensional lattice. Let *x* represent allelic status at each locus, the allelic status at each

locus of the *i*-th focal plant and *j*-th neighboring plants can be designated as $x_{i(j)} \in \{-1, +1\}$.

Based on a two-dimensional Ising model, we can define a phenotype value of *i*-th focal

individual plant y_i as

$$y_i = \beta_1 x_i + \beta_2 \sum_{j=1}^{L} x_i x_j \qquad \text{[eq. 1]},$$

where β_1 and β_2 denote self-genotype and neighbor effects, respectively and *L* is the number

of neighboring plants to refer. If two neighboring plants shared the same allele at a given

locus, the product $x_i x_j$ turned into (-1)×(-1)=+1 or (+1)×(+1)=+1. If two neighbors had

different alleles, the product $x_i x_j$ became (-1)×(+1)=-1 or (+1)×(-1)=-1. Accordingly, the

effects of neighbor genotypic identity on a phenotype depended on the coefficient β_2 and the

p. 5

number of two alleles in a neighborhood. If the numbers of identical and different alleles

were the same near a focal plant, these neighbors offset the sum of the products $\sum_{j=1}^{L} x_i x_j$ and

exerted no effects on a phenotype. When we summed up the phenotype values for the total

number of plants n and replaced it as $E = -\beta_2$, $H = -\beta_1$ and $\epsilon_I = \sum y_i$, eq. 1 can be

transformed as $\epsilon_I = -E \sum_{<i,j>} x_i x_j - H \sum x_i$, which defines the interaction energy of a two-

dimensional ferromagnetic Ising model (McCoy and Maillard 2012). The neighbor effect β_2

and self-genotype effect β_1 were interpreted as the energy coefficient E and external magnetic

field H, respectively. An individual plant represented a spin and the two allelic states of each

locus corresponded to a north or south dipole. The positive or negative value of $\sum x_i x_j$

indicated a ferromagnetism or paramagnetism, respectively. In this study, we did not consider

the effects of allele dominance because this model was applied to inbred *A. thaliana*.

However, heterozygotes could be processed if the neighbor covariate $x_i x_j$ was weighted by an

estimated degree of dominance in the self-genotypic effects on a phenotype.

Mixed model. For association mapping, we needed to determine β_1 and β_2 from

observed phenotypes and consider a confounding sample structure as advocated by previous

GWAS (e.g., Kang et al. 2008; Korte and Farlow 2013). Extending the basic model eq. 1, we

described a linear mixed model at an individual level as

$$y_i = \beta_0 + \beta_1 x_i + \frac{\beta_2}{L} \sum_{j=1}^{L} x_i x_j + u_i + e_i \qquad \text{[eq. 2]},$$

where β_0 indicates the intercept, and the term $\beta_1 x_i$ represents fixed self-genotype effects as

tested in conventional GWAS; β_2 is the coefficient of fixed neighbor effects, and the neighbor

covariate $\sum_{j=1}^{L} x_i x_j$ is scaled by the number of neighboring plants, L. Variance components

due to a sample structure in self and neighbor effects was modeled by a random effect

$u_i \sim \text{Norm}(0, \sigma_S^2 K_S + \sigma_N^2 K_N)$. The residual was expressed as $e_i \sim \text{Norm}(0, \sigma_e^2)$.

The $n \times n$ variance-covariance matrices represented the similarity in self-genotypes

(i.e., kinship) and neighbor covariates among n individual plants as:

$$K_S = \frac{1}{q\text{-}1} X_S^T X_S \text{ and}$$

$$K_N = \frac{1}{q\text{-}1} X_N^T X_N,$$

where q indicates the number of markers. As we defined $x_{i(j)} \in \{-1, +1\}$, the elements of the

kinship matrix K_S are scaled to represent the proportion of marker loci shared among $n \times n$

plants such that $K_S = (0.5 k_{S,ij} + 0.5)$ and $k_{S,ij} = [0, 1]$; σ_S^2 and σ_N^2 indicates variance

component parameters for the self and neighbor effects.

The n plants $\times$ q markers matrix X_S and X_N are explanatory variables for the self

and neighbor effects as $X_S = (x_i)$ and $X_N = \left(\frac{\sum_{j=1}^{L} x_i x_j}{L} \right)$. The individual-level formula eq. 2 could

also be converted into a conventional matrix form as:

$$y = X\beta + Zu + e \qquad \text{[eq. 3]},$$

where y is $n \times 1$ vector of phenotypes. X is a matrix of fixed effects, including mean, self-

genotype x_i, neighbor covariate $(\sum_{j=1}^{L} x_i x_j)/L$, and other confounding covariates for n plants;

β is a vector that represents coefficients of the fixed effects; Z is a design matrix allocating

individuals to a genotype, and becomes an identity matrix if all plants are different

accessions; u is the random effect with Var(u) = $\sigma_S^2 K_S + \sigma_N^2 K_N$, and e is residual as Var(e) =

$\sigma_e^2 I$. In such a mixed model, the proportion of phenotypic variation explained (PVE) by the

self and neighbor effects could be calculated as $\text{PVE}_{\text{self}} = \sigma_S^2 / (\sigma_S^2 + \sigma_N^2 + \sigma_e^2)$ and $\text{PVE}_{\text{nei}} =$

$\sigma_N^2 / (\sigma_S^2 + \sigma_N^2 + \sigma_e^2)$, respectively. The line of extensions to incorporate neighbor effects into

GWAS is referred to as "neighbor GWAS" hereafter.

Power simulation

To examine the power to detect neighbor effects, we applied the neighbor GWAS to

simulated phenotypes. Phenotypes were simulated using a part of the real genotypes of *A.*

thaliana. To evaluate the power of the simple linear model, we assumed a complex ecological

form of neighbor effects with multiple variance components controlled. The power was

evaluated by the receiver operating characteristic (ROC) curve on the association score of -

$\log_{10}(p$-value) (e.g., Gage et al. 2018). All analyses were performed using R version 3.4.0 (R

Core Team 2017).

Genotype data. To consider a realistic genetic structure in the simulation, we used

part of the *A. thaliana* RegMap panel (Horton et al. 2012). The genotype data on 1307

accessions were downloaded from the Joy Bergelson laboratory website

(http://bergelson.uchicago.edu/?page_id=790 accessed on 9 February 2017). We extracted the

chromosome 1 and 2 data with the minor allele frequency (MAF) at >0.1, providing a matrix

of 1307 plants with 65,226 single nucleotide polymorphisms (SNPs). Pairwise linkage

disequilibrium (LD) among the loci was $r^2 = 0.003$ [0.00-0.06: median with upper and lower

95 percentiles]. Before generating a phenotype, each locus was centered by its mean and

scaled by its standard deviation. Subsequently, we randomly selected 1296 accessions (= 36 ×

36 accessions) without any replacements for each iteration, and placed them in a 36 × 72

checkered space following *Arabidopsis* experimental settings (see Fig. 2b).

Phenotype simulation. To address ecological issues specific to plant neighborhood

effects, we considered two extensions, namely asymmetric neighbor effects and spatial

scales. Firstly, previous studies showed that such plant-plant interactions are sometimes

asymmetric between two accessions in herbivory (e.g., Bergvall et al. 2006; Verschut et al.

2016; Sato and Kudoh 2017) and height competition (Weiner 1990). Such asymmetric

p. 8

neighbor effects can be tested by statistical interactions terms in a linear model (Bergvall et

al. 2006; Sato and Kudoh 2017). Secondly, several studies showed that the strength of

neighbor effects depended on spatial scales (Hambäck et al. 2014) and the scale of neighbors

to be analyzed relied on the dispersal ability of causative organisms (see Hambäck et al.

2009; Sato and Kudoh 2015; Vershut et al. 2018; Ida et al. 2018 for insect and mammal

herbivores; Rieux et al. 2014 for pathogen dispersal) or the size of competing plants (Weiner

1990). We assumed the distance decay at s-th sites from a focal individual i with the decay

coefficient α as $w(s, \alpha) = e^{-\alpha(s-1)}$, since such an exponential distance decay has been widely

adopted in empirical studies (Devaux et al. 2007; Carrasco et al. 2010; Rieux et al. 2014; Ida

et al. 2018). Therefore, we assumed a more complex model for simulated phenotypes than the

model for neighbor GWAS as follows:

$$y_i = \beta_0 + \beta_1 x_i + \frac{\beta_2}{L}\sum_{j=1}^{L} w(s, \alpha)x_i x_j + \beta_{12}\frac{x_i}{L}\sum_{j=1}^{L} w(s, \alpha)x_i x_j + u_i + e_i \quad [\text{eq. 4}],$$

where β_{12} is the coefficient for asymmetry in neighbor effects. Total variance components due

to the three background effects i.e., the self, neighbor, and self-by-neighbor effects is defined

as $u_i \sim \text{Norm}\left(0, \sigma_S^2 K_S + \sigma_N^2 K_N + \sigma_{SxN}^2 K_{SxN}\right)$. The three variance component parameters σ_S^2, σ_N^2,

and σ_{SxN}^2, determined the relative importance of self-genotype, neighbor, and asymmetric

neighbor effects in u_i. Given the n plants $\times$ q marker explanatory matrix with

$X_{S\times N} = (\frac{x_i}{L}\sum_{j=1}^{L} w(s, \alpha)x_i x_j)$, the similarity in asymmetric neighbor effects was calculated as

$K_{S\times N} = \frac{1}{q\text{-}1}X_{S\times N}^{\text{T}}X_{S\times N}$. To control phenotypic variations, we further partitioned the proportion

of phenotypic variation into those explained by major-effect genes and variance components

$\text{PVE}_\beta + \text{PVE}_u$, major-effect genes alone PVE_β, and residual error PVE_e, where $\text{PVE}_\beta + \text{PVE}_u$

$+ \text{PVE}_e = 1$. The *optimize* function in R was used to adjust simulated phenotypes to given

amounts of PVE.

p. 9

Parameter setting. Fifteen phenotypes were simulated for each combination of the

distance decay α, the proportion of phenotypic variance explained by major-effect genes

PVE$_{\beta}$, variance components PVE$_{u}$, and the relative importance of multiple variance

components $\sigma_S^2{:}\sigma_N^2{:}\sigma_{SxN}^2$ as: $\alpha = 0.25$, 1.0 or 3.0, $\sigma_S^2{:}\sigma_N^2{:}\sigma_{SxN}^2 = 6{:}3{:}1$, 4:4:1, or 3:6:1, PVE$_{\beta} =$

0.1, 0.3 or 0.6, and PVE$_{\beta}$ + PVE$_{u}$ = 0.4 or 0.8. The maximum reference scale was fixed at $s =$

3. The line of simulations was repeated for the number of causal SNPs at 20 or 200 to

examine cases of an oligogenic and polygenic control for a trait. The non-zero coefficients for

the causal SNPs were randomly sampled from a uniform distribution, Unif($|0.5|$, $|2.0|$), and

assigned as some causal SNPs were responsible for both the self and neighbor effects. Of the

total number of causal SNPs, 15% had all self, neighbor, and asymmetric neighbor effects

(i.e., $\beta_1 \neq 0$ and $\beta_2 \neq 0$ and $\beta_{12} \neq 0$); another 15% had both the self and neighbor effects, but no

asymmetry in the neighbor effects ($\beta_1 \neq 0$ and $\beta_2 \neq 0$ and $\beta_{12} = 0$); another 35% had self-

genotypic effects only ($\beta_1 \neq 0$); and the remaining 35% had neighbor effects alone ($\beta_2 \neq 0$).

Given its biological significance, we assumed that some loci having neighbor signals

possessed asymmetric interactions between neighbors ($\beta_2 \neq 0$ and $\beta_{12} \neq 0$) while the others

had symmetric ones ($\beta_2 \neq 0$ and $\beta_{12} = 0$). Therefore, the number of causal SNPs in β_{12} was

smaller than that in the main neighbor effects β_2. According to this assumption, the variance

component σ_{SxN}^2 was also assumed to be smaller than σ_N^2.

Summary statistics. The simulated phenotypes were fitted by eq. 2 to test the

significance of coefficients β_1 and β_2, and to estimate the variance component due to self and

neighbor effects PVE$_{self}$ and PVE$_{nei}$. The stepwise likelihood ratio tests were performed first

between the null model and model with a self-genotype effect alone, and then between the

self-genotype effect model and model with both self and neighbor effects. The likelihood

ratio was calculated as the difference in deviance i.e., -2 × log-likelihood, which is

p. 10

asymptotically χ^2 distributed with one degree of freedom. The variance components, PVE_{self}

and PVE_{nei}, were estimated using the average information restricted maximum likelihood

(AI-REML) algorithm implemented in the *lmm.aireml* function in the *gaston* package of R

(Perdry and Dandine-Roulland 2018). Subsequently, the two variance parameters σ_S^2 and

σ_N^2 were replaced with their estimates $\hat{\sigma}_S^2$ and $\hat{\sigma}_N^2$ by the AI-REML, and association tests

were performed by solving linear mixed models with a fast approximation by an eigen value

decomposition (implemented in the *lmm.diago* function: Perdry and Dandine-Roulland

2018). The likelihood was computed using the *lmm.diago.profile.likelihood* function to test β_1

or β_2. True or false positive rates were evaluated by ROC curves and area under the ROC

curves (AUC) (Gage et al. 2018). An AUC of 0.5 would indicate that GWAS has no power to

detect true signals, while an AUC of 1.0 would indicate that all the top signals predicted by

GWAS agree with true signals. The roc function in the pROC package (Robin et al. 2011)

was used to calculate AUC from $-\log_{10}(p\text{-value})$. The AUC and variance components were

calculated from $s = 1$ (the first nearest neighbors) to $s = 3$ (up to the third nearest neighbors)

cases. The AUCs were also calculated using standard linear models without any random

effects to examine whether the linear mixed models were superior to the linear models.

***Arabidopsis* herbivory data**

We applied the neighbor GWAS to field data of *Arabidopsis* herbivory. This field experiment

followed our previous publication on a summer herbivory on field-grown *A. thaliana* (Sato et

al. 2019). We used 200 worldwide accessions comprising the RegMap (Horton et al. 2012)

and 1001 Genomes project (Alonso-Blanco et al. 2016), of which most were overlapped with

a previous GWAS of biotic interactions (Horton et al. 2014) and half were included by a

GWAS of glucosinolates (Chan et al. 2010). Eight replicates of the 200 accessions were first

prepared in a laboratory and then transferred to the outdoor garden at the Center for

Ecological Research, Kyoto University, Japan (Otsu, Japan: 35º 06' N, 134º 56' E, alt. ca. 200

m: Fig. 2a). Seeds were sown on Jiffy-seven pots (33-mm diameter), and stratified under 4 ℃

for a week. Seedlings were cultivated for 1.5 months under a short-day condition (8 h light:

16 h dark, 20 ℃). Plants were then separately potted in plastic pots (6 cm in diameter) filled

with mixed soil of agricultural composts (Metro-mix 350, SunGro Co., USA) and perlites at a

3:1 L ratio. In the field setting, 200 accessions were randomly assigned in a checkered

manner within a block (Fig. 2b). Eight replicates of these blocks were set >2 m apart from

each other (Fig. 2c). Potted plants were exposed to the field environment for 3 wk in June

2017. At the end of experiment, we scored leaves eaten as 0 for no visible damage, 1 for

$\leq$10%, 2 for >10% and $\leq$ 25%, 3 for > 25% and $\leq$ 50%, 4 for >50% and $\leq$ 75%, and 5 for

>75% of the leaf area eaten. All plants were scored by a single person to avoid observer bias.

The most predominant herbivore in this field trial was the diamond back moth *Plutella*

xylostella, followed by the small white butterfly *Pieris rapae*. We also recorded the initial

plant size and the presence of inflorescence to incorporate them as covariates. Initial plant

size was evaluated by the length of the largest rosette leaf (mm) at the beginning of the field

experiment and the presence of inflorescence was recorded 2 wk after transplanting.

We estimated the variance components and performed the association tests for the

leaf damage score with the neighbor covariate at $s = 1$ and 2. These two scales corresponded

to $L = 4$ (the nearest four neighbors) and $L = 12$ (up to the second nearest neighbors),

respectively, in the *Arabidopsis* dataset. The variation partitioning and association tests were

performed using the *gaston* package, as mentioned above. To determine the significance of

variance component parameters, we compared the likelihood between mixed models with one

or two random effects. For the genotype data, we used an imputed SNP matrix of all the 2029

accessions studied by the RegMap (Horton et al. 2012) and 1001 Genomes project (Alonso-

Blanco et al. 2016). Missing genotypes were imputed using BEAGLE (Browning and

Browning 2009), as described by Togninalli et al. (2018) and updated on the AraGWAS

Catalog (https://aragwas.1001genomes.org). Of the 10,709,466 SNPs from the full imputed

matrix, we used 1,242,128 SNPs with MAF at > 0.05 and LD of adjacent SNPs at $r^2 < 0.8$. We

considered the initial plant size, presence of inflorescence, and experimental blocks as fixed

covariates. After the association mapping, we searched candidate genes within ~10 kb around

target SNPs, based on the Araport11 gene model with the latest TAIR annotation (accessed on

7 September 2019). Gene ontology (GO) enrichment analysis was applied to the candidate

genes near the top 0.1% SNP score. GO categories including >20 and <200 genes were tested

by Fisher's exact probability tests and adjusted by false discovery rate (FDR: Benjamini and

Hochberg 1995). The GO.db package (Carlson et al. 2018) and the latest TAIR AGI code

annotation were used for the GO enrichment analysis. The R source codes, accession list, and

phenotype data are available at the GitHub repository

(https://github.com/naganolab/NeighborGWAS).

RESULTS

Power simulation

A set of phenotypes were simulated from real genotype data following a complex model eq.

4, and then fitted by a simplified model eq. 2. Analyzing the factors affecting AUCs, we

found that the proportion of phenotypic variation (PVE) explained by major-effect genes

PVE_{β} and distance decay of neighbor effects were the most influential on the power to detect

neighbor signals (Table 1b, d). In addition to PVE_{β}, the amount of variance components PVE_{u}

p. 13

also significantly affected the AUCs of the self and neighbor effects, but these additional

effects were less significant compared to those of PVE_β alone (Table 1). In contrast, the

AUCs of neither self nor neighbor effects were significantly affected by the ratio of three

variance components $\sigma_S^2 : \sigma_N^2 : \sigma_{SxN}^2$ (Table 1).

Notably, there was a clear relationship between the distance decay α and the

proportion of phenotypic variation explained by neighbor effects PVE_{nei} or AUCs at different

spatial scales (Fig. 3). If the distance decay was weak and the effective range of neighbor

effects was broad, PVE_{nei} and AUCs increased linearly as the reference spatial scale was

broadened (Fig. 3a). On the other hand, if the distance decay was strong and the effective

scale of neighbor effects was narrow, PVE_{nei} saturated at the scale of the first nearest

neighbors (Fig. 3c) or AUCs did not increase (Fig. 3b, c). These results remained the same

between the number of causal SNPs = 20 and 200 (Fig. 3 and Fig. S1). The line of simulation

results indicated that the effective spatial scales could be estimated by calculating PVE_{nei}

across different spatial scales.

In the case of the number of causal SNPs = 20, signals of major-effect genes were

well detected as AUC ranged from moderate (>0.7) to high (>0.9) (Fig. S2). For the case of

the number of causal SNPs = 200, it became relatively difficult to detect the major-effect

genes as AUCs were ≤ 0.75 (Fig. S2). The line of simulations indicated that neighbor effects

were detectable when a target trait was governed by several major genes and the range of

neighbor effects was spatially limited. Additionally, linear mixed models outperformed

standard linear models in the detection of self and neighbor signals ($\Delta AUC_{self} = 0.105$ [0.101

− 0.109], $\Delta AUC_{nei} = 0.024$ [0.021 − 0.026]: 10,000-times bootstrap mean with 95%

confidence intervals). This indicated that the mixed models were more appropriate for the

neighbor GWAS to deal with spurious associations due to a sample structure.

p. 14

Arabidopsis herbivory data

The variation partitioning of leaf damage showed that the PVE by neighbor effects were

larger than PVE by self-genotypic effects ($PVE_{self} = 0.026$, $\chi_1^2 = 0.151$, p-value = 0.70;

$PVE_{nei} = 0.218$, $\chi_1^2 = 7.17$, p-value = 0.0074: Fig. 4a). Heritability, namely PVE_{self} without

neighbor effects, was 0.159 ($\chi_1^2 = 8.73$, p-value = 0.003: Fig. S3). This range of heritability

was overlapped with PVE by neighbor effects alone (PVE_{nei} without self-effects = 0.24 at

scale $s = 1$, $\chi_1^2 = 15.7$, p-value < 0.0001: Fig. S3), indicating that there was an intersection

between PVE by self and neighbor effects on the leaf damage variation. Phenotypic variation

explained by neighbor effects on leaf damage did not increase when the neighbor scale was

referred up to the nearest and second nearest individuals ($PVE_{self} = 0.083$, $\chi_1^2 = 1.03$, p-value

= 0.311; $PVE_{nei} = 0.13$, $\chi_1^2 = 1.29$, p-value = 0.256: Fig. 4a); therefore, the variation

partitioning was stopped at $s = 2$. These results indicated a narrow effective scale and

significant contribution of neighbor effects to the leaf damage score.

Association mapping of the self-genotype effects on the leaf damage found a SNP

with the largest $-\log_{10}(p$-values) score at "chr1-23149476". This SNP was located within ~10

kb of the AT1G62540 locus that encoded flavin-monooxygenase glucosinolate S-oxygenase 2

(FMO GS-OX2), though this was not above a threshold of Bonferroni correction. Gene

ontology annotation of "cellular response to extracellular stimulus" was marginally enriched

among genes within ~10 kb around SNPs with the top 0.1% $-\log_{10}(p$-values) score which

corresponded to p-values at < 0.00096 (FDR<0.1: Table 2a). A QQ-plot did not exhibit an

inflation of p-values for the self-genotype effects (Fig. S4).

We found a marginally significant SNP for neighbor effects at the second and third

chromosome (Fig. 4c), of which the second chromosomal region had higher association

p. 15

scores than expected by the QQ-plot (Fig. S4). A locus encoding FAD-binding Berberine

family protein (AT2G34810 named *BBE16*) were located within the ~10 kb window near the

SNP with the largest -log$_{10}$(*p*-values) at the second chromosome, which are known to be up-

regulated by methyl jasmonate (Devoto et al. 2005). Three transposable elements and a

pseudogene of lysyl-tRNA synthetase 1 were located near the most significant SNP at the

third chromosome. These top ten SNPs significantly related to the neighbor effects exhibited

positive estimates of β_1 and β_2. Three defense-related GO annotations of "killing cells of

other organisms" and "disruption of cells of other organism" were significantly enriched

among genes within ~10 kb around SNPs with the top 0.1% score of -log$_{10}$(*p*-values)

(FDR<0.05: Table 2b). Of the genes with these GO annotations, we found 22 low-molecular

weight cysteine-rich proteins or plant defensin family proteins (Table S2).

Based on the estimated coefficients $\hat{\beta}_1$ and $\hat{\beta}_2$, we ran a post hoc simulation to infer

a spatial arrangement that minimizes a population sum of the leaf damage $\sum y_i = \beta_1 \sum x_i +$

$\beta_2 \sum_{<i,j>} x_i x_j$. The constant intercept β_0, the variance component u_i, and residual e_i were not

considered because they were not involved in deterministic dynamics of the model. Figure 5

shows three representatives and a neutral expectation. For example, a mixture of a

dimorphism was expected to decrease the total leaf damage for a SNP at "chr2-14679190"

near the *BBE16* locus ($\hat{\beta}_2$>0: Fig. 5a). On the other hand, a clustered distribution of a

dimorphism was expected to decrease the total damage for a SNP at "chr2-9422409" near the

AT2G22170 locus encoding a lipase/lipooxygenase PLAT/LH2 family protein ($\hat{\beta}_1$≈0, $\hat{\beta}_2$<0:

Fig. 5b). Furthermore, near monomorphism was expected to decrease the leaf damage for a

SNP at "chr5-19121831" near the AT5G47075 and AT5G47077 loci encoding low-molecular

cysteine-rich proteins, LCR20 and LCR6 ($\hat{\beta}_1$>0, $\hat{\beta}_2$<0: Fig. 5c). No self and neighbor effects

p. 16

led to a random distribution and no mitigation of damage i.e., $\sum y_i \approx 0$ (Fig. 5d). These post

hoc simulations suggested a potential applicability of neighbor GWAS in optimizing spatial

arrangements in field cultivation.

DISCUSSION

Spatial and genetic factors affecting the power to detect signals

Benchmark tests using simulated phenotypes revealed that appropriate spatial scales could be

estimated by variation partitioning of observed phenotypes. When the scale of neighbor

effects was narrow or moderate (α = 1.0 or 3.0), the scale of the first nearest neighbors would

be optimum for the power to detect neighbor signals. In terms of neighbor effects in plant

defense, mobile animals, such as mammalian browsers and flying insects, often select a

cluster of plant individuals (e.g., Bergvall et al. 2006; Hambäck et al. 2009; Vershut et al.

2016); however, neighbor effects could not be detected among individual plants within a

cluster (Hambäck et al. 2014; Sato and Kudoh 2015). This case was represented by the

exponential distance decay of α = 0.25; only in such a special case should more than the first

nearest be referred to gain the power.

Neighbor GWAS could retain its power as long as neighbor effects were spatially

limited and several major-effect genes governed a trait. In contrast, when hundreds of causal

variants involved a single trait and less than half of phenotypic variation was attributable to

neighbor effects, we observed a reasonable power down of neighbor GWAS. In GWAS, false

positive rates can be reduced using linear mixed models that deal with kinship structure as a

random effect (Korte and Farlow 2013). Indeed, mixed models were superior to standard

linear models in this simulation. Our simulation also adjusted the three variance components

p. 17

σ_S^2, σ_N^2, and σ_{SxN}^2, but their relative contribution did not have significant effects on the power.

This was likely due to the fact that the self-genotypic variable x_i was encompassed into the

neighbor variable $\sum x_i x_j / L$, and thus the kinship matrix $\boldsymbol{K}_S$ was partially redundant with the

similarity matrix of neighbor effects $\boldsymbol{K}_N$. Indeed, elemental-wise correlations between $\boldsymbol{K}_S$

and $\boldsymbol{K}_N$ were strong in our simulations ($R^2 > 0.7$). Thus, linear mixed models gain the power

to detect neighbor effects if signals are strong, but likelihood ratio tests are reliable enough to

deal with the correlated variables.

Candidate genes related to field herbivory on *Arabidopsis*

Our *Arabidopsis* data successfully detected known defense-related genes involved in the self-

genotype effects on leaf damage. Aliphatic glucosinolates are a major chemical defense

against insect herbivory (Brachi et al. 2015; Kerwin et al. 2017). Specifically, FMO GS-OX2

is involved in aliphatic glucosinolate biosynthesis by catalyzing the conversion from

methylthioalkyl to corresponding methylsulfinylalkyl glucosinolates (Li et al. 2008).

Furthermore, previous GWAS reported methionine synthase 2 (AT3G03780), disease

resistance protein (TIR-NBS-LRR class) family (AT4G16950), and monodehydroascorbate

reductase 4 (AT3G27820) as candidate genes involved in self-resistance to the white butterfly

Pieris rapae (Davila-Olivas et al. 2017; Nallu et al. 2018). In this field experiment, we

observed larvae of *P. rapae* largely feeding on *A. thaliana*, and the GWAS of self-genotype

effects on leaf damage detected the above three candidate genes near SNPs with the top 0.1%

association score. Thus, our GWAS results seemed convincing in terms of the detection of

known defense-related genes in the self-genotypic effects on herbivory.

Notably, the neighbor effects in herbivory were relevant to candidate genes

disrupting cells of other organisms. Plant defensins are stable and cysteine-rich peptides that

confer plant resistance by killing cells of other organisms (Stotz 2009). While anti-fungal

resistance is a well-known function of plant defensins (Stotz 2009), they can also act as

protease inhibitors against insect herbivores (Bloch and Richardson 1991; Pelegrini et al.

2008; Choi et al. 2009). Typical examples of neighbor effects may be a direct induction of

plant defense via volatile organic chemicals (e.g., Schuman et al. 2015; Dahlin et al. 2018),

but ecological studies have shown that herbivore host choice is one of the most important

processes leading indirect neighbor effects to anti-herbivore defenses (Bergvall et al. 2006;

Verschut et al. 2016; Sato et al. 2017). The findings of our neighbor GWAS suggest a putative

role of plant defensins in modulating insect feeding behaviors and thus neighbor effects in

herbivory.

Conclusion and applicability

Based on the newly proposed methodology, we suggest that neighbor effects are a more

important source of phenotypic variation in field-grown plants than currently appreciated. To

date, regional-scale interactions among plants have been analyzed using a genome-

environment association study of plant community composition (Frachon et al. 2019), but

fine-scale neighbor effects have yet to be examined. Using tens of *A. thaliana* accessions and

their experimentally mixed populations, Wuest and Niklaus (2018) recently showed that a

single genomic region drives neighbor effects in plant growth via soil improvements, and this

genetic effect shapes a positive relationship between plant genotype diversity and

productivity. Our newly proposed methodology of neighbor GWAS could be a powerful tool

to identify such a key genetic variant responsible for neighbor effects and resulting

biodiversity effects.

Neighbor GWAS may also potentially help determine an optimal spatial arrangement

p. 19

in plant cultivation, as suggested by the post hoc simulation. The Ising model is well

established in statistical physics (McCoy and Maillard 2012) and is now applied to a

machine-learning pipeline that deals with high-dimensionality in genomics data (Fisher and

Mehta 2015). Genome-wide polymorphism data are useful not only to seek causal genes in

GWAS, but also to predict breeding values of crop plants in genomic selection (e.g., Jannink

et al. 2010; Hamblin et al. 2011; Yamamoto et al. 2017). Although it is still challenging to

determine β_1 and β_2 for all SNPs efficiently, the linear model of neighbor GWAS could also

be implemented as a genomic selection at a population level. Thus, our study provides an

avenue for future studies to predict population-level phenotypes in spatially structured

environments.

ACKNOWLEDGEMENTS

The authors thank Ü. Seren, A. Korte, and M. Nordborg for kindly providing the full imputed SNP data; and T. Tsuchimatsu and K. Iwayama for discussions. This study was supported by the Japan Science and Technology Agency (JST) PRESTO (Grant number, JPMJPR17Q4 and JPMJPR16Q9) to Y.S. and E.Y.; Japan Society for the Promotion of Science (JSPS) Postdoctoral Fellowship (16J30005) to Y.S.; MEXT KAKENHI (18H04785) and the Swiss National Science Foundation to K.K.S.; and JST CREST (JPMJCR15O2 and JPMJCR16O3) to A.J.N. and K.K.S. The field experiment was supported by the Joint Usage/Research Grant of Center for Ecological Research, Kyoto University, Japan.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interests in this study.

DATA ARCHIVING

The leaf damage data on *A. thaliana* are included in the supporting information (Table S1). The R source codes used in this study are available at the GitHub repository (https://github.com/naganolab/NeighborGWAS).

REFERENCES

1. Alonso-Blanco C, Andrade J, Becker C, Bemm F, Bergelson J, Borgwardt KM. et al. (2016). 1,135 genomes reveal the global pattern of polymorphism in *Arabidopsis thaliana*. Cell 166: 481–491. doi:10.1016/j.cell.2016.05.063

2. Atwell S, Huang YS, Vilhjálmsson BJ, Willems G, Horton M, Li Y. et al. (2010). Genome-wide association study of 107 phenotypes in *Arabidopsis thaliana* inbred lines.

p. 21

487  Nature 465: 627–631.

488 3. Azaele S, Muneepeerakul R, Rinaldo A, Rodriguez-Iturbe I. (2010). Inferring plant

489  ecosystem organization from species occurrences. J Theor Biol **262**: 323–329.

490  doi:10.1016/j.jtbi.2009.09.026

491 4. Barbosa P, Hines J, Kaplan I, Martinson H, Szczepaniec A, Szendrei Z. (2009).

492  Associational resistance and associational susceptibility: Having right or wrong

493  neighbors. Ann Rev Ecol Evol Sys 40: 1–20.

494 5. Benjamini Y, Hochberg Y. (1995). Controlling the false discovery rate: a practical and

495  powerful approach to multiple testing. J Royal Stat Soc B 57: 289–300.

496  doi:10.1111/j.2517-6161.1995.tb02031.x

497 6. Bergvall UA, Rautio P, Kesti K, Tuomi J, Leimar O. (2006). Associational effects of

498  plant defences in relation to within- and between-patch food choice by a mammalian

499  herbivore: neighbour contrast susceptibility and defence. Oecologia 147: 253–260.

500  doi:10.1007/s00442-005-0260-8

501 7. Bloch C, Richardson M. (1991). A new family of small (5 kDa) protein inhibitors of

502  insect α-amylases from seeds or sorghum (*Sorghum bicolor* (L) Moench) have sequence

503  homologies with wheat γ-purothionins. FEBS letters 279: 101-104.

504 8. Brachi B, Meyer CG, Villoutreix R, Platt A, Morton TC, Roux F, Bergelson J. (2015).

505  Coselected genes determine adaptive variation in herbivore resistance throughout the

506  native range of *Arabidopsis thaliana*. Proc Natl Acad Sci USA 112: 4032–4037.

507 9. Browning BL, Browning SR. (2009). A unified approach to genotype imputation and

508  haplotype-phase inference for large data sets of trios and unrelated individuals. Am J

509  Hum Genet 84: 210-223.

510 10. Carlson M. (2018). GO.db: A set of annotation maps describing the entire Gene

p. 22

511  Ontology. R package version 3.7.0.

512 11. Carrasco LR, Harwood TD, Toepfer S, MacLeod A, Levay N, Kiss J, Baker RHA,

513   Mumford JD, Knight JD. (2010). Dispersal kernels of the invasive alien western corn

514   rootworm and the effectiveness of buffer zones in eradication programmes in Europe.

515   Ann Appl Biol 156: 63–77. doi:10.1111/j.1744-7348.2009.00363.x

516 12. Chan EKF, Rowe HC, Kliebenstein DJ. (2010). Understanding the evolution of defense

517   metabolites in *Arabidopsis thaliana* using genome-wide association mapping. Genetics

518   185: 991–1007.

519 13. Choi MS, Kim YH, Park HM, Seo BY, Jung JK, Kim ST. et al. (2009). Expression of

520   BrD1, a plant defensin from *Brassica rapa*, confers resistance against brown planthopper

521   (*Nilaparvata lugens*) in transgenic rices. Mol Cells 28: 131–137. doi:10.1007/s10059-

522   009-0117-9.

523 14. Dahlin I, Rubene D, Glinwood R, Ninkovic V. (2018). Pest suppression in cultivar

524   mixtures is influenced by neighbor-specific plant-plant communication. Ecol Appl 28:

525   2187–2196. doi:10.1002/eap.1807

526 15. Devaux C, Lavigne C, Austerlitz F, Klein EK. (2007). Modelling and estimating pollen

527   movement in oilseed rape (*Brassica napus*) at the landscape scale using genetic markers.

528   Mol Ecol 16: 487–499. doi:10.1111/j.1365-294X.2006.03155.x

529 16. Davila-Olivas NH, Kruijer W, Gort G, Wijnen CL, van Loon JJA, Dicke M. (2017).

530   Genome-wide association analysis reveals distinct genetic architectures for single and

531   combined stress responses in *Arabidopsis thaliana*. New Phytol 213: 838–851.

532   doi:10.1111/nph.14165

533 17. Devoto A, Ellis C, Magusin A, Chang HS, Chilcott C, Zhu T, Turner JG. (2005).

534   Expression profiling reveals *COI1* to be a key regulator of genes involved in wound- and

p. 23

535  methyl jasmonate-induced secondary metabolism, defence, and hormone interactions.

536  Plant Mol Biol 58: 497–513.

537 18. Dicke M, Baldwin IT. (2010). The evolutionary context for herbivore-induced plant

538   volatiles: beyond the 'cry for help'. Trends Plant Sci 15: 167–175.

539   doi:10.1016/j.tplants.2009.12.002

540 19. Erb M. (2018). Volatiles as inducers and suppressors of plant defense and immunity -

541   origins, specificity, perception and signaling. Curr Opin Plant Biol 44: 117–121.

542   doi:10.1016/j.pbi.2018.03.008

543 20. Frachon L, Mayjonade B, Bartoli C, Hautekèete NC, Roux F. (2019). Adaptation to plant

544   communities across the genome of *Arabidopsis thaliana*. Mol Biol Evol 36: 1442–1456.

545 21. Fisher CK, Mehta P. (2015). Bayesian feature selection for high-dimensional linear

546   regression via the Ising approximation with applications to genomics. Bioinformatics **31**:

547   1754–1761.

548 22. Gage JL, De Leon N, Clayton MK. (2018). Comparing genome-wide association study

549   results from different measurements of an underlying phenotype. G3: Genes, Genomes,

550   Genetics 8: 3715–3722.

551 23. Genung MA, Bailey JK, Schweitzer JA. (2012). Welcome to the neighbourhood:

552   interspecific genotype by genotype interactions in *Solidago* influence above- and

553   belowground biomass and associated communities. Ecol Lett 15: 65–73.

554 24. Gondro C, van der Werf J, Hayes B. (eds.) (2013). Genome-wide association studies and

555   genomic prediction. Methods Mol Biol. Humana Press, New York.

556 25. Hambäck PA, Björkman M, Rämert B, Hopkins RJ. (2009). Scale-dependent responses

557   in cabbage herbivores affect attack rates in spatially heterogeneous systems. Basic Appl

558   Ecol 10: 228–236. doi:10.1016/j.baae.2008.06.004

p. 24

26. Hambäck PA, Inouye BD, Andersson P, Underwood N. (2014). Effects of plant

neighborhoods on plant-herbivore interactions: resource dilution and associational

effects. Ecology 95: 1370–1383. doi:10.1890/13-0793.1

27. Hamblin MT, Buckler ES, Jannink JL. (2011). Population genetics of genomics-based

crop improvement methods. Trends Genet 27: 98–106.

28. Hauser MT, Harr B, Schlötterer C. (2001). Trichome distribution in *Arabidopsis thaliana*

and its close relative *Arabidopsis lyrata*: molecular analysis of the candidate gene

GLABROUS1. Mol Biol Evol 18: 1754–1763.

29. Horton MW, Hancock AM, Huang YS, Toomajian C, Atwell S, Auton A, et al. (2012).

Genome-wide patterns of genetic variation in worldwide *Arabidopsis thaliana* accessions

from the RegMap panel. Nat Genet 44: 212–216. doi:10.1038/ng.1042

30. Horton MW, Bodenhausen N, Beilsmith K, Meng D, Muegge BD, Subramanian S, et al.

(2014). Genome-wide association study of *Arabidopsis thaliana* leaf microbial

community. Nat Commun 5: 5320. doi:10.1038/ncomms6320

31. Ida TY, Takanashi K, Tamura M, Ozawa R, Nakashima Y, Ohgushi T. (2018). Defensive

chemicals of neighboring plants limit visits of herbivorous insects: Associational

resistance within a plant population. Ecol Evol 8: 12981–12990. doi:10.1002/ece3.4750

32. Ising E. (1925). Beitrag zur theorie des ferromagnetismus. *Zeitschrift für Physik* 31: 253–

258.

33. Jannink JL, Lorenz AJ, Iwata H. (2010). Genomic selection in plant breeding: from

theory to practice. Brief Funct Genomic 9: 166–177.

34. Kang HM, Zaitlen NA, Wade CM, Kirby A, Heckerman D, Daly MJ, Eskin E. (2008).

Efficient control of population structure in model organism association mapping.

Genetics 178: 1709–1723.

35. Kerwin RE, Feusier J, Muok A, Lin C, Larson B, Copeland D. et al. (2017). Epistasis ×

environment interactions among *Arabidopsis thaliana* glucosinolate genes impact

complex traits and fitness in the field. New Phytol 215: 1249–1263.

36. Kizaki S, Katori M. (1999). Analysis of canopy-gap structures of forests by Ising-Gibbs

states-equilibrium and scaling property of real forests. J Phys Soc Japan **68**: 2553-2560.

37. Korte A, Farlow A. (2013). The advantages and limitations of trait analysis with GWAS:

a review. Plant Methods 9: 29.

38. Li J, Hansen BG, Ober JA, Kliebenstein DJ, Halkier BA. (2008). Subclade of flavin-

monooxygenases involved in aliphatic glucosinolate biosynthesis. Plant Physiol 148:

1721–1733.

39. McCoy BM, Maillard JM. (2012). The importance of the Ising model. Progress Theor

Phys 127: 791-817.

40. Mundt CC. (2002). Use of multiline cultivars and cultivar mixtures for disease

management. Ann Rev Phytopathol 40: 381-410.

41. Nallu S, Hill JA, Don K, Sahagun C, Zhang W, Meslin C. et al. (2018). The molecular

genetic basis of herbivory between butterflies and their host plants. Nat Ecol Evol 2:

1418-1427. doi:10.1038/s41559-018-0629-9

42. Pelegrini PB, Lay FT, Murad AM, Anderson MA, Franco OL. (2008). Novel insights on

the mechanism of action of A-amylase inhibitors from the plant defensin family.

Proteins: Struct Funct Bioinformatics 73: 719–729. doi:10.1002/prot.22086.

43. Perdry H, Dandine-Roulland C. (2018). gaston: Genetic Data Handling (QC, GRM, LD,

PCA) & Linear Mixed Models. R package version 1.5.4. https://CRAN.R-

project.org/package=gaston

44. R Core Team. (2017). R: A language and environment for statistical computing. R

p. 26

Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

45. Rieux A, Soubeyrand S, Bonnot F, Klein EK, Ngando JE, Mehl A. et al. (2014). Long-

distance wind-dispersal of spores in a fungal plant pathogen: estimation of anisotropic

dispersal kernels from an extensive field experiment. PLoS ONE 9: e103225.

doi:10.1371/journal.pone.0103225

46. Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC, Muller M. (2011).

pROC: an open-source package for R and S+ to analyze and compare ROC curves. *BMC*

Bioinformatics **12**: 77. doi:10.1186/1471-2105-12-77

47. Schlicht R, Iwasa Y. (2004). Forest gap dynamics and the Ising model. J Theoret Biol

230: 65–75. doi:10.1016/j.jtbi.2004.04.027

48. Sato Y. (2018). Associational effects and the maintenance of polymorphism in plant

defense against herbivores: review and evidence. Plant Species Biol 33: 91–108.

doi:10.1111/1442-1984.12201

49. Sato Y, Kudoh H. (2015). Tests of associational defence provided by hairy plants for

glabrous plants of *Arabidopsis halleri* subsp. *gemmifera* against insect herbivores. Ecol

Entomol 40: 269–279. doi:10.1111/een.12179

50. Sato Y, Kudoh H. (2017). Herbivore-mediated interaction promotes the maintenance of

trichome dimorphism through negative frequency-dependent selection. Am Nat 190:

E67–E77. doi:10.1086/692603

51. Sato Y, Ito K, Kudoh H. (2017). Optimal foraging by herbivores maintains

polymorphism in defence in a natural plant population. Funct Ecol 31: 2233–2243.

doi:10.1111/1365-2435.12937

52. Sato Y, Shimizu-Inatsugi R, Yamazaki M, Shimizu KK, Nagano AJ. (2019). Plant

trichomes and a single gene *GLABRA1* contribute to insect community composition on

field-grown *Arabidopsis thaliana*. BMC Plant Biol 19:163. doi:10.1186/s12870-019-

1705-2

53. Schuman MC, Allmann S, Baldwin IT. (2015). Plant defense phenotypes determine the

consequences of volatile emission for individuals and neighbors. eLife 4: e04490

54. Stotz HU, Thomson J, Wang Y. (2009). Plant defensins: defense, development and

application. Plant Signal Behav 4: 1010–1012.

55. Tahvanainen JO, Root RB. (1972). The influence of vegetational diversity on the

population ecology of a specialized herbivore, *Phyllotreta cruciferae* (Coleoptera:

Chrysomelidae). Oecologia 10: 321–346.

56. Togninalli M, Seren Ü, Meng D, Fitz J, Nordborg M, Weigel D. et al. (2018). The

AraGWAS Catalog: a curated and standardized *Arabidopsis thaliana* GWAS catalog.

Nucleic Acids Res 46: D1150–D1156.

57. Underwood N, Inouye BD, Hambäck PA. (2014). A conceptual framework for

associational effects: when do neighbors matter and how would we know? Q Rev Biol

89: 1–19.

58. Verschut TA, Becher PG, Anderson P, Hambäck PA. (2016). Disentangling associational

effects: both resource density and resource frequency affect search behaviour in complex

environments. Funct Ecol 30: 1826-1833.

59. Wang M, Roux F, Bartoli C, Huard-Chauveau C, Meyer C, Lee H. et al. (2018). Two-

way mixed-effects methods for joint association analysis using both host and pathogen

genomes. Proc Natl Acad Sci USA 115: E5440–E5449. doi:10.1073/pnas.1710980115

60. Weiner J. (1990). Asymmetric competition in plant populations. Trends Ecol Evol 5: 360-

364.

61. Wuest SE, Niklaus PA. (2018). A plant biodiversity effect resolved to a single

chromosomal region. Nat Ecol Evol 2: 1933–1939.

62. Yamamoto E, Matsunaga H, Onogi A, Ohyama A, Miyatake K, Yamaguchi H. et al.

(2017). Efficiency of genomic selection for breeding population design and phenotype

prediction in tomato. Heredity 118: 202–209.

63. Zeller SL, Kalinina O, Flynn DFB, Schmid B. (2012). Mixtures of genetically modified

wheat lines outperform monocultures. Ecol Appl 22: 1817–1826.

TABLES & FIGURES

Table 1. Factors affecting the power to detect signals in simulated phenotypes. The response variable was the maximum Area Under the ROC Curve (AUC) of the spatial scales from $s = 1$ to $s = 3$. ANOVA tables show the degree of freedom (df), sum of squares (SS), F-statistics, and p-values.

(a) AUC_{self}, No. of causal SNPs = 20

Factors	df	SS	F	p-value
$\sigma_S^2 : \sigma_N^2 : \sigma_{SxN}^2$	2	0.0110	1.91	0.149
α	1	0.0003	0.10	0.750
PVE_β	1	0.410	142.7	< 2.2e-16
$PVE_\beta + PVE_u$	1	0.0196	6.81	0.00933
Residuals	534	1.53		

(c) AUC_{self}, No. of causal SNPs = 200

Factors	df	SS	F	p-value
$\sigma_S^2 : \sigma_N^2 : \sigma_{SxN}^2$	2	0.0000	0.01	0.9909
α	1	0.0044	6.05	0.0142
PVE_β	1	0.798	1106	<2.2e-16
$PVE_\beta + PVE_u$	1	0.0115	16.0	7.29E-05
Residuals	534	0.385		

(b) AUC_{nei}, No. of causal SNPs = 20

Factors	df	SS	F	p-value
$\sigma_S^2 : \sigma_N^2 : \sigma_{SxN}^2$	2	0.013	1.4	0.259
α	1	1.006	205	< 2.2e-16
PVE_β	1	1.101	225	< 2.2e-16
$PVE_\beta + PVE_u$	1	0.060	12.2	0.000520
Residuals	174	2.62		

(d) AUC_{nei}, No. of causal SNPs = 200

Factors	df	SS	F	p-value
$\sigma_S^2 : \sigma_N^2 : \sigma_{SxN}^2$	2	0.0011	0.81	0.445
α	1	0.195	295	< 2.2e-16
PVE_β	1	0.248	375	< 2.2e-16
$PVE_\beta + PVE_u$	1	0.0050	7.60	0.00605
Residuals	534	0.352		

p. 30

679 **Table 2.** GO enrichment analysis of the leaf damage score with Fisher's exact probability

680 tests at FDR < 0.1. Candidate genes within ~10 kb around SNPs with the top 0.1%

681 association score -$\log_{10}$(p-values) were subject to the GO analysis.

(a) Self, β_1

GO	FDR	Description
GO:0043531	0.0071	ADP binding
GO:0009267	0.0083	cellular response to starvation
GO:0031669	0.0083	cellular response to nutrient levels
GO:0050662	0.0127	coenzyme binding
GO:0031668	0.0546	cellular response to extracellular stimulus
GO:0042594	0.0546	response to starvation
GO:0071496	0.0673	cellular response to external stimulus
GO:0009605	0.0829	response to external stimulus
GO:0004553	0.0829	hydrolase activity, hydrolyzing O-glycosyl compounds
GO:0016798	0.0829	hydrolase activity, acting on glycosyl bonds
GO:0031667	0.0969	response to nutrient levels
GO:0005618	0.0969	cell wall
GO:0030312	0.0969	external encapsulating structure
GO:0000166	0.0982	nucleotide binding
GO:1901265	0.0982	nucleoside phosphate binding

(b) Neighbor, β_2

GO	FDR	Description
GO:0004857	0.0271	enzyme inhibitor activity
GO:0031640	0.0271	killing of cells of other organism
GO:0001906	0.0271	cell killing
GO:0044364	0.0271	disruption of cells of other organism
GO:0043531	0.0271	ADP binding
GO:0035821	0.0307	modification of morphology or physiology of other organism
GO:0010393	0.0307	galacturonan metabolic process
GO:0045488	0.0307	pectin metabolic process
GO:0042545	0.0341	cell wall modification
GO:0044419	0.0595	interspecies interaction between organisms

p. 31

Figure 1. Relationship between Neighbor GWAS and Ising model. Upper panels show spatial

arrangements expected by a 2-D Ising model $\sum y_i = \beta_1 \sum x_i + \beta_2 \sum_{<i,j>} x_i x_j$. (a) If $\beta_2 > 0$, mixed

patterns give the argument of the minimum for a population sum of phenotype values Σy_i. (b)

If $\beta_2 < 0$, clustered patterns give the argument of the minimum for Σy_i. (c) In addition, β_1

determines overall patterns favoring -1 or +1 states. Shown are outcomes from a random 100

$\times$ 100 lattice after 1000 iterations of Gibbs sampling. Conversely, the neighbor GWAS was

implemented as an inverse problem of the 2-D Ising model, where genotypes and its spatial

arrangement, x_i and $x_i x_j$, are given while the coefficients β_1 and β_2 are to be estimated from

observed phenotypes y_i. In addition, the variance component due to self and neighbor effects

was considered a random effect in a linear mixed model, such that $u_i \sim \text{Norm}(0, \sigma_S^2 K_S + \sigma_N^2 K_N)$.

Once β_1 and β_2 are determined, we could simulate a genotype distribution that maximizes or

minimizes Σy_i.

Figure 2. Experimental setting in the *Arabidopsis* herbivory data. (a) Photograph of the field

site. Each 0.6×2.6 m block included a replicate of 200 accessions, where 5×40 plants were

assigned to a row and column, respectively. (b) *Arabidopsis thaliana* plants were arranged in

a checkered manner. Yellow lines represent s-th neighbor scales from a focal i-th plant. (c) A

graphical explanation of the experimental area. A meadow (green) was separately covered

with weed-masking sheets (grey).

Figure 3. Scale dependence of neighbor effects on simulated phenotypes among all iterations

with the number of causal SNPs = 20. The broad (a), intermediate (b), and narrow (c)

effective range of neighbor effects are represented by weak, moderate, and strong distance

decay, respectively. The proportion of phenotypic variation explained by neighbor effects

(PVE_{nei}) and the area under the ROC curve (AUC) of neighbor effects are shown along the

spatial scale from the first nearest ($s = 1$) to the third nearest ($s = 3$) neighbors. An AUC at 1.0

indicates a perfect detection of signals. Boxplots show center line: median, box limits: upper

and lower quartiles, whiskers: 1.5× interquartile range, and points: outliers. The case for the

number of causal SNPs = 200 is given in Figure S1.

p. 34

Figure 4. Neighbor GWAS of the leaf damage score on field-grown *Arabidopsis thaliana*. (a)

The proportion of leaf damage variation explained by self-genotype effects PVE_{self} (=

$\sigma_S^2/(\sigma_S^2+\sigma_N^2+\sigma_e^2)$: blue fraction), neighbor effects PVE_{nei} (= $\sigma_N^2/(\sigma_S^2+\sigma_N^2+\sigma_e^2)$: red fraction), and

residuals at the spatial scale of $s = 1$ and $s = 2$. Asterisks highlight a significant fraction with

likelihood ratio tests: ***p*-value < 0.01. (b, c) Manhattan plots for the self and neighbor effects

on the leaf damage score. Different colors highlight the first to fifth chromosomes of *A.*

thaliana. Lighter plots indicate smaller MAF. Results of neighbor effects are shown at $s = 1$.

(a) Chr 2, Position 14679190: $\beta_1 = 0.15$, $\beta_2 = 0.26$, $\Sigma y_i = -176$

(b) Chr 2, Position 9422409: $\beta_1 = -0.003$, $\beta_2 = -0.18$, $\Sigma y_i = -290$

(c) Chr 5, Position 19121831: $\beta_1 = 0.13$, $\beta_2 = -0.24$, $\Sigma y_i = -650$

(d) No effects: $\beta_1 = 10^{-6}$, $\beta_2 = 10^{-6}$, $\Sigma y_i = 10^{-4}$

729 **Figure 5.** Post hoc simulations exemplifying a spatial arrangement of two alleles expected by

730 the estimated self and neighbor effects, β_1 and β_2, on the leaf damage score of *Arabidopsis*

731 *thaliana*. Population sum of the leaf damage $\sum y_i = \beta_1 \sum x_i + \beta_2 \sum_{<i,j>} x_i x_j$ was minimized

732 using 1000 iterations of Gibbs sampling from a random distribution of two alleles in a 10 ×

733 40 space.