Neighbor GWAS: incorporating neighbor genotypic identity into genome-

wide association studies of field herbivory on *Arabidopsis thaliana*

Yasuhiro Sato[1,2], Eiji Yamamoto[1,3], Kentaro K. Shimizu[4,5*], Atsushi J. Nagano[6*]

[1]PRESTO, Japan Science and Technology Agency, Kawaguchi 332-0012, Japan

[2]Research Institute for Food and Agriculture, Ryukoku University, Yokotani 1-5, Seta Oe-

cho, Otsu, Shiga 520-2194, Japan

[3]Kazusa DNA Research Institute, Kazusa-kamatari 2-6-7, Kisarazu, Chiba 292-0818, Japan

[4]Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Winterthurerstrasse 190, 8057 Zurich, Switzerland

[5]Kihara Institute for Biological Research, Yokohama City University, 641-12 Maioka, 244-

0813 Totsuka-ward, Yokohama, Japan

[6]Department of Plant Life Sciences, Faculty of Agriculture, Ryukoku University, Yokotani 1-

5, Seta Oe-cho, Otsu, Shiga 520-2194, Japan

*Co-correspondence:

Atsushi J. Nagano

anagano@agr.ryukoku.ac.jp

Kentaro K. Shimizu

kentaro.shimizu@ieu.uzh.ch

Running head: Neighbor GWAS of herbivory

Word count for main text (excluding refences, tables and figures): 6589

p. 1

ABSTRACT

An increasing number of field studies show that the phenotype of an individual plant depends not only on its genotype but also on neighboring plants; however, this fact is not taken into consideration in genome-wide association studies (GWAS). Based on the Ising model of ferromagnetism, we incorporated neighbor genotypic identity into a regression model. The proposed method, named "neighbor GWAS," was applied to simulated and real phenotypes using *Arabidopsis thaliana* accessions. Our simulations showed that phenotypic variation explained by neighbor effects approached a plateau when an effective spatial scale became narrow. Thus, the effective scale of neighbor effects could be estimated by patterns of the phenotypic variation explained. The power to detect causal variants of neighbor effects was moderate to strong when a trait was governed by tens of variants. In contrast, the power decreased when hundreds of variants underlay a single trait. We applied the neighbor GWAS to field herbivory data on 199 accessions of *A. thaliana*, and found a significant contribution of the neighbor effects to the observed damage variation. Interestingly, a locus responsive to methyl jasmonate (AT2G34810 called *BBE16*) was located near the second top-scoring single nucleotide polymorphism (SNP) of the neighbor effects on the herbivory, while self-genotype effects had the third top-scoring SNP near a locus encoding flavin-monooxygenase glucosinolate S-oxygenase 2 (FMO GS-OX2). Overall, the neighbor GWAS highlights the overlooked role of plant neighborhood effects in shaping phenotypic variation, thereby providing a novel and powerful tool to analyze complex traits in spatially structured environments.

p. 2

INTRODUCTION

Plants are immobile and thus cannot escape their neighbors. In natural and agricultural

systems, individual phenotypes depend not only on the plants' own genotype but also on

other plants in a neighborhood (Tahvanainen and Root 1972; Barbosa et al. 2009; Underwood

et al. 2014). This phenomenon has been termed neighbor effects or associational effects in

plant ecology (Barbosa et al. 2009; Underwood et al. 2014; Sato 2018). Such neighbor effects

were initially reported as a form of interspecific interaction among different plant species

(Tahvanainen and Root 1972), but many studies have illustrated that neighbor effects occur

among different genotypes within a plant species with respect to: (i) herbivory (Schuman et

al. 2015; Sato 2018; Ida et al. 2018), (ii) pathogen infections (Mundt 2002; Zeller et al.

2012), and (iii) pollinator visitations (Underwood et al. 2020). Although neighbor effects are

of considerable interest in plant science (Dicke and Baldwin 2010; Erb 2018) and agriculture

(Zeller et al. 2012; Dahlin et al. 2018), these effects are not often considered in quantitative

genetics of field-grown plants.

Complex mechanisms underlie neighbor effects through direct competition (Weiner

1990), herbivore and pollinator movement (Bergvall et al. 2006; Verschut et al. 2016;

Underwood et al. 2020), and volatile communication among plants (Schuman et al. 2015;

Dahlin et al. 2018). For example, lipoxygenase (*LOX*) genes govern jasmonate-mediated

volatile emissions that induce defenses of neighboring plants in a wild tobacco (*Nicotiana*

attenuata) (Schuman et al. 2015). Even if direct plant–plant communications are absent,

herbivores can mediate indirect interactions between plant genotypes (Sato and Kudoh 2017;

Ida et al. 2018): *GLABRA1* (*GL1*) polymorphism determines hairy or glabrous phenotypes in

Arabidopsis plants (Hauser et al. 2001) and allow a flightless leaf beetle (*Phaedon brassicae*)

to avoid hairy plants when adult beetles encountered hairy plants at a low frequency (Sato

p. 3

and Kudoh 2017; Sato et al. 2017). Yet, there are few hypothesis-free approaches to seek key

genetic variants responsible for plant neighborhood effects.

Genome-wide association studies (GWAS) have been increasingly adopted to resolve

the genetic architecture of complex traits in the model plant, *Arabidopsis thaliana* (Atwell et

al. 2010; Togninalli 2018), and crop species (Hamblin et al. 2011). The interactions of plants

with herbivores (Brachi et al. 2015; Nallu et al. 2018), microbes (Horton et al. 2014; Wang et

al. 2018), and other plant species (Frachon et al. 2019), are examples of such complex traits

investigated through the lens of GWAS. To distinguish causal variants from the genome

structure, GWAS often employs a linear mixed model with kinship considered as a random

effect (Kang et al. 2008; Korte and Farlow 2013). However, because of combinatorial

explosion, it is generally impossible to test the full set of inter-genomic locus-by-locus

interactions (Gondro et al. 2013); thus, some feasible and reasonable approach should be

developed for GWAS of neighbor effects.

To incorporate neighbor effects into GWAS, we focused on a theoretical model of

neighbor effects in magnetic fields, known as the Ising model (Ising 1925; McCoy and

Maillard 2012), which has been applied to forest gap dynamics (Kizaki and Katori 1999;

Schlicht and Iwasa 2004) and community assembly (Azaele et al. 2010) in plant ecology.

Using the Ising analogy, we compare individual plants to a magnet: the two alleles at each

locus correspond to the north and south dipoles, and genome-wide multiple testing across all

loci is analogous to a number of parallel two-dimensional layers. The Ising model has a clear

advantage in its interpretability, such that: (i) the optimization problem for a population sum

of trait values can be regarded as an inverse problem of a simple linear model, (ii) the sign of

neighbor effects determines a model trend to generate a clustered or checkered spatial pattern

of the two states, and (iii) the self-genotypic effect determines the general tendency to favor

p. 4

one allele over another (Fig. 1).

In this study, we proposed a new methodology integrating GWAS and the Ising

model, named "neighbor GWAS." The method was applied to simulated phenotypes and real

data of field herbivory on *A. thaliana*. We addressed two specific questions: (i) what spatial

and genetic factors influenced the power to detect causal variants? and (ii) were neighbor

effects significant sources of leaf damage variation in field-grown *A. thaliana*? Based on the

simulation and application, we determined the feasibility of our approach to detect neighbor

effects in field-grown plants.

MATERIALS & METHODS

Neighbor GWAS

Basic model. We analyzed neighbor effects in GWAS as an inverse problem of the two-

dimensional Ising model (Fig. 1). We considered a situation where a plant accession has

either of two alleles at each locus, and a number of accessions occupied a finite set of field

sites in a two-dimensional lattice. The allelic status at each locus was represented by x, and

so the allelic status at each locus of the *i*-th focal plant and the *j*-th neighboring plants was

designated as $x_{i(j)} \in \{-1, +1\}$. Based on a two-dimensional Ising model, we defined a

phenotype value of *i*-th focal individual plant y_i as:

$$y_i = \beta_1 x_i + \beta_2 \sum_{<i,j>} x_i x_j \qquad (1)$$

where β_1 and β_2 denoted self-genotype and neighbor effects, respectively. If two neighboring

plants shared the same allele at a given locus, the product $x_i x_j$ turned into (-1)×(-1)=+1 or

(+1)×(+1)=+1. If two neighbors had different alleles, the product $x_i x_j$ became (-1)×(+1)=-1

or (+1)×(-1)=-1. Accordingly, the effects of neighbor genotypic identity on a phenotype

p. 5

depended on the coefficient β_2 and the number of two alleles in a neighborhood. If the

numbers of identical and different alleles were the same near a focal plant, these neighbors

offset the sum of the products between the focal plant i and all j neighbors $\sum_{<i,j>} x_i x_j$ and

exerted no effects on a phenotype. When we summed up the phenotype values for the total

number of plants n and replaced it as $E = -\beta_2$, $H = -\beta_1$ and $\epsilon_I = \sum y_i$, eq. 1 could be

transformed as $\epsilon_I = -E \sum_{<i,j>} x_i x_j - H \sum x_i$, which defined the interaction energy of a two-

dimensional ferromagnetic Ising model (McCoy and Maillard 2012). The neighbor

effect β_2 and self-genotype effect β_1 were interpreted as the energy coefficient E and external

magnetic effects H, respectively. An individual plant represented a spin and the two allelic

states of each locus corresponded to a north or south dipole. The positive or negative value

of $\sum x_i x_j$ indicated a ferromagnetism or paramagnetism, respectively. In this study, we did not

consider the effects of allele dominance because this model was applied to inbred *A. thaliana*.

However, heterozygotes could be processed if the neighbor covariate $x_i x_j$ was weighted by

an estimated degree of dominance in the self-genotypic effects on a phenotype.

Mixed model. For association mapping, we needed to determine β_1 and β_2 from

observed phenotypes and considered a confounding sample structure as advocated by

previous GWAS (e.g., Kang et al. 2008; Korte and Farlow 2013). Extending the basic model

eq. 1, we described a linear mixed model at an individual level as:

$$y_i = \beta_0 + \beta_1 x_i + \frac{\beta_2}{L} \sum_{<i,j>}^s x_i x_{j,s} + u_i + e_i \qquad (2)$$

140   where β_0 indicated the intercept, and the term $\beta_1 x_i$ represented fixed self-genotype effects as

141   tested in conventional GWAS; β_2 was the coefficient of fixed neighbor effects. The neighbor

142   covariate $\sum_{<i,j>}^s x_i x_{j,s}$ indicated a sum of products for all combinations between the i-th focal

143   plant and the j-th neighbor at the s-th spatial scale from the focal plant i, and was scaled by

p. 6

144   the number of neighboring plants, L. Variance components due to a sample structure in self

145   and neighbor effects was modeled by a random effect $u_i \sim \text{Norm}(0, \sigma_1^2 K_1 + \sigma_2^2 K_2)$. The

146   residual was expressed as $e_i \sim \text{Norm}(0, \sigma_e^2)$.

147   The $n \times n$ variance-covariance matrices represented the similarity in self-genotypes

148   (i.e., kinship) and neighbor covariates among n individual plants as:

149                          $$K_1 = \frac{1}{q-1} X_1^T X_1 \quad \text{and}$$

150                          $$K_2 = \frac{1}{q-1} X_2^T X_2$$

151   where q indicated the number of markers. As we defined $x_{i(j)} \in \{+1, -1\}$, the elements of the

152   kinship matrix K_1 were scaled to represent the proportion of marker loci shared among $n \times n$

153   plants such that $K_1 = \left(\frac{k_{ij}+1}{2}\right)$ and $k_{ij} = [0, 1]$; σ_1^2 and σ_2^2 indicated variance component

154   parameters for the self and neighbor effects.

155   The elements of n plants $\times$ q markers matrix X_1 and X_2 consisted of explanatory

156   variables for the self and neighbor effects as $X_1 = (x_i)$ and $X_2 = \left(\frac{\sum_{<i,j>}^{s} x_i x_{j,s}}{L}\right)$. The

157   individual-level formula eq. 2 could also be converted into a conventional matrix form as:

158                          $$y = X\beta + Zu + e$$

where y was $n \times 1$ vector of phenotypes; X was a matrix of fixed effects, including mean,

self-genotype x_i, neighbor covariate $\frac{\sum_{<i,j>}^{s} x_i x_{j,s}}{L}$, and other confounding covariates for n plants;

β was a vector that represents coefficients of the fixed effects; Z was a design matrix

allocating individuals to a genotype, and became an identity matrix if all plants were different

accessions; u was the random effect with $\text{Var}(u) = \sigma_1^2 K_1 + \sigma_2^2 K_2$; and e was residual as

$\text{Var}(e) = \sigma_e^2 I$. In such a mixed model, the proportion of phenotypic variation explained (PVE)

by the self and neighbor effects could be calculated as $\text{PVE}_{\text{self}} = \sigma_1^2 / (\sigma_1^2 + \sigma_2^2 + \sigma_e^2)$ and

$\mathrm{PVE_{nei}} = \sigma_2^2 / (\sigma_1^2 + \sigma_2^2 + \sigma_e^2)$, respectively. The line of extensions to incorporate neighbor

effects into GWAS was hereafter referred to as "neighbor GWAS." Note also that the standard

GWAS was a subset of the neighbor GWAS. These two models became equivalent at $s = 0$

and $\sigma_2^2 = 0$. When testing the self-genotype effect β_1, we recommended that neighbor effects

and its variance component σ_2^2 should be excluded; otherwise, the standard GWAS failed to

correct a sample structure because of the additional variance component at $\sigma_2^2 \neq 0$.

Power simulation

To examine the power to detect neighbor effects, we applied the neighbor GWAS to

simulated phenotypes. Phenotypes were simulated using a part of the real genotypes of *A.*

thaliana. To evaluate the power of the simple linear model, we assumed a complex ecological

form of neighbor effects with multiple variance components controlled. The power was

evaluated by the receiver operating characteristic (ROC) curve on the association score of -

$\log_{10}(p$-value) (e.g., Gage et al. 2018). All analyses were performed using R version 3.6.0 (R

Core Team 2019).

181   ***Genotype data.*** To consider a realistic genetic structure in the simulation, we used

part of the *A. thaliana* RegMap panel (Horton et al. 2012). The genotype data on 1307

accessions were downloaded from the Joy Bergelson laboratory website

(http://bergelson.uchicago.edu/?page_id=790 accessed on February 9, 2017). We extracted

the chromosome 1 and 2 data with the minor allele frequency (MAF) at >0.1, providing a

matrix of 1307 plants with 65 226 single nucleotide polymorphisms (SNPs). Pairwise linkage

disequilibrium (LD) among the loci was $r^2 = 0.003$ [0.00-0.06: median with upper and lower

95 percentiles]. Before generating a phenotype, each locus was centered by its mean and

scaled by its standard deviation. Subsequently, we randomly selected 1296 accessions (= 36 ×

p. 8

36 accessions) without any replacements for each iteration, and placed them in a 36 × 72

checkered space following *Arabidopsis* experimental settings (see Fig. 2b).

Phenotype simulation. To address ecological issues specific to plant neighborhood

effects, we considered two extensions, namely asymmetric neighbor effects and spatial

scales. Firstly, previous studies have shown that such plant–plant interactions are sometimes

asymmetric between two accessions in herbivory (e.g., Bergvall et al. 2006; Verschut et al.

2016; Sato and Kudoh 2017) and height competition (Weiner 1990). Such asymmetric

neighbor effects can be tested by statistical interaction terms in a linear model (Bergvall et al.

2006; Sato and Kudoh 2017). Secondly, several studies have shown that the strength of

neighbor effects depends on spatial scales (Hambäck et al. 2014), and that the scale of

neighbors to be analyzed relies on the dispersal ability of causative organisms (see Hambäck

et al. 2009; Sato and Kudoh 2015; Verschut et al. 2016; Ida et al. 2018 for insect and mammal

herbivores; Rieux et al. 2014 for pathogen dispersal) or the size of competing plants (Weiner

1990). We assumed the distance decay at *s*-th sites from a focal individual *i* with the decay

coefficient α as $w(s, \alpha) = \mathrm{e}^{-\alpha(s-1)}$, since such an exponential distance decay has been

widely adopted in empirical studies (Devaux et al. 2007; Carrasco et al. 2010; Rieux et al.

2014; Ida et al. 2018). Therefore, we assumed a more complex model for simulated

phenotypes than the model for neighbor GWAS as follows:

$$y_i = \beta_0 + \beta_1 x_i + \frac{\beta_2}{L}\sum_{<i,j>}^{s} w(s, \alpha)x_i x_{j,s} + \beta_{12}\frac{x_i}{L}\sum_{<i,j>}^{s} w(s, \alpha)x_i x_{j,s} + u_i + e_i \quad (3)$$

where β_{12} was the coefficient for asymmetry in neighbor effects. Total variance components

resulting from three background effects (i.e., the self, neighbor, and self-by-neighbor effects)

was defined as $u_i \sim \mathrm{Norm}(0, \sigma_1^2 \mathbf{K}_1 + \sigma_2^2 \mathbf{K}_2 + \sigma_{12}^2 \mathbf{K}_{12})$. The three variance component

parameters σ_1^2, σ_2^2, and σ_{12}^2, determined the relative importance of self-genotype, neighbor,

and asymmetric neighbor effects in u_i. Given the elements of the *n* plants × *q* marker

p. 9

explanatory matrix with $X_{12} = (\frac{x_i}{L}\sum_{<i,j>}^{s} w(s,\alpha)x_i x_{j,s})$, the similarity in asymmetric

neighbor effects was calculated as $K_{12} = \frac{1}{q-1}X_{12}^T X_{12}$. To control phenotypic variations, we

further partitioned the proportion of phenotypic variation into those explained by major-effect

genes and variance components $PVE_\beta + PVE_u$, major-effect genes alone PVE_β, and residual

error PVE_e, where $PVE_\beta + PVE_u + PVE_e = 1$. The *optimize* function in R was used to adjust

simulated phenotypes to the given amounts of PVE.

Parameter setting. Ten phenotypes were simulated for each combination of the

distance decay α, the proportion of phenotypic variance explained by major-effect genes

PVE_β, variance components PVE_u, and the relative importance of multiple variance

components $\sigma_1^2 : \sigma_2^2 : \sigma_{12}^2$ as: $\alpha = 0.25$, 1.0 or 3.0, $\sigma_1^2 : \sigma_2^2 : \sigma_{12}^2 = 6:3:1$, 4:4:1, or 3:6:1, PVE_β and

$PVE_\beta + PVE_u = 0.1$ and 0.4, 0.3 and 0.4, 0.3 and 0.8, or 0.6 and 0.8. The maximum reference

scale was fixed at $s = 3$. The line of simulations was repeated for the number of causal SNPs

at 10, 50, or 300 to examine cases of an oligogenic and polygenic control for a trait. The non-

zero coefficients for the causal SNPs were randomly sampled from a uniform distribution,

Unif(|0.5|, |2.0|), and assigned as some causal SNPs were responsible for both the self and

neighbor effects. Of the total number of causal SNPs, 15% had all self, neighbor, and

asymmetric neighbor effects (i.e., $\beta_1 \neq 0$ and $\beta_2 \neq 0$ and $\beta_{12} \neq 0$); another 15% had both the

self and neighbor effects, but no asymmetry in the neighbor effects ($\beta_1 \neq 0$ and $\beta_2 \neq$

0 and $\beta_{12} = 0$); another 35% had self-genotypic effects only ($\beta_1 \neq 0$); and the remaining

35% had neighbor effects alone ($\beta_2 \neq 0$). Given its biological significance, we assumed that

some loci having neighbor signals possessed asymmetric interactions between neighbors

($\beta_2 \neq 0$ and $\beta_{12} \neq 0$) while the others had symmetric interactions ($\beta_2 \neq 0$ and $\beta_{12} = 0$).

Therefore, the number of causal SNPs in β_{12} was smaller than that in the main neighbor

p. 10

effects β_2. According to this assumption, the variance component σ_{12}^2 was also assumed to be

smaller than σ_2^2.

Summary statistics. The simulated phenotypes were fitted by eq. 2 to test the

significance of coefficients β_1 and β_2, and to estimate the variance component due to self and

neighbor effects PVE_{self} and PVE_{nei}. The stepwise likelihood ratio tests were performed first

between the null model and model with a self-genotype effect alone, and then between the

self-genotype effect model and model with both self and neighbor effects. The likelihood

ratio was calculated as the difference in deviance, i.e., -2 × log-likelihood, which is

asymptotically χ^2 distributed with one degree of freedom. The variance

components, σ_1^2 and σ_2^2, were estimated using a linear mixed model without any fixed effects.

To solve the mixed model with the two random effects, we used the average information

restricted maximum likelihood (AI-REML) algorithm implemented in the *lmm.aireml*

function in the *gaston* package of R (Perdry and Dandine-Roulland 2018). Subsequently, we

replaced the two variance parameters σ_1^2 and σ_2^2 in eq. 2 into their estimates $\hat{\sigma}_1^2$ and $\hat{\sigma}_2^2$ by the

AI-REML, and performed association tests by solving a linear mixed model with a fast

approximation by an eigen value decomposition (implemented in the *lmm.diago* function:

Perdry and Dandine-Roulland 2018). We computed the stepwise likelihood using the

lmm.diago.profile.likelihood function to first test β_1 in the standard GWAS, and then to

test β_2 in the neighbor GWAS. True or false positive rates were evaluated by ROC curves and

area under the ROC curves (AUC) (Gage et al. 2018). An AUC of 0.5 would indicate that

GWAS has no power to detect true signals, while an AUC of 1.0 would indicate that all the

top signals predicted by GWAS agree with true signals. The roc function in the pROC

package (Robin et al. 2011) was used to calculate AUC from $-\log_{10}(p\text{-value})$. Factors

affecting the AUCs were tested by analysis-of-variance (ANOVA). The AUCs had

proportional values, and thus were arcsine-square-root transformed to improve the Shapiro–

Wilk normality statistic before ANOVA. The AUC and variance components were calculated

from $s = 1$ (the first nearest neighbors) to $s = 3$ (up to the third nearest neighbors) cases. The

AUCs were also calculated using standard linear models without any random effects to

examine whether the linear mixed models were superior to the linear models. We also tested

the neighbor GWAS model incorporating neighbor phenotype $y_{j,s}$ instead of $x_{j,s}$.

***Arabidopsis* herbivory data**

We applied the neighbor GWAS to field data of *Arabidopsis* herbivory. The procedure of the

present field experiment followed our previous experiment on summer herbivory on field-

grown *A. thaliana* (Sato et al. 2019). We selected 199 worldwide accessions from 2029

accessions sequenced by the RegMap (Horton et al. 2012) and 1001 Genomes project

(Alonso-Blanco et al. 2016). Of the 199 accessions, most were overlapped with a previous

GWAS of biotic interactions (Horton et al. 2014) and half were included by a GWAS of

glucosinolates (Chan et al. 2010). Eight replicates of each of the 199 accessions were first

prepared in a laboratory and then transferred to the outdoor garden at the Center for

Ecological Research, Kyoto University, Japan (Otsu, Japan: 35°06′N, 134°56′E, alt. ca. 200

m: Fig. 2a). Seeds were sown on Jiffy-seven pots (33-mm diameter), and stratified at a

temperature of 4 °C for a week. Seedlings were cultivated for 1.5 months under a short-day

condition (8 h light: 16 h dark, 20 °C). Plants were then separately potted in plastic pots (6

cm in diameter) filled with mixed soil of agricultural compost (Metro-mix 350, SunGro Co.,

USA) and perlite at a 3:1 ratio. Potted plants were set in plastic trays (10 × 40 cells) in a

checkered pattern (Fig. 2b). In the field setting, a set of 199 accessions and an additional Col-

0 accession were randomly assigned to each block without replacement (Fig. 2b). Eight

p. 12

replicates of these blocks were set >2 m apart from each other (Fig. 2c). Potted plants were

exposed to the field environment for 3 wk in June 2017. At the end of experiment, we scored

the percentage of foliage eaten as: 0 for no visible damage, 1 for ≤10%, 2 for >10% and ≤

25%, 3 for > 25% and ≤ 50%, 4 for >50% and ≤ 75%, and 5 for >75% of the foliage area

eaten. All plants were scored by a single person to avoid observer bias. The most

predominant herbivore in this field trial was the diamond back moth (*Plutella xylostella*),

followed by the small white butterfly (*Pieris rapae*). We also recorded the initial plant size

and the presence of inflorescence to incorporate them as covariates. Initial plant size was

evaluated by the length of the largest rosette leaf (mm) at the beginning of the field

experiment and the presence of inflorescence was recorded 2 wk after transplanting.

We estimated the variance components and performed the association tests for the

leaf damage score with the neighbor covariate at $s = 1$ and 2. These two scales corresponded

to $L = 4$ (the nearest four neighbors) and $L = 12$ (up to the second nearest neighbors),

respectively, in the *Arabidopsis* dataset. The variation partitioning and association tests were

performed using the *gaston* package, as mentioned above. To determine the significance of

variance component parameters, we compared the likelihood between mixed models with one

or two random effects. For the genotype data, we used an imputed SNP matrix of all the 2029

accessions studied by the RegMap (Horton et al. 2012) and 1001 Genomes project (Alonso-

Blanco et al. 2016). Missing genotypes were imputed using BEAGLE (Browning and

Browning 2009), as described by Togninalli et al. (2018) and updated on the AraGWAS

Catalog (https://aragwas.1001genomes.org). Of the 10 709 466 SNPs from the full imputed

matrix, we used 1 242 128 SNPs with MAF at > 0.05 and LD of adjacent SNPs at r^2 <0.8. We

considered the initial plant size, presence of inflorescence, experimental blocks, and the edge

or center within a block as fixed covariates; these factors explained 12.5% of the leaf damage

p. 13

variation (1.2% by initial plant size, Wald test, Z = 3.53, p-value<0.001; 2.4% by the

presence of inflorescence, Z = -5.69, p-value<10^{-8}; 8.3% by the experimental blocks,

likelihood ratio test, χ^2 = 152.8, df = 7, p-value<10^{-28}; 0.5% by the edge or center, Z = 3.11,

p-value = 0.002). After the association mapping, we searched candidate genes within ~10 kb

around target SNPs, based on the Araport11 gene model with the latest annotation of The

Arabidopsis Information Resource (TAIR) (accessed on 7 September 2019). Gene-set

enrichment analysis was performed using the Gowinda algorithm that enables unbiased

analysis of GWAS results (Kofler and Schlotterer 2012). We tested SNPS with the top 0.1% -

$\log_{10}(p$-value) score with the option "--gene-definition undownstream10000," "--min-genes

20," and "--mode snp." The GO.db package (Carlson et al. 2018) and the latest TAIR AGI

code annotation were used to build input files. The R source codes, accession list, and

phenotype data are available at the GitHub repository

(https://github.com/naganolab/NeighborGWAS).

RESULTS

Power simulation

A set of phenotypes were simulated from real genotype data following a complex model (eq.

3), and then fitted by a simplified model (eq. 2). Analyzing the factors affecting AUCs, we

found that the number of causal SNPs, the proportion of phenotypic variation (PVE)

explained by major-effect genes PVE$_{\boldsymbol{\beta}}$, and distance decay of neighbor effects were the most

influential factors affecting the power of the proposed method to detect neighbor signals

(Table 1). In addition to PVE$_{\boldsymbol{\beta}}$, the amount of variance components PVE$_{\boldsymbol{u}}$ also significantly

affected the AUCs of the self and neighbor effects, but these additional effects were less

p. 14

significant compared to those of PVE$_\beta$ alone (Table 1). The power of the standard GWAS to

detect self-genotype effects depended on the number of causal SNPs and PVE$_\beta$, and was not

significantly influenced by the distance decay of neighbor effects (Table 1). Neither the

power to detect self nor neighbor effects were significantly affected by the ratio of three

variance components $\sigma_1^2 : \sigma_2^2 : \sigma_{12}^2$ (Table 1).

The power to detect self-genotype signals changed from strong (AUC>0.9) to weak

(AUC<0.6) as PVE$_\beta$ decreased and the number of causal SNPs increased (Fig. 3a; Fig. S1a).

Compared to the self-genotype effects, it was relatively difficult to detect neighbor effects

(Fig. 3b; Fig. S1b). In the case of the number of causal SNPs = 10, the power to detect

neighbor signals decreased from high (AUC>0.9) to moderate (AUC>0.7) with the

decreasing PVE$_\beta$ and distance decay (Fig. 3b; Fig. S1b). The power to detect neighbor effects

was very weak (AUC> 0.5 and AUC<0.6) in the case of the number of causal SNPs = 300

(Fig. 3b; Fig. S1b). The result of simulations indicated that neighbor effects were detectable

when a target trait was governed by several major genes and the range of neighbor effects

was spatially limited. Additionally, linear mixed models outperformed standard linear models

as shown by 9.3% and 1.7% increases in the power to detect self and neighbor signals,

respectively (AUC$_{self}$, paired t-test, mean of the difference = 0.093, p-value<10^{-16}; AUC$_{nei}$,

mean of the difference = 0.017, p-value<10^{-16}). When the neighbor phenotype $y_{j,s}$ was

incorporated instead of the genotype $x_{j,s}$, the power to detect neighbor effects was very weak,

such that AUC decreased to near 0.5.

Notably, there was a clear relationship between the distance decay α and the

proportion of phenotypic variation explained by neighbor effects PVE$_{nei}$ or AUCs at different

spatial scales (Fig. 4). If the distance decay was weak and the effective range of neighbor

effects was broad, PVE$_{nei}$ and AUCs increased linearly as the reference spatial scale was

p. 15

broadened (Fig. 4a). On the other hand, if the distance decay was strong and the effective

scale of neighbor effects was narrow, PVE_{nei} saturated at the scale of the first nearest

neighbors (Fig. 4c) and AUCs did not increase (Fig. 4b, c). The simulation results indicated

that the effective spatial scales could be estimated by calculating PVE_{nei} across different

spatial scales, and that the first nearest scale was enough to detect neighbor signals unless the

distance decay was very weak.

363   Examination of variance components provided further insights into the PVE

estimates (Fig. S2; Table S2). Comparing the true and estimated PVE, we found that the

estimated PVE by self and neighbor effects was biased as a consequence of the ratio of

variance components $\sigma_1^2 : \sigma_2^2 : \sigma_{12}^2$, the total amount of PVE, and the magnitude of distance

decay (Table S2). The standard heritability namely, PVE_{self} at $s = 0$, was overestimated when

neighbor effects involved a simulated phenotype. The bias of estimated PVE_{self} decreased as

the reference spatial scale increased (Fig. S2). In contrast, PVE_{nei} was underestimated and its

bias decreased as the reference scale increased (Fig. S2). These results indicated that neighbor

effects and its variance component should be tested by comparing the standard GWAS and

neighbor GWAS, because some fraction of PVE was mutually non-exclusive between the two

factors.

***Arabidopsis* herbivory data**

The variation partitioning of leaf damage showed that the PVE by neighbor effects was larger

than PVE by self-genotypic effects ($PVE_{self} = 0.041$, $\chi_1^2 = 0.36$, p-value $= 0.54$; $PVE_{nei} =$

0.214, $\chi_1^2 = 7.23$, p-value $= 0.004$: Fig. 5). Phenotypic variation explained by neighbor effects

on leaf damage did not increase when the neighbor scale was referred up to the nearest and

second nearest individuals ($PVE_{self} = 0.091$, $\chi_1^2 = 1.19$, p-value $= 0.200$; $PVE_{nei} = 0.14$, $\chi_1^2 =$

p. 16

1.41, p-value = 0.166: Fig. 5); therefore, the variation partitioning was stopped at $s = 2$.

Heritability, namely PVE_{self} at $s = 0$, was 0.173 and significant ($\chi_1^2 = 10.1$, p-value = 0.005:

Fig. 5). This range of heritability was overlapped with PVE by neighbor effects alone (PVE_{nei}

without self-effects = 0.248 and 0.243 at $s = 1$ and 2), indicating that some fraction of PVE_{self}

and PVE_{nei} was mutually non-exclusive. We additionally incorporated the neighbor

phenotype $y_{j,s}$ instead of the neighbor genotype $x_{j,s}$ in eq. 2, but PVE_{nei} did not increase

(PVE_{nei} = 0.066 and 0.068 at $s = 1$ and 2, respectively). These results indicated a narrow

effective scale and significant contribution of neighbor effects to the leaf damage score,

where the neighbor genotype rather than the neighbor phenotype was likely a source of

phenotypic variation.

Standard GWAS of the self-genotype effects on the leaf damage detected SNPs with

the second and third largest -$\log_{10}(p$-values) score on the first chromosome (chr1), though

they were not above a threshold of Bonferroni correction (Fig. 6a; Table S3). The second SNP

at chr1-21694386 was located within ~10 kb of three loci encoding a disease resistance

protein (CC-NBS-LRR class) family. The third SNP at chr1-23149476 was located within

~10 kb of the AT1G62540 locus that encoded flavin-monooxygenase glucosinolate S-

oxygenase 2 (FMO GS-OX2), This gene's GO annotations, "glucosinolate biosynthetic

process" and "oxidation-reduction process," were not significantly enriched among genes

within ~10 kb around SNPs with the top 0.1% -$\log_{10}(p$-values) score. In contrast, the

"sesquiterpene metabolic process" was significantly enriched with four unique genes (false

discovery rate = 0.003; Table S4). A QQ-plot did not exhibit an inflation of p-values for the

self-genotype effects (Fig. S3).

Regarding the neighbor effects on the leaf damage, we found non-significant but

weak peaks on the second and third chromosomes (Fig. 6b; Table S3). The second

p. 17

chromosomal region had higher association scores than predicted by the QQ-plot (Fig. S4). A

locus encoding FAD-binding Berberine family protein (AT2G34810 named *BBE16*) was

located within the ~10 kb window near SNPs with the top eleven -log$_{10}$(p-values) score at the

second chromosome, which are known to be up-regulated by methyl jasmonate (Devoto et al.

2005). Three transposable elements and a pseudogene of lysyl-tRNA synthetase 1 were

located near the most significant SNP on the third chromosome. Gene ontology categories

involved in wounding responses were not significantly enriched among genes within ~10 kb

around SNPs with the top 0.1% score of -log$_{10}$(p-values); however, "glucosinolate catabolic

process" was significantly enriched with seven unique genes (false discovery rate = 0.014:

Table S4). We additionally tested asymmetric neighbor effects β_{12} in the real dataset on field

herbivory, but the top 0.1% SNPs of neighbor effects β_2 did not overlap with those of

asymmetric neighbor effects β_{12} (Table S3).

Based on the estimated coefficients $\hat{\beta}_1$ and $\hat{\beta}_2$, we ran a post hoc simulation to infer a

spatial arrangement that minimizes a population sum of the leaf damage $\sum y_i = \beta_1 \sum x_i +$

$\beta_2 \sum_{<i,j>} x_i x_j$. The constant intercept β_0, the variance component u_i, and residual e_i were not

considered because they were not involved in deterministic dynamics of the model. Figure 7

shows three representatives and a neutral expectation. For example, a mixture of a

dimorphism was expected to decrease the total leaf damage for a SNP at chr2-14679190 near

the *BBE16* locus ($\hat{\beta}_2 > 0$: Fig. 7a). On the other hand, a clustered distribution of a

dimorphism was expected to decrease the total damage for a SNP at chr2-9422409 near the

AT2G22170 locus encoding a lipase/lipooxygenase PLAT/LH2 family protein ($\hat{\beta}_1 \approx 0, \hat{\beta}_2 <$

0: Fig. 7b). Furthermore, near monomorphism was expected to decrease the leaf damage for a

SNP at chr5-19121831 near the AT5G47075 and AT5G47077 loci encoding low-molecular

cysteine-rich proteins, LCR20 and LCR6 ($\hat{\beta}_1 > 0, \hat{\beta}_2 < 0$: Fig. 7c). If self and neighbor

p. 18

coefficients had no effects, we observed a random distribution and no mitigation of damage

i.e., $\sum y_i \approx 0$ (Fig. 7d). These post hoc simulations suggested a potential applicability of

neighbor GWAS in optimizing spatial arrangements in field cultivation.

DISCUSSION

Spatial and genetic factors underlying simulated phenotypes

Benchmark tests using simulated phenotypes revealed that appropriate spatial scales could be

estimated by variation partitioning of observed phenotypes. When the scale of neighbor

effects was narrow or moderate ($\alpha = 1.0$ or 3.0), the scale of the first nearest neighbors would

be optimum for the power to detect neighbor signals. In terms of neighbor effects in the

context of plant defense, mobile animals (e.g., mammalian browsers and flying insects) often

select a cluster of plant individuals (e.g., Bergvall et al. 2006; Hambäck et al. 2009; Vershut

et al. 2016); however, previous studies have not detected neighbor effects among individual

plants within a cluster (Hambäck et al. 2014; Sato and Kudoh 2015). The exponential

distance decay of $\alpha = 0.25$ represented situations in which the effective range of neighbor

effects was too broad to be detected; only in such situations should more than the nearest

neighbors be referred to gain the power to detect neighbor effects.

Neighbor GWAS could retain its power as long as neighbor effects were spatially

limited, and several major-effect genes governed a trait. In contrast, when hundreds of causal

variants involved a single trait and less than half of phenotypic variation was attributable to

neighbor effects, we observed a decrease in the power to detect neighbor signals. In GWAS,

false positive rates can be reduced using linear mixed models that deal with kinship structure

as a random effect (Korte and Farlow 2013). Indeed, mixed models were superior to standard

p. 19

linear models in this simulation. Our simulation also adjusted the three variance

components σ_1^2, σ_2^2, and σ_{12}^2, but their relative contribution did not have significant effects on

the power to detect neighbor signals. In addition, the power to detect self-genotype signals

largely depended on the number of causal SNPs and PVE by major-effect genes. This result

suggests that the power for the standard GWAS to detect the self-genotype effects was

independent from neighbor effects, though its heritability may confound PVE by neighbor

effects.

Neighbor GWAS of the field herbivory on *Arabidopsis*

Our genetic analysis of neighbor effects is of ecological interest because how host plant

genotypes shape variation in plant–herbivore interactions is a long-standing question in

population ecology (e.g., Karban 1992; Underwood and Rausher 2000; Utsumi et al. 2011).

Despite the low heritability and several confounding factors under field conditions, the

present study illustrated the significant contribution of neighbor genotypic identity to spatial

variation in herbivory on *A. thaliana*. As we did not find any significant peaks above the

threshold of genome-wide multiple testing, such neighbor effects on herbivory could be

under polygenic control. Glucosinolates and sesquiterpene-mediated defense might jointly

contribute to variation in herbivory among individual plants, as suggested by a gene-set

enrichment analysis.

Regarding the self-genotype effects, we detected *GS-OX2*, a key gene involved in

glucosinolate biosynthesis, near the third top-scoring SNP on the first chromosome. Aliphatic

glucosinolates are well known as a chemical defense against insect herbivory (van Poecke

2007; Brachi et al. 2015), where GS-OX2 catalyzes the conversion from methylthioalkyl to

corresponding methylsulfinylalkyl glucosinolates (Li et al. 2008). Specifically, Geiselhardt et

p. 20

al. (2013) reported that *GS-OX2* was up-regulated in response to feeding by larvae of the

large white butterfly (*Pieris brassicae*) and such an inducible defense was suppressed by

prior oviposition by adult butterflies. The present GWAS supported the relevance of *GS-OX2*

to herbivory by *Pieris* butterflies, because we observed larvae of *P. rapae* largely feeding on

A. thaliana. Furthermore, our experiment was conducted under the field condition where

insect movements and microclimates were not controlled. Thus, the present results provide

ecological insights into the gene function of *GS-OX2* from the laboratory to the field

environment.

Remarkably, the second top-scoring SNP of the neighbor effects was located near the

BBE16 locus responsive to methyl jasmonate. Methyl jasmonate is well known as a volatile

organic chemical that is emitted from damaged tissue and elicits defense response of other

plants (Reymond and Farmer 1998; van Poecke 2007). Volatile-mediated plant–plant

communication is a key driver of neighbor effects in herbivory (e.g., Schuman et al. 2015;

Erb 2018; Dahlin et al 2018), while a molecular function of *BBE16* remains unexplored.

More recent studies have shown that berberine bridge enzyme-like proteins are involved in

damage-associated molecular patterns (Benedetti et al. 2018), which might be a subject of

future studies on inducible plant defense and neighbor effects.

Potential applicability and limitation

GWAS has often been applied to crop plants (Jannink et al. 2010; Hamblin et al. 2011), where

genotypes are known and individuals are evenly transplanted in space. Considering this

outlook for agriculture, we provided an example of neighbor GWAS across a lattice space in

this study. However, wild plant populations sometimes exhibit more complex spatial patterns

than expected by the Ising model (e.g., Kizaki and Katori 1999; Schlicht and Iwasa 2004). A

p. 21

variation partitioning approach could also be applied to uneven distribution if the neighbor

GWAS method will be elaborated to handle distance decay across a complex spatial structure.

Now that a vast number of plant genomes are being sequenced (Exposito-Alonso et al. 2020),

further studies on spatial statistics are encouraged to enable the neighbor GWAS in wild

plants.

We should also note that some fraction of phenotypic variation cannot be separated

into self or neighbor effects, as shown by the shift in PVE from $s = 0$ to $s = 1$ in the simulated

and real data. Such an overlapped fraction might arise because of a strong correlation

between K_1 and K_2 (e.g., $r = 0.60$ and 0.78 at $s = 1$ and 2 in the real data analysis). In eq. 2,

the self-genotypic variable x_i was incorporated into the neighbor variable $\sum x_i x_j / L$, and thus

the kinship matrix K_1 was partially redundant with the similarity matrix of neighbor

effects K_2. For a proper test of neighbor effects β_2 and its variance component σ_2^2, it is

necessary to compare the likelihood ratio between the standard and neighbor GWAS. Once

neighbor effects are detected by the stepwise likelihood ratio test, the heritability and PVE

should be carefully interpreted regarding the potential bias along the reference spatial scale s.

Conclusion

Based on the newly proposed methodology, we suggest that neighbor effects are an

overlooked source of phenotypic variation in field-grown plants. To date, regional-scale

interactions among plants have been analyzed using a genome–environment association study

of plant community composition (Frachon et al. 2019), but fine-scale neighbor effects have

yet to be examined. Recent studies have shown that a single genomic region or a single gene

drives neighbor effects in plant growth, and such a genetic effect shapes a positive

relationship between plant genotype diversity and yield (Wuest and Niklaus 2018; McGale et

p. 22

al. 2020). Our newly proposed methodology of neighbor GWAS could be an efficient tool to

identify such a key genetic variant responsible for neighbor effects and resulting biodiversity

effects.

Neighbor GWAS may also potentially help determine an optimal spatial arrangement

in plant cultivation, as suggested by the post hoc simulation. The Ising model is well

established in statistical physics (McCoy and Maillard 2012) and is now applied to a

machine-learning pipeline that deals with high-dimensional genome data (Fisher and Mehta

2015). Genome-wide polymorphism data are useful not only to seek causal genes in GWAS,

but also to predict breeding values of crop plants in genomic selection (e.g., Jannink et al.

2010; Hamblin et al. 2011; Yamamoto et al. 2017). Given that the neighbor GWAS consists of

a marker-based regression, this methodology could also be implemented as a genomic

selection at a population level. Thus, our study provides an avenue for future studies to

predict population-level phenotypes in spatially structured environments.

ACKNOWLEDGEMENTS

The authors thank Ü. Seren, A. Korte, and M. Nordborg for kindly providing the full imputed

SNP data; and T. Tsuchimatsu and K. Iwayama for discussions. This study was supported by

the Japan Science and Technology Agency (JST) PRESTO (Grant number, JPMJPR17Q4 and

JPMJPR16Q9) to Y.S. and E.Y.; Japan Society for the Promotion of Science (JSPS)

Postdoctoral Fellowship (16J30005) to Y.S.; MEXT KAKENHI (18H04785) and the Swiss

National Science Foundation to K.K.S.; and JST CREST (JPMJCR15O2 and JPMJCR16O3)

to A.J.N. and K.K.S. The field experiment was supported by the Joint Usage/Research Grant

of Center for Ecological Research, Kyoto University, Japan.

p. 23

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA ARCHIVING

The leaf damage data on *A. thaliana* are included in the supporting information (Table S1).

The R source codes used in this study are available at the GitHub repository

(https://github.com/naganolab/NeighborGWAS).

REFERENCES

Alonso-Blanco C, Andrade J, Becker C, Bemm F, Bergelson J, Borgwardt KM et al. (2016)

1,135 genomes reveal the global pattern of polymorphism in *Arabidopsis thaliana*. *Cell* **166**:

481–491. doi:10.1016/j.cell.2016.05.063

Atwell S, Huang YS, Vilhjálmsson BJ, Willems G, Horton M, Li Y et al. (2010) Genome-

wide association study of 107 phenotypes in *Arabidopsis thaliana* inbred lines. *Nature* **465**:

627–631. doi:10.1038/nature08800

Azaele S, Muneepeerakul R, Rinaldo A, Rodriguez-Iturbe I (2010) Inferring plant ecosystem

organization from species occurrences. *J Theor Biol* **262**: 323–329.

doi:10.1016/j.jtbi.2009.09.026

Barbosa P, Hines J, Kaplan I, Martinson H, Szczepaniec A, Szendrei Z (2009) Associational

resistance and associational susceptibility: Having right or wrong neighbors. *Ann Rev Ecol*

Evol Sys **40**: 1–20. doi:10.1146/annurev.ecolsys.110308.120242

Benedetti M, Verrascina I, Pontiggia D, Locci F, Mattei B, De Lorenzo G, Cervone F (2018)

Four *Arabidopsis* berberine bridge enzyme-like proteins are specific oxidases that inactivate

the elicitor-active oligogalacturonides. *Plant J* **94**: 260–273. doi:10.1111/tpj.13852

p. 24

Bergvall UA, Rautio P, Kesti K, Tuomi J, Leimar O (2006) Associational effects of plant

defences in relation to within- and between-patch food choice by a mammalian herbivore:

neighbour contrast susceptibility and defence. *Oecologia* **147**: 253–260. doi:10.1007/s00442-

005-0260-8

Brachi B, Meyer CG, Villoutreix R, Platt A, Morton TC, Roux F, Bergelson J (2015)

Coselected genes determine adaptive variation in herbivore resistance throughout the native

range of *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* **112**: 4032–4037.

doi:10.1073/pnas.1421416112

Browning BL, Browning SR (2009) A unified approach to genotype imputation and

haplotype-phase inference for large data sets of trios and unrelated individuals. *Am J Hum*

Genet **84**: 210-223. doi:10.1016/j.ajhg.2009.01.005

Carlson M (2018) GO.db: A set of annotation maps describing the entire Gene Ontology. R

package version 3.7.0. doi:10.18129/B9.bioc.GO.db

Carrasco LR, Harwood TD, Toepfer S, MacLeod A, Levay N, Kiss J et al. (2010) Dispersal

kernels of the invasive alien western corn rootworm and the effectiveness of buffer zones in

eradication programmes in Europe. *Ann Appl Biol* **156**: 63–77. doi:10.1111/j.1744-

7348.2009.00363.x

Chan EKF, Rowe HC, Kliebenstein DJ (2010) Understanding the evolution of defense

metabolites in *Arabidopsis thaliana* using genome-wide association mapping. *Genetics* **185**:

991–1007. doi:10.1534/genetics.109.108522

Dahlin I, Rubene D, Glinwood R, Ninkovic V (2018) Pest suppression in cultivar mixtures is

influenced by neighbor-specific plant-plant communication. *Ecol Appl* **28**: 2187–2196.

doi:10.1002/eap.1807

Devaux C, Lavigne C, Austerlitz F, Klein EK (2007) Modelling and estimating pollen

p. 25

movement in oilseed rape (*Brassica napus*) at the landscape scale using genetic markers. *Mol Ecol* **16**: 487–499. doi:10.1111/j.1365-294X.2006.03155.x

Devoto A, Ellis C, Magusin A, Chang HS, Chilcott C, Zhu T, Turner JG (2005) Expression profiling reveals *COI1* to be a key regulator of genes involved in wound- and methyl jasmonate-induced secondary metabolism, defence, and hormone interactions. *Plant Mol Biol* **58**: 497–513. doi:10.1007/s11103-005-7306-5

Dicke M, Baldwin IT (2010) The evolutionary context for herbivore-induced plant volatiles: beyond the 'cry for help'. *Trends Plant Sci* **15**: 167–175. doi:10.1016/j.tplants.2009.12.002

Erb M (2018) Volatiles as inducers and suppressors of plant defense and immunity - origins, specificity, perception and signaling. *Curr Opin Plant Biol* **44**: 117–121. doi:10.1016/j.pbi.2018.03.008

Exposito-Alonso M, Drost H, Burbano HA, Weigel D (2020) The Earth BioGenome project: opportunities and challenges for plant genomics and conservation. *Plant J* doi:10.1111/tpj.14631

Frachon L, Mayjonade B, Bartoli C, Hautekèete NC, Roux F (2019) Adaptation to plant communities across the genome of *Arabidopsis thaliana*. *Mol Biol Evol* **36**: 1442–1456. doi:10.1093/molbev/msz078

Fisher CK, Mehta P (2015) Bayesian feature selection for high-dimensional linear regression via the Ising approximation with applications to genomics. *Bioinformatics* **31**: 1754–1761. doi:×

Gage JL, De Leon N, Clayton MK (2018) Comparing genome-wide association study results from different measurements of an underlying phenotype. *G3: Genes, Genomes, Genetics* **8**: 3715–3722. doi:10.1534/g3.118.200700

Geiselhardt S, Yoneya K, Blenn B, Drechsler N, Gershenzon J, Kunze R, Hilker M (2013)

p. 26

Egg laying of cabbage white butterfly (*Pieris brassicae*) on *Arabidopsis thaliana* affects

subsequent performance of the larvae. *PLoS ONE* **8**: e59661.

doi:10.1371/journal.pone.0059661

Gondro C, van der Werf J, Hayes B (eds.) (2013) Genome-wide association studies and

genomic prediction. *Methods Mol Biol*. Humana Press, New York. doi:10.1007/978-1-62703-

447-0

Hambäck PA, Björkman M, Rämert B, Hopkins RJ (2009). Scale-dependent responses in

cabbage herbivores affect attack rates in spatially heterogeneous systems. *Basic Appl Ecol* **10**:

228–236. doi:10.1016/j.baae.2008.06.004

Hambäck PA, Inouye BD, Andersson P, Underwood N (2014) Effects of plant neighborhoods

on plant-herbivore interactions: resource dilution and associational effects. *Ecology* **95**:

1370–1383. doi:10.1890/13-0793.1

Hamblin MT, Buckler ES, Jannink JL (2011) Population genetics of genomics-based crop

improvement methods. *Trends Genet* **27**: 98–106. doi:10.1016/j.tig.2010.12.003

Hauser MT, Harr B, Schlötterer C (2001) Trichome distribution in *Arabidopsis thaliana* and

its close relative *Arabidopsis lyrata*: molecular analysis of the candidate gene *GLABROUS1*.

Mol Biol Evol **18**: 1754–1763. doi:10.1093/oxfordjournals.molbev.a003963

Horton MW, Hancock AM, Huang YS, Toomajian C, Atwell S, Auton A et al. (2012)

Genome-wide patterns of genetic variation in worldwide *Arabidopsis thaliana* accessions

from the RegMap panel. *Nat Genet* **44**: 212–216. doi:10.1038/ng.1042

Horton MW, Bodenhausen N, Beilsmith K, Meng D, Muegge BD, Subramanian S et al.

(2014) Genome-wide association study of *Arabidopsis thaliana* leaf microbial community.

Nat Commun **5**: 5320. doi:10.1038/ncomms6320

Ida TY, Takanashi K, Tamura M, Ozawa R, Nakashima Y, Ohgushi T (2018) Defensive

chemicals of neighboring plants limit visits of herbivorous insects: Associational resistance

within a plant population. *Ecol Evol* **8**: 12981–12990. doi:10.1002/ece3.4750

Ising E (1925) Beitrag zur theorie des ferromagnetismus. *Zeitschrift für Physik* **31**: 253–258.

Jannink JL, Lorenz AJ, Iwata H (2010) Genomic selection in plant breeding: from theory to

practice. *Brief Funct Genomic* **9**: 166–177. doi:10.1093/bfgp/elq001

Kang HM, Zaitlen NA, Wade CM, Kirby A, Heckerman D, Daly MJ, Eskin E (2008)

Efficient control of population structure in model organism association mapping. *Genetics*

178: 1709–1723. doi:10.1534/genetics.107.080101

Karban R (1992) Plant variation: its effects on populations of herbivorous insects. In: Fritz

RS, Simms EL (eds) *Plant resistance to herbivores and pathogens: ecology, evolution, and*

genetics, University of Chicago Press: Chicago, pp. 195–215.

Kizaki S, Katori M (1999) Analysis of canopy-gap structures of forests by Ising-Gibbs states-

equilibrium and scaling property of real forests. *J Phys Soc Japan* **68**: 2553-2560.

doi:10.1143/JPSJ.68.2553

Kofler R, Schlotterer C (2012) Gowinda: unbiased analysis of gene set enrichment for

genome-wide association studies. *Bioinformatics* **28**: 2084–2085.

doi:10.1093/bioinformatics/bts315

Korte A, Farlow A (2013) The advantages and limitations of trait analysis with GWAS: a

review. *Plant Methods* **9**: 29. doi:10.1186/1746-4811-9-29

Li J, Hansen BG, Ober JA, Kliebenstein DJ, Halkier BA (2008) Subclade of flavin-

monooxygenases involved in aliphatic glucosinolate biosynthesis. *Plant Physiol* **148**: 1721–

1733. doi:10.1104/pp.108.125757

McCoy BM, Maillard JM (2012) The importance of the Ising model. Progress *Theor Phys*

127: 791-817. doi:10.1143/PTP.127.791

p. 28

McGale E, Valim H, Mittal D, Jimenez JM, Halitschke R, Schuman MC, Baldwin IT (2020)

Determining the scale at which variation in a single gene changes population yields. *eLife* **9**.

doi:10.7554/eLife.53517

Mundt CC (2002) Use of multiline cultivars and cultivar mixtures for disease management.

Ann Rev Phytopathol **40**: 381-410. doi:10.1146/annurev.phyto.40.011402.113723

Nallu S, Hill JA, Don K, Sahagun C, Zhang W, Meslin C et al. (2018) The molecular genetic

basis of herbivory between butterflies and their host plants. *Nat Ecol Evol* **2**: 1418-1427.

doi:10.1038/s41559-018-0629-9

Perdry H, Dandine-Roulland C (2018) gaston: Genetic Data Handling (QC, GRM, LD, PCA)

& Linear Mixed Models. R package version 1.5.4. https://CRAN.R-

project.org/package=gaston

R Core Team (2019) R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Reymond P, Farmer EE (1998) Jasmonate and salicylate as global signals for defense gene

expression. *Curr Opin Plant Biol* **1**: 404–411. doi:10.1016/s1369-5266(98)80264-1

Rieux A, Soubeyrand S, Bonnot F, Klein EK, Ngando JE, Mehl A et al. (2014) Long-distance

wind-dispersal of spores in a fungal plant pathogen: estimation of anisotropic dispersal

kernels from an extensive field experiment. *PLoS ONE* **9**: e103225.

doi:10.1371/journal.pone.0103225

Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC, Muller M (2011) pROC: an

open-source package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics*

12: 77. doi:10.1186/1471-2105-12-77

Sato Y (2018) Associational effects and the maintenance of polymorphism in plant defense

against herbivores: review and evidence. *Plant Species Biol* **33**: 91–108. doi:10.1111/1442-

1984.12201

Sato Y, Kudoh H (2015) Tests of associational defence provided by hairy plants for glabrous

plants of *Arabidopsis halleri* subsp. *gemmifera* against insect herbivores. *Ecol Entomol* **40**:

269–279. doi:10.1111/een.12179

Sato Y, Kudoh H (2017) Herbivore-mediated interaction promotes the maintenance of

trichome dimorphism through negative frequency-dependent selection. *Am Nat* **190**: E67–

E77. doi:10.1086/692603

Sato Y, Ito K, Kudoh H (2017) Optimal foraging by herbivores maintains polymorphism in

defence in a natural plant population. *Funct Ecol* **31**: 2233–2243. doi:10.1111/1365-

2435.12937

Sato Y, Shimizu-Inatsugi R, Yamazaki M, Shimizu KK, Nagano AJ (2019) Plant trichomes

and a single gene *GLABRA1* contribute to insect community composition on field-grown

Arabidopsis thaliana. *BMC Plant Biol* **19**: 163. doi:10.1186/s12870-019-1705-2

Schlicht R, Iwasa Y. (2004) Forest gap dynamics and the Ising model. *J Theoret Biol* **230**:

65–75. doi:10.1016/j.jtbi.2004.04.027

Schuman MC, Allmann S, Baldwin IT (2015) Plant defense phenotypes determine the

consequences of volatile emission for individuals and neighbors. *eLife* **4**: e04490.

doi:10.7554/eLife.04490

Tahvanainen JO, Root RB (1972) The influence of vegetational diversity on the population

ecology of a specialized herbivore, *Phyllotreta cruciferae* (Coleoptera: Chrysomelidae).

Oecologia **10**: 321–346. doi:10.1007/BF00345736

Togninalli M, Seren Ü, Meng D, Fitz J, Nordborg M, Weigel D et al. (2018) The AraGWAS

Catalog: a curated and standardized *Arabidopsis thaliana* GWAS catalog. *Nucleic Acids Res*

46: D1150–D1156. doi:10.1093/nar/gkx954

p. 30

Underwood N, Rausher MD (2000) The effects of host-plant genotype on herbivore

population dynamics. *Ecology* **81**: 1565-1576. doi:10.1890/0012-

9658(2000)081[1565:TEOHPG]2.0.CO;2

Underwood N, Inouye BD, Hambäck PA (2014) A conceptual framework for associational

effects: when do neighbors matter and how would we know? *Quart Rev Biol* **89**: 1–19.

doi:10.1086/674991

Underwood N, Hambäck PA, Inouye BD (2020) Pollinators, herbivores, and plant

neighborhood effects. *Quart Review Biol* **95**: 37–57. doi:10.1086/707863

Utsumi S, Ando Y, Craig TP, Ohgushi T (2011) Plant genotypic diversity increases population

size of a herbivorous insect. *Proc Royal Soc B* **278**: 3108–3115. doi:10.1098/rspb.2011.0239

van Poecke RM (2007) Arabidopsis-insect interactions. *The Arabidopsis Book* **5**: e0107.

doi:10.1199/tab.0107

Verschut TA, Becher PG, Anderson P, Hambäck PA (2016) Disentangling associational

effects: both resource density and resource frequency affect search behaviour in complex

environments. *Funct Ecol* **30**: 1826-1833. doi:10.1111/1365-2435.12670

Wang M, Roux F, Bartoli C, Huard-Chauveau C, Meyer C, Lee H et al. (2018) Two-way

mixed-effects methods for joint association analysis using both host and pathogen genomes.

Proc Natl Acad Sci USA **115**: E5440–E5449. doi:10.1073/pnas.1710980115

Weiner J (1990) Asymmetric competition in plant populations. *Trends Ecol Evol* **5**: 360-364.

doi:10.1016/0169-5347(90)90095-U

Wuest SE, Niklaus PA (2018) A plant biodiversity effect resolved to a single chromosomal

region. *Nat Ecol Evol* **2**: 1933–1939. doi:10.1038/s41559-018-0708-y

Yamamoto E, Matsunaga H, Onogi A, Ohyama A, Miyatake K, Yamaguchi H et al. (2017)

Efficiency of genomic selection for breeding population design and phenotype prediction in

p. 31

741   tomato. *Heredity* **118**: 202–209. doi:10.1038/hdy.2016.84

742   Zeller SL, Kalinina O, Flynn DFB, Schmid B (2012) Mixtures of genetically modified wheat

743   lines outperform monocultures. *Ecol Appl* **22**: 1817–1826. doi:10.1890/11-0876.1

TABLES & FIGURES

Table 1. Factors affecting the power to detect major-effect genes in simulated phenotypes.
ANOVA tables show the degree of freedom (df), sum of squares (SS), *F*-statistics, and *p*-
values. For the neighbor effects, the response variable was the maximum Area Under the
ROC Curve (AUC) of the spatial scales from $s = 1$ to $s = 3$. AUCs were arcsine square-root
transformed to improve normality. NA means not available.

Response	Factors	df	SS	F	p-value
AUC_{self}	No. of causal SNPs	1	30.2	2172.5	< 2e-16
	PVE_β	1	2.24	160.7	< 2e-16
	$PVE_\beta + PVE_u$	1	0.08	5.40	0.020
	$\sigma_1^2 : \sigma_2^2 : \sigma_{12}^2$	2	0.00	0.14	0.869
	α	1	0.02	1.71	0.192
	Residuals	1073	14.93	NA	NA
AUC_{nei}	No. of causal SNPs	1	24.0	1343.4	< 2e-16
	PVE_β	1	1.88	105.4	< 2e-16
	$PVE_\beta + PVE_u$	1	0.11	6.10	0.014
	$\sigma_1^2 : \sigma_2^2 : \sigma_{12}^2$	2	0.00	0.12	0.886
	α	1	1.46	81.8	< 2e-16
	Residuals	1073	19.20	NA	NA

p. 33

Figure 1. Relationship between Neighbor GWAS and Ising model. Upper panels show spatial

arrangements expected by a 2-D Ising model $\sum y_i = \beta_1 \sum x_i + \beta_2 \sum_{<i,j>} x_i x_j$. (a) If $\beta_2 > 0$,

mixed patterns give the argument of the minimum for a population sum of phenotype

values $\sum y_i$. (b) If $\beta_2 < 0$, clustered patterns give the argument of the minimum for $\sum y_i$. (c) In

addition, β_1 determines overall patterns favoring -1 or +1 states. The figure shows outcomes

from a random 100×100 lattice after 1000 iterations of simulated annealing. Conversely, the

neighbor GWAS was implemented as an inverse problem of the 2-D Ising model, where

genotypes and its spatial arrangement, x_i and $x_i x_j$, were given while the

coefficients β_1 and β_2 were to be estimated from observed phenotypes y_i. In addition, the

variance component due to self and neighbor effects was considered a random effect in a

linear mixed model, such that $u_i \sim \text{Norm}(0, \sigma_1^2 K_1 + \sigma_2^2 K_2)$. Once β_1 and β_2 were determined,

we could simulate a genotype distribution that maximizes or minimizes $\sum y_i$.

Figure 2. Experimental setting in the *Arabidopsis* herbivory data. (a) Photograph of the field

site. Each 0.6 × 2.6 m block included a replicate of 199 accessions and an additional Col-0.

(b) *Arabidopsis thaliana* plants were arranged in a 10 × 40 cell plastic tray. The plants were

placed in a checkered pattern so that alternate cells were left empty; therefore, each of the 40

rows contained 5 plants and each of the 10 columns contained 20 plants. Yellow lines

represent *s*-th neighbor scales from a focal *i*-th plant. (c) A graphical explanation of the

experimental area. A meadow (green) was separately covered with weed-masking sheets

(grey).

Figure 3. Receiver operating characteristic (ROC) curves representing the sensitivity (true positive rate) and the specificity (1 - false positive rate) in simulated phenotypes. The upper and lower rows give the results of self and neighbor effects with the number of causal SNPs at 10, 50, and 300. The shade of the lines indicates the degree of PVE_β: $PVE_\beta = 0.1$ (light gray), 0.3 (dark gray), 0.6 (black). The thickness of the lines indicates distance decay: $\alpha = 0.25$ (thin), 1 (medium), 3 (thick).

Figure 4. Scale dependence of neighbor effects on simulated phenotypes among all

iterations. The broad (a), intermediate (b), and narrow (c) effective range of neighbor effects

are represented by weak, moderate, and strong distance decay, respectively. The proportion of

phenotypic variation explained by neighbor effects (PVE_{nei}) and the area under the ROC

curve (AUC) of neighbor effects are shown along the spatial scale from the first nearest ($s =$

1) to the third nearest ($s = 3$) neighbors. Boxplots show center line: median, box limits: upper

and lower quartiles, whiskers: 1.5× interquartile range, and points: outliers.

p. 37

Figure 5. Proportion of phenotypic variation explained (PVE) by self-genotype or neighbor

effects on the leaf damage score of field-grown *Arabidopsis thaliana*. The PVEs at the spatial

scale of $s = 0$ (without any neighbor effects), $s = 1$ and $s = 2$ are given with different colors.

The red fraction indicates PVE by self effects as $\mathrm{PVE}_{\mathrm{self}} = \sigma_1^2/(\sigma_1^2 + \sigma_2^2 + \sigma_e^2)$, while the

blue fraction indicates PVE by neighbor effects as $\mathrm{PVE}_{\mathrm{nei}} (= \sigma_2^2/(\sigma_1^2 + \sigma_2^2 + \sigma_e^2)$. The white

fraction indicates residuals. Asterisks highlight a significant fraction with stepwise likelihood

ratio tests from simpler to complex models: [**]p-value < 0.01.

(a) Self, β_1

(b) Neighbor, β_2

Figure 6. Manhattan plots for the self (a) and neighbor (b) effects on the leaf damage score of

field-grown *Arabidopsis thaliana*. The first to fifth chromosomes of *A. thaliana* are

differently colored, where lighter plots indicate smaller MAF. Horizontal dashed lines

indicate the threshold after Bonferroni correction at *p*-value < 0.05. The red vertical line in

panel (a) indicates a SNP position near the *GS-OX2* locus, while three circles highlighted by a

black outline in panel (b) indicates the variants subject to the post hoc simulation (Fig. 7).

Results of the self and neighbor effects are shown at $s = 0$ (i.e., standard GWAS) and $s = 1$,

respectively.

(a) Chr 2, Position 14679190: $\beta_1 = 0.19$, $\beta_2 = 0.31$, $\Sigma y_i = -237.2$

(b) Chr 2, Position 9422409: $\beta_1 = -0.001$, $\beta_2 = -0.18$, $\Sigma y_i = -296.2$

(c) Chr 5, Position 19121831: $\beta_1 = 0.12$, $\beta_2 = -0.22$, $\Sigma y_i = -593.8$

(d) No effects: $\beta_1 = 10^{-6}$, $\beta_2 = 10^{-6}$, $\Sigma y_i = 10^{-4}$

Figure 7. Post hoc simulations exemplifying a spatial arrangement of two alleles expected by the estimated self and neighbor effects, $\hat{\beta}_1$ and $\hat{\beta}_2$, on the leaf damage score of *Arabidopsis thaliana*. Population sum of the leaf damage $\sum y_i = \beta_1 \sum x_i + \beta_2 \sum_{<i,j>} x_i x_j$ was minimized using 1000 iterations of simulated annealing from a random distribution of two alleles in a 10 × 40 space.