1 **Neighbor GWAS: incorporating neighbor genotypic identity**

2 **into genome-wide association studies of field herbivory on**

3 ***Arabidopsis thaliana***

4

5 Yasuhiro Sato[1,2], Eiji Yamamoto[1,3], Kentaro K. Shimizu[4,5*], Atsushi J.

6 Nagano[6*]

7

8 [1]PRESTO, Japan Science and Technology Agency, Kawaguchi 332-0012,

9 Japan

10 [2]Research Institute for Food and Agriculture, Ryukoku University, Yokotani

11 1-5, Seta Oe-cho, Otsu, Shiga 520-2194, Japan

12 [3]Graduate School of Agriculture, Meiji University, Higashi-Mita 1-1-1,

13 Tama-ku, Kawasaki, Kanagawa 214-8571, Japan

14 [4]Department of Evolutionary Biology and Environmental Studies,

15 University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

16 [5]Kihara Institute for Biological Research, Yokohama City University, 641-

17 12 Maioka, 244-0813 Totsuka-ward, Yokohama, Japan

18 [6]Faculty of Agriculture, Ryukoku University, Yokotani 1-5, Seta Oe-cho,

19 Otsu, Shiga 520-2194, Japan

20 *Co-correspondence:

21 Atsushi J. Nagano

22 anagano@agr.ryukoku.ac.jp

Kentaro K. Shimizu

kentaro.shimizu@ieu.uzh.ch

Running head: Neighbor GWAS of herbivory

ABSTRACT

An increasing number of field studies show that the phenotype of an

individual plant depends not only on its genotype but also on that of

neighboring plants; however, this fact is not taken into consideration in

genome-wide association studies (GWAS). Based on the Ising model of

ferromagnetism, we incorporated neighbor genotypic identity into a

regression model, named "Neighbor GWAS". Our simulations showed that

phenotypic variation explained by neighbor effects approached a plateau

when an effective spatial scale became narrow. Thus, the effective range of

neighbor effects could be estimated based on the scale at which an increase

in the explained variation peaked. The power to detect causal variants of

neighbor effects was moderate to strong when a trait was governed by tens

of variants. In contrast, the power decreased when hundreds of variants

underlay a single trait. We applied neighbor GWAS to field herbivory data

from 199 accessions of *Arabidopsis thaliana* and found a significant

contribution of the neighbor effects to the observed damage variation.

p. 2

Interestingly, a locus responsive to methyl jasmonate (AT2G34810 called

BBE16) was located near the second top-scoring single nucleotide

polymorphism (SNP) of the neighbor effects on the herbivory, while self-

genotype effects had the third top-scoring SNP near a locus encoding flavin-

monooxygenase glucosinolate S-oxygenase 2 (FMO GS-OX2). Overall,

neighbor GWAS highlight the overlooked role of plant neighborhood effects

in shaping phenotypic variation. The neighbor GWAS method is available as

an R package at CRAN (https://cran.r-

project.org/package=rNeighborGWAS), providing a novel tool to analyze

complex traits in spatially structured environments.

INTRODUCTION

Plants are immobile and thus cannot escape their neighbors. In natural and

agricultural systems, individual phenotypes depend not only on the plants'

own genotype but also on the genotypes of other neighboring plants

(Tahvanainen and Root 1972; Barbosa et al. 2009; Underwood et al. 2014).

This phenomenon has been termed neighbor effects or associational effects

in plant ecology (Barbosa et al. 2009; Underwood et al. 2014; Sato 2018).

Such neighbor effects were initially reported as a form of interspecific

interaction among different plant species (Tahvanainen and Root 1972), but

many studies have illustrated that neighbor effects occur among different

p. 3

genotypes within a plant species with respect to: (i) herbivory (Schuman et

al. 2015; Sato 2018; Ida et al. 2018), (ii) pathogen infections (Mundt 2002;

Zeller et al. 2012), and (iii) pollinator visitations (Underwood et al. 2020).

Although neighbor effects are of considerable interest in plant science

(Dicke and Baldwin 2010; Erb 2018) and agriculture (Zeller et al. 2012;

Dahlin et al. 2018), these effects are often not considered in quantitative

genetics of field-grown plants.

Complex mechanisms underlie neighbor effects through direct

competition (Weiner 1990), herbivore and pollinator movement (Bergvall et

al. 2006; Verschut et al. 2016; Underwood et al. 2020), and volatile

communication among plants (Schuman et al. 2015; Dahlin et al. 2018). For

example, lipoxygenase (*LOX*) genes govern jasmonate-mediated volatile

emissions in wild tobacco (*Nicotiana attenuata*) that induce defenses of

neighboring plants (Schuman et al. 2015). Even if direct plant–plant

communications are absent, herbivores can mediate indirect interactions

between plant genotypes (Sato and Kudoh 2017; Ida et al. 2018). *GLABRA1*

gene is known to determine hairy or glabrous phenotypes in *Arabidopsis*

plants (Hauser et al. 2001), where a flightless leaf beetle (*Phaedon*

brassicae) prefers glabrous plants to hairy ones (Sato et al. 2017).

Consequently, hairy plants escape herbivory when surrounded by glabrous

plants (Sato and Kudoh 2017). Yet, there are few hypothesis-free approaches

to seek key genetic variants responsible for plant neighborhood effects.

Genome-wide association studies (GWAS) have been increasingly

adopted to resolve the genetic architecture of complex traits in the model

plant, *Arabidopsis thaliana* (Atwell et al. 2010; Togninalli et al. 2018), and

crop species (Hamblin et al. 2011). The interactions of plants with

herbivores (Brachi et al. 2015; Nallu et al. 2018), microbes (Horton et al.

2014; Wang et al. 2018), and other plant species (Frachon et al. 2019) are

examples of such complex traits investigated through the lens of GWAS. To

distinguish causal variants from the genome structure, GWAS often employs

a linear mixed model with kinship considered as a random effect (Kang et

al. 2008; Korte and Farlow 2013). However, because of combinatorial

explosion, it is generally impossible to test the full set of inter-genomic

locus-by-locus interactions (Gondro et al. 2013); thus, some feasible and

reasonable approach should be developed for GWAS of neighbor effects.

To incorporate neighbor effects into GWAS, we focused on a

theoretical model of neighbor effects in magnetic fields, known as the Ising

model (Ising 1925; McCoy and Maillard 2012), which has been applied to

forest gap dynamics (Kizaki and Katori 1999; Schlicht and Iwasa 2004) and

community assembly (Azaele et al. 2010) in plant ecology. Using the Ising

analogy, we compare individual plants to a magnet: the two alleles at each

locus correspond to the north and south dipoles, and genome-wide multiple

testing across all loci is analogous to a number of parallel two-dimensional

layers. The Ising model has a clear advantage in its interpretability, such

p. 5

that: (i) the optimization problem for a population sum of trait values can be

regarded as an inverse problem of a simple linear model, (ii) the sign of

neighbor effects determines a model trend to generate a clustered or

checkered spatial pattern of the two states, and (iii) the self-genotypic effect

determines the general tendency to favor one allele over another (Fig. 1).

In this study, we proposed a new methodology integrating GWAS

and the Ising model, named "neighbor GWAS." The method was applied to

simulated phenotypes and real data of field herbivory on *A. thaliana*. We

addressed two specific questions: (i) what spatial and genetic factors

influenced the power to detect causal variants? and (ii) were neighbor

effects significant sources of leaf damage variation in field-grown *A.*

thaliana? Based on the simulation and application, we determined the

feasibility of our approach to detect neighbor effects in field-grown plants.

MATERIALS & METHODS

Neighbor GWAS

Basic model. We analyzed neighbor effects in GWAS as an inverse problem

of the two-dimensional Ising model, named "neighbor GWAS" hereafter

(Fig. 1). We considered a situation where a plant accession has either of two

alleles at each locus, and a number of accessions occupied a finite set of

field sites in a two-dimensional lattice. The allelic status at each locus was

p. 6

represented by x, and so the allelic status at each locus of the i-th focal plant

and the j-th neighboring plants was designated as $x_{i(j)} \in \{-1, +1\}$. Based on

a two-dimensional Ising model, we defined a phenotype value of i-th focal

individual plant y_i as:

$$y_i = \beta_1 x_i + \beta_2 \sum_{<i,j>} x_i x_j \qquad (1)$$

where β_1 and β_2 denoted self-genotype and neighbor effects, respectively. If

two neighboring plants shared the same allele at a given locus, the

product $x_i x_j$ turned into (-1)×(-1)=+1 or (+1)×(+1)=+1. If two neighbors

had different alleles, the product $x_i x_j$ became (-1)×(+1)=-1 or (+1)×(-1)=-1.

Accordingly, the effects of neighbor genotypic identity on a phenotype

depended on the coefficient β_2 and the number of two alleles in a

neighborhood. If the numbers of identical and different alleles were the

same near a focal plant, these neighbors offset the sum of the products

between the focal plant i and all j neighbors $\sum_{<i,j>} x_i x_j$ and exerted no

effects on a phenotype. When we summed up the phenotype values for the

total number of plants n and replaced it as $E = -\beta_2$, $H = -\beta_1$ and $\epsilon_I =$

$\sum y_i$, eq. 1 could be transformed as $\epsilon_I = -E \sum_{<i,j>} x_i x_j - H \sum x_i$, which

defined the interaction energy of a two-dimensional ferromagnetic Ising

model (McCoy and Maillard 2012). The neighbor effect β_2 and self-

genotype effect β_1 were interpreted as the energy coefficient E and external

magnetic effects H, respectively. An individual plant represented a spin and

the two allelic states of each locus corresponded to a north or south dipole.

p. 7

The positive or negative value of $\sum x_i x_j$ indicated a ferromagnetism or

paramagnetism, respectively. In this study, we did not consider the effects of

allele dominance because this model was applied to inbred *A. thaliana*.

However, heterozygotes could be processed if the neighbor

covariate $x_i x_j$ was weighted by an estimated degree of dominance in the

self-genotypic effects on a phenotype.

Association tests. For association mapping, we needed to

determine β_1 and β_2 from observed phenotypes and considered a

confounding sample structure as advocated by previous GWAS (e.g., Kang

et al. 2008; Korte and Farlow 2013). Extending the basic model eq. 1, we

described a linear mixed model at an individual level as:

$$y_i = \beta_0 + \beta_1 x_i + \frac{\beta_2}{L} \sum_{<i,j>}^{L} x_i x_j^{(s)} + u_i + e_i \qquad (2)$$

where β_0 indicated the intercept, and the term $\beta_1 x_i$ represented fixed self-

genotype effects as tested in standard GWAS; β_2 was the coefficient of fixed

neighbor effects. The neighbor covariate $\sum_{<i,j>}^{L} x_i x_j^{(s)}$ indicated a sum of

products for all combinations between the *i*-th focal plant and the *j*-th

neighbor at the *s*-th spatial scale from the focal plant *i*, and was scaled by

the number of neighboring plants, *L*. The number of neighboring

plants *L* was dependent on the spatial scale *s* to be referred. Variance

components due to a sample structure in self and neighbor effects were

modeled by a random effect $u_i \sim \text{Norm}(0, \sigma_1^2 K_1 + \sigma_2^2 K_2)$. The residual was

p. 8

expressed as $e_i \sim \text{Norm}(0, \sigma_e^2)$.

Variation partitioning. To estimate the proportion of phenotypic

variation explained (PVE) by self and neighbor effects, we utilized variance

component parameters in linear mixed models. The $n \times n$ variance-

covariance matrices represented the similarity in self-genotypes (i.e.,

kinship) and neighbor covariates among n individual plants as $K_1 =$

$\frac{1}{q-1} X_1^T X_1$ and $K_2 = \frac{1}{q-1} X_2^T X_2$, where q indicated the number of markers. As

we defined $x_{i(j)} \in \{+1, -1\}$, the elements of the kinship matrix K_1 were

scaled to represent the proportion of marker loci shared among $n \times n$ plants

such that $K_1 = \left(\frac{k_{ij}+1}{2}\right)$; σ_1^2 and σ_2^2 indicated variance component parameters

for the self and neighbor effects.

The elements of n plants $\times$ q markers matrix X_1 and X_2 consisted of

explanatory variables for the self and neighbor effects as $X_1 =$

(x_i) and $X_2 = \left(\frac{\sum_{<i,j>}^{L} x_i x_j^{(s)}}{L}\right)$. The individual-level formula eq. 2 could also be

converted into a conventional matrix form as:

191                        $$y = X\beta + Zu + e \qquad (3)$$

192   where y was $n \times 1$ vector of phenotypes; X was a matrix of fixed effects,

193   including mean, self-genotype x_i, neighbor covariate $\frac{\sum_{<i,j>}^{L} x_i x_j^{(s)}}{L}$, and other

194   confounding covariates for n plants; β was a vector that represents

195   coefficients of the fixed effects; Z was a design matrix allocating individuals

p. 9

to a genotype, and became an identity matrix if all plants were different

accessions; $\boldsymbol{u}$ was the random effect with $\mathrm{Var}(\boldsymbol{u}) = \sigma_1^2 \boldsymbol{K}_1 + \sigma_2^2 \boldsymbol{K}_2$; and $\boldsymbol{e}$ was

residual as $\mathrm{Var}(\boldsymbol{e}) = \sigma_e^2 \boldsymbol{I}$. The proportion of phenotypic variation explained

(PVE) by self and neighbor effects is calculated using a single-random

effect model defined by 'single' $\mathrm{PVE}_{\mathrm{self}} = \sigma_1^2 / (\sigma_1^2 +$

$\sigma_e^2)$ when s and σ_2^2 were set at 0; or 'single' $\mathrm{PVE}_{\mathrm{nei}} = \sigma_2^2 / (\sigma_2^2 +$

$\sigma_e^2)$ when σ_1^2 was set at 0 in eq. 3. Furthermore, we could partial out either

of the two variance components using a two-random effect model and

define these metrics as 'partial' $\mathrm{PVE}_{\mathrm{self}} = \sigma_1^2 / (\sigma_1^2 + \sigma_2^2 + \sigma_e^2)$ and 'partial'

$\mathrm{PVE}_{\mathrm{nei}} = \sigma_2^2 / (\sigma_1^2 + \sigma_2^2 + \sigma_e^2)$. The partial $\mathrm{PVE}_{\mathrm{self}}$ was equivalent to the

single $\mathrm{PVE}_{\mathrm{self}}$ when s was set at 0.

We should be cautious when applying the model structure

underlying neighbor GWAS to data. The standard GWAS is a subset of the

neighbor GWAS and these two models become equivalent at $s = 0$ and $\sigma_2^2 =$

0. When testing the self-genotype effect β_1, we recommend that neighbor

effects and its variance component σ_2^2 should be excluded; otherwise, the

standard GWAS fails to correct a sample structure because of the additional

variance component at $\sigma_2^2 \neq 0$. Furthermore, if the neighbor effects range

was broad enough to encompass an entire field (i.e., $s \to \infty$), the neighbor

covariate and self-genotype x_i would become colinear according to the

following equation: $\left(\sum_{<i,j>}^{L} x_i x_j^{(s)}\right)/L = x_i \left(\sum_{j=1}^{L} x_j^{(s)}\right)/L = x_i \bar{x}_j,$

p. 10

where $\bar{x}_j$ indicates a population-mean of neighbor genotypes and

corresponds to a population-mean of self-genotype values $\bar{x}_i$, if $s \to \infty$.

Such a collinearity makes distinguishing self and neighbor effects for

neighbor GWAS challenging, when the effective range is broad. Therefore,

the maximum range of neighbor effects was restricted during simulation as

described below.

Power simulation

To examine the power to detect neighbor effects, we applied the neighbor

GWAS to simulated phenotypes. Phenotypes were simulated using a part of

the real genotypes of *A. thaliana*. To evaluate the power of the simple linear

model, we assumed a complex ecological form of neighbor effects with

multiple variance components controlled. The power was evaluated by the

receiver operating characteristic (ROC) curve on the association score of -

$\log_{10}(p\text{-value})$ (e.g., Gage et al. 2018). All analyses were performed using R

version 3.6.0 (R Core Team 2019).

Genotype data. To consider a realistic genetic structure in the

simulation, we used part of the *A. thaliana* RegMap panel (Horton et al.

2012). The genotype data on 1,307 accessions were downloaded from the

Joy Bergelson laboratory website

(http://bergelson.uchicago.edu/?page_id=790 accessed on February 9,

2017). We extracted the chromosome 1 and 2 data with the minor allele

p. 11

frequency (MAF) at >0.1, providing a matrix of 1,307 plants with 65,226

single nucleotide polymorphisms (SNPs). Pairwise linkage disequilibrium

(LD) among the loci was $r^2 = 0.003$ [0.00-0.06: median with upper and

lower 95 percentiles]. Before generating a phenotype, each locus was

standardized to have a mean of zero and a variance of 1. Subsequently, we

randomly selected 1,296 accessions (= 36 × 36 accessions) without any

replacements for each iteration and placed them in a 36 × 72 checkered

space following *Arabidopsis* experimental settings (see Fig. 2b).

Phenotype simulation. To address ecological issues specific to

plant neighborhood effects, we considered two extensions, namely

asymmetric neighbor effects and spatial scales. Firstly, previous studies

have shown that such plant–plant interactions are sometimes asymmetric

between two accessions in herbivory (e.g., Bergvall et al. 2006; Verschut et

al. 2016; Sato and Kudoh 2017) and height competition (Weiner 1990).

Such asymmetric neighbor effects can be tested by statistical interaction

terms in a linear model (Bergvall et al. 2006; Sato and Kudoh 2017).

Secondly, several studies have shown that the strength of neighbor effects

depends on spatial scales (Hambäck et al. 2014), and that the scale of

neighbors to be analyzed relies on the dispersal ability of causative

organisms (see Hambäck et al. 2009; Sato and Kudoh 2015; Verschut et al.

2016; Ida et al. 2018 for insect and mammal herbivores; Rieux et al. 2014

for pathogen dispersal) or the size of competing plants (Weiner 1990). We

p. 12

assumed the distance decay at s-th sites from a focal individual i with the

decay coefficient α as $w(s, \alpha) = \mathrm{e}^{-\alpha(s-1)}$, since such an exponential

distance decay has been widely adopted in empirical studies (Devaux et al.

2007; Carrasco et al. 2010; Rieux et al. 2014; Ida et al. 2018). Therefore, we

assumed a more complex model for simulated phenotypes than the model

for neighbor GWAS as follows:

$$y_i = \beta_0 + \beta_1 x_i + \frac{\beta_2}{L}\sum_{<i,j>}^{L} w(s,\alpha)x_i x_j^{(s)} + \beta_{12}\frac{x_i}{L}\sum_{<i,j>}^{L} w(s,\alpha)x_i x_j^{(s)} +$$

$$u_i + e_i \quad (4)$$

where β_{12} was the coefficient for asymmetry in neighbor effects. By

incorporating an asymmetry coefficient, the model eq. 4 can deal with cases

where neighbor effects are one-sided or occur irrespective of a focal

genotype. Total variance components resulting from three background

effects (i.e., the self, neighbor, and self-by-neighbor effects) was defined

as $u_i \sim \mathrm{Norm}(0, \sigma_1^2 K_1 + \sigma_2^2 K_2 + \sigma_{12}^2 K_{12})$. The three variance component

parameters σ_1^2, σ_2^2, and σ_{12}^2, determined the relative importance of self-

genotype, neighbor, and asymmetric neighbor effects in u_i. Given the

elements of the n plants $\times$ q marker explanatory matrix with $X_{12} =$

$(\frac{x_i}{L}\sum_{<i,j>}^{L} w(s,\alpha)x_i x_j^{(s)})$, the similarity in asymmetric neighbor effects was

calculated as $K_{12} = \frac{1}{q-1}X_{12}^T X_{12}$. To control phenotypic variations, we

further partitioned the proportion of phenotypic variation into those

explained by major-effect genes and variance components $PVE_\beta + PVE_u$,

p. 13

major-effect genes alone PVE*β*, and residual error PVE*e*, where PVE*β* +

PVE*u* + PVE*e* = 1. The *optimize* function in R was used to adjust simulated

phenotypes to the given amounts of PVE.

Parameter setting. Ten phenotypes were simulated for each

combination of the distance decay coefficient α, the proportion of

phenotypic variation explained by major-effect genes PVE*β*, the proportion

of phenotypic variation explained by major-effect genes and variance

components PVE*β* + PVE*u*, and the relative importance of multiple variance

components $\sigma_1^2 : \sigma_2^2 : \sigma_{12}^2$ as: $\alpha = 0.25$, 1.0 or 3.0, $\sigma_1^2 : \sigma_2^2 : \sigma_{12}^2 = 6{:}3{:}1$, 4:4:1, or

3:6:1, PVE*β* and PVE*β* + PVE*u* = 0.1 and 0.4, 0.3 and 0.4, 0.3 and 0.8, or 0.6

and 0.8. The maximum reference scale was fixed at $s = 3$. The line of

simulations was repeated for the number of causal SNPs at 10, 50, or 300 to

examine cases of an oligogenic and polygenic control for a trait. The non-

zero coefficients for the causal SNPs were randomly sampled from -1 or 1

digit and then assigned as some causal SNPs were responsible for both the

self and neighbor effects. Of the total number of causal SNPs, 15% had all

self, neighbor, and asymmetric neighbor effects (i.e., $\beta_1 \neq 0$ and $\beta_2 \neq$

0 and $\beta_{12} \neq 0$); another 15% had both the self and neighbor effects, but no

asymmetry in the neighbor effects ($\beta_1 \neq 0$ and $\beta_2 \neq 0$ and $\beta_{12} = 0$);

another 35% had self-genotypic effects only ($\beta_1 \neq 0$); and the remaining

35% had neighbor effects alone ($\beta_2 \neq 0$). Given its biological significance,

we assumed that some loci having neighbor signals possessed asymmetric

p. 14

interactions between neighbors ($\beta_2 \neq 0$ and $\beta_{12} \neq 0$) while the others had

symmetric interactions ($\beta_2 \neq 0$ and $\beta_{12} = 0$). Therefore, the number of

causal SNPs in β_{12} was smaller than that in the main neighbor effects β_2.

According to this assumption, the variance component σ_{12}^2 was also assumed

to be smaller than σ_2^2.

Summary statistics. The simulated phenotypes were fitted by eq. 2

to test the significance of coefficients β_1 and β_2, and to estimate single or

partial PVE_{self} and PVE_{nei}. The stepwise likelihood ratio tests were

performed first between the null model and model with a self-genotype

effect alone, and then between the self-genotype effect model and model

with both self and neighbor effects. The likelihood ratio was calculated as

the difference in deviance, i.e., -2 × log-likelihood, which is asymptotically

χ^2 distributed with one degree of freedom. The variance

components, σ_1^2 and σ_2^2, were estimated using a linear mixed model without

any fixed effects. To solve the mixed model with the two random effects, we

used the average information restricted maximum likelihood (AI-REML)

algorithm implemented in the *lmm.aireml* function in the *gaston* package of

R (Perdry and Dandine-Roulland 2018). Subsequently, we replaced the two

variance parameters σ_1^2 and σ_2^2 in eq. 2 into their estimates $\hat{\sigma}_1^2$ and $\hat{\sigma}_2^2$ by the

AI-REML, and performed association tests by solving a linear mixed model

with a fast approximation by an eigen value decomposition (implemented in

the *lmm.diago* function: Perdry and Dandine-Roulland 2018). The model

p. 15

likelihood was computed using the *lmm.diago.profile.likelihood* function.

We evaluated self and neighbor effects for association mapping based on

forward selection of the two fixed effects β_1 and β_2 as described below:

1. Computed the null likelihood with $\sigma_1^2 \neq 0$ and $\sigma_2^2 = 0$ in eq. 2

2. Tested the self-effect, β_1, by comparing with the null likelihood

3. Computed the self-likelihood with $\hat{\sigma}_1^2$, $\hat{\sigma}_2^2$, and β_1 using eq. 2

4. Tested the neighbor effects, β_2, by comparing with the self-likelihood

We also calculated PVE using the mixed models eq. 3 without β_1 and β_2 as

follows:

1. Calculated single PVE_{self} or single PVE_{nei} by setting either σ_1^2 or σ_2^2 at 0.

2. Tested the single PVE_{self} or single PVE_{nei} by likelihood ratio between the

null and one-random effect model

3. Calculated partial PVE_{self} and partial PVE_{nei} by

estimating σ_1^2 and σ_2^2 simultaneously

4. Tested the partial PVE_{self} and partial PVE_{nei} by likelihood ratio between

the two- and one-random effect model

True or false positive rates were evaluated by ROC curves and area

under the ROC curves (AUC) (Gage et al. 2018). An AUC of 0.5 would

indicate that GWAS has no power to detect true signals, while an AUC of

1.0 would indicate that all the top signals predicted by GWAS agree with

true signals. The roc function in the pROC package (Robin et al. 2011) was

used to calculate AUC from $-\log_{10}(p\text{-value})$. Factors affecting the AUC were

p. 16

tested by analysis-of-variance (ANOVA) for the self or neighbor effects

(AUC$_{self}$ or AUC$_{nei}$). The AUC had proportional values, and thus was

arcsine-square-root transformed to improve the Shapiro–Wilk normality

statistic before ANOVA. The AUC and PVE were calculated from $s = 1$ (the

first nearest neighbors) to $s = 3$ (up to the third nearest neighbors) cases. The

AUC was also calculated using standard linear models without any random

effects to examine whether the linear mixed models were superior to the

linear models. We also tested the neighbor GWAS model incorporating

neighbor phenotype $y_j^{(s)}$ instead of $x_j^{(s)}$.

***Arabidopsis* herbivory data**

We applied the neighbor GWAS to field data of *Arabidopsis* herbivory. The

procedure of the present field experiment followed our previous experiment

on summer herbivory on field-grown *A. thaliana* (Sato et al. 2019). We

selected 199 worldwide accessions from 2029 accessions sequenced by the

RegMap (Horton et al. 2012) and 1001 Genomes project (Alonso-Blanco et

al. 2016). Of the 199 accessions, most were overlapped with a previous

GWAS of biotic interactions (Horton et al. 2014) and half were included by

a GWAS of glucosinolates (Chan et al. 2010). Eight replicates of each of the

199 accessions were first prepared in a laboratory and then transferred to the

outdoor garden at the Center for Ecological Research, Kyoto University,

Japan (Otsu, Japan: 35°06′N, 134°56′E, alt. ca. 200 m: Fig. 2a). Seeds were

p. 17

sown on Jiffy-seven pots (33-mm diameter), and stratified at a temperature

of 4 °C for a week. Seedlings were cultivated for 1.5 months under a short-

day condition (8 h light: 16 h dark, 20 °C). Plants were then separately

potted in plastic pots (6 cm in diameter) filled with mixed soil of

agricultural compost (Metro-mix 350, SunGro Co., USA) and perlite at a

3:1 ratio. Potted plants were set in plastic trays (10 × 40 cells) in a

checkered pattern (Fig. 2b). In the field setting, a set of 199 accessions and

an additional Col-0 accession were randomly assigned to each block without

replacement (Fig. 2b). Eight replicates of these blocks were set >2 m apart

from each other (Fig. 2c). Potted plants were exposed to the field

environment for 3 wk in June 2017. At the end of experiment, we scored the

percentage of foliage eaten as: 0 for no visible damage, 1 for ≤10%, 2 for

>10% and ≤ 25%, 3 for > 25% and ≤ 50%, 4 for >50% and ≤ 75%, and 5 for

>75% of the foliage area eaten. All plants were scored by a single person to

avoid observer bias. The most predominant herbivore in this field trial was

the diamond back moth (*Plutella xylostella*), followed by the small white

butterfly (*Pieris rapae*). We also recorded the initial plant size and the

presence of inflorescence to incorporate them as covariates. Initial plant size

was evaluated by the length of the largest rosette leaf (mm) at the beginning

of the field experiment and the presence of inflorescence was recorded 2 wk

after transplanting.

We estimated the variance components and performed the

p. 18

association tests for the leaf damage score with the neighbor covariate at $s =$

1 and 2. These two scales corresponded to $L = 4$ (the nearest four neighbors)

and $L = 12$ (up to the second nearest neighbors), respectively, in the

Arabidopsis dataset. The variation partitioning and association tests were

performed using the *gaston* package, as mentioned above. To determine the

significance of variance component parameters, we compared the likelihood

between mixed models with one or two random effects. For the genotype

data, we used an imputed SNP matrix of all the 2029 accessions studied by

the RegMap (Horton et al. 2012) and 1001 Genomes project (Alonso-

Blanco et al. 2016). Missing genotypes were imputed using BEAGLE

(Browning and Browning 2009), as described by Togninalli et al. (2018) and

updated on the AraGWAS Catalog (https://aragwas.1001genomes.org). Of

the 10,709,466 SNPs from the full imputed matrix, we used 1,242,128 SNPs

with MAF at > 0.05 and LD of adjacent SNPs at $r^2 < 0.8$. We considered the

initial plant size, presence of inflorescence, experimental blocks, and the

edge or center within a block as fixed covariates; these factors explained

12.5% of the leaf damage variation (1.2% by initial plant size, Wald test, $Z =$

3.53, p-value<0.001; 2.4% by the presence of inflorescence, $Z = -5.69$, p-

value$<10^{-8}$; 8.3% by the experimental blocks, likelihood ratio test, $\chi^2 =$

152.8, df $= 7$, p-value$<10^{-28}$; 0.5% by the edge or center, $Z = 3.11$, p-value $=$

0.002). After the association mapping, we searched candidate genes within

~10 kb around target SNPs, based on the Araport11 gene model with the

p. 19

latest annotation of The Arabidopsis Information Resource (TAIR)

(accessed on 7 September 2019). Gene-set enrichment analysis was

performed using the Gowinda algorithm that enables unbiased analysis of

GWAS results (Kofler and Schlotterer 2012). We tested SNPS with the top

0.1% -$\log_{10}$(p-value) score with the option "--gene-definition

undownstream10000," "--min-genes 20," and "--mode gene." The GO.db

package (Carlson et al. 2018) and the latest TAIR AGI code annotation were

used to build input files. To assess implementation, we performed standard

GWAS using GEMMA version 0.98 (Zhou and Stephens 2012) and

confirmed that the neighbor GWAS yielded the same -$\log_{10}$(p-values) scores

for self-effects as the GEMMA program ($r = 0.9999$ among all the SNPs).

The R source codes, accession list, and phenotype data are available at the

GitHub repository (https://github.com/naganolab/NeighborGWAS).

R package, "rNeighborGWAS"

To increase the availability of the new method, we have developed the

neighbor GWAS into an R package, which is referred to as

"rNeighborGWAS". In addition to the genotype and phenotype data, the

package requires a spatial map indicating positions of individuals across a

space. In this package, we generalized the discrete space example into a

continuous two-dimensional space, allowing the handling of any spatial

distribution along the x- and y-axes. Based on the three input files, the

p. 20

rNeighborGWAS package estimates the effective range of neighbor effects

by calculating partial PVE_{nei} and performs association mapping of neighbor

effects using the linear mixed models described earlier. Details and usage

are described in the help files and vignette of the rNeighborGWAS package

available via CRAN at https://cran.r-project.org/package=rNeighborGWAS.

RESULTS

Power simulation

A set of phenotypes were simulated from real genotype data following a

complex model (eq. 4), and then fitted by a simplified model (eq. 2).

Analyzing the factors affecting the AUC, we found that the number of

causal SNPs, the proportion of phenotypic variation (PVE) explained by

major-effect genes PVE_{β}, and distance decay coefficient α were the most

influential factors affecting the power of the proposed method to detect

neighbor signals (Table 1). In addition to PVE_{β}, the amount of variance

components PVE_{u} also significantly affected the AUC of the neighbor

effects, but these additional effects were less significant compared to those

of PVE_{β} alone (Table 1). The power of the standard GWAS to detect self-

genotype effects depended on the number of causal SNPs and PVE_{β}, but

was not significantly influenced by the distance decay coefficient of

p. 21

neighbor effects (Table 1). Neither the power to detect self nor neighbor

effects were significantly affected by the ratio of three variance

components $\sigma_1^2 : \sigma_2^2 : \sigma_{12}^2$ (Table 1).

The power to detect self-genotype signals changed from strong

(AUC$_{\text{self}}$>0.9) to weak (AUC$_{\text{self}}$<0.6) as PVE$_\beta$ decreased and the number of

causal SNPs increased (Fig. 3a; Fig. S1a). Compared to the self-genotype

effects, it was relatively difficult to detect neighbor effects (Fig. 3b; Fig.

S1b). In the case of the number of causal SNPs = 10, the power to detect

neighbor signals decreased from high (AUC$_{\text{nei}}$>0.9) to moderate

(AUC$_{\text{nei}}$>0.7) with the decreasing PVE$_\beta$ and distance decay coefficient (Fig.

3b; Fig. S1b). The power to detect neighbor effects was very weak

(AUC$_{\text{nei}}$>0.5 and AUC$_{\text{nei}}$<0.6) in the case of the number of causal SNPs =

300 (Fig. 3b; Fig. S1b). The result of simulations indicated that neighbor

effects were detectable when a target trait was governed by several major

genes and the range of neighbor effects was spatially limited. Additionally,

linear mixed models outperformed standard linear models as shown by 9.7%

and 1.7% increases in the power to detect self and neighbor signals,

respectively (AUC$_{\text{self}}$, paired t-test, mean of the difference = 0.097, p-

value<10^{-16}; AUC$_{\text{nei}}$, mean of the difference = 0.017, p-value<10^{-16}). When

the neighbor phenotype $y_j^{(s)}$ was incorporated instead of the genotype $x_j^{(s)}$,

the power to detect neighbor effects was very weak, such that AUC$_{\text{nei}}$

decreased to near 0.5.

Notably, there was a clear relationship between the distance decay

coefficient α and the proportion of phenotypic variation explained by

neighbor effects partial PVE_{nei} or AUC_{nei} at different spatial scales (Fig. 4).

If the distance decay was weak (small value of decay coefficient α) and the

effective range of neighbor effects was broad, partial PVE_{nei} and AUC_{nei}

increased linearly as the reference spatial scale was broadened (Fig. 4a). On

the other hand, if the distance decay was strong (large value of decay

coefficient α) and the effective scale of neighbor effects was narrow, partial

PVE_{nei} saturated at the scale of the first nearest neighbors (Fig. 4c) and

AUC_{nei} did not increase (Fig. 4b, c). The simulation results indicated that the

effective spatial scales could be estimated by the scale s at which the partial

PVE_{nei} peaked, and that the first nearest scale was enough to detect neighbor

signals unless the distance decay was very weak.

Examination of variance components provided further insights into

PVE_{nei} estimation (Fig. S2). Comparing the true and estimated PVE, we

found that the sum of partial PVE_{self} and partial PVE_{nei} matched the total

PVE attributable to the model at the nearest neighbor scale (Fig. S2a).

Single PVE_{self} or single PVE_{nei} overestimated the respective true PVE_{self} or

PVE_{nei} (Fig. S2b, c). In terms of total PVE, a joint mixed model including

the partial PVE_{self} and partial PVE_{nei} outperformed that including the single

PVE_{self} alone and compensated for the remaining total PVE unexplained by

the single PVE_{self} (Fig. S2a). The partial PVE_{nei} underestimated the true

p. 23

PVE_{nei} depending on the distance decay coefficient (Fig. S2e); if the

distance decay was strong ($\alpha = 3$), the underestimation of true PVE_{nei} was

moderate even at the nearest scale ($s = 1$: Fig. S2e), but if the distance decay

was weak ($\alpha = 0.25$), the bias was large at $s = 1$ and the broader scale ($s = 3$)

should be referred (Fig. S2e). Once the effective range is determined by

increasing patterns of the partial PVE_{nei} across s, the partial PVE_{nei} provides

a conservative proxy to true PVE_{nei}.

***Arabidopsis* herbivory data**

To estimate PVE_{self} and PVE_{nei}, we applied a linear mixed model (eq. 3) for

the leaf damage score of field-grown *A. thaliana*. Variation partitioning

showed a significant contribution of neighbor effects to phenotypic variation

in the leaf damage at the nearest scale (single $PVE_{nei} = 0.248$, $\chi_1^2 = 17.8$, p-

value $= 10^{-5}$; partial $PVE_{nei} = 0.214$, $\chi_1^2 = 7.23$, p-value $= 0.004$ at $s = 1$: Fig.

5). The proportion of phenotypic variation explained by neighbor effects did

not increase when the neighbor scale was referred up to the nearest and

second nearest individuals (single $PVE_{nei} = 0.243$, $\chi_1^2 = 11.1$, p-value $=$

0.0005; partial $PVE_{nei} = 0.14$, $\chi_1^2 = 1.41$, p-value $= 0.166$ at $s = 2$: Fig. 5);

therefore, the effective scale of neighbor effects was estimated at $s = 1$ and

variation partitioning was stopped at $s = 2$. Notably, these neighbor effects

explained the leaf damage variation better than self-genotype effects, since

single PVE_{nei} was larger than single PVE_{self} at $s = 0$ (single $PVE_{self} = 0.173$,

p. 24

$\chi_1^2 = 10.1$, p-value = 0.005) and partial PVE_{nei} was larger than the partial

PVE_{self} (partial $PVE_{self} = 0.041$, $\chi_1^2 = 0.36$, p-value = 0.54 at $s = 1$; partial

$PVE_{self} = 0.091$, $\chi_1^2 = 1.19$, p-value = 0.200 at $s = 2$: Fig. 5). We additionally

incorporated the neighbor phenotype $y_j^{(s)}$ instead of the neighbor

genotype $x_j^{(s)}$ in eq. 2, but partial PVE_{nei} did not increase (partial $PVE_{nei} = $

0.066 and 0.068 at $s = 1$ and 2, respectively). These results indicated a

narrow effective scale and 15-20% variation, which was explained by

neighbor effects on the leaf damage.

Standard GWAS of the self-genotype effects on the leaf damage

detected SNPs with the second and third largest -$\log_{10}(p$-values) score on

the first chromosome (chr1), though they were not above a threshold of

Bonferroni correction (Fig. 6a; Table S2). The second SNP at chr1-

21694386 was located within ~10 kb of three loci encoding a disease

resistance protein (CC-NBS-LRR class) family. The third SNP at chr1-

23149476 was located within ~10 kb of the AT1G62540 locus that encoded

flavin-monooxygenase glucosinolate S-oxygenase 2 (FMO GS-OX2). No

GOs were significantly enriched for the self-effects on herbivory (false

discovery rate > 0.08). A QQ-plot did not exhibit an inflation of p-values for

the self-genotype effects (Fig. S3).

Regarding the neighbor effects on the leaf damage, we found non-

significant but weak peaks on the second and third chromosomes (Fig. 6b;

Table S2). The second chromosomal region had higher association scores

p. 25

than predicted by the QQ-plot (Fig. S3). A locus encoding FAD-binding

Berberine family protein (AT2G34810 named *BBE16*) was located within

the ~10 kb window near SNPs with the top eleven $-\log_{10}(p$-values) score at

the second chromosome, which are known to be up-regulated by methyl

jasmonate (Devoto et al. 2005). Three transposable elements and a

pseudogene of lysyl-tRNA synthetase 1 were located near the most

significant SNP on the third chromosome. No GOs were significantly

enriched for the self-effects on herbivory (false discovery rate > 0.9). We

additionally tested asymmetric neighbor effects β_{12} in the real dataset on

field herbivory, but the top 0.1% SNPs of neighbor effects β_2 did not

overlap with those of asymmetric neighbor effects β_{12} (Table S2).

Based on the estimated coefficients $\hat{\beta}_1$ and $\hat{\beta}_2$, we ran a post hoc

simulation to infer a spatial arrangement that minimizes a population sum of

the leaf damage $\sum y_i = \beta_1 \sum x_i + \beta_2 \sum_{<i,j>} x_i x_j$. The constant intercept β_0,

the variance component u_i, and residual e_i were not considered because they

were not involved in deterministic dynamics of the model. Figure 7 shows

three representatives and a neutral expectation. For example, a mixture of a

dimorphism was expected to decrease the total leaf damage for a SNP at

chr2-14679190 near the *BBE16* locus ($\hat{\beta}_2 > 0$: Fig. 7a). On the other hand, a

clustered distribution of a dimorphism was expected to decrease the total

damage for a SNP at chr2-9422409 near the AT2G22170 locus encoding a

lipase/lipooxygenase PLAT/LH2 family protein ($\hat{\beta}_1 \approx 0, \hat{\beta}_2 < 0$: Fig. 7b).

p. 26

Furthermore, near monomorphism was expected to decrease the leaf

damage for a SNP at chr5-19121831 near the AT5G47075 and AT5G47077

loci encoding low-molecular cysteine-rich proteins, LCR20 and LCR6

($\hat{\beta}_1 > 0, \hat{\beta}_2 < 0$: Fig. 7c). If self and neighbor coefficients had no effects,

we observed a random distribution and no mitigation of damage i.e., $\sum y_i \approx$

0 (Fig. 7d). These post hoc simulations suggested a potential applicability of

neighbor GWAS in optimizing spatial arrangements in field cultivation.

DISCUSSION

Spatial and genetic factors underlying simulated phenotypes

Benchmark tests using simulated phenotypes revealed that appropriate

spatial scales could be estimated by variation partitioning of observed

phenotypes. When the scale of neighbor effects was narrow or moderate (α

= 1.0 or 3.0), the scale of the first nearest neighbors would be optimum for

the power to detect neighbor signals. In terms of neighbor effects in the

context of plant defense, mobile animals (e.g., mammalian browsers and

flying insects) often select a cluster of plant individuals (e.g., Bergvall et al.

2006; Hambäck et al. 2009; Sato and Kudoh 2015; Verschut et al. 2016). In

this case, neighbor effects could not be observed among individual plants

p. 27

within a cluster (Sato and Kudoh 2015). The exponential distance decay at α

= 0.25 represented situations in which the effective range of neighbor effects

was too broad to be detected; only in such situations should more than the

nearest neighbors be referred to gain the power to detect neighbor effects.

Neighbor GWAS could retain its power if neighbor effects were

spatially limited, and several major-effect genes governed a trait. In contrast,

when hundreds of causal variants involved a single trait and less than half of

phenotypic variation was attributable to neighbor effects, we observed a

decrease in the power to detect neighbor signals. In GWAS, false positive

rates can be reduced using linear mixed models that deal with kinship

structure as a random effect (Korte and Farlow 2013). Indeed, mixed models

were superior to standard linear models in this simulation. Our simulation

also adjusted the three variance components σ_1^2, σ_2^2, and σ_{12}^2, but their

relative contribution did not have significant effects on the power to detect

neighbor signals. In addition, the power to detect self-genotype signals

largely depended on the number of causal SNPs and PVE by major-effect

genes. This result suggests that the power for the standard GWAS to detect

the self-genotype effects was independent from neighbor effects.

Neighbor GWAS of the field herbivory on *Arabidopsis*

Our genetic analysis of neighbor effects is of ecological interest because

how host plant genotypes shape variation in plant–herbivore interactions is a

p. 28

long-standing question in population ecology (e.g., Karban 1992;

Underwood and Rausher 2000; Utsumi et al. 2011). Despite the low PVE

and several confounding factors under field conditions, the present study

illustrated the significant contribution of neighbor genotypic identity to

spatial variation in herbivory on *A. thaliana*. Surprisingly, the degree of leaf

damage variation explained by neighbor effects was greater than that by

self-genotype effects, although joint contribution of self and neighbor

effects was not doubled compared to the self-genotype effects. As we did

not find any significant peaks above the threshold of genome-wide multiple

testing, such neighbor effects on herbivory could be under polygenic control

and is exemplified by the simulation scenario where a number of variants

and strong distance decay caused moderate total phenotypic variation.

Regarding the self-genotype effects, we detected *GS-OX2*, a key

gene involved in glucosinolate biosynthesis, near the third top-scoring SNP

on the first chromosome. Aliphatic glucosinolates are well known as a

chemical defense against insect herbivory (van Poecke 2007; Brachi et al.

2015), where GS-OX2 catalyzes the conversion from methylthioalkyl to

corresponding methylsulfinylalkyl glucosinolates (Li et al. 2008).

Specifically, Geiselhardt et al. (2013) reported that *GS-OX2* was up-

regulated in response to feeding by larvae of the large white butterfly (*Pieris*

brassicae) and such an inducible defense was suppressed by prior

oviposition by adult butterflies. The present GWAS supported the relevance

p. 29

634   of *GS-OX2* to herbivory by *Pieris* butterflies, because we observed larvae of

635   *P. rapae* largely feeding on *A. thaliana*.

636   Remarkably, the second top-scoring SNP of the neighbor effects

was located near the *BBE16* locus responsive to methyl jasmonate. Methyl

jasmonate is well known as a volatile organic chemical that is emitted from

damaged tissue and elicits defense response of other plants (Reymond and

Farmer 1998; van Poecke 2007). Volatile-mediated plant–plant

communication is a key driver of neighbor effects in herbivory (e.g.,

Schuman et al. 2015; Erb 2018; Dahlin et al 2018), while a molecular

function of *BBE16* remains unexplored. More recent studies have shown

that berberine bridge enzyme-like proteins are involved in damage-

associated molecular patterns (Benedetti et al. 2018), which might be a

subject of future studies on inducible plant defense and neighbor effects.

Potential applicability and limitation

GWAS has often been applied to crop plants (Jannink et al. 2010; Hamblin

et al. 2011), where genotypes are known, and individuals are evenly

transplanted in space. Considering this outlook for agriculture, we provided

an example of neighbor GWAS across a lattice space in this study. However,

wild plant populations sometimes exhibit more complex spatial patterns

than expected by the Ising model (e.g., Kizaki and Katori 1999; Schlicht and

Iwasa 2004). In the rNeighborGWAS package, we allowed neighbor GWAS

p. 30

656   a continuous two-dimensional space. While its application has now been

657   limited to experimental populations, neighbor GWAS has potential

658   compatibility with an emerging discipline of landscape genomics (Bragg et

659   al. 2015). In this context, the additional R package could help future studies

660   to analyze neighbor effects using a wide variety of plant species.

661   Neighbor effects do not necessarily explain phenotypic variation

twice as much as conventional self-genotype effects, since some fractions of

PVE are not independent of self and neighbor effects, which is due to scale-

dependent collinearity between the elements K_1 and K_2 (e.g., $r = 0.60$ and

0.78 at $s = 1$ and 2 in the herbivory data analysis). Nevertheless, for each

explanatory variable, there was no strong correlation and the magnitude of

collinearity varied between the self-genotypic variable x_i and the neighbor

variable $\sum x_i x_j / L$ (mean ± SD among all SNPs, $r = -0.49 \pm 0.54$ and $-0.56 \pm$

0.63 at $s = 1$ and 2 in the herbivory data analysis). For an appropriate test of

the neighbor effects β_2 and its variance component σ_2^2, it is necessary to

compare the likelihood ratio between the standard and neighbor GWAS. In

such a case, PVE_{nei} is estimated by partial PVE_{nei} using a mixed model

including the two variance components into random effects. Given that the

collinearity becomes strong when s is large (see subsection "Neighbor

GWAS" in the method), we should be cautious when applying neighbor

GWAS to small sampling areas or long-range neighbor effects.

677   The present neighbor GWAS focused on single-locus effects and

p. 31

did not incorporate locus-by-locus interactions. Although it is difficult to

consider such an epistasis into GWAS (Gondro et al. 2013), it is possible

that multiple combinations among different variants govern neighbor

effects. For example, neighbor effects on insect herbivory may occur due to

the joint action of volatile-mediated signaling and the accumulation of

secondary metabolites (Dicke and Baldwin 2010; Erb 2018). The linear

mixed model could be extended as exemplified by the asymmetric neighbor

effects; however, we need to reconcile multiple criteria including the

collinearity of explanatory variables, inflation of p-values, and

computational costs. Further customization is warranted when analyzing

more complex forms of neighbor effects.

Conclusion

Based on the newly proposed methodology, we suggest that neighbor effects

are an overlooked source of phenotypic variation in field-grown plants. To

date, regional-scale interactions among plants have been analyzed using a

genome–environment association study of plant community composition

(Frachon et al. 2019), but fine-scale neighbor effects have yet to be

examined. Recent studies have shown that a single genomic region or a

single gene drives neighbor effects in plant growth, and such a genetic effect

shapes a positive relationship between plant genotype diversity and yield

p. 32

(Wuest and Niklaus 2018; McGale et al. 2020). Our newly proposed

methodology of neighbor GWAS could be an efficient tool to identify such a

key genetic variant responsible for neighbor effects and resulting

biodiversity effects.

Neighbor GWAS may also potentially help determine an optimal

spatial arrangement in plant cultivation, as suggested by the post hoc

simulation. The Ising model is well established in statistical physics

(McCoy and Maillard 2012) and is now applied to a machine-learning

pipeline that deals with high-dimensional genome data (Fisher and Mehta

2015). Genome-wide polymorphism data are useful not only to seek causal

genes in GWAS, but also to predict breeding values of crop plants in

genomic selection (e.g., Jannink et al. 2010; Hamblin et al. 2011; Yamamoto

et al. 2017). Given that the neighbor GWAS consists of a marker-based

regression, this methodology could also be implemented as a genomic

selection at a population level. Thus, our study provides an avenue for future

studies to predict population-level phenotypes in spatially structured

environments.

ACKNOWLEDGEMENTS

The authors would like to thank Ü. Seren, A. Korte, and M. Nordborg for

kindly providing the full imputed SNP data prior to being publicly available;

p. 33

T. Tsuchimatsu, K. Iwayama, and J. Bascompte for discussions; and

Dynacom Co., Ltd. for technical assistance with the R package

development. This study was supported by the Japan Science and

Technology Agency (JST) PRESTO (Grant number, JPMJPR17Q4 and

JPMJPR16Q9) to Y.S. and E.Y.; Japan Society for the Promotion of Science

(JSPS) Postdoctoral Fellowship (16J30005) to Y.S.; MEXT KAKENHI

(18H04785) and the Swiss National Science Foundation to K.K.S.; URPP

Global Change and Biodiversity of the University of Zurich to Y.S. and

K.S.S.; and JST CREST (JPMJCR15O2 and JPMJCR16O3) to A.J.N. and

K.K.S. The field experiment was supported by the Joint Usage/Research

Grant of Center for Ecological Research, Kyoto University, Japan.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA ARCHIVING

The leaf damage data on *A. thaliana* are included in the supporting

information (Table S1). The source codes used in this study are available at

the GitHub repository (https://github.com/naganolab/NeighborGWAS). R

package version of the neighbor GWAS method is available at CRAN

(https://cran.r-project.org/package=rNeighborGWAS).

REFERENCES

Alonso-Blanco C, Andrade J, Becker C, Bemm F, Bergelson J, Borgwardt

KM et al. (2016) 1,135 genomes reveal the global pattern of polymorphism

in *Arabidopsis thaliana*. *Cell* **166**: 481–491. doi:10.1016/j.cell.2016.05.063

Atwell S, Huang YS, Vilhjálmsson BJ, Willems G, Horton M, Li Y et al.

(2010) Genome-wide association study of 107 phenotypes in *Arabidopsis*

thaliana inbred lines. *Nature* **465**: 627–631. doi:10.1038/nature08800

Azaele S, Muneepeerakul R, Rinaldo A, Rodriguez-Iturbe I (2010) Inferring

plant ecosystem organization from species occurrences. *J Theor Biol* **262**:

323–329. doi:10.1016/j.jtbi.2009.09.026

Barbosa P, Hines J, Kaplan I, Martinson H, Szczepaniec A, Szendrei Z

(2009) Associational resistance and associational susceptibility: Having

right or wrong neighbors. *Ann Rev Ecol Evol Sys* **40**: 1–20.

doi:10.1146/annurev.ecolsys.110308.120242

Benedetti M, Verrascina I, Pontiggia D, Locci F, Mattei B, De Lorenzo G,

Cervone F (2018) Four *Arabidopsis* berberine bridge enzyme-like proteins

are specific oxidases that inactivate the elicitor-active oligogalacturonides.

Plant J **94**: 260–273. doi:10.1111/tpj.13852

Bergvall UA, Rautio P, Kesti K, Tuomi J, Leimar O (2006) Associational

effects of plant defences in relation to within- and between-patch food

choice by a mammalian herbivore: neighbour contrast susceptibility and

defence. *Oecologia* **147**: 253–260. doi:10.1007/s00442-005-0260-8

Brachi B, Meyer CG, Villoutreix R, Platt A, Morton TC, Roux F, Bergelson

J (2015) Coselected genes determine adaptive variation in herbivore

resistance throughout the native range of *Arabidopsis thaliana. Proc Natl*

Acad Sci USA **112**: 4032–4037. doi:10.1073/pnas.1421416112

Bragg JG, Supple MA, Andrew RL, Borevitz JO (2015) Genomic variation

across landscapes: insights and applications. *New Phytol* **207**: 953–967.

Browning BL, Browning SR (2009) A unified approach to genotype

imputation and haplotype-phase inference for large data sets of trios and

unrelated individuals. *Am J Hum Genet* **84**: 210–223.

doi:10.1016/j.ajhg.2009.01.005

Carlson M (2018) GO.db: A set of annotation maps describing the entire

Gene Ontology. R package version 3.7.0. doi:10.18129/B9.bioc.GO.db

Carrasco LR, Harwood TD, Toepfer S, MacLeod A, Levay N, Kiss J et al.

(2010) Dispersal kernels of the invasive alien western corn rootworm and

the effectiveness of buffer zones in eradication programmes in Europe. *Ann*

Appl Biol **156**: 63–77. doi:10.1111/j.1744-7348.2009.00363.x

Chan EKF, Rowe HC, Kliebenstein DJ (2010) Understanding the evolution

of defense metabolites in *Arabidopsis thaliana* using genome-wide

association mapping. *Genetics* **185**: 991–1007.

doi:10.1534/genetics.109.108522

Dahlin I, Rubene D, Glinwood R, Ninkovic V (2018) Pest suppression in

cultivar mixtures is influenced by neighbor-specific plant-plant

communication. *Ecol Appl* **28**: 2187–2196. doi:10.1002/eap.1807

Devaux C, Lavigne C, Austerlitz F, Klein EK (2007) Modelling and

estimating pollen movement in oilseed rape (*Brassica napus*) at the

landscape scale using genetic markers. *Mol Ecol* **16**: 487–499.

doi:10.1111/j.1365-294X.2006.03155.x

Devoto A, Ellis C, Magusin A, Chang HS, Chilcott C, Zhu T, Turner JG

(2005) Expression profiling reveals *COI1* to be a key regulator of genes

involved in wound- and methyl jasmonate-induced secondary metabolism,

defence, and hormone interactions. *Plant Mol Biol* **58**: 497–513.

doi:10.1007/s11103-005-7306-5

Dicke M, Baldwin IT (2010) The evolutionary context for herbivore-

induced plant volatiles: beyond the 'cry for help'. *Trends Plant Sci* **15**: 167–

175. doi:10.1016/j.tplants.2009.12.002

Erb M (2018) Volatiles as inducers and suppressors of plant defense and

immunity - origins, specificity, perception and signaling. *Curr Opin Plant*

Biol **44**: 117–121. doi:10.1016/j.pbi.2018.03.008

Frachon L, Mayjonade B, Bartoli C, Hautekèete NC, Roux F (2019)

Adaptation to plant communities across the genome of *Arabidopsis*

thaliana. *Mol Biol Evol* **36**: 1442–1456. doi:10.1093/molbev/msz078

Fisher CK, Mehta P (2015) Bayesian feature selection for high-dimensional

linear regression via the Ising approximation with applications to genomics.

Bioinformatics **31**: 1754–1761. doi:×

Gage JL, De Leon N, Clayton MK (2018) Comparing genome-wide

association study results from different measurements of an underlying

phenotype. *G3: Genes, Genomes, Genetics* **8**: 3715–3722.

doi:10.1534/g3.118.200700

Geiselhardt S, Yoneya K, Blenn B, Drechsler N, Gershenzon J, Kunze R,

Hilker M (2013) Egg laying of cabbage white butterfly (*Pieris brassicae*) on

Arabidopsis thaliana affects subsequent performance of the larvae. *PLoS*

ONE **8**: e59661. doi:10.1371/journal.pone.0059661

Gondro C, van der Werf J, Hayes B (eds.) (2013) Genome-wide association

studies and genomic prediction. *Methods Mol Biol*. Humana Press, New

York. doi:10.1007/978-1-62703-447-0

Hambäck PA, Björkman M, Rämert B, Hopkins RJ (2009). Scale-dependent

responses in cabbage herbivores affect attack rates in spatially

heterogeneous systems. *Basic Appl Ecol* **10**: 228–236.

doi:10.1016/j.baae.2008.06.004

Hambäck PA, Inouye BD, Andersson P, Underwood N (2014) Effects of

plant neighborhoods on plant-herbivore interactions: resource dilution and

associational effects. *Ecology* **95**: 1370–1383. doi:10.1890/13-0793.1

Hamblin MT, Buckler ES, Jannink JL (2011) Population genetics of

p. 38

genomics-based crop improvement methods. *Trends Genet* **27**: 98–106.

doi:10.1016/j.tig.2010.12.003

Hauser MT, Harr B, Schlötterer C (2001) Trichome distribution in

Arabidopsis thaliana and its close relative *Arabidopsis lyrata*: molecular

analysis of the candidate gene *GLABROUS1*. *Mol Biol Evol* **18**: 1754–1763.

doi:10.1093/oxfordjournals.molbev.a003963

Horton MW, Hancock AM, Huang YS, Toomajian C, Atwell S, Auton A et

al. (2012) Genome-wide patterns of genetic variation in worldwide

Arabidopsis thaliana accessions from the RegMap panel. *Nat Genet* **44**:

212–216. doi:10.1038/ng.1042

Horton MW, Bodenhausen N, Beilsmith K, Meng D, Muegge BD,

Subramanian S et al. (2014) Genome-wide association study of *Arabidopsis*

thaliana leaf microbial community. *Nat Commun* **5**: 5320.

doi:10.1038/ncomms6320

Ida TY, Takanashi K, Tamura M, Ozawa R, Nakashima Y, Ohgushi T (2018)

Defensive chemicals of neighboring plants limit visits of herbivorous

insects: Associational resistance within a plant population. *Ecol Evol* **8**:

12981–12990. doi:10.1002/ece3.4750

Ising E (1925) Beitrag zur theorie des ferromagnetismus. *Zeitschrift für*

Physik **31**: 253–258.

Jannink JL, Lorenz AJ, Iwata H (2010) Genomic selection in plant breeding:

from theory to practice. *Brief Funct Genomic* **9**: 166–177.

p. 39

doi:10.1093/bfgp/elq001

Kang HM, Zaitlen NA, Wade CM, Kirby A, Heckerman D, Daly MJ, Eskin

E (2008) Efficient control of population structure in model organism

association mapping. *Genetics* **178**: 1709–1723.

doi:10.1534/genetics.107.080101

Karban R (1992) Plant variation: its effects on populations of herbivorous

insects. In: Fritz RS, Simms EL (eds) *Plant resistance to herbivores and*

pathogens: ecology, evolution, and genetics, University of Chicago Press:

Chicago, pp. 195–215.

Kizaki S, Katori M (1999) Analysis of canopy-gap structures of forests by

Ising-Gibbs states-equilibrium and scaling property of real forests. *J Phys*

Soc Japan **68**: 2553-2560. doi:10.1143/JPSJ.68.2553

Kofler R, Schlotterer C (2012) Gowinda: unbiased analysis of gene set

enrichment for genome-wide association studies. *Bioinformatics* **28**: 2084–

2085. doi:10.1093/bioinformatics/bts315

Korte A, Farlow A (2013) The advantages and limitations of trait analysis

with GWAS: a review. *Plant Methods* **9**: 29. doi:10.1186/1746-4811-9-29

Li J, Hansen BG, Ober JA, Kliebenstein DJ, Halkier BA (2008) Subclade of

flavin-monooxygenases involved in aliphatic glucosinolate biosynthesis.

Plant Physiol **148**: 1721–1733. doi:10.1104/pp.108.125757

McCoy BM, Maillard JM (2012) The importance of the Ising model.

Progress *Theor Phys* **127**: 791-817. doi:10.1143/PTP.127.791

p. 40

McGale E, Valim H, Mittal D, Jimenez JM, Halitschke R, Schuman MC,

Baldwin IT (2020) Determining the scale at which variation in a single gene

changes population yields. *eLife* **9**. doi:10.7554/eLife.53517

Mundt CC (2002) Use of multiline cultivars and cultivar mixtures for

disease management. *Ann Rev Phytopathol* **40**: 381-410.

doi:10.1146/annurev.phyto.40.011402.113723

Nallu S, Hill JA, Don K, Sahagun C, Zhang W, Meslin C et al. (2018) The

molecular genetic basis of herbivory between butterflies and their host

plants. *Nat Ecol Evol* **2**: 1418-1427. doi:10.1038/s41559-018-0629-9

Perdry H, Dandine-Roulland C (2018) gaston: Genetic Data Handling (QC,

GRM, LD, PCA) & Linear Mixed Models. R package version 1.5.4.

https://CRAN.R-project.org/package=gaston

R Core Team (2019) R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria.

https://www.R-project.org/

Reymond P, Farmer EE (1998) Jasmonate and salicylate as global signals

for defense gene expression. *Curr Opin Plant Biol* **1**: 404–411.

doi:10.1016/s1369-5266(98)80264-1

Rieux A, Soubeyrand S, Bonnot F, Klein EK, Ngando JE, Mehl A et al.

(2014) Long-distance wind-dispersal of spores in a fungal plant pathogen:

estimation of anisotropic dispersal kernels from an extensive field

experiment. *PLoS ONE* **9**: e103225. doi:10.1371/journal.pone.0103225

Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC, Muller M

(2011) pROC: an open-source package for R and S+ to analyze and compare

ROC curves. *BMC Bioinformatics* **12**: 77. doi:10.1186/1471-2105-12-77

Sato Y (2018) Associational effects and the maintenance of polymorphism

in plant defense against herbivores: review and evidence. *Plant Species Biol*

33: 91–108. doi:10.1111/1442-1984.12201

Sato Y, Kudoh H (2015) Tests of associational defence provided by hairy

plants for glabrous plants of *Arabidopsis halleri* subsp. *gemmifera* against

insect herbivores. *Ecol Entomol* **40**: 269–279. doi:10.1111/een.12179

Sato Y, Kudoh H (2017) Herbivore-mediated interaction promotes the

maintenance of trichome dimorphism through negative frequency-

dependent selection. *Am Nat* **190**: E67–E77. doi:10.1086/692603

Sato Y, Ito K, Kudoh H (2017) Optimal foraging by herbivores maintains

polymorphism in defence in a natural plant population. *Funct Ecol* **31**:

2233–2243. doi:10.1111/1365-2435.12937

Sato Y, Shimizu-Inatsugi R, Yamazaki M, Shimizu KK, Nagano AJ (2019)

Plant trichomes and a single gene *GLABRA1* contribute to insect community

composition on field-grown *Arabidopsis thaliana*. *BMC Plant Biol* **19**: 163.

doi:10.1186/s12870-019-1705-2

Schlicht R, Iwasa Y. (2004) Forest gap dynamics and the Ising model. *J*

Theoret Biol **230**: 65–75. doi:10.1016/j.jtbi.2004.04.027

Schuman MC, Allmann S, Baldwin IT (2015) Plant defense phenotypes

p. 42

determine the consequences of volatile emission for individuals and

neighbors. *eLife* **4**: e04490. doi:10.7554/eLife.04490

Tahvanainen JO, Root RB (1972) The influence of vegetational diversity on

the population ecology of a specialized herbivore, *Phyllotreta cruciferae*

(Coleoptera: Chrysomelidae). *Oecologia* **10**: 321–346.

doi:10.1007/BF00345736

Togninalli M, Seren Ü, Meng D, Fitz J, Nordborg M, Weigel D et al. (2018)

The AraGWAS Catalog: a curated and standardized *Arabidopsis thaliana*

GWAS catalog. *Nucleic Acids Res* **46**: D1150–D1156.

doi:10.1093/nar/gkx954

Underwood N, Rausher MD (2000) The effects of host-plant genotype on

herbivore population dynamics. *Ecology* **81**: 1565-1576. doi:10.1890/0012-

9658(2000)081[1565:TEOHPG]2.0.CO;2

Underwood N, Inouye BD, Hambäck PA (2014) A conceptual framework

for associational effects: when do neighbors matter and how would we

know? *Quart Rev Biol* **89**: 1–19. doi:10.1086/674991

Underwood N, Hambäck PA, Inouye BD (2020) Pollinators, herbivores, and

plant neighborhood effects. *Quart Review Biol* **95**: 37–57.

doi:10.1086/707863

Utsumi S, Ando Y, Craig TP, Ohgushi T (2011) Plant genotypic diversity

increases population size of a herbivorous insect. *Proc Royal Soc B* **278**:

3108–3115. doi:10.1098/rspb.2011.0239

p. 43

van Poecke RM (2007) Arabidopsis-insect interactions. *The Arabidopsis*

Book **5**: e0107. doi:10.1199/tab.0107

Verschut TA, Becher PG, Anderson P, Hambäck PA (2016) Disentangling

associational effects: both resource density and resource frequency affect

search behaviour in complex environments. *Funct Ecol* **30**: 1826-1833.

doi:10.1111/1365-2435.12670

Wang M, Roux F, Bartoli C, Huard-Chauveau C, Meyer C, Lee H et al.

(2018) Two-way mixed-effects methods for joint association analysis using

both host and pathogen genomes. *Proc Natl Acad Sci USA* **115**: E5440–

E5449. doi:10.1073/pnas.1710980115

Weiner J (1990) Asymmetric competition in plant populations. *Trends Ecol*

Evol **5**: 360-364. doi:10.1016/0169-5347(90)90095-U

Wuest SE, Niklaus PA (2018) A plant biodiversity effect resolved to a single

chromosomal region. *Nat Ecol Evol* **2**: 1933–1939. doi:10.1038/s41559-

018-0708-y

Yamamoto E, Matsunaga H, Onogi A, Ohyama A, Miyatake K, Yamaguchi

H et al. (2017) Efficiency of genomic selection for breeding population

design and phenotype prediction in tomato. *Heredity* **118**: 202–209.

doi:10.1038/hdy.2016.84

Zeller SL, Kalinina O, Flynn DFB, Schmid B (2012) Mixtures of

genetically modified wheat lines outperform monocultures. *Ecol Appl* **22**:

1817–1826. doi:10.1890/11-0876.1

Zhou X, Stephens M (2012) Genome-wide efficient mixed-model analysis

for association studies. *Nat Genet* **44**: 821–824.

TABLES & FIGURES

Table 1. Factors affecting the power to detect major-effect genes in simulated phenotypes. ANOVA tables show the degree of freedom (df), sum of squares (SS), F-statistics, and p-values. Explanatory variables are the number of causal SNPs, proportion of phenotypic variation explained (PVE) by major-effect genes (PVE_β), total PVE by major-effect genes and variance components ($\text{PVE}_\beta + \text{PVE}_u$), ratio of variance component parameters ($\sigma_1^2 : \sigma_2^2 : \sigma_{12}^2$), and distance decay coefficient α. For the neighbor effects, the response variable was the maximum Area Under the ROC Curve (AUC) of the spatial scales from $s = 1$ to $s = 3$. The AUC was arcsine square-root transformed to improve normality. NA means not available.

Response	Factors	df	SS	F	p-value
AUC_{self}	No. of causal SNPs	1	37.70	2840.2	< 2e-16
	PVE_β	1	2.23	168.2	< 2e-16
	$\text{PVE}_\beta + \text{PVE}_u$	1	0.04	2.64	0.104
	$\sigma_1^2 : \sigma_2^2 : \sigma_{12}^2$	2	0.01	0.30	0.738
	α	1	0.00	0.01	0.942
	Residuals	1073	14.24	*NA*	*NA*
AUC_{nei}	No. of causal SNPs	1	30.70	1500.4	< 2e-16
	PVE_β	1	3.01	147.3	< 2e-16
	$\text{PVE}_\beta + \text{PVE}_u$	1	0.12	5.65	0.01764
	$\sigma_1^2 : \sigma_2^2 : \sigma_{12}^2$	2	0.00	0.08	0.925
	α	1	1.87	91.25	< 2e-16
	Residuals	1073	21.96	*NA*	*NA*

p. 46

(a) $\beta_2 = 0.2$, $\beta_1 = 0.0$ **(b)** $\beta_2 = -0.2$, $\beta_1 = 0.0$ **(c)** $\beta_2 = -0.2$, $\beta_1 = 0.05$

2-D Ising model   *post hoc simulation*

$$\sum y_i = \beta_1 \sum x_i + \beta_2 \sum_{<i,j>} x_i x_j$$

$$y_i = \beta_0 + \beta_1 x_i + \frac{\beta_2 \sum_{<i,j>}^{L} x_i x_j^{(s)}}{L} + u_i + e_i$$

inverse problem   **Neighbor GWAS**
                                               (linear mixed model)

Figure 1. Relationship between Neighbor GWAS and Ising model. Upper

panels show spatial arrangements expected by a 2-D Ising model $\sum y_i =$

$\beta_1 \sum x_i + \beta_2 \sum_{<i,j>} x_i x_j$. (a) If $\beta_2 > 0$, mixed patterns give the argument of the

minimum for a population sum of phenotype values $\sum y_i$. (b) If $\beta_2 < 0$,

clustered patterns give the argument of the minimum for $\sum y_i$. (c) In

addition, β_1 determines overall patterns favoring -1 or +1 states. The figure

shows outcomes from a random 100×100 lattice after 1000 iterations of

simulated annealing. Conversely, the neighbor GWAS was implemented as

an inverse problem of the 2-D Ising model, where genotypes and its spatial

arrangement, x_i and $x_i x_j$, were given while the coefficients β_1 and β_2 were

to be estimated from observed phenotypes y_i. In addition, the variance

component due to self and neighbor effects was considered a random effect

p. 47

995   in a linear mixed model, such that $u_i \sim \text{Norm}(0, \sigma_1^2 \mathbf{K}_1 + \sigma_2^2 \mathbf{K}_2)$.

996   Once β_1 and β_2 were determined, we could simulate a genotype distribution

997   that maximizes or minimizes $\sum y_i$.

Figure 2. Experimental setting in the *Arabidopsis* herbivory data. (a)

Photograph of the field site. Each 0.6 × 2.6 m block included a replicate of

199 accessions and an additional Col-0. (b) *Arabidopsis thaliana* plants

were arranged in a 10 × 40 cell plastic tray. The plants were placed in a

checkered pattern so that alternate cells were left empty; therefore, each of

the 40 rows contained 5 plants and each of the 10 columns contained 20

plants. Yellow lines represent s-th neighbor scales from a focal i-th plant. (c)

A graphical explanation of the experimental area. A meadow (green) was

separately covered with weed-masking sheets (grey).

(a) Self, β_1

(b) Neighbor, β_2

Figure 3. Receiver operating characteristic (ROC) curves representing the

sensitivity (true positive rate) and the specificity (1 - false positive rate) in

simulated phenotypes. The upper and lower rows give the results of self and

neighbor effects with the number of causal SNPs at 10, 50, and 300. The

shade of the lines indicates the proportion of phenotypic variation explained

by major-effect genes as $PVE_\beta = 0.1$ (light gray), 0.3 (dark gray), or 0.6

(black). The thickness of the lines indicates distance decay coefficients as α

= 0.25 (thin), 1 (medium), or 3 (thick). Entire results are summarized using

the area under the ROC curve (AUC) in Figure S1.

(a) $\alpha = 0.25$ **(b)** $\alpha = 1.0$ **(c)** $\alpha = 3.0$

scale, s

1020

Figure 4. Scale dependence of neighbor effects on simulated phenotypes

among all iterations. The broad (a), intermediate (b), and narrow (c)

effective range of neighbor effects are represented by weak, moderate, and

strong distance decay, respectively. The proportion of phenotypic variation

explained by neighbor effects (partial PVE_{nei}) and the area under the ROC

curve (AUC) of neighbor effects are shown along the spatial scale from the

p. 51

1027   first nearest ($s = 1$) to the third nearest ($s = 3$) neighbors. Boxplots show

1028   center line: median, box limits: upper and lower quartiles, whiskers: 1.5×

1029   interquartile range, and points: outliers.

1030

1031

Figure 5. Proportion of phenotypic variation explained (PVE) by self-

genotype or neighbor effects on the leaf damage score of field-grown

Arabidopsis thaliana. The partial or single PVE is shown at the spatial scale

of $s = 0$ (without any neighbor effects), $s = 1$ and $s = 2$, as defined by the

partial $\text{PVE}_{\text{self}} = \sigma_1^2/(\sigma_1^2 + \sigma_2^2 + \sigma_e^2)$; partial $\text{PVE}_{\text{nei}} = \sigma_2^2/(\sigma_1^2 + \sigma_2^2 + $

$\sigma_e^2)$; single $\text{PVE}_{\text{self}} = \sigma_1^2/(\sigma_1^2 + \sigma_e^2)$; and single $\text{PVE}_{\text{nei}} = \sigma_2^2/(\sigma_2^2 + \sigma_e^2)$.

The red or blue fraction indicates PVE by self or neighbor effects,

respectively. The white fraction indicates residuals. Asterisks highlight a

significant fraction with stepwise likelihood ratio tests from simpler to

complex models: [**]p-value < 0.01: [***]p-value < 0.001.

1042

1043

p. 53

1044

Figure 6. Manhattan plots for the self (a) and neighbor (b) effects on the

leaf damage score of field-grown *Arabidopsis thaliana*. The first to fifth

chromosomes of *A. thaliana* are differently colored, where lighter plots

indicate smaller MAF. Horizontal dashed lines indicate the threshold after

Bonferroni correction at *p*-value < 0.05. The red vertical line in panel (a)

indicates a SNP position near the *GS-OX2* locus, while three circles

highlighted by a black outline in panel (b) indicates the variants subject to

the post hoc simulation (Fig. 7). Results of the self and neighbor effects are

shown at $s = 0$ (i.e., standard GWAS) and $s = 1$, respectively.

1054

(a) Chr 2, Position 14679190: β_1 = 0.19, β_2 = 0.31, Σy_i = -237.2

(b) Chr 2, Position 9422409: β_1 = -0.001, β_2 = -0.18, Σy_i = -296.2

(c) Chr 5, Position 19121831: β_1 = 0.12, β_2 = -0.22, Σy_i = -593.8

(d) No effects: β_1 = 10^{-6}, β_2 = 10^{-6}, Σy_i = 10^{-4}

Figure 7. Post hoc simulations exemplifying a spatial arrangement of two alleles expected by the estimated self and neighbor effects, $\hat{\beta}_1$ and $\hat{\beta}_2$, on the leaf damage score of *Arabidopsis thaliana*. Population sum of the leaf damage $\sum y_i = \beta_1 \sum x_i + \beta_2 \sum_{<i,j>} x_i x_j$ was minimized using 1000 iterations of simulated annealing from a random distribution of two alleles in a 10 × 40 space.