Nick-seq for single-nucleotide resolution genomic maps of DNA modifications and damage

Bo Cao[1,2,3,4,†,*], Xiaolin Wu[2,3,5,†], Jieliang Zhou[6], Hang Wu[2,7], Michael S. DeMott[2,8], Chen Gu[2,9],

Lianrong Wang[5], Delin You[4,*], Peter C. Dedon[2,3,8,*]

[1] College of Life Sciences, Qufu Normal University, Qufu, Shandong, 273165, China

[2] Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, MA

02139, USA

[3] Singapore-MIT Alliance for Research and Technology, Antimicrobial Drug Resistance

Interdisciplinary Research Group, Singapore 138602, Singapore

[4] State Key Laboratory of Microbial Metabolism, Joint International Research Laboratory of

Metabolic and Developmental Sciences, and School of Life Sciences & Biotechnology, Shanghai

Jiao Tong University, Shanghai, 200030, China

[5] Key Laboratory of Combinatorial Biosynthesis and Drug Discovery, Ministry of Education and

School of Pharmaceutical Sciences, Wuhan University, Wuhan, Hubei, 430071, China

[6] KK Research Center, KK Women's and Children's Hospital, 229899, Singapore

[7] School of Life Sciences, Anhui University, Hefei, Anhui, 230601, China

[8] Center for Environmental Health Sciences, Massachusetts Institute of Technology, Cambridge,

MA 02139, USA

[9] Current affiliation: Merck Research Laboratories, Merck & Co., Inc., Boston, MA 02115, USA

[†] These authors contributed equally to this work.

Corresponding authors: B.C. (ORCID ID: 0000-0002-7011-1676) (caobo@qfnu.edu.cn), D.Y.

(dlyou@sjtu.edu.cn), and P.C.D. (ORCID ID: 0000-0003-0011-3067) (pcdedon@mit.edu)

Abstract

Here we present the Nick-seq platform for quantitative mapping of DNA modifications and

damage at single-nucleotide resolution across genomes. Pre-existing breaks are blocked and

DNA structures converted to strand-breaks for 3'-extension by nick-translation to produce

nuclease-resistant oligonucleotides, and 3'-capture by terminal transferase tailing. Libraries from

both products are subjected to next-generation sequencing. Nick-seq is a generally applicable

method illustrated with quantitative profiling of single-strand-breaks, phosphorothioate

modifications, and DNA oxidation.

Main text (1,000~1,500 word including figure legend)

Genomic mapping of specific DNA modifications[1] and damage[2] can be achieved with methods

such as bisulfite sequencing for 5-methylcytidine[3], chromatin immunoprecipitation (ChIP)

coupled with next-generation sequencing (NGS)[4,5], and single-molecule real-time (SMRT)[6] and

nanopore[7] sequencing. However, all are limited to specific modifications or suffer from low

sensitivity and specificity. Here we describe Nick-seq for highly sensitive quantitative genomic

mapping of any type of DNA modification or damage that can be converted to a strand-break. As

shown in **Figure 1a**, purified genomic DNA is subjected to sequencing-compatible fragmentation

and the resulting 3'-OH ends are blocked with dideoxyNTPs. The DNA modification is then

converted to a strand-break by enzyme or chemical treatment, followed by capture of the 3'- and

5'-ends of resulting strand-breaks using two complementary strategies. One portion of DNA is

subjected to nick translation (NT) with α-thio-dNTPs to generate 100-200 nt phosphorothioate-

containing oligonucleotides that are resistant to subsequent hydrolysis of the bulk of the genomic

DNA by exonuclease III and RecJ$_f$. The purified PT-protected fragment is used to generate an

NGS library with the modification of interest positioned at the 5'-end of the PT-labeled fragment.

A second portion of the same DNA sample is used for terminal transferase (TdT)-dependent

poly(dT) tailing of the 3'-end of the strand-break, with the tail used to create a sequencing library

by reverse transcriptase template switching[8]. Subsequent NGS positions the modification of

interest 5'-end of the poly(dT) tail.

The workflow for sequencing data processing (**Fig. 1b**) uses the NT-derived reads as the

primary dataset for developing a rough modification map, with TdT-derived reads as

complementary corrective data. This hybrid approach exploits the fact that NT is agnostic to the

base identify at the damage site but generates a high background of false positive sites, while TdT

cannot be used with modifications occurring at dT due to loss of the poly(dT) tail during data

analysis. The TdT reads are used to correct NT false-positive reads. For example, if NT maps a

strand-break at T 1000 in the genome, then TdT reads are examined for a strand-break at position

999 and 1001. This 1 position shift accommodates poly(dT) tail removal during data processing

and validates the NT map. If TdT does not call a strand-break at 999 or 1001, then the NT result

is considered a false positive. In other cases, if the NT-detected site occurs at a G, C, or A, then

TdT valdiates the same site. The use of both methods increases the sensitivity and specificity of

the resulting map.

We validated Nick-seq by mapping DNA single-strand-breaks caused by the site-specific

endonuclease, Nb.BsmI, at the 2,681 G/CATTC motifs in the *E.coli* genome. Purified DNA was

treated with Nb.BsmI and the Nick-seq-processed library sequenced using the Illumina NextSeq

platform with an average of 10^7 raw sequencing reads for each sample (**Fig. 2b**). Paired-end

sequencing confirmed that >80% of reads uniquely aligned to the *E.coli* genome (**Supplementary**

Table 1). For subequent reads enrichment (**Fig. 1b**), we calculated position-wise coverage values

using the 5'-end of sequencing reads (NT read 1, TdT read 2) and defined Nick-seq peaks as

having >5 reads and 2-times more reads than sites located one-nucleotide up- and down-stream.

We then calculated the coverage ratio of the peaks to corresponding sites in an untreated DNA

control. To identify the optimal minimal coverage ratio, we varied the ratio and calculated the

number of identified sites at each ratio value (**Fig. 2b**). As the coverage ratio increased from 2 to

7, the number of identified sites decreased from 92% to 59% of 2,681 expected ("sensitivity"),

while the accuracy (identified sites/expected sites; "specificity") only increased from 98% to

99.5%. To maximize sensitivity, we chose a coverage ratio of 2, which allowed identification of

2,462 (97.5%) of the predicted Nb.BsmI sites. Another 1% of called sites (27) occurred in

sequences differing from the consensus by one nucleotide (**Supplementary Table 2**). These sites

showed lower average sequencing coverage (75 vs 1318) and likely represent Nb.BsmI "star"

activity. Another validation experiment with Nb.BsrDI, which cuts at N/CATTGC, showed no

evident 3'-end sequence bias for DNA break site detection (**Supplementary Table 3)**. Thus, Nick-

seq showed high accuracy and sensitivity for single-nucleotide genomic mapping of DNA strand-

breaks.

The validated Nick-seq was applied to map the naturally-occurring phosphorothioate (PT)

DNA modifications in *Salmonella enterica* serovar Cerro 87, using iodine to oxidize PTs to

produce DNA strand-breaks[9] (**Fig. 2c**). We previously established by SMRT sequencing that PTs

occurred as bistranded modifications at 10-15% of the GAAC/GTTC motifs in *S. enterica*[9]. Nick-

seq recognized 12,239 PT sites (**Fig. 2c, Supplementary Table 4**), of which 11,684 (96%)

occurred at $G_{PS}AAC/G_{PS}TTC$, with 8,568 (73%) modified on both strands and 27% modified on

one strand (**Fig. 2c**). This agrees with our previous observations using an orthogonal sequencing

method[9]. In addition to GAAC/GTTC motifs, Nick-seq also revealed less abundant PTs at $G_{PS}TAC$

(168), $G_{PS}ATC$ (30), and $G_{PS}AAAC$ or $G_{PS}AAAAC$ (19), with half of $G_{PS}TAC$ and $G_{PS}ATC$ sites

modified on only one strand (**Supplementary Table 4**). These results indicate that Nick-seq has a

higher sensitivity to detect rare PT modifications than other methods.[9, 10]

Finally, we applied Nick-seq to DNA modifications not previously subjected to genomic

mapping: oxiatively-induced abasic sites. Apurinic and apyrimidinic (AP) sites represent a

prevalent and toxic form of DNA damage that blocks DNA replication and transcription[11, 12]. AP

sites arise as intermediates in base excision DNA repair, in which damaged bases are excised by

glycosylases and the resulting AP sites cleaved by AP endonucleases[13]. AP sites can also arise

by oxidation of DNA on both the nucleobase and 2'-deoxyribose moietities[14, 15], as well as by

demethylation of m^5C epigenetic marks.[16] In spite of the importance of AP sites, little is known

about their formation, persistence, and distribution in genomic DNA. Here we used Nick-seq to

profile AP sites in *E. coli* exposed to H_2O_2 at a non-lethal dose of 0.2 mM (LD_{50} ~5 mM) (**Fig.**

2d). Following DNA purification, AP sites were expressed as strand-breaks using endonuclease

IV (EndoIV), which cleaves both native and oxidized AP sites[14, 15]. Nick-seq identified 1,519

EndoIV-sensitive sites, as well as 82 sites in the endogenous plasmid, with an unexposed control

showing 11 and 8 sites, respectively (**Fig. 2d, Supplementary Tables 5, 6**). Considering the

nucleobase precursor of the AP site, there was a weak preference for thymine (33%) followed by

adenine (25%), cytosine (24%), and guanine (18%), with a similar distribution in the plasmid

(**Supplementary Tables 5, 6**). This suggests that H_2O_2-derived DNA oxidizing agents either do

not selectively oxidize guanine as predicted[17] or that the predominant form of damage is DNA

sugar oxidation. However, there was a more pronounced sequence context effect. Analysis of 15

bp up- and down-stream of the AP sites revealed a strong preference for cytosine (47%) at -1

relative to the AP sites. The distribution of AP sites on the plasmid was also non-random (**Fig.**

2d), with clustering in three regions related to DNA replication and transcription: the F1 origin,

pUC origin, and Amp^R gene (**Supplementary Table 7**). AP site clustering near DNA replication

sites was observed previously by immunostaining[18], suggesting that the transcriptionally active

and single-strand DNA are vulnerable to oxidatively-induced AP sites. We tested this by analyzing

the distribution of AP sites in the *E. coli* genome relative to origins of replication (OriC), coding

sequences, and non-coding sequences. While there was an average of 0.32 Aps/kbp (1,519 AP per

4,686,000 bp), the 20 kb region around OriC showed 0.70 Aps/kbp. Nick-seq also revealed 1401

APs in the 4.1×10^6 bp coding sequence region (0.34 AP/kbp), and 118 APs (0.20 AP/kbp) in the

0.58×10^6 bp non-coding region. These results suggest a preference for AP sites in DNA

undergoing replication or transcription during H_2O_2 stress.

Nick-seq thus provides an efficient, label-free approach to quantive mapping of DNA

damage and modifications in genomes, with applications in DNA damage and repair, epigenetics,

restriction-modification systems, and DNA metabolism.

Methods – see the online Methods Section

Acknowledgements

The authors thank the MIT BioMicro Center, MIT Center for Environmental Health Science,

Singapore-MIT Alliance for Research and Technology (SMART) for use of their facilities, and

funding support from the National Natural Science Foundation of China (31630002), National

Science Foundation of the USA (CHE-1709364), National Research Foundation of Singapore

through the SMART Infectious Disease and Antimicrobial IRGs, National Institute of

Environmental Health Sciences (P30-ES002109), and Fundamental Research Funds for the

Central Universities of China (2015306020202). X.W. was supported by a fellowship from the

China Scholarship Council (201606270163).

Author Contributions

P.C.D. and B.C. designed Nick-seq. B.C., X.W., H.W. performed Nick-seq experiments. P.C.D,

B.C., X.W., and J.Z. constructed the bioinformatics pipeline. M.S.D., C.G., D.Y., and L.W.,

contributed vital reagents. All authors discussed the results and contributed to the final manuscript.

Competing Interests Statement

B.C., M.S.D., and P.C.D are co-inventors on a PCT patent (PCT/US2019/013714) and US Patent

(US 2019/0284624 A1) relating to the published work.

Data Availability

Sequencing data has been deposited in NCBI GEO database under accession numbers GSE138070,

GSE138173, and GSE138476.

Software and Code Availability

Custom scripts for processing the sequencing data are described in Methods and are available at

https://github.com/BoCao2019/Nick-seq: .gitignore,   NT_negative_strand.R,

NT_positive_strand.R,   TdT_negative+NT_positive.R,   TdT_negative_strand.R,

TdT_positive+NT_negative.R, and TdT_positive_strand.R.

References

1.   Chen, Y. et al. *Chem Soc Rev* **46**, 2844-2872 (2017).

2.   Roos, W.P., Thomas, A.D. & Kaina, B. *Nat Rev Cancer* **16**, 20-33 (2016).

3.   Li, Q., Hermanson, P.J. & Springer, N.M. *Methods Mol Biol* **1676**, 185-196 (2018).

4.   Hu, J., Selby, C.P., Adar, S., Adebali, O. & Sancar, A. *J Biol Chem* (2017).

5.   Ding, Y., Fleming, A.M. & Burrows, C.J. *J Am Chem Soc* **139**, 2569-2572 (2017).

6.   Clark, T.A., Spittle, K.E., Turner, S.W. & Korlach, J. *Genome Integr* **2**, 10 (2011).

7.   Schibel, A.E. et al. *J Am Chem Soc* **132**, 17992-17995 (2010).

8.   Zhu, Y.Y., Machleder, E.M., Chenchik, A., Li, R. & Siebert, P.D. *Biotechniques* **30**, 892-
173   897 (2001).

174   9.   Cao, B. et al. *Nat Commun* **5** (2014).

175   10.   Li, J. et al. *PLoS Genet* **15**, e1008026 (2019).

176   11.   Clauson, C.L., Oestreich, K.J., Austin, J.W. & Doetsch, P.W. *Proc Natl Acad Sci U S A*
177   **107**, 3657-3662 (2010).

178   12.   Galhardo, R.S., Almeida, C.E., Leitao, A.C. & Cabral-Neto, J.B. *J Bacteriol* **182**, 1964-
179   1968 (2000).

180   13.   Dianov, G.L., Sleeth, K.M., Dianova, II & Allinson, S.L. *Mutat Res* **531**, 157-163 (2003).

181   14.   Xu, Y.J., Kim, E.Y. & Demple, B. *J Biol Chem* **273**, 28837-28844 (1998).

182   15.   Greenberg, M.M., Weledji, Y.N., Kim, J. & Bales, B.C. *Biochemistry* **43**, 8178-8183
183   (2004).

16.   Wu, X. & Zhang, Y. *Nat Rev Genet* **18**, 517-534 (2017).

17.   Dedon, P.C. & Tannenbaum, S.R. *Arch Biochem Biophys* **423**, 12-22 (2004).

18.   Chastain II, P.D. et al. *FASEB J* **24**, 3674-3680 (2010).

Figure Legends

Figure. 1. Nick-seq library preparation (a) and data analysis workflow (b).

Figure. 2. **Nick-seq validation and application**. (**a**) Mapping single-strand breaks produced by

Nb.BsmI in *E. coli* genomic DNA. Middle panel: representative view of sequencing reads

distirbuted in one genomic region. Red and green peaks mark reads mapped to forward and reverse

strands of the genome, respectively. Lower panel: amplification of the region surrounding one peak,

with read pile ups for TdT and NT sequencing converging on the site of the strand-break. (**b**)

Nb.BsmI mapping data were used to define data mining parameters for accuracy and sensitivity of

Nick-seq. In general, higher ratios yield greater accuracy but lower sensitivity. (**c**) Mapping PTs

across the *S. enterica* genome by Nick-seq. (**d**) Application of Nick-seq to quantify abasic sites

generated by H_2O_2 exposure in *E. coli*.

Supplementary Information

Supplementary Figure 1. Mapping Nb.BsrDI induced DNA strand-break sites in *E.coli* genomic

DNA by Nick-seq. (a) The Nb.BsrDI cleavage motif derived from Nick-seq data. (**b**) A Venn

diagram depicting the overlap of Nick-seq detected sites and Nb.BsrDI motif sites in the *E.coli*

genome.

Supplementary Figure 2. Flanking sequence frequency analysis of H_2O_2-induced EndoIV-

specific DNA damage sites based on Nick-seq data for genomic DNA (**a**) and plasmid (**b**) from

cells exposed to a sublethal H_2O_2 dose. Site 0 represents a Nick-seq detected site.

Supplementary Figure 3. Detection of H_2O_2-induced EndoIV-specific DNA damage sites on

genomic DNA (a) and an endogenous plasmid (b) in *E. coli* by Nick-seq.

Supplementary Figure 4. Distributions of H_2O_2-induced EndoIV-specific DNA damage sites on

E.coli genomic DNA and plasmid. Outward from the center, circles represent: 0 and 0.2 mM H_2O_2

induced EndoIV-specific DNA damage sites.

Supplementary Table 1: Statistical analysis of paired-end DNA sequencing reads from Nick-seq.

Supplementary Table 2: Nb. BsmI sites identified by Nick-seq on the *E.coli* genome.

Supplementary Table 3: Nb. BsrDI sites identified by Nick-seq on the *E.coli* genome.

Supplementary Table 4: Phosphorothioate sites identified by Nick-seq on Salmonella genome.

Supplementary Table 5: AP sites identified by Nick-seq on *E.coli* genome.

Supplementary Table 6: AP sites identified by Nick-seq on plasmid in *E.coli*.

Supplementary Table 7: Regional distribution of AP sites identified by Nick-seq.

Online Methods Section

Materials. Nicking endonuclease Nb. BsmI, Nb. BspQI, Nb. BsrDI, Endonuclease IV, DNA polymerase I, OneTaq DNA polymerase, dNTPs, Nci I, Exonuclease III and RecJ$_f$ were purchased from New England Biolabs. All DNA oligos were synthesized by Integrated Device Technology, Inc. (IDT). ddNTPs and alpha-thio-dNTPs were purchased from TriLink BioTech. Agilent Bioanalyzer 2100 was used for size analysis of DNA fragments. Other chemicals were of molecular biology grade. All cell lines used in this work are readily available from the authors.

Cell growth and preparation of DNA. The PT-containing strain *Salmonella enterica* serovar Cerro 87 and its genomic DNA were prepared as described previously[9]. *E.coli* DH10B was used for nicking enzyme and H$_2$O$_2$-induced DNA damage mapping studies. A single colony of *E. coli* DH10B was grown in 5 mL LB medium overnight at 37 °C. 1mL cells were harvested by centrifuge at ambient temperature (unless indicated otherwise). Cells were resuspended and diluted with fresh 10 LB medium to a starting optical density at 600 nm (OD600) of 0.1, followed

by growth at 37°C, 230 rpm until OD600=0.8 for DNA extraction or H_2O_2 treatment. 10 µL diluted

H_2O_2 solution were added to the culture with a final concentration 0.1, 0.5, 1 and 2 mM. As un-

exposed control, 10 µL sterile water was used instead of H_2O_2. After sitting at ambient temperature

for 30 min, 10 µL of the cells were used for lethal dose (LD) analysis by counting the colony

formation unit on LB agar plate. All the rest cells were harvested for DNA extraction with

OMEGA bacterial genomic DNA or plasmid isolation kit by following the manufacture's protocol.

Mapping of modification/damage sites on DNA by NT-dependent method. These

studies were initiated by random fragmentation of purified genomic DNA (1 µg) in each of three

separate digestions with NciI, or HindIII and XhoI, or SalI, XbaI and NdeI. RNase A was also

added to each reaction to remove trace of contaminating RNA. After digestion, the DNA was

purified using a Qiagen PCR Purification Kit. The three purified DNA samples were mixed for the

blocking step. Blocking of pre-existing strand-break sites was achieved in a reaction mixture

(40 µl) containing 4 µl of reaction buffer (NEBcutsmart buffer), 1 µl of shrimp alkaline

phosphatase (NEB), and 1 µg of template genomic DNA, with incubation at 37 °C for 30 min to

remove phosphate at 3' end of the strand-breaks. The phosphatase was then inactivated by heating

at 70°C for 10 min. After cooling, 2 µl of ddNTPs (2.5 mM each, TriLink) and 1 µl of DNA

polymerase I (10 U, NEB) was added to the reaction with incubation at 37 °C for 40 min to block

any pre-existing strand-break sites. Shrimp alkaline phosphatase (1 µl) was then added at 37 °C

for 30 min to degraded excess ddNTPs and the reaction was terminated by heating at 75 °C for

10 min. Following de-salting using a DyeEx column (QIAGEN), the DNA was ready for one of

the following nick creation or conversion procedures.

Nicking *E. coli* DH10B genomic DNA with Nb. BsmI and Nb. BsrDI was accomplished

in a 50 µl reaction mixture containing 1 µl Nb. BsmI (10 U, NEB), 1 µl Nb. BsrDI (10 U, NEB),

1 µg genomic DNA, and 1X NEBcutsmart buffer incubated at 65 °C for 1 h. The reaction was

terminated by heating at 80 °C for 20 min and cooled down to 4 °C at the rate of 0.1 °C/s. The

reaction product was used for NT- or TdT- reactions as described below with no further

purification.

For mapping PT modifications, 40 µl of blocked DNA from *S. enterica* was mixed with 5

µl of dibasic sodium phosphate buffer (500 mM, pH 9.0) and 2 µl of iodine solution (0.1 N,

FLUKA). After incubation at 65 °C for 5 min and cooling to 4 °C, the reaction product was purified

using a DyeEX column (QIAGEN) to remove salts and iodine. The purified product was treated

with shrimp alkaline phosphatase by adding 5 µl of NEBcutsmart buffer and 1 µl of phosphatase

to remove 3'-phosphates arising from iodine cleavage. After incubation at 37 °C for 20 min and

75 °C for another 10 min, the product was kept on ice for the following NT- or TdT- reactions with

no additional purification.

For mapping H_2O_2-induced AP sites, genomic DNA was extracted from H_2O_2-treated *E.*

coli and the AP sites converted to strand-breaks in a 50 µl reaction mixture containing 1 µg

genomic DNA, endonuclease IV (20 U), and 1X NEBcutsmart buffer, with incubation at 37 °C for

60 min. The reaction mixture was then kept on ice and used for the following NT- or TdT- reactions

with no further inactivation or purification.

Following splitting of the nicked sample into two portions for NT- and TdT- reactions, the

NT- reaction was achieved by further splitting the DNA sample into two parts: one for NT-reaction

and the other as a negative control. NT-reaction was performed in a 50 µl reaction system

containing 2 µl of α-thio-dNTPs (2.5 mM each, TRILINK), 1X of NEBcutsmart buffer, 2 µl of

DNA polymerase I, and the DNA template. The negative control consisted of H_2O instead of DNA

polymerase I. The reaction mixture was incubated at 15 °C for 90 min and then terminated by

heating at 75 °C for 20 min. The product was ready of template DNA digestion after the

purification by DyeEx. The template DNA digestion reaction was performed in a 50 μl reaction

system containing 200 U of exonuclease III, 5 μl NEBcutsmart buffer and DNA sample by

incubating at 37 °C for 60 min. The DNA was then denatured by heating at 95 °C for 3 min and

crashing on ice. RecJ$_f$ (60 U) was then added to the reaction mixture with incubation at 37 °C for

60 min. For some DNA samples with high complexity of structure and/or modifications, digestion

with an additional 60 units of RecJ$_f$ might be necessary. After digestion, the enzymes were

inactivated by incubation at 80 °C for 10 min. The DNA product was then purified using a Zymo

Oligo Clean & Concentrator kit (Zymo) following the manufacturer's protocol. The purified

product was ready for Illumina library preparation.

Illumina library preparation was performed by the Clontech SMART ChIp-seq kit

(Clontech) by following the manufacturer's protocol. 12 cycles were used in the final step of PCR

amplification. The PCR product of each sample was combined with its corresponding negative

control and then size selected using AMPure XP beads (NEB). The purified library was submitted

to Illumina NextSeq 500 instrument for 75 bp paired-end sequencing.

Mapping of modification and damage sites by TdT-dependent method. The steps of

DNA fragmentation, blocking, and nick conversion are the same as described above in NT-

method.100 ng of the nick-converted DNA was denatured by heating at 95 °C for 3 min in 20 μl

of H$_2$O, followed by adding A poly(T) tail to the ssDNA in a 30 μl reaction system containing 3

μl DNA SMART buffer (Clontech), 1 μl Terminal Deoxynucleotidyl Transferase (TdT, Clontech)

and 1 μl DNA SMART T-Tailing Mix (Clontech) by incubating at 37 °C for 20 min and

terminating the reaction at 70 °C for 10 min. The primer annealing and template switching reaction

was then performed with the Clontech SMART ChIp-seq kit (Clontech) by following the

manufacturer's protocol. The final step of PCR was perfomed using the Illumina primers provided

in ChIp-seq kit and 12 cycles were used for amplification. The PCR product of each sample, with

unique sequencing barcode, was combined with its corresponding negative control and then size

selected using AMPure XP beads (NEB). The purified library was submitted to Illumina NextSeq

500 instrument for 75 bp paired-end sequencing.

Data analysis. Sequencing results were processed on the Galaxy web platform

(https://usegalaxy.org/). Initially, the paired-end reads were pre-processed by Trim Galore! to

remove adapters, as well as trimming the first 3 bp on the 5' end of read 1. All the reads were

aligned to the corresponding genome using Bowtie 2. A custom method for peak calling of

sequencing data was developed with BamTools, BEDTools and Rstudio. Briefly, the BamTools

results were filtered based on R1 (selected for NT data) or R2 (selected for TdT data). The 5'

coverage (experiment sample and controls) or full coverage (controls) on each position were

calculated based on the filtered BamTools results by BEDTools (positive and negative strand

separately). For both NT or TdT data on each strand, three ".tabular" files containing the genome

position and their corresponding read coverage (sample_coverage_5.tabular,

control_coverage_5.tabular, control_coverage_full.tabular) were analyzed and exported from

Galaxy after the previous BEDTools. These data were used to normalize the read coverage by the

sequencing depth and then calculate the read coverage ratio of specific position compared to its

up- downstream position in the same sample and the same sample in negative controls. Three ratios

were calculated at each position by RStudio for modification site calling: coverage of position N

(sample)/coverage of position N-1(sample), coverage of position N(sample)/coverage of position

N+1(sample), and coverage of position N(sample)/coverage of position N(control). Positions with

a ratio>1 were retained using the following R scripts: TdT_positive_strand.R

TdT_negative_strand.R NT_positive_strand.R NT_negative_strand.R From these datasets, the

intersection of the datasets from the NT and TdT methods were calculated using the following R

scripts: TdT_positive+NT_negative.R TdT_negative+NT_positive.R The output files (CSV files;

Excel format) contain the read coverage ratio information for the putative nick sites. The ratio

cutoffs can be varied in the Excel spreadsheet as needed. For example, for site-specific nicking by

Nb. BsmI and Nb. BsrDI, we determined that a ratio >2 was adequate to capture nearly all sites,

while for variable sites (PT) or unknown samples (H_2O_2), the ratio was increased to 5-10.

a

Fragmentation

3'-OH Blocking

Convert **X** to strand break

α-S-dNTP NT   TdT **dTTP**

Nuclease

Library prep

Sequencing   Data mining

NT peaks

5'   3'
3'   5'

TdT peaks

Site 1   Site 2   Site 3

b

- *Filter raw reads for seq coverage ≥5*

- *Parameters for refining filtered reads*
  - x = coverage at N/coverage at N-1
  - y = coverage at N/coverage at N+1
  - z = coverage at N/coverage at N neg ctl

- *Steps for refining the filtered reads*
  1. Filter NT data for x>1, y>1, z>0
  2. Filter TdT data for x>2, y>2, z>2
  3. Intersect step #1 and #2 datasets
  4. Merged data used for final map

TdT Coverage ≥5   **NT** Coverage ≥5

x>2   x>1
y>2   y>1
z>2   z>0

Merged data for map