1
2
3

Alpha 6 Integrins Regulate the Expression of Laminin-511 and CXCR4 to Promote Endothelial Tubular Morphogenesis

6
7

Hao Xu, Kevin Pumiglia and Susan E. LaFlamme*

Department of Regenerative and Cancer Cell Biology
Albany Medical College
Albany NY 12208

13
14

Running Title: Endothelial morphogenesis

Corresponding author: laflams@amc.edu

Key words: Integrin, laminin, CXCR4, angiogenesis

Summary statement

Endothelial-secreted laminin-511 and its receptors, α6 integrins promote tubular morphogenesis

by regulating the expression of the chemokine receptor, CXCR4. Additionally, the depletion of

α6 integrins from established tubes results in the loss of tubular integrity and laminin-511

expression.

Abstract (180 words)

During angiogenesis, endothelial cells engage components of the extracellular matrix through

integrin-mediated adhesion. Endothelial cells express laminin-411 and laminin-511 that bind to

integrins, including the α6 integrins, α6β1 and α6β4. However, little is known about the

contribution of these laminins to endothelial tubular morphogenesis and stability. We used two

organotypic angiogenesis assays in conjunction with RNAi approaches to demonstrate that

endothelial depletion of either the α4 chain of laminin-411 or the α5 chain of laminin-511

inhibited sprouting and tube formation. Depletion of α6 integrins resulted in similar phenotypes.

Interestingly, depletion of α6 integrins also inhibited the expression of laminin-511, which

correlated with the loss of tubular stability. Loss of either α6 integrins or laminin-511 resulted in

the inhibition of the expression of CXCR4, a gene previously associated with sprouting

endothelial cells. Pharmacological inhibition of CXCR4 signaling suppressed endothelial

sprouting and morphogenesis, suggesting that α6 integrins and laminin-511 promotes

endothelial tubular morphogenesis in part by regulating the expression of CXCR4. Taken

together, our results suggest that α6 integrins regulate gene expression to promote both early

events in tubular morphogenesis, as well as the stability of established endothelial tubes.

Introduction

Angiogenesis is a process by which new vessels sprout from pre-existing vasculature,

anastomose with neighboring sprouts to form new networks, which then remodel and mature

(Carmeliet, 2003). Angiogenesis contributes to both normal and pathological processes,

including tissue repair, tumor progression, and inflammation (Carmeliet, 2005). Thus,

understanding the mechanisms that regulate various aspects of this process remains an

important objective.

Endothelial cells interact with components of the extracellular matrix (ECM), including

fibronectin, to promote early stages of angiogenesis, whereas the binding of endothelial cells to

components of the basement membrane, such as laminin, contributes to vessel stability (Song

et al., 2017; Thyboll et al., 2002). Although endothelial cells can adhere to ECM components

provided by other cell types, such as those present in the provisional matrix that is formed

during tissue repair (Eming et al., 2007), endothelial cells themselves secrete both ECM and

basement membrane components, including fibronectin and laminins (Hallmann et al., 2005;

Turner et al., 2017).

Laminins are heterotrimeric proteins, containing α, β and γ chains (Yurchenco, 2011).

Endothelial cells express laminin-411, which contains the α4, β1 and γ1 chains and laminin-511,

which contains the α5, β1 and γ1 chains (Hallmann et al., 2005). Current data suggests that

these laminins support vessel stability. The global deletion of the laminin-α4 (Lm- α4) chain

resulted in embryonic hemorrhaging (Thyboll et al., 2002), whereas the endothelial-specific

deletion of Lm- α5 chain reduced endothelial barrier function (Song et al., 2017). Both of these

phenotypes suggest that these laminin isoforms promote vessel stability; however, whether

these laminins are required for endothelial tubular morphogenesis has not been examined.

Integrins are α/β heterodimeric receptors that bind to ECM proteins and components of the

basement membrane, including laminins, to mediate adhesion and to activate signaling

pathways that cooperate with growth factor receptors to regulate cell behavior (Danen and

Sonnenberg, 2003; Streuli and Akhtar, 2009). Endothelial cells express several integrin

heterodimers that allow their interaction with components of the extracellular matrix including

fibronectin and collagen, as well as laminins present in the endothelial basement membrane

(Avraamides et al., 2008; Davis and Senger, 2005). Our current study focuses on determining

whether laminin-411 and laminin-511 and their integrin receptors, α6 integrins (α6β1 and α6β4)

contribute to early steps in endothelial tubular morphogenesis (Hallmann et al., 2005;

Kortesmaa et al., 2000).

82   Other labs have previously analyzed the role of α6 integrins in angiogenesis. Mice with the

83 Tie 1-dependent endothelial deletion of the α6 integrin subunit showed that the depletion of α6

84 integrins promoted angiogenesis (Germain et al., 2010), presumably due in part to the reduced

85 association of endothelial cells with their underlying basement membrane. Interestingly, the Tie

86 2-dependent endothelial deletion of α6 integrins resulted in reduced angiogenesis; this was

87 explained by the loss of α6 integrins on endothelial progenitor cells and Tie-2 expressing

88 macrophages (Bouvard et al., 2010; Bouvard et al., 2014). Notably, *in vitro* and explant

89 angiogenesis assays demonstrated a requirement for α6 integrins; however, this requirement

90 was dependent upon the ECM microenvironment (Bouvard et al., 2010; Bouvard et al., 2014;

91 Lee et al., 2006; Primo et al., 2010; Samarelli et al., 2014).

92   The onset of angiogenesis requires the disruption of stable interactions of endothelial cells

93 with their basement membrane. Thus, to isolate the contribution of laminins and their integrin

94 receptors to early steps in endothelial tubular morphogenesis, we used two organotypic co-

95 culture models that do not require the destabilization of existing tubes. We asked whether the

96 endothelial expression of laminin-411 and/or laminin-511 is required for endothelial tubular

97 morphogenesis, and if so, whether α6 integrins were also required. Our data demonstrate that

98 depleting endothelial cells of the LM-α4 chain, the LM-α5 chain, or the integrin α6 subunit

99 inhibits endothelial morphogenesis. This is the first report demonstrating that endothelial-

100 secreted laminins can contribute to this process. Additionally we show that the expression of

101 laminin-511 and α6 integrins regulate the expression of the proangiogenic gene CXCR4

102 (Salcedo and Oppenheim, 2003; Salvucci et al., 2002; Tachibana et al., 1998; Unoki et al.,

103 2010). Furthermore, the pharmacological inhibition of CXCR4 impairs tubulogenesis

104 demonstrating that the regulation of CXCR4 by α6 integrins and laminin-511 contributes to

105 tubular morphogenesis. Lastly, we show that α6 integrins regulate the stability of endothelial

106 tubes at least in part by regulating the expression of the α5 chain of laminin-511. Taken together,

107 our data suggests that the interaction of α6 integrins and laminin-511 contributes to

108 angiogenesis by regulating the expression of CXCR4 in early stages of endothelial

109 morphogenesis, and that once tubes have formed the interaction of α6 integrins and laminin-

110 511 promotes the stability of endothelial tubes.

Results

Endothelial laminins regulate endothelial tubulogenesis

To examine the contribution of endothelial laminin-411 and laminin-511 in tubular

morphogenesis, we employed RNAi technology in conjunction with two organotypic

angiogenesis assays: the planar co-culture model and the bead sprout assay. In the planar co-

culture model, human endothelial cells (HUVECs) are plated at very low density on a confluent

layer of human dermal fibroblasts (Bishop et al., 1999). Endothelial cells then proliferate and

migrate to form cell trains/cords, which then form lumenized tubes over time (Bishop et al.,

1999; Mavria et al., 2006). In the bead sprout assay, endothelial cells are adhered to cytodex

beads and embedded in a fibrin gel, which is then covered with a confluent layer of human

dermal fibroblasts. Endothelial cells sprout out from the beads into the fibril gel and form

lumenized tubes over time (Nakatsu and Hughes, 2008; Nakatsu et al., 2003). Importantly,

endothelial cells express laminin-411 and laminin-511 in addition to Col IV in both organotypic

models as demonstrated by immunofluorescence microscopy (Fig. 1A-B)

To determine whether the expression of either laminin-411 or laminin-511 was necessary

for endothelial morphogenesis, we inhibited their expression with siRNA targeting either the α4

chain of laminin-411 or the α5 chain of laminin-511. Data was analyzed from three independent

experiments. The efficiency of knockdown in each experiment was determined by qPCR (Fig.

1C-D). Interestingly, knockdown of either LM-α5 or α4 chains caused defective endothelial

morphogenesis in planar co-culture with a more dramatic phenotype resulting from inhibiting the

expression of the α5 chain (Fig. 1E-F). Quantitation of the lengths of cell trains/cords formed

after two days showed a mean length of 340 μm in control, whereas depletion of either the Lm-

α4 or-α5 chain resulted in an average length of 175 μm and 95 μm, respectively. Similarly,

depletion of either laminin chain inhibited sprouting in the bead-sprout assay (Fig. 1G-I). Both

the number and lengths of sprouts were decreased (Fig. 1H-I). Our data indicate that both

laminin-411 and laminin-511 contribute to endothelial sprouting and tube formation and that the

expression of one isoform cannot compensate for the loss of the other suggesting that each

laminin isoform may play distinct role during tubular morphogenesis.

Endothelial α6 integrins promote tubular morphogenesis

Since α6 integrins have been shown to bind to both laminin-411 and laminin-511 (Kortesmaa et

al., 2000), we were interested to test the requirement for α6 integrins in our organotypic models

and to determine whether depletion of α6 integrins phenocopied the loss of laminin-411 or

laminin-511. As expected, endothelial cells express α6 integrins in co-culture (Fig. 2A). To

determine the effect of depleting endothelial cells of α6 integrins, we employed lentiviral vectors

for the doxycycline inducible expression of a non-targeting shRNA or shRNAs targeting the

integrin α6 subunit. Induction of these shRNAs results in the co-expression of a GFP reporter.

Endothelial cells were transduced with these lentiviral vectors and the expression of shRNAs

was induced by the addition of doxycycline. Effects of α6 depletion on endothelial

morphogenesis in the planar co-culture were analyzed in three independent experiments. The

efficiency of knockdown from these experiments was analyzed by western blot (Fig. 2B, left

panel). Effects on endothelial cord/tube length were quantified every two days for a total of 10

days (Fig. 2B, middle panel). Images of representative co-cultures at day 6 are shown in Figure

2B (right panel). Taken together, the data show that morphogenesis is inhibited by day 2 and

remained suppressed with very little lengthening of individual endothelial cords in the α6-

depleted condition at day 10. Similar results were obtained with two additional shRNA targeting

sequences (Supplemental Fig. 1).

The depletion of α6 integrins had a similar inhibitory effect in the bead sprout assay (Fig.

2C). Data was obtained from three independent experiments in which α6 expression was

inhibited by the induction of α6 targeting shRNA (Fig. 2C, left panel). Measurements of sprout

length overtime revealed that endothelial sprouting remained inhibited throughout the 8-day

assay (Fig. 2C, middle panel). Images of representative bead sprouts formed at day 8 by control

and α6 depleted endothelial cells are shown in Fig. 2C (right panel). The depletion of α6

integrins phenocopies the depletion of the LM-α5 chain and LM-α4 chain, suggesting that the

interaction between α6 integrins and laminins secreted by endothelial cells is crucial for tubular

morphogenesis.

The expression of α6 integrins promotes endothelial tube integrity

Given our results that α6 integrins are required for tubulogenesis in our organotypic assays,

we asked whether these integrins are also required for the stability of established tubes. To

accomplish this, planar co-cultures were set up with endothelial cells that were transduced with

lentiviruses carrying either α6-targeting or non-targeting shRNAs. Co-cultures were incubated in

the absence of doxycycline for 10 days at which time normal endothelial tubes were well

established. Co-cultures were then treated with doxycycline to induce the expression of non-

targeting or α6-targeting shRNAs and imaged after 8 and 12 days of doxycycline treatment.

Endothelial tubes expressing α6 shRNA progressively became destabilized over time (Fig. 3A),

which was quantified as changes in the morphology of endothelial structures (Fig. 3B).

Alpha-6 integrins regulate the expression of laminin-511

Since laminin-411 and laminin-511 have been associated with vessel maturation and

stability, we asked whether the loss of endothelial tube integrity after the depletion of α6

integrins was accompanied by changes in the expression of these laminins. We examined the

expression of the α5 and α4 laminin chains in planar co-culture assays in which we induced the

expression of either non-targeting or α6 targeting shRNAs after the formation of tubes as

described in Figure 3A. Our results indicate that the expression of the α5 chain of laminin-511 is

lost after the depletion of α6 integrins (Fig. 3C). In contrast, the LM α4 chain is still expressed,

but appears diffuse compared to control (Fig. 3C). The loss of laminin-511 may in part be due to

decreased levels of LM-α5 RNA (Fig. 4A). These results suggest that α6 integrins and laminin-

511 promote endothelial tube integrity.

Laminin-511 and α6 integrins regulate the expression of the pro-angiogenic genes

CXCR4 and ANGPT2

To gain further insight into the mechanisms by which endothelial laminins and α6 integrins

regulate tubular morphogenesis, we asked whether laminin-511 and α6 integrins regulate the

same set of angiogenesis-associated genes. Since depleting endothelial cells of laminin-411

also inhibited sprouting, we tested whether expression of the same or a distinct set of genes

was affected. We focused on the expression of genes previously associated with sprouting

angiogenesis. These include VEGFR2, CXCR4, ANGPT2, Dll4, PDGFB, NRP1, JAG1, and

MMP14 (De Smet et al., 2009; del Toro et al., 2010; Strasser et al., 2010). RNA was isolated

from non-targeting and α6 targeting shRNA expressing endothelial cells and gene expression

was analyzed by qPCR. Interestingly, depletion of either laminin-511 or α6 integrins led to

significant decreases in RNA transcripts for CXCR4 and ANGPT2 (Fig. 4A&C). Depleting

endothelial cells of laminin-411 had little effect on the expression of these genes (Fig. 4B).

CXCR4 signaling is required for endothelial tubular morphogenesis

Since the best characterized mechanism of action for angiopoietin-2, the product of the

ANGPT2 gene, is to antagonize the effects of angiopoietin-1 secreted by neighboring mural

cells (Carmeliet and Jain, 2011), we did not analyze the contribution of ANGPT2 expression in

our model. However, we did analyze the contribution of the chemokine receptor, CXCR4 to

endothelial morphogenesis in our organotypic models. CXCR4 has been previously implicated

in vascular development and angiogenesis (Salcedo and Oppenheim, 2003; Salvucci et al.,

2002; Tachibana et al., 1998; Unoki et al., 2010). Stromal-derived factor-1 (SDF-1) is a ligand

for CXCR4, and both endothelial cells (Salvucci et al., 2002) and fibroblasts (Nagasawa, 2014;

Quan et al., 2015) express SDF-1. Thus, it seemed possible that CXCR4 signaling is required

downstream of α6 integrins and laminin-511 for endothelial tubular morphogenesis. To test this

possibility, we used the pharmacological inhibitor AMD3100, which blocks CXCR4 activity

(Hatse et al., 2002 65; Kalatskaya et al., 2009). In dose response experiments, AMD3100

inhibited both the number and length of sprouts (Fig. 5A). Significant effects on sprout numbers

were observed at 500 nM, whereas sprout length was inhibited at concentrations as low as 100

nM (Fig. 5A). Similarly, endothelial morphogenesis was inhibited in planar co-culture assays in a

dose-dependent manner with significant inhibition starting at 100 nM of the inhibitor (Fig. 5B).

These data suggest that laminin-511 and α6 integrins regulate CXCR4 expression to promote

endothelial tubular morphogenesis.

Discussion

We used two organotypic co-culture models to demonstrate (1) that the endothelial

expression of laminin-411, laminin-511, and α6 integrins is required for tubular morphogenesis;

(2) that α6 integrins regulate the expression of laminin-511; (3) that laminin-511 and α6 integrins

regulate the expression of CXCR4 and ANGPT2; and (4) that CXCR4 signaling is a critical

component in the α6/laminin-511 morphogenesis response.

Several matrix proteins and integrin heterodimers can promote endothelial morphogenesis

and angiogenesis (Avraamides et al., 2008; Hynes, 2007; Senger and Davis, 2011). The

contribution of specific matrix proteins and integrins are context-dependent. Cell culture and

aortic ring explant cultures have been employed to examine the role of α6 integrins in tubular

morphogenesis and new vessel growth. Alpha-6 integrins were required when these assays

were performed in Matrigel, which is rich in laminin-111, a ligand for α6 integrins (Bouvard et al.,

2012; Primo et al., 2010). However, in collagen gels, α6 integrins were not required (Bouvard et

al., 2012; Primo et al., 2010). These findings suggest that the composition of the ECM

microenvironment can dictate usage of specific integrin heterodimers. However, endothelial

cells can secrete fibronectin, and endothelial expressed fibronectin and the αvβ3 and α5β1

fibronectin-binding integrins are essential for developmental retinal angiogenesis (Turner et al.,

2017). It is not clear why fibronectin expressed by endothelial cells does not override the

requirement for α6 integrins in assays performed in Matrigel. It is possible that the ECM

components present in the microenvironment together with the available angiogenic and other

growth factors dictate the role or regulate the expression of endothelial-secreted fibronectin,

suggesting that the contribution of endothelial fibronectin is context-dependent. This is

supported by the finding that endothelial expressed fibronectin and its integrin receptors are not

required for tumor angiogenesis (Murphy et al., 2015). Similarly, the reliance of α6 integrins and

endothelial-secreted laminin may also be context dependent. Both organotypic models present

an ECM microenvironment similar to that observed in wounds (Eming et al., 2007). In the planar

co-culture assay, there is a well-established fibronectin matrix secreted by fibroblasts, and the

bead sprout assay is performed in fibrin gels. The reliance of α6 integrins and endothelial-

secreted laminins could be due to a combination of the available matrix proteins and the growth

factor milieu present in these assays.

We analyzed the expression of a number of genes that are regulated during angiogenesis,

some of which are enriched in tips cells of sprouting vessels (De Smet et al., 2009; del Toro et

al., 2010; Strasser et al., 2010). The expression of the majority of genes interrogated did not

change upon depletion of either the LM-α5 chain or the integrin α6 subunit. However, the levels

of both CXCR4 and ANGPT2 were reduced upon the depletion of either α6 integrins or the LM-

α5 chain, suggesting that α6 integrins and LM-511 promotes their expression. Both the global

and endothelial specific knockout of CXCR4 resulted in defects in vascularization (Ara et al.,

2005). In our models, we demonstrated that tubular morphogenesis is regulated by CXCR4,

whose expression is regulated by α6 integrins and laminin-511. It is possible that CXCR4

signaling is a major regulator of endothelial morphogenesis in these models; thus, the

dependence on α6 integrins and laminin-511.

Our results also show that α6 integrins promote the expression of laminin-511 at least

partially through regulation of LM- α5 RNA. Furthermore, the loss of α6 integrins and the

concomitant loss of laminin-511 in established endothelial tubes resulted in the loss of their

structural integrity, suggesting that α6 integrins and laminin-511 contribute to vessel stability.

This is consistent with *in vivo* data discussed above demonstrating that the loss of expression of

α6 integrins in endothelial cells enhances tumor angiogenesis.

In contrast to the depletion of laminin-511, there was no overlap between the genes

regulated by loss of expression of the laminin-411 and the loss of α6 integrins. We anticipated

the down regulation of the Notch ligand, DLL4 upon LM-α4 depletion, as previous studies

demonstrated that DLL4 expression is down regulated in LM-α4 null mice during developmental

retinal angiogenesis (Stenzel et al., 2011). We also did not observe an obvious connection

between laminin-411 expression and the expression of α6 integrins, suggesting that another

integrin heterodimer engages laminin-411 to promote endothelial morphogenesis. The α2β1

integrin is a potential contributor as previous *in vitro* studies demonstrated that α2β1 plays a

central role in endothelial cell adhesion to laminin-411 (Stenzel et al., 2011).

In summary, the characterization of the contributions of laminins and α6 laminin-binding

integrins during angiogenesis in the adult is complicated by their roles in maintaining vessel

integrity (Miner et al., 1998; Thyboll et al., 2002; Turner et al., 2017). We were able to analyze

the role of endothelial-expressed laminins in early steps in tubular morphogenesis in the planar

co-culture and bead sprout assays, as endothelial morphogenesis in these models does not

require the disruption of established endothelial tubes. Our data support the conclusion that α6

integrins and laminin-511 positively regulates the expression of CXCR4 to promote early events

in tubular morphogenesis and also protects the integrity of established endothelial tubes.

Materials and methods

Cell Culture

Human umbilical vein endothelial cells (HUVECs) were from Lonza (Allendale, NJ) and were

cultured in in EGM-2 (Lonza, CC-3162). Adult human dermal fibroblasts (HDFs) were isolated

and characterized as previously described {Varney, 2016 #54; Zheng, 2019 #55) and

generously provided by the Van De Water laboratory (Albany Medical College). Human

embryonic kidney epithelial 293FT cells (HEK293FT) were a kind gift from Dr. Alejandro Pablo

Adam lab (AMC). HDFs and HEK293FT cells were cultured in DMEM (Sigma D6429) containing

10% FBS (Atlanta Biologicals), 100 units/ml penicillin (Life Technologies), 100 µg/ml

streptomycin (Life Technologies), and 2.92 µg/ml L-glutamine (GE LifeSciences). All cells were

cultured at 37°C in 5% CO2.

Antibodies and Reagents

Antibodies used in the study are from the following sources. Mouse monoclonal antibody to

CD31 (JC70A) was from Dako (Santa Clara, CA). Rabbit polyclonal antibody to CD31 (M-20)

was from Santa Cruz Biotechnology Inc. (Santa Cruz, CA). FITC-conjugated UEA lectin (L9006)

was from Sigma Aldrich (St. Louis, MO). Rat monoclonal to the α6 integrin subunit (GoH3) was

from BD Biosciences (Billerica, MA). Rabbit polyclonal antibody to Collagen IV (ab6586), rabbit

polyclonal antibody to laminin-111 (ab11575), mouse monoclonal antibody to the laminin α4

chain (ab205568), and mouse monoclonal antibody to the laminin α5 chain (ab77175) were

from Abcam (Cambridge, MA). Donkey anti-rat IgG Alexa Fluor 488 (A21208), donkey anti-

mouse IgG Alexa Fluor 488 (A21202), donkey anti-mouse IgG Alexa Fluor 568 (A10037), and

goat anti-rabbit IgG Alexa Fluor 488 (A11034) were from ThermoFisher Scientific (Waltham,

MA).

siRNA

HUVECs were plated in 6-well tissue culture plates and transfected with siRNA at a 50 nM

concentration with RNAiMAX (ThermoFisher) using the protocol provider by the manufacturer.

HUVECs transfected with siRNA were assayed for knockdown and used in planar co-cultures

(described below) 48 h after transfection. HUVECs transfected with siRNA and used in bead

sprout assays (described below) were transfected during bead coating and assayed for

knockdown at the end of experiment. siRNAs targeting were purchased from Sigma Aldrich (St.

Louis, MO). The non-targeting control was MISSION siRNA Universal Negative Control #1

(SIC001). The nucleotide sequences of targeting siRNAs are as follows: laminin α4 chain (5'-

GCAGAATCCTGCTATAGGA- 3'), laminin α5 chain (5'-CTGATCACCTGCACGGGCA-3'), and

the integrin α6 subunit (5'-GGATTGTTCGTGTAGAGCA-3').

Inducible shRNA

Doxycycline-inducible lentiviral (SMART) vectors harboring shRNAs targeting the α6 integrin subunit or a non-targeting (NT) shRNA were purchased from Dharmacon (Lafayette, CO). Lentiviruses were produced by co-transfection of HEK293FT cells with the shRNA expression vector together with the packaging plasmid, psPAX2, coding for Gag, Pol, Rev, Tat (#12260, Addgene), and the envelope plasmid, pMD2.G, coding for VSV-G (#12259, Addgene). HUVECs were transduced with filtered viral supernatant plus 8 μg/ml polybrene. Cells were induced with doxycycline (100 ng/ml) for 48 h prior to adding HUVECs to co-cultures or in some experiments after 10 days in co-culture as described below. The nucleotide sequences of the shRNAs are as follows: NT (5'-TGGTTTACATGTTGTGTGA-3'), α6 targeting sequence 1 (5'-AAAACCACGGCTCCACTGT-3'), α6 targeting sequence 2 (5'-CTCCATGCACACTTTCTGT-3'), and α6 targeting sequence 3 (5'-AACTCGATCCGCGTGCATG-3').

Quantitative PCR (qPCR)

TRIzol (ThermoFisher) was used to isolate RNA from siRNA transfected HUVECs, as well as, shRNA expressing HUVECs. Extraction of RNA from bead sprout assays (described below) using TRIzol was performed after the removal of HDFs with trypsin-EDTA Solution 10X (59418C, Sigma). cDNA was synthesized with iScript Reverse Transcription Supermix (BioRad) using 1 μg of RNA. Equal amounts of cDNA were used in qPCR reactions performed with iQ SYBR Green Supermix (BioRad). The nucleotide sequences of the qPCR primers used are listed in Table 1.

Western blotting

Western blotting was used to confirm RNAi induced knockdown. Cells were lysed in mRIPA buffer (50 mM Tris pH 7.4, 1% NP-40, 0.25% Na Deoxycholate, 150 mM NaCl, 1 mM EDTA) containing both phosphatase (Sigma, #4906837001) and protease inhibitor cocktails (ThermoFisher, 78440). Equal amounts of protein (20 to 40 μg) were separated by SDS-PAGE and transferred to nitrocellulose for antibody probing. Imaging was performed with a ChemiDoc XRS+ (BioRad) and quantitation with Image Lab (BioRad).

Organotypic culture assays

Planar co-culture: As one organotypic culture, we utilized the planar co-culture model developed by Bishop and colleagues (Bishop et al., 1999) and modified by the Pumiglia lab (Bajaj et al.,

2012). This model reconstitutes some of the complex interactions that occur during

angiogenesis among endothelial cells, the ECM and supporting cells. HDFs were seeded in

tissue cultures dishes ± glass coverslips and cultured to confluence. The medium was changed

to EGM-2. HUVECs, expressing targeting or non-targeting RNAi, were then seeded 16 h later at

a density of 20,000 cells per 9 cm^2 and cultured up to 10 days. ShRNA-mediated knockdown in

pre-formed tubes was accomplished by culturing HUVECs expressing doxycycline-inducible

lentiviral (SMART) vectors on HDFs for 10 days, in the absence of doxycycline with medium

changed every 48 h. Doxycycline was then introduced on day 11, at a concentration of 100

ng/ml and refreshed every 48 h for up to 16 days. Endothelial morphogenesis and changes in

tube structure were analyzed by immunofluorescence microscopy.

Bead sprout assay: To study endothelial sprouting, we employed the bead sprout assay as

described by Nakatsu and Hughes (Nakatsu and Hughes, 2008). Cytodex 3 beads (GE) were

coated at ~1000 HUVECs per bead inside of a 2 ml microcentrifuge tube for 4 at 37°C, mixing

gently by inverting the tubes every 20 min and transferred to a T25 flask and incubated at 37°C,

overnight. Beads were then washed 3X with EGM-2 medium and re-suspended in PBS

containing 3 mg/ml of fibrinogen (Sigma, #F8630) and 0.15 U/ml of aprotinin (Sigma, #A6279).

Thrombin (Sigma, #T4648) was added at a final concentration of 0.125 U/ml and the mixture

was plated in wells of and 8-well slide (Corning, #3-35411). The mixture was allowed to clot for

30 min at 37°C. HDFs were then added to the top surface of the fibrin gel in EGM-2 medium at

a concentration of 30,000 cells per well. The formation of sprouts and sprout lengths were

assayed by either immunofluorescence of phase contrast microscopy.

Immunofluorescence Microscopy

Cells were fixed cells with 4% PFA (Electron Microscopy Sciences) for 15 min, permeabilized

with 0.5% Triton X-100 in PBS for 15 min, and then blocked with 2% BSA in PBST for 1 h at RT.

Antibodies were diluted in 2% BSA in PBST and incubated with cells overnight at 4°C. Samples

were then washed 4X with PBST at RT over the course of 4 h, and then incubated for 1 h with

the appropriate secondary antibodies (1:1000 dilution). Following secondary antibody staining,

samples were washed 3X with PBST at RT for 1 h and mounted with SlowFade Gold antifade

reagent (ThermoFisher). Samples were analyzed using a Nikon inverted TE2000-E microscope

equipped with phase contrast and epifluorescence, a digital CoolSNAP HQ camera, a Prior

ProScanII motorized stage and a Nikon C1 confocal system and EZC1 and NIS-Elements

acquisition software. Images were acquired with Plan Fluor 4X/0.13, Plan Fluor 10X/0.30, Plan

Fluor ELWD 20X/0.45, Plan Apo 40X/1.0 oil, and Plan Apo 100X/1.4 oil objectives and analyzed

with either NIS elements (Nikon) or IMARIS software as indicated. Contrast and/or brightness

were adjusted for some images to assist in visualization.

Image analysis

For planar co-cultures, were endothelial cells initially form cell trains/cord, which mature into

endothelial tubes, cord/tube lengths in planar co-cultures were measured by tracing tubes within

each field using NIS Elements (Nikon). Any tubes that extend beyond the field were excluded

from analysis. Tube widths in planar co-cultures were calculated with AngioTool (NIH) by

dividing total tube area by total tube lengths per field. Sprout lengths in bead sprout assays

were measured by tracing each sprout using NIS elements (Nikon) and sprouts per bead were

counted manually.

Statistical analysis

Statistical analysis was performed with GraphPad Prism software using Student's t-test or one-

way ANOVA. P value of $p < 0.05$ was considered to be statistically significant.

Acknowledgements

The authors thank Drs. Livingston Van De Water and C Michael DiPersio for critically reading

this manuscript, Dr. Van De Water for providing human adult dermal fibroblasts, and Debbie

Moran for assistance in the preparation of the figures.

Competing interests

No competing interests declared.

Funding

This work was supported by Institutional seed funds to SEL and by funding from the David E

Bryant Foundation to KP

Data availability

All reagents generated in this study will be made available upon request.

References

Ara, T., K. Tokoyoda, R. Okamoto, P.A. Koni, and T. Nagasawa. 2005. The role of CXCL12 in the organ-specific process of artery formation. *Blood*. 105:3155-3161.

Avraamides, C.J., B. Garmy-Susini, and J.A. Varner. 2008. Integrins in angiogenesis and lymphangiogenesis. *Nat Rev Cancer*. 8:604-617.

Bajaj, A., Q.F. Li, Q. Zheng, and K. Pumiglia. 2012. Loss of NF1 expression in human endothelial cells promotes autonomous proliferation and altered vascular morphogenesis. *PLoS One*. 7:e49222.

Bishop, E.T., G.T. Bell, S. Bloor, I.J. Broom, N.F. Hendry, and D.N. Wheatley. 1999. An in vitro model of angiogenesis: basic features. *Angiogenesis*. 3:335-344.

Bouvard, C., A. De Arcangelis, B. Dizier, I. Galy-Fauroux, A.M. Fischer, E. Georges-Labouesse, and D. Helley. 2012. Tie2-dependent knockout of alpha6 integrin subunit in mice reduces post-ischaemic angiogenesis. *Cardiovasc Res*. 95:39-47.

Bouvard, C., B. Gafsou, B. Dizier, I. Galy-Fauroux, A. Lokajczyk, C. Boisson-Vidal, A.M. Fischer, and D. Helley. 2010. alpha6-integrin subunit plays a major role in the proangiogenic properties of endothelial progenitor cells. *Arterioscler Thromb Vasc Biol*. 30:1569-1575.

Bouvard, C., Z. Segaoula, A. De Arcangelis, I. Galy-Fauroux, L. Mauge, A.M. Fischer, E. Georges-Labouesse, and D. Helley. 2014. Tie2-dependent deletion of alpha6 integrin subunit in mice reduces tumor growth and angiogenesis. *Int J Oncol*. 45:2058-2064.

Carmeliet, P. 2003. Angiogenesis in health and disease. *Nat Med*. 9:653-660.

Carmeliet, P. 2005. Angiogenesis in life, disease and medicine. *Nature*. 438:932-936.

Carmeliet, P., and R.K. Jain. 2011. Molecular mechanisms and clinical applications of angiogenesis. *Nature*. 473:298-307.

Danen, E.H., and A. Sonnenberg. 2003. Integrins in regulation of tissue development and function. *J Pathol*. 201:632-641.

Davis, G.E., and D.R. Senger. 2005. Endothelial extracellular matrix: biosynthesis, remodeling, and functions during vascular morphogenesis and neovessel stabilization. *Circ Res*. 97:1093-1107.

De Smet, F., I. Segura, K. De Bock, P.J. Hohensinner, and P. Carmeliet. 2009. Mechanisms of vessel branching: filopodia on endothelial tip cells lead the way. *Arterioscler Thromb Vasc Biol*. 29:639-649.

del Toro, R., C. Prahst, T. Mathivet, G. Siegfried, J.S. Kaminker, B. Larrivee, C. Breant, A.
Duarte, N. Takakura, A. Fukamizu, J. Penninger, and A. Eichmann. 2010. Identification
and functional analysis of endothelial tip cell-enriched genes. *Blood*. 116:4025-4033.
Eming, S.A., B. Brachvogel, T. Odorisio, and M. Koch. 2007. Regulation of angiogenesis:
wound healing as a model. *Prog Histochem Cytochem*. 42:115-170.
Germain, M., A. De Arcangelis, S.D. Robinson, M. Baker, B. Tavora, G. D'Amico, R. Silva, V.
Kostourou, L.E. Reynolds, A. Watson, J.L. Jones, E. Georges-Labouesse, and K.
Hodivala-Dilke. 2010. Genetic ablation of the alpha 6-integrin subunit in Tie1Cre mice
enhances tumour angiogenesis. *J Pathol*. 220:370-381.
Hallmann, R., N. Horn, M. Selg, O. Wendler, F. Pausch, and L.M. Sorokin. 2005. Expression
and function of laminins in the embryonic and mature vasculature. *Physiol Rev*. 85:979-
1000.
Hatse, S., K. Princen, G. Bridger, E. De Clercq, and D. Schols. 2002. Chemokine receptor
inhibition by AMD3100 is strictly confined to CXCR4. *FEBS Lett*. 527:255-262.
Hynes, R.O. 2007. Cell-matrix adhesion in vascular development. *J Thromb Haemost*. 5 Suppl
1:32-40.
Kalatskaya, I., Y.A. Berchiche, S. Gravel, B.J. Limberg, J.S. Rosenbaum, and N. Heveker. 2009.
AMD3100 is a CXCR7 ligand with allosteric agonist properties. *Mol Pharmacol*. 75:1240-
1247.
Kortesmaa, J., P. Yurchenco, and K. Tryggvason. 2000. Recombinant laminin-8
(alpha(4)beta(1)gamma(1)). Production, purification,and interactions with integrins. *J Biol
Chem*. 275:14853-14859.
Lee, T.H., S. Seng, H. Li, S.J. Kennel, H.K. Avraham, and S. Avraham. 2006. Integrin regulation
by vascular endothelial growth factor in human brain microvascular endothelial cells: role
of alpha6beta1 integrin in angiogenesis. *J Biol Chem*. 281:40450-40460.
Mavria, G., Y. Vercoulen, M. Yeo, H. Paterson, M. Karasarides, R. Marais, D. Bird, and C.J.
Marshall. 2006. ERK-MAPK signaling opposes Rho-kinase to promote endothelial cell
survival and sprouting during angiogenesis. *Cancer Cell*. 9:33-44.
Miner, J.H., J. Cunningham, and J.R. Sanes. 1998. Roles for laminin in embryogenesis:
exencephaly, syndactyly, and placentopathy in mice lacking the laminin alpha5 chain. *J
Cell Biol*. 143:1713-1723.
Murphy, P.A., S. Begum, and R.O. Hynes. 2015. Tumor angiogenesis in the absence of
fibronectin or its cognate integrin receptors. *PLoS One*. 10:e0120872.

Nagasawa, T. 2014. CXC chemokine ligand 12 (CXCL12) and its receptor CXCR4. *J Mol Med*
(Berl). 92:433-439.

Nakatsu, M.N., and C.C. Hughes. 2008. An optimized three-dimensional in vitro model for the
analysis of angiogenesis. *Methods Enzymol*. 443:65-82.

Nakatsu, M.N., R.C. Sainson, J.N. Aoto, K.L. Taylor, M. Aitkenhead, S. Perez-del-Pulgar, P.M.
Carpenter, and C.C. Hughes. 2003. Angiogenic sprouting and capillary lumen formation
modeled by human umbilical vein endothelial cells (HUVEC) in fibrin gels: the role of
fibroblasts and Angiopoietin-1. *Microvasc Res*. 66:102-112.

Primo, L., G. Seano, C. Roca, F. Maione, P.A. Gagliardi, R. Sessa, M. Martinelli, E. Giraudo, L.
di Blasio, and F. Bussolino. 2010. Increased expression of alpha6 integrin in endothelial
cells unveils a proangiogenic role for basement membrane. *Cancer Res*. 70:5759-5769.

Quan, C., M.K. Cho, Y. Shao, L.E. Mianecki, E. Liao, D. Perry, and T. Quan. 2015. Dermal
fibroblast expression of stromal cell-derived factor-1 (SDF-1) promotes epidermal
keratinocyte proliferation in normal and diseased skin. *Protein Cell*. 6:890-903.

Salcedo, R., and J.J. Oppenheim. 2003. Role of chemokines in angiogenesis: CXCL12/SDF-1
and CXCR4 interaction, a key regulator of endothelial cell responses. *Microcirculation*.
10:359-370.

Salvucci, O., L. Yao, S. Villalba, A. Sajewicz, S. Pittaluga, and G. Tosato. 2002. Regulation of
endothelial cell branching morphogenesis by endogenous chemokine stromal-derived
factor-1. *Blood*. 99:2703-2711.

Samarelli, A.V., E. Riccitelli, L. Bizzozero, T.N. Silveira, G. Seano, M. Pergolizzi, G. Vitagliano, I.
Cascone, G. Carpentier, A. Bottos, L. Primo, F. Bussolino, and M. Arese. 2014.
Neuroligin 1 induces blood vessel maturation by cooperating with the alpha6 integrin. *J*
Biol Chem. 289:19466-19476.

Senger, D.R., and G.E. Davis. 2011. Angiogenesis. *Cold Spring Harb Perspect Biol*. 3:a005090.

Song, J., X. Zhang, K. Buscher, Y. Wang, H. Wang, J. Di Russo, L. Li, S. Lutke-Enking, A.
Zarbock, A. Stadtmann, P. Striewski, B. Wirth, I. Kuzmanov, H. Wiendl, D. Schulte, D.
Vestweber, and L. Sorokin. 2017. Endothelial Basement Membrane Laminin 511
Contributes to Endothelial Junctional Tightness and Thereby Inhibits Leukocyte
Transmigration. *Cell Rep*. 18:1256-1269.

Stenzel, D., C.A. Franco, S. Estrach, A. Mettouchi, D. Sauvaget, I. Rosewell, A. Schertel, H.
Armer, A. Domogatskaya, S. Rodin, K. Tryggvason, L. Collinson, L. Sorokin, and H.
Gerhardt. 2011. Endothelial basement membrane limits tip cell formation by inducing
Dll4/Notch signalling in vivo. *EMBO Rep*. 12:1135-1143.

Strasser, G.A., J.S. Kaminker, and M. Tessier-Lavigne. 2010. Microarray analysis of retinal
endothelial tip cells identifies CXCR4 as a mediator of tip cell morphology and branching.
Blood. 115:5102-5110.
Streuli, C.H., and N. Akhtar. 2009. Signal co-operation between integrins and other receptor
systems. *Biochem J*. 418:491-506.
Tachibana, K., S. Hirota, H. Iizasa, H. Yoshida, K. Kawabata, Y. Kataoka, Y. Kitamura, K.
Matsushima, N. Yoshida, S. Nishikawa, T. Kishimoto, and T. Nagasawa. 1998. The
chemokine receptor CXCR4 is essential for vascularization of the gastrointestinal tract.
Nature. 393:591-594.
Thyboll, J., J. Kortesmaa, R. Cao, R. Soininen, L. Wang, A. Iivanainen, L. Sorokin, M. Risling, Y.
Cao, and K. Tryggvason. 2002. Deletion of the laminin alpha4 chain leads to impaired
microvessel maturation. *Mol Cell Biol*. 22:1194-1202.
Turner, C.J., K. Badu-Nkansah, and R.O. Hynes. 2017. Endothelium-derived fibronectin
regulates neonatal vascular morphogenesis in an autocrine fashion. *Angiogenesis*.
20:519-531.
Unoki, N., T. Murakami, K. Nishijima, K. Ogino, N. van Rooijen, and N. Yoshimura. 2010. SDF-
1/CXCR4 contributes to the activation of tip cells and microglia in retinal angiogenesis.
Invest Ophthalmol Vis Sci. 51:3362-3371.
Yurchenco, P.D. 2011. Basement membranes: cell scaffoldings and signaling platforms. *Cold
Spring Harb Perspect Biol*. 3.
548
549

Figure Legends

Figure 1. Laminin-411 and laminin-511 regulate endothelial tubular morphogenesis

(A-B) Endothelial basement membrane components are expressed in co-culture.

Immunofluorescence staining of 3-day planar co-cultures **(A)** or 6-day bead sprouts **(B)** shows

endothelial expression of the laminin α4 (LM-α4) and α5 (LM-α5) chains, the laminin (LM) β1

and γ1 chains, as well as COL IV. Scale bar = 50 μm and 100 μm respectively. **(C-I)** Depletion

of either the LM-α4 or LM-α5 chain inhibits endothelial morphogenesis. **(C-D)** The efficiency of

siRNA-mediated knockdown of the LM-α4 chain **(C)** and LM-α5 chain **(D)** in endothelial cells

used in functional studies was assayed by qPCR for each of the three independent experiments

and plotted as the mean ± s.d. **(E)** Representative images of RFP-expressing endothelial cells

at day 2 of planar co-culture depleted of either the LM-α4 or LM-α5 chain compared to control.

Scale = 100 μm. **(F)** Quantitation of endothelial cord length at day 2 for the different conditions.

Plotted is length of endothelial cords measured from 10 random fields in 3 independent

experiments. Data was analyzed by one-way ANOVA and plotted as the mean length ± s.d. **(G)**

Representative images of endothelial cells depleted of either the Lm α4 or α5 chain in a 5-day

bead sprout assay showing decreased sprout length and sprouts number per bead compared to

control. Arrows indicate examples of sprouts. Scale = 250 μm. **(H-I)** Quantitation of number of

sprouts **(H)** or sprout length **(I)** from 6-8 beads in each of 10 randomly selected fields in 3

independent experiments. Plotted are the **(H)** number of sprouts on individual beads or **(I)** or the

length of individual sprouts **(I)**. Data was analyzed by one-way ANOVA. The mean ± s.d. is

indicated.

Figure 2. Depletion of endothelial α6 integrins inhibits endothelial tubulogenesis. (A)

Immunofluorescence staining of 2-day planar co-cultures indicating the endothelial expression

of α6 integrins. **(B)** Alpha-6 integrins were depleted in endothelial cells by shRNA and the

effects were assayed in 3 independent experiments. Efficiency of knockdown was assayed by

western blot (left panel) Statistical analysis was performed with a two-tailed Student's t-test. The

mean ± s.d is indicated. The effect of α6 depletion on cord/tube length in planar co-culture was

assayed every 2 days for 10 days and compared to control. Cord/tube length was quantified in

three independent experiments. Plotted with the length of individual cords/tubes with each

experiment represent by a different color (middle panel). Representative images are shown from

the 6-day co-cultures (right panel). Scale = 100 μm. **(C)** Similar effects of α6 knockdown were

observed in bead sprout assays (left panel). Efficiency of α6 knockdown was determined by

western blot. Statistical analysis was performed using a two-tailed Student's t-test. The mean ±

s.d. is indicated (left panel). Sprout length was measured every 2 days for 8 days. Plotted is the

length of individual sprouts from 3 independent experiments with each experiment represented

by a different color. Representative images of 8-day bead sprouts assays are shown in the Left

Panel. Scale = 500 μm. *p ≤ 0.05, **p ≤ 0.01.

Figure 3. Depletion of α6 integrins from established endothelial tubes disrupts tube

integrity and inhibits the expression of laminin-511. (A-B) Depleting endothelial tubes of α6

integrins inhibits tube stability. Endothelial tubes were formed for 10 days in the planar co-

culture assay and the expression of α6 or non-targeting shRNA was induced with doxycycline.

(A) Representative images of endothelial tubes after 8 and 12 days of induction of either the α6

or non-targeting (NT) shRNA. **(B)** The disruption of endothelial tubes resulted in altered

morphology best characterized by an increase in the width of the resulting endothelial structures.

Plotted is the average width of individual tubes/field (NT-shRNA) and endothelial structures (α6-

shRNA) measured from ten randomly selected fields in two independent experiments. n=22.

Student's t was used to compare conditions. The mean ± s.d. is indicated. ***p ≤ 0.001.   **(C)**

Comparison of laminin-411 and laminin-511 in endothelial tubes after the depletion of α6

integrins. The depletion of α6 integrins from stable endothelial tubes was performed as

described above. Both endothelial laminins are expressed by stable endothelial tubes as shown

in the NT control. Depletion of α6 resulted in the loss of laminin-511, but not laminin-411. Scale

= 50 μm.

Figure 4. The expression of the CXCR4 and ANGPT2 genes are positively regulated by α6

integrins and laminin-511. (A-B) RNA was isolated from 6-day bead sprouts and analyzed by

qPCR. **(A**) Shown is the efficiency of α6 depletion and the effects of α6 depletion on the

expression of the LM-α4, LM- α5, CXCR4 and ANGPT2 normalized to non-targeting control. **(B)**

Shown is the efficiency of depletion of the LM-α4 chain and the effects of this depletion of the

expression of CXCR4 and ANGPT2. **(C)** Shown is the efficiency of depletion of the LM-α5 chain

and the effects of this depletion of the expression of CXCR4 and ANGPT2. n = 3, **p ≤ 0.01,

***p ≤ 0.001, ****p ≤ 0.0001.

Figure 5. Pharmacological inhibition of the CXCR4 receptor inhibits endothelial tubular

morphogenesis organotypic culture. (A) The dose-dependent reduction in the number of

endothelial sprouts, as well as sprout length in 6-day bead sprout assays in response to the

CXCR4 inhibitor, AMD3100. Left panel - Plotted is the number of sprouts on individual beads

from 3 fields containing 7-10 beads from each of two independent experiments. The mean and

s.d. are indicated. Right panel- Plotted is the length of individual sprouts on these same beads

(B) AMD3100 inhibition of CXCR4 in 6-day planar co-culture causes defective morphogenesis

in a dose-dependent manner. Representative images are shown in the Right Panel. Scale = 100

µm. Left Panel- Plotted is the average length of cords/tubes fields. Six fields were analyzed in

each of two independent experiments. The mean ± s.d. is indicated. One-way ANOVA was used

to compare sprouts or tube lengths. $**p \leq 0.01$, $***p \leq 0.001$ $****p \leq 0.0001$.

Figure 1

Figure 2

A

Days with doxcyline treatment
After tube formation

8   12

NT

α6 shRNA

B

***

Average cord/tube diameter(μm)/field

NT   α6
shRNA

C

Lectin   LM α5 chain   Lectin   LM α4 chain

NT

α6 shRNA

Figure 3

Figure 4

Figure 5

Supplemental Figure 1. (A) Depletion of endothelial α6 integrins using 3 independent shRNA targeting sequences shows similar inhibition of tubular morphogenesis compared to control. Cord/tube lengths were collected from 10 randomly selected fields in 3 independent experiments. **(B)** Knockdown of α6 was confirmed using qPCR. Mean RNA expression is from 3 independent experiments. Alpha 6 targeting sequence 2 was used in Figures 2, 3, and 4 of main text.

A   Gene expression in α6 integrin depleted sprouts

B   Gene expression in LM-511 depleted sprouts

C   Gene expression in LM-411 depleted sprouts

Supplemental Figure 2. Expression of angiogenesis-associated genes in 5-day bead sprout assay compared to control **(A)** alpha 6 integrin-depleted endothelial cells **(B)** laminin-511-depleted endothelial cells and **(C)** laminin-411-depleted endothelial cells were measured by qPCR. Data are plotted as the mean ± s.d. from 3 independent experiments. Expression of each gene was compared to NT control using two-way Student's T-test. *p ≤ 0.05.