Quantification of the Resilience and Vulnerability of HIV-1 Native Glycan Shield at Atomistic Detail

Srirupa Chakraborty[1,2,5], Zachary T. Berndsen[3,4,5], Nicolas W. Hengartner[1], Bette T. Korber[1], Andrew B. Ward[3,4,6], and S. Gnanakaran[1,6]

[1]Theoretical Biology & Biophysics, Los Alamos National Laboratory, Los Alamos, NM 87545, USA
[2]Center for Non-Linear Studies, Los Alamos National Laboratory, Los Alamos, NM 87545, USA
[3]Department of Integrative Structural and Computational Biology, The Scripps Research Institute, La Jolla, CA 92037, USA
[4]IAVI Neutralizing Antibody Center and Collaboration of AIDS Vaccine Discovery, The Scripps Research Institute, La Jolla, CA 92037, USA

[5]These authors contributed equally

[6]Lead Contacts: andrew@scripps.edu, gnana@lanl.gov

Summary

Dense surface glycosylation on the HIV-1 envelope (Env) protein acts as a shield from the adaptive immune system. However, the molecular complexity and flexibility of glycans make experimental studies a challenge. Here we have integrated high-throughput atomistic modeling of fully glycosylated HIV-1 Env with graph theory to capture immunologically important features of the shield topology. This is the first complete all-atom model of HIV-1 Env SOSIP glycan shield that includes both oligomannose and complex glycans, providing results which are physiologically more relevant than the previous models with uniform glycosylation. This integrated approach including quantitative comparison with cryo-electron microscopy data provides hitherto unexplored details of the native shield architecture and its difference from the high-mannose glycoform. We have also derived a measure to quantify the shielding effect over the antigenic protein surface that defines regions of relative vulnerability and resilience of the shield and can be harnessed for rational immunogen design.

Introduction

Protein glycosylation is an essential aspect of post-translational modification, with 50-70% of human proteins been estimated to be glycosylated to some degree(An et al., 2009). These glycans play significant roles in numerous biological processes, such as cellular signaling, recognition and adhesion, protein folding, structural stability, and immune system interactions(Dennis et al., 1999; Dwek, 1996; Moremen et al., 2012). Additionally, envelope proteins from several high-risk viral pathogens which hijack the host protein production and glycosylation machineries, such as HIV (lentivirus)(Burton and Mascola, 2015), Coronavirus (Vankadari and Wilce, 2020; Walls et al., 2016), Lassa (arenavirus)(Sommerstein et al., 2015), Hepatitis C (flavivirus)(Zhang et al., 2017), Epstein Barr (herpesvirus)(Szakonyi et al., 2006), Ebola (filovirus)(Ilinykh et al., 2018; Lennemann et al., 2014), and Influenza(Wang et al., 2009) are also heavily glycosylated. These surface-expressed viral proteins are important immunological targets for neutralizing antibodies that can block viral infection of cells, and form the primary focus of vaccine studies(Amanat et al., 2018; Saphire et al., 2018). However, the dynamic and dense glycan coating can effectively act as a shield for the underlying envelope protein, masking antigenic surfaces, barricading it from the adaptive immune system, and defending against immune surveillance(Crispin et al., 2018; Doores, 2015; Karsten and Alter, 2017; Ringe et al., 2019; Stewart-Jones et al., 2016; Wagh et al., 2018). A deeper molecular level understanding of the glycan shields in these pathogenic viruses may help inform vaccine design strategies that can overcome this protective barrier.

The HIV-1 Envelope Glycoprotein (Env) is a heterodimeric trimer composed of proteins gp120 and gp41, and is responsible for the molecular recognition of the host receptor and fusion into host target cells. A number of obstacles hinder traditional vaccine design methods in case of HIV-1 – namely its remarkable sequence diversity, conformational plasticity, dramatic shifts in position and number of glycans(Zhang et al., 2004), and extremely dense glycosylation making up to approximately half the mass of the entire Env molecule(Lasky et al., 1986). As a result, there has been only limited success in eliciting broadly neutralizing antibodies (bNAbs) to Env vaccine immunogens(Bradley et al., 2016; Haynes et al., 2014; Walker et al., 2010). Structures of bNAbs in complex with Env indicate that these antibodies need to extend through the glycan shield in order to engage with the epitopes at the protein surface(Huang et al., 2014; Julien et al., 2013;

Lyumkis et al., 2013; Pancera et al., 2014). Some bNAbs have evolved to include conserved

glycans as part of the epitope(Crispin and Doores, 2015; Falkowska et al., 2014; Mouquet et al.,

2012; Walker et al., 2009). Moreover, these surface glycans are also critical for Env folding, viral

assembly, and infectivity.

Almost all the glycans on the Env are N-linked and occur at potential N-glycosylation sites

(PNGS, given by the sequon Asn-X-[Ser or Thr], where X is not proline)(Stanley et al., 2015).

Such glycans usually have about 10-20 pyranose rings covalently connected in a tree-like structure

to the asparagine(N) residue of the sequon. These can further undergo additional

processing(Crispin and Doores, 2015) into complex sugars from simple oligomannose, depending

on their spatial location, local protein content, crowding, and producer cell type. A high degree of

processing is indicative of exposure or accessibility of sugars to enzymes. At regions with dense

crowding of glycans, steric constraints limit the activities of the carbohydrate processing

enzymes(Pritchard et al., 2015a; Pritchard et al., 2015b) as the proteins fold and translocate

through endoplasmic reticulum and Golgi apparatus.

The extreme dynamics and conformational heterogeneity stemming from large variance of

possible constituent sugars, linkage, branching patterns, and rotamer flexibility, make the study of

these glycans immensely challenging(Imberty and Perez, 2000; Walsh, 2010). Traditional X-ray

crystallography methods are rendered ineffectual(Chang et al., 2007) since this heterogeneity

prevents proper crystallization of the glycoproteins. Inherent glycan flexibility also complicates

analysis by ensemble techniques like Nuclear Magnetic Resonance (NMR) and single-particle

electron cryomicroscopy (cryo-EM) where the signal from glycans is often too noisy for structural

interpretation at the resolutions necessary for building atomic structure. (Chang et al., 2007; Davis

and Crispin, 2010; Slynko et al., 2009; Woods et al., 1998). Within the ~300 experimental

structures of HIV-1 envelope glycoprotein in the Protein Databank (PDB) (Berman et al., 2000),

only a quarter of the total glycan content has been structurally resolved and are mostly in high

mannose form. Recently there have been a few cryo-EM and X-ray structures published of natively

glycosylated Env(Barnes et al., 2018; Gristick et al., 2016; Lee et al., 2016). Since most of these

structurally resolved glycans were stabilized by interaction with antibodies, they provide little

information about the structural and dynamic details of the individual and collective glycans.

Furthermore, all computational studies of HIV-1 Env glycosylation comprise of oligomannose

glycan moieties ranging from mannose-5 to mannose-9(Ferreira et al., 2018; Lemmin et al., 2017;

Liang et al., 2016; Stewart-Jones et al., 2016; Yang et al., 2017). The main drawback affecting the

quality of such MD simulations is the robustness of conformational sampling. Due to the

intrinsically dynamic nature of glycans, to effectively sample a biologically relevant energy

landscape of the glycoprotein, long and often multiple trajectories need to be run, preferably with

enhanced sampling techniques(Yang et al., 2017).

We had previously established a high throughput pipeline to robustly build atomistic

models of glycans, sample the glycan conformational space, construct the glycan network

topology, and extract molecular level description of the viral glycan shield (Berndsen et al., 2019).

We had also set up a protocol to experimentally validate the models by generating synthetic cryo-

EM datasets and comparing the resulting maps in a quantitative manner. Here, we have employed

a similar quantitative comparison using computational and experimental single-particle cryo-EM

data and show that the dynamics of high-mannose and native glycosylation remains practically

indistinguishable around the glycan stem β-mannose (BMA) residue. This provides further

rationale towards the need for computational modeling of complete glycans in a natively

glycosylated shield. Presently this seems to be the only way to elucidate the ramifications of such

complex sugars on the shield architecture. Therefore, we have built a glycosylated Env having

both oligomannose and complex glycans(Behrens et al., 2016; Cao et al., 2017) based on the

information provided by mass spectrometry (MS)(Behrens et al., 2016). To our knowledge, this is

the first time that a complete computational model of Env SOSIP was generated to include

processed glycoforms, thus obtaining results which are immunologically more relevant.

Using these atomistic models, we have employed graph theory to capture the glycan shield

topological network, pinpoint potential interaction pathways, and identify concerted behavior of

the glycans. The glycan-glycan and glycan-protein interactions influence the behavior of the shield

as a whole and can affect distant sites in the glycan network(Behrens et al., 2016; Ferreira et al.,

2018; Stewart-Jones et al., 2016). Analyses of various network attributes, such as relative centrality

of different glycans and critical subnetwork properties, have aided in detailed examination of the

native glycan shield in the context of bNAb interactions. Important global and local feature

differences due to native-like glycosylation pattern come to light as compared to all high-

oligomannose glycans, arising from the presence of charged sialic acids at the tips and fucose at

the base. Both the simulated models and cryo-EM maps identify a change in projection angle

relative to the peptide surface for fucosylated glycans. We have also been successful in quantifying

the glycan shield based on the number of sugar heavy atoms encountered over the antigen surface

that can be implemented to define regions of relative vulnerability and regions where the glycan

shield blocks access. Due to the rapid yield rate of this method, it can be applied to a large number

of diverse HIV-1 sequences or can be seamlessly extended to model glycan shields of other viruses

in a rapid manner.

Results

Selection of site-specific glycans for native glycosylation of HIV-1 Env

The soluble, recombinant BG505 SOSIP.664 (here on referred simply as BG505) trimer

has been well characterized as a native-like, Env-mimetic model, and serves as the prototypical

immunogen in several vaccine development programs(Sanders et al., 2013; Sanders et al.,

2015). For our study, we have used this system in the pre-fusion closed conformation, for which

several structures have been determined(Barnes et al., 2018; Gristick et al., 2016; Lee et al.,

2016; Stewart-Jones et al., 2016). A single protomer of this BG505 protein contains 28 N-linked

glycosylation sites. While all N-glycans share a common core sugar sequence, they are broadly

classified into three types **(Figure 1)**: (i) oligomannose, in which only mannose residues are

attached to the core; (ii) complex, in which differently branching "antennae" are attached to the

core; and (iii) hybrid, which is a mix of the first two types. Recent novel MS-based approaches

have been successful in identifying site-specific glycosylation profile in multiple HIV-1

trimers(Cao et al., 2017; Go et al., 2017). It is known that each glycosylation site at BG505 has

a distribution of multiple possible glycan species, depending on various factors, such as type of

cell lines, purification methods and sources, structural constraints, and so on. We used a site-

specific distribution of glycan type at each PNGS that was obtained by Behrens *et. al*(Behrens

et al., 2016) and Cao *et. al.*(Cao et al., 2017), to identify the most likely glycan species at each

site. The glycan species at each site was selected based on the highest relative abundance per

the MS studies (Figure 1 in reference (Behrens et al., 2016)), and is listed in Supplementary

Information (SI) **Table S1**. The structure schematic for each of the selected glycan species have been shown in **Figure 1A**.

High-throughput atomistic modeling of the native glycan shield

In the static pictures as obtained from fully glycosylated structures(Gristick et al., 2016; Stewart-Jones et al., 2016) and even single snapshots from computational models, a significant fraction of the protein surface appears to be exposed, with each glycan taking up a particular conformation (**Figure 1B**). However, due to their flexible and dynamic nature, these glycans are not confined to a particular conformation, instead sampling a large volume in space. A previous simulation study suggested(Guvench et al., 2011) that the root mean squared deviations (RMSD) for the carbohydrate regions are more than 4 times larger than that of the underlying protein loops. As a result of this conformational variability, the cumulative effect of the glycans over time is like that of a cloud of glycan atoms that shield the underlying surface from any approaching protein probe (an antibody for example) as illustrated in **Figure 1C**.

Here we have incorporated experimental information-driven conformational sampling in the flexible unstructured protein regions to model the entire glycan shield. Details of the modeling procedure are given in the Methods section, and are briefly reiterated here. A robust ensemble of BG505 glycoprotein atomic models was generated by utilizing a template-free glycan modeling pipeline including a sequence of refinement steps with restraints to enforce proper stereochemistry. The underlying protein scaffold was built by utilizing templates from a number of experimentally determined Env structures deposited in the Protein Data Bank by assimilating the differences in local structural regions including loops, that arise due to flexibility. The glycans were modeled *ab initio* on several distinct protein scaffolds by implementing the ALLOSMOD(Guttman et al., 2013; Weinkam et al., 2012) package of MODELLER(Eswar et al., 2006; Sali, 1995) in a streamlined pipeline.

Due to the initial randomization of the glycan orientations, this integrated technique can sufficiently sample a physiologically relevant conformational space accessible to carbohydrates in a very short time, leading to the overall spatial shielding effect (**Figure 1C**). The torsion angle distributions of 10 different inter-glycan linkages within the ensemble was compared with those

obtained from different glycan structures available in the PDB database using GlyTorsion

webserver(Lütteke et al., 2005) and is shown in **SI Figure S1**. These distributions match very well

between our generated ensemble and the PDB structures (see SI for details), indicating that the

modeled glycans sample a stereo-chemically relevant landscape. In order to understand the effects

just due to native glycans, we have also built a similar ensemble with mannose-9 glycans at all

glycosylation sites, (hereon referred as all-man9 model) for comparison.

Quantitative comparison of simulated models with cryo-EM data indicates glycoform-independent glycan stem dynamics

We generated simulated cryo-EM density maps for both the native and all-man9 ensembles

by utilizing the method that we established earlier to convert *in silico* atomistic models to synthetic

cryo-EM data(Berndsen et al., 2019). Then, we quantitatively compared them to the experimental

cryo-EM maps of BG505.SOSIP.664 from HEK 293F cells which produce a native-like glycan

shield including both oligomannose and complex glycans, and from HEK 293S cells which do not

produce complex glycans(Berndsen et al., 2019). The experimental and simulated maps show

similar features in the shield, with only the first few sugar residues at each site defined at high

resolution **(Figure 2A).** A scale-space analysis developed in our earlier study(Berndsen et al.,

2019) also shows a similar trend to the experimental data **(Figure 2B)**. Progressive smoothing of

the map is accompanied by a large expansion in glycan volume that plateaus around 1.75-2 SD.

The effect of the glycans is best appreciated in comparison to a map generated from a non-

glycosylated ensemble **(Figure 2B - dashed line)**. This suggests the simulation is capturing native-

like dynamics in the glycan shield, at least globally. The 1.75 SD Gaussian filtered maps are

visualized at high and low thresholds in **Figure 2C**, revealing a similar overall shape and topology.

At the level of individual glycan dynamics, comparison of the normalized mean intensity at the

location of each BMA residue for the simulated map with that of the experimental BG505_293F

map reveals a strong positive correlation (**Figure 2D-E**), with a correlation coefficient of ~0.82

(p=6.82e-6). We also confirmed that local map intensity around BMA residues accurately captures

relative differences in Root Mean Squared Fluctuations (RMSF) at each glycan **(SI Figure S2)** for

the native ensemble.

An interesting outcome of the above comparison is that the correlation coefficients remain
similarly strong not only when we compare the all-man9 model to BG505_293S data, but also
when the native model is compared to high-oligomannose experiments and vice versa **(SI Figure**
S2). Therefore, while these comparisons give credence to the validity of the model in accurately
capturing the relative position and flexibility of the glycan stems, they also indicate that the glycan
dynamics are practically identical within the observational limits of cryo-EM analysis between
complex and oligomannose glycans. The native glycosylation pattern on the shield topology only
manifest at the level of the flexible antennae beyond the stem and can therefore only be observed
in detail using computational models.

Fucose-mediated changes in glycan orientation distinguish native glycan shield

While the glycan stem shows indistinguishable dynamics at the BMA sugar, an important
difference between complete models of complex and all-man9 arise due to the presence of the
fucose ring at the core of the complex sugars **(Figure 3)**. By structurally aligning the first core
sugar and the three residues at the base of the glycan (n-1, n, n+1; where n is the glycosylated
asparagine residue) for all the structures in the ensemble, the oligomannose glycans are found to
be spaced symmetrically around the core, while complex sugars have a distinct bend away from
the side where the fucose ring is present **(Figure 3A)**. The change in projection angle of the
principal axis of the glycan relative to the protein surface is predicted to be ~10° in our models.
This difference in orientation preference, along with the presence of the negatively charged sialic
acid tips of the antennae, play notable roles that govern the differences between high oligomannose
and native glycosylation.

The change in projection angle from the fucose ring manifests as a reasonably large change
at the tips of the antennae, but the overall change in position at the core N-acetylglycosamine
(NAG) is too small to be captured in the high-resolution cryo-EM maps **(Figure 3B)**. However, a
close examination of the low-resolution difference map between the BG505_293F and
BG505_293S maps (Berndsen et al., 2019) does reveal some evidence for such changes in
projection angle at sites with complex glycans **(Figure 3C-E)**. For instance, the difference signal
around the glycans at N611 and N618 is asymmetric around the glycan stem and is therefore likely
not the result of differences in occupancy. The difference signal at N611 is located primarily on

the underside of the glycan stem, on the face opposite where we identified density likely belonging

to the fucose ring **(Figure 3D)**, which is suggestive of a change in projection angle similar to what

we observed in the simulations. In addition, examination of the glycan stem models built and

relaxed into the Gaussian filtered BG505_293F and BG505_293S maps used for assessing local

map intensity **(Figure 3E)** show a slight change in projection angle, further confirming this fucose-

mediated glycan reorientation.

<u>Network analysis of glycan topology explicates the shield connectivity</u>

Network analyses has historically been used to study protein allosteric frameworks and

evolutionary paths (Beleva Guthrie et al., 2018; Eargle and Luthey-Schulten, 2012; Huang et al.,

2017; Sethi et al., 2013; Skjaerven et al., 2014), and only recently begun to be applied to in

glycoprotein structural characterization (Lemmin et al., 2017). While each glycan exerts effects in

its immediate vicinity, due to the inter-glycan interactions, their influence can percolate over long

distances across the surface of protein(Berndsen et al., 2019). Long-range glycan interactions can

occur, as with perturbation of a glycan at one site can affect the processing and antibody

interactions of another glycan at a distant site(Behrens et al., 2016; Bricault et al., 2019; Doores,

2015; Stewart-Jones et al., 2016). We have previously shown that understanding long-range

interactions between glycans is important for characterizing global properties of the glycan

shield(Berndsen et al., 2019).

Now, we implement the graph-theory based approach to native glycan shield. Details of

modeling the network are recapitulated in the Methods section. The natively glycosylated BG505

trimer, like all HIV-1 Envs, has a highly dense glycosylation pattern and each glycan samples a

particular region in space (**SI Figure S3A-B**), such that neighboring glycans can occupy

overlapping regions (**Figure 1D**). The fraction of volume overlap gives a measure of the

interaction probability between the constituent glycans (**SI Figure S3C**). Within a protomer there

are three main regions of overlap – the apex, the gp41 base, and the glycan-dense central high-

mannose patch (HMP). Inter-protomer overlaps are primarily due to V1 and V2 loop glycans near

the trimer apex. **Figure 4A** shows the obtained network of BG505 native glycosylation unfolded

and laid out in two dimensions for the ease of visualization. This is a force-directed layout, which

uses attractive forces between adjacent nodes and repulsive forces between distant nodes to reach

an optimum distribution of the node points in space(Fruchterman and Reingold, 1991). For
conceptualization, the network relative to the BG505 structure is provided in **SI Figure S3D.**

The nodes in the central region around the V4 loop, in the HMP, are very highly inter-
connected. Glycans in the V1, V2 apex are also reasonably well-connected, but the connections at
the base near gp41 are relatively sparse, both within the locality, and globally with the rest of the
network. Glycans at positions 234, 355, 462 and 276 connect the sparse base region of the network
with the dense apex and central regions. Glycan 156, 160, and those in the V2 loop, enable inter-
protomer glycan interactions at the apex, stabilizing the protein quaternary trimer
structure(Berndsen et al., 2019). Glycan 197, crucial in the binding of CD4bs and V3-specific
antibodies(Li et al., 2008; Liang et al., 2016), connects the central crowded region of the network
between neighboring protomers. gp41 glycans 611 and 637 also contribute to inter-protomer
pathways, with glycan 637 communicating with glycan 448 directly in the central mannose patch.

<u>Utilizing the glycosylation network to extract functionally relevant topological features</u>

The native glycosylation network is well-connected as indicated by the network degree, or
the number of connections a node has to other nodes (**SI Figure S3E**). Due to the high structural
fluctuations of the V2 loop, the glycans on and flanking V2 can interact with a number of other
neighboring nodes, having the highest degrees of connectivity. There is dense crowding of glycans
around the high mannose patch with a number of inter-connections, resulting in the glycans of this
patch being less processed because of reduced accessibility to enzymes(Coss et al., 2016; Doores
et al., 2010; Pritchard et al., 2015a). The network degree values are relatively lower around the
base of the protein on and around gp41, reflective of the sparse architecture of glycan topology in
this region. The site N332 is central in the glycan interaction network, and is known to have both
direct and subtle long-range influences over a number of bNAb sites(Behrens et al., 2016; Bricault
et al., 2019). We calculated the shortest paths to all glycans from the N332 glycan site using the
Floyd-Warshall algorithm(Floyd, 1962; Warshall, 1962), where the glycan distances were edge-
weighted (**Figure 4B**). Our calculation shows the most probable pathway over which each glycan
feels the influence of the glycan at site N332, providing useful insights into how these inter-glycan
communications can occur.

To elucidate the relative influence of each of the glycans on the network, we calculated the
eigenvector centrality, or eigencentrality, of the nodes to measure its connectivity to the network.
A large relative value indicates that the node is well connected to the network and thus is
"centrally" located, whereas a low relative value indicates that the node is on the periphery of the
network. Operationally, the eigencentrality is the eigenvector associated with the largest
eigenvalue of the adjacency matrix, that in position (i,j) reports the overlap of the ensemble of
glycan at sites i and j. Value of that particular matrix element will be zero if the glycans in position
i and j do not physically interact (details given in Methods section). The normalized eigencentrality
of the glycans are projected on the network as a colormap in **Figure 4A**. The eigencentrality
increases towards the middle of the graph, with the glycans at the crowded HMP with a large
number of connections between each other having the highest centrality values. When we increase
the threshold of volume overlap needed to form an edge, those glycans with the least centrality
leave the network first. Glycan 611 is the first to be eliminated, succeeded by some of the inter-
protomer interactions and the other gp41 glycans, following the relative centrality scores. The
HMP with high eigencentrality values persists throughout even with high overlap thresholds. We
have previously confirmed for all-man9(Berndsen et al., 2019) that successive enzymatic digestion
of glycans from Env loosely follows a pattern that matches with the network centrality. Those
glycans which are sparsely connected to each other, having lower centrality, are eliminated earlier
during the process of endoH digestion. On the other hand, the glycans having higher network
centrality, such as those in the HMP, takes longer to be eliminated by the digestive enzymes. Thus,
the eigenvector centrality provides a measure of the crowding of the highly central glycans which
makes them difficult for endoglycosidase to access.

Influence of complex glycans on overall network topology

The degree of connectivity decreases almost throughout the network for the native
glycosylation (**SI Figure S4A**) compared to all-man9. The number of stable glycan-glycan
interactions at several glycan sites decreases because spatially proximal glycans avoid unfavorable
charge-charge interactions. For example, the uncharged glycans 234 and 355 takes up a position
central to all the neighboring charged sugars, increasing the distances between them, and the
charged N406 glycan buries itself in the middle of the surrounding high-mannose glycans (**Figure
4A**), clustering together with them. In fact, removal of this charged glycan N406 glycan can

increase the processing of the neighboring high-mannose moieties(Cao et al., 2018). Further

evidence for such alterations in glycan connectivity due to the presence of complex glycans can

be observed in the comparisons of the RMSF variations. A detailed analysis has been presented in

the SI and illustrated in **SI Figure S5**. Combination of the network node location differences and

RMSF variations suggest that the presence of the charged sialic acid at the tips of these complex

glycans can dictate the interactions between neighbors, increasing their structural variations if

surrounded by other charged glycans, or reducing them if a stable conformation buried between

uncharged high-mannose patches can be found. This overall decrease in connectivity slightly

increases the diameter of the native network, increasing the mean number of hops for the shortest

path from 2.4 hops (and distance 0.19) in the all-man9 model to 3 hops (and distance 0.27).

The distribution of centrality also shifts between the two glycosylation models (**SI Figure**

S4B). The centralities of the glycans at 137, 234, 355, 398, 406, 462 and those at gp41 increase

due to native glycosylation. On the other hand, the V1 and V2 loop glycans as well as 262, 295,

363 decrease from the all-man9 model. **Figure 5A** illuminates the difference in adjacency matrices

between the two models, with blue color indicating decrease in edge weight, and red indicating

increase in edge weight in native network, as compared to all-man9. Overall connectivity goes

down in the apex and the central patch for the native, and increases within gp41 glycans, as

compared to all-man9. The differences between the two networks (with at least 10% edge-weight

change) are shown in **Figure 5B** and **C**. The pathways around 197, as well as the inter-protomer

interactions involving 156, 185e and 637 become stronger in native glycosylation. (**Figure 5B**).

Some of these 197 connections, such as those with 276 and the V5 loop glycans occur across the

CD4 binding site. It was previously shown that in predominantly high-mannose Env structures,

CD4bs targeting antibody VRC01 has very little interaction with glycans 197 and 276(Stewart-

Jones et al., 2016). Conversely, in another Env structure with fully processed native glycans, a

VRC01-like antibody called IOMA interacted extensively with both 197 and 276(Gristick et al.,

2016). This matches our observation of increased orientation of glycan 197 over the CD4bs in

native glycosylation pattern.   However, the paths connecting 137, 156, 301, and 197 (of

neighboring protomer) are significantly weakened in the native model (**Figure 5C**). Subnetworks

involving V4 loop glycans and some of the central mannose patch become weaker due to the

presence of native glycans.

Community analysis to determine breaches in glycan shield

In order to interact and neutralize the Env protein, antibodies look for breaches in the shield
where glycan densities are sparse. This information of relative sparsity is encoded within the
glycan network, in terms of edge clustering. We examine this clustering of glycan interactions
within the network by defining communities using a modularity maximization approach that
divides the network into sub-modules or groups (see Methods). Communities have high density of
edges between the nodes within a module and comparatively sparse edges connecting different
modules (**Figure 5D inset**). This analysis identifies five distinct communities within the BG505
native glycan network (**Figure 5D and E**). The apex glycans around loop V2 from the three
protomers together form a single community (1, green). Right below that, glycans 262, 295, 301,
332 and 448 (2, blue) forms a community around the V3 and alpha 2 helix. Glycans 133, 137,
197, 363, and 386 (3, yellow) and 339, 392, 398, 406 and 411 (4, red) form two distinct
communities involving the glycans on and surrounding V4 loop region. Rest of the glycans 88,
234, 276, 355 and 462 from gp120 and all four glycans from gp41 form the fifth community (5,
cyan), though the modularity value is lower due to sparser connections. The possibility of an
approaching probe reaching the protein surface through these strongly connected communities is
low. A similar study clustering glycans from microsecond simulation runs of BG505 SOSIP was
performed by Lemmin *et. al.*(Lemmin et al., 2017), identifying 4 glycan microdomains that
roughly correspond to our modules 1, 2, (3+4) and 5. However, in that study, mannose-5 glycoform
was used at all sites, and due to the smaller length of these glycans, some of the interactions,
including inter-protomer interactions were not observed. Remarkably, in that study, junctions
between microdomains were found to indicate regions of relative vulnerability. The communities
we identified, also demarcate the regions where the glycan shield can be penetrated. Broadly
neutralizing antibodies whose binding epitopes are known target these community boundaries
(**Figure 5D**). Thus, glycan community dynamics can help to determine susceptible regions of the
glycan shield, and can be further used for guiding immunological studies.

Quantifying the vulnerability of glycan shield for antibody-based neutralization

380   To capture the impact of the glycan shield acting as an immunological barrier, we
381 quantified the sugar barrier over the Env protein surface using the ensemble of structures that have

been generated. Towards this goal, we have defined the glycan encounter factor (GEF) as the number of glycan heavy atoms encountered by an external probe approaching the protein surface. This parameter is calculated at each residue on the surface of the protein within a probe of diameter 6Å calculated using our ensemble. The highest value obtained is 14 for the native model, and is located in the HMP. Based on existing structures, the main interaction points between Env and bNAbs are often hairpin-like loop regions. Even large-scale atomistic simulations suggest that the first line of contact between Env and an antibody is through a loop region(Schmidt et al., 2013). Accordingly, for our analysis, we have used a probe size of 6Å, which is the typical diameter of a hairpin loop. At each residue present on the surface of the protein, the approaching probe was considered in three directions (**Figure 6A**): perpendicular to the surface **z**, and then the **x** and **y** directions spanning the plane parallel to the surface. The geometric mean of the **x**, **y** and **z** values provide the GEF per residue on the Env surface. This value will go to zero when the glycan encounter factor is zero from any one of the three cardinal directions. Thus, for any point on the surface which has a dense glycan covering, such as D in **Figure 6B**, has a high glycan encounter factor value, versus a point such as R where the glycan covering is sparse, which will have a low GEF.

Previous evidence suggests that 70% of the Env ECD surface area is covered by glycans(Pancera et al., 2014). Based on this, we determined the lower cut-off of GEF below which we can define glycan holes. **Figure 6C** shows GEFs mapped onto the trimer protein surface. The calculated solvent-exposed surface area of the protein part of our modeled BG505 structure without considering glycans is 86,055 Å^2. Excluding the exposed region at the base of the soluble SOSIP, this reduces to 79,672 Å^2. Varying the lower cut-off of GEF, we find that a cut-off of 1.5 GEF leads to 30% of the surface to be exposed. Regions of BG505 surface having a GEF less than 1.5 is colored in black in **Figure 6D**. From the figure, it is clear that the typical glycan holes targeted by bNAbs in BG505, such as the CD4 binding site, the V3-loop epitope and the fusion peptide binding region fall below this GEF cut-off. GEF tracks with epitopes that are relatively generic to a broad range of Env strains(Burton et al., 2012; Xu et al., 2018).

Antibodies elicited by BG505.SOSIP.664 are mainly biased towards the missing 241 and 289 glycan holes (GH) and the cleft-of-trimer (COT) epitope regions as demonstrated by cryo-EM and immunogenicity assays(Bianchi et al., 2018; McCoy et al., 2016) (see Figure 1 in reference

(Bianchi et al., 2018)). The GEF parameter identifies these BG505-specific epitopes as breaches

in the glycan shield (**Figure 6B**, "R" region; **Figure 6D** yellow dashed circle). On the other hand,

the densely glycosylated regions around V2, V4 loop and alpha 2 helix have high values of GEF.

At each point, the GEF value is given by a combination of all glycans in the vicinity that can come

in the way of the approaching probe. At the same time, we can analyze the extent of influence of

each glycan on the protein surface. Thus, we are now equipped with a tool to quantify the barrier

effect of the glycans individually or as a group for immunogen design, as further

demonstrated(Bianchi et al., 2018; McCoy et al., 2016) in the following section.

Combining network topology and glycan encounter factor to inform on local and global effects of neutralization

The proposed network is useful in deciphering the impact of addition or deletion of glycan

on neutralization. Removal of the high conserved glycan at 197 by mutating the sequon leads to

enhanced neutralization sensitivity to a variety of CD4bs and V3-specific antibodies(Huang et al.,

2012; Li et al., 2008; Townsley et al., 2016). This glycan is situated proximal to the CD4 binding

site, and the tip of V3 loop, directly affecting the binding of these antibodies. However, past

experimental evidence also suggests that the deletion of this glycan at N197 increases the binding

affinity of two antibodies PG9 and PGT145 (Behrens et al., 2016) which target the trimer apex of

Env. The epitope of PG9 as determined from the PDB structure 5VJ6 include residues 160, 161,

167-173, 185, 305 and 307 (**Figure 7A**) and the epitope of PGT145 as determined from PDB

structure 5V8L include residues 123, 124, 127, 160-162, and 167-169 (**Figure 7A**). The footprint

of glycan 197 as per our GEF model is shown in **Figure 7B**. Residues of the antibody epitopes do

not overlap with those regions directly covered by glycan 197. Yet, removal of glycan 197 results

in significant reduction in the glycan encounter factor over the V2 antibody epitope as evidenced

in **Figure 7C** and **D**.

The removal of glycan 197 affects the glycans that were originally acting as barriers over

the epitopes to resample the available space in such a way that the barrier over the epitopes is now

reduced. Glycans 156, 160, 185e and 185h from the neighboring protomers directly shield the PG9

and PGT145 epitopes. Looking at the shortest paths of communication between residue 197 from

either of the protomers to all other glycans (**SI Figure S6A, B**) demonstrate that while the shortest

path between glycans 160 and 197 is direct, other glycans covering the epitope regions
communicate with 197 across neighboring protomers or via a series of inter-glycan interactions.
Deletion of 197 does not only affect the V1/V2 loop region glycans. The difference in adjacency
matrix due to this perturbation is illustrated in **SI Figure S7A**. There is a slight modification of the
community structure within the network as well, with glycan 392 leaving module 4 (red) and
entering module 3 (yellow) instead (**SI Figure S7B**). Interactions such as those between glycans
N88 – N625, N133 - N363, and interprotomer N156 – N185e are reduced more than 10%. On the
other hand, glycan 276, which was originally interacting strongly with 197 now forms new
interactions with N295, N332 and N411 glycans (**SI Figure S7C, D**).

Discussion

Dense arrangement of N-glycans masking antigenic surfaces on Env acts as a dynamic
shield from the adaptive immune system. Understanding the structure and dynamics of the glycan
shield as a whole is therefore important for characterizing the immunologically inert regions of
Env. While high-resolution X-ray, NMR and cryo-EM structures have supplied a number of
important molecular details about the Env glycoprotein, they do not account for the high flexibility
and dynamics of these glycans that leads to the glycan shield. Our integrated method for capturing
dynamics of the glycans beyond the core β-mannose, and quantification of the barrier resilience
and vulnerability as a function of glycosylation pattern can add new perspective to current HIV
vaccine design efforts.

Previous computational studies have generalized glycan moieties as simple oligo-
mannoses such as mannose-5 and mannose-9 (Ferreira et al., 2018; Lemmin et al., 2017; Liang et
al., 2016; Stewart-Jones et al., 2016; Yang et al., 2017) for the purpose of modeling. For the first
time, we have incorporated native glycosylation, and included complex sugars based on site-
specific mass spectrometry results. While the overall aspects of the network are consistent between
mannose-9 and native-like glycosylation, there are critical differences. The most noteworthy are a
consequence of the presence of the bulky fucose ring at the base or the negatively charged sialic
acids at the antennae tips. These lead to overall rearrangement of glycan orientations affecting its

microenvironment, ultimately influencing the shield topology. Thus, our comprehensive

characterization of the shield is capable of capturing individual glycan effects which are

physiologically and immunologically more relevant.

Our network-based approach illuminates the collective behavior of glycans. We compare

the relative centrality of glycans, identify potential interaction pathways and detect stable

communities. The centrality or importance of glycans correlates well with experimental cryo-EM

data(Amanat et al., 2018; Berndsen et al., 2019). Glycans with lower centrality have lesser

influence on the graph, and are the first to be eliminated from the network, if the adjacency

threshold is increased. Similarly, the most central region of the network is the most resilient to

enzymatic action. Such centrality measures can help determine the ease of targeting glycans and

their modifications, guiding the process of immunogen development in the context of distinct

Envs. While the shield resilience is the main consideration for epitope exposure, some antibodies

have also evolved to capitalize on specific interactions with a number of the conserved glycans

during their molecular engagement with the epitope. Due to the stability of glycan interactions

within the detected communities, this help determine the antibody angle of approach that has been

shown to influence the breadth of bNAbs (Moore and Williamson, 2016).

There are known common and uncommon 'glycan holes' that can open up in the shield to

make the virus more vulnerable, or conversely get covered resulting in immune escape, as a result

of evolutionary addition/deletion or shift of glycans(Wagh et al., 2018). While some advances

have been made previously in identifying these breaches based on the area of influence of each

glycan(Wagh et al., 2018), here we have found that this area can vary, depending on the glycan

type, charge, neighbors, etc. Amino acid signature analyses suggest that even minor perturbations

such as single site mutations could potentially change the shielding effect over certain

epitopes(Bricault et al., 2019). Therefore, we have derived a measure to quantify the shielding

effect based on the encounter of glycans over the antigenic protein surface. This tool allows us to

define regions of relative vulnerability and resilience in the glycan shield.

The method we developed here for the structural modeling of the glycoprotein atomistic

ensemble and the subsequent development of the network is high-throughput compared to

traditional sampling by MD simulations. Due to the ease of this fast and efficient pipeline we are

now equipped with a tool to perform comparative structural studies due to glycan additions,
deletions or modifications, as well as other variations in the Env sequence. This is important to
understand the evolution of the glycan shield over longitudinal sampling of lineages. The structural
basis of addition or removal of glycans that are known to drive antibody maturation and
neutralization activities(LaBranche et al., 2018; Umotoy et al., 2019) can also be easily
investigated. Capitalizing on the relatively low computational overhead, our pipeline can be
integrated into a polyclonal epitope mapping assay to track the glycan shield as a function of
hierarchical antibody response, by modeling the shield network with the presence of different
antibodies and observing how the topology is altered.

Beyond HIV-1 Env and other viral envelope proteins, the significance of glycoproteins in
a vast array of biological processes from protein folding, cellular communication to immune-
regulation, make them a fast-emerging field of interest in biomedical research. Changes in these
glycosylation patterns have been associated with various diseases, including rheumatoid
arthritis(Nakagawa et al., 2007), and cancers(Taniguchi and Kizuka, 2015). Additionally, many of
the current therapeutic antibodies in the market are N-linked glycoproteins, and the significance
of N-glycans is becoming increasingly evident (Dalziel et al., 2014). The ease of modeling of
glycan network utilizing our approach makes it translatable to other systems, and can assist in
determining the role of these glycans in conjugation with the underlying proteins at a molecular
level. In comparison to N-glycans, O-linked glycosylations are relatively more complex in terms
of structural variations and our understanding is limited. Generalization our methodology to
encompass O-glycosylated systems can provide molecular insights on such refractory systems,
like Ebola that have both N- and O-linked glycosylations, and explore the pathological
implications of dense O-glycans in mucin associated cancers(Bhatia et al., 2019).

In summary, HIV and the human immune system are at war constantly, and the virus uses
the Env glycan shield to mask the human immune surveillance. One battlefront in this war is the
glycan shield: while the virus evolves to develop resilience, the immune response counteracts by
looking for vulnerabilities. As efforts are underway to aid the immune system overcome this race
by conditioning it with engineered immunogens, there is a need to quantify the resistance and
vulnerability of glycan shield in a more quantitative manner. This is that first time that the native
glycan network and shielding has been spatially quantified. Our derivation of the Glycan

Encounter Factor measures the relative barrier over the Env protein surface, and can potentially
aid to distinguish subtle differences on the shield due to variations in the glycosylation or even
protein sequences. We are therefore armed with a set of validated *in silico* tools with which to help
the anti-HIV war efforts and guide immunogen design.

STAR Methods

Contact for Reagent and Resource Sharing

535 Further information and requests for resources and reagents should be directed to and will be
536 fulfilled by the Lead Contacts S. Gnanakaran (gnana@lanl.gov) and Andrew B. Ward
537 (andrew@scripps.edu).

Method Details

High-throughput conformational modeling

540 The ensemble of BG505 glycoprotein 3D conformations were built in atomistic detail by
541 implementing the ALLOSMOD(Guttman et al., 2013; Weinkam et al., 2012) package of
542 MODELLER(Eswar et al., 2006; Sali, 1995) [RRID: SCR_008395]. For ease of understanding,
543 the modeling protocol has been described as a flow schematic in **SI figure S8.** The protocol
544 consists of the following steps:

(a) ***Base template selection for protein scaffold:*** The BG505 protein scaffold was homology
modeled, by threading the protein sequence into available crystal structure templates.
Ligand-free SOSIP structure with PDB accession ID 4ZMJ(Kwon et al., 2015) was used
as the base template for gp120, that also guided the three-fold symmetry of the trimer.
5CEZ(Garces et al., 2015) was used as the base template for gp41 and since it has the least
number of missing residues among the available structures.

(b) ***Identifying flexible regions from available experimental PDB structures to guide
conformational sampling:*** There are a large number of unstructured flexible regions in the

553	protein. In order to incorporate the fluctuations as dictated by the available structural data
554	we devised the following protocol. We aimed to construct 10 distinct protein models for
555	these flexible regions as starting scaffolds for glycan building. In order to model these, 34
556	different BG505 SOSIP experimental structures were selected below a resolution cut-off
557	of 4Å (see **SI Table S2**). Open-like, chimeric and synthetic models were not considered.
558	The set of 34 were structurally aligned using PyMOL(Schrödinger, 2015) and their residue-
559	wise backbone RMSF was calculated (**SI Figure S9A**). The base-level of 0.1 was used as
560	a threshold in the normalized RMSF measure. The unstructured residues (coils and turns)
561	were identified on 4ZMJ and 5CEZ using the STRIDE program(Frishman and Argos,
562	1995). All the identified regions with RMSF higher than the threshold (**SI Figure S9A**)
563	were considered for modeling based on multiple templates. These include residue ranges
564	64 – 82, 133 – 153, 184 – 189, 298 – 303, 312 – 314, 324 – 329, 396 – 413, 425 – 431, 457
565	– 464, and 609 – 618 (in HXB2 numbering).

(c) ***Template selection for identified flexible regions:*** Some of these identified regions have missing residues in many of the PDBs (**SI Figure S9B**). Wherever enough models were available, 10 distinct conformations were randomly selected as templates. If this number 10 could not be reached for some regions (for example, in loops V2 and V4) due to missing residues (**SI Figure S9B**), those residues were modeled *ab initio*. All known disulphide bonds were kept as additional restraints. We made sure these short-listed regions for *ab initio* modeling were less than MODELLER-prescribed 14 residues in length. This is because, modeling of such long loops are usually considered to be error-prone(Fiser et al., 2000). The loop region 546-569 in gp41 is present only in the structure 5CEZ. We did not model this region *ab initio*, and used 5CEZ as the only template for all models. The 10 models thus built were further tested for proper stereochemistry, based on MODELLER optimization scores and PROCHECK scores(Laskowski et al., 1993). This results in different starting orientations of the hypervariable loops, within the experimentally determined limits of fluctuations.

(d) ***Modeling of glycans and their ensemble sampling:*** For each of the 10 selected protein structures, glycans were initially added with random orientation, at the known glycosylation sites, based on ideal geometries as dictated by CHARMM36(Best et al., 2012; Huang and MacKerell Jr, 2013) force field internal coordinates, followed by a 1Å

randomization added to the overall atomic coordinates as described by Guttman et.
al.(Guttman et al., 2013). Once all the glycans were added, ensuing refinement steps of the
glycoprotein system optimized an energy function given by a combination of template-
based spatial restraints, CHARMM36 forcefield terms, and a soft sphere-like truncated
Gaussian term to prevent collisions. The structures were relaxed with 1000 steps of
conjugate gradient minimization followed by a short molecular dynamics equilibration of
500ps. Further refining with five rounds of simulated annealing was performed between
1,300K to 300K in 8 steps. The glycans and the loop regions were kept flexible during the
refinement steps. 100 fully glycosylated structures were modeled from each of the 10
selected protein models, resulting in the final ensemble containing 1000 different poses.
Template restraints were removed from the hypervariable regions with glycans in order for
them to sample a wider range of conformations during the glycoprotein relaxation phase.
These residues by HXB2 numbering are as follows: 143 to 152 (V1 loop), BG505-specific
insert residues 185A to 185I and 186 to 189 (V2 loop), 309 to 315 and 325 to 329 (V3
loop), 400 to 410 (V4 loop) and 458 to 464 (V5 loop) as determined from the LANL HIV
database (https://www.hiv.lanl.gov/).

Glycan root mean square fluctuations and volume calculation

601   For both the native and all-man9 glycosylated models, the root mean square fluctuations
602   (RMSF) of each glycan (with index n) was calculated as an average over all its heavy atoms, by
603   the following equation:

$$RMSF_n = \frac{1}{K} \sum_{k=1}^{K} \sqrt{\frac{1}{M} \sum_{m=1}^{M} |\vec{r}_{mnk} - \langle \vec{r}_{nk} \rangle|^2}$$

where $\vec{r}_{mnk}$ is the atomic position of heavy atom k of glycan n in snapshot m, $\langle \vec{r}_{nk} \rangle =$
$(\frac{1}{M}) \sum_{m=1}^{M} \vec{r}_{mnk}$ is the average atomic position of heavy atom k in glycan n. K is the total number
of heavy atoms in the glycan. It is 127 for mannose-9, and varies depending on the type of glycan.
The ensemble for each model contains 1000 snapshots, making $M = 1000$ snapshots for each of

the two models. The standard deviations were obtained by dividing the 1000 snapshots into 4 sets

of M=250, and calculating the four sets of RMSF values **(SI Figure S5).**

The 1000 structures of the total ensemble are built from 10 initial starting protein

conformations, as described above. The main difference between these 10 conformations are the

variations in the loop regions, due to the missing residues in the templates. In order to remove the

effects of the loop fluctuations and consider the RMSF coming from the glycans alone, the reduced

RMSF values were also calculated in each of these 10 sub-models, and their average and s.d.

calculated (**SI Figure S5B**). The RMSF difference between the models **(SI Figure S5E)** were

obtained by subtracting the reduced RMSF values of all-man9 from native model (native minus

all-man9). Since, the average reduced RMSF value is ~4Å, only those RMSF difference values are

statistically significant which are above 0.2 Å, which corresponds to a p-value of 0.05, rejecting

those values within the null hypothesis. Ensemble of individual glycans were converted to atomic

densities at 1 Å gridsize in order to calculate the volume. Volume calculation was done using

CHIMERA package(Pettersen et al., 2004).

Glycan overlap and network analysis

The inter-glycan overlap is calculated as the total fraction of heavy atoms from the two

glycans that come within 5Å of each other. Let us consider the example of mannose-9 to illustrate

the parameter of overlap. A single mannose-9 glycan has 127 heavy atoms. Since our ensemble is

composed of 1000 possible structures, there are effectively 127,000 heavy atoms per ensemble of

mannose-9 at one position. The fraction of the total number of heavy atoms from two neighboring

ensembles that come within contact distance defines the overlap fraction. Since mannose-9 is the

most commonly occurring glycoform in our system, we have used it as our reference for

normalization of the overlap probability. An overlap greater than or equal to 50% of heavy atoms

from two neighboring mannose-9 glycans is assigned as 1. This overlap matrix is used to define

the adjacency matrix for our network analysis. Each glycan functions as a node of the graph

(Figure 4A inset), and two nodes are connected by an edge if there is at least 5% overlap as per

our overlap definition given above. The edge length is inversely proportional to the overlap value,

i.e., the larger the overlap, the closer two nodes (glycans) are in the graph. Only those glycans

from the neighboring protomers are considered, that have an inter-protomer edge. All graph theory

and network analyses were performed using Python(Rossum, 1995) [RRID: SCR_008394] and
Matlab_R2018a packages(MATLAB, 2018) [RRID: SCR_001622].

Eigenvector centrality calculation:

For a given graph, G, with adjacency matrix $\mathbf{A}=(a_{v,t})$ where $a_{v,t}$ is the edge weight
connecting nodes v and t ($a_{v,t}=0$ when there is no connection), the relative centrality score x, of
node v can be defined as:

644
$$x_v = \frac{1}{\lambda} \sum_{t \in N(v)} x_t$$

645 where $N(v)$ is the set of neighbors directly connected to v, and λ is a constant. From the definition
646 of the adjacency matrix where the elements go to zero if two nodes are not connected, the above
647 equation can be expressed as:

648
$$x_v = \frac{1}{\lambda} \sum_{t \in G} a_{v,t} x_t$$

This has the form of the eigenvector equation $\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$. With the added constraint of the
eigencentrality values needing to be non-negative, by the Perron-Frobenius theorem(Pillai et al.,
2005), the eigenvector corresponding to the largest eigenvalue gives the desired measure of
centrality. The eigenvector is a unit vector and therefore the centrality values add up to one. For
the purpose of this work, we have normalized the centrality values with respect to the node with
the highest centrality assigned at 1, to obtain the relative centrality values.

Modularity maximization for community detection

Community detection within the glycan network was performed using the modularity
maximization approach given by Newmann and Girvan(Newman, 2006; Newman and Girvan,
2004). Modularity Q is calculated as the difference between the fraction of edges that fall within
a module and the expected fraction if the edges were distributed in random.

$$Q = \sum_{i=1}^{k}(e_{ii} - a_i^2)$$

Where e_{ii} is the fraction of edges in module i; a_i is the fraction of edges with at least one end in

module i. It is calculated by a greedy heuristic, beginning with the trivial system of each node

being a cluster, and merging two clusters that will increase modularity by the largest value,

stopping when any further merge would decrease the modularity. This approach is known to work

well for small networks similar to our system. The calculations were implemented through a

standard algorithm(Blondel et al., 2008) in Matlab R2018a.

Visualization of Structures: All visualizations of structures for figures were generated using

the Visual Molecular Dynamics package(Humphrey et al., 1996) [RRID: SCR_001820].

Data Availability

671 All data needed to evaluate the conclusions in the paper are present in the paper and/or the

672 Supplementary Materials, and further deposited in the library:

673 http://dx.doi.org/10.17632/yds8rtfjjy.1

674 Additional data related to this paper may be requested from the authors.

Supplemental Information

676 Supplemental Information includes details of torsion angle distributions, flexibility and volume

677 sampling analysis of modeled glycan structures, along with 2 Tables and 9 Figures.

Acknowledgements

680 S.C., B.T.K and S.G were funded by grants from the NIH (Center for HIV/AIDS Vaccine

681 Immunology and immunogen Discovery, UM1 AI100645; Consortia for HIV/AIDS Vaccine

Development, UM1 AI144371). Z.T.W and A.B.W were supported by the National Institute of Allergy and Infectious Diseases grants: UM1 AI100663 and UM1 AI144462, and Collaboration for AIDS Vaccine Discovery (CAVD) grant OPP1115782. S.C was also partially supported by the Center for Nonlinear Studies (CNLS) at Los Alamos National Laboratory (LANL). We acknowledge Timothy Travers and Cesar Lopez for suggestions on glycan modeling, Kshitij Wagh for insights on glycosylation holes in HIV, Kevin Weihe and Rory Henderson from Duke University for helpful discussions on Env immune response. S.C and S.G acknowledge the LANL High Performance Computing Division for providing computational facilities.

Author Contributions

Conceptualization, S.C., Z.T.B, N.W.H., B.T.K., A.B.W. and S.G; Methodology, S.C., Z.T.B, N.W.H., A.B.W. and S.G.; Modeling and Simulations, S.C.; Cryo-EM experiments, Z.T.B.; Formal Analysis, S.C. and Z.T.B.; Visualization, S.C. and Z.T.B.; Manuscript Preparation, S.C., Z.T.B, N.W.H., B.T.K., A.B.W. and S.G.; Supervision, A.B.W. and S.G.

Competing Interests

The authors declare no competing interests.

References

Amanat, F., Duehr, J., Oestereich, L., Hastie, K.M., Ollmann Saphire, E., and Krammer, F. (2018). Antibodies to the Glycoprotein GP2 Subunit Cross-React between Old and New World Arenaviruses. mSphere. 3(3). DOI: 10.1128/mSphere.00189-18.
An, H.J., Froehlich, J.W., and Lebrilla, C.B. (2009). Determination of glycosylation sites and site-specific heterogeneity in glycoproteins. Curr Opin Chem Biol. 13(4), 421-426. DOI: 10.1016/j.cbpa.2009.07.022.
Barnes, C.O., Gristick, H.B., Freund, N.T., Escolano, A., Lyubimov, A.Y., Hartweger, H., West, A.P., Jr., Cohen, A.E., Nussenzweig, M.C., and Bjorkman, P.J. (2018). Structural characterization of a highly-potent V3-glycan broadly neutralizing antibody bound to natively-glycosylated HIV-1 envelope. Nat Commun. 9(1), 1251. DOI: 10.1038/s41467-018-03632-y.

Behrens, A.J., Vasiljevic, S., Pritchard, L.K., Harvey, D.J., Andev, R.S., Krumm, S.A., Struwe,
W.B., Cupo, A., Kumar, A., Zitzmann, N., et al. (2016). Composition and Antigenic Effects of
Individual Glycan Sites of a Trimeric HIV-1 Envelope Glycoprotein. Cell Rep. 14(11), 2695-
2706. DOI: 10.1016/j.celrep.2016.02.058.
Beleva Guthrie, V., Masica, D.L., Fraser, A., Federico, J., Fan, Y., Camps, M., and Karchin, R.
(2018). Network analysis of protein adaptation: Modeling the functional impact of multiple
mutations. Molecular biology and evolution. 35(6), 1507-1519.
Berman, H.M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T.N., Weissig, H., Shindyalov, I.N.,
and Bourne, P.E. (2000). The Protein Data Bank. Nucleic Acids Res. 28(1), 235-242. DOI:
10.1093/nar/28.1.235.
Berndsen, Z., Chakraborty, S., Wang, X., Cottrell, C., Torres, J., Diedrich, J.K., Lopez, C.A.,
Yates III, J.R., van-Gills, M.J., Paulson, J.C., et al. (2019). Visualization of the HIV-1 Env
Glycan Shield Across Scales. bioRxiv. 10.1101/839217.
Best, R.B., Zhu, X., Shim, J., Lopes, P.E., Mittal, J., Feig, M., and MacKerell Jr, A.D. (2012).
Optimization of the additive CHARMM all-atom protein force field targeting improved sampling
of the backbone ϕ, ψ and side-chain χ1 and χ2 dihedral angles. Journal of chemical theory and
computation. 8(9), 3257-3273.
Bhatia, R., Gautam, S.K., Cannon, A., Thompson, C., Hall, B.R., Aithal, A., Banerjee, K., Jain,
M., Solheim, J.C., Kumar, S., et al. (2019). Cancer-associated mucins: role in immune
modulation and metastasis. Cancer Metastasis Rev. 38(1-2), 223-236. DOI: 10.1007/s10555-018-
09775-0.
Bianchi, M., Turner, H.L., Nogal, B., Cottrell, C.A., Oyen, D., Pauthner, M., Bastidas, R.,
Nedellec, R., McCoy, L.E., Wilson, I.A., et al. (2018). Electron-Microscopy-Based Epitope
Mapping Defines Specificities of Polyclonal Antibodies Elicited during HIV-1 BG505 Envelope
Trimer Immunization. Immunity. 49(2), 288-300 e288. DOI: 10.1016/j.immuni.2018.07.009.
Blondel, V.D., Guillaume, J.-L., Lambiotte, R., and Lefebvre, E. (2008). Fast unfolding of
communities in large networks. Journal of statistical mechanics: theory and experiment.
2008(10), P10008.
Bradley, T., Fera, D., Bhiman, J., Eslamizar, L., Lu, X., Anasti, K., Zhang, R., Sutherland, L.L.,
Scearce, R.M., Bowman, C.M., et al. (2016). Structural Constraints of Vaccine-Induced Tier-2
Autologous HIV Neutralizing Antibodies Targeting the Receptor-Binding Site. Cell Rep. 14(1),
43-54. DOI: 10.1016/j.celrep.2015.12.017.
Bricault, C.A., Yusim, K., Seaman, M.S., Yoon, H., Theiler, J., Giorgi, E.E., Wagh, K., Theiler,
M., Hraber, P., Macke, J.P., et al. (2019). HIV-1 Neutralizing Antibody Signatures and
Application to Epitope-Targeted Vaccine Design. Cell Host Microbe. 25(1), 59-72 e58. DOI:
10.1016/j.chom.2018.12.001.
Burton, D.R., Ahmed, R., Barouch, D.H., Butera, S.T., Crotty, S., Godzik, A., Kaufmann, D.E.,
McElrath, M.J., Nussenzweig, M.C., Pulendran, B., et al. (2012). A Blueprint for HIV Vaccine
Discovery. Cell Host Microbe. 12(4), 396-407. DOI: 10.1016/j.chom.2012.09.008.
Burton, D.R., and Mascola, J.R. (2015). Antibody responses to envelope glycoproteins in HIV-1
infection. Nat Immunol. 16(6), 571-576. DOI: 10.1038/ni.3158.
Cao, L., Diedrich, J.K., Kulp, D.W., Pauthner, M., He, L., Park, S.R., Sok, D., Su, C.Y.,
Delahunty, C.M., Menis, S., et al. (2017). Global site-specific N-glycosylation analysis of HIV
envelope glycoprotein. Nat Commun. 8, 14954. DOI: 10.1038/ncomms14954.
Cao, L., Pauthner, M., Andrabi, R., Rantalainen, K., Berndsen, Z., Diedrich, J.K., Menis, S., Sok,
D., Bastidas, R., Park, S.R., et al. (2018). Differential processing of HIV envelope glycans on the

757 virus and soluble recombinant trimer. Nat Commun. 9(1), 3693. DOI: 10.1038/s41467-018-
758 06121-4.
759 Chang, V.T., Crispin, M., Aricescu, A.R., Harvey, D.J., Nettleship, J.E., Fennelly, J.A., Yu, C.,
760 Boles, K.S., Evans, E.J., Stuart, D.I., et al. (2007). Glycoprotein structural genomics: solving the
761 glycosylation problem. Structure. 15(3), 267-273. DOI: 10.1016/j.str.2007.01.011.
762 Coss, K.P., Vasiljevic, S., Pritchard, L.K., Krumm, S.A., Glaze, M., Madzorera, S., Moore, P.L.,
763 Crispin, M., and Doores, K.J. (2016). HIV-1 Glycan Density Drives the Persistence of the
764 Mannose Patch within an Infected Individual. J Virol. 90(24), 11132-11144. DOI:
765 10.1128/JVI.01542-16.
766 Crispin, M., and Doores, K.J. (2015). Targeting host-derived glycans on enveloped viruses for
767 antibody-based vaccine design. Curr Opin Virol. 11, 63-69. DOI: 10.1016/j.coviro.2015.02.002.
768 Crispin, M., Ward, A.B., and Wilson, I.A. (2018). Structure and Immune Recognition of the HIV
769 Glycan Shield. Annu Rev Biophys. DOI: 10.1146/annurev-biophys-060414-034156.
770 Dalziel, M., Crispin, M., Scanlan, C.N., Zitzmann, N., and Dwek, R.A. (2014). Emerging
771 principles for the therapeutic exploitation of glycosylation. Science. 343(6166), 1235681. DOI:
772 10.1126/science.1235681.
773 Davis, S.J., and Crispin, M. (2010). Solutions to the glycosylation problem for low-and high-
774 throughput structural glycoproteomics. In: Functional and Structural Proteomics of
775 Glycoproteins, (Springer), pp. 127-158.
776 Dennis, J.W., Granovsky, M., and Warren, C.E. (1999). Protein glycosylation in development
777 and disease. Bioessays. 21(5), 412-421. DOI: 10.1002/(SICI)1521-1878(199905)21:5<412::AID-
778 BIES8>3.0.CO;2-5.
779 Doores, K.J. (2015). The HIV glycan shield as a target for broadly neutralizing antibodies. The
780 FEBS journal. 282(24), 4679-4691.
781 Doores, K.J., Bonomelli, C., Harvey, D.J., Vasiljevic, S., Dwek, R.A., Burton, D.R., Crispin, M.,
782 and Scanlan, C.N. (2010). Envelope glycans of immunodeficiency virions are almost entirely
783 oligomannose antigens. Proc Natl Acad Sci U S A. 107(31), 13800-13805. DOI:
784 10.1073/pnas.1006498107.
785 Dwek, R.A. (1996). Glycobiology: Toward Understanding the Function of Sugars. Chem Rev.
786 96(2), 683-720.
787 Eargle, J., and Luthey-Schulten, Z. (2012). NetworkView: 3D display and analysis of
788 protein.RNA interaction networks. Bioinformatics. 28(22), 3000-3001. DOI:
789 10.1093/bioinformatics/bts546.
790 Eswar, N., Webb, B., Marti-Renom, M.A., Madhusudhan, M., Eramian, D., Shen, M.y., Pieper,
791 U., and Sali, A. (2006). Comparative protein structure modeling using Modeller. Current
792 protocols in bioinformatics. 15(1), 5.6. 1-5.6. 30.
793 Falkowska, E., Le, K.M., Ramos, A., Doores, K.J., Lee, J.H., Blattner, C., Ramirez, A., Derking,
794 R., van Gils, M.J., Liang, C.H., et al. (2014). Broadly neutralizing HIV antibodies define a
795 glycan-dependent epitope on the prefusion conformation of gp41 on cleaved envelope trimers.
796 Immunity. 40(5), 657-668. DOI: 10.1016/j.immuni.2014.04.009.
797 Ferreira, R.C., Grant, O.C., Moyo, T., Dorfman, J.R., Woods, R.J., Travers, S.A., and Wood,
798 N.T. (2018). Structural Rearrangements Maintain the Glycan Shield of an HIV-1 Envelope
799 Trimer After the Loss of a Glycan. Sci Rep. 8(1), 15031. DOI: 10.1038/s41598-018-33390-2.
800 Fiser, A., Do, R.K., and Sali, A. (2000). Modeling of loops in protein structures. Protein Sci.
801 9(9), 1753-1773. Published online 2000/10/25 DOI: 10.1110/ps.9.9.1753.
802 Floyd, R.W. (1962). Algorithm 97: shortest path. Communications of the ACM. 5(6), 345.

Frishman, D., and Argos, P. (1995). Knowledge-based protein secondary structure assignment.
Proteins. 23(4), 566-579. DOI: DOI 10.1002/prot.340230412.
Fruchterman, T.M., and Reingold, E.M. (1991). Graph drawing by force-directed placement.
Software: Practice and experience. 21(11), 1129-1164.
Garces, F., Lee, J.H., de Val, N., de la Pena, A.T., Kong, L., Puchades, C., Hua, Y., Stanfield,
R.L., Burton, D.R., Moore, J.P., et al. (2015). Affinity Maturation of a Potent Family of HIV
Antibodies Is Primarily Focused on Accommodating or Avoiding Glycans. Immunity. 43(6),
1053-1063. DOI: 10.1016/j.immuni.2015.11.007.
Go, E.P., Ding, H., Zhang, S., Ringe, R.P., Nicely, N., Hua, D., Steinbock, R.T., Golabek, M.,
Alin, J., Alam, S.M., et al. (2017). Glycosylation Benchmark Profile for HIV-1 Envelope
Glycoprotein Production Based on Eleven Env Trimers. J Virol. 91(9). DOI: 10.1128/JVI.02428-
16.
Gristick, H.B., von Boehmer, L., West, A.P., Jr., Schamber, M., Gazumyan, A., Golijanin, J.,
Seaman, M.S., Fatkenheuer, G., Klein, F., Nussenzweig, M.C., et al. (2016). Natively
glycosylated HIV-1 Env structure reveals new mode for antibody recognition of the CD4-
binding site. Nat Struct Mol Biol. 23(10), 906-915. DOI: 10.1038/nsmb.3291.
Guttman, M., Weinkam, P., Sali, A., and Lee, K.K. (2013). All-atom ensemble modeling to
analyze small-angle x-ray scattering of glycosylated proteins. Structure. 21(3), 321-331. DOI:
10.1016/j.str.2013.02.004.
Guvench, O., Mallajosyula, S.S., Raman, E.P., Hatcher, E., Vanommeslaeghe, K., Foster, T.J.,
Jamison, F.W., and MacKerell Jr, A.D. (2011). CHARMM additive all-atom force field for
carbohydrate derivatives and its utility in polysaccharide and carbohydrate–protein modeling.
Journal of chemical theory and computation. 7(10), 3162-3180.
Haynes, B.F., Moody, M.A., Alam, M., Bonsignori, M., Verkoczy, L., Ferrari, G., Gao, F.,
Tomaras, G.D., Liao, H.X., and Kelsoe, G. (2014). Progress in HIV-1 vaccine development. J
Allergy Clin Immunol. 134(1), 3-10; quiz 11. DOI: 10.1016/j.jaci.2014.04.025.
Huang, J., Kang, B.H., Pancera, M., Lee, J.H., Tong, T., Feng, Y., Imamichi, H., Georgiev, I.S.,
Chuang, G.Y., Druz, A., et al. (2014). Broad and potent HIV-1 neutralization by a human
antibody that binds the gp41-gp120 interface. Nature. 515(7525), 138-142. DOI:
10.1038/nature13601.
Huang, J., and MacKerell Jr, A.D. (2013). CHARMM36 all-atom additive protein force field:
Validation based on comparison to NMR data. Journal of computational chemistry. 34(25),
2135-2145.
Huang, L., Liao, L., and Wu, C.H. (2017). Evolutionary analysis and interaction prediction for
protein-protein interaction network in geometric space. PloS one. 12(9), e0183495.
Huang, X., Jin, W., Hu, K., Luo, S., Du, T., Griffin, G.E., Shattock, R.J., and Hu, Q. (2012).
Highly conserved HIV-1 gp120 glycans proximal to CD4-binding region affect viral infectivity
and neutralizing antibody induction. Virology. 423(1), 97-106. DOI:
10.1016/j.virol.2011.11.023.
Humphrey, W., Dalke, A., and Schulten, K. (1996). VMD: Visual molecular dynamics. J Mol
Graph Model. 14(1), 33-38. DOI: Doi 10.1016/0263-7855(96)00018-5.
Ilinykh, P.A., Santos, R.I., Gunn, B.M., Kuzmina, N.A., Shen, X., Huang, K., Gilchuk, P., Flyak,
A.I., Younan, P., Alter, G., et al. (2018). Asymmetric antiviral effects of ebolavirus antibodies
targeting glycoprotein stem and glycan cap. PLoS Pathog. 14(8), e1007204. DOI:
10.1371/journal.ppat.1007204.

Imberty, A., and Perez, S. (2000). Structure, conformation, and dynamics of bioactive
oligosaccharides: theoretical approaches and experimental validations. Chem Rev. 100(12),
4567-4588.
Julien, J.P., Cupo, A., Sok, D., Stanfield, R.L., Lyumkis, D., Deller, M.C., Klasse, P.J., Burton,
D.R., Sanders, R.W., Moore, J.P., et al. (2013). Crystal structure of a soluble cleaved HIV-1
envelope trimer. Science. 342(6165), 1477-1483. DOI: 10.1126/science.1245625.
Karsten, C.B., and Alter, G. (2017). The HIV-1 Glycan Shield: Strategically Placed Kinks in the
Armor Improve Antigen Design. Cell Rep. 19(4), 669-670. DOI: 10.1016/j.celrep.2017.04.033.
Kwon, Y.D., Pancera, M., Acharya, P., Georgiev, I.S., Crooks, E.T., Gorman, J., Joyce, M.G.,
Guttman, M., Ma, X., Narpala, S., et al. (2015). Crystal structure, conformational fixation and
entry-related interactions of mature ligand-free HIV-1 Env. Nat Struct Mol Biol. 22(7), 522-531.
DOI: 10.1038/nsmb.3051.
LaBranche, C.C., McGuire, A.T., Gray, M.D., Behrens, S., Kwong, P.D.K., Chen, X., Zhou, T.,
Sattentau, Q.J., Peacock, J., Eaton, A., et al. (2018). HIV-1 envelope glycan modifications that
permit neutralization by germline-reverted VRC01-class broadly neutralizing antibodies. PLoS
Pathog. 14(11), e1007431. DOI: 10.1371/journal.ppat.1007431.
Laskowski, R.A., MacArthur, M.W., Moss, D.S., and Thornton, J.M. (1993). PROCHECK: a
program to check the stereochemical quality of protein structures. Journal of applied
crystallography. 26(2), 283-291.
Lasky, L.A., Groopman, J.E., Fennie, C.W., Benz, P.M., Capon, D.J., Dowbenko, D.J.,
Nakamura, G.R., Nunes, W.M., Renz, M.E., and Berman, P.W. (1986). Neutralization of the
AIDS retrovirus by antibodies to a recombinant envelope glycoprotein. Science. 233(4760), 209-
212. DOI: 10.1126/science.3014647.
Lee, J.H., Ozorowski, G., and Ward, A.B. (2016). Cryo-EM structure of a native, fully
glycosylated, cleaved HIV-1 envelope trimer. Science. 351(6277), 1043-1048. DOI:
10.1126/science.aad2450.
Lemmin, T., Soto, C., Stuckey, J., and Kwong, P.D. (2017). Microsecond Dynamics and
Network Analysis of the HIV-1 SOSIP Env Trimer Reveal Collective Behavior and Conserved
Microdomains of the Glycan Shield. Structure. 25(10), 1631-1639 e1632. DOI:
10.1016/j.str.2017.07.018.
Lennemann, N.J., Rhein, B.A., Ndungo, E., Chandran, K., Qiu, X., and Maury, W. (2014).
Comprehensive functional analysis of N-linked glycans on Ebola virus GP1. MBio. 5(1),
e00862-00813. DOI: 10.1128/mBio.00862-13.
Li, Y., Cleveland, B., Klots, I., Travis, B., Richardson, B.A., Anderson, D., Montefiori, D.,
Polacino, P., and Hu, S.L. (2008). Removal of a single N-linked glycan in human
immunodeficiency virus type 1 gp120 results in an enhanced ability to induce neutralizing
antibody responses. J Virol. 82(2), 638-651. DOI: 10.1128/JVI.01691-07.
Liang, Y., Guttman, M., Williams, J.A., Verkerke, H., Alvarado, D., Hu, S.L., and Lee, K.K.
(2016). Changes in Structure and Antigenicity of HIV-1 Env Trimers Resulting from Removal of
a Conserved CD4 Binding Site-Proximal Glycan. J Virol. 90(20), 9224-9236. DOI:
10.1128/JVI.01116-16.
Lütteke, T., Frank, M., and von der Lieth, C.-W. (2005). Carbohydrate Structure Suite (CSS):
analysis of carbohydrate 3D structures derived from the PDB. Nucleic acids research.
33(suppl_1), D242-D246.

Lyumkis, D., Julien, J.P., de Val, N., Cupo, A., Potter, C.S., Klasse, P.J., Burton, D.R., Sanders,
R.W., Moore, J.P., Carragher, B., et al. (2013). Cryo-EM structure of a fully glycosylated soluble
cleaved HIV-1 envelope trimer. Science. 342(6165), 1484-1490. DOI: 10.1126/science.1245627.
MATLAB, 2018. MATLAB, Version R2018a. The MathWorks Inc Natick, MA.
McCoy, L.E., van Gils, M.J., Ozorowski, G., Messmer, T., Briney, B., Voss, J.E., Kulp, D.W.,
Macauley, M.S., Sok, D., Pauthner, M., et al. (2016). Holes in the Glycan Shield of the Native
HIV Envelope Are a Target of Trimer-Elicited Neutralizing Antibodies. Cell Rep. 16(9), 2327-
2338. DOI: 10.1016/j.celrep.2016.07.074.
Moore, P.L., and Williamson, C. (2016). Approaches to the induction of HIV broadly
neutralizing antibodies. Current Opinion in HIV and AIDS. 11(6), 569.
Moremen, K.W., Tiemeyer, M., and Nairn, A.V. (2012). Vertebrate protein glycosylation:
diversity, synthesis and function. Nat Rev Mol Cell Biol. 13(7), 448-462. DOI:
10.1038/nrm3383.
Mouquet, H., Scharf, L., Euler, Z., Liu, Y., Eden, C., Scheid, J.F., Halper-Stromberg, A.,
Gnanapragasam, P.N., Spencer, D.I., Seaman, M.S., et al. (2012). Complex-type N-glycan
recognition by potent broadly neutralizing HIV antibodies. Proc Natl Acad Sci U S A. 109(47),
E3268-3277. DOI: 10.1073/pnas.1217207109.
Nakagawa, H., Hato, M., Takegawa, Y., Deguchi, K., Ito, H., Takahata, M., Iwasaki, N.,
Minami, A., and Nishimura, S. (2007). Detection of altered N-glycan profiles in whole serum
from rheumatoid arthritis patients. J Chromatogr B Analyt Technol Biomed Life Sci. 853(1-2),
133-137. DOI: 10.1016/j.jchromb.2007.03.003.
Newman, M.E. (2006). Modularity and community structure in networks. Proceedings of the
national academy of sciences. 103(23), 8577-8582.
Newman, M.E., and Girvan, M. (2004). Finding and evaluating community structure in
networks. Physical review E. 69(2), 026113.
Pancera, M., Zhou, T., Druz, A., Georgiev, I.S., Soto, C., Gorman, J., Huang, J., Acharya, P.,
Chuang, G.Y., Ofek, G., et al. (2014). Structure and immune recognition of trimeric pre-fusion
HIV-1 Env. Nature. 514(7523), 455-461. DOI: 10.1038/nature13808.
Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., and
Ferrin, T.E. (2004). UCSF chimera - A visualization system for exploratory research and
analysis. Journal of Computational Chemistry. 25(13), 1605-1612. DOI: 10.1002/jcc.20084.
Pillai, S.U., Suel, T., and Cha, S. (2005). The Perron-Frobenius theorem: some of its
applications. IEEE Signal Processing Magazine. 22(2), 62-75.
Pritchard, L.K., Spencer, D.I., Royle, L., Bonomelli, C., Seabright, G.E., Behrens, A.J., Kulp,
D.W., Menis, S., Krumm, S.A., Dunlop, D.C., et al. (2015a). Glycan clustering stabilizes the
mannose patch of HIV-1 and preserves vulnerability to broadly neutralizing antibodies. Nat
Commun. 6, 7479. DOI: 10.1038/ncomms8479.
Pritchard, L.K., Vasiljevic, S., Ozorowski, G., Seabright, G.E., Cupo, A., Ringe, R., Kim, H.J.,
Sanders, R.W., Doores, K.J., Burton, D.R., et al. (2015b). Structural Constraints Determine the
Glycosylation of HIV-1 Envelope Trimers. Cell Rep. 11(10), 1604-1613. DOI:
10.1016/j.celrep.2015.05.017.
Ringe, R.P., Pugach, P., Cottrell, C.A., LaBranche, C.C., Seabright, G.E., Ketas, T.J.,
Ozorowski, G., Kumar, S., Schorcht, A., van Gils, M.J., et al. (2019). Closing and Opening
Holes in the Glycan Shield of HIV-1 Envelope Glycoprotein SOSIP Trimers Can Redirect the
Neutralizing Antibody Response to the Newly Unmasked Epitopes. J Virol. 93(4). DOI:
10.1128/JVI.01656-18.

Rossum, G. (1995). Python reference manual.

Sali, A. (1995). Comparative protein modeling by satisfaction of spatial restraints. Mol Med
Today. 1(6), 270-277.

Sanders, R.W., Derking, R., Cupo, A., Julien, J.P., Yasmeen, A., de Val, N., Kim, H.J., Blattner,
C., de la Pena, A.T., Korzun, J., et al. (2013). A next-generation cleaved, soluble HIV-1 Env
trimer, BG505 SOSIP.664 gp140, expresses multiple epitopes for broadly neutralizing but not
non-neutralizing antibodies. PLoS Pathog. 9(9), e1003618. DOI: 10.1371/journal.ppat.1003618.

Sanders, R.W., van Gils, M.J., Derking, R., Sok, D., Ketas, T.J., Burger, J.A., Ozorowski, G.,
Cupo, A., Simonich, C., Goo, L., et al. (2015). HIV-1 VACCINES. HIV-1 neutralizing
antibodies induced by native-like envelope trimers. Science. 349(6244), aac4223. DOI:
10.1126/science.aac4223.

Saphire, E.O., Schendel, S.L., Fusco, M.L., Gangavarapu, K., Gunn, B.M., Wec, A.Z.,
Halfmann, P.J., Brannan, J.M., Herbert, A.S., Qiu, X., et al. (2018). Systematic Analysis of
Monoclonal Antibodies against Ebola Virus GP Defines Features that Contribute to Protection.
Cell. 174(4), 938-952 e913. DOI: 10.1016/j.cell.2018.07.033.

Schmidt, A.G., Xu, H., Khan, A.R., O'Donnell, T., Khurana, S., King, L.R., Manischewitz, J.,
Golding, H., Suphaphiphat, P., Carfi, A., et al. (2013). Preconfiguration of the antigen-binding
site during affinity maturation of a broadly neutralizing influenza virus antibody. Proc Natl Acad
Sci U S A. 110(1), 264-269. DOI: 10.1073/pnas.1218256109.

Schrödinger, L.L.C. (2015). The PyMOL molecular graphics system, Version 1.8. Technical
Report.

Sethi, A., Tian, J., Derdeyn, C.A., Korber, B., and Gnanakaran, S. (2013). A mechanistic
understanding of allosteric immune escape pathways in the HIV-1 envelope glycoprotein. PLoS
Comput Biol. 9(5), e1003046. DOI: 10.1371/journal.pcbi.1003046.

Skjaerven, L., Yao, X.Q., Scarabelli, G., and Grant, B.J. (2014). Integrating protein structural
dynamics and evolutionary analysis with Bio3D. BMC Bioinformatics. 15, 399. DOI:
10.1186/s12859-014-0399-6.

Slynko, V., Schubert, M., Numao, S., Kowarik, M., Aebi, M., and Allain, F.H. (2009). NMR
structure determination of a segmentally labeled glycoprotein using in vitro glycosylation. J Am
Chem Soc. 131(3), 1274-1281. DOI: 10.1021/ja808682v.

Sommerstein, R., Flatz, L., Remy, M.M., Malinge, P., Magistrelli, G., Fischer, N., Sahin, M.,
Bergthaler, A., Igonet, S., Ter Meulen, J., et al. (2015). Arenavirus Glycan Shield Promotes
Neutralizing Antibody Evasion and Protracted Infection. PLoS Pathog. 11(11), e1005276. DOI:
10.1371/journal.ppat.1005276.

Stanley, P., Taniguchi, N., and Aebi, M. (2015). N-Glycans. In: Essentials of Glycobiology, rd,
A. Varki, R.D. Cummings, J.D. Esko, P. Stanley, G.W. Hart, M. Aebi, A.G. Darvill, T.
Kinoshita, N.H. Packer, et al. eds. (Cold Spring Harbor (NY)), pp. 99-111.

Stewart-Jones, G.B., Soto, C., Lemmin, T., Chuang, G.Y., Druz, A., Kong, R., Thomas, P.V.,
Wagh, K., Zhou, T., Behrens, A.J., et al. (2016). Trimeric HIV-1-Env Structures Define Glycan
Shields from Clades A, B, and G. Cell. 165(4), 813-826. DOI: 10.1016/j.cell.2016.04.010.

Szakonyi, G., Klein, M.G., Hannan, J.P., Young, K.A., Ma, R.Z., Asokan, R., Holers, V.M., and
Chen, X.S. (2006). Structure of the Epstein-Barr virus major envelope glycoprotein. Nat Struct
Mol Biol. 13(11), 996-1001. DOI: 10.1038/nsmb1161.

Taniguchi, N., and Kizuka, Y. (2015). Glycans and cancer: role of N-glycans in cancer
biomarker, progression and metastasis, and therapeutics. Adv Cancer Res. 126, 11-51. DOI:
10.1016/bs.acr.2014.11.001.

Townsley, S., Li, Y., Kozyrev, Y., Cleveland, B., and Hu, S.L. (2016). Conserved Role of an N-
Linked Glycan on the Surface Antigen of Human Immunodeficiency Virus Type 1 Modulating
Virus Sensitivity to Broadly Neutralizing Antibodies against the Receptor and Coreceptor
Binding Sites. J Virol. 90(2), 829-841. DOI: 10.1128/JVI.02321-15.
Umotoy, J., Bagaya, B.S., Joyce, C., Schiffner, T., Menis, S., Saye-Francisco, K.L., Biddle, T.,
Mohan, S., Vollbrecht, T., Kalyuzhniy, O., et al. (2019). Rapid and Focused Maturation of a
VRC01-Class HIV Broadly Neutralizing Antibody Lineage Involves Both Binding and
Accommodation of the N276-Glycan. Immunity. 51(1), 141-154 e146. DOI:
10.1016/j.immuni.2019.06.004.
Vankadari, N., and Wilce, J.A. (2020). Emerging WuHan (COVID-19) coronavirus: glycan
shield and structure prediction of spike glycoprotein and its interaction with human CD26.
Emerg Microbes Infect. 9(1), 601-604. Published online 2020/03/18 DOI:
10.1080/22221751.2020.1739565.
Wagh, K., Kreider, E.F., Li, Y., Barbian, H.J., Learn, G.H., Giorgi, E., Hraber, P.T., Decker,
T.G., Smith, A.G., Gondim, M.V., et al. (2018). Completeness of HIV-1 Envelope Glycan Shield
at Transmission Determines Neutralization Breadth. Cell Rep. 25(4), 893-908 e897. DOI:
10.1016/j.celrep.2018.09.087.
Walker, L.M., Phogat, S.K., Chan-Hui, P.Y., Wagner, D., Phung, P., Goss, J.L., Wrin, T., Simek,
M.D., Fling, S., Mitcham, J.L., et al. (2009). Broad and potent neutralizing antibodies from an
African donor reveal a new HIV-1 vaccine target. Science. 326(5950), 285-289. DOI:
10.1126/science.1178746.
Walker, L.M., Simek, M.D., Priddy, F., Gach, J.S., Wagner, D., Zwick, M.B., Phogat, S.K.,
Poignard, P., and Burton, D.R. (2010). A limited number of antibody specificities mediate broad
and potent serum neutralization in selected HIV-1 infected individuals. PLoS Pathog. 6(8),
e1001028. DOI: 10.1371/journal.ppat.1001028.
Walls, A.C., Tortorici, M.A., Frenz, B., Snijder, J., Li, W., Rey, F.A., DiMaio, F., Bosch, B.J.,
and Veesler, D. (2016). Glycan shield and epitope masking of a coronavirus spike protein
observed by cryo-electron microscopy. Nat Struct Mol Biol. 23(10), 899-905. Published online
2016/09/13 DOI: 10.1038/nsmb.3293.
Walsh, G. (2010). Biopharmaceutical benchmarks 2010. Nat Biotechnol. 28(9), 917-924. DOI:
10.1038/nbt0910-917.
Wang, C.C., Chen, J.R., Tseng, Y.C., Hsu, C.H., Hung, Y.F., Chen, S.W., Chen, C.M., Khoo,
K.H., Cheng, T.J., Cheng, Y.S., et al. (2009). Glycans on influenza hemagglutinin affect receptor
binding and immune response. Proc Natl Acad Sci U S A. 106(43), 18137-18142. DOI:
10.1073/pnas.0909696106.
Warshall, S., 1962. A theorem on boolean matrices, Journal of the ACM. Citeseer.
Weinkam, P., Pons, J., and Sali, A. (2012). Structure-based model of allostery predicts coupling
between distant sites. Proc Natl Acad Sci U S A. 109(13), 4875-4880. DOI:
10.1073/pnas.1116274109.
Woods, R.J., Pathiaseril, A., Wormald, M.R., Edge, C.J., and Dwek, R.A. (1998). The high
degree of internal flexibility observed for an oligomannose oligosaccharide does not alter the
overall topology of the molecule. Eur J Biochem. 258(2), 372-386.
Xu, K., Acharya, P., Kong, R., Cheng, C., Chuang, G.Y., Liu, K., Louder, M.K., O'Dell, S.,
Rawi, R., Sastry, M., et al. (2018). Epitope-based vaccine design yields fusion peptide-directed
antibodies that neutralize diverse strains of HIV-1. Nat Med. 24(6), 857-867. DOI:
10.1038/s41591-018-0042-6.

Yang, M., Huang, J., Simon, R., Wang, L.X., and MacKerell, A.D., Jr. (2017). Conformational
Heterogeneity of the HIV Envelope Glycan Shield. Sci Rep. 7(1), 4435. DOI: 10.1038/s41598-
017-04532-9.
Zhang, M., Gaschen, B., Blay, W., Foley, B., Haigwood, N., Kuiken, C., and Korber, B. (2004).
Tracking global patterns of N-linked glycosylation site variation in highly variable viral
glycoproteins: HIV, SIV, and HCV envelopes and influenza hemagglutinin. Glycobiology.
14(12), 1229-1246. DOI: 10.1093/glycob/cwh106.
Zhang, X., Jia, R., Shen, H., Wang, M., Yin, Z., and Cheng, A. (2017). Structures and functions
of the envelope glycoprotein in flavivirus infections. Viruses. 9(11), 338.
1039

 Figures and Figure Legends

1043

Figure1: Native glycosylation and shielding of HIV-1 Env. (A). Schematic representation of N-
glycan types. The common core of Man3GlcNAc2Asn is indicated. The glycan species selected
for different sites of the BG505 native model (as given in **Table S1**). M9 ≡ mannose-9, FA2 ≡
fucosylated two-antennae, FA3 ≡ fucosylated three-antennae, FH ≡ fucosylated hybrid. Complex
glycans in gp41 are different from those in gp120, as per site-specific mass spectroscopy
experiments. (B) Single snapshot of modeled natively glycosylated BG505 structure, with each
glycan taking up a particular conformation. (C) Cumulative shielding effect over time due to the
dynamic nature of the glycans. 100 randomly selected models from the 1000-structure ensemble
is shown here. (D) Neighboring glycans within close proximity can sample overlapping volumes.
Glycan ensembles N88 shown in green and N625 shown in orange.

Figure 2: Comparison of glycan dynamics between experimental and simulated cryo-EM maps. (A) Experimental BG505_293F (top) and simulated BG505_native (bottom) cryo-EM maps sharpened and filtered at the global resolution (~3Å), with maps of de-glycosylated trimers displayed underneath to highlight the glycan density. The BG505_293F data is reproduced from (Berndsen et al., 2019) and density corresponding to the three RM20A3 Fabs bound to the base of the SOSIP were deleted for the sake of aesthetics (B) Noise threshold and map volume at the noise threshold as a function of Gaussian filter width (standard deviation - SD) of the BG505_native and de-glycosylated maps showing the progressive emergence of glycan signal at lower resolution. (C) 1.75 SD Gaussian filtered maps from panel A displayed at high and low isosurface thresholds. (D) Gaussian filtered BG505_native map with relaxed atomic model used for measuring local glycan dynamics along with a close up of a single glycan with a spherical probe around the BMA residue. **(E)** Bar plot showing relationship between the normalized site-specific mean local intensity from the BG505_293F (experimental) and BG505_native (simulated) maps along with the absolute difference at each site along with the Pearson correlation coefficient.

1070

Figure 3: Model and Experimental cryo-EM data reveals subtle change in orientation due to core fucose. (A) Difference in overall glycan orientation between all-man9 and fucosylated complex glycan in simulated model. The first core sugar and three residues at the base of the glycan (n-1, n, n+1; where n is the glycosylated asparagine residue) were aligned for a single sugar ensemble. Oligomannose samples a space symmetrically around the core, while, complex sugars have a distinct bend away from the side where the fucose ring is present. (B) Cryo-EM maps and refined atomic models for BG505_293F and BG505_293S (add citation) showing the 4 glycans from gp41. Only the core NAG residues can be confidently built into the sharpened maps. (C) Gaussian filtered BG505_293S map (gray) and BG505_293F-BG505_293S difference map (purple). (D) Close up of the glycan at N611 showing asymmetric difference signal around the glycan stem. (E) Full glycan stalks built and relaxed into both Gaussian filtered maps showing more pronounced change in projection angle.

1083

Figure 4: Spatial network of BG505 native glycosylation. (A) Visual representation of network in 2-dimensions, based on a force-directed layout. Each glycan forms a node-point on the graph, two nodes are connected by an edge (inset, from reference (Berndsen et al., 2019), Figure 8). The nodes are colored by the normalized eigenvector centrality of the glycans. (B) Shortest path of communication between glycan N332 and all other glycans in the network.

1089

Figure 5: Network difference and community analysis. (A) Difference in adjacency matrices between native and all-man9 models (native minus all-man9). Blue color indicates decrease in edge weight, and red indicates increase in edge weight in native network, as compared to all-man9. (B) Increase and (C) decrease in connectivity due to native glycosylation in comparison with all-man9 model (only those edges that change by 10% or more are shown). (D) 5 different communities were identified based on modularity maximization. The sub-community junctions identify susceptible regions in the shield where the shown HIV antibodies tend to bind. (E) Location of each community projected on the Env surface.

1098

Figure 6: Glycan Encounter Factor (GEF) for quantifying shielding effect. (A) At each residue
present on the surface of the protein, the approaching probe is considered in three directions **x, y**
and **z.** (B) Any point on the surface which has a dense glycan covering, such as D, has a high
glycan encounter factor value, versus a point such as R where the glycan covering is sparse, which
will have a low GEF. (C) Representation of GEF on Env surface, given by a colormap. Regions
of BG505 surface having a GEF less than 0.14 is colored in black (D). Typical known antibody
epitopes are indicated by arrows. BG505-specific GH and COT epitope region demarcated by
yellow dashed circle.

1107

Figure 7: Deletion of glycan N197 decreases glycan shielding at PG9 and PGT145 epitope.
(A) Top view of Env showing epitope regions of antibodies PG9 and PGT145 at the apex. (B)
Footprint of glycan N197 on Env surface, colored by glycan encounter factor contributed by N197
alone. (C) Glycan encounter factor over the PG9 and (D) PGT145 epitope residues. GEF decreases
for both the epitopes due to deletion of N197 glycan.