**1** **HIV-1 genome trafficking initiates APOBEC3G packaging in the cytosol**

3 Jordan T. Becker, Edward L. Evans III, Bayleigh E. Benner, Stephanie L. Fricke, Laura
4 E. Smith, Andra E. Bates, & Nathan M. Sherer*

6
7

McArdle Laboratory for Cancer Research, Institute for Molecular Virology, and Carbone
Cancer Center
University of Wisconsin – Madison
1525 Linden Drive, Madison, WI 53706

Short title: HIV-1 RNA genome regulates A3G trafficking

*To whom correspondence should be addressed: 501 Robert M. Bock Lab, 1525 Linden
Drive, Madison, WI 53706. Tel: (608) 890-2551. Email: nsherer@wisc.edu

**ABSTRACT**

HIV-1 RNA genomes interact with diverse RNA binding proteins in the cytoplasm

including antiviral factor APOBEC3G (A3G) that, in the absence of viral Vif proteins, is

packaged into virions. When and where genome-A3G interactions are initiated in the host

cell is unknown. Here we use quantitative long-term (>24 h) live cell fluorescence video

microscopy and a new in-cell RNA-protein interaction assay (the "IC-IP") to describe

subcellular viral and A3G trafficking behaviors over the entire HIV-1 productive phase.

Among other findings, we demonstrate that genome-A3G interactions are initiated in the

cytosol soon if not immediately after genome nuclear export; that A3G-genome

interactions are sufficiently strong so that tethering either factor to membranes inhibits

trafficking of the reciprocal binding partner; and that selective recognition of genomes

promotes consistent delivery of A3G to sites of virion assembly. Further elucidation of

RNA signature(s) detected by A3G may inform development of RNA-targeted antivirals.

**INTRODUCTION**

The human immunodeficiency virus type 1 (HIV-1) hijacks a diverse set of host cellular RNA binding proteins (RBPs) to carry out viral RNA transcription, nuclear export, translation, and trafficking[1–4]. Select host RBPs are packaged into virions and exhibit antiviral properties, with the best-characterized example being members of the Apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3 (APOBEC3) family. A subset of APOBEC3 proteins (A3F, A3G, and A3H) are packaged into virions to abolish infectivity by deaminating cytidines on the nascent minus-sense DNA strand of the viral genome, thereby generating G-to-A mutations in the DNA provirus[5–8]. During productive infection, however, APOBEC3 proteins are counteracted by viral infectivity factor (Vif) proteins that facilitate their proteasome-mediated degradation prior to the onset of virus particle production[9–11].

How A3G proteins is delivered to virions in the absence of Vif remains a poorly understood aspect of the cell-intrinsic host defense. HIV-1 virion assembly is driven by viral Gag capsid proteins multimerizing on an RNA scaffold consisting of host-derived RNA molecules and two dimerized viral RNA genomes, with Gag-RNA binding mediated by Gag's C-terminal Nucleocapsid (NC) domain [reviewed in [12]]. In the absence of viral genomes, Gag expression is sufficient to drive formation of non-infectious particles containing cellular RNAs, packaged in proportion to their relative abundance in the cell[13,14]. A3G's incorporation into virions has been shown, for most cell types, to be both RNA- and NC-dependent[15–20]. However, the relative contributions of genomes vs. host RNAs to this process remains controversial. On the one hand, A3G-RNA binding is relatively promiscuous, with A3G incorporated into virus particles even when packageable

genomes are not expressed[17,19]. On the other hand, A3G incorporation levels are

moderately enhanced when genomes are present[16,21]; and A3G has been shown to

exhibit selective RNA-binding characteristics[19,22] including a reported preference for G-

rich segments of the HIV-1 genome[20].

Where and when A3G interfaces with Gag and/or genomes in the cell has also been

under investigation for some time with conflicting results. At steady-state, A3G is

distributed throughout the cytosol (the aqueous phase of the cytoplasm) and accumulates

to high levels at non-membranous cytoplasmic sites of mRNA decay known as processing

bodies (PBs)[23,24]. Cytosolic A3G is also rapidly re-localized to sites of translational

repression known as stress granules (SGs) in response to heat shock or oxidative

stress[23,24]. Initial studies proposed a functional link between PBs and A3G's antiviral

activity[23,25,26]. By contrast, more recent work has shown visible PBs to have little to no

discernible relevance to the HIV-1 life cycle[27,28].

The dynamic and complex nature of cytoplasmic RNA trafficking during HIV-1 virion

genesis may have obscured consistent prior observations made in fixed cells,

emphasizing a need for real-time characterization the host RBP response to HIV-1

infection at subcellular resolution. Accordingly, herein we set out to comprehensively

define the behaviors of HIV-1 RNA genomes, Gag, A3G, and additional PB and SG

markers over the entire viral productive phase using long-term (>24h) live cell video

microscopy. Our results expose single-cell A3G degradation and trafficking behaviors in

the presence or absence of Vif; and show that HIV-1 infection has little to no net effect on

cytoplasmic PBs or SG formation. Fluorescence recovery after photobleaching (FRAP)

and a new in-cell RNA-protein interaction assay indicate that A3G-genome interactions

are strong, specific, Gag-independent, and initiated in a non-localized fashion throughout

the cytosol long before the onset of virus particle assembly. Importantly, A3G-genome

interactions were sufficiently strong so that artificially tethering A3G to membranes

sequestered genomes in the cytoplasm and inhibited HIV-1 virus particle production.

Finally, using single virion analysis, we demonstrate that while genomes are not essential

for A3G packaging, their presence promotes a more consistent per virion delivery of A3G

to sites of virus particle assembly. We discuss the potential implications of A3G's selective

recognition of RNA in the host cytosol, and how these principles might be exploited to

suppress genome trafficking and virus replication *in vivo*.

## RESULTS

**Tracking HIV-1 replication using a YFP-A3G biosensor cell line.** To study real-time RBP responses to HIV-1 in single cells over an entire round of viral replication, we generated "biosensor" HeLa cell lines stably expressing fluorescent protein-tagged versions of A3G (YFP-A3G) (Fig 1a) or relevant cytoplasmic RBPs including TIA1-YFP (a marker of SGs) (Fig. 2a) and CFP-DCP1A (a marker of PBs) (Fig. 2e)[24,29]. Cells were infected with either of two versions of an HIV-1$_{NL4-3}$ reporter virus modified to express cyan fluorescent protein (CFP) from the *nef* locus (as a marker of early viral gene expression); either; (1) Vif-competent (Vif+) virus; or (2) a virus rendered Vif-minus (Vifxx) due to insertion of two stop codons in the *vif* reading frame (see cartoon depiction in Fig 1b). In the absence of virus, YFP-A3G was localized to the cytoplasm and accumulated in bright foci consistent with PBs, as anticipated (Fig. 1a, left panel). Induction of oxidative stress using 250 µM sodium arsenite (Ars)[30] caused YFP-A3G to accumulate in SGs, confirming an intact stress response (Fig. 1a, right panel). We also confirmed that our YFP-A3G fusion protein was packaged into Vifxx virus particles and retained antiviral activity using single-round assembly and infectivity assays (Fig. 1c).

In our first set of live cell imaging experiments, we infected Hela.YFP-A3G cells with Vif+ virus at a MOI of <1 in order to visualize both infected and uninfected cells side-by-side in the same field. Viral gene expression (based on detection of CFP) was initiated 12-16 hours post-infection (hpi), followed by a gradual loss of YFP-A3G signal consistent with Vif-dependent YFP-A3G proteasomal degradation (decay rate of 24.5% / h +/-6.0%; n=33; Fig. 1d, Video 1, and quantification in Fig. 1e). We observed YFP-A3G to be depleted from both the diffuse cytosolic and PB pools simultaneously, with levels

stabilized at near-background with little to no recovery, even at the latest time points

(Fig.1d shows a 48h time point, with Fig.1e showing kinetics over the entire 48h time

course). We also observed cell rounding at the latest time points (Fig. 1d and Video 1,

see 48 h time point) consistent with real-time visualization of Vif-induced cell cycle

arrest[31]. Taken together, these movies validated our use of YFP-A3G as a proxy sensor

for detecting HIV-1 infection and tracking multivariate single-cell responses to Vif. Further,

they exposed Vif-A3G kinetic details including: (1) that Vif's capacity to degrade A3G is

potent in every single infected cell; (2) that Vif operates ubiquitously throughout the

cytoplasm; and (3) that Vif's effects are remarkably persistent on a per cell basis, *i.e.*, the

YFP-A3G biosensor demonstrated no discernible fluctuations to single-cell Vif activity

post-YFP-A3G degradation, even over time frames of greater than 24 h.

**HIV-1 infection in the absence of Vif re-localizes YFP-A3G to virion assembly sites**

**at the plasma membrane.** We next visualized YFP-A3G responses after HeLa.YFP-A3G

infection with Vif-deleted (Vifxx) virus. As expected, YFP-A3G expression was maintained

in the cytoplasm and at PBs throughout the entire course of infection, and cell cycle arrest

was not observed (Fig. 1f and Video 2). Strikingly, however, in infected cells we observed

YFP-A3G coalescing in large (up to 4 µm diameter) clusters at 30-48 hpi (Fig. 1f, white

arrows highlight growing clusters). These clusters were unlikely to be PBs or SGs based

on their proximity to the cell surface and real-time observations of cluster release and

transfer to neighboring cells (Fig. 1f, 32h time point, and tracked in Video 2). Instead, we

hypothesized that clusters represented recruitment of YFP-A3G from the cytoplasm to

sites of virus particle assembly at the plasma membrane. Consistent with this hypothesis,

YFP-A3G clusters stained positive for both HIV-1 Gag/Gag-Pol and viral genome but not

*YFP-A3G* mRNA (as a negative control) detected using a 4-color combined anti-p24$_{Gag}$

immunofluorescence (IF) and RNA fluorescence in situ hybridization (FISH) protocol (Fig.

1g). Thin section electron microscopy also confirmed clustering of virus particles at the

surface of HeLa.YFP-A3G cells at 48 hpi for both the Vif+ and Vifxx conditions (Fig. 1h).

Based on these results, we concluded that even in the absence of Vif, YFP-A3G serves

as a useful proxy biosensor for detecting HIV-1 virus particle assembly through monitoring

YFP-A3G's virus-driven translocation from the cytoplasm to the plasma membrane.

**HIV-1 replication has little to no effect on PB integrity or SG formation.** Because

HIV-1 drastically alters YFP-A3G subcellular trafficking behaviors in both the presence

and absence of Vif expression, we thought it important to ascertain if HIV-1 impacted the

net integrity of A3G's major cytoplasmic sites of accumulation, *i.e.*, SGs and PBs. To

address this question, we monitored Vif+ and Vifxx HIV-1 infection of two biosensor cell

lines engineered to stably express either YFP-TIA1 (Fig. 2a-2c) or CFP-DCP1a (Fig. 2e);

validated markers of PBs and SGs, respectively [reviewed in 29]. We first confirmed that

our YFP-TIA1 "stress" biosensor responded to stress similar to YFP-A3G (see Fig. 1a) by

treating cells with 250 µM Ars [30] (Fig. 2a, right panel) or infecting cells with rhinovirus A16

(RVA16), a positive-strand RNA virus we had studied previously[32] and discovered to

cause rampant SG formation as early as 4h post-infection (Fig. 2b and Video 3).

Unlike RVA16, neither Vif+ nor Vifxx HIV-1 triggered SG formation in either YFP-

TIA1 or YFP-A3G cells over 48h of continuous imaging, a time window sufficient to

encompass an entire round of viral replication (Fig. 2c with quantification in 2d). HIV-1

also had no discernible effects on PB number or morphology in CFP-DCP1A biosensor

cells; with PB number being remarkable consistent and stable for all individual cells and

conditions (typically 6-9 PBs per cell, Fig. 2e with quantification in 2f). Based on these

largely negative results, we concluded that despite HIV-1's marked effects on YFP-A3G

as shown in Fig. 1, net modulation of PBs or induction of SGs are unlikely to be intrinsic

features of the HIV-1 replication cycle.

Because HIV-1 Gag was recently reported to inhibit SG formation[33,34], we also

tested the effects of treating HeLa.YFP-A3G or HeLa.YFP-TIA1 biosensor cells with Ars

at 31 hpi when NL4-3 Gag expression in HeLa cells is relatively high (*e.g.*, see [31]).

Contrary to our expectations, both YFP-A3G and YFP-TIA1 accumulated in SGs in Ars-

treated cells infected with either Vif+ or Vifxx viruses (Fig. 2g with quantification in 2h). To

localize Gag and genome under these conditions, we again performed combined Gag IF

and genome FISH, finding that the bulk of the HIV-1 genome signal co-localized with

YFP-A3G in Ars-induced SGs at this time point (Fig. 2i, *gag-pol* FISH). By contrast, the

cytoplasmic pool of Gag was largely excluded from these complexes (Fig. 2i, anti-p24).

SGs are induced in response to protein kinase R-mediated phosphorylation of

translation initiation factor eIF2a, triggering aggregation of scaffolding proteins that

include TIA1, TIAR, and G3BP1 in complex with mRNAs bound to translation initiation

factors and polysome-associated proteins [reviewed in [29,35]]. Accordingly, co-localization

of YFP-A3G with genomes in SGs could suggest stronger recognition of genomes by

YFP-A3G relative to Gag in association with free ribosomes in the cytosol.

**HIV-1 RNA genomes but not Gag modulate A3G subcellular mobility; evidence for genome-A3G interactions in the cytosol.** So that we could also monitor Gag and genome trafficking dynamics in living HeLa.YFP-A3G cells, we next engineered two-color, HIV-1 genomes that encoded CFP-tagged Gag from a "self-tagging" genome bearing 24 copies of MS2 RNA binding loop (located in the *gag-pol* open reading frame; ORF) and expressing an RFP-tagged MS2 bacteriophage (MS2-RFP) coat protein from the viral *nef* locus (Gag-CFP/MS2-RFP virus, see cartoon depictions in Figs. 3a and 3c). Gag-CFP allowed for single cell measurements of viral late gene expression, with genomes tracked based on their binding to the MS2-RFP protein. Expression of Vif+ or Vifxx Gag-CFP/MS2-RFP viruses recapitulated effects seen during infection with CFP reporter viruses (Figs. 1 and 2); including rapid, Vif-dependent down-regulation of YFP-A3G (decay rate = 25.9% per hour +/- 7.0%; n = 31, Fig. 3b, green, with quantification in Figs. 3e and 3f, Video 4) and, for Vif-deficient (Vifxx) conditions, co-clustering of YFP-A3G, MS2-RFP-tagged genomes, and Gag-CFP to assembly sites at the plasma membrane (Fig. 3d, arrows, Video 5; and see Fig. 4d for an image recorded at higher magnification). We observed Vif-mediated YFP-A3G degradation to occur prior to the onset of virus particle assembly (Fig. 3b, compare 2 h and 3 h time points), consistent with a prior report[36], but at >1h after genomes had populated the cytoplasm (Fig. 3d, compare 1h and 4h time points).

Both prior to and during virus particle assembly, Vifxx genomes and YFP-A3G were co-distributed in a non-localized fashion throughout the cytoplasm, with YFP-A3G but not genome or Gag-CFP accumulating at PBs (Fig. 3d, 2 and 3 h time points post-detection of MS2-RFP). To test if genomes and YFP-A3G were interacting in the fluid compartment,

we performed fluorescence recovery after photobleaching (FRAP) analysis to measure

rates of YFP-A3G recovery with or without genomes and in the presence or absence of

Gag (Figs. 3g, 3h, and quantification in 3i). Infection with Vifxx HIV-1 mCherry reporter

virus reduced rates of YFP-A3G recovery ($t_{1/2}$ = 34.4 s +/- 5.3 s; $n$=8) relative to an

uninfected "YFP-A3G alone" control ($t_{1/2}$ = 27.0 s +/- 13.5 s; $n$=14). Interestingly, we also

observed incomplete recovery of YFP-A3G at the latest time points in infected cells (~75%

recovery compared to nearly 90% for untreated control cells Fig. 3i); suggesting virus-

induced immobilization of YFP-A3G [37].

Expression of genomes in the absence of Gag using a "genome only" Vifxx MS2-

RFP genome, wherein we mutated the *gag* start codon to abolish Gag synthesis (see Fig.

3g), yielded markedly slower rates of YFP-A3G recovery ($t_{1/2}$ = 70.8 s +/- 27.6 s; $n$=6)

relative to both the "YFP-A3G Alone" condition, and a "Gag Only" condition wherein we

expressed Gag-mCherry from an mRNA modified to lack 5' and 3' viral regulatory RNA

sequences and codon-optimized to further reduce potential contributions of viral *cis*-

acting RNA elements located in the *gag* ORF (COGag-mCh; $t_{1/2}$ = 30 s +/- 8.9 s; $n$=7)

(Fig. 3i). Taken together, because the expression of genomes but not Gag altered YFP-

A3G movements in the cytoplasm based on FRAP analysis, genome-YFP-A3G

interactions are likely initiated in the cytoplasm independently of virus particle assembly;

and potentially even prior to the onset of Gag synthesis.

**Genome-A3G interactions are selective, Gag-independent, and initiated prior to the**

**onset of virus particle assembly.** Because FRAP was unable to distinguish between

direct or indirect genome-A3G interactions, we next developed a single cell assay to allow

us to measure the strength and specificity of RNA-RBP interactions in the cytoplasm head

on; dubbed the In-Cell RNA-Protein Interaction Protocol (IC-IP) (depicted in Fig. 4a). The

IC-IP was based on the principle of immobilizing genomes at unnatural sites in the

cytoplasm (*e.g.*, membranes or the actin cytoskeleton) using an MS2-based tethering

strategy that we have described previously[38]. Should an RBP such as YFP-A3G be

strongly associated with the tethered genome, its localization will be also shifted with clear

evidence of RNA-RBP co-localization at the unnatural site of tethering.

For genome-A3G IC-IPs, MSL-bearing genomes served as bait for YFP-A3G, as

depicted in Fig. 4a. Genome retargeting was achieved by co-expressing "targeter" MS2

coat proteins that were modified to tether MSL-tagged genomes to either membranes

(using Src-MS2; bearing a 10 amino-acid membrane targeting signal) or the actin

cytoskeleton (using Lifeact-MS2; bearing a 17 amino acid F-actin targeting signal) (Fig.

4b). The Src-MS2 and Lifeact-MS2 targeters were fused to iRFP670 so that we could co-

visualize both the targeter and the genome (labelled with the self-tagging MS2-RFP

"tracker" protein, see cartoon depiction in Fig. 4c) without experiencing spectral overlap

(see Fig. 4d).

Co-expression of two-color Vifxx Gag-CFP/MS2-RFP genomes with the Src-MS2-

iRFP or Lifeact-MS2-iRFP targeters caused moderate re-localization of Gag-CFP

proteins from the cytosolic pool to intracellular vesicles (Fig. 4d, compare panels iv. and

ix.) or F-actin filaments (Fig. 4d, compare panels iv. to xiv.), respectively; consistent with

a subset of Gag proteins co-trafficking with its genome substrate in the cytosol (as we

have previously shown[38]). Remarkably, however, re-localization of YFP-A3G was more

striking, with >50% of the net YFP-A3G per cell fluorescent signal now associated with

intracellular vesicles or F-actin filaments (Fig. 4d, compare panels ii., viii., and xiii., and

see YFP-A3G regions-of-interest; ROI; with quantification in 4e and 4f). Consistent with

our results from FRAP analyses (Figs. 3g and 3h), a "genome only" bait was sufficient to

recruit >50% of YFP-A3G to intracellular vesicles as directed by Src-MS2, suggesting that

Gag plays no role in genome-A3G cytosolic interactions (Figs. 5a and 5b). By contrast, a

control MSL-bearing mRNA encoding codon-optimized Gag-CFP ("Gag Only") had no

effect on YFP-A3G cytoplasmic distribution (Figs. 5c and 5d). Collectively, these results

indicated that HIV-1 genome-A3G interactions in the cytoplasm are strong, specific, and

occur in the presence or absence of Gag.

A particularly useful feature of the IC-IP is that it could be coupled to video

microscopy in order to determine the timing of YFP-A3G-genome interactions timed to

the onset of viral gene expression. Live cell imaging of cells co-expressing Src-MS2-

iRFP670 with a "Genome Only" RNA bait demonstrated that YFP-A3G was relocalized to

vesicles within two hours of first detecting genome expression in the cytoplasm (Fig. 5e,

compare 1h and 2h time points, and see Video 6 for additional detail). Accordingly, this

experiment indicated that strong genome-A3G interactions are initiated concomitantly

with or very soon after genome nuclear export.

**Tethering A3G to membranes arrests HIV-1 genome trafficking and reduces virus**

**particle production.** We reasoned that if A3G's interactions with genomes were

sufficiently strong, then tethering A3G to membranes would, as a corollary, arrest HIV-1

genome mobility in the cytosol and abolish genome trafficking to sites of virus particle

assembly. To test this idea, we engineered a "reciprocal" IC-IP experiment (Fig. 6a)

wherein YFP-A3G was modified so that it would bind membranes due to its bearing the

same N-terminal Src-derived myristoylation signal we used for our Src-MS2 genome

retargeting constructs (Figs. 2b-2e). This fusion protein (Src-YFP-A3G) was co-

expressed with two-color (Gag-CFP/MS2-RFP) Vifxx (Fig. 6b) or a CFP-tagged variant

(Src-CFP-A3G) co-expressed with one-color (MS2-RFP) "Genome Only" Vifxx genomes

(Fig. 6c). As would be anticipated for a strong interaction, Src-FP-A3G expression

induced the marked clustering of MS2-RFP-tagged genomes to perinuclear vesicles, with

genomes co-localizing with Src-FP-A3G (Figs. 6b and 6c). To address the more relevant

scenario of infected cells, we also generated a cell line constitutively expressing Src-YFP-

A3G (HeLa.Src-YFP-A3G cells) and infected these cells with a Vifxx reporter virus (Fig.

6d). As expected, we observed genomes (and, to a lesser extent, Gag) co-clustering with

Src-YFP-A3G at cytoplasmic vesicles at 24 hpi, detected using combined IF/FISH (Fig.

6d, bottom, white arrows).

We hypothesized that arrest of genome subcellular trafficking due to Src-YFP-A3G

expression would reduce Gag synthesis and net virus particle production. Consistent with

this hypothesis, co-expression of Src-YFP-A3G or a Src-MS2 control with MSL-tagged 2-

color genome (Gag-BFP in this experiment) yielded a dose-dependent reduction to

cytosolic Gag level and net efficiency of virus particle release (Fig. 6e). Taken together,

these experiments indicated that A3G-genome interactions in the cytosol are sufficiently

strong that tethering A3G to membranes can effectively repurpose this protein as an

inhibitor of genome trafficking and net virion production.

**291** **The presence of genomes promotes more consistent per virion delivery of A3G to**

**292** **sites of virus particle assembly.** Combined, the above experiments suggested that

**293** A3G has evolved to preferentially recognize one or more HIV-1 genome RNA signatures.

**294** However, A3G is encapsidated by Gag into virus particles even in the absence of

**295** packageable genomes[17–19]. In an effort to better rationalize prior observations of A3G

**296** genome specificity vs. promiscuity, we performed a comparative single virion analysis

**297** (SVA) of A3G delivery into virus particles; based on a technique originally pioneered by

**298** the Hu and Pathak groups wherein fluorescently-labeled virus-like particles (VLPs) are

**299** harvested from HEK293T cells (that produce a greater quantity of particles relative to

**300** HeLa) and subjected to sub-micron quantitative multicolor fluorescence imaging to

**301** measure relative levels of per particle genome and/or A3G incorporation[39,40] (Fig. 7). For

**302** our analysis, we compared per particle levels of YFP-A3G encapsidation for four

**303** independent Gag/genome scenarios (depicted in Fig. 7a); "Gag-Only" mRNAs encoding

**304** codon-optimized Gag-CFP (COGag-CFP = "Gag-Only"), viral Vif+ or Vifxx 2-color HIV

**305** Gag-CFP/MS2-RFP genomes, or mRNAs encoding codon-optimized Gag-CFP wherein

**306** the NC RNA-binding domain was replaced by a leucine zipper (ΔNCzip) to abolish Gag-

**307** RNA binding (based on [41]). As expected, YFP-A3G packaging was only observed for

**308** COGag and the Vifxx HIV-1 conditions (Fig. 7b).

**309** Using SVA, Vif+ and Vifxx viruses exhibited high efficiency MS2-RFP incorporation

**310** (~89 and ~88% of Gag-CFP particles scoring positive for MS2-RFP, respectively), very

**311** consistent with a prior report of HIV-1 genome packaging efficiency[39]. Vifxx Gag-

**312** CFP/MS2-RFP particles exhibited a similar frequency of YFP-A3G incorporation (86% of

**313** total Gag-CFP particles) (Fig. 7c and quantification in 7d). However, for "Gag-Only"

(COGag-CFP) particles, the frequency of detectable YFP-A3G incorporation was lower

(57% efficiency), intermediate to Vifxx HIV-1 vs. the ΔNCzip "no-RNA" negative control

(Fig. 7c and quantification in 7d). COGag particles also exhibited reduced per virion levels

of YFP-A3G fluorescence when compared to full-length, Vifxx HIV (Fig. 7c and

quantification in 7e). In sum, these data indicated that, although genomes are not

essential for YFP-A3G delivery to virus particles, selective genome-A3G interactions

promote a more consistent and more enriched per virion delivery of A3G to virus particle

assembly sites.

**DISCUSSION**

Herein we studied the HIV-1 life cycle from the host cell perspective, combining

several complementary live cell imaging approaches with functional assays to

characterize the coordinated behaviors A3G, PB or SG proteins, and viral elements (Gag

and genome) over an entire round of viral replication (summarized in Fig. S1).

Our YFP-A3G biosensor allowed for indirect measurements of HIV-1 replication

kinetics in single cells and exposed A3G behaviors including; (1) that Vif-mediated A3G

degradation occurs from all subcellular pools simultaneously (Fig. 1d and Movie 1); (2)

that A3G suppression by Vif is remarkably stable and maintained over the entire late

phase of infection (lasting hours to days, see Figs. 1d, Movie 1, and 1e); and (3) that A3G

degradation is completed prior to the onset of virus particle assembly (very consistent

with prior report from Holmes *et al*.,[36] but shown here to be after genome nuclear export)

(Fig. 3b). In the absence of Vif, we could also monitor virus particle assembly, observing

large quantities of YFP-A3G being re-localized from the cytoplasm to the plasma

membrane during this process (Fig. 1f and Video 2). Taken together, on a technical level

there should be great utility in the general strategy of tracking RBP biosensors to detect

virus replication dynamics and study host cell responses (*e.g.*, see also RVA16-induced

stress in Fig. 2b). Moreover, now validated, HeLa.YFP-A3G cells should serve as useful

multivariate reporter system for further dissection of the viral and host machineries

governing A3G degradation and packaging.

Despite HIV-1's marked effects on YFP-A3G behaviors, we observed little to no

discernible effects of HIV-1 on SGs or PBs monitored using our YFP-TIA1 or CFP-DCP1A

biosensors, with or without Vif expression (Fig. 2). These movies reinforce that

modulation of PB number or architecture or SG induction is unlikely to be an intrinsic

feature of the HIV-1 life cycle[27,28]. Overall, the underlying significance of A3G's localization

to PBs and SGs remains is unknown; but predicted to spatially coupled A3G to RNP

complexes with active roles in viral mRNA surveillance (*i.e.*, PBs are sites miRNA-

dependent mRNA decay and SGs can suppress viral translation)[23,24,26]. In this context,

we speculate that all successful retroviruses and endogenous elements must adapt to

subvert, suppress, or avoid PB- or SG-associated activities. Indeed, HIV-1[42,43], other

retroviruses[44,45], and endogenous retroelements including LINEs[46] and yeast Ty

elements[47] are thought to exploit non-membranous RNP complexes related to PBs/SGs

in order to compartmentalize activities including mRNA translation and genome

packaging. Regarding suppression, intriguing recent work from Mouland and colleagues

has shown HIV-1 to actively inhibit SG formation, proposed to reduce the potential for

stress signaling triggered by Gag NC interacting with host RNA species[33,34]. While we did

not observe overt SG suppression in this study (Figs. 2g-i), the dynamics of this process

warrant additional investigation and it seems reasonable to posit that retroviruses prevent

SG formation to promote a fluid flow of genomes from the nucleus to polysomes and then

on to capsid assembly sites. Overall, however, our data support the notion that HIV-1's

general *modus operandi* is to largely avoid association with visible PBs or induction of

SGs.

At a more molecular level, our imaging assays also suggest that A3G is able to

selectively compete for genome binding in the cytosol. First, HIV-1 genomes but not Gag

rapidly accumulated at YFP-A3G-enriched SGs in infected cells after Ars treatment (Fig

2i). Second, based on FRAP analysis, the presence of genomes, but not Gag, reduces

A3G mobility in the cytoplasm (Fig. 3g and 3h). Third, and most convincingly, our IC-IP

experiments demonstrated that >50% of the cytosolic YFP-A3G is recruited to

membranes or F-actin by Src- or Lifeact-MS2-tethered genomes, respectively, (Fig. 4f);

effects that were far less evident for Gag-CFP; not observed in the absence of genomes

(Fig. 5d); and detected soon if not immediately after genome nuclear export (Fig. 5e). As

to *why* A3G competes for genome binding, single virion analyses (Fig. 7) revealed an

~90% frequency of A3G incorporation into genome-containing virions relative to "Gag-

only" particles wherein the frequency of A3G incorporation was less than 60%. This result

suggests to us that, although A3G-RNA binding is, overall, relatively promiscuous[19], it has

evolved to preferentially target genomes in a way that ensures consistent and efficient

delivery to virions, largely consistent with a model recently proposed by Bieniasz and

colleagues wherein Gag and A3G are adapted to compete for similar RNA sequences[20].

In this way, A3G maximizes its antiviral potential and thus necessitated evolution of the

Vif antagonist. Endeavors to further delineate the relevant protein constituents of A3G-

genome "surveillance" complexes, and to determine whether or not A3G exhibits

differential detection of genomes destined for packaging vs. translation (as *gag-pol*

mRNAs), are ongoing.

To summarize, our studies reinforce that the core, essential nature of genome

trafficking in the cytosol is diffusion in dynamic association with RNP complexes, with

some proteins (*e.g.*, A3G) more strongly associated with genomes than others. While we

have yet to define the RNA signature(s) that specify genome detection by A3G, our

observations may be informative in the context of recent CLIP-seq studies demonstrating

A3G's RNA-binding preference to be relatively sequence non-specific[19,20] coupled to

compelling recent work showing that HIV-1 genomes are selectively enriched in selective

post-transcriptional regulatory marks including $N_6$ methyladenosine (m6a)[48–50], 5-

methylcytosine (m5c), and 2'O-methylation[51]. Artificially tethering A3G to membranes

restricts HIV-1 genome trafficking and virus particle assembly (Fig. 6). Based on these

effects, we predict it will be feasible to design inhibitory molecules or biologicals capable

of achieving similar strong, selective inhibition of HIV-1 RNA trafficking, translation, and

genome packaging in the context of antiviral strategies.

**ACKNOWLEDGEMENTS**

We thank Randall Massey at the University of Wisconsin SMPH Electron Microscopy Facility for assistance with EM sample preparation. We are grateful to Michael Malim (King's College London), Chad Swanson, (King's College London), and Sarah Gallois-Montbrun (Institut Cochin) for plasmid reagents and advice. The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: HIV-1 p24 hybridoma (183-H12-5C) (from Bruce Chesebro)[52]. We thank Kelly Watters, Marchel Hill, and Ann Palmenberg for RVA16[32]. This study was supported by NIH grants RO1AI110221 and U54AI150470; the Wisconsin Partnership Program New Investigator Program (ID 2830); the Greater Milwaukee Foundation's Shaw Scientist Program; and the UW-Madison UW2020 Program to N.M.S. J.T.B. was supported by National Science Foundation Graduate Research Fellow Program grant DGE-1256259 and both a Research Competition Award and a Dissertation Completion Fellowship from the UW—Madison office of the Vice Chancellor of Research and Graduate Education. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation. E.L.E. III received support from NIH training grant CA009135 and an Advance Opportunity Fellowship from the UW-Madison SciMed/GRS program. B.E.B. was supported by National Science Foundation Graduate Research Fellow Program grant DGE-1256259 and NIH training grant GM008349. S.L.F. received support from a UW-Madison Hilldale Undergraduate Research Fellowship.

**AUTHOR CONTRIBUTIONS:**

JTB and NMS conceived the project, designed reagents, and established methods. JTB carried out all experiments and wrote the first version of the manuscript. ELE, BEB, SLF, LES, and AEB contributed reagents and generated preliminary data. JTB, ELE, and NMS processed and analyzed data. All authors contributed to the final revised manuscript and approved submission.

**METHODS**

**Cell culture, plasmids, and stable cell lines.** Human HeLa and HEK293T cell lines (obtained from ATCC) were cultured in DMEM (Sigma-Aldrich) supplemented with 10% fetal bovine serum (heat-inactivated, filter-sterilized), 1% L-glutamine, and 1% penicillin-streptomycin. HeLa.YFP-A3G, HEK293T.YFP-A3G, HeLa.Src-YFP-A3G, HeLa.YFP-TIA1, and HeLa.CFP-DCP1a were generated as previously described[38]. Briefly, YFP-A3G, Src-YFP-A3G, YFP-TIA1, and CFP-DCP1a cDNA[24] were inserted into a MIGR1-derived retroviral vector (pCMS28) upstream of sequence encoding an internal ribosome entry site (IRES) and a second reading frame encoding Puromycin-N-acetyltransferase[48]. High performance clones were selected by limiting dilution and maintained in 2µg/mL puromycin. YFP, YFP-A3G, and Src-YFP-A3G as well as CFP versions of construct were also generated using the pcDNA3.1 transient expression vector backbone. HIV-1 reporter virus plasmids were derived from the pNL4-3 molecular clone[53] bearing inactivating mutations in *env*, *vpr*, and expressing a green fluorescent protein (GFP) reporter from the *nef* locus (E-R-/GFP)[54]; with the GFP reporter replaced with either cerulean fluorescent protein (CFP) or mCherry (mCh). Vif-minus HIV-1 reporter virus plasmids (Vifxx CFP and Vifxx mCherry) were generated by changing *vif* codons 26 and 27 to stop codons[55] using overlapping PCR and inserted into Vif+ CFP and Vif+ mCherry plasmids. Two-color fluorescent HIV-1 reporter viruses (Gag-CFP/MS2-RFP) were generated by replacing the *gag* reading frame in Vif+ and Vifxx CFP reporter viruses with *gag-CFP* just upstream of a cassette encoding 24 copies of the MS2 bacteriophage RNA stem loop (MSL, a kind gift of Robert Singer, Albert Einstein University, New York, NY, USA)[56], and then subsequently replacing CFP with sequence encoding the MS2 coat protein fused to

mApple or mCherry and bearing an SV40 nuclear localization signal (MS2-RFP). Gag

was also fused to mTagBFP2 for select experiments (*e.g.*, Fig. 6e) in order to avoid

CFP/YFP cross-detection when immunoblotting for Src-YFP-A3G. Mutated versions of

full-length HIV-1 were generated using overlapping PCR as previously described[38]. Src-

MS2-CFP, Src-MS2-RFP, and Src-MS2-iRFP targeting constructs encoding an amino-

terminal membrane targeting signal derived from the Src kinase (MGSSKSKPKD) were

generated by overlapping PCR and subcloned into pcDNA3.1. mTurquoise2 (CFP) and

mApple (RFP) were gifts of Michael Davidson (Addgene # 54843 and 54747,

respectively). iRFP670 cDNA[57] was amplified from the ColorfulCell expression plasmid,

a gift of Pierre Neveu (Addgene plasmid # 62449)[58]. mTagBFP2[59] was a gift of Michal

Davidson (Addgene plasmid # 55302).

**Retroviral assembly and infectivity assays**. Cells at 30-40% confluency were

transfected with 2µg DNA in six well plates using polyethylenimine (PEI; #23966,

Polysciences Inc.). pcDNA3.1 or pBlueScript were used as empty vector controls. Culture

media were replaced at 24 hours post-transfection and cell lysates and supernatants were

harvested for immunoblot analysis at 48 hours as previously described[60](Sherer et al.

2011). Briefly, 1mL of harvested culture supernatant was filtered, underlaid with 20%

sucrose (w/v) in PBS, subjected to centrifugation at >21,000g for two hours at 4ºC, and

viral pellets were resuspended in 35µL dissociation buffer (62.5 mM Tris-HCl, pH 6.8,

10% glycerol, 2% sodium dodecyl sulfate [SDS], 5% β-mercaptoethanol). Cells were

harvested in 500µL radioimmunoprecipitation assay (RIPA) lysis buffer (10 mM Tris-HCl,

pH 7.5, 150 mM NaCl, 1 mM EDTA, 0.1% SDS, 1% Triton X-100, 1% sodium

deoxycholate), homogenized by passage through a 26G needle, subjected to

centrifugation at 1,500g for 20 minutes at 4ºC, and liquid supernatant fraction was

combined 1:1 with 2X dissociation buffer. Proteins were resolved by sodium dodecyl

sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to 0.2 μm

nitrocellulose membranes. Gag was detected using a mouse monoclonal antibody

recognizing HIV-1 capsid/p24 (183-H12-5C; 1:1000 dilution) from Dr. Bruce Chesebro

and obtained from the NIH AIDS Research and Reference Reagent Program (Bethesda,

MD, USA)[52] and anti-mouse secondary antibodies conjugated to an infrared fluorophore

(IRDye680LT, 1:10000 dilution, Li-Cor Biosciences) for quantitative immunoblotting. As

loading controls, heat shock protein 90A/B (HSP90) was detected using a rabbit

polyclonal antibody (H-114, 1:2500 dilution, Santa Cruz Biotechnology, Santa Cruz, CA,

USA) or glyceraldehyde 3-phosphate dehydrogenase (GAPDH) was detected using a

mouse monoclonal antibody (6C5, 1:2500 dilution, Santa Cruz Biotechnology) and anti-

rabbit or anti-mouse secondary secondary antibodies conjugated to an infrared

fluorophore (IRDye800CW, 1:7500 dilution, Li-Cor Biosciences). YFP-containing proteins

(*e.g.*, YFP-A3G) were detected using a rabbit polyclonal antibody recognizing GFP (FL

sc-8334, 1:1000 dilution, Santa Cruz Biotechnology) and anti-rabbit secondary antibodies

conjugated to an infrared fluorophore (IRDye800CW). For infectivity assays,

supernatants containing single-round infectious HIV-1 virions (pseudo-typed with VSV-G)

were filtered and then added to target HeLa cells in the presence of 10μg/mL polybrene.

Approximately 36 hours later, target HeLa cells were scanned using a BioTek Cytation5

to detect reporter CFP (455nm/510nm excitation/emission filter) expression following

infection.

**Microscopy, immunofluorescence, and FISH.** Cells were plated in 24-well glass-bottom plates (Eppendorf) or 8-well microslides (IBIDI) and transfected using PEI or infected with HIV-1 reporter viruses. Transfection mixes contained 1µg (24-well) or 333ng (IBIDI) total plasmid DNA, respectively. Infections were performed with VSV-G pseudo-typed HIV-1 reporter viruses produced in HEK293T cells and titered on HeLa cells to determine MOI of 0.5-1, thereby ensuring that 33-66% of the cells would be infected per experiment. Fixed-cell experiments were performed on a Nikon Ti-Eclipse inverted wide-field microscope (Nikon Corporation, Melville, NY, USA) using a 100x Plan Apo oil objective (numerical aperture [NA], 1.45). Transfected cells were fixed 24 – 32 hours post-transfection and infected cells were fixed 42 hours post-infection using 4% paraformaldehyde. If treated with 250 µM sodium arsenite (Sigma-Aldrich), drug or an equivalent volume of dimethyl sulfoxide were added to cell culture wells at approximately one hour prior to fixation and returned to incubation at 37ºC. Live-cell imaging experiments were also performed on a Nikon Ti-Eclipse inverted wide-field microscope using a 20x Plan Apo objective lens (NA, 0.75) with images acquired every 60 minutes (except where otherwise indicated) over a time course of 16 to 90 hours. Images were acquired using an ORCA-Flash4.0 CMOS camera (Hammatsu Photonics) and the following excitation/emission filter sets (nanometer ranges): BFP (402/455), CFP (430/470), YFP (510/535), mApple (555/605), mCherry (572/632), and iRFP (645/705). All images were processed and analyzed using FIJI/ImageJ2.

For fixed-cell experiments using FISH, cells were plated as described as above and previously[38]. At 42 hours post-infection or after 24 hours plated (for uninfected cells),

cells were washed, fixed in 4% formaldehyde, and permeabilized in 70% ethanol for at

least 4 hours at 4ºC. Custom Stellaris FISH probes were designed to recognize NL4-3

HIV-1 *gag-pol* reading frame nucleotides 386-4456 using the Stellaris RNA FISH Probe

Designer 4.1 (Biosearch Technologies, Inc.) available online. To detect *yfp* mRNAs, we

used a DesignReady Probe set specific to eGFP. Both probe sets were labeled with CAL

Fluor Red 590 Dye (Biosearch Technologies, Inc.). Samples were hybridized with the

*gag/gag-pol* probe set according to the manufacturer's instructions available online.

Simultaneous immunofluorescence to detect Gag used a mouse monoclonal antibody

specific to p24 (#24-2, a gift from Dr. Michael Malim). Imaging experiments were

performed as describe above on a Nikon Ti-Eclipse inverted wide-field microscope using

a 100x Plan Apo objective lens.

FRAP experiments were performed using a Nikon Ti-Eclipse inverted A1R+

resonant/galvano hybrid confocal line-scanning microscope. Images were captured using

a 20x Plan Apo objective lens (NA, 0.75) and a GaAsP multi-detector for 488 and 560nm

channels. YFP-A3G was imaged using the 488nm laser at a low arbitrary intensity and

photobleached using the laser's maximum intensity. MS2-RFP and COGag-mCherry

were imaged using the 560nm laser at a low arbitrary intensity. Cells were maintained in

an environmental chamber (Pathology Devices, Inc.) at 37ºC, 5% $CO_2$, and 50% humidity.

Cells were imaged every 30 seconds for 4 frames prior to photobleaching (3 rapid

ablations of cytoplasmic ROIs 10µm in diameter) followed by imaging every 5 seconds

for a minute, 15 seconds for four minutes, and 30 seconds for 5 minutes. This time frame

was sufficient for fluorescence recovery to reach a plateau. FRAP analysis was performed

using the FIJI plugin FRAP Profiler[61] that adjusts for incidental field photobleaching

outside the ROI.

Single virion analyses (SVA) were performed as previously described[39]. Virus

particles were produced as described above for western blotting in HEK293T.YFP-A3G

cells to maintain consistent levels of YFP-A3G. Filtered culture supernatants (750µL)

were purified by sucrose centrifugation as described above and resuspended in 100µL

1x PBS (Sigma-Aldrich). 24-well plates were pre-coated with 2% FBS diluted in 1x PBS

for at least 30 minutes, this solution was removed, and the resuspended concentrated

virus particles were added to wells. Images were acquired on a Nikon Ti-Eclipse inverted

wide-field microscope using a 100x Plan Apo objective lens (NA, 1.45). These images

were processed and analyzed using Analyze Particles plugin in FIJI/ImageJ2[62].

**Thin-section electron microscopy.** HeLa.YFP-A3G were infected with Vif+ CFP or

Vifxx CFP HIV-1 viruses and fixed at 48 hours for chemical processing as previously

described[63]. Samples were sectioned into 100nm slices and with sections collected on

copper thin-bar grids. Sections were observed with a Phillips CM120 transmission

electron microscope, and images were collected with a MegaView III (Olympus-SIS,

Lakewood, CO, USA) side-mounted digital camera. All images were processed and

analyzed using FIJI/ImageJ2[62].

**Statistics.** For assembly assays (Figs. 1b and 6e), results were obtained from three

biological replicates as defined as cells plated in six well plates transfected and processed

on separate days. Graphs plot the mean value with error bars representing the standard

deviation of the mean with the exception of Fig. 7e, a violin plot showing all data points

with the mean (solid line) and quartiles (dashed lines) indicated. All statistical

comparisons were carried out using the two-tailed Student's *t* test and performed using

Microsoft Excel or Graphpad Prism.

**REFERENCES**

1. Bolinger, C. & Boris-Lawrie, K. Mechanisms employed by retroviruses to exploit host
factors for translational control of a complicated proteome. *Retrovirology* **6**, 8 (2009).
2. Cochrane, A. W., McNally, M. T. & Mouland, A. J. The retrovirus RNA trafficking
granule: from birth to maturity. *Retrovirology* **3**, 18 (2006).
3. Karn, J. & Stoltzfus, C. M. Transcriptional and posttranscriptional regulation of HIV-1
gene expression. *Cold Spring Harb. Perspect. Med.* **2**, a006916 (2012).
4. Swanson, C. M. & Malim, M. H. Retrovirus RNA trafficking: from chromatin to
invasive genomes. *Traffic Cph. Den.* **7**, 1440–1450 (2006).
5. Harris, R. S. *et al.* DNA deamination mediates innate immunity to retroviral infection.
*Cell* **113**, 803–809 (2003).
6. Malim, M. H. APOBEC proteins and intrinsic resistance to HIV-1 infection. *Philos.*
*Trans. R. Soc. Lond. B. Biol. Sci.* **364**, 675–687 (2009).
7. Refsland, E. W. & Harris, R. S. The APOBEC3 family of retroelement restriction
factors. *Curr. Top. Microbiol. Immunol.* **371**, 1–27 (2013).
8. Sheehy, A. M., Gaddis, N. C., Choi, J. D. & Malim, M. H. Isolation of a human gene
that inhibits HIV-1 infection and is suppressed by the viral Vif protein. *Nature* **418**,
646–650 (2002).
9. Mariani, R. *et al.* Species-specific exclusion of APOBEC3G from HIV-1 virions by Vif.
*Cell* **114**, 21–31 (2003).
10. Sheehy, A. M., Gaddis, N. C. & Malim, M. H. The antiretroviral enzyme APOBEC3G
is degraded by the proteasome in response to HIV-1 Vif. *Nat. Med.* **9**, 1404–1407
(2003).
11. Stopak, K., de Noronha, C., Yonemoto, W. & Greene, W. C. HIV-1 Vif blocks the
antiviral activity of APOBEC3G by impairing both its translation and intracellular
stability. *Mol. Cell* **12**, 591–601 (2003).
12. Freed, E. O. HIV-1 assembly, release and maturation. *Nat. Rev. Microbiol.* **13**, 484–
496 (2015).
13. Eckwahl, M. J. *et al.* Analysis of the human immunodeficiency virus-1 RNA
packageome. *RNA N. Y. N* **22**, 1228–1238 (2016).
14. Rulli, S. J. *et al.* Selective and nonselective packaging of cellular RNAs in retrovirus
particles. *J. Virol.* **81**, 6623–6631 (2007).
15. Zennou, V., Perez-Caballero, D., Göttlinger, H. & Bieniasz, P. D. APOBEC3G
incorporation into human immunodeficiency virus type 1 particles. *J. Virol.* **78**,
12058–12061 (2004).
16. Khan, M. A. *et al.* Analysis of the contribution of cellular and viral RNA to the
packaging of APOBEC3G into HIV-1 virions. *Retrovirology* **4**, 48 (2007).
17. Cen, S. *et al.* The interaction between HIV-1 Gag and APOBEC3G. *J. Biol. Chem.*
**279**, 33177–33184 (2004).
18. Alce, T. M. & Popik, W. APOBEC3G is incorporated into virus-like particles by a
direct interaction with HIV-1 Gag nucleocapsid protein. *J. Biol. Chem.* **279**, 34083–
34086 (2004).
19. Apolonia, L. *et al.* Promiscuous RNA binding ensures effective encapsidation of
APOBEC3 proteins by HIV-1. *PLoS Pathog.* **11**, e1004609 (2015).
20. York, A., Kutluay, S. B., Errando, M. & Bieniasz, P. D. The RNA Binding Specificity

of Human APOBEC3 Proteins Resembles That of HIV-1 Nucleocapsid. *PLoS*
*Pathog.* **12**, e1005833 (2016).
21. Svarovskaia, E. S. *et al.* Human apolipoprotein B mRNA-editing enzyme-catalytic
polypeptide-like 3G (APOBEC3G) is incorporated into HIV-1 virions through
interactions with viral and nonviral RNAs. *J. Biol. Chem.* **279**, 35822–35828 (2004).
22. Gallois-Montbrun, S. *et al.* Comparison of cellular ribonucleoprotein complexes
associated with the APOBEC3F and APOBEC3G antiviral proteins. *J. Virol.* **82**,
5636–5642 (2008).
23. Wichroski, M. J., Robb, G. B. & Rana, T. M. Human retroviral host restriction factors
APOBEC3G and APOBEC3F localize to mRNA processing bodies. *PLoS Pathog.* **2**,
e41 (2006).
24. Gallois-Montbrun, S. *et al.* Antiviral protein APOBEC3G localizes to
ribonucleoprotein complexes found in P bodies and stress granules. *J. Virol.* **81**,
2165–2178 (2007).
25. Chable-Bessia, C. *et al.* Suppression of HIV-1 replication by microRNA effectors.
*Retrovirology* **6**, 26 (2009).
26. Nathans, R. *et al.* Cellular microRNA and P bodies modulate host-HIV-1 interactions.
*Mol. Cell* **34**, 696–709 (2009).
27. Izumi, T. *et al.* Mov10 and APOBEC3G localization to processing bodies is not
required for virion incorporation and antiviral activity. *J. Virol.* **87**, 11047–11062
(2013).
28. Phalora, P. K., Sherer, N. M., Wolinsky, S. M., Swanson, C. M. & Malim, M. H. HIV-1
replication and APOBEC3 antiviral activity are not regulated by P bodies. *J. Virol.* **86**,
11712–11724 (2012).
29. Ivanov, P., Kedersha, N. & Anderson, P. Stress Granules and Processing Bodies in
Translational Control. *Cold Spring Harb. Perspect. Biol.* **11**, (2019).
30. Kedersha, N. *et al.* Dynamic shuttling of TIA-1 accompanies the recruitment of
mRNA to mammalian stress granules. *J. Cell Biol.* **151**, 1257–1268 (2000).
31. Evans, E. L., Becker, J. T., Fricke, S. L., Patel, K. & Sherer, N. M. HIV-1 Vif's
Capacity To Manipulate the Cell Cycle Is Species Specific. *J. Virol.* **92**, (2018).
32. Watters, K. *et al.* Differential Disruption of Nucleocytoplasmic Trafficking Pathways
by Rhinovirus 2A Proteases. *J. Virol.* **91**, (2017).
33. Cinti, A., Le Sage, V., Ghanem, M. & Mouland, A. J. HIV-1 Gag Blocks Selenite-
Induced Stress Granule Assembly by Altering the mRNA Cap-Binding Complex.
*mBio* **7**, e00329 (2016).
34. Valiente-Echeverría, F. *et al.* eEF2 and Ras-GAP SH3 domain-binding protein
(G3BP1) modulate stress granule assembly during HIV-1 infection. *Nat. Commun.* **5**,
4819 (2014).
35. Protter, D. S. W. & Parker, R. Principles and Properties of Stress Granules. *Trends*
*Cell Biol.* **26**, 668–679 (2016).
36. Holmes, M., Zhang, F. & Bieniasz, P. D. Single-Cell and Single-Cycle Analysis of
HIV-1 Replication. *PLoS Pathog.* **11**, e1004961 (2015).
37. Ishikawa-Ankerhold, H. C., Ankerhold, R. & Drummen, G. P. C. Advanced
fluorescence microscopy techniques--FRAP, FLIP, FLAP, FRET and FLIM. *Mol.*
*Basel Switz.* **17**, 4047–4132 (2012).
38. Becker, J. T. & Sherer, N. M. Subcellular Localization of HIV-1 gag-pol mRNAs

Regulates Sites of Virion Assembly. *J. Virol.* **91**, (2017).

39. Chen, J. *et al.* High efficiency of HIV-1 genomic RNA packaging and heterozygote
formation revealed by single virion analysis. *Proc. Natl. Acad. Sci. U. S. A.* **106**,
13535–13540 (2009).

40. Desimmie, B. A. *et al.* APOBEC3 proteins can copackage and comutate HIV-1
genomes. *Nucleic Acids Res.* **44**, 7848–7865 (2016).

41. Accola, M. A., Strack, B. & Göttlinger, H. G. Efficient particle production by minimal
Gag constructs which retain the carboxy-terminal domain of human
immunodeficiency virus type 1 capsid-p2 and a late assembly domain. *J. Virol.* **74**,
5395–5402 (2000).

42. Reed, J. C. *et al.* HIV-1 Gag co-opts a cellular complex containing DDX6, a helicase
that facilitates capsid assembly. *J. Cell Biol.* **198**, 439–456 (2012).

43. Barajas, B. C. *et al.* Identifying the assembly intermediate in which Gag first
associates with unspliced HIV-1 RNA suggests a novel model for HIV-1 RNA
packaging. *PLoS Pathog.* **14**, e1006977 (2018).

44. Soto-Rifo, R. *et al.* HIV-2 genomic RNA accumulates in stress granules in the
absence of active translation. *Nucleic Acids Res.* **42**, 12861–12875 (2014).

45. Bann, D. V., Beyer, A. R. & Parent, L. J. A murine retrovirus co-Opts YB-1, a
translational regulator and stress granule-associated protein, to facilitate virus
assembly. *J. Virol.* **88**, 4434–4450 (2014).

46. Goodier, J. L., Zhang, L., Vetter, M. R. & Kazazian, H. H. LINE-1 ORF1 protein
localizes in stress granules with other RNA-binding proteins, including components
of RNA interference RNA-induced silencing complex. *Mol. Cell. Biol.* **27**, 6469–6483
(2007).

47. Larsen, L. S. Z. *et al.* Ty3 nucleocapsid controls localization of particle assembly. *J.*
*Virol.* **82**, 2501–2514 (2008).

48. Kennedy, E. M. *et al.* Posttranscriptional m(6)A Editing of HIV-1 mRNAs Enhances
Viral Gene Expression. *Cell Host Microbe* **19**, 675–685 (2016).

49. Lichinchi, G. *et al.* Dynamics of the human and viral m(6)A RNA methylomes during
HIV-1 infection of T cells. *Nat. Microbiol.* **1**, 16011 (2016).

50. Tirumuru, N. *et al.* N(6)-methyladenosine of HIV-1 RNA regulates viral infection and
HIV-1 Gag protein expression. *eLife* **5**, (2016).

51. Courtney, D. G. *et al.* Epitranscriptomic Addition of m5C to HIV-1 Transcripts
Regulates Viral Gene Expression. *Cell Host Microbe* **26**, 217-227.e6 (2019).

52. Chesebro, B., Wehrly, K., Nishio, J. & Perryman, S. Macrophage-tropic human
immunodeficiency virus isolates from different patients exhibit unusual V3 envelope
sequence homogeneity in comparison with T-cell-tropic isolates: definition of critical
amino acids involved in cell tropism. *J. Virol.* **66**, 6547–6554 (1992).

53. Adachi, A. *et al.* Production of acquired immunodeficiency syndrome-associated
retrovirus in human and nonhuman cells transfected with an infectious molecular
clone. *J. Virol.* **59**, 284–291 (1986).

54. Connor, R. I., Chen, B. K., Choe, S. & Landau, N. R. Vpr is required for efficient
replication of human immunodeficiency virus type-1 in mononuclear phagocytes.
*Virology* **206**, 935–944 (1995).

55. Simon, J. H. *et al.* The regulation of primate immunodeficiency virus infectivity by Vif
is cell species restricted: a role for Vif in determining virus host range and cross-

species transmission. *EMBO J.* **17**, 1259–1267 (1998).

56. Femino, A. M., Fay, F. S., Fogarty, K. & Singer, R. H. Visualization of single RNA
transcripts in situ. *Science* **280**, 585–590 (1998).

57. Shcherbakova, D. M. & Verkhusha, V. V. Near-infrared fluorescent proteins for
multicolor in vivo imaging. *Nat. Methods* **10**, 751–754 (2013).

58. Sladitschek, H. L. & Neveu, P. A. MXS-Chaining: A Highly Efficient Cloning Platform
for Imaging and Flow Cytometry Approaches in Mammalian Systems. *PloS One* **10**,
e0124958 (2015).

59. Subach, O. M., Cranfill, P. J., Davidson, M. W. & Verkhusha, V. V. An enhanced
monomeric blue fluorescent protein with the high chemical stability of the
chromophore. *PloS One* **6**, e28674 (2011).

60. Sherer, N. M. *et al.* Evolution of a species-specific determinant within human CRM1
that regulates the post-transcriptional phases of HIV-1 replication. *PLoS Pathog.* **7**,
e1002395 (2011).

61. Hardin, J. Imaging embryonic morphogenesis in C. elegans. *Methods Cell Biol.* **106**,
377–412 (2011).

62. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*
*Methods* **9**, 676–682 (2012).

63. Garcia-Miranda, P. *et al.* Stability of HIV Frameshift Site RNA Correlates with
Frameshift Efficiency and Decreased Virus Infectivity. *J. Virol.* **90**, 6906–6917
(2016).

 **FIGURE LEGENDS**

**Fig. 1. Tracking the HIV-1 replication cycle using a YFP-A3G biosensor cell line. a**

HeLa.YFP-A3G cells showing diffuse and punctate (processing bodies, PB, white arrows)

distribution of YFP-A3G at steady-state (left) and accumulation into stress granules (SGs,

magenta arrows) following arsenite treatment (right, Ars). **b** Schematic of Vif-competent

(Vif+) or deficient (Vifxx) HIV-1 NL4-3 Env-, Vpr-, expressing CFP as a reporter of

infection. **c** Single-round infection confirms YFP-A3G dependent restriction of HIV-1 in

the absence of Vif. 293T.YFP-A3G cells were transfected with the indicated HIV-1

reporter viruses with or without co-expression of a plasmid encoding Vif-mCherry. Cells

and virus particles ($n$=3) were harvested at 48 h post-transfection and analyzed by

quantitative immunoblot using anti-p24$_{Gag}$, anti-YFP, and anti-HSP90 (loading control)

antisera. **d** Representative images from long-term time-lapse imaging showing

degradation of YFP-A3G upon HIV-1 Vif+ CFP infection and eventual Vif-dependent

G2/M arrest. **e** Quantification of movies (IntDen) represented in d (Vif+, solid lines) and **f**

(Vifxx, dashed lines). Cells from 3 independent experiments ($n$=33) with ≥16 hours per

cell (1 image/hour). **f** Representative images from long-term time-lapse imaging of YFP-

A3G upon HIV-1 Vifxx CFP infection. Magenta boxes highlight zoomed regions of interest

(ROIs) shown below and white arrows highlight YFP-A3G clustering at PM. **g** Fixed cell

images show Gag and *gagpol* mRNA (viral RNA genomes) accumulating with YFP-A3G

at cell periphery. White arrows highlight cell-peripheral clusters of YFP-A3G, Gag, and

genome. **h** TSEM images showing virus particles clustering in the extracellular space

after budding from HeLa.YFP-A3G cells infected with either Vif+ or Vifxx viruses. All scale

bars in fluorescent images = 10μm and in TSEM = 100nm. Error bars represent standard

deviation of the mean.

**Fig. 2. HIV-1 infection has little to no impact on P-bodies or stress granules**. **a**

HeLa.YFP-TIA1 cells showing steady-state distribution (vehicle, left) and accumulation

into SGs (magenta arrows) following arsenite treatment (right, Ars). **b** Representative

images of time-lapse imaging showing induction of SGs during infection by RVA16

(MOI=10). **c** Time-lapse images showing no effect on YFP-TIA1 distribution following

Vifxx CFP infection, quantified in **d** for >400 HeLa.YFP-A3G (black bars) and HeLa.YFP-

TIA1 (gray bars) cells per condition, combined from three independent experiments. **e**

Time-lapse images showing no effect on CFP-DCP1a following Vifxx mCh infection,

quantified in **f** for 30 cells from 3 movies at ~18 hpi (YFP-A3G, black bars; CFP-DCP1a,

white bars). **g** HeLa.YFP-A3G infected with Vifxx CFP and treated with Ars to induce SGs

at 24 hpi and quantified in **h** for at least 100 infected cells from three independent

experiments. Magenta arrows highlight SGs present in infected cells. **i** Simultaneous

FISH/IF to detect *gagpol* mRNA and Gag show HIV RNA genomes accumulating in SGs

in HIV infected cells following Ars treatment. Magenta arrows highlight SGs present in

infected cells. All scale bars in fluorescent images = 10μm. Error bars represent standard

deviation of the mean.

**Fig. 3. HIV-1 RNA genomes but not Gag regulate A3G subcellular localization. a and**

**c** Schematic Vif+ and Vifxx two-color self-tagging virus encoding Gag-CFP, 24xMSL, and

MS2-RFP. **b** and **d** Time-lapse images of cells expressing Vif+ and Vifxx two-color HIV

constructs showing nuclear export of MS2-RFP tagged genomes, YFP-A3G degradation

(**b,** Vif+) or localization with MS2-RFP (**d,** Vifxx), and Gag-CFP expression and

accumulation into virus particles at the PM (white arrows). **e** and **f** Quantification of YFP-

A3G degradation and MS2-RFP cytoplasmic accumulation (**e**) and Gag-CFP expression

(**f**) during expressing of Vif+ two-color HIV construct. Cells (*n*=31) from three independent

experiments wherein individual cells that could be tracked for at least 17 h (1 image/hour).

Black lines highlight the ~3 h window of YFP-A3G degradation. T=0 represents onset of

MS2-RFP expression. **g** Schematic of "genome only" and "Gag only" constructs. **h**

Representative images from before (top) and after (before) photobleaching in HeLa.YFP-

A3G cells expressing "genome only" construct. White dashed circles represent regions of

targeted photobleaching. **i** Quantification of YFP-A3G fluorescence recovery experiments

and zoomed inset of time period over first 80 seconds after photobleaching. All scale bars

in fluorescent images = 10µm. Error bars represent standard deviation of the mean.

**Fig. 4. Viral RNA Genome-A3G interactions are sufficiently strong to redistribute**

**A3G to unnatural subcellular locales. a** Schematic of In Cell RNA-protein Interaction

Protocol (IC-IP). **b** Schematic of MS2-iRFP constructs used for "targeting" (top) and

representative images of MS2-iRFP and YFP-A3G localization in the absence of mRNAs

encoding MSL cassette. White arrows point to characteristic subcellular targets of Src-

(vesicles) and Lifeact- (filamentous actin) MS2-iRFP. **c** Schematic of IC-IP RNA Bait –

Vifxx two-color self-tagging HIV construct. **d** Representative images showing Free MS2-

iRFP-NLS, MS2-RFP-NLS, YFP-A3G, and Gag-CFP accumulating at PM-adjacent

assembly sites (top). (Middle) Src-MS2-iRFP accumulates at intracellular/perinuclear

vesicles, recruits MS2-RFP-tagged RNA genomes, YFP-A3G, and Gag-CFP. (Bottom)

Lifeact-MS2-iRFP induces similar accumulation at actin filaments. Region of interest

(ROI) shows YFP-A3G signal in dashed white boxes with arrows highlighting assembly

sites (top), vesicles (middle), and linear filaments (bottom). **e** Quantification of re-

localization of YFP-A3G phenotypes ($n$ = > 90 cells per condition) in the presence of NLS,

Src-, and Lifeact-MS2 targeting constructs and **f** percentages of YFP-A3G per cell signal

(IntDen) relocalized by Src- and Lifeact-MS2 constructs ($n$ > 6). All scale bars in

fluorescent images = 10µm, except in A3G ROI insets scale bars = 1µm. Error bars

represent standard deviation of the mean.

**Fig. 5. Recruitment of YFP-A3G is specific to HIV-1 RNA genome, Gag-independent,**

**and occurs long before the onset of virus particle assembly. a and c** Schematics of

"genome only" and "Gag only" IC-IP bait constructs, respectively. **b** Representative

images showing normal co-distribution of MS2-RFP tagged RNA genomes and YFP-A3G

(top) and retargeting of YFP-A3G to intracellular vesicles with RNA genomes by Src-MS2-

iRFP (bottom). **d** Representative images showing Src-MS2-RFP accumulating at

intracellular vesicles and a complete lack of YFP-A3G at these same sites (white arrow).

**e** Time-lapse images of cells expressing "genome only" construct retargeted by Src-MS2-

iRFP over a period of 4 hours. T=0 represents the first detection of MS2-RFP genome

proxy. Arrow shows accumulation of YFP-A3G at Src-MS2-iRFP+ vesicles as early as 2

hours after the onset of MS2-RFP expression. All scale bars in fluorescent images =

10µm.

**Fig. 6. Membrane-targeted APOBEC3G recruits HIV-1 RNA genomes and can inhibit**

**virus particle production. a** Schematic of Reciprocal IC-IP using Src-YFP-A3G targeted

to intracellular membranes. **b** and **c** Schematic of Vifxx two-color HIV (**b**) and "genome

only" (**c**) constructs and representative images showing Src-YFP-A3G (**b**) or Src-CFP-

A3G (**c**) recruiting MS2-RFP tagged genomes to intracellular vesicles (white arrows). **d**

Schematic of Vifxx Luc HIV infectious single-cycle virus used to infect HeLa.Src-YFP-

A3G cells and representative images of FISH/IF detecting *gag-pol* mRNA and Gag p-24,

respectively. White arrows highlight sites where Src-YFP-A3G has recruited *bona fide*

HIV RNA genomes. **e** Western blot showing dose-dependent inhibition of HIV virus

particle assembly and production from HEK293T cells transfected with Vifxx two-color

construct, Src-MS2-YFP (lanes 1 and 2), YFP-A3G (lane 3) and Src-YFP-A3G (lanes 4

and 5). Graph shows relative release factor as the ratio of Gag in virus-like particles

(VLPs) divided by Gag in cellular lysates quantified from three independent experiments.

All scale bars in fluorescent images = 10μm. Error bars represent standard deviation of

the mean.

**Fig. 7. Cytoplasmic A3G-genome interactions promote consistent per virion**

**delivery of A3G into progeny virions. a** Schematics of constructs used in single virion

analysis (SVA) studies. **b** Western blot showing cellular expression, YFP-A3G packaging,

and Gag assembly and release of constructs depicted in a. **c** Representative images of

single fluorescent virion images harvested from HEK293T.YFP-A3G cell line. Scale bar

= 500nm. **d** Quantification of SVA for each of the four constructs showing percent of

virions with MS2-RFP signal (pink) and YFP-A3G (green). Error bars represent standard

deviation of the mean ($n$ > 13,000 virions per condition). **e** Relative YFP-A3G signal per

virion or virus-like particle derived from Vifxx two-color or COGag conditions. Relative

values are normalized to the mean value for Vifxx with means (solid lines) and 25th and

75th quartiles (dashed lines) indicated.

**SUPPLEMENTARY MATERIAL**

**Fig. S1. Graphical abstract.** Summary of single-cell live imaging approaches used and

key findings.

**Video 1. Time-lapse imaging of HeLa-.FP-A3G cells infected with Vif+ HIV-1/CFP**

**virus.** Multi-channel images of HeLa.YFP-A3G cells were acquired every 60 minutes for

49h after infection. In the highlighted cell, viral gene expression (cyan) is detected ~16hpi

with YFP-A3G degradation green0 at 21hpi. This cell rounds up at 28 hpi consistent with

Vif-induced G2/M cell cycle arrest.

**Video 2. Time-lapse imaging of HeLa.YFP-A3G cells infected with Vifxx HIV-1/CFP**

**virus.** Multi-channel images were acquired every 60 minutes for 49h after infection. In the

highlighted cell, viral gene expression (cyan) is detected ~19hpi with detection of YFP-

A3G (green) at virus particles (arrows) starting at ~25hpi. An example of a transfer event

wherein a large cluster of YFP-A3G+ virus particles is released from an infected cell is

highlighted (arrows) at 32-36hpi.

**Video 3. Time-lapse imaging of HeLa.YFP-TIA1 cells infected with Rhinovirus A16.**

Multi-channel images were acquired every 30 minutes after addition of RVA16 (MOI~10).

The first instances of SG formation (YFP-TIA1 in green) are evident at ~11hpi.

**Video 4. Time-lapse imaging of HeLa.YFP-A3G cells expressing the two-color Vif+ Gag-CFP/MS2-RFP HIV-1 reporter virus.** Multi-channel images were acquired every 60 minutes for 24h beginning ~4h post-transfection. Left panel shows genome tracked using MS2-RFP (magenta), with nuclear export detected at T=4h just prior to the onset of Gag-CFP (cyan) expression (right panel) and YFP-A3G (green) degradation (center panel).

**Video 5. Time-lapse imaging of HeLa.YFP-A3G cells expressing the two-color Vifxx Gag-CFP/MS2-RFP HIV-1 reporter virus.** Multi-channel images were acquired every 60 minutes for 24 hours beginning ~4 hours post-transfection. Left panel shows genome tracked using MS2-RFP (magenta), with nuclear export detected at T=6h just prior to the onset of Gag-CFP (cyan) expression (right panel) and co-accumulation of YFP-A3G (green, center panel) with genome (MS2-RFP) and Gag-CFP at the plasma membrane (at putative sites of virus particle assembly).

**Video 6. Time-lapse imaging of HeLa.YFP-A3G cells co-expressing "genome only" MS2-RFP HIV-1 reporter virus with the Src-MS2-iRFP targeter (orange).** Multi-channel images were acquired every 60 minutes for 15h starting ~4h post-transfection. MS2-RFP tracked genomes (magenta, central panel) are first observed at T=0, with both genomes and YFP-A3G (green, right panel) immediately recruited to perinuclear vesicles, co-localizing with the Src-MS2-iRFP targeter (yellow, left panel). Video demonstrates that genomes have marked effects on YFP-A3G trafficking in the cytosol even at the earliest time points post-genome nuclear export.

# FIGURE 1

**a** HeLa.YFP-A3G

PBs

SGs

+250μM Ars

**b** NL4-3 E-R- CFP (Vif+/xx CFP)

*gag* Ψ *pol* *vif+/xx vpu* *tat rev* RRE *cfp* AAAA

**c**

% Relative Infectivity

*** p = 0.00174

Virions — p24 / YFP-A3G

Lysates — Gag / YFP-A3G / HSP90

Mock / Vif+ CFP / Vifxx CFP / Vifxx CFP + Vif

**d** 0 18 22 48 hpi

YFP-A3G

Vif+ CFP — Arrest

**e**

YFP-A3G Fluorescence / CFP Fluorescence

Vif+ — Vifxx

Times (hours)

**f** 0 21 32 48 hpi

YFP-A3G

Vifxx CFP

21 32 48 hpi

ROI YFP-A3G — 1μm / 2μm / 3μm / 4μm

**g** YFP-A3G / Gag α-p24 / *gagpol* FISH / *yfp-a3g* FISH

Vifxx

Merge — 3μm

**h** Vif+ HIV / Vifxx HIV — pm / ex

# FIGURE 2

# FIGURE 3

**a** Vif+ HIV Gag-CFP/MS2-RFP

**c** Vifxx HIV Gag-CFP/MS2-RFP

**g** genome only: ΔGag Vifxx HIV MS2-RFP

Gag only: COGag-mCh

**i**

Immobile Fraction

YFP-A3G alone — YFP-A3G + ΔGag/Vifxx HIV MS2-RFP (gRNA only)

YFP-A3G + COGag-mCh (Gag only) — YFP-A3G + Vifxx HIV mCherry (infection)

# FIGURE 4

**a** *in cellulo* RNA-protein Interaction Protocol (IC-IP)

Normal Bait | Retargeted Bait | Retargeted Bait

Normal Prey | Recruited Prey | Non-specific Prey

Colocalization | Interaction | No Interaction

**b** CONTROL (FREE) | MEMBRANE TARGETER | ACTIN TARGETER

MS2-iRFP-NLS | Src-MS2-iRFP | Lifeact-MS2-iRFP

MS2-FP

~Vesicles | Filaments

YFP-A3G

**c** IC-IP RNA Bait: Vifxx HIV Gag-CFP/MS2-RFP

*gag-cfp* / Ψ / MSL / ̶v̶i̶f̶ / vpu / tat rev / RRE / *MS2-RFP* / AAAA

**d**

Targeter: MS2-iRFP | Tracker: MS2-RFP | Prey: YFP-A3G | Gag-CFP | Merge

TARGET

Free (NLS): i. | ii. | iii. | iv. | v.

Membrane (Src): vi. | vii. | viii. | ix. | x.

Actin (Lifeact): xi. | xii. | xiii. | xiv. | xv.

A3G ROI

Free (NLS): Assembly Sites

Membrane (Src): Vesicles

Actin (Lifeact): Filaments

**e** % Cells with A3G phenotype

** p = 0.0485 | *** p = 0.000139

NLS | Src | Lifeact

■ Diffuse  ■ Vesicles  □ Actin

**f** % A3G Relocalized

Src | Lifeact

# FIGURE 5

**a** IC-IP Bait: Gagx Vifxx HIV MS2-RFP (Genome Only)

*gag*  Ψ  *MSL*  *vif*  *vpu*  *tat/rev*  **MS2-RFP** AAAA  RRE

**c** IC-IP Bait: COGag-CFP mRNA (Gag Only)

*cogag-cfp*  *MSL* AAAA

**b**

| | Targeter: MS2-iRFP | Tracker: MS2-RFP | Prey: YFP-A3G | A3G ROI |
|---|---|---|---|---|

TARGET
- Free (NLS)
- Membrane (Src)

Vesicles

**d**

| | Targeter: MS2-RFP | Prey: YFP-A3G | Gag-CFP | Merge |
|---|---|---|---|---|

TARGET
- Free (NLS)
- Membrane (Src)

**e**  0  1  2  3  4 hours

Targeter: Membranes (Src)

Tracker: MS2-RFP

YFP-A3G

Early YFP-A3G Recruitment to Vesicles by Genomes

T = 0

T = 5 h

# FIGURE 6

**a** Reciprocal *in cellulo* RNA-protein Interaction Protocol (IC-IP)

Normal A3G  Retargeted A3G  Retargeted A3G

Normal Prey  Recruited Prey  Non-specific Prey

Colocalization  Interaction  No Interaction

**b** Vifxx Gag-CFP/MS2-RFP

*gag-cfp*  ~~vif~~  vpu  tat rev  *MS2-RFP* AAAA
Ψ  MSL  RRE

MS2-RFP  Src-YFP-A3G  Gag-CFP  Merge

**c** Genome Only Vifxx

~~gag~~  ~~vif~~  vpu  tat rev  *MS2-RFP* AAAA
Ψ  MSL  RRE

MS2-RFP  YFP-A3G  Src-CFP-A3G  Merge

**d** Vifxx Luc

*gag*  *vifxx*  *vpu*  *tat rev*  *luc* AAAA
Ψ  *pol*  RRE

*gRNA* FISH  Src-YFP-A3G

Gag α-p24  Merge

**e**

Release Factor

*** p = 0.00141    *** p = 0.00265

Src-MS2  YFP-A3G  Src-A3G

VLPs  Gag-BFP
YFP

Lysates  Gag-BFP
YFP
GAPDH

1  2  3  4  5

# FIGURE 7