Sexual conflict mediated by ecological sex differences can generate diverse patterns of transgenerational plasticity

ABSTRACT

Transgenerational plasticity (TGP) occurs when the environment experienced by parents induces changes in traits of offspring and/or subsequent generations. Such effects can be adaptive or non-adaptive and are increasingly recognised as key determinants of health, cognition, development and performance across a wide range of taxa, including humans. While the conditions that favour maternal TGP are well understood, rapidly accumulating evidence indicates that TGP can be maternal or paternal, and offspring responses can be sex-specific. However, the evolutionary mechanisms that drive this diversity are unknown. We used an individual-based model to investigate the evolution of TGP when the sexes experience different ecologies. We find that adaptive TGP rarely evolves when alleles at loci that determine offspring responses to environmental information originating from the mother and father are subject to sexually antagonistic selection. By contrast, duplication and sex-limitation of such loci can allow for the evolution of a variety of sex-specific responses, including non-adaptive sex-specific TGP when sexual selection is strong. Sexual conflict could therefore help to explain why adaptive TGP evolves in some species but not others, why sons and daughters respond to parental signals in different ways, and why complex patterns of sex-specific TGP may often be non-adaptive.

Keywords:
Sex-specific ecology, parental effects, transgenerational plasticity, intralocus sexual conflict, individual-based model

INTRODUCTION

Theory predicts the evolution of anticipatory TGP when environmental conditions experienced by parents and offspring are correlated (Marshall and Uller 2007; Burgess and Marshall 2014). Environmental predictability across generations is thought to allow parents experiencing conditions that fluctuate in space or time to adaptively match the phenotype of their offspring to expected conditions through the transmission of relevant environmental information (Mousseau and Fox 1998; Marshall and Uller 2007; Dey et al. 2016). However, like other forms of plasticity, TGP is not always adaptive. Anticipatory TGP can be costly if the environment of offspring fails to match the parental environment (Crean and Marshall 2009), and some environmental factors appear to induce phenotypic changes that are invariably maladaptive or pathological (Anway et al. 2005; Marshall et al. 2008). Furthermore, mother-offspring conflict can result in induced phenotypes that optimise mothers' long-term fitness but are suboptimal for individual offspring, and *vice versa* (Uller 2008; Uller and Pen 2011; Kuijper and Johnstone 2018). Although this diversity of effects might explain why evidence for adaptive TGP across studies is generally weak (Uller et al. 2013), existing explanations fail to account for other important patterns of variation in TGP.

An increasing number of studies suggests that TGP is often sex-specific, both in terms of the parent that transfers environmental information and the offspring that responds. Paternal and maternal environments are known to have contrasting effects on offspring (Galloway 2001; Ducatez et al. 2012; Akkerman et al. 2016; Guillaume et al. 2016), and sons and daughters often respond differently to maternal versus paternal information (Pembrey et al. 2006; Dunn et al. 2011; Emborski and Mikheyev 2019; Hellmann et al. 2019), leading to diverse sex-specific patterns of TGP such as mother-daughter, father-son, mother-son and/or father-daughter effects. For example, early-life smoking in human fathers induces larger body mass in sons but not in daughters (Pembrey et al. 2006). In three-spined sticklebacks, predator-exposed mothers produce more cautious offspring regardless of sex, whereas fathers that are similarly exposed produce bolder sons but not bolder daughters (Hellmann et al. 2019). While there is growing interest in the importance of sex-specific TGP in ecology (Uller 2008), conservation biology (Donelson et al. 2018), gerontology (Benayoun et al. 2015), psychology (Bale et al. 2010), neurology (McCarthy et al. 2009), and epidemiology (Skinner et al. 2010), existing models consider only maternal TGP (i.e., environment-induced maternal

effects) and typically ignore fathers and sons (e.g., see Uller and Pen 2011; Kuijper and

Johnstone 2018). Thus, it remains unclear why maternal and paternal environments often

have contrasting phenotypic effects on offspring, and why sons and daughters often respond

differently depending on the sex of the parent.

Sex-specific responses to parental environments may reflect underlying differences in the

ecology of the sexes. In dioecious organisms, males and females are often selected to utilise

environments in different ways as a consequence of sexual dimorphism (Slatkin 1983; Shine

1989; Butler 2007; Butler et al. 2007), which can lead to ecological sex-differences in

aggregation (Ruckstuhl and Neuhaus 2006), dispersal (Pusey 1987), foraging (Schoener

1967, 1968; Vitt and Cooper Jr. 1985; Lewis et al. 2002), predation (Schoener and Schoener

1982; Magnhagen 1991), parasitism (Zuk and McKean 1996), nutrient intake (Bearhop et al.

2006; Dalu et al. 2017) and physiological stress (Bale and Epperson 2015). Environmental

change can have sex-specific effects as well (Olsson and Van der Jeugd 2002), and such sex-

differences are expected to be consistent across generations. Thus, environmental

perturbations experienced by mothers (fathers) may predict perturbations experienced by

daughters (sons). While correlations between parental and offspring environments are a key

condition for the evolution of adaptive TGP (Burgess and Marshall 2014), the implications of

sex-specific correlation of environments across generations have not been investigated.

Environmental fluctuations that affect the sexes differently could favour the evolution of sex-

specific TGP. If females and males transmit contrasting information to their offspring about

the phenotype they should adopt, progeny that pay attention to information from their same-

sex parent may gain a fitness benefit because their induced phenotype will match the

expected environment for their sex. However, progeny that pay attention to their opposite-sex

parent may incur a fitness cost resulting from a mismatch between their phenotype and

environment. Assuming that the sexes share a common genetic architecture for response to

parental environments, loci that control such responses may therefore be subject to sexually

antagonistic selection. Many genetically determined traits that are expressed in both sexes

have separate male and female fitness optima (Bonduriansky and Chenoweth 2009). This

form of genetic conflict between the sexes ("intralocus sexual conflict") can result in either

an evolutionary stalemate, where each sex expresses the trait sub-optimally (Chippindale et

al. 2001), or a resolution, where sex-linked modifiers or duplicated genes (Bonduriansky and

Chenoweth 2009; Gallach and Betrán 2011; Vankuren and Long 2018) allow for optimal,

sex-specific trait expression (i.e., optimal sexual dimorphism; Ellegren and Parsch 2007). The

outcome of conflict at loci responsive to parental information (i.e., stalemate or resolution)

could have profound consequences for resulting patterns of TGP. For example, ongoing

intralocus conflict might constrain the sexes' ability to optimise developmental responses,

resulting in polymorphic patterns of TGP within species that show little sex specificity.

However, resolution of such conflict might generate sexually dimorphic patterns of TGP by

releasing the sexes from the constraints of a shared genetic architecture, allowing sons and

daughters to attain their sex-specific optima simultaneously.

The potential for sexual conflict mediated by sex-differences in ecology to explain the

diversity of observed patterns of TGP remains unknown. Here, using an individual-based

model, we investigate the role of sexually antagonistic selection in the evolution of sex-

independent and sex-specific TGP under scenarios where loci responsive to environmental

information originating from parents are (1) subject to persistent intralocus sexual conflict

('ongoing conflict' model); or (2) sex-specific and sex-limited and therefore not subject to

sexually antagonistic selection ('resolved conflict' model). Each version of the model

considers a scenario typical of organisms that show anticipatory TGP in which environmental

perturbation—such as higher predation risk (Agrawal et al. 1999), increased temperature

(Guillaume et al. 2016), or greater food scarcity (Barrès and Zierath 2016)—causes costly

physiological stress. To assess the role of sex-specific ecology in the evolution of TGP, we

model scenarios where environmental perturbation affects the sexes in either similar or

different ways, such that males, females, both sexes, or neither sex have a higher likelihood

of experiencing stress. We assume that individuals cannot alter their own phenotype in

response to stress but, instead, transfer information about the changed environment to their

offspring via a nongenetic factor. Offspring that carry 'listening' alleles can respond to this

information from their parents by developing a stress-resistant phenotype that enhances

fitness in stressful conditions. Thus, our model adopts an "offspring's eye view" of TGP,

whereby selection acts on sons' and daughters' responses (i.e., listening/not listening) to

nongenetic information received from the father and mother. (For a similar modelling

approach in the context of mother-offspring conflict, see Kuijper and Johnstone 2018). Our

model shows how intralocus sexual conflict and its resolution can generate the diversity of

patterns of TGP seen in nature, including adaptive, non-adaptive, sex-specific and sex-

independent effects, thereby providing a new, unifying framework for the evolution of TGP.

THE MODEL

Life cycle and stress ecology

Our individual-based model considers a dioecious, bi-sexual population with discrete generations distributed over 21 x 21 patches in a torus-shaped world, with no limitation on the number of individuals per patch. Patches vary in the amount of food they hold, which is determined at the start of each generation by assigning a randomly sampled integer to each patch from the discrete uniform distribution $U\{0, 2\}$. Food resources are fixed for the duration of each generation and cannot be reduced or depleted. Each generation begins with the simultaneous emergence of adult individuals with 20-timestep lifespans, distributed randomly across patches. In every timestep, adults move randomly to one of the eight patches bordering their current patch and consume food resources, with the amount consumed dependent on the amount of resources available in that patch. The cumulative amount of resources consumed over the lifetime contributes to condition at mating/reproduction. Individuals also experience the density of conspecifics in each patch visited, and this density contributes to costly stress (with individual susceptibility to stress determined by inherited nongenetic factors, sex, and 'listening' genotype, as described below). For simplicity, we assume that adults do not alter their behaviour in response to food or conspecific density. In the final time-step, adults mate, reproduce, and die. Reproductive success is affected by condition at the end of the lifetime, which is an increasing function of the amount of resources accumulated and a decreasing function of the extent of phenotypic mismatch to stressful conditions, as described below. Figure 1 provides a visual summary of the lifecycle.

Our model assumes that adults experience stress when interacting with conspecifics (Champagne 2010; Gudsnuk and Champagne 2012; Dantzer et al. 2013), and that males and females encounter different levels of stress due to one of the sexes being more likely than the other to interact with surrounding individuals in the patch. That is, we assume that the density of conspecifics encountered by individuals within a patch is random, but that the rate of interaction with the individuals encountered is sex-specific. Consistent with the physiology of stress tolerances in animal systems (Pörtner 2002), we assume that stress (a binary state) is triggered when the cumulative number of intra-patch conspecific encounters that an individual has experienced in its life so far (K_{ind}) surpasses a threshold (α); that is, when $K_{ind} \geq \alpha_m$ (or $K_{ind} \geq \alpha_f$), where α_m (α_f) is the number of encounters that males (females)

can withstand without becoming stressed. Thus, once triggered, stress persists for the rest of

the lifetime. Thresholds α_m and α_f are allowed to vary for a fixed high or low value (80 and

8, respectively), such that the sexes can have a similar or different probability of experiencing

stress. Thus, a high α_m and low α_f threshold would represent a scenario where males rarely

interact with surrounding conspecifics (and therefore infrequently experience costly stress),

while females regularly interact with conspecifics (and therefore often experience costly

stress). Such sex-differences in the probability of conspecific interaction are likely to arise in

natural systems as a consequence of innate sex-differences in life-history strategies, such as

females seeking high quality food resources, or males searching for mates, aggregating in

leks, or avoiding competitors (Clutton-Brock 1989). While we focus on TGP triggered by

high rates of interaction with conspecifics, our model generalizes to any type of adaptive

TGP that evolves in response to environmental challenges that may be sex-specific, such as

predation, parasitism, thermal stress, nutrient intake, or starvation.

Models of anticipatory TGP often assume periodic fluctuations between environmental states

(Uller and Pen 2011; Kuijper and Johnstone 2018). For simplicity, we consider a single

period of sex-specific stress. However, simulations with multiple periodic shifts in stress

produce very similar results (see Supplementary Figure S1).

Transgenerational plasticity

Given the importance of constraints on phenotypic plasticity for the evolution of TGP

(Kuijper and Hoyle 2015), we assume that adults cannot plastically change their own

phenotype to cope with stress but transmit to offspring a stress-induced, nongenetic factor

that has the potential to alter offspring development. We make no assumptions about the type

of factor involved or whether stressed mothers and fathers necessarily transmit the same

factor. Our model simply assumes that changes in nongenetic factors—whether transcription

factors (Tao et al. 2017), small RNAs (Chen et al. 2016), DNA methylation (Weaver et al.

2004; Herman and Sultan 2016), histone modification (Morgan et al. 2005), nutrients (Blount

et al. 2000), hormones (Groothuis *et al.* 2005), antibodies (Hasselquist & Nilsson 2009),

parental care (Eggert et al. 1998), or something else—are consistently triggered by stress, that

offspring faithfully inherit these factors from stressed parents, and that such factors carry

information about the stressfulness of the parental environment. Offspring that inherit stress-

induced factors and also carry appropriate 'listening' alleles develop into individuals with a

stress-adapted phenotype (see Genetic architecture, below). We assume that offspring are

capable of responding differently to factors received from the mother versus the father, as

suggested by reported examples of sex-specific TGP (Pembrey et al. 2006; Zizzari et al.

2016; Hellmann et al. 2019), and that the phenotypic effect of receiving factors from both

parents is the same as receiving a factor from one (i.e., no additive or multiplicative effects),

such that offspring that receive conflicting information listen to the signal of the stressed

parent. However, offspring that do not carry listening alleles are unable to alter their

development in response to nongenetic information, and instead develop the default, non-

resistant phenotype.

Since mismatch between phenotype and environment is often modelled as a fitness cost of

TGP (e.g., Kuijper and Johnstone 2018), we include this cost in our model but allow it to be

sex-specific. Individuals with the default phenotype are considered well-matched if they

experience benign conditions (i.e., $K_{ind} < \alpha_m$ or $K_{ind} < \alpha_f$) and mismatched if they

experience stressful conditions (i.e., $K_{ind} \geq \alpha_m$ or $K_{ind} \geq \alpha_f$); conversely, individuals with

the stress-resistant phenotype are considered well-matched if they experience stressful

conditions (i.e., $K_{ind} \geq \alpha_m$ or $K_{ind} \geq \alpha_f$) and mismatched if they experience benign

conditions (i.e., $K_{ind} < \alpha_m$ or $K_{ind} < \alpha_f$). Thus, the level of match (∂_{ind}) to stressful

conditions is phenotype-dependent, such that $\partial_{ind} = 1 - \delta_{ind}$ for default phenotypes and

$\partial_{ind} = \delta_{ind}$ for stress-resistant phenotypes, where ∂_{ind} is the proportion of the lifetime spent

well-matched, and δ_{ind} is the proportion of the lifetime spent in a stressed state. Because

phenotype-environment mismatch can be maladaptive (Kettlewell 1956; Zimova et al. 2016),

we assume that phenotypic match affects an individual's ability to accumulate resources, and

therefore its condition at mating/reproduction (C_{ind}), such that $C_{ind} = \partial_{ind} \cdot \varphi_{ind}$, where

φ_{ind} is the total quantity of resources encountered during the lifetime. Thus, the greater the

proportion of time spent matched, the greater the amount of resources an individual is able to

accumulate, and the higher its condition at mating/reproduction. Condition determines female

fecundity (W_{ind}) and male competitiveness (Q_{ind}), such that $W_{ind} = C_{ind}$ and $Q_{ind} = C_{ind}$.

These positive linear relationships reflect the strong condition dependence of male secondary

sexual traits (Andersson 1982) and female fecundity (Honěk 1993) in natural systems.

Selection, mating and reproduction

Selection is sex-specific, with sexual selection on males occurring during the mating phase,

and fecundity selection on females occurring during the reproduction phase. In both cases,

phenotypic match (∂_{ind}), which determines Q_{ind} and W_{ind}, is the phenotypic trait directly

subject to selection.

At the end of each generation, females mate once with the male in their mating

neighbourhood that has the highest condition (i.e., the 'best male'), or a random male as a

control ('random mating'). Thus, selection acts on both sexes under best male settings, but

only on females when mating is random. The explicit spatial structure of our model allows us

to investigate the consequences of sexual selection intensity by varying the size of the mating

neighbourhood (1 patch, 9 patches, 25 patches, 49 patches), with larger neighbourhoods

generating a larger skew in male mating success (i.e., more intense sexual selection).

Reproduction occurs immediately following mating, within the same timestep. For simplicity,

we assume no male contribution to fecundity. Females produce a total of $W_{ind} = C_{ind}$

offspring (to the nearest integer), with sons and daughters equally likely to be produced. The

total number of offspring, n_{gen}, produced at the end of a generation is always more than can

survive (i.e., greater than the global carrying capacity, P). To maintain a stable population

size, $n_{gen} - P$ offspring are killed randomly before eggs hatch. We assume $P = 1000$ for all

simulations. The lifecycle repeats following the death of all parents and the simultaneous

emergence of offspring.

Genetic architecture
257

Although the genetic architecture of offspring response to nongenetic factors is poorly

understood, our assumption of genetic control of this response is motivated by empirical

findings that TGP can depend on offspring genotype (Cayuela et al. 2019). Our ongoing

conflict model assumes two listening loci that determine offspring responses to information

from stress-exposed fathers (locus *A*) and mothers (locus *B*). Phenotypic responses of

offspring are necessarily sex-independent in this version of the model because males and

females share the same genetic architecture for listening. Two alleles segregate at each locus

(wildtype: *a* and *b*; mutants: *A* and *B*), with additive effects on the probability of listening

(0% chance of listening: *aa* and *bb*; 50% chance of listening: *Aa* and *Bb*; 100% chance of

listening: *AA* and *BB*) (see Table 1 for a summary of phenotypic effects). Note that listening

alleles determine whether offspring respond to nongenetic factors, not whether parents

transmit those factors.

The resolved conflict model assumes that listening loci are duplicated and sex-limited: locus

C controls sons' listening to fathers, locus *D* controls sons' listening to mothers, locus *E*

controls daughters' listening to fathers, and locus *F* controls daughters' listening to mothers.

The loci are unlinked, and two alleles segregate at each locus (wildtype: *c*, *d*, *e*, and *f*; mutant:

C, *D*, *E*, and *F*), with additive effects on the probability of listening (0% chance of listening:

cc, *dd*, *ee* and *ff*; 50% chance of listening: *Cc*, *Dd*, *Ee* and *Ff*; 100% chance of listening: *CC*,

DD, *EE* and *FF*) (see Table 1 for a summary of phenotypic effects).

Simulations
280

In all simulations, mutations at listening loci are introduced haphazardly at the start of

generation 25 following a short burn-in period to allow population dynamics to stabilise, such

that wildtype alleles are replaced by listening alleles, and *vice versa*, at an ongoing per-locus

per-timestep rate r (fixed at $r = 0.001$).

To assess the role of sexual conflict in the evolution of sex-independent and sex-specific

listening, we varied parameters controlling male and female sensitivity to conspecific

encounter, α_m and α_f, and the intensity of sexual selection on males, for both the ongoing

conflict and resolved conflict models. We ran 40 simulations per parameter combination for

1000 generations and recorded allele frequencies at each listening locus at the end of each

run. We considered listening to have evolved if the frequency of a listening allele was ≥ 0.95

at the end of a simulation run, and polymorphisms in listening to have occurred if the

frequency of a listening allele remained stable and consistently greater than 0 and less than

0.95 for the duration of simulations. To understand how selection was acting on the sexes, we

also recorded the sex-specific distribution of phenotypic matching for each parameter

combination at the end of each run.

RESULTS
299

Ongoing conflict

The introduction of listening alleles at loci that control embryos' ability to respond to the

epigenetic signal from their father (locus A) and mother (locus B) leads to four evolutionary

outcomes: paternal TGP (allele A fixes), maternal TGP (allele B fixes), both types of TGP

(alleles A and B fix), or no TGP (neither A nor B fixes). The relative level of stress

experienced by each sex largely determines the probability of these outcomes (see Figure 2a).

When males experience higher stress than females, listening to fathers (allele A) is favoured

in sons but disfavoured in daughters because the stress-induced phenotype that listening

permits improves the phenotypic match (and, hence, fitness) of sons but decreases the match

of daughters. Listening to mothers (allele B) is neutral in this scenario, since mothers rarely

experience stress. However, despite the evolutionary benefit to males of listening to fathers,

sexual conflict at locus A largely inhibits the fixation of the A allele, which codes for paternal

TGP, because the benefit of paternal TGP to sons is not enough to overcome the cost of

paternal TGP to daughters (hence, the paucity of outcomes where the A allele approaches

fixation in Figure 2a). A similar but converse situation occurs when females experience more

stress than males (Figure 2a). Thus, despite the benefit that listening provides to the stressed

sex, sexual conflict at listening loci constrains the evolution of TGP because of the costs that

listening imposes on the opposite sex.

The sexes' evolutionary interests in listening diverge when one sex experiences higher stress

than the other and, in our model, this conflict manifests as sex-differences in ∂_{ind} (see Figure

3a). Sexual selection on males has a strong influence on the distribution of ∂_{ind} and hence

patterns of listening under sexual conflict. Indeed, the relative strength of selection on the

sexes modulates the effect of intralocus conflict on listening outcomes. In the absence of

sexual selection, fecundity selection maximises female matching regardless of the parent that

experiences stress (see Supplementary Figure S2), which leads to optimal listening outcomes

for females (i.e., the evolution of maternal TPG when females experience higher stress, and

no TGP when males experience higher stress; see Supplementary Figure S3). However,

sexual selection on males counteracts this effect. When sexual selection and fecundity

selection are similar in strength but act in opposite ways on stress-resistance, stable sex-

specific listening polymorphisms evolve (see Figure 4). These intermediate allele frequencies

are associated with bimodal sex-differences in ∂_{ind} whereby each sex exhibits both high and

low matching (see Figure 3a), reflecting an evolutionary stalemate between the sexes—a

predicted outcome of intralocus sexual conflict (Bonduriansky and Chenoweth 2009).

Increasing the intensity of sexual selection gives males a greater edge in the conflict, as

evidenced by the reduced number of simulations ending in maternal TGP when females

experience greater stress (Figure 2a).

In simulations in which neither parent is stressed, listening alleles are neutral and there is no

selection for listening (Figure 2a). However, in cases where mothers and fathers are both

stressed, listening to either parent is favoured in both sons and daughters, and the particular

form of TGP that evolves (paternal, maternal or both) is determined by whichever alleles

happen to spread faster by chance (Figure 2a).

Resolved conflict

When intralocus sexual conflict is resolved through the duplication and sex-limitation of

listening loci (Gallach and Betrán 2011), the sexes can pursue optimal listening strategies

free from the constraints of a shared genetic architecture. This resolution allows for

simulations of the resolved conflict model to end in any combination of sex-specific maternal

and paternal TGP.

The resolution of intralocus conflict is characterised by higher values of ∂_{ind} (Figure 3b) and

optimal sex-specific listening outcomes (Figure 2b) for both sexes. When females experience

higher stress than males, fecundity selection drives daughters to listen exclusively to mothers

(Figure 2b). Conversely, when males experience higher stress than females, sexual selection

drives sons to listen exclusively to fathers (Figure 2b). These patterns evolve because sexual

selection and fecundity selection are able to act independently on son-specific and daughter-

specific listening alleles, respectively. The resolution of sexual conflict allows for the

adaptive evolution of these simple mother-daughter and father-son effects in a similar way to

parent-of-origin effects on gene expression (Day and Bonduriansky 2004).

However, our model also shows how more complex patterns of TGP involving multiple

parent-offspring effects can evolve. Diverse sex-specific listening outcomes can evolve when

stress is experienced by (1) both parents (representing adaptive effects) or (2) neither parent

(representing non-adaptive effects) (see Figure 2b). In the first case (1), diverse combinations

result from the random fixation of equally beneficial alleles: if both parents send similar

stress signals, sons and daughters can reap similar benefits by listening to either parent. Very

few simulations in this parameter space end in only one sex listening because sexual selection

and fecundity selection simultaneously favour listening in both males and females. In the

second case (2), stochastic patterns of sex-specific listening evolve via random fixation of

neutral alleles driven by sexual selection. As expected, no TGP evolves if both parents are

unstressed and sexual selection is weak (Figure 2b). However, intensifying sexual selection

leads to increased diversity of sex-specific combinations of TGP. Indeed, 11 out of the 15

possible listening combinations evolve at the highest sexual selection intensity (Figure 2b).

This diversity occurs because the skew in male mating success generated by intense sexual

selection substantially reduces effective population size, causing neutral alleles carried by the

most successful males to reach fixation via an effect analogous to hitchhiking.

DISCUSSION

We modelled the evolution of environmentally induced TGP for stress-resistance within a

heterogeneous environment where mothers and fathers can experience different levels of

stress that are correlated within sex across generations. We then asked how sex-differences in

stress ecology might drive selection on alleles that allow offspring to respond ('listen') to

parental information and develop a stress-resistant phenotype when the genetic architecture

of listening is either shared between the sexes or independent of sex. Our model allowed us to

investigate the extent to which intralocus sexual conflict and its resolution could explain the

diverse patterns of sex-specific and sex-independent TGP in natural systems. Our model also

enabled us to ask how selection acting more strongly on one sex than the other affects

listening outcomes.

Our simulations produced five key findings. First, if both sexes share the same genetic

architecture for listening and one sex experiences a greater probability of stress than the

other, intralocus sexual conflict will inhibit the spread of listening alleles. The resulting allele

frequencies for each sex will then be determined by the relative strength of selection on males

and females: stronger selection on one sex will skew listening outcomes more toward that

sex's optimum, whereas selection of similar strength will generate bimodal fitnesses and

stable listening polymorphisms for each sex. Second, sex-independent listening outcomes,

where offspring listen to parents regardless of their own sex, will occur most frequently when

sons and daughters gain equal benefit from listening (i.e., when both sexes experience stress

and therefore have the same evolutionary interest in listening to stress signals). Third, the

resolution of intralocus sexual conflict through duplication and sex-limitation of listening

loci, which allows each sex to optimise listening independent of the other sex, will generate

daughter-specific maternal TGP when females experience greater stress and son-specific

paternal TGP when males experience greater stress. Fourth, complex patterns of sex-specific

listening, whereby sons and daughters independently listen to one or both parents, can evolve

if conflict is resolved and stress is high for both sexes. Under these settings, all listening

alleles have equal benefit to either sex, and the allele that spreads most quickly by chance

will go to fixation. Fifth, complex listening combinations can also evolve if conflict is

resolved and neither sex experiences stress. In this situation, all listening alleles are neutral

and strong sexual selection can drive random combinations of listening alleles to fixation via

an effect analogous to genetic hitchhiking.

Our results suggest that a range of phenomena reported in the empirical TGP literature could

be explained by sexual conflict. Maternal and paternal environments are known to affect the

phenotypes of sons and daughters in different ways (Pembrey et al. 2006; Von Engelhardt et

al. 2006; Dunn et al. 2011; Kruuk et al. 2015; Emborski and Mikheyev 2019; Hellmann et al.

2019), and our model provides a potential explanation for this intriguing pattern of TGP. The

key assumption of our model that generates sex-biased outcomes is the sex-specificity of

environmental stress. Existing models of TGP also assume environmental correlations

spanning multiple generations (Motro 1983; Godfray and Parker 1991; Kilner and Hinde

2008; Uller and Pen 2011; Kuijper and Johnstone 2018), but such models overlook the

contribution of the paternal environment to offspring phenotypes (i.e., paternal effects; Rando

2012; Crean and Bonduriansky 2014; Soubry et al. 2014), as well as the sex-specificity of

offspring responses to parental information (e.g., Pembrey et al. 2006; Zizzari et al. 2016;

Hellmann et al. 2019). By incorporating both sexes and sex-specific environments, our

model shows how differences in parental signals driven by sex-differences in ecology can

lead to the evolution of sex-biased TGP.

Our simulations also suggest that sex-specific patterns of TGP may often be non-adaptive.

One of the surprising results from our model was the diversity of sex-specific outcomes that

evolved when sex-limited listening alleles were selectively neutral. Sexual selection on males

facilitated the spread of neutral listening alleles via a kind of hitchhiking effect, resulting in

diverse combinations of sex-specific TGP. This result suggests that organisms characterised
by intense sexual selection may be more likely to exhibit complex patterns of sex-specific
TGP. Other processes that promote genetic drift, such as genetic bottlenecks, founder effects
and genetic hitchhiking, could generate similar results. The diversity of sex-specific
outcomes that evolved in our model suggests that much of the sex-specific variation in TGP
in natural systems (e.g., Pembrey et al. 2006; Dunn et al. 2011; Emborski and Mikheyev
2019) may be non-adaptive. A finding of diverse and unpredictable sex-specific TGP
following experimental manipulation of the intensity of sexual selection would lend empirical
support to this non-adaptive explanation.
445
Our model may additionally help to explain why many studies fail to observe adaptive TGP
despite *a priori* predictions of an adaptive benefit (see reviews by Uller et al. 2013; Heard
and Martienssen 2014). Existing models suggest that mother-offspring conflict—where the
fitness optima of mothers and offspring with respect to offspring phenotype are different and
cannot be achieved simultaneously (Trivers 1974)—can lead to the breakdown of TGP under
certain conditions (Uller and Pen 2011; Kuijper and Johnstone 2018). Our simulations show
that conflict between the sexes—which is ubiquitous in bisexual taxa (Arnqvist and Rowe
2005)—could also inhibit the fixation of alleles for TGP. Intralocus sexual conflict could
therefore be a powerful and taxonomically widespread explanation for the absence of
adaptive TGP in many taxa, just as intralocus sexual conflict can constrain the evolution of
sexual dimorphism in many other traits (Bonduriansky and Chenoweth 2009). Indeed, the
ubiquity of sex-specific ecologies in natural systems (Schoener 1967, 1968; Schoener and
Schoener 1982; Vitt and Cooper Jr. 1985; Pusey 1987; Shine 1989, 1991; Magnhagen 1991;
Zuk and McKean 1996; Temeles et al. 2000; Mysterud 2000; Lewis et al. 2002; Olsson and
Van der Jeugd 2002; Bearhop et al. 2006; Ruckstuhl and Neuhaus 2006; Butler 2007; Butler
et al. 2007; Wearmouth and Sims 2008; Bale and Epperson 2015; Dalu et al. 2017; Fryxell et
al. 2019), and the conflicting signals that parents in such ecologies are likely to send to
offspring, could prevent the evolution of adaptive TGP, unless the intersexual genetic
correlation for listening can be reduced through changes in genetic architecture involving
locus duplication or sex-linked modifiers that enable sex-specific listening. Our model also
shows that intralocus sexual conflict could maintain genetic variation for listening via the
evolution of stable listening polymorphisms. This is a predicted outcome of non-random
mating with respect to traits subject to sexual conflict (Härdling and Bergsten 2006), which
explains why polymorphisms evolve in our simulations only when sexual selection is present.

Empirical studies indicate that TGP can be induced by a variety of environment factors,

including temperature (Salinas and Munch 2012), light (Galloway and Etterson 2007), food

availability (Vijendravarma et al. 2010), food quality (Bonduriansky and Head 2007), water

availability (Herman and Sultan 2016), predation risk (Sheriff et al. 2010), pathogen exposure

(Sadd et al. 2005) and parasite load (Lefèvre et al. 2010). Although we specifically model

environmental perturbation as stressful, this assumption does not reduce the generality of our

conclusions as the sexes may experience benign or resource-rich environments differently as

well. For example, offspring traits such as body size that can be enhanced by high parental

condition (Qvarnström and Price 2001; Crean and Bonduriansky 2014) often have sex-

specific optima (Badyaev and Martin 2000), and therefore alleles for listening to such cues

may also be subject to sexual conflict. Our findings also highlight the importance of

accounting for sex-specific variation in ecology when motivating TGP experiments and

interpreting results, as failure to do so could lead to potentially misleading conclusions (see

also Kruuk et al. 2015).

Our model could be tested empirically by assessing the expected match between fitness

outcomes of phenotypic responses for each sex and the type of TGP that is observed. For

example, an absence of a clear signal of TGP associated with contrasting fitnesses between

the sexes with respect to offspring phenotype would suggest that fixation of alleles at loci

responsive to parental environments is inhibited by ongoing intralocus conflict. Further

support for such a conclusion would be found if experimentally decreasing the intensity of

selection on one sex results in the evolution of patterns of TGP that are more aligned to the

fitness interests of the opposite sex. Assessment of the genetic architecture of TGP in natural

systems could also prove valuable. There is evidence that TGP has a genetic basis (Stjernman

and Little 2011; Herman and Sultan 2016; Cayuela et al. 2019), but the extent to which the

expression of TGP depends specifically on genes that regulate sex-specific development

remains an open question. Genetic analyses could determine whether sex-specific TGP,

where sons (daughters) respond exclusively to paternal (maternal) environments (e.g.,

Pembrey et al. 2006; Zizzari et al. 2016; Hellmann et al. 2019), is mediated by listening

alleles that are sex-limited in offspring.

The strong inhibiting effect of intralocus conflict on offspring listening that we observed in

our model suggests that TGP may be especially widespread in organisms for which sexual

conflict is absent or weak. For example, TGP may be particularly common in asexual

lineages in which adaptive maternal effects can evolve in the absence of males. However,

mother-offspring conflict may be a more prevalent factor in such lineages (Kuijper and

Johnstone 2018). Similarly, monogamous species may have a greater capacity for TGP if

sexual conflict is relatively weak in such taxa (Holland and Rice 1999; Martin and Hosken

2003) and if monogamy is associated with reduced sex-specificity of ecology (Fryxell et al.

2019; Giery and Layman 2019). Comparative analyses of the incidence of TGP in sexual

versus asexual taxa, and monogamous versus polyandrous taxa, could shed light on these

questions.

Future work could usefully extend our model by investigating how the evolution of

alternative strategies for coping with environmental stress in parents, such as phenotypic

plasticity in development (Denver 1997), morphology (Gratani 2014) or behaviour (Badyaev

2005), or offspring sex-ratio adjustment (Nager et al. 1999; Rosenfeld and Roberts 2004),

modulates or inhibits the evolution of TGP. Assessment of the effect of temporal

heterogeneity in sex-differences, and error in nongenetic signals and offspring responses,

may also be fruitful.

Our model suggests a number of novel predictions. First, species in which the sexes

experience substantially different ecologies may be more likely to exhibit non-adaptive or

polymorphic TGP due to ongoing intralocus conflict at listening loci, or, alternatively,

adaptive sex-specific TGP if conflict is resolved through the evolution of sex-specific

listening loci, with mother-daughter (father-son) effects most likely when females (males)

experience greater stress. Second, sex-independent TGP may be common in species in which

males and females experience similar ecologies and no sexual conflict over listening. Third,

complex patterns of sex-specific TGP are likely to be non-adaptive, and may be most

common in species in which males are subject to strong sexual selection. Future studies could

test these predictions by investigating the genetic architecture of loci involved in TGP and

selection on expression of parentally transmitted information in both sexes. The diversity of

listening outcomes generated by our model is consistent with the diverse patterns of TGP

seen in nature across a wide range of taxa (Mousseau and Fox 1998; Agrawal et al. 1999;

Galloway 2001; Anway et al. 2005; Pembrey et al. 2006; Galloway and Etterson 2007; Dunn

et al. 2011; Salinas and Munch 2012; Ducatez et al. 2012; Babenko et al. 2015; Akkerman et

al. 2016; Guillaume et al. 2016; Emborski and Mikheyev 2019). Our model therefore

provides a unifying framework for understanding the origin and maintenance of diversity in

observed patterns of TGP.

DATA ACCESSIBILITY

The simulation data that support the findings of this study are available on Dryad Data

Repository:

https://datadryad.org/stash/share/a_c-2-t-hugOmUf0IdBKmw6gcL8ujoC_eDKVovj39fA

The IBM code that generated the data is also available via this link.

REFERENCES

Agrawal, A. A., C. Laforsch, and R. Tollrian. 1999. Transgenerational induction of defences
     in animals and plants. Nature 401:60–63.
Akkerman, K. C., A. Sattarin, J. K. Kelly, and A. G. Scoville. 2016. Transgenerational
     plasticity is sex-dependent and persistent in yellow monkeyflower (Mimulus guttatus).
     Environ. Epigenetics 2:dvw003.
Andersson, M. 1982. Sexual selection, natural selection and quality advertisement. Biol. J.
     Linn. Soc. 17:375–393. Oxford University Press.
Anway, M. D., A. S. Cupp, N. Uzumcu, and M. K. Skinner. 2005. Epigenetic
     transgenerational actions of endocrine disruptors and male fertility. Science (80-.).
     308:1466–1469.
Arnqvist, G., and L. Rowe. 2005. Sexual Conflict. Princeton University Press, Princeton.
Babenko, O., I. Kovalchuk, and G. A. S. Metz. 2015. Stress-induced perinatal and
     transgenerational epigenetic programming of brain development and mental health.
     Neurosci. Biobehav. Rev. 48:70–91. Elsevier Ltd.
Badyaev, A. V. 2005. Stress-induced variation in evolution: from behavioural plasticity to
     genetic assimilation. Proc. R. Soc. B Biol. Sci. 272:877–886.
Badyaev, A. V., and T. E. Martin. 2000. Sexual dimorphism in relation to current selection in
     the house finch. Evolution (N. Y). 54:987–997.
Bale, T. L., T. Z. Baram, A. S. Brown, J. M. Goldstein, T. R. Insel, M. M. McCarthy, C. B.
     Nemeroff, T. M. Reyes, R. B. Simerly, E. S. Susser, and E. J. Nestler. 2010. Early life
     programming and neurodevelopmental disorders. Biol. Psychiatry 68:314–319. Elsevier
     Inc.
Bale, T. L., and C. N. Epperson. 2015. Sex differences and stress across the lifespan. Nat.
     Neurosci. 18:1413. Nature Publishing Group.
Barrès, R., and J. R. Zierath. 2016. The role of diet and exercise in the transgenerational
     epigenetic landscape of T2DM. Nat. Rev. Endocrinol. 12:441–451. Nature Publishing
     Group.
Bearhop, S., R. A. Phillips, R. McGill, Y. Cherel, D. A. Dawson, and J. P. Croxall. 2006.
     Stable isotopes indicate sex-specific and long-term individual foraging specialisation in
     diving seabirds. Mar. Ecol. Prog. Ser. 311:157–164.

Benayoun, B. A., E. A. Pollina, and A. Brunet. 2015. Epigenetic regulation of ageing: linking environmental inputs to genomic stability. Nat. Rev. Mol. Cell Biol. 16:593–610. Nature Publishing Group.

Blount, J. D., D. C. Houston, and A. P. Moller. 2000. Why egg yolk is yellow. Trends Ecol. Evol. 15:47–49.

Bonduriansky, R., and S. F. Chenoweth. 2009. Intralocus sexual conflict. Trends Ecol. Evol. 24:280–288.

Bonduriansky, R., and M. Head. 2007. Maternal and paternal condition effects on offspring phenotype in Telostylinus angusticollis (Diptera: Neriidae). J. Evol. Biol. 20:2379–2388. Wiley Online Library.

Burgess, S. C., and D. J. Marshall. 2014. Adaptive parental effects: The importance of estimating environmental predictability and offspring fitness appropriately. Oikos 123:769–776.

Butler, M. A. 2007. Vive le difference! Sexual dimorphism and adaptive patterns in lizards of the genus Anolis. Integr. Comp. Biol. 47:272–284.

Butler, M. A., S. A. Sawyer, and J. B. Losos. 2007. Sexual dimorphism and adaptive radiation in Anolis lizards. Nature 447:202–205.

Cayuela, H., S. Jacob, N. Schtickzelle, R. Verdonck, H. Philippe, M. Laporte, M. Huet, L. Bernatchez, and D. Legrand. 2019. Transgenerational dispersal plasticity and its fitness consequences are under genetic control. bioRxiv 791210.

Champagne, F. A. 2010. Epigenetic influence of social experiences across the lifespan. Dev. Psychobiol. 52:299–311.

Chen, Q., W. Yan, and E. Duan. 2016. Epigenetic inheritance of acquired traits through sperm RNAs and sperm RNA modifications. Nat. Rev. Genet. 17:733–743. Nature Publishing Group.

Chippindale, A. K., J. R. Gibson, and W. R. Rice. 2001. Negative genetic correlation for adult fitness between sexes reveals ontogenetic conflict in Drosophila. Proc. Natl. Acad. Sci. U. S. A. 98:1671–1675.

Clutton-Brock, T. H. 1989. Mammalian mating system. Proc. R. Soc. B Biol. Sci. 236:339–372.

Crean, A. J., and R. Bonduriansky. 2014. What is a paternal effect? Trends Ecol. Evol. 29:554–559. Elsevier Ltd.

Crean, A. J., and D. J. Marshall. 2009. Coping with environmental uncertainty: dynamic bet hedging as a maternal effect. Philos. Trans. R. Soc. B Biol. Sci. 364:1087–1096.

Dalu, T., R. J. Wasserman, T. J. F. Vink, and O. L. F. Weyl. 2017. Sex and species specific isotopic niche specialisation increases with trophic complexity: Evidence from an ephemeral pond ecosystem. Sci. Rep. 7:1–12. Nature Publishing Group.

Dantzer, B., A. E. M. Newman, R. Boonstra, R. Palme, S. Boutin, M. M. Humphries, and A. G. McAdam. 2013. Density triggers maternal hormones that increase adaptive offspring growth in a wild mammal. Science (80-.). 340:1215–1217.

Day, T., and R. Bonduriansky. 2004. Intralocus sexual conflict can drive the evolution of genomic imprinting. Genetics 167:1537–1546.

Denver, R. J. 1997. Environmental stress as a developmental cue: corticotropin-releasing hormone is a proximate mediator of adaptive phenotypic plastiity in amphibian metamorphosis. Horm. Behav. 31:169–179.

Dey, S., S. R. Proulx, and H. Teotónio. 2016. Adaptation to temporally fluctuating environments by the evolution of maternal effects. PLoS Biol. 14:1–29.

Donelson, J. M., S. Salinas, P. L. Munday, and L. N. S. Shama. 2018. Transgenerational plasticity and climate change experiments: Where do we go from here? Glob. Chang. Biol. 24:13–34.

Ducatez, S., M. Baguette, V. M. Stevens, D. Legrand, and H. Fréville. 2012. Complex interactions between paternal and maternal effects: parental experience and age at reproduction affect fecundity and offspring performance in a butterfly. Evolution (N. Y). 66:3558–3569.

Dunn, G. A., C. P. Morgan, and T. L. Bale. 2011. Sex-specificity in transgenerational epigenetic programming. Horm. Behav. 59:290–295.

Eggert, A. K., M. Reinking, and J. K. Müller. 1998. Parental care improves offspring survival and growth in burying beetles. Anim. Behav. 55:97–107.

Ellegren, H., and J. Parsch. 2007. The evolution of sex-biased genes and sex-biased gene expression. Nat. Rev. Genet. 8:689–698.

Emborski, C., and A. S. Mikheyev. 2019. Ancestral diet transgenerationally influences offspring in a parent-of-origin and sex-specific manner. Philos. Trans. R. Soc. B Biol. Sci. 374.

Fryxell, D. C., D. E. Weiler, M. T. Kinnison, and E. P. Palkovacs. 2019. Eco-Evolutionary Dynamics of Sexual Dimorphism. Trends Ecol. Evol. 34:591–594. Elsevier Ltd.

Gallach, M., and E. Betrán. 2011. Intralocus sexual conflict resolved through gene duplication. Trends Ecol. Evol. 26:222–228.

Galloway, L. F. 2001. The effect of maternal and paternal environments on seed characters in the herbaceous plant Campanula americana (Campanulaceae). Am. J. Bot. 88:832–840.

Galloway, L. F., and J. R. Etterson. 2007. Transgenerational plasticity is adaptive in the wild. Science (80-.). 318:1134–1136.

Giery, S. T., and C. A. Layman. 2019. Ecological consequences of sexually selected traits: an eco-evolutionary perspective. Q. Rev. Biol. 94:29–74.

Godfray, H. C. J., and G. A. Parker. 1991. Clutch size, fecundity and parent-offspring conflict. Philos. Trans. - R. Soc. London, B 332:67–79.

Gratani, L. 2014. Plant phenotypic plasticity in response to environmental factors. Adv. Bot. 2014:1–17.

Groothuis, T. G. G., W. Müller, N. Von Engelhardt, C. Carere, and C. Eising. 2005. Maternal hormones as a tool to adjust offspring phenotype in avian species. Neurosci. Biobehav. Rev. 29:329–352.

Gudsnuk, K., and F. A. Champagne. 2012. Epigenetic influence of stress and the social environment. ILAR J. 53:279–288.

Guillaume, A. S., K. Monro, and D. J. Marshall. 2016. Transgenerational plasticity and environmental stress: do paternal effects act as a conduit or a buffer? Funct. Ecol. 30:1175–1184.

Härdling, R., and J. Bergsten. 2006. Nonrandom mating preserves intrasexual polymorphism and stops population differentiation in sexual conflict. Am. Nat. 167:401–409. University of Chicago Press.

Hasselquist, D., and J. Å. Nilsson. 2009. Maternal transfer of antibodies in vertebrates: Trans-generational effects on offspring immunity. Philos. Trans. R. Soc. B Biol. Sci. 364:51–60.

Heard, E., and R. A. Martienssen. 2014. Transgenerational epigenetic inheritance: myths and mechanisms. Cell 157:95–109. Elsevier Inc.

Hellmann, J. K., S. A. Bukhari, J. Deno, and A. M. Bell. 2019. Sex-specific transgenerational plasticity in threespined sticklebacks. bioRxiv https://do.

Herman, J. J., and S. E. Sultan. 2016. DNA methylation mediates genetic variation for adaptive transgenerational plasticity. Proc. R. Soc. B Biol. Sci. 283.

Holland, B., and W. R. Rice. 1999. Experimental removal of sexual selection reverses intersexual antagonistic coevolution and removes a reproductive load. Proc. Natl. Acad. Sci. U. S. A. 96:5083–5088.

Honěk, A. 1993. Intraspecific Variation in Body Size and Fecundity in Insects: A General
Relationship. Oikos 66:483.
Kettlewell, H. D. 1956. Further selection experiments on industrial melanism in the
Lepidoptera. Heredity (Edinb). 4:287–301.
Kilner, R. M., and C. A. Hinde. 2008. Information warfare and parent-offspring conflict.
Adv. Study Behav. 38:283–336.
Kruuk, L. E. B., J. Livingston, A. Kahn, and M. D. Jennions. 2015. Sex-specific maternal
effects in a viviparous fish. Biol. Lett. 11.
Kuijper, B., and R. B. Hoyle. 2015. When to rely on maternal effects and when on
phenotypic plasticity? Evolution (N. Y). 69:950–968.
Kuijper, B., and R. A. Johnstone. 2018. Maternal effects and parent–offspring conflict.
Evolution (N. Y). 72:220–233.
Lefèvre, T., L. Oliver, M. D. Hunter, and J. C. De Roode. 2010. Evidence for trans-
generational medication in nature. Ecol. Lett. 13:1485–1493.
Lewis, S., S. Benvenuti, L. Dall–Antonia, R. Griffiths, L. Money, T. N. Sherratt, S. Wanless,
and K. C. Hamer. 2002. Sex-specific foraging behaviour in a monomorphic seabird.
Proc. R. Soc. London. Ser. B Biol. Sci. 269:1687–1693. The Royal Society.
Magnhagen, C. 1991. Predation risk as a cost of reproduction.
Marshall, D. J., R. M. Allen, and A. J. Crean. 2008. The ecological and evolutionary
importance of maternal effects in the sea. Oceanogr. Mar. Biol. 46:203–250.
Marshall, D. J., and T. Uller. 2007. When is a maternal effect adaptive? Oikos 116:1957–
1963.
Martin, O. Y., and D. J. Hosken. 2003. Costs and benefits of evolving under experimentally
enforced polyandry or monogamy. Evolution (N. Y). 57:2765–2772.
McCarthy, M. M., A. P. Auger, T. L. Bale, G. J. De Vries, G. A. Dunn, N. G. Forger, E. K.
Murray, B. M. Nugent, J. M. Schwarz, and M. E. Wilson. 2009. The epigenetics of sex
differences in the brain. J. Neurosci. 29:12815–12823.
Morgan, H. D., F. Santos, K. Green, W. Dean, and W. Reik. 2005. Epigenetic reprogramming
in mammals. Hum. Mol. Genet. 14:47–58.
Motro, U. 1983. Optimal rates of dispersal. III. Parent-offspring conflict. Theor. Popul. Biol.
23:159–168.
Mousseau, T. A., and C. W. Fox. 1998. Maternal Effects as Adaptations. Oxford University
Press.
Mysterud, A. 2000. The relationship between ecological segregation sexual body size
dimorphism large herbivores. Oecologia 124:40–54.
Nager, R. G., P. Monaghan, R. Griffiths, D. C. Houston, and R. Dawson. 1999. Experimental
demonstration that offspring sex ratio varies with maternal condition. Proc. Natl. Acad.
Sci. U. S. A. 96:570–573.
Olsson, O., and H. P. Van der Jeugd. 2002. Survival in king penguins Aptenodytes
patagonicus: temporal and sex-specific effects of environmental variability. Oecologia
132:509–516.
Pembrey, M. E., L. O. Bygren, G. Kaati, S. Edvinsson, K. Northstone, M. Sjöström, and J.
Golding. 2006. Sex-specific, male-line transgenerational responses in humans. Eur. J.
Hum. Genet. 14:159–166.
Pörtner, H. O. 2002. Climate variations and the physiological basis of temperature dependent
biogeography: systemic to molecular hierarchy of thermal tolerance in animals. Comp.
Biochem. Physiol. - A Mol. Integr. Physiol. 132:739–761.
Pusey, A. 1987. Sex-biased dispersal and inbreeding avoidance in birds and mammals.
Trends Ecol. Evol. 2:295–299.
Qvarnström, A., and T. D. Price. 2001. Maternal effects, paternal effects and sexual selection.

Trends Ecol. Evol. 16:95–100.

Rando, O. J. 2012. Daddy issues: paternal effects on phenotype. Cell 151:702–708. Elsevier
Inc.

Rosenfeld, C. S., and R. M. Roberts. 2004. Maternal diet and other factors affecting offspring
sex ratio: a review. Biol. Reprod. 71:1063–1070.

Ruckstuhl, K., and P. Neuhaus. 2006. Sexual segregation in vertebrates. Cambridge
University Press.

Sadd, B. M., Y. Kleinlogel, R. Schmid-Hempel, and P. Schmid-Hempel. 2005. Trans-
generational immune priming in a social insect. Biol. Lett. 1:386–388.

Salinas, S., and S. B. Munch. 2012. Thermal legacies: transgenerational effects of
temperature on growth in a vertebrate. Ecol. Lett. 15:159–163.

Schoener, T. W. 1968. The Anolis lizards of Bimini: resource partitioning in a complex
fauna. Ecology 49:704–726.

Schoener, T. W. 1967. The ecological significance of sexual dimorphism in size in the lizard
Anolis conspersus. Science (80-.). 155:474–477.

Schoener, T. W., and A. Schoener. 1982. The ecological correlates of survival in some
Bahamian Anolis lizards. Oikos 39:1–16.

Sheriff, M. J., C. J. Krebs, and R. Boonstra. 2010. The ghosts of predators past: population
cycles and the role of maternal programming under fluctuating predation risk. Ecology
91:2983–2994.

Shine, R. 1989. Ecological causes for the evolution of sexual dimorphism: a review of the
evidence. Q. Rev. Biol. 64:419–461.

Shine, R. 1991. Intersexual dietary divergence and the evolution of sexual dimorphism in
snakes. Am. Nat. 138:103–122.

Skinner, M. K., M. Manikkam, and C. Guerrero-Bosagna. 2010. Epigenetic transgenerational
actions of environmental factors in disease etiology. Trends Endocrinol. Metab. 21:214–
222. Elsevier Ltd.

Slatkin, M. 1983. Ecological causes of sexual dimorphism. Evolution (N. Y). 38:622–630.

Soubry, A., C. Hoyo, R. L. Jirtle, and S. K. Murphy. 2014. A paternal environmental legacy:
Evidence for epigenetic inheritance through the male germ line. BioEssays 36:359–371.

Stjernman, M., and T. J. Little. 2011. Genetic variation for maternal effects on parasite
susceptibility. J. Evol. Biol. 24:2357–2363.

Tao, Z., L. Shen, X. Gu, Y. Wang, H. Yu, and Y. He. 2017. Embryonic epigenetic
reprogramming by a pioneer transcription factor in plants. Nature 551:124–128. Nature
Publishing Group.

Temeles, E. J., I. L. Pan, J. L. Brennan, and J. N. Horwitt. 2000. Evidence for ecological
causation of sexual dimorphism in a hummingbird. Science (80-.). 289:441–443.

Trivers, R. L. 1974. Parent-offspring conflict. Am. Zool. 14:249–264.

Uller, T. 2008. Developmental plasticity and the evolution of parental effects. Trends Ecol.
Evol. 23:432–438.

Uller, T., S. Nakagawa, and S. English. 2013. Weak evidence for anticipatory parental effects
in plants and animals. J. Evol. Biol. 26:2161–2170.

Uller, T., and I. Pen. 2011. A theoretical model of the evolution of maternal effects under
parent-offspring conflict. Evolution (N. Y). 65:2075–2084.

Vankuren, N. W., and M. Long. 2018. Gene duplicates resolving sexual conflict rapidly
evolved essential gametogenesis functions. Nat. Ecol. Evol. 2:705–712. Springer US.

Vijendravarma, R. K., S. Narasimha, and T. J. Kawecki. 2010. Effects of parental larval diet
on egg size and offspring traits in Drosophila. Biol. Lett. 6:238–241.

Vitt, L. J., and W. E. Cooper Jr. 1985. The evolution of sexual dimorphism in the skink
Eumeces laticeps : an example of sexual selection . Can. J. Zool. 63:995–1002.

Von Engelhardt, N., C. Carere, C. Dijkstra, and T. G. G. Groothuis. 2006. Sex-specific
effects of yolk testosterone on survival, begging and growth of zebra finches. Proc. R.
Soc. B Biol. Sci. 273:65–70.
Wearmouth, V. J., and D. W. Sims. 2008. Sexual segregation in marine fish, reptiles, birds
and mammals: behaviour patterns, mechanisms and conservation implications. Adv.
Mar. Biol. 54:107–170. Elsevier.
Weaver, I. C. G., N. Cervoni, F. A. Champagne, A. C. D'Alessio, S. Sharma, J. R. Seckl, S.
Dymov, M. Szyf, and M. J. Meaney. 2004. Epigenetic programming by maternal
behavior. Nat. Neurosci. 7:847–854.
Zimova, M., L. S. Mills, and J. J. Nowak. 2016. High fitness costs of climate change-induced
camouflage mismatch. Ecol. Lett. 19:299–307.
Zizzari, Z. V., N. M. van Straalen, and J. Ellers. 2016. Transgenerational effects of nutrition
are different for sons and daughters. J. Evol. Biol. 29:1317–1327.
Zuk, M., and K. A. McKean. 1996. Sex differences in parasite infections: patterns and
processes. Int. J. Parasitol. 26:1009–1024. Elsevier.
795

   TABLE 1

809   **Phenotypic effects of listening genotypes**

810   When sexual conflict is ongoing, listening alleles have sex-independent effects on offspring

811   responses to maternal and paternal information; whereas when sexual conflict is resolved,

812   listening alleles have sex-specific effects on offspring responses. Note that an individual's

813   overall genotype is determined by the alleles it carries at each locus.

	Locus	Genotype	Phenotype
Ongoing conflict	A	aa	Offspring do not listen to father's information
		Aa	Offspring listen to father's information 50% of the time
		AA	Offspring listen to father's information 100% of the time
	B	bb	Offspring do not listen to mother's information
		Bb	Offspring listen to mother's information 50% of the time
		BB	Offspring listen to mother's information 100% of the time
Resolved conflict	C	cc	Sons do not listen to father's information
		Cc	Sons listen to father's information 50% of the time
		CC	Sons listen to father's information 100% of the time
	D	dd	Sons do not listen to mother's information
		Dd	Sons listen to mother's information 50% of the time
		DD	Sons listen to mother's information 100% of the time
	E	ee	Daughters do not listen to father's information
		Ee	Daughters listen to father's information 50% of the time
		EE	Daughters listen to father's information 100% of the time
	F	ff	Daughters do not listen to mother's information
		Ff	Daughters listen to mother's information 50% of the time
		FF	Daughters listen to mother's information 100% of the time

FIGURE 1

Lifecycle of organisms in the ongoing conflict model

At the start of the lifecycle, adult individuals emerge with a default phenotype suited to
benign conditions that is fixed for the lifetime (1). A stress-resistant phenotype is also
possible but must be induced transgenerationally. Individuals experience stress when they
interact with too many conspecifics. Stress induces the production of a nongenetic factor (2),
which is later transferred to offspring. The extent to which an individual's phenotype matches
its experience of stress determines its condition (3). For males, condition determines mating
success (4), whereas for females, condition determines fecundity (5). Females mate with the
most competitive male in their mating neighbourhood, or with a random male as a control. At
reproduction, both parents faithfully pass on their nongenetic factors (6) as well as listening
alleles (7). Offspring that carry listening alleles utilise nongenetic information from parents
(8) to develop a stress-resistant phenotype (9). Offspring without listening alleles ignore
parental information and develop a default phenotype suited to benign conditions (1). The
same lifecycle applies to the resolved conflict model except that sex-specific listening loci C,
D, E and F replace loci A and B.

FIGURE 2

Sex-independent listening (a) and sex-specific listening (b)

Graphs on the left (a) show the proportion of runs of the best male version of the ongoing

conflict model ending in sex-independent paternal TGP (dark grey), maternal TGP (medium

grey), both types of TGP (black), and no TGP (light grey) at generation 1000. Graphs on the

right (b) show the proportion of simulation runs of the best male version of the resolved

conflict model ending in sex-specific TGP at generation 1000. Note that sexual selection

occurs under best male settings and increases in intensity with number of patches in the

mating neighbourhood. Colours in the legend represent all possible combinations of listening.

The first letter in each double-letter code represents the sex of the offspring that does the

listening (M = males (i.e., sons); F = females (i.e., daughters); B = both sexes), and the

second letter represents the sex of the parent that is listened to (M = males (i.e., fathers); F =

females (i.e., mothers)). Thus, for example, BM indicates a sex-independent paternal effect

(i.e., sons and daughters both listening to fathers), and MF indicates a son-specific maternal

effect (i.e., sons listening to mothers). Colours with more than one double-letter code indicate

runs where more than one combination of sex-independent or sex-specific TGP evolves.

Runs were counted as ending in TGP if listening allele frequencies were ≥ 0.95. In (a), when

one parent experiences higher stress than the other, intralocus conflict is generated between

sons and daughters over whether to pay attention to information received from the stressed or

unstressed parent. In (b), sex-specific listening loci allow sexual conflict to be resolved via

the evolution of sex-specific TGP.

FIGURE 3

Patterns of phenotypic matching (∂_{ind})

Histograms show the outcome of selection on males (blue) and females (red) at the end of a
single simulation run of the best male version of the ongoing conflict model (a) and resolved
conflict model (b) at generation 1000. Note that sexual selection occurs under best male
settings. In (a), sex-differences in the distribution of matching is indicative of intralocus
sexual conflict and occurs when one parent experiences higher stress than the other. These
divergent distributions become bimodal due to the opposing action of sexual selection on
males and fecundity selection on females. The resolution of sexual conflict in (b) allows each
sex to achieve maximal matching at the same time. For the unstressed sex in both (a) and (b),
distributions of ∂_{ind} show little variation around the peak because individuals of this sex
rarely encounter more conspecifics than their sex's encounter threshold (α_m or α_f), and so
are almost always uniformly well-matched. Other settings: mating neighbourhood size = 25
patches.

FIGURE 4

Opposing selection on the sexes generates listening polymorphisms

Graphs show the frequencies of allele *A* (paternal TGP; blue) and allele *B* (maternal TGP;

red) for a single simulation run of the best male version of the ongoing conflict model when

males experience higher stress (a) and females experience higher stress (b). Note that sexual

selection occurs under best male settings. Listening polymorphisms are consistently

maintained for hundreds of generations. Other settings: mating neighbourhood size = 25

patches.

SUPPLEMENTARY FIGURE S1

Sex-independent listening (a) and sex-specific listening (b) when environmental stress is

periodic

Fluctuating stress generates patterns of TGP that are very similar to those resulting from a

single, extended bout of elevated stress (see main text). The plot shows the proportion of 40

simulation runs ending in listening outcomes at generation 1000 when conditions alternate

between benign and stressful every 25 generations, causing females or both sexes to

experience fluctuating stress. Other settings are: 'best male' version of each model; mating

neighbourhood size = 25 patches.

(a) Ongoing conflict   (b) Resolved conflict

The sex that experiences stress: Neither, Father, Mother, Both

Proportion of lifetime spent phenotypically matched (∂_{ind})

Phenotypic matching: □ In males □ In females

SUPPLEMENTARY FIGURE S2

Patterns of phenotypic matching (∂_{ind}) when mating is random

Histograms show the outcome of selection on males (blue) and females (red) at the end of a single simulation run of the 'random mating' version of the ongoing conflict model (a) and resolved conflict model (b) at generation 1000. The absence of sexual selection on males allows fecundity selection to maximise female matching in (a) regardless of which parent experiences stress, and limits maximal matching in males in (b). For the unstressed sex in both (a) and (b), distributions of ∂_{ind} show little variation around the peak because individuals of this sex rarely encounter more conspecifics than their sex's encounter threshold (α_m or α_f), and so are almost always uniformly well-matched. Other settings: mating neighbourhood size = 25 patches.

SUPPLEMENTARY FIGURE S3

Sex-independent listening (a) and sex-specific listening (b) when mating is random

Graphs on the left (a) show the proportion of runs of the random mating version of the ongoing conflict model ending in sex-independent paternal TGP (dark grey), maternal TGP (medium grey), both types of TGP (black), and no TGP (light grey) at generation 1000. Graphs on the right (b) show the proportion of simulation runs of the random mating version of the resolved conflict model ending in sex-specific TGP at generation 1000. Colours in the legend represent all possible combinations of listening. The first letter in each double-letter code represents the sex of the offspring that does the listening (M = males (i.e., sons); F = females (i.e., daughters); B = both sexes), and the second letter represents the sex of the parent that is listened to (M = males (i.e., fathers); F = females (i.e., mothers)). Thus, for example, BM indicates a sex-independent paternal effect (i.e., sons and daughters both listening to fathers), and MF indicates a son-specific maternal effect (i.e., sons listening to mothers). Colours with more than one double-letter code indicate runs where more than one combination of sex-independent or sex-specific TGP evolves. Runs were counted as ending in TGP if listening allele frequencies were ≥ 0.95.