Spatial attention and expectations rely on modality-specific and multisensory

mechanisms

Authors: Arianna Zuanazzi[1,2], Uta Noppeney[1,3]

[1]Computational Neuroscience and Cognitive Robotics Centre, University of Birmingham,

UK; [2]Department of Psychology, New York University, New York, NY, USA; [3]Donders

Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, The

Netherlands

Corresponding author: Arianna Zuanazzi, Department of Psychology, New York University,

New York, NY, USA, email: az1864@nyu.edu

Number of words: 8732

Abstract

In our natural environment, the brain needs to combine signals from multiple sensory modalities into a coherent percept. While spatial attention guides perceptual decisions by prioritizing processing of signals that are task-relevant, expectations encode the probability of signals over space. Previous studies have shown that behavioral effects of spatial attention generalize across sensory modalities. However, because they manipulated spatial attention as signal probability over space, these studies could not dissociate attention and expectation.

In two experiments, we orthogonally manipulated spatial attention (i.e., task-relevance) and expectation (i.e., signal probability) selectively in one sensory modality (i.e., primary) (experiment 1: audition, experiment 2: vision) and assessed their effects on primary and secondary sensory modalities in which attention and expectation were held constant.

Our results show behavioral effects of spatial attention and expectation that are comparable for audition or vision as primary modalities; yet, spatial expectations were formed more slowly in audition. Critically, spatial attention and expectation affected responses more strongly in the primary modality where they were manipulated and generalized to the secondary modality only in an attenuated fashion. Collectively, our results suggest that both spatial attention and expectation rely on modality-specific and multisensory mechanisms.

Keywords

Multisensory, space, attention, expectation, perceptual inference

Public significance of the study

In our natural environment, our senses are bombarded with myriads of signals. How does the brain allocate spatial attention and form spatial expectations across the senses? Previous studies have shown that behavioral effects of spatial attention generalize across sensory

modalities. Yet, these studies conflated attention and expectation, because they manipulated spatial attention as signal probability over space.

This psychophysics study orthogonally manipulated spatial attention (i.e., task-relevance) and expectation (i.e., signal probability) selectively in the auditory (resp. visual) modality and evaluated their effects on target detection in the visual modality (i.e., audition to vision) and vice versa (i.e., vision to audition). We show that spatial attention and expectation operate similarly in audition and vision; yet, spatial expectations are learnt more slowly in audition than vision. Our results suggest that spatial attention and expectation rely on modality-specific and multisensory mechanisms that enable partial generalization from audition to vision and vice versa.

Introduction

Spatial attention is a top-down mechanism that is critical for the selection of task-relevant information. It facilitates perception (e.g., faster reaction times, greater accuracy) of signals presented at the attended location (Carrasco, 2011; van Ede et al., 2012). By contrast, spatial expectation (signal probability) facilitates perception by encoding the statistical structure of the environment (Summerfield & Egner, 2009; Rohenkohl et al., 2014). In everyday life, spatial attention and expectation are closely intertwined. For instance, observers often allocate attentional resources to locations in space where events are more likely to occur (Summerfield & Egner, 2009; Feldman & Friston, 2010 for further discussion within the predictive coding framework).

Importantly, in our natural multisensory environment our brain is constantly exposed to auditory and visual signals. This raises the critical question of whether allocation of attention and encoding of signal probability are performed in a modality-specific fashion or interactively across sensory modalities. Previous research has suggested that spatial attention relies on cognitive resources that are partially shared across sensory modalities (Eimer & Driver, 2001; Wahn & König 2015, 2017). For instance, Spence & Driver (1996) manipulated spatial attention by presenting signals with a higher probability in the attended relative to unattended hemifield in one modality only (i.e., primary modality). They showed behavioral facilitation for signals presented at the attended location not only for the primary modality (e.g., audition) but also for the secondary modality (e.g., vision) in which spatial attention was not explicitly manipulated. Likewise, neuroimaging studies showed increased activations for signals presented at the attended location not only in the primary modality in which attention was manipulated but also in the secondary modality (Eimer & Schröger, 1998; Eimer, 1999; Macaluso et al., 2002; Santangelo et al., 2009; Zuanazzi & Noppeney,

2019). Crucially, in this past work the attentional effects were greater in the primary than in the secondary modality (Spence & Driver, 1996; Mondor & Amirault, 1998). The attenuated generalization across sensory modalities suggests that attentional resources are not supramodal, but partially shared (Driver & Spence, 1998).

However, this past research conflated attention and expectation by manipulating spatial attention via probabilistic spatial cues or changes in signal probability (Posner, 1980; Spence & Driver, 1996, 1997; Macaluso et al., 2002; Kincade et al., 2005; Bressler et al., 2008; Santangelo et al., 2009). Notably, the Posner cuing paradigm shifts observers' attention via spatial cues that indicate whether a target is for instance more likely to appear the left or right hemifield. This raises the question of whether spatial attention and/or expectation generalize across sensory modalities to a similar extent when they are manipulated independently. A recent neuroimaging study for instance suggested that spatial attention relies mainly on frontoparietal cortices for both primary and secondary modalities, while spatial expectations are formed in sensory systems only for the primary modality (Zuanazzi & Noppeney, 2019). As a consequence, we would expect spatial signal probability to be encoded selectively for the primary modality and generalize to a secondary modality only to a very limited degree.

A second unresolved question is whether the generalization across sensory modalities depends on whether attention and expectation are manipulated in the auditory or visual modalities as primary manipulation modality – i.e., on the direction of cross-sensory generalization. Previous studies of multisensory attention have indeed shown asymmetric multisensory generalization, depending on which modality was manipulated as primary modality (Ward at al., 2000; Greene et al., 2001; Molholm et al., 2007). Moreover, one may expect differences in cross-sensory generalization from vision to audition and vice versa because spatial representations and expectations are learnt and encoded differently in audition and vision. First, visual and auditory systems encode space via different reference frames

(i.e., eye-centered vs head-centered) and representational formats. In the visual system spatial

location is directly encoded in the sensory epithelium and later in a place code, i.e., via

retinotopic organization of primary and higher order visual cortices (e.g., Sereno et al., 1995;

Maier & Groh, 2009). In the auditory system, spatial locations are computed from binaural

and monoaural cues in the brain stem and are represented in a hemifield code in primary

auditory cortices (e.g., Lauter et al., 1985; Maier & Groh, 2009). Further, in everyday life

under normal lighting conditions vision usually provides more reliable spatial information

than audition and therefore often dominates spatial perception (Spence & Driver, 1997; Aller

et al., 2015; Odegaard et al., 2015; Rohe & Noppeney, 2015a, 2015b. 2016, 2018; Aller &

Noppeney, 2019; Jones et al., 2019; Meijer et al., 2019). As a result, we would expect the

generalization of spatial attention and expectation to depend on whether attention and

expectation are manipulated primarily in vision or audition.

To summarize this study was designed to address two questions: First, whether spatial

attention and expectations (when manipulated independently) rely on amodal or modality-

specific mechanisms. Second, whether the cross-sensory generalization differ when operating

from vision to audition or vice versa, if spatial expectations rely on partially shared

mechanisms. In two experiments (within participants), we orthogonally manipulated spatial

attention and expectation selectively in one sensory modality as primary modality

(experiment 1: audition, experiment 2: vision). We assessed the effects of spatial attention

and expectation in this primary modality and how they generalize crossmodally to signals in a

secondary sensory modality, in which spatial attention and expectation were not explicitly

manipulated (experiment 1: vision, experiment 2: audition). We expected that both spatial

attention and expectations rely on mechanisms that are partially shared across sensory

modalities and that the generalization of spatial expectations depends on whether they are

manipulated in vision or audition as primary modalities.

Materials and Methods

Participants

Twenty-eight healthy subjects (19 females; mean age = 25.57 years; 24 right-handed)

participated in the study (experiment 1 and experiment 2, within participants). A power

analysis (G*, Faul et al., 2009) showed that at least 14 and 26 subjects were needed to detect

effect sizes (i.e., Cohen's d_{av}, calculated as M_{diff} / ((SD$_1$ + SD$_2$) / 2)) of 0.18 and 0.20

respectively, with a power (1-β) of 0.8 and α = 0.05. These effect sizes were estimated in a

previous study that we conducted with the same experimental design as experiment 1

(Zuanazzi & Noppeney, 2018) and correspond to the simple main effects of expectation in

the attended and unattended conditions.

All participants had normal or corrected to normal vision and reported normal hearing. All

participants provided written informed consent and were naïve to the aim of the study. The

study was approved by the local ethics committee of the University of Birmingham (Science,

Technology, Mathematics and Engineering (STEM) Ethical Review Committee) and the

experiment was conducted in accordance with these guidelines and regulations.

Stimuli and Apparatus

171 Spatial auditory stimuli of 100 ms duration were created by convolving bursts of white noise

172 (with 5 ms onset and offset ramps) with spatially selective head-related transfer functions

173 (HRTFs) based on the KEMAR dummy head of the MIT Media Lab

174 (http://sound.media.mit.edu/resources/KEMAR.html). Visual stimuli ('flashes') were white

175 discs (radius: 0.88º visual angle, luminance: 196 cd/m2) of 100 ms duration presented on a

grey background. Both auditory and visual stimuli were presented at ±10º of horizontal visual angle along the azimuth (0º of vertical visual angle). Throughout the entire experiment, a fixation cross was presented in the center of the screen.

Prior to the beginning of the study, participants were tested for their ability to discriminate left and right auditory stimuli on a brief series of 20 trials. They indicated their spatial discrimination response (i.e., 'left' vs 'right') via a two-choice key press (group mean accuracy was $99\% \pm 0.4\%$ [across subjects mean ± SEM]).

During the experiment, participants rested their chin on a chinrest with the height held constant across all the participants. Auditory stimuli were presented at approximately 72 dB SPL, via HD 280 PRO headphones (Sennheiser, Germany). Visual stimuli were displayed on a gamma-corrected LCD monitor (2560 x 1600 resolution, 60 Hz refresh rate, 30" Dell UltraSharp U3014, USA), at a viewing distance of approximately 50 cm from the participant's eyes. Stimuli were presented using Psychtoolbox version 3 (Brainard, 1997; www.psychtoolbox.org), running under Matlab R2014a (Mathworks Inc., Natick, MA, USA) on a Windows machine. Participants' responses were recorded via one key of a small keypad (Targus, USA). Throughout the study, participants' eye-movements and fixation were monitored using Tobii Eyex eyetracking system (Tobii EyeX, Tobii, Sweden, ~60 Hz sampling rate).

Study overview: rationale and analysis strategy

This study included two experiments. Each experiment conforms to a four-factorial design. Because the two experiments were performed within the same participants the study as a whole could also be treated as a five factorial within-subject experiment. However, because (1) experiment 1 and 2 were completed on different days, (2) experiment 1 was a replication

of our previous study and (3) the understanding of results of five factorial designs is rather

complex, we will initially analyse each of the two experiments separately.

The separate analyses of experiments 1 and 2 allow us to address our first question, i.e.,

whether spatial attention and expectations rely on modality-specific or at least partially

shared mechanisms. While experiment 1 is intended to replicate our findings reported in our

previous research, experiment 2 is intended to extend them and demonstrate that this pattern

of results does not depend on whether audition or vision is used as a primary manipulation

modality. Moreover, showing the same profile across the two experiments also resolves the

ambiguity of our previous research, in which a smaller spatial expectation effect for the

secondary (i.e. visual) modality in experiment 1 could potentially be explained by differences

in sensory modality rather than attenuated cross-sensory generalization.

In a second step we will directly combine data from experiment 1 and 2 to address our second

question, i.e. whether the spatial expectation effect generalizes differently from audition to

vision than from vision to audition. To address this question, we need to compare the

expectation effects for auditory and visual stimuli in the attended hemifield between the two

experiments (i.e. this question cannot be addressed by any of the two experiments alone).

Please also note that the Design and Procedure section mostly overlaps with that of our

previous paper (Zuanazzi & Noppeney, 2018) to enable the reader to quickly compare our

different studies and obtain a convergent picture across all results.

Design and Procedure

Experiment 1 investigated the effect of auditory spatial attention and expectation on detection

of auditory (i.e., primary modality) and visual (i.e., secondary modality) targets using a 2

(auditory spatial attention: left vs right hemifield) x 2 (auditory spatial expectation: left vs

right hemifield) x 2 (stimulus modality: auditory vs visual) x 2 (stimulus location: left vs

right hemifield) factorial design. Hence, experiment 1 manipulated spatial attention and

expectation selectively in audition and assessed their direct effects on auditory stimulus

processing and indirect generalization to visual stimuli. In experiment 2 primary and

secondary modality were reversed (i.e., primary modality: vision; secondary modality:

audition); design and procedural details were otherwise comparable to experiment 1. For the

data analysis we pooled over stimulus locations (left/right) leading to a 2 (auditory spatial

attention: attended vs unattended) x 2 (auditory spatial expectation: expected vs unexpected)

x 2 (stimulus modality: auditory vs visual) factorial design.

Spatial attention was manipulated for the primary modality as task-relevance, i.e., the

requirement to respond to an auditory (experiment 1) or a visual (experiment 2) target in the

left vs right hemifield. Prior to each run a cue (duration: 2000 ms) informed the observer

whether to respond to targets in either their left or right hemifield.

Spatial expectation was manipulated as spatial signal probability for signals in the primary

modality across experimental sessions that were performed on different days. Auditory (i.e.,

primary modality in experiment 1) or visual (i.e., primary modality in experiment 2) signals

were presented with a ratio of 2.33/1 (i.e., 70%/30%) in the expected/unexpected hemifield.

Observers were not informed about those probabilities but learned them implicitly.

Importantly, spatial attention and expectation were not directly manipulated in the secondary

modality, allowing us to assess their cross-sensory generalization. As a result, participants

needed to respond to all visual targets that were presented with equal probabilities in their

spatial hemifields in experiment 1 (i.e., ratio 1/1 in the expected/unexpected hemifields) (Fig.

1A and 1C). Likewise, they had to respond to all auditory targets that were presented with

equal probabilities in experiment 2.

Each experiment included two sessions (i.e., spatial expectation left vs right on different

days). Hence, subjects participated in the two experiments on four days: 2 sessions for

experiment 1 and 2 sessions for experiment 2 = 4 sessions in total for each participant. Each session included 12 attention runs. Runs were of two types: in run type A (Fig. 1A, 1C and 1D) spatial attention and expectation were congruent (i.e., spatial attention was directed to the hemifield with higher stimulus frequency); in run type B spatial attention and expectation were incongruent (i.e., spatial attention was directed to the hemifield with less frequent stimuli). The overall probability to respond (i.e., response probability) was greater when attention and expectation were congruent and directed to the same hemifield (85%, runs of type A) than when they were directed to different hemifields (65%, runs of type B) (Fig. 1D). The order of experiments 1 vs 2 and of expectation sessions (i.e., left vs right) was counterbalanced across participants; the order of attention runs (i.e., left vs right) was counterbalanced within and across participants and the order of stimulus locations (i.e., left vs right) and stimulus modalities (sound vs flash) was pseudo-randomized within each participant. Brief breaks were included after every two attention runs to provide feedback to participants about their performance accuracy (averaged across all conditions) in the target detection task and about their eye-movements (i.e., fixation maintenance).

Overall, each experiment included 80 trials x 12 attention runs (6 runs of type A and 6 runs of type B, duration: 3 mins/run) x 2 expectation sessions = 1920 trials in total (and 3840 for the whole study). Specifically, each run type included i. 336 targets presented in the expected hemifield (pooled over left and right) and 144 targets in the unexpected hemifield (pooled over left and right) for the primary modality and ii. 240 targets presented in the expected hemifield and 240 targets in the unexpected hemifield (pooled over left and right) for the secondary modality. For further details see Fig. 1C which shows the absolute number of trials for each condition and run type.

Each trial (SOA: 2300 ms) included three time windows: i. fixation cross alone (700 ms duration), ii. brief sound or flash (stimulus duration: 100 ms) and iii. fixation cross alone

(1500 ms as response interval, see Fig. 1B). Participants responded to the targets in the

primary modality presented in the attended hemifield and to all targets in the secondary

modality irrespective of hemifield via key press, with their index finger (i.e., the same

response for all auditory and visual stimuli) as fast and accurately as possible (Fig. 1B).

Prior to each session, participants were familiarized with the stimuli in brief practice runs

(with equal spatial signal probability) and trained on target detection performance and

fixation (i.e., a warning signal was shown when the disparity between the central fixation

cross and the eye-data samples exceeded 2.5 degrees).

After the final session (i.e., experiment 1 for 14 participants and experiment 2 for the other 14

participants), participants indicated in a questionnaire whether they thought the stimuli in the

primary modality were presented more frequently in one of the two spatial hemifields. Ten

out of 14 participants in experiment 1 and 11 out of 14 participants in experiment 2 correctly

identified the expectation manipulation in the primary modality. Moreover, 13 out of 14

participants in experiment 1 and 13 out of 14 participants in experiment 2 correctly reported

that stimuli in the secondary modality were presented with equal probabilities across the two

hemifields. These data suggest that the majority of participants were aware of the

manipulation of signal probability at least at the end of the fourth session.

Figure 1: Design and example trials of experiment 1 (audition to vision)

A. Experiment 1: auditory spatial attention and expectation (i.e., signal probability) were manipulated in a 2 (auditory modality – dark orange, vs visual modality – light blue) x 2 (attended hemifield vs unattended hemifield) x 2 (expected hemifield vs unexpected hemifield) factorial design. For illustration purposes, stimulus locations (left/right) were collapsed. Presence vs absence of response requirement is indicated by the hand symbol, spatial signal probability manipulation is indicated by the %. **B.** Experiment 1: example of two trials in a session where auditory stimuli were presented with a probability of 0.7 in the left hemifield and 0.3 in the right hemifield. At the beginning of each run (i.e., 80 trials), a cue informed participants whether to attend and respond to auditory signals selectively in their left or right hemifield throughout the entire run. On each trial participants were presented with an auditory or visual stimulus (100 ms duration) either in their left or right hemifield. They were instructed to respond to auditory stimuli only in the attended hemifield

and to all visual stimuli irrespective of the hemifield as fast and accurately as possible with their index finger. The response window was limited to 1500 ms. Participants were not explicitly informed that auditory signals were more likely to appear in one of the two hemifields. Instead, spatial expectation was implicitly learnt within a session (i.e., day).

C. Experiment 1: number of auditory (dark orange) and visual (light blue) trials in the 2 (attended vs unattended hemifield) x 2 (expected vs unexpected hemifield) design (pooling over left/right stimulus location). Presence vs absence of response requirement is indicated by the hand symbol. The fraction of the area indicated by the 'Response' hand symbol pooled over the two bars of one particular run type (e.g., run type A) represents the response related expectation (i.e., general response probability: the overall probability that a response is required on a particular trial); general response probability is greater for run type A (85%), where attention and expectation are congruent, than for run type B (65%), where attention and expectation are incongruent, as indicated in **D**.

Note. Design and procedure of experiment 2 were comparable to that of experiment 1, with the only difference that vision was the primary modality and audition was the secondary modality. In other words, in experiment 2 attention and expectation were manipulated selectively in vision.

Data analysis

Eye movement: exclusion criteria

We excluded trials where participants did not successfully fixate the central cross based on a dispersion criterion (i.e., distance of fixation from subject's center of fixation, as defined in calibration trials, > 1.3 degrees for three subsequent samples, Blignaut, 2009). Our eyetracking data confirmed that participants successfully maintained fixation in both experiments with only a small number of trials to be excluded (experiment 1: excluded auditory response trials $1.8\% \pm 0.5\%$ [across subjects mean $\pm$ SEM]; excluded visual response trials $1.7\% \pm 0.5\%$ [across subjects mean $\pm$ SEM]; experiment 2: excluded visual response trials $2.7\% \pm 1\%$ [across subjects mean $\pm$ SEM]; excluded auditory response trials $2.7\% \pm 0.9\%$ [across subjects mean $\pm$ SEM]).

Response time analysis - separately for experiments 1 and 2

We initially analysed response times separately for primary and secondary modalities and independently for experiments 1 and 2.

Response time (RT) analysis was limited to trials with RT within the 1500 ms response window and was performed after pooling over stimulus location (left/right).

For the primary modality (i.e., experiment 1 = audition, experiment 2 = vision), subject-specific median RT were entered into a two-sided paired-sample t-test with spatial expectation (expected vs unexpected stimulus) as factor (observers did not respond to targets in the primary modality in the 'unattended' hemifield). For the secondary modality (i.e., experiment 1 = vision, experiment 2 = audition), subject-specific median response time were entered into a 2 (spatial attention: attended vs unattended stimulus) x 2 (spatial expectation: expected vs unexpected stimulus) repeated measures analysis of variance (ANOVA).

For both experiments 1 and 2, the mean hit rates were very high (> 99% in all conditions, Table 1), indicating that participants accurately performed the detection task. Because of the absence of a substantial number of misses, hit rates were not further analyzed.

Response time analysis - combined for experiments 1 and 2

Next, we investigated whether the effect of spatial expectation (i.e., expected vs unexpected) in the attended hemifield (no response was required for stimuli in the primary modalities presented in the unattended hemifield) depended on i. whether targets were presented in the primary or secondary modalities (i.e., the extent to which spatial expectations generalize across the senses) and ii. the multisensory generalization direction, from audition to vision and from vision to audition (i.e., whether spatial expectations generalize differently depending on whether audition or vision is the primary modality). Hence, we first computed the difference in median RT (ΔRT_{Exp}) between unexpected and expected stimuli presented in the attended hemifield (which corresponds to ΔRT_{Exp} between attended stimuli in run type B and run type A) for targets in each experiment, yielding four conditions: i. auditory targets as primary modality (experiment 1), ii. visual targets as secondary modality (experiment 1), iii. visual targets as primary modality (experiment 2), iv. auditory targets as secondary modality (experiment 2). ΔRT_{Exp} were entered into a 2 (multisensory generalization direction: audition to vision vs vision to audition) x 2 (manipulation: primary/direct vs secondary/indirect modality) repeated measures ANOVA. Please note that the interaction then reflects the difference between targets in the auditory and visual modality.

Time course of response times - combined for experiment 1 and 2

Finally, we assessed how these effects of multisensory generalization direction and direct/indirect manipulation evolved over time. For this, we computed the difference in

median RT (ΔRT_{Exp}) between unexpected and expected stimuli, as in the previous analysis,

but now separately for the first and second half of the experiment (i.e., one half = 430 trials).

These ΔRT_{Exp} (for each half) were entered in a 2 (multisensory generalization direction:

audition to vision vs vision to audition) x 2 (manipulation: primary/direct vs

secondary/indirect modality) x 2 (time: first vs second half of the experiment) repeated

measures ANOVA.

Unless otherwise indicated, we only report effects that are significant at $p < 0.05$

(Greenhouse-Geisser corrected where appropriate).

Results

Generalization of attention and expectation effects across modalities – separately for

experiment 1 and 2

In experiment 1, participants responded to auditory targets presented in their attended

hemifield and to all visual targets. In experiment 2, participants responded to visual targets

presented in their attended hemifield and to all auditory targets.

We observed qualitatively similar effects across experiments 1 and 2. Table 1 shows RT

(across participants' mean ± SEM) for targets in the auditory and visual modalities for the

two experiments.

For the primary modality, the two-sided paired-sample t-tests showed significantly faster RT

in the attended hemifield, when this hemifield was expected than unexpected (experiment 1,

auditory modality: $t(27) = -2.83$, $p = 0.009$, Cohen's d_{av} [95% CI] = -0.16 [-0.27 , -0.04];

experiment 2, visual modality: $t(27) = -9.62$, $p < 0.001$, Cohen's d_{av} [95% CI] = -0.35 [-0.47 ,

-0.23]) (Table 1, Fig. 2A and 2B).

For the secondary modality, the 2 (attended vs unattended) x 2 (expected vs unexpected)

repeated measures ANOVAs revealed a significant main effect of attention (experiment 1,

visual modality: $F(1, 27) = 72.08$, $p < 0.001$, η_p^2 [90% CI] = 0.73 [0.55, 0.80]; experiment 2,

auditory modality: $F(1, 27) = 36.91$, $p < 0.001$, η_p^2 [90% CI] = 0.58 [0.34, 0.70]). Results

showed that participants responded faster to targets presented in their attended than

unattended hemifields.

Moreover, a significant crossover interaction between attention and expectation was observed

(experiment 1, visual modality: $F(1, 27) = 10.09$, $p = 0.004$, η_p^2 [90% CI] = 0.27 [0.06, 0.46];

experiment 2, auditory modality: $F(1, 27) = 44.10$, $p < 0.001$, η_p^2 [90% CI] = 0.62 [0.40,

0.73]) (Table 1, Fig. 2A and 2B). The simple main effects showed that participants responded

significantly faster to targets in their attended hemifield, when this hemifield was expected

than unexpected (experiment 1, visual modality: $t(27) = -2.81$, $p = 0.009$, Cohen's d_{av} [95%

CI] = -0.08 [-0.15 , -0.02]; experiment 2, auditory modality: $t(27) = -5.96$, $p < 0.001$, Cohen's

d_{av} [95% CI] = -0.14 [-0.19 , -0.08]). This significant simple main effect demonstrates that

the effects of spatial attention and expectation generalized from primary to secondary

modalities, where neither attention not expectation were explicitly manipulated. By contrast,

participants responded significantly more slowly to targets in the secondary modality in the

unattended hemifield, when this hemifield was expected than unexpected (experiment 1,

visual modality: $t(27) = 2.56$, $p = 0.016$, Cohen's d_{av} [95% CI] = 0.09 [0.02, 0.17];

experiment 2, auditory modality: $t(27) = 5.53$, $p < 0.001$, Cohen's d_{av} [95% CI] = 0.18 [0.10 ,

0.26]) (Table 1, Fig. 2A and 2B). We suggest that simple main effects for expectation show

opposite directions because of response inhibition. In the attended hemifield observers need

to respond to the stimuli in the primary modality. Hence, if stimuli from the primary modality

are frequent (i.e., expected) in the attended hemifield, observers need to respond on a large

percentage of trials. By contrast, in the unattended hemifield observers should not respond to

the stimuli in the primary modality. Hence, if stimuli in the primary modality are frequent (i.e., expected) in the unattended hemifield, observers need to inhibit their response on a large percentage of trials.

Table 1: Group mean hit rates and reaction times (RT) and for each stimulus modality in each condition that required a response for experiment 1 (primary modality: audition, secondary modality: vision) and experiment 2 (primary modality: vision, secondary modality: audition). In experiment 1, participants responded only to attended auditory targets (and to all visual targets); in experiment 2 participants responded only to attended visual targets (and to all auditory targets). Standard errors (SEM) are given in parentheses.

	Auditory modality				**Visual modality**			
Experiment 1	+att +exp	+att -exp	-att +exp	-att -exp	+att +exp	+att -exp	-att +exp	-att -exp
Hit rate (%) (SEM)	99.5 (0.19)	99.5 (0.15)	/	/	99.7 (0.07)	99.7 (0.07)	99.3 (0.16)	99.4 (0.13)
RT (ms) (SEM)	599.7 (20.1)	616.2 (19.2)	/	/	514.4 (16.9)	522.1 (16.9)	583.8 (18.5)	574.4 (18.9)
Experiment 2	+att +exp	+att -exp	-att +exp	-att -exp	+att +exp	+att -exp	-att +exp	-att -exp
Hit rate (%) (SEM)	99.8 (0.08)	99.9 (0.03)	99.2 (0.22)	99.4 (0.18)	99.7 (0.09)	99.7 (0.09)	/	/
RT (ms) (SEM)	527.4 (20.2)	542.1 (20)	605.2 (25.7)	581.3 (24.2)	526.3 (17.9)	561.8 (19.9)	/	/

The effects of spatial expectation – combined for experiment 1 and 2

In the previous analysis we showed that the effects of expectation on response times generalized from the primary to the secondary modality. Next, we directly compared the effects of spatial expectation across the two experiments, in which either audition was the primary and vision the secondary modality or vice versa. The 2 (multisensory generalization direction: audition to vision vs vision to audition) x 2 (manipulation: primary/direct vs secondary/indirect modality) repeated measures ANOVA revealed a significant main effect of manipulation ($F(1, 27) = 18.03$, $p < .001$, η_p^2 [90% CI] $=0.40$ [0.16, 0.56]), showing overall greater expectation effects for primary than secondary modalities (dark vs light bars in Fig. 2C). Moreover, we observed a significant main effect of crossmodal generalization direction ($F(1, 27) = 20.71$, $p < .001$, η_p^2 [90% CI] $= 0$ .43, [0.19, 0.59]), with a greater expectation effect (i.e., greater ΔRT_{Exp}) for vision to audition than for audition to vision (experiment 2 vs experiment 1 in Fig. 2C). Critically, this generalization effect may be greater from vision to audition than vice versa, because the expectation effect is stronger when vision is the primary modality (dark blue bar in Fig. 2C). Alternatively, the expectation effect generalizes more effectively from vision to audition than vice versa (light orange bar in Fig. 2C).

Time course of the effects of spatial expectation - combined for experiment 1 and 2

To disentangle between these two possibilities, we investigated how signal probability is learnt over time when audition (experiment 1) or vision (experiment 2) are the primary modality by repeating the previous analysis with the additional factor of time (i.e., first vs second half of experiment). The 2 (multisensory generalization direction: audition to vision vs vision to audition) x 2 (manipulation: primary/direct vs secondary/indirect modality) x 2 (time: first vs second half of the experiment) repeated measures ANOVA performed on

ΔRT_{Exp} revealed significant main effects of multisensory generalization direction ($F(1, 27)$ =

22.70, $p < 0.001$, η_p^2 [90% CI] = 0.46 [0.21, 0.61]), manipulation ($F(1, 27)$ = 13.77, $p <$

0.001, η_p^2 [90% CI] = 0.34 [0.10, 0.51]) as well as a significant interaction between

multisensory generalization direction x manipulation x time ($F(1, 27)$ = 11.38, $p = 0.002$, η_p^2

[90% CI] = 0.29 [0.07, 0.48]).

We unpacked the 3-way ANOVA into two 2-ways ANOVAs for further analysis, one for

each half of the experiment, with factors multisensory generalization direction (audition to

vision vs vision to audition) and manipulation (primary/direct vs secondary/indirect

modality).

For the first half of the experiment, our results revealed a significant main effect of

multisensory generalization direction ($F(1, 27)$ = 8.34, $p = 0.008$, η_p^2 [90% CI] = 0.24 [0.04,

0.42]), manipulation ($F(1, 27)$ = 28.84, $p < 0.001$, η_p^2 [90% CI] = 0.52 [0.27, 0.65]) and a

significant interaction between multisensory generalization direction and manipulation ($F(1,$

27) = 12.83, $p = 0.001$, η_p^2 [90% CI] = 0.32 [0.09, 0.50]) (left bar plot in Fig. 2D). Post-hoc

comparisons indicated that ΔRT_{Exp} in the primary modality of experiment 2 (i.e., vision) were

significantly greater than ΔRT_{Exp} in the primary modality of experiment 1 (i.e., audition)

($t(27)$ = 5.72, $p < 0.001$, Cohen's d_{av} [95% CI] = 1.05, [0.59, 1.50], dark blue vs dark orange

bars in the left bar plot of Fig. 2D), and greater than the effects of expectation in the

secondary modality of experiment 2 (i.e., audition) ($t(27)$ = 4.97, $p < 0.001$, Cohen's d_{av}

[95% CI] = 1.03, [0.53, 1.51], dark blue vs light orange bars in the left bar plot of Fig. 2D).

Moreover, the effects of expectation in the secondary modality of experiment 2 (i.e.,

audition) were significantly greater than those in the secondary modality of experiment 1

(i.e., vision) ($t(27)$ = 2.14, $p = 0.042$, Cohen's d_{av} [95% CI] = 0.48, [0.02, 0.94], light orange

vs light blue bars in the left bar plot of Fig. 2D).

For the second half of the experiment, our results only revealed a significant main effect of manipulation ($F(1, 27) = 11.41$, $p = 0.002$, η_p^2 [90% CI] = 0.30 [0.07, 0.48]) (right bar plot in Fig. 2D) but no significant main effect of multisensory generalization direction ($p = 0.20$) or significant interaction between multisensory generalization direction and manipulation ($p = 0.65$) was found.

To summarize, these results show that: (1) an effect of generalization direction (i.e., audition to vision vs vision to audition) was found only for the first half of the experiment. Here, the effects of expectation generalized crossmodally in an attenuated fashion only from vision to audition in experiment 2 (i.e., primary modality: vision) but no difference between audition and vision was found in experiment 1 (i.e., primary modality: audition). By contrast, in the second half of the experiment we did not observe an effect of multisensory generalization direction. Instead, the effects of expectation generalized from the primary to the secondary modality in an attenuated fashion similarly when vision or audition were the primary modality. As shown in figure 2D, this difference between first and second halves can be explained by the fact that observers form spatial expectations (i.e., learn signal probability over space) more slowly in audition than vision. Yet, once expectations are learnt in audition, the crossmodal generalization is comparable for audition and vision. In other words, the effect of generalization direction that we observed in our analysis that did not yet account for learning effects (i.e., our second analysis) can be explained away by the speed with which observers form spatial expectations in their primary modality. Spatial expectations are formed faster in vision than audition, but once they are formed, they generalize similarly from vision to audition and vice versa.

Figure 2: Behavioural results of experiment 1 and 2.

Bar plots represent across subjects' mean (±SEM) RT for each of the six conditions with response requirements for experiment 1 (primary modality: audition; secondary modality: vision, **A**) and 2 (primary modality: vision; secondary modality: audition, **B**), pooling over left/right stimulus location. Overall slower RT are observed for runs type B than runs type A, which reflects differences in general response probability (see Fig. 1D) **C.** Bar plots represent across subjects' mean (±SEM) ΔRT for effects of spatial expectation (attended unexpected – attended expected hemifield) in the primary (dark bars) and secondary modalities (light bars) for experiment 1 and 2. **D.** Effects of response probability (attended unexpected – attended

521   expected hemifield) over time (i.e., first and second half) for audition and vision as primary

522   (dark bars) or secondary (light bars) modality. Brackets and stars indicate significance of

523   main effects and interactions. *$p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. Audition: orange; vision:

524   blue.

Discussion

The current study investigated how observers allocate attention and form expectations over space across audition and vision. We orthogonally manipulated spatial attention as response requirement and expectation as stimulus probability over space selectively in the primary modality and assessed their effects on behavioral responses to targets presented in the primary and secondary modalities. Across two experiments we alternated the assignment of vision and audition to primary or secondary modality. This allowed us to compare behavioral effects of spatial attention and expectation in audition and vision and their crossmodal generalization.

Regardless of sensory modality we observed a significant main effect of spatial attention for targets in the secondary modality, in which attention was not directly manipulated. Auditory spatial attention partially generalized to the visual modality and vice versa. These findings converge with a large body of behavioral and neuroimaging work suggesting that attentional resources are allocated interactively across the senses (Spence & Driver, 1996; Eimer & Schröger, 1998; Eimer, 1999; Macaluso et al., 2002; Santangelo et al., 2009; Zuanazzi & Noppeney, 2019).

Likewise, in the attended hemifield observers were faster at target detection in the primary and secondary modalities when the hemifield was expected than unexpected (i.e., high > low signal probability). Again, this response facilitation for expected (relative to unexpected) spatial locations were observed irrespective of whether vision or audition served as primary modality. Yet, in the unattended hemifield, in which we could assess effects of expectation only for the secondary modality, we observed the opposite pattern, i.e., observers were slower at target detection for expected than unexpected hemifields. Combining these two results, we observed a significant interaction between spatial attention and expectation, for both vision and audition as secondary modalities. In a previous study (Zuanazzi & Noppeney, 2018) we

argued that this interaction profile between spatial attention and expectation is explained by attention and expectation jointly co-determining general response probability (i.e., the probability to respond regardless of the hemifield in which the signal is presented). More specifically, in runs in which attention and expectation are directed to the same hemifield (runs of type A, Fig. 1A, 1C and 1D), participants have to respond to 85% of trials of the entire run, but only to 65% trials in runs of type B in which attention and expectation are directed to different hemifields (i.e., general response probability, Fig. 1A, 1C and 1D). Hence, faster response times may result from an increase in alertness, arousal or motor preparation that is needed to respond on a large proportion of trials (i.e., attended/expected and unattended/unexpected conditions, run type A, Fig. 2A and 2B) (Mars et al., 2007; Bestmann et al., 2008). By contrast, in runs in which attention and expectation are directed to different hemifields (runs of type B, Fig. 1A, 1C and 1D), observers need to inhibit their response to the frequent stimuli of the primary modality in the expected hemifield and hence respond more slowly to targets in the secondary modality.

Critically, response probability does not depend on whether the signal is auditory or visual, but it is calculated as the probability that any signal is responded to. If the expectation effects result purely from amodal mechanisms (e.g., general alertness, arousal, motor preparation etc.) associated with changes in response probability, the expectation effects in the attended hemifield should be equal for primary and secondary modalities. By contrast, if expectations (i.e., auditory or visual signal probability) are formed at least partially in a modality-specific fashion, we should observe expectation effects (i.e., ΔRT_{Exp}) that are greater for the primary modality (where expectation was explicitly manipulated) than for the secondary modality (i.e., an attenuated crossmodal generalization).

To arbitrate between these two hypotheses, we analyzed ΔRT_{Exp} in a 2 (multisensory generalization direction) x 2 (manipulation) repeated measures ANOVA. This analysis

revealed a significant main effect of 'manipulation', i.e., whether the stimulus was presented in the primary modality (in which expectation was explicitly manipulated) or in the secondary modality. Consistent with additional modality-specific mechanisms of expectation, observers showed a greater expectation effect for targets in the primary than the secondary modality. These results strongly suggest that spatial expectations modulate perceptual decision making via both modality-specific and amodal response mechanisms. This duality of modality-specific and amodal mechanisms converge with recent neuroimaging findings which showed effects of expectation selective for auditory stimuli as primary modality in auditory cortices and higher-order frontoparietal systems (Zuanazzi & Noppeney, 2019). Potentially, the activation increases in auditory cortices may reflect prediction error signals based on modality-specific expectations (Friston, 2005), while higher frontoparietal systems may be associated with additional response-related processes.

Surprisingly, this repeated measures ANOVA also revealed a main effect of multisensory generalization direction with greater effects in experiment 2 (i.e., vision to audition) than experiment 1 (i.e., audition to vision). Our time course analysis showed that this difference between experiments does not reflect genuine differences in the effectiveness with which spatial expectations generalize from audition to vision and vice versa but differences in the speed with which spatial expectations are learnt in audition and vision. In the second half of the experiment in which the expectation effect in the primary modality is comparable between auditory (experiment 1) and visual (experiment 2) targets, we no longer observe a significant effect of multisensory generalization direction. In other words, observers are slower at learning spatial expectations (i.e., signal probability) in audition than in vision. But, once they have formed spatial expectations of comparable precision for audition and vision, they also generalize similarly from audition to vision and vice versa.

The difference in learning rates between vision and audition may result from how the brain forms spatial representations in vision and audition (Neumann et al., 1986). While visual cortices are retinotopically organized and hence directly represent spatial location in a place code (i.e., based on space, Sereno et al., 1995; Maier & Groh, 2009), primary auditory cortices are tonotopically organized (i.e., based on frequency, Lauter et al., 1985, Middlebrooks &, 1991; Maier & Groh, 2009). In audition, spatial locations are computed only indirectly from binaural amplitude and latency differences and from monoaural filtering cues. Moreover, visual objects tend to be more permanent across time, whereas source sounds are often transient and dynamic (Neisser, 1976; Neumann et al., 1986). Most importantly, in everyday life vision provides typically (i.e., under optimal lighting conditions) more reliable spatial information than audition (Dacey et al., 1992; Knudsen & Brainard, 1995; Stephen et al., 2002; Talsma et al., 2008; Mengotti et al., 2018; see also Molholm et al., 2007). In the current study, the high spatial reliability of the visual stimulus (i.e., a white disc) may also have contributed to the shorter time for participants to become aware of the manipulation of signal probability in vision. Critically, participants' awareness of such manipulation is evidenced by our questionnaires' results. Alternatively, even in absence of explicit awareness, the distribution of events or targets across space could have been learnt faster for more reliable visual than auditory signals (Miller & Pachella, 1973; Jabar & Anderson, 2015). To better understand the fine-grained temporal aspects of spatial expectation or signal probability learning across sensory modalities, future neuroimaging studies will need to characterize the time course of spatial learning across sensory systems.

In summary, our results suggest that the brain allocates spatial attention and forms spatial expectation to some extent interactively across audition and vision (Eimer & Schröger, 1998; Eimer, 1999; Macaluso, 2010; Zuanazzi & Noppeney, 2019). With respect to spatial attention, our results corroborate previous research. With respect to spatial

expectation, we show that they rely on modality-specific and amodal mechanisms. In support of modality-specific mechanisms we demonstrate that spatial expectations in the attended hemifield generalize from the primary to the secondary modality only in an attenuated fashion. In support of amodal response-related mechanisms, we demonstrate that, for both primary and secondary modalities, response times are closely related to the general response probability and associated processes of arousal, alertness and motor preparation. Critically, our learning analysis suggests that observers learn spatial probabilities more slowly in audition than vision, which may be related to their different spatial reliabilities. Once observers have formed comparable spatial expectations in audition, these generalize equally effectively from audition to vision as from vision to audition.

Declaration of Conflicting Interests

The authors declared that they had no conflicts of interest with respect to their authorship or the publication of this article.

Funding

This research was funded by ERC-2012-StG_20111109 multsens

References

Aller, M., Giani, A., Conrad, V., Watanabe, M., & Noppeney, U. (2015). A spatially collocated sound thrusts a flash into awareness. *Frontiers in Integrative Neuroscience*, 9(February), 1–8. https://doi.org/10.3389/fnint.2015.00016

Aller, M., & Noppeney, U. (2019). To integrate or not to integrate: Temporal dynamics of hierarchical Bayesian causal inference. *PLoS biology* (Vol. 17). https://doi.org/10.1371/journal.pbio.3000210

Bestmann, S., Harrison, L. M., Blankenburg, F., Mars, R. B., Haggard, P., Friston, K. J., & Rothwell, J. C. (2008). Influence of uncertainty and surprise on human corticospinal excitability during preparation for action. *Current Biology*, 18(10), 775–780. http://doi.org/10.1016/j.cub.2008.04.051

Blignaut, P. (2009). Fixation identification: the optimum threshold for a dispersion algorithm. Attention, *Perception & Psychophysics*, 71(4), 881–895.

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, 10(4), 433–436. http://doi.org/10.1163/156856897X00357

Bressler, S. L., Tang, W., Sylvester, C. M., Shulman, G. L., & Corbetta, M. (2008). Top-down control of human visual cortex by frontal and parietal cortex in anticipatory visual spatial attention. *The Journal of Neuroscience*, 28, 10056–10061. https://doi.org/10.1523/JNEUROSCI.1776-08.2008

Carrasco, M. (2011). Visual attention: The past 25 years. *Vision Research*, 51(13), 1484–1525. http://doi.org/10.1016/j.visres.2011.04.012

Dacey, D. M., & Petersen, M. R. (1992). Dendritic field size and morphology of midget and parasol ganglion cells of the human retina. *Proceedings of the National Academy of Sciences of the United States of America*, 89(20), 9666–9670. http://doi.org/10.1073/pnas.89.20.9666

Driver, J., & Spence, C. (1998). Attention and the crossmodal construction of space. *Trends*

in Cognitive Sciences, 2(7), 254–262. http://doi.org/10.1016/S1364-6613(98)01188-7

Eimer, M. (1999). Can attention be directed to opposite locations in different modalities? An

ERP study. *Clinical Neurophysiology*, 110(7), 1252–1259. https://doi.org/10.1016/S1388-

2457(99)00052-8

Eimer, M., & Driver, J. (2001). Crossmodal links in endogenous and exogenous spatial

attention: evidence from event-related brain potential studies. *Neuroscience and*

Biobehavioral Reviews, 25(6), 497–511. http://doi.org/S0149-7634(01)00029-X [pii]

Eimer, M., & Schroger, E. (1998). ERP effects of intermodal attention and cross-modal links

in   spatial   attention.   *Psychophysiology*,   35(3),   313–27.

https://doi.org/10.1017/S004857729897086X

Feldman, H., & Friston, K. J. (2010). Attention, uncertainty, and free-energy. *Frontiers in*

Human Neuroscience, 4(December), 1–23. http://doi.org/10.3389/fnhum.2010.00215

Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses using

G*Power 3.1: tests for correlation and regression analyses. *Behavior Research Methods*,

41(4), 1149–60. http://doi.org/10.3758/BRM.41.4.1149

Friston, K. (2005). A theory of cortical responses. *Philosophical Transactions of the Royal*

Society   of   London.   Series   B,   Biological   Sciences,   360(1456),   815–836.

https://doi.org/10.1098/rstb.2005.1622

Greene, A. J., Easton, R. D., & LaShell, L. S. R. (2001). Visual-auditory events: Cross-modal

perceptual priming and recognition memory. *Consciousness and Cognition*, 10(3), 425–435.

https://doi.org/10.1006/ccog.2001.0502

Jabar, S. B., & Anderson, B. (2015). Probability shapes perceptual precision: A study in

orientation estimation. *Journal of Experimental Psychology: Human Perception and*

Performance, 41(6), 1666–1679. http://doi.org/10.1037/xhp0000121

Jones, S. A., Beierholm, U., Meijer, D., & Noppeney, U. (2019). Older adults sacrifice

response speed to preserve multisensory integration performance. *Neurobiology of Aging*, 84,

148–157. https://doi.org/10.1016/j.neurobiolaging.2019.08.017

Kincade, J. M., Abrams, R. A., Astafiev, S. V., Shulman, G. L., & Corbetta, M. (2005). An

event-related functional magnetic resonance imaging study of voluntary and stimulus-driven

orienting of attention. *The Journal of Neuroscience*, 25(18), 4593–4604.

https://doi.org/10.1523/JNEUROSCI.0236-05.2005

Knudsen, E., & Brainard, M. S. (1995). Creating a unified representation of visual and

auditory space in the brain. *Annual Review of Neuroscience*, 18(1), 19–43.

http://doi.org/10.1146/annurev.neuro.18.1.19

Lauter, J. L., Herscovitch, P., Formby, C., & Raichle, M. E. (1985). Tonotopic organization

in human auditory cortex revealed by positron emission tomography. *Hearing Res*, 20, 199–

205.

Macaluso, E., Frith, C. D., & Driver, J. (2002). Supramodal effects of covert spatial orienting

triggered by visual or tactile events. *Journal of Cognitive Neuroscience*, 14(3), 389–401.

https://doi.org/10.1162/089892902317361912

Macaluso, E. (2010). Orienting of spatial attention and the interplay between the senses.

Cortex, 46(3), 282–297. http://doi.org/10.1016/j.cortex.2009.05.010

Maier, J. X., & Groh, J. M. (2009). Multisensory guidance of orienting behavior. *Hearing*

Research, 258(1–2), 106–112. https://doi.org/10.1016/j.heares.2009.05.008

Mars, R. B., Bestmann, S., Rothwell, J. C., & Haggard, P. (2007). Effects of motor

preparation and spatial attention on corticospinal excitability in a delayed-response paradigm.

Experimental Brain Research, 182(1), 125–129. http://doi.org/10.1007/s00221-007-1055-4

Meijer, D., Veselič, S., Calafiore, C., & Noppeney, U. (2019). Integration of audiovisual spatial signals is not consistent with maximum likelihood estimation. *Cortex*. https://doi.org/10.1016/j.cortex.2019.03.026

Middlebrooks, J. C., & Green, D. M. (1991). Sound localization by human listeners. *Annu. Rev. Psychol.*, (42), 135–159.

Mengotti, P., Boers, F., Dombert, P. L., Fink, G. R., & Vossel, S. (2018). Integrating modality-specific expectancies for the deployment of spatial attention. *Scientific Reports*, 8(1), 1210. https://doi.org/10.1038/s41598-018-19593-7

Miller, J. O., & Pachella, R. G. (1973). Locus of the stimulus probability effect. *Journal of Experimental Psychology*, 101(2), 227–231. http://doi.org/10.1037/h0035214

Molholm, S., Martinez, A., Shpaner, M., & Foxe, J. J. (2007). Object-based attention is multisensory: Co-activation of an object's representations in ignored sensory modalities. *European Journal of Neuroscience*, 26(2), 499–509. https://doi.org/10.1111/j.1460-9568.2007.05668.x

Mondor, T. A., & Amirault, K. J. (1998). Effect of same- and different-modality spatial cues on auditory and visual target identification. *Journal of Experimental Psychology*, 24(3), 745–755.

Neisser, U. (1977). Cognition and reality. *The American Journal of Psychology*, 90(3), 541–543.

Neumann, O., Van der Heijden, A. H., & Allport, D. A. (1986). Visual selective attention: Introductory remarks. *Psychological Research*, 48(4), 185–188. http://doi.org/10.1007/bf00309082

Odegaard, B., Wozny, D. R., & Shams, L. (2015). Biases in visual, auditory, and audiovisual perception of space. *PLoS Computational Biology*, 11(12), 1–23. https://doi.org/10.1371/journal.pcbi.1004649

Posner, M. I. (1980). Orienting of attention. *Q J Exp Psychol*, 32(1), 3–25.

http://doi.org/10.1080/00335558008248231

Rohe, T., & Noppeney, U. (2015a). Cortical hierarchies perform Bayesian Causal Inference

in multisensory perception. *PLOS Biology*, 13(2), e1002073.

https://doi.org/10.1371/journal.pbio.1002073

Rohe, T., & Noppeney, U. (2015b). Sensory reliability shapes perceptual inference via two

mechanisms. *Journal of Vision*, 15, 1–38. https://doi.org/10.1167/15.5.22.doi

Rohe, T., & Noppeney, U. (2016). Distinct computational principles govern multisensory

integration in primary sensory and association cortices. *Current Biology*, 26(4), 509–514.

https://doi.org/10.1016/j.cub.2015.12.056

Rohe, T., & Noppeney, U. (2018). Reliability-weighted integration of audiovisual signals can

be modulated by top-down control. *Eneuro*, 5, 1–20. http://doi.org/10.1523/ENEURO.0315-

17.2018

Rohenkohl, G., Gould, I. C., Pessoa, J., & Nobre, A. C. (2014). Combining spatial and

temporal expectations to improve visual perception. *Journal of Vision*, 14(4), 1–13.

http://doi.org/10.1167/14.4.8

Santangelo, V., Olivetti Belardinelli, M., Spence, C., & Macaluso, E. (2009). Interaction

between voluntary and stimulus-driven spatial attention mechanism across sensory

modalities. *Journal of Cognitive Neuroscience*, 21(12), 2384–2397.

https://doi.org/10.1162/jocn.2008.21178

Sereno, M. I., Dale, a M., Reppas, J. B., Kwong, K. K., Belliveau, J. W., Brady, T. J., …

Tootell, R. B. H. (1995). Borders of multiple visual areas in humans revealed by functional

magnetic resonance imaging borders of multiple visual areas in humans revealed by

functional magnetic resonance imaging. *Science*, 268(May), 889–893.

http://doi.org/10.1126/science.7754376

Spence, C., & Driver, J. (1996). Audiovisual links in endogenous covert spatial attention. *Journal of Experimental Psychology: Human Perception and Performance*, 22(4), 1005–1030. http://doi.org/10.1037/0096-1523.22.4.1005

Spence, C., & Driver, J. (1997). Audiovisual links in exogenous covert spatial orienting. *Perception & Psychophysics*, 59(1), 1–22. http://doi.org/10.3758/BF03206843

Stephen, J. M., Aine, C. J., Christner, R. F., Ranken, D., Huang, M., & Best, E. (2002). Central versus peripheral visual field stimulation results in timing differences in dorsal stream sources as measured with MEG. *Vision Research*, 42(28), 3059–3074. http://doi.org/10.1016/S0042-6989(02)00415-7

Summerfield, C., & Egner, T. (2009). Expectation (and attention) in visual cognition. *Trends in Cognitive Sciences*, 13(9), 403–409. http://doi.org/10.1016/j.tics.2009.06.003

Talsma, D., Kok, A., Slagter, H. A., & Cipriani, G. (2008). Attentional orienting across the sensory modalities. *Brain and Cognition*, 66(1), 1–10. https://doi.org/10.1016/j.bandc.2007.04.005

van Ede, F., de Lange, F. P., & Maris, E. (2012). Attentional cues affect accuracy and reaction time via different cognitive and neural processes. *The Journal of Neuroscience*, 32(30), 10408–10412. https://doi.org/10.1523/JNEUROSCI.1337-12.2012

Wahn, B., & König, P. (2015). Audition and vision share spatial attentional resources, yet attentional load does not disrupt audiovisual integration. *Frontiers in Psychology*, 6(July), 1–12. http://doi.org/10.3389/fpsyg.2015.01084

Wahn, B., & König, P. (2017). Is attentional resource allocation across sensory modalities task-dependent? *Advances in Cognitive Psychology*, 13(1), 83–96.

Ward, L. M., McDonald, J. J., & Lin, D. (2000). On asymmetries in cross-modal spatial attention orienting. *Perception & Psychophysics*, 62(6), 1258–1264. http://doi.org/10.3758/BF03212127

Zuanazzi, A., & Noppeney, U. (2018). Additive and interactive effects of spatial attention

and expectation on perceptual decisions. *Scientific Reports*, 1–12.

http://doi.org/10.1038/s41598-018-24703-6

Zuanazzi, A., & Noppeney, U. (2019). Distinct neural mechanisms of spatial attention and

expectation guide perceptual inference in a multisensory world. *The Journal of Neuroscience*,

39(12), 2873–18. https://doi.org/10.1523/jneurosci.2873-18.2019.