1 *Article*

Rewetting of Three Drained Peatlands Drives Congruent Compositional Changes in Pro- and Eukaryotic Soil Microbiomes Through Environmental Filtering

Micha Weil[1], Haitao Wang[1], Mia Bengtsson[1], Daniel Köhn[2], Anke Günther[2], Gerald Jurasinski[2], John Couwenberg[3], Wakene Negassa[2], Dominik Zak [4, 5], Tim Urich[1*]

[1] Institute of Microbiology, University of Greifswald, Felix-Hausdorff-Str. 8, 17487 Greifswald, Germany

[2] Faculty of Agriculture and Environmental Sciences, University of Rostock, Justus-von-Liebig-Weg 6, 18059 Rostock, Germany

[3] Institute of Botany and Landscape Ecology, University of Greifswald, Soldmannstraße 15, 17487 Greifswald, Germany, partner of the Greifswald Mire Center

[4] Department of Chemical Analytics and Biogeochemistry, Leibniz-Institute of Freshwater Biology and Inland Fisheries, Müggelseedamm 301, 12587 Berlin, Germany

[5] current address: Department of Bioscience, Aarhus University, Vejlsøvej 25, 8600 Silkeborg, Denmark

*corresponding author, Institute of Microbiology, University of Greifswald, Felix-Hausdorff-Str. 8, 17487 Greifswald, Germany; Tel.: +49 (0)3834 420 5904; E-Mail: tim.urich@uni-greifswald.de

Abstract:

Drained peatlands are significant sources of the greenhouse gas (GHG) carbon dioxide. Rewetting is a proven strategy to protect carbon stocks; however, it can lead to increased emissions of the potent GHG methane. The response to rewetting of soil microbiomes as drivers of these processes is poorly understood, as are biotic and abiotic factors that control community composition.

We analyzed the pro- and eukaryotic microbiomes of three contrasting pairs of minerotrophic fens subject to decade-long drainage and subsequent rewetting. Also, abiotic soil properties including moisture, dissolved organic matter, methane fluxes and ecosystem respiration rates.

The composition of the microbiomes was fen-type-specific, but all rewetted sites showed higher abundance of anaerobic taxa compared to drained sites. Based on multi-variate statistics and network analyses we identified soil moisture as major driver of community composition. Furthermore, salinity drove the separation between coastal and freshwater fen communities. Methanogens were more than tenfold more abundant in rewetted than in drained sites, while their abundance was lowest in the coastal fen, likely due to competition with sulfate reducers. The microbiome compositions were reflected in methane fluxes from the sites. Our results shed light on the factors that structure fen microbiomes via environmental filtering.

Keywords: peatland management; soil microbiome; methanogens; sulfate reducers; methanotrophic bacteria; greenhouse gas; methane

1. Introduction

Peatlands cover 3% of the global land surface and contain about 30% of the global soil organic
carbon (SOC) pool in the form of peat [1]. During the agricultural industrialization in the 20th
century, many temperate mires in Europe but also other parts of the world were drained to be used
as grassland or arable farming. Globally, drained or generally disturbed peatlands cover 0.3% of the
global land surface but contribute approx. 5% of all anthropogenic greenhouse gas (GHG) emissions
including carbon dioxide (CO_2), methane (CH_4) and nitrous oxide (N_2O). The latter holds also for
Germany, where drained peatlands cover approx. 5 % of the surface [2,3]. In the German federal state
of Mecklenburg-West-Pomerania, peatlands cover 13% of the land surface, with their majority being
drained (approx. 90%) contributing to about 30% of the total GHG budget in the region [4,5].

Peat accumulates under water-logged conditions, when more plant biomass is formed by
growth than is mineralized by the prokaryotic and eukaryotic soil (micro-)biome. Water is a key
factor, limiting decomposition by cutting off the supply of oxygen and other alternative terminal
electron acceptors (TEA) to mineralizing microbiota [6]. Under anoxic conditions plant litter
accumulates and forms peat with a high content of soil organic matter that accumulates over
millennia. Changes in ecology and hydrology of peatlands can lead to substantial changes in GHG
fluxes by changing the peatland biogeochemistry [7]. If peatlands are drained, formerly anoxic soil
layers become aerated and carbon is released as CO_2 during aerobic mineralization by peat
microbiota [8].

In the past three decades, several 100,000 ha of European peatlands were rewetted for
environmental protection and to recover some of their ecological functions [9,10]. Rewetting is a
proven strategy to protect the large SOC stocks; however, it can also lead to increased emissions of
the potent GHG CH_4 and to the release of dissolved organic matter (DOM). While effects of peatland
rewetting on GHG and DOM fluxes are comparably well studied [11–15], the impact on the peat
microbiota, the primary driver of GHG production and emissions, is poorly understood, since it has
been not well studied in temperate fens (see an exception in [16]). The major players in soil organic
matter (SOM) decomposition in peat soils are microorganisms of the bacterial, archaeal and
eukaryotic (e.g. fungi) domains of life, participating in a cascade of SOM degradation steps [9,17],
eventually resulting in emissions of CH_4 and CO_2. Hydrolytic extracellular enzymes catalyze the
initial decomposition of polymeric SOM. Under anoxic conditions, this step has been considered as a
major bottleneck and the one determining the rate of downstream processes [8]. Further steps in the
degradation are carried out through anaerobic respiratory microorganisms (like denitrifiers, iron
reducers and sulfate reducers), fermentative and methanogenic microorganisms (Dean *et al.*, 2018),
while methanotrophic microorganisms constitute the biological filter for CH_4 emissions from peat
(e.g. [18]). Remarkably, many novel players in soil CH_4 cycling have been identified in the past few
years [19–22]. Thus, despite more than half a century of research on methanogens, their diversity,
physiology and interactions within the microbiome are still poorly understood.

It is crucial to obtain a mechanistic understanding of microbial responses and dynamics upon
rewetting to suggest robust managing practices for climate-optimal peatland protection and
utilization (e.g. paludiculture, [23]). Environmental filtering, i.e. the selection of certain taxa through
abiotic environmental conditions, is considered to be a major mechanism structuring communities
[24]. In this context, identifying the main environmental drivers of both total microbiome
composition, as well as the abundance of relevant functional groups involved in carbon cycling and
methanogenesis is essential.

In this study we explore the effect of rewetting on temperate fen prokaryotic and eukaryotic
microbial communities. For this purpose, three pairs of drained (dry; terms in parenthesis are used
in the following to refer to these ecosystems and their states) and rewetted (wet) fens with contrasting
water tables and vegetation were investigated: a brackish coastal fen (Coast), a percolation fen (Perco)
and an alder carr (Alder). We hypothesize that despite the obvious structural and geo genetic
differences, congruent effects of rewetting will be detectable in the microbiomes of the three peatland
types because of environmental filtering. Possible controlling factors of community composition, like
DOM quantity and quality, soil moisture, soil/water salinity, were also determined to allow for

analyzing the effect of rewetting on key players of SOM mineralization and CH_4-cycling microorganisms.

A suite of cultivation-independent microbiological and geochemical methods was applied, including DOM profiling as well as qPCR of the methanogenesis key enzyme *mcrA* and 16S and 18S rRNA gene amplicon-based microbiome analysis. Our results may contribute to better understanding of how microbiomes respond to rewetting and to the relevant controlling environmental factors. Eventually. This could pave the way for microbiome-based proxies for predicting CH_4 emissions from rewetted peatlands.

2. Materials and Methods

2.1 Sites and sampling

All sampling sites, alder carr, coastal fen and percolation fen, are located in north-eastern Germany (Figure 1). The alder carr is part of the Recknitz river valley, was drained at least from 1786 on and has been used as a managed forest. The rewetted sub-site ($Alder_{wet}$) (54°13'N 12°49'E) was unintentionally flooded in 1999 due to a blocked drain and has remained wet since then. In the drained sub-site ($Alder_{dry}$) (54°13'N 12°51'E) long-term deep drainage has caused subsidence of the soil surface of about 1 m.

The coastal fen, located on a peninsula, was naturally episodically flooded by brackish seawater coming from the Bay of Greifswald of the Baltic Sea, typically during winter storm surges. The entire area was diked in 1850 to allow for pasture and other agricultural use. In 1993, the dykes around the rewetted sub site ($Coast_{wet}$) (54°16'N 13°39'E) were removed and, subsequently, the site was episodically flooded again. The drained sub-site ($Coast_{dry}$) (54°16'N 13°39'E) in contrast remains behind the dyke and, thus, cut off from the seawater influence. Both, $Coast_{wet}$ and $Coast_{dry}$ are used as cattle pastures.

The percolation fens, representing the third type of peatlands, are located in the catchment areas of the rivers Trebel and Recknitz. Both sites were deeply drained for intensive land-use and used as deep-drained grassland in the 20[th] century. In 1998, the water table at the rewetted sub-site ($Perco_{wet}$) (54°10'N 12°74'E) in the Trebel valley was raised again to 5-10 cm above ground level as part of a large rewetting project (EU LIFE project) (Ministerium für Bau, Landesentwicklung und Umwelt, M-V 1998). In contrast, the water table of the still drained sub-site ($Perco_{dry}$) (54°13'N 12°63'E) in the Recknitz valley remains several decimeters below the surface and is mown 1-2 times per year.

At all sites, peat samples were taken in April 2017 as triplicate cores at 5-10 cm, 15-20 cm and 25-30 cm depths (Figure S4) using a gouge. Each sample was homogenized and stored on ice in 50-ml reaction tubes over night until further processing.

Figure 1: Overview of sampling sites. A: alder carr; P: percolation fen; C: coastal fen; D: drained; W: rewetted.

2.2 Soil physico-chemical properties

Soil moisture was measured gravimetrically by drying 2-3 g of soil over night at 90°C until mass constancy. Soil moisture is expressed as the percentage of lost water weight to wet soil weight. The potential soil pH was measured at room temperature with a digital pH meter (pH 540 GLP, WTW, Weilheim, Germany) in 0.01 M CaCl$_2$ solution with a 1: 2.5 soil to solution ratio. Concentrations of total C, N and S were measured using a CNS analyzer (Vario MICRO cube – Elementar Analysensysteme GmbH Langenselbold, Germany). DOM was measured in soil extracts derived from mixing 3 g of previously frozen soil with 30 ml 0.1 M NaCl in 50 ml reaction tubes with subsequent shaking (vortex, 180 rpm) for 30 min. Extracts were filtered through 0.45 µm (pore size) sodium acetate filters, which were prewashed with 50 µl deionized H$_2$O to remove soluble acetate. The concentrations and the composition of DOM based on size categories were determined using a size-exclusion chromatography (SEC) with organic carbon and organic nitrogen detection (LC-OCD-OND analyzer, DOC-Labor Huber, Karlsruhe, Germany) [25]. The DOM was classified into three major sub-categories: (i) 'biopolymers', i.e. non-humic high molecular weight substances (> 10 kDa) of hydrophilic character and no unsaturated structures like polysaccharides and proteins, (ii) aromatic 'humic or humic-like substances' including building blocks, and (iii) 'low molecular-weight substances' including low molecular weight acids and low molecular weight neutral substances. Fractions were assigned based on standards of the International Humic Substances Society. Detection limits for each fraction was 0.01 mg C L^{-1}. Before analysis, all samples were stored at 5°C for less than two weeks to avoid significant changes of the DOM composition [26].

Soluble reactive phosphorus was determined with the ammonium molybdate spectrometric method (DIN EN 1189 D11) using a Cary 1E Spectrophotometer (Varian). N-NH$_4^+$ and N-NO$_3^-$ were determined calorimetrically using the photometry CFA method (Skalar SAN, Skalar Analytical B.V., The Netherlands) according to the guidelines in EN ISO 11732 (DEV-E 23) and EN ISO 13395 (DEV, D 28), respectively.

2.3 DNA extraction and PCR/qPCR

DNA was extracted from 0.25 g soil using the DNeasy PowerSoil Kit (QIAGEN, Hilden, Germany) according to the manufacturer´s recommendation. The bead beating step was performed using a FastPrep®-24 5G instrument (MP Biomedicals, Santa Ana, USA), with intensity 5.0 for 45 s. DNA concentrations were measured with Qubit®2.0 dsDNA High Sensitivity and dsDNA Broad Range assays (Thermo Fisher, Waltham, USA). DNA size was analyzed using 1% agarose gels.

Quantitative polymerase chain reaction was used to measure the abundances of the *mcrA* gene, coding for methyl coenzyme M reductase subunit A. Quantification of *mcrA* gene copies in peat soil DNA extracts was performed on a qTOWER 2.2 (Analytic Jena, Jena, Germany). The primer pair of forward mlas-mod and reverse *mcrA*- rev [27] was used. For each reaction, 15 µl of PCR mixture contained 7.5 µl of innuMIX qPCR MasterMix SyGreen (Analytic Jena, Jena, Germany), 0.75 µl of each primer (10 pmol/µl), 5 µl of ddH2O and 1 µl of template DNA. Duplicate measurements of two concentrations (1 and 2 ng/µl) were performed for each DNA sample. Assay condition was 95°C for 5 min, 35 cycles of 95°C for 30 s, 55°C for 45 s and 72°C for 45°C, followed by melting curve analysis to confirm PCR-product specificity. *mcrA* gene copy numbers were calculated from a standard curve obtained by serially diluting a standard from 10^6 to 10^1 *mcrA* gene copies per µl. The standard was created with above-mentioned primers by amplifying *mcrA* genes from cow rumen fluid [19] and cloning them into the pGEM®-Teasy vector system (Promega, Mannheim, Germany). Amplicons for *mcrA* standard were generated with vector-specific primers sp6 and T7 and the resulting PCR product was cleaned with DNA purification kit (Biozym, Hessisch Oldendorf, Germany), quality-controlled by agarose gel electrophoresis and quantified with Qubit®2.0 dsDNA High Sensitivity assay. The qPCR assay had the following parameters: slope: 3.41-3.52, efficiency: 0.92-0.96, R^2: > 0.995.

2.4 Amplicon sequencing and bioinformatics

16S and 18S rRNA gene amplicons were prepared and sequenced using 300 bp paired-end read Illumina MiSeq V3 sequencing by LGC Genomics (Berlin, Germany). This comprised 16S and 18S rRNA gene PCR amplification, Illumina MiSeq library preparation and sequencing using the primer pair 515YF (GTG YCA GCM GCC GCG GTA A)/ B806R (GGA CTA CNV GGG TWT CTA AT) [28] for prokaryotes and 1183F (AATTTGACTCAACRCGGG)/ 1443R (GRGCATCACAGACCTG) [29] for eukaryotes. The sequences have been submitted to European Nucleotide Archive (ENA), Project number PRJEB35436, accession number ERP118476, project name "Changes in peatland microbiome through rewetting".

The data were processed using R version 3.5.1 [30]. 16S rRNA and 18S rRNA gene amplicon sequences were quality filtered, de-replicated and clustered into amplicon sequence variants (ASVs) using the *dada2* package [31]. The chimeric sequences were de-novo checked and removed with *dada2*. Then the representative sequence of each ASV was assigned to taxonomy against a modified version of the SILVA SSUref_NR_128 data base, containing an updated taxonomy for *Archaea* [32], using the programs BLASTn [33] and Megan 5 [34]. ASVs that were assigned to Chloroplast or mitochondria were filtered. Due to the low number of sequences (<500) in one sample (Coast$_{dry}$, core 2, 15-20 cm depth) for both 16S and 18S rRNA genes, this sample was discarded from further analysis. After filtering, 1,134,953 and 3,990,919 sequences, sequences were retained and were clustered into 7697 and 5548 ASVs for 16S and 18S rRNA genes, respectively. Plant sequences in 18S rRNA genes, which accounted for 288 ASVs, were filtered for downstream analysis to increase the resolution of other eukaryotic taxa.

Alpha diversity (Shannon entropy) of the microbial community was calculated using the package *phyloseq* [35]. ASV count table was then normalized using metagenomeSeq's cumulative sum scaling (CSS) [36]. NMDS was performed to analyze the community composition using the package *vegan* [37] with defaults. Pairwise Spearman's rank correlations were conducted among soil properties, DNA concentrations, qPCR data, NMDS axes and methanogen relative abundances using the package *Hmisc* to detect the significance of the impact of the factors on microbial communities [38]. The soil properties with significant impact were fitted into NMDS ordination using the envfit function in the *vegan* package.

The correlations were visualized by heatmap using *corrplot*. All *P*-values for multiple
comparisons were adjusted by the false discovery rate (FDR) method and the null hypothesis was
rejected when *P*-values were less than 0.05.
The microbial functional groups in focus, i.e. fermenters, methanogens, methanotrophs and
sulfate reducers, were predicted from 16S rRNA amplicons using FAPROTAX [39]. Also,
methanogens, methanotrophs and sulfate reducers were identified based on literature knowledge;
e.g. ANME2-D group within *Methanosarcina* known as methanotrophic, using the methanogenic
pathway in reverse, was excluded from methanogens. All plots were created using package *ggplot2*
[40].

2.4.1 Co-occurrence network
A co-occurrence network was constructed to explore the potential interactions between species. Plant
ASVs were included in network analysis because plant residues are a major part of peat and they
might be important in interactions between microbes. To eliminate the influence of rare taxa, ASVs
with relative abundances lower than 0.05% were discarded. The pairwise Spearman's rank
correlations were conducted with the package *Hmisc* and all *P*-values were adjusted by the FDR. The
cut-offs of adjusted *P*-value and correlation coefficient were 0.01 and 0.7, respectively. The network
was visualized using package *igraph* [41]. A network showing the connections between eukaryotes
and prokaryotes was sub-selected from the original network by keeping the edges connecting these
two groups. ASVs that were statistically more associated to a certain fen type were defined as
indicator taxa. The indicators of each group or group combination of a factor (site or water condition)
were identified using *indicspecies* package.

2.5 Gas flux measurements
CH_4 exchange and ecosystem respiration (CO_2) from soil and non-woody vegetation were measured
in July 2017 with opaque, height-adjustable closed chambers in non-steady-state through-flow mode
[42]. Chambers are constructed of flexible polyurethane walls and are connected to circular collars
(n=5 per site, 0.63 m diameter) during the measurements. Collars were installed in the soil down to
10 cm depth more than one month prior to measurements. During chamber placement, headspace
CO_2 and CH_4 concentrations were monitored with portable analyzers (GasScouter®, Picarro and
Ultraportable Gas Analyzer, Los Gatos Research) [43].
231
232

233 **3. Results**

234 **3.1 Study sites and soil properties**

235 Our investigations took place in three pairs of drained and rewetted temperate peatlands in
236 Northern Germany (see Table 1, Figure1 and materials and methods for details).

237 **Table 1: Sampling site characteristics**

Abbr.	Site	State (year of rewetting)	Water table	Peat thickness	Dominant plant species
Alder	Alder carr	dry	-100 cm	60 cm	Black Alder (*Alnus glutinosa)*, Ground Elder (*Aegopodium podagraria*) and Common Nettle (*Urtica dioica*)
		wet (1999)	+15 cm	> 1 m	Black Alder (*Alnus glutinosa, Greater Pond Sedge (Carex riparia)*
Coast	Coastal fen	dry	-70 cm	70 cm	Creeping Bentgrass (*Agrostis stolonifera*)
		wet (1993)	-5 cm	30 cm	Creeping Bentgrass (*Agrostis stolonifera*)
Perco	Percolation fen	dry	-20 cm	6 m	Creeping Buttercup (*Ranunculus repens*), Tufted Hairgrass (*Deschampsia cespitosa*)
		wet (1998)	+10 cm	6 m	Lesser Pond Sedge (*Carex acutiformis*)

238
In general, the gravimetric water content was higher in the rewetted than in the drained sites
(Table S1). However, comparing the water content mean values of each site (mean of n=9, ± standard
deviation (SD)), there was only a minor difference between Perco$_{dry}$ (74.2% ± 5.1) and Perco$_{wet}$
(75.4% ± 5.3), because of a high water-table near the soil surface in Perco$_{dry}$ at the time of sampling. In
all of the drained sites and in Coast$_{wet}$, the water content increased with depth. In Perco$_{wet}$ and Alder$_{wet}$
(78.7 ± 4.6) with water table 10 cm above ground, the water content increased slightly with depth or
did not change. The lower gravimetric water contents in Coast$_{dry}$ (45.9% ± 6.7), Coast$_{wet}$
(53.8% ± 10.03), and Alder$_{dry}$ (46.3% ± 7.0) corresponded with higher amounts of mineral soil
compounds (sand and clay).
The sampling sites were generally acidic to sub neutral; pH values varied between fen types but
increased with depth in all drained sites. Lower pH values were detected in Coast$_{dry}$ (3.9 - 4.0) and
Alder$_{dry}$ (4.2 - 4.7), while pH values were higher in Perco$_{dry}$ (5.0-5.2) and Coast$_{wet}$ (4.7 - 6.1). In Alder$_{wet}$
(5.1 - 5.1) and Perco$_{wet}$ (5.4 – 5.4), the pH values did not change with depth (Table S1).
Salinity in the freshwater fens was generally low (0.08 - 0.49‰). The influence of the nearby
brackish bay water with a salinity of 8‰ led to a higher salinity in Coast$_{dry}$ (1.32‰) and in Coast$_{wet}$
(5.2‰) (Table S1).
DOM concentrations in the soil extracts of Alder$_{wet}$ and Coast$_{wet}$ were higher than in the drained
sites. In Perco$_{wet}$ DOM values measured near the surface were higher than in Perco$_{dry}$, but they
showed a strong decrease with increasing depth. The distribution of separate DOM fractions, i.e.
biopolymers, humic or humic-like substances and low molecular organic matter (lCOM) followed
the overall pattern of total DOM (Table S1).
Total carbon (TC) and total nitrogen (TN) content of peat decreased with depth in Alder$_{dry}$,
Coast$_{dry}$ and Coast$_{wet}$. In contrast, TC increased with depth in Alder$_{wet}$, Perco$_{dry}$ and Perco$_{wet}$. In
Alder$_{wet}$, Perco$_{wet}$ and Perco$_{dry}$ TN did not change with depth. Total sulfur (TS) content increased with
depth in Alder$_{dry}$, Alder$_{wet}$ and Perco$_{wet}$ while it decreased with depth in Coast$_{dry}$ and Coast$_{wet}$. In
Perco$_{dry}$, no change was observed over depth (Table S1).
265
266

3.2 Microbiome analysis

3.2.1 Diversity of prokaryotes and eukaryotes

Alpha diversity of prokaryotes, calculated from 16S rRNA gene amplicon sequence variants (ASVs), was significantly higher in the two freshwater peatlands than in the coastal fen (Figure S1, p < 0.001). Alpha diversity was slightly higher in rewetted than in drained sites, but this was significant only in the coastal fen (p < 0.001). Alpha diversity showed a positive correlation with soil water content, TC, TN and DNA content (Figure S2).

Similarly, alpha diversity of eukaryotes in freshwater peatlands was higher than in the coastal fen (*p* = 0.11). In contrast to prokaryotes, eukaryote alpha diversity was significantly lower in Perco$_{wet}$ than in Perco$_{dry}$ (*p* = 0.015), while no significant difference was observed in Alder and Coast. Alpha diversity of eukaryotes decreased with depth in the rewetted sites. For both pro- and eukaryotes, alpha diversity was negatively correlated with salinity (Figure S2).

3.2.2 Microbiota composition

Community composition in 16S rRNA and 18S rRNA sequencing analysis of prokaryotes and eukaryotes showed remarkably similar patterns in the non-metric multidimensional scaling (NMDS) plot (Figure 2), with a significant clustering of microbiota from the same sampling sites (PERMANOVA *p*= 0.001). While the communities of freshwater fens Alder and Perco showed some overlap, the microbiota in Coast were clearly separated along NMDS axis 1, possibly driven by the differences in abiotic factors, such as salinity, total nitrogen (TN), total carbon (TC), C/N ratio and DOM compounds (Figure 2). Nevertheless, the pro- and eukaryotic microbiota of the dry and wet sites of each fen type were consistently separated along NMDS axis 2, indicating that the water table of the sites influenced the community composition (PERMANOVA *p*= 0.001, R^2 = 0.267). Depth showed less impact on community composition (PERMANOVA *p* = 0.161, R^2 = 0.046), although some clustering according to sample depth was observed.

Figure 2: **NMDS plots showing (a) prokaryotic and (b) eukaryotic community compositions**. Symbols show triplicates of cores and indicate the study sites; the color indicates the depth in cm (see legend). Environmental vectors with a significance of *p* < 0.05 are shown in black. TN: total nitrogen, hCOM: humic like carbon, TC: total carbon, PolyC: Carbon polymers, DOM: dissolved organic matter, lCOM: low molecular organic matter, TS: total Sulphur, C/N: carbon/nitrogen ratio.

Pro- and eukaryotic community compositions were compared between fen types, water conditions (dry/wet) and sampling depths (Figures 3a, 5, A1). We detected a higher relative abundance of Acidobacteria in drained sites, while Betaproteobacteria, Deltaproteobacteria (mainly comprised of Myxobacteria and sulfate reducers), Chloroflexi and Bathyarchaeota showed higher values in all rewetted sites and Perco$_{dry}$. SPAM and Latescibacteria were not detected in Coast, but they occurred in the freshwater fens, whereas Actinobacteria showed higher values in Coast than in Alder and Perco.

The eukaryotic community varied considerably between fen types and water condition (Figure
3b). Among the fungi, ASVs of Ascomycota (which contain saprophytes, yeasts and several plant
pathogens) showed high relative abundance in each site, except Perco_dry. ASV values of
Basidiomycota decreased with depth in Alder and Coast, and showed higher relative abundance in
these two sites than in Perco. ASV values of Glomerycota, which form arbuscular mycorrhiza with
plants, comprising mostly Mortirellales were high in Alder_dry and Perco_dry.
Among the protists, ASV values of Apicomplexa were high (up to 40%) in Perco_dry, but
accounted for less than 5% in other sites. Cercozoa, which contain micro-predators among other life
styles, had higher values in the freshwater sites Alder and Perco. Relative abundance of Lobosa and
Ciliophora, also potential micro-predators, was high in Alder_wet and in sites with high water content
(rewetted sites and Perco_dry) respectively.
Among the Metazoa, Nematoda showed highest relative abundance, followed by Arthropoda.
The latter accounted for 5-10% of total amplicons in Coast and Perco_dry and the 5-10 cm layer of
Alder_dry, while their values were lower in Alder_wet and Perco_wet.
320

322   **Figure 3: Composition of prokaryotic and eukaryotic microbial communities of drained and**
323   **rewetted fens**. a) Taxonomic composition of prokaryotic 16S rRNA genes displayed on phylum level,
324   with Proteobacteria shown on class level. b) Taxonomic composition of eukaryotic 18S rRNA genes,
325   on phylum level. Bars show triplicate of cores; phyla with < 1% abundance are displayed as "other".
326   The sample ID on x-axis displays fen type and depth in cm.

3.2.3 Co-occurrence network analysis

A co-occurrence network was constructed to explore the potential interactions between pro- and
eukaryotic species. The network shows two distinct clusters of ASVs (Figure 4). Color coding of the
nodes according to association with sites shows that ASVs within the same site were more connected
than between sites (Figure 4b). However, ASVs associated with rewetted sites show little connections
with those from the drained sites (Figure 4c), highlighting the significant effect of water conditions
on the co-occurrence patterns. Similarly, the indicator taxa from Coast were mostly connected with
each other, and indicator taxa from Perco and Alder were also distinctly connected in the network
(Figure 4b). However, there were quite some shared indicator taxa from both Perco and Alder,
indicating similarities between these two habitats compared with Coast, which we already detected
in community composition (Figures 2 and 3). The number of a nodes adjacent edges (degree) and the
number of shortest paths going through a node (closeness centrality) of the indicator taxa were higher
in the wet sites than in the dry sites (Figures 4c, S4), particularly in Coast, followed by Alder, and

Perco (Figures 4b, c, S4). The probability that the neighbors of a node are connected (transitivity) was
higher among Coast indicators compared with those from Alder (Figure S4).

Figure 4: Co-occurrence networks. a) Co-occurrence network showing connections between
prokaryotes and eukaryotes corresponding to the ASVs of 16S and 18S rRNA amplicon sequencing.
ASVs with relative abundances lower than 0.05% were discarded.   b) Co-occurrence network of
ASVs associated with (A) alder carr, (C) coastal fen and (P) percolation fen, or with both alder carr
and coastal fen (AC), both alder carr and percolation fen (AP) and both coastal fen and percolation
fen (CP). c) Co-occurrence of ASVs associated with drained (D) and rewetted (W) sites. ASVs were
associated with sites using the *indicspecies* package [44]. Solid lines indicate significant positive
Spearman's rank correlations (r > 0.7. *p* < 0.01), dashed lines indicate significant negative correlations
(r < -0.7. *p* < 0.01). Nodes are sized according to the number of connected neighbors (degree) and
color-coded according to site type or water condition (drained/rewetted). The relative abundance of
prokaryotes and eukaryotes associated with the different site types is shown on the right-hand side
of b) and c).

The prokaryotic indicator taxa in drained sites mainly comprised ASVs from Acidobacteria and
Alpha- and Gamma-Proteobacteria, whereas the main indicator taxa of rewetted sites were Beta- and
Delta-Proteobacteria. Beta- and Delta-Proteobacteria were also the main phyla in both hydrological
status (Figure 3). Plants (Tracheophyta) accounted for the largest proportion of the eukaryotic
indicator taxa for both water condition and site. Nematoda ASVs accounted for a larger proportion
in the drained than in the rewetted sites; Ascomycota showed the reverse trend.
In the network with only connections between eukaryotes and prokaryotes (Figure 4a),
eukaryotes showed higher degrees of connection compared with prokaryotes (Figure 4). Among the
eukaryotes, fungi (Ascomycota, Basidiomycota and Glomeromycota) possessed the most connections
with the prokaryotes.

3.2.4 Functional groups
365
• **Methanogens**
The abundance of methanogens in the samples, assessed by qPCR of the *mcrA* gene, ranged from
2.1×10^5 to 5.1×10^7 gene copies g^{-1} dry soil (Figure 5). The *mcrA* gene abundance was significantly
higher in the three rewetted sites compared with the drained sites (Kruskal-Wallis test, H = 16.5, *p* <
0.001). The difference was largest in the top soils, where *mcrA* abundances were much higher in
rewetted sites. In Alder$_{wet}$ and Coast$_{wet}$ methanogen abundance decreased significantly with depth,
while it was rather constant in Perco$_{wet}$. In contrast, methanogen abundance in Alder$_{dry}$ was rather
constant or increased with depth (Coast$_{dry}$ and Perco$_{dry}$). Positive correlations were observed between
mcrA abundance and water, DNA and DOM content (strongest correlation with biopolymers and
lCOM; Figure S2). Furthermore, TC and TN as well as pH value were positively correlated with *mcrA*
abundance (Figure S2).
377

378
379   **Figure 5: Abundance of methanogens**. *mcrA* gene copy numbers per gram dry soil based on
380   qPCR. X-axis shows mcrA copies per g dry soil, drained sites in white, rewetted sites in black bars,
381   error bars show standard deviation of triplicate soil cores.
382

The relative abundance of methanogens in the prokaryotic microbiota was low. Methanogen-
affiliated ASVs in the amplicon sequence datasets were on average below 0.25% of total 16S rRNA
amplicons (Figure A1). In most of the drained fens, methanogen ASVs were even absent, although
they were detectable by *mcrA* qPCR (Figure 5). Upper soil samples of Alder~wet~ and Coast~wet~ showed
highest abundances, while methanogens were more abundant in the lower samples of Perco~wet~.
Methanobacteriales and Methanosarcinales (including the families Methanosaetaceae and
Methanosarcinaceae) were the dominant orders in Alder~wet~ and Coast~wet~, while methanogens were
more diverse in Perco~wet~, where Methanomassiliicoccales and Methanomicrobiales were also present.
Despite apparently missing out on some methanogens in the amplicon data, the relative
abundance calculated from *mcrA* gene copies per g dry soil (qPCR) shows a clear positive correlation
with the relative abundance of methanogenic archaea in 16S rRNA amplicon (Spearman correlation,
R= 0.578; $p < 0.001$).

Figure 6: Relative abundance of functional groups in 16S rRNA gene amplicon datasets.
Boxplots shows relative abundance in depth and triplicate samples. X-axes represent fen type: alder
carr (A), coastal fen (C), percolation fen (P), hydrological state drained (D) and rewetted (W), depth
in cm (05, 15, 25). Functional groups were assigned from 16S rRNA gene amplicon data by
FAPROTAX analysis.

• **Methanotrophs**
16S rRNA genes of known methanotrophic prokaryotes as assigned via FAPROTAX analysis,
were found in all sites (Figure 6), but never exceeded 1% of all prokaryotic 16S rRNA genes on
average. Their relative abundance and taxon composition differed both between fen types and
dry/wet. In drained fens ASVs affiliated with *Ca.* Methylacidiphilum (phylum Verrucomicrobia)

were dominant across the soil profile (Figure A1). In contrast, ASVs of Methylococcaceae (class Gammaproteobacteria) were the most abundant methanotrophs in Coast_wet and Perco_wet. Putative anaerobic methanotrophs affiliated with *Ca.* Methanoperedens (ANME-2d, archaea) and with *Ca.* Methylomirabilis (NC10 phylum), were the dominant methanotroph guild at the Alder_wet site.

- **Sulfate reducers**

ASVs assigned as sulfate reducers via FAPROTAX analysis were generally more abundant than methanogens and methanotrophs (between 0.1% and 5% on average, Figure 6). Like methanogens, their relative abundance was higher in rewetted sites than in drained sites. Highest relative abundance was observed in the Coast_wet, which is exposed to recurrent influx of sea water. ASVs of Desulfobacterales were particularly abundant in this site (Figure A1). Desulfarculales and Desulfosporosinus were only detected in rewetted sites, while Desulfuromonadales occurred in all sites. Relative abundance of sulfate reducers was positively correlated to salinity, but not to total sulfur (Figure S2).

3.3 Greenhouse gas emissions

In all drained peatlands a net uptake of CH_4 was measured, however, an average CH_4 emission of 15.8 mg m^{-2} h^{-1} CH_4 was found for Perco_wet with high variability (Tab. 2). Ecosystem respiration was highest in coastal peatlands. In all sites the drained peat soil emitted higher amounts of CO_2.

Table 2: Fluxes of CH_4 and ecosystem respiration (measured as CO_2 emissions) on all study sites compared to methanogen abundance (mcrA_DW). Negative flux values represent uptake from the atmosphere. Table shows mean values (n=5 for CH_4 and ecosystem respiration fluxes, n=9 for mcrA_DW) and standard deviation. Significance was tested with Kruskal-Wallis test.

Site	CH$_4$ flux (mg m^{-2} h^{-1})		Ecosystem respiration (CO$_2$) flux (mg m^{-2} h^{-1})		mcrA_DW (copies g^{-1} soil dry weight)	
Alder_dry	-0.08	± 0.07	493	± 406	8.5E+05	± 2.9E+05
Alder_wet	0.88*	± 0.51	47†	± 70	2.1E+07	± 2.6E+07
Coast_dry	-0.11	± 0.14	2024	± 585	3.5E+05	± 2.5E+05
Coast_wet	0.00*	± 0.01	1420	± 311	1.8E+06	± 2.1E+06
Perco_dry	-0.12	± 0.05	783	± 152	4.3E+06	± 3.5E+06
Perco_wet	15.80*	± 23.79	550†	± 232	3.4E+07	± 1.5E+07

* significantly higher than in drained site ($p < 0.05$)
† significantly lower than in drained site ($p < 0.05$)

4. Discussion

To our knowledge, this is the first comprehensive study investigating the impact of long-term rewetting (>> 10 years) on freshwater and coastal fen microbiomes. The parallel study of three different fen types allowed for the identification of overarching factors next to fen type-specific factors. The NMDS plots of both prokaryotes and eukaryotes were almost superimposable. It is remarkable that rewetting caused highly congruent effects on beta-diversity of all three domains of life and it is conceivable that water table and salinity were the main environmental filters among the physicochemical controlling factors. Rewetting resulted in more anoxic conditions in the rewetted sites compared with the drained sites. Increased anoxia is supported by several lines of evidence, including (1) a higher moisture content and (2) higher proportions of presumably fermentative microorganisms and (3) higher proportion of taxa with different types of anaerobic respiration, such as methanogenesis and sulfate reduction.

Environmental filtering effect on pro- and eukaryotic community composition

The network analysis confirmed the conclusions we drew from the community composition: there are more taxa indicative for each of the different fen types than for water status. Consequently, site location has a more significant effect on the community assemblages. The higher degree and closeness centrality of indicators in rewetted sites indicates more connections in these sites. This might be due to that higher water flow rates in soils create greater homogeneity, and hence weak niche differentiation, contributing to stronger interactions between soil microbes. Further, it might reflect the close metabolic interactions in anaerobic microbiomes during plant biomass degradation [18,45]. Main taxa of indicator ASVs in rewetted sites were Chloroflexi and Delta- and Betaproteobacteria, which contained ASVs known from anoxic environments, also highly abundant in the wet communities (Figure 3 and Figure 4) [46]. In the drained sites, most indicator taxa were from Acidobacteria, the predominant phylum, which would correspond with the general acidic to sub-neutral site conditions. Drainage and subsequent oxidation processes in peat soils are associated with acid production and depending on the buffer capacity of soils pH may go down by one or two units compared with pre-drained conditions [47]. In contrast, rewetting with base-rich water will increase alkalinity and pH [48]. Especially the two dominant classes Acidobacteria (subdivision 1) and Solibacteres (subdivision 3) decreased in relative abundance upon rewetting (Supplementary Figure A1), which is in line with the obligatory aerobic lifestyle of most characterized member species [49]. Members of the aerobic Acidobacteria play important roles in carbohydrate degradation, but some subdivisions also contain facultative or obligate anaerobic species [49,50]. Other putative anaerobic phyla such as Ignavibacter and Bathyarchaeota occurred only in rewetted sites and deep layers of drained sites. Obviously, soil moisture acted as environmental filter leading to community compositional changes from oxygen-dependent taxa (drained) to anaerobic and fermentative taxa in the rewetted peat soils (Figure 6). A second likely environmental filter in our study is salinity. The higher connectivity and transitivity of indicators from Coast may result from environmental filtering through higher salinity in these sites, leading to a change in microbial community with increases of Actinobacteria and sulfate reducers and decrease of methanogens ([51] and see below). Remarkably, the largest proportion of eukaryotic indicator ASVs originated from Tracheophyta (plants), indicating that the interaction between plants and the soil microbiota is very important. Similar to the prokaryotes, the interaction was even stronger in rewetted sites than in drained sites. In Coast with the lowest moisture content, Tracheophyta ASVs showed lower values than in Perco and Alder. Among the animals, Nematoda indicator ASVs accounted for a larger proportion in drained sites, likely because they are dependent on oxic conditions.

In the network, we also observed that eukaryotes had higher degrees than prokaryotes. The prokaryotes mostly connected with fungi, like Ascomycota, Basidiomycota and Glomeromycota, suggesting that especially fungi may drive the composition of prokaryotic communities. We found fewer eukaryotic than prokaryotic ASVs, suggesting that one eukaryote was probably connected with multiple prokaryotes, as implied by higher degrees of connection for eukaryotes. Similarly, the richness of eukaryotes was far less than the richness of prokaryotes, which is common for soil microbiota, as shown before in e.g. [18]. Therefore, our results potentially suggest this as a universal rule of pro-eukaryote interactions.

The influence of rewetting on methanogens

The important group of methanogenic archaea was hardly detected in the analysis of microbial networks and played no role in the determination of indicator taxa. The reason was the low relative abundance of methanogens in the amplicon datasets of the microbiota, being less than three per mil in all samples. Although their relative abundance in the microbiota was low even under rewetted conditions, their abundance per gram soil was high, especially in the rewetted sites. Wet peatlands are known to contribute a large share of the global CH_4 emissions [1,52]. The higher *mcrA* abundance after rewetting (Figure 5) could lead to increased CH_4 emissions from peatlands, as is the case in rice paddy fields [16,53]. Interestingly, their abundance was high in the rewetted top soils, at least in Alder_wet and Perco_wet, and not in the subsoils. This is in contrast to general assumptions about their

vertical distribution in the soil profile (reviewed in [6]) and corroborates recent findings from other
ecosystems [54,55]. This observation makes peatland management practices such as top soil removal
attractive for mitigation of CH_4 emissions from peatlands [56,57]. The abundance of methanogens
was strongly correlated with soil water content (Figure S2), which drives anoxia in the soil. However,
many other factors like substrate availability of DOM, especially lCOM, C and N content of soil,
degradation status of the peat and sulfate and nitrate concentration all positively correlated with
methanogen abundance per gram dry soil as well (Figure S2). One significant factor was salinity,
resulting from brackish water intrusions from the Baltic Sea in Coast$_{wet}$, which led to large differences
between freshwater fens (Alder and Perco) and Coast. In Coast$_{wet}$ the abundance of methanogens per
gram soil was at least 10x lower than in the freshwater sites Alder$_{wet}$ and Perco$_{wet}$. Thus, this
environmental filtering through salinity likely leads to lower CH_4 emissions, also shown before in
several studies [46,58,59].

The strong positive correlation of *mcrA* gene abundance in qPCR with the relative abundance of
methanogenic archaea in 16S rRNA gene amplicon datasets confirms the validity of the measured
results. However, in most of the drained peats (Perco$_{dry}$, most Alder$_{dry}$ and Coast$_{dry}$), methanogenic
taxa were not present in the dataset, although *mcrA* genes were quantified by qPCR. This can be
explained by the rather shallow average sequencing depth of 20,000 reads per sample, giving a taxon
detection limit of about 0.05‰ in the microbiota, and thus taxa with lower abundances could not be
detected. Still, the distribution of methanogenic taxa led to the assumption that substrate availability
was different in the sites. While hydrogenotrophic Methanobacteriales were dominant in Coast$_{wet}$,
Coast$_{dry}$ and Alder$_{wet}$, Perco$_{wet}$ showed a high variability in methanogenic groups. Here the
Methanosarcinales families of Methanosaetaceae (obligate acetoclastic) and Methanosarcinaceae
(flexible substrate spectrum) occurred, as did the Methanomicrobiales and the
Methanomassiliicoccales (that use the H_2 dependent methylotrophic pathway).
Methanomassiliicoccales were detected in every site with Methanomassiliicoccales specific qPCR
primers AS1/AS2 [60] (data not shown). Their high abundance in Perco$_{wet}$ indicates that besides the
well reported hydrogenotrophic and acetotrophic pathways [15], methylotrophic methanogenesis
may contribute more to peatland CH_4 emissions than currently assumed [19,61].

Interactions between methanogens, methanotrophs and sulfate reducers
The second important player in peatland CH_4 emissions is the heterogenic group of methanotrophic
Bacteria and Archaea, acting as biological CH_4 filter through aerobic or anaerobic oxidation of the
produced CH_4 (reviewed in [52,62]). Do the methanogens influence abundance of methanotrophs?
The methanotrophic groups differed with site, and were generally lower abundant in the drained
sites where *Ca.* Methylacidiphilum aerobic CH_4 oxidizers were dominant. These organisms were
originally isolated from acidic and volcanic environments, but have recently also been found in pH
neutral environments [63,64]. They are characterized by growth on low CH_4 concentrations, and are
possibly atmospheric CH_4 oxidizers, which would make them independent of substrate supply by
methanogens. As found in other wetland studies (e.g. [65]) aerobic CH_4 oxidizers of
Methylococcaceae dominated in the wet sites. The anaerobic group *Ca.* Methylomirabilis [64], which
uses oxygen from nitrite to reduce CH_4 appeared in the sites with highest water content (Alder$_{wet}$,
Perco$_{dry}$ and Perco$_{wet}$). The anaerobic CH_4 oxidizing ANME-2d (*Ca.* Methanoperedens) which reduce
CH_4 by the reverse methanogenic pathway [20] were detected in the water-covered fen Alder$_{wet}$.
Although we did not find a clear correlation between methanogen and methanotroph abundance (see
[16] for similar findings), we did find a strong distinction of anaerobic and aerobic methanotrophs
depending on water content and therefore oxygen availability, between sites and depths.
What is the basis of the environmental filter leading to the low methanogen abundance in the
coastal fen? It is known that sulfate reducers can have an impact on methanogens, because
methanogens and sulfate reducers not only share the anaerobic niche, but also compete for the same
substrates, such as hydrogen [46,66,67] and acetate [68]. Due to their higher substrate affinity, sulfate
reducers can outcompete methanogens, leading to lower methanogen abundance in the presence of
sulfate reducers [59,68]. In fact, Desulfobacterales, known competitors for H_2 as a substrate for

methanogenesis, occurred in the coastal site Coast_{wet} with a large proportion, which was associated with a more than tenfold lower *mcrA* gene abundance than in Alder_{wet} and Perco_{wet}. The influence of salinity in tidal marshes is also known to decrease CH_4 emission compared to freshwater sites [59,69].

These inferences based on microbial analysis and soil parameters were confirmed by GHG flux measurements, conducted three months after microbial sampling. For instance, ecosystem respiration in the drained sites were significantly higher than in the wet sites, in line with the higher abundance of aerobic pro- and eukaryotes that presumably oxidize more soil organic carbon. As suggested by the higher abundance of methanogens (Table 2 and Figure 5), CH_4 emissions were higher in wet sites compared to the drained sites. Interestingly, all drained sites as well as Coast_{wet} were CH_4 sinks in the summer, presumably fostered by the activity of the methanotrophs. In contrast, the wet sites Alder_{wet} and Perco_{wet} with mcrA abundances $< 10^7$ g^{-1} dry weight at the time of sampling acted as CH_4 sources.

5. Conclusions

In conclusion, this study provides microbiome-based insights into the environmental filtering effects of rewetting of fen ecosystems and suggests that, in terms of minimizing CH_4 emissions, increased salinity and/or top soil removal might be promising options. Future analysis of seasonal microbiome dynamics and measurements of CH_4 fluxes will provide evidence, whether these predictions hold true.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1 Figure S1: Alpha diversity of fen microbiomes, Figure S2: Heatmap showing Spearman correlation between microbial community factors and soil parameters, Figure S3 Relative abundance of Acidobacteria in 16S rRNA gene amplicon datasets, Figure S4: Node features of co-occurrence network analysis, Table S1: Biotic and abiotic soil parameters

Author Contributions: Conceptualization, M.W. and T.U.; methodology, M.W., H.W., D.K., A.G., W.N., D.Z. and T.U.; data curation, M.W. and H.W.; formal analysis, M.W., H.W. and M.B.; funding acquisition, G.J., J.C., T.U.; investigation, M.W. and T.U.; resources, M.W., G.J., W.N., D.Z. and T.U.; visualization, M.W. and H.W.; writing—original draft, M.W., H.W., T.U.; writing—review & editing, M.W., H.W., M.B., A.G., G.J., J.C., W.N., D.Z. and T.U. All authors have read and agreed to the published version of the manuscript.

Funding: The European Social Fund (ESF) and the Ministry of Education, Science and Culture of Mecklenburg-Western Pomerania (Germany) funded this work within the scope of the project WETSCAPES (ESF/14-BM-A55-0034/16 and ESF/14-BM-A55-0030/16).

Acknowledgments: The authors would like to thank Florian Beyer for providing the maps and arrangement of figure S4.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Appendix A

Figure A1: Relative abundance of functional groups in 16S rRNA gene amplicon datasets. a) Methanogens, b) methanotrophic bacteria and archaea, c) sulfate reducers. Barplots show relative abundance. X-axes represent fen type: alder carr (A), coastal fen (C), percolation fen (P), hydrological state drained (D) and rewetted (W), depth in cm (05, 15, 25). Data are shown as mean values of triplicate soil cores.

References

1. Joosten, H.; Sirin, A.; Couwenberg, J.; Laine, J.; Smith, P. The role of peatlands in climate regulation. *Peatl. Restor. Ecosyst. Serv. Sci. Policy Pract.* **2016**, 63–76.

2. Freibauer, A.; Drösler, M.; Gensior, A.; Schulze, E.D. (2009) Das Potenzial von Wäldern und Mooren für den Klimaschutz in Deutschland und auf globaler Ebene. Natur Landsch 84(1):20-25. *Natur und Landschaft* **2009**.

3. Joosten, H.; Couwenberg, J. Are emission reductions from peatlands measurable, reportable and verifiable? *Wetlands* **2009**, 14.

4.　Ziebarth, M.; Lenschow, U.; Permien, T. Konzept zum Schutz und zur Nutzung der Moore -
Fortschreibung des Konzeptes zur Bestandssicherung und zur Entwicklung der Moore. **2009**, 109.

5.　Greifswald Mire Center Moorbodenschutz als naturbasierte Lösung im Klimaschutzprogramm 2030 -
schnelle Einstellung von Moor-Entwässerung für wirkungsvollen Klimaschutz nötig ! **2019**, 2017–2020.

6.　Dean, J.F.; Middelburg, J.J.; Röckmann, T.; Aerts, R.; Blauw, L.G.; Egger, M.; Jetten, M.S.M.; de Jong,
A.E.E.; Meisel, O.H.; Rasigraf, O.; et al. Methane Feedbacks to the Global Climate System in a Warmer
World. *Rev. Geophys.* **2018**, *56*, 207–250.

7.　Parish, F.; Sirin, A.; Charman, D.; Joosten, H.; Minayeva, T.; Silvius, M.; Stringer, L. *Assessment on
Peatlands, Biodiversity and Climate Change: Main Report*; 2008; ISBN 978-983-43751-0-2.

8.　Conrad, R. Soil Microorganisms as Controllers of Atmospheric Trace Gases (H 2 , CO , CH 4 , OCS , N
2 O , and NO). **1996**, *60*, 609–640.

9.　Andersen, R.; Chapman, S.J.; Artz, R.R.E. Microbial communities in natural and disturbed peatlands: A
review. *Soil Biol. Biochem.* **2013**, *57*, 979–994.

10.　Chimner, R.A.; Cooper, D.J.; Wurster, F.C.; Rochefort, L. An overview of peatland restoration in North
America: where are we after 25 years? *Restor. Ecol.* **2017**, *25*, 283–292.

11.　Couwenberg, J.; Thiele, A.; Tanneberger, F.; Augustin, J.; Bärisch, S.; Dubovik, D.; Liashchynskaya, N.;
Michaelis, D.; Minke, M.; Skuratovich, A.; et al. Assessing greenhouse gas emissions from peatlands
using vegetation as a proxy. *Hydrobiologia* **2011**, *674*, 67–89.

12.　Hahn, J.; Köhler, S.; Glatzel, S.; Jurasinski, G. Methane exchange in a coastal fen in the first year after
flooding - A systems shift. *PLoS One* **2015**, *10*.

13.　Estop-Aragonés, C.; Zajac, K.; Blodau, C. Effects of extreme experimental drought and rewetting on CO2
and CH4 exchange in mesocosms of 14 European peatlands with different nitrogen and sulfur
deposition. *Glob. Chang. Biol.* **2016**, *22*, 2285–2300.

14.　Wilson, D. Blain, D. Cowenberg, J. Evans, C.D. Murdiyarso, D. Page, S. E. Renou-Wilson, F. Rieley, J.O.
Sirin, A. Strack. M. Tuittila, E.-S. Greenhouse gas emission factors associated with rewetting of organic
soils. *Mires Peat* **2016**, *17*, 1–28.

15.　Lyu, Z.; Shao, N.; Akinyemi, T.; Whitman, W.B. Methanogenesis. *Curr. Biol.* 2018, *28*, R727–R732.

16.　Wen, X.; Unger, V.; Jurasinski, G.; Koebsch, F.; Horn, F.; Rehder, G.; Sachs, T.; Zak, D.; Lischeid, G.;
Knorr, K.H.; et al. Predominance of methanogens over methanotrophs in rewetted fens characterized by
high methane emissions. *Biogeosciences* **2018**, *15*, 6519–6536.

17.　Yarwood, S.A. The role of wetland microorganisms in plant-litter decomposition and soil organic matter
formation: a critical review. *FEMS Microbiol. Ecol.* **2018**, *94*, 1–17.

18.　Tveit, A.T.; Urich, T.; Frenzel, P.; Svenning, M.M. Metabolic and trophic interactions modulate methane

production by Arctic peat microbiota in response to warming. *Proc. Natl. Acad. Sci.* **2015**, *112*, E2507–
E2516.

19. Söllinger, A.; Schwab, C.; Weinmaier, T.; Loy, A.; Tveit, A.T.; Schleper, C.; Urich, T. Phylogenetic and
genomic analysis of Methanomassiliicoccales in wetlands and animal intestinal tracts reveals clade-
specific habitat. *FEMS Microbiol. Ecol.* **2016**, *92*, 1–12.

20. Welte, C.U.; Rasigraf, O.; Vaksmaa, A.; Versantvoort, W.; Arshad, A.; Op den Camp, H.J.M.; Jetten,
M.S.M.; Li¿½ke, C.; Reimann, J.; Lüke, C.; et al. Nitrate- and nitrite-dependent anaerobic oxidation of
methane. *Environ. Microbiol. Rep.* **2016**, *8*, 941–955.

21. Sorokin, D.Y.; Makarova, K.S.; Abbas, B.; Ferrer, M.; Golyshin, P.N.; Galinski, E.A.; Ciordia, S.; Mena,
M.C.; Merkel, A.Y.; Wolf, Y.I.; et al. Discovery of extremely halophilic, methyl-reducing euryarchaea
provides insights into the evolutionary origin of methanogenesis. *Nat. Microbiol.* **2017**, *2*, 17081.

22. Singleton, C.; McCalley, C.; Woodcroft, B.; Boyd, J.; Evans, P.; Hodgkins, S.; Chanton, J.; Frolking, S.;
Crill, P.; Saleska, S.; et al. Methanotrophy across a natural permafrost thaw environment. *ISME J.* **2018**,
1.

23. Bonn, A.; Allott, T.; Evans, M.; Joosten, H.; Stoneman, R. *Peatland restoration and ecosystem services: Science,*
policy and practice; 2016; ISBN 9781139177788.

24. Cadotte, M.W.; Tucker, C.M. Should Environmental Filtering be Abandoned? *Trends Ecol. Evol.* **2017**, *32*,
429–437.

25. Huber, S.A.; Balz, A.; Abert, M.; Pronk, W. Characterisation of aquatic humic and non-humic matter
with size-exclusion chromatography - organic carbon detection - organic nitrogen detection (LC-OCD-
OND). *Water Res.* **2011**, *45*, 879–885.

26. Heinz, M.; Zak, D. Storage effects on quantity and composition of dissolved organic carbon and nitrogen
of lake water, leaf leachate and peat soil water. *Water Res.* **2018**, *130*, 98–104.

27. Steinberg, L.M.; Regan, J.M. mcrA-targeted real-time quantitative PCR method to examine methanogen
communities. *Appl. Environ. Microbiol.* **2009**, *75*, 4435–4442.

28. Walters, W.; Hyde, E.R.; Berg-lyons, D.; Ackermann, G.; Humphrey, G.; Parada, A.; Gilbert, J. a; Jansson,
J.K. Transcribed Spacer Marker Gene Primers for Microbial Community Surveys. *mSystems* **2015**, *1*,
e0009-15.

29. Ray, J.L.; Althammer, J.; Skaar, K.S.; Simonelli, P.; Larsen, A.; Stoecker, D.; Sazhin, A.; Ijaz, U.Z.; Quince,
C.; Nejstgaard, J.C.; et al. Metabarcoding and metabolome analyses of copepod grazing reveal feeding
preference and linkage to metabolite classes in dynamic microbial plankton communities. *Mol. Ecol.*
2016, *25*, 5585–5602.

30. The R Core Team *R: A language and environment for statistical computing.*; 2018; ISBN 3_900051_00_3.

31. Callahan, B.J.; McMurdie, P.J.; Rosen, M.J.; Han, A.W.; Johnson, A.J.A.; Holmes, S.P. DADA2: High-

resolution sample inference from Illumina amplicon data. *Nat. Methods* **2016**, *13*, 581–583.

32. Lanzén, A.; Jørgensen, S.L.; Huson, D.H.; Gorfer, M.; Grindhaug, S.H.; Jonassen, I.; Øvreås, L.; Urich, T.
CREST - Classification Resources for Environmental Sequence Tags. *PLoS One* **2012**, *7*.

33. Altschul, S.F.; Gish, W.; Miller, W.; Myers, E.W.; Lipman, D.J. Basic local alignment search tool. *J. Mol.*
Biol. **1990**.

34. Huson, D.H.; Beier, S.; Flade, I.; Górska, A.; El-Hadidi, M.; Mitra, S.; Ruscheweyh, H.J.; Tappu, R.
MEGAN Community Edition - Interactive Exploration and Analysis of Large-Scale Microbiome
Sequencing Data. *PLoS Comput. Biol.* **2016**, *12*, 1–12.

35. McMurdie, P.J.; Holmes, S. Phyloseq: An R Package for Reproducible Interactive Analysis and Graphics
of Microbiome Census Data. *PLoS One* **2013**, *8*.

36. Paulson, J.N.; Colin Stine, O.; Bravo, H.C.; Pop, M. Differential abundance analysis for microbial marker-
gene surveys. *Nat. Methods* **2013**.

37. Oksanen, J.; Blanchet, F.G.; Kindt, R.; Legendre, P.; Minchin, P.R.; O'Hara, R.B.; Simpson, G.L.; Solymos,
P.; Stevens, M.H.H.; Wagner, H. vegan: Community Ecology Package. *http://cran.r-*
project.org/package=vegan 2016.

38. Harrell, F.E.; Dunpont, C. Package "Hmisc" Version 4.2-0. **2019**, 428.

39. Louca, S.; Wegener Parfrey, L.; Doebeli, M. Decoupling function and taxonomy in the global ocean
microbiome. *Science* **2016**, *353*, 1272–1277.

40. Valero-Mora, P.M. *ggplot2: Elegant Graphics for Data Analysis* ; Springer-Verlag: New York, 2010; Vol. 35;
ISBN ISBN 978-3-319-24277-4.

41. Package, R. Package ' igraph ' R topics documented : **2019**.

42. Livingston, G.P.; Hutchinson, G.L. Enclosure-based measurement of trace gas exchange: applications
and sources of error. In *Biogenic Trace Gases: Measuring Emissions from Soil and Water*; Matson, P.A.,
Harris, R.C., Eds.; 1995; pp. 14–53.

43. Günther, A.; Schenzle, E.; Jurasinski, G. *Flexible chambers for greenhouse gas measurements in tall vegetation.*;
2014;

44. De Cáceres, M.; Legendre, P. Associations between species and groups of sites: Indices and statistical
inference. *Ecology* **2009**, *90*, 3566–3574.

45. Schink, B. Energetics of syntrophic cooperation in methanogenic degradation. *Microbiol. Mol. Biol. Rev.*
1997, *61*, 262–80.

46. He, S.; Malfatti, S. a; Mcfarland, J.W.; Anderson, F.E.; Pati, A.; Huntemann, M.; Tremblay, J.; Glavina, T.;
Waldrop, M.P.; Windham-myers, L.; et al. Patterns in Wetland Microbial Community Composition and

Functional Gene Repertoire Associated with Methane Emissions. *Mbio.Asm.Org* **2015**, *6*, 1–15.

47.   Lamers, L.P.M.; van Roozendaal, S.M.E.; Roelofs, J.G.M. A_1005083526455(1).pdf. *Water. Air. Soil Pollut.*
701   **1998**, *105*, 95–106.

702   48.   Cusell, C.; Lamers, L.P.M.; van Wirdum, G.; Kooijman, A. Impacts of water level fluctuation on
703   mesotrophic rich fens: Acidification vs. eutrophication. *J. Appl. Ecol.* **2013**, *50*, 998–1009.

704   49.   Eichorst, S.A.; Trojan, D.; Roux, S.; Herbold, C.; Rattei, T.; Woebken, D. Genomic insights into the
705   Acidobacteria reveal strategies for their success in terrestrial environments. *Environ. Microbiol.* **2018**, *20*,
706   1041–1063.

707   50.   Hausmann, B.; Pelikan, C.; Herbold, C.W.; Köstlbacher, S.; Albertsen, M.; Eichorst, S.A.; Glavina Del Rio,
708   T.; Huemer, M.; Nielsen, P.H.; Rattei, T.; et al. Peatland Acidobacteria with a dissimilatory sulfur
709   metabolism. *ISME J.* **2018**, *12*, 1729–1742.

710   51.   Dang, C.; Morrissey, E.M.; Virginia, W.; Neubauer, S.C.; Franklin, R.B. Novel microbial community
711   composition and carbon biogeochemistry emerge over time following saltwater intrusion in wetlands.
712   **2018**.

713   52.   Nazaries, L.; Murrell, J.C.; Millard, P.; Baggs, L.; Singh, B.K. Methane, microbes and models:
714   Fundamental understanding of the soil methane cycle for future predictions. *Environ. Microbiol.* 2013, *15*,
715   2395–2417.

716   53.   Reim, A.; Hernández, M.; Klose, M.; Chidthaisong, A.; Yuttitham, M.; Conrad, R. Response of
717   methanogenic microbial communities to desiccation stress in flooded and rain-fed paddy soil from
718   Thailand. *Front. Microbiol.* **2017**, *8*, 1–17.

719   54.   Hahn-Schöfl, M.; Zak, D.; Minke, M.; Gelbrecht, J.; Augustin, J.; Freibauer, A. Organic sediment formed
720   during inundation of a degraded fen grassland emits large fluxes of CH4 and CO2. *Biogeosciences* **2011**,
721   *8*, 1539–1550.

722   55.   Angle, J.C.; Morin, T.H.; Solden, L.M.; Narrowe, A.B.; Smith, G.J.; Borton, M.A.; Rey-Sanchez, C.; Daly,
723   R.A.; Mirfenderesgi, G.; Hoyt, D.W.; et al. Methanogenesis in oxygenated soils is a substantial fraction
724   of wetland methane emissions. *Nat. Commun.* **2017**, *8*, 1567.

725   56.   Harpenslager, S.F.; van den Elzen, E.; Kox, M.A.R.; Smolders, A.J.P.; Ettwig, K.F.; Lamers, L.P.M.
726   Rewetting former agricultural peatlands: Topsoil removal as a prerequisite to avoid strong nutrient and
727   greenhouse gas emissions. *Ecol. Eng.* **2015**, *84*, 159–168.

728   57.   Zak, D.; Goldhammer, T.; Cabezas, A.; Gelbrecht, J.; Gurke, R.; Wagner, C.; Reuter, H.; Augustin, J.;
729   Klimkowska, A.; McInnes, R. Top soil removal reduces water pollution from phosphorus and dissolved
730   organic matter and lowers methane emissions from rewetted peatlands. *J. Appl. Ecol.* **2018**, *55*, 311–320.

731   58.   Neubauer, S.C.; Franklin, R.B.; Berrier, D.J. Saltwater intrusion into tidal freshwater marshes alters the
732   biogeochemical processing of organic carbon. *Biogeosciences* **2013**, *10*, 8171–8183.

59.   van Dijk, G.; Nijp, J.J.; Metselaar, K.; Lamers, L.P.M.; Smolders, A.J.P. Salinity-induced increase of the
hydraulic conductivity in the hyporheic zone of coastal wetlands. *Hydrol. Process.* **2017**, *31*, 880–890.

60.   Mihajlovski, A.; Doré, J.; Levenez, F.; Alric, M.; Brugère, J.F. Molecular evaluation of the human gut
methanogenic archaeal microbiota reveals an age-associated increase of the diversity. *Environ. Microbiol.*
Rep. **2010**, *2*, 272–280.

61.   Zalman, C.A.; Meade, N.; Chanton, J.; Kostka, J.E.; Bridgham, S.D.; Keller, J.K. Methylotrophic
methanogenesis in Sphagnum-dominated peatland soils. *Soil Biol. Biochem.* **2017**, *118*, 156–160.

62.   Knief, C. Diversity and habitat preferences of cultivated and uncultivated aerobic methanotrophic
bacteria evaluated based on pmoA as molecular marker. *Front. Microbiol.* **2015**, *6*.

63.   van Teeseling, M.C.F.; Pol, A.; Harhangi, H.R.; van der Zwart, S.; Jetten, M.S.M.; Op den Camp, H.J.M.;
van Niftrik, L. Expanding the verrucomicrobial methanotrophic world: Description of three novel
species of Methylacidimicrobium gen. nov. *Appl. Environ. Microbiol.* **2014**, *80*, 6782–6791.

64.   Vaksmaa, A.; Lüke, C.; van Alen, T.; Valè, G.; Lupotto, E.; Jetten, M.S.M.; Ettwig, K.F. Distribution and
activity of the anaerobic methanotrophic community in a nitrogen-fertilized Italian paddy soil. *FEMS*
Microbiol. Ecol. **2016**, *92*, 1–11.

65.   Tveit, A.T.; Urich, T.; Svenning, M.M. Metatranscriptomic analysis of arctic peat soil microbiota. *Appl.*
Environ. Microbiol. **2014**, *80*, 5761–5772.

66.   Kristjansson, J.K.; Schönheit, P. Why do sulfate-reducing bacteria outcompete methanogenic bacteria for
substrates? *Oecologia* **1983**, *60*, 264–266.

67.   Thauer, R.K.; Kaster, A.-K.K.; Seedorf, H.; Buckel, W.; Hedderich, R. Methanogenic archaea: Ecologically
relevant differences in energy conservation. *Nat. Rev. Microbiol.* **2008**, *6*, 579–591.

68.   Kristjansson, J.K.; Schönheit, P.; Thauer, R.K. Different Ks values for hydrogen of methanogenic bacteria
and sulfate reducing bacteria: An explanation for the apparent inhibition of methanogenesis by sulfate.
Arch. Microbiol. **1982**.

69.   Poffenbarger, H.J.; Needelman, B.A.; Megonigal, J.P. Salinity influence on methane emissions from tidal
marshes. *Wetlands* **2011**, *31*, 831–842.

a)

Prokaryote
- Deltaproteobacteria
- Actinobacteria
- Epsilonbacteraeota
- Chloroflexi
- Betaproteobacteria
- Gammaproteobacteria
- Ignavibacteriae
- Acidobacteria
- Verrucomicrobia
- Gemmatimonadetes
- Planctomycetes
- Alphaproteobacteria
- Nitrospirae
- Chlorobi
- NC10
- SPAM
- Ca. Latescibacteria (WS3)
- Ca. Parcubacteria
- Thaumarchaeota
- Bacteroidetes
- Ca. Bathyarchaeota
- Ca. Gracilibacteria
- Spirochaetes

Eukaryote
- Ascomycota
- Basidiomycota
- Glomeromycota
- Ciliophora
- Tracheophyta
- Chytridiomycetes
- Mortierellales
- Apicomplexa
- Nematoda
- Unclassified
- Labyrinthulomycetes
- Arthropoda
- Lobosa
- Cercozoa
- Peronosporomycetes
- Dinoflagellata
- Cryptomycota
- Annelida
- Mollusca

b)

● Prokaryote
▲ Eukaryote

- None-indicator
- A
- C
- P
- AC
- AP
- CP

c)

- Rewetted
- Drained
- None-indicator

● Prokaryote
▲ Eukaryote