Rewetting of three drained peatlands drives congruent compositional changes in

pro- and eukaryotic microbiomes through environmental filtering

Micha Weil[1], Haitao Wang[1], Mia Bengtsson[1], Anke Günther[2], Gerald Jurasinski[2], John Couwenberg[3],

Wakene Negassa[2], Dominik Zak [4, 5], Tim Urich[1*]

[1] Institute of Microbiology, University of Greifswald, Felix-Hausdorff-Str. 8, 17487 Greifswald, Germany

[2] Faculty of Agriculture and Environmental Sciences, University of Rostock, Justus-von-Liebig-Weg 6, 18059

Rostock, Germany

[3] Institute of Botany and Landscape Ecology, University of Greifswald, Soldmannstraße 15, 17487 Greifswald,

Germany, partner of the Greifswald Mire Center

[4] Department of Chemical Analytics and Biogeochemistry, Leibniz-Institute of Freshwater Biology and Inland

Fisheries, Müggelseedamm 301, 12587 Berlin, Germany

[5] current address: Department of Bioscience, Aarhus University, Vejlsøvej 25, 8600 Silkeborg, Denmark

*corresponding author, Institute of Microbiology, University of Greifswald, Felix-Hausdorff-Str. 8, 17487

Greifswald, Germany; Tel.: +49 (0)3834 420 5904; E-Mail: tim.urich@uni-greifswald.de

Running Title: Changes in peatland microbiome through rewetting

Summary

Drained peatlands are significant sources of the greenhouse gas (GHG) carbon dioxide. Rewetting is a proven strategy to protect the large carbon stocks; however, it can also lead to increased emissions of the potent GHG methane. The response to rewetting of soil microbiomes as drivers of these processes is poorly understood, as are the biotic and abiotic factors that control community composition.

We analyzed the soil microbiomes of three contrasting pairs of minerotrophic fens subject to decade-long drainage and subsequent rewetting. Analyses comprised qPCR of methanogens, amplicon sequencing of prokaryotic and eukaryotic rRNA genes, and abiotic soil properties including moisture and dissolved organic matter.

The composition of pro- and eukaryotic communities was fen-type-specific, but all rewetted site showed higher abundance of anaerobic taxa. Based on multi-variate statistics and network analyses we identified soil moisture as major driver of community composition. Furthermore, salinity drove the separation between coastal and freshwater fen communities. Methanogens were more than tenfold more abundant in rewetted than in drained sites while their abundance was lowest in the coastal fen, likely due to competition with sulfate reducers. These results shed light on the factors that structure fen microbiomes via environmental filtering.

1. Introduction

Peatlands cover 3% of the global land surface and contain about 30% of the global soil organic carbon (SOC) pool in the form of peat (Joosten *et al.*, 2016). During the agricultural industrialization in the 20th century, many temperate mires in Europe but also other parts of the world were drained to be used as grassland or arable farming. Globally, drained or generally disturbed peatlands cover 0.3% of the global land surface but contribute approx. 5% of all anthropogenic greenhouse gas (GHG) emissions including carbon dioxide (CO_2), methane (CH_4) and nitrous oxide (N_2O). The latter holds also for Germany, where drained peatlands cover approx. 5 % of the surface (Freibauer *et al.*, 2009; Joosten and Couwenberg, 2009). In the German federal state of Mecklenburg-West-Pomerania, peatlands cover 13% of the land surface, their large majority being drained (approx. 90%) and emitting about 30% of the total GHG budget in the region (Ziebarth *et al.*, 2009; Greifswald Mire Center, 2019).

Peat accumulates under water-logged conditions, when more plant biomass is formed by growth than is mineralized by the prokaryotic and eukaryotic soil (micro-)biome. Water is a key factor, limiting decomposition by cutting off the supply of oxygen and other alternative terminal electron acceptors (TEA) to mineralizing microbiota (Dean *et al.*, 2018). Under anoxic conditions plant litter accumulates and forms peat with a high content of soil organic matter that accumulates over millennia. Changes in ecology and hydrology of peatlands can lead to substantial changes in GHG fluxes by changing the peatland biogeochemistry (Parish *et al.*, 2008). If peatlands are drained, formerly anoxic soil layers become aerated and carbon is released as CO_2 during aerobic mineralization by peat microbiota (Conrad, 1996).

In the past three decades, several 100 000 ha of European peatlands were rewetted for environmental protection and to recover some of their ecological functions (Andersen *et al.*, 2013; Chimner *et al.*, 2017). Rewetting is a proven strategy to protect the large SOC stocks; however, it can also lead to increased emissions of the potent GHG CH_4 and to the release of dissolved organic matter (DOM). While effects of peatland rewetting on GHG and DOM fluxes are comparably well studied (Couwenberg *et al.*, 2011; Hahn *et al.*, 2015; Estop-Aragonés *et al.*, 2016; Wilson, D. Blain, D. Cowenberg, J. Evans,

C.D. Murdiyarso, D. Page, S. E. Renou-Wilson, F. Rieley, J.O. Sirin, A. Strack. M. Tuittila, 2016; Lyu *et al.*,

2018), the impact on the peat microbiota, who is the primary driver of GHG production and thus

emissions is poorly understood, because it has been not well studied in temperate fens (see an

exception in (Wen *et al.*, 2018). The major players in soil organic matter (SOM) decomposition in peat

soils are microorganisms of the bacterial, archaeal and eukaryotic (e.g. fungi) domains of life,

participating in a cascade of SOM degradation steps (Andersen *et al.*, 2013; Yarwood, 2018), eventually

resulting in emissions of CH_4 and CO_2. Hydrolytic extracellular enzymes catalyze the initial

decomposition of polymeric SOM. Under anoxic conditions, this step has been considered as a major

bottleneck and the one determining the rate of downstream processes (Conrad, 1996). Further steps

in the degradation are carried out through anaerobic respiratory microorganisms (like denitrifiers, iron

reducers and sulfate reducers), fermentative and methanogenic microorganisms (Dean *et al.*, 2018),

while methanotrophic microorganisms constitute the biological filter for CH_4 emissions from peat (e.g.

Tveit et al., 2013). Remarkably, many novel players in soil CH_4 cycling have been identified in the past

few years (Söllinger *et al.*, 2016; Welte *et al.*, 2016; Sorokin *et al.*, 2017; Singleton *et al.*, 2018). Thus,

despite more than half a century of research on methanogens, their diversity, physiology and

interactions within the microbiom are still poorly understood.

It is crucial to obtain a mechanistic understanding of microbial responses and dynamics upon rewetting

to suggest robust managing practices for climate-optimal peatland protection and utilization (e.g.

paludiculture, Joosten et al., 2016). Environmental filtering, i.e. the selection of certain taxa through

abiotic environmental conditions, is considered to be a major mechanism structuring communities

(Cadotte and Tucker, 2017). In this context, identifying the main environmental drivers of both total

microbiome composition, as well as the abundance of relevant functional groups involved in carbon

cycling and methanogenesis is essential.

In this study we explore the effect of rewetting on temperate fen prokaryotic and eukaryotic microbial

communities. For this purpose, three pairs of drained (dry; terms in parenthesis are used in the

following to refer to these ecosystems and their states) and rewetted (wet) fens with contrasting water

tables and vegetation were investigated: a brackish coastal fen (Coast), a percolation fen (Perco) and

an alder carr (Alder). We hypothesize that despite the obvious structural and geo genetic differences,

congruent effects of rewetting will be detectable in the microbiomes of the three peatland types

because of environmental filtering. Possible controlling factors of community composition, like DOM

quantity and quality, soil moisture, soil/water salinity, were also determined to allow for analyzing the

effect of rewetting on key players of SOM mineralization and CH_4-cycling microorganisms.

A suite of cultivation-independent microbiological and geochemical methods was applied, including

DOM profiling as well as qPCR of the methanogenesis key enzyme *mcrA* and 16S and 18S rRNA gene

amplicon based microbiome analysis. Our results may contribute to better understand how

microbiomes respond to rewetting and to the relevant controlling environmental factors. Microbiome-

based proxies for predicting CH_4 emissions from rewetted peatlands eventually derive.

2. Results

101 **Study site and their soil properties**

102 Our investigations took place in three pairs of drained and rewetted peatlands (see table 1 and

103 materials and methods for details).

104 Table 1: Sampling site characteristics

Abbreviation	Site	State (year of rewetting)	Water table	Peat thickness	Dominant plant species
Alder	Alder carr	dry	-100 cm	60 cm	Black Alder (*Alnus glutinosa),* Ground Elder (*Aegopodium podagraria*) and Common Nettle (*Urtica dioica*)
		wet (1999)	+15 cm	> 1 m	Black Alder (*Alnus glutinosa, Greater Pond Sedge (Carex riparia)*
Coast	Coastal fen	dry	-70 cm	70 cm	Creeping Bentgrass (*Agrostis stolonifera*)
		wet (1993)	-5 cm	30 cm	Creeping Bentgrass (*Agrostis stolonifera*)
Perco	Percolation fen	dry	-20 cm	6 m	Creeping Buttercup (*Ranunculus repens*), Tufted Hairgrass (*Deschampsia cespitosa*)
		wet (1998)	+10 cm	6 m	Lesser Pond Sedge (*Carex acutiformis*)

In general, the gravimetric water content was higher in the rewetted than in the drained sites (Table S1). However, there was only a minor difference between $Perco_{dry}$ (74.2% ± 5.1) and $Perco_{wet}$ (75.4% ± 5.3), because of a high water-table near the soil surface in $Perco_{dry}$ at the time of sampling. In all of the drained sites and in $Coast_{wet}$, the water content increased with depth. In $Perco_{wet}$ and $Alder_{wet}$ (78.7 ± 4.6) with water table 10 cm above ground, the water content increased slightly with depth or did not change. The lower gravimetric water contents in $Coast_{dry}$ (45.9% ± 6.7), $Coast_{wet}$ (53.8% ± 10.03), and $Alder_{dry}$ (46.3% ± 7.0) corresponded with higher amounts of mineral soil compounds.

The sampling sites were generally acidic; pH values varied between fen types but increased with depth in all drained sites. Lower pH values were detected in $Coast_{dry}$ (3.9 - 4.0) and $Alder_{dry}$ (4.2 - 4.7), while pH values were higher in $Perco_{dry}$ (5.0-5.2) and $Coast_{wet}$ (4.7 - 6.1). In $Alder_{wet}$ (5.1 - 5.1) and $Perco_{wet}$ (5.4 – 5.4), the pH values did not change with depth (Table S1).

Salinity in the freshwater fens was generally low (0.08 - 0.49‰). The influence of the nearby brackish bay water with a salinity of 8‰ led to a higher salinity in $Coast_{dry}$ (1.32‰) and in $Coast_{wet}$ (5.2‰) (Table S1).

DOM concentrations in the soil extracts of $Alder_{wet}$ and $Coast_{wet}$ were higher than in the drained sites. In $Perco_{wet}$ DOM values measured near the surface were higher than in $Perco_{dry}$, but they showed a strong decrease with increasing depth. The distribution of separate DOM fractions, i.e. biopolymers, humic or humic-like substances and low molecular organic matter (lCOM) followed the overall pattern of total DOM (Table S1).

Total carbon (TC) and total nitrogen (TN) content of peat decreased with depth in $Alder_{dry}$, $Coast_{dry}$ and $Coast_{wet}$. In contrast, TC increased with depth in $Alder_{wet}$, $Perco_{dry}$ and $Perco_{wet}$. In $Alder_{wet}$, $Perco_{wet}$ and $Perco_{dry}$ TN did not change with depth. Total sulfur (TS) content increased with depth in $Alder_{dry}$, $Alder_{wet}$ and $Perco_{wet}$ while it decreased with depth in $Coast_{dry}$ and $Coast_{wet}$. In $Perco_{dry}$, no change was observed over depth (Table S1).

Diversity of prokaryotes and eukaryotes

Alpha diversity of prokaryotes, calculated from 16S rRNA gene amplicon sequence variants (ASVs), was significantly higher in the two freshwater peatlands than in the coastal fen (Fig. S1, $p < 0.001$). Alpha diversity was slightly higher in rewetted than in drained sites, but this was significant only in the coastal fen ($p < 0.001$). Alpha diversity showed a positive correlation with soil water content, TC, TN and DNA content (Fig. S2).

Similarly, alpha diversity of eukaryotes in freshwater peatlands was higher than in the coastal fen ($p = 0.11$). In contrast to prokaryotes, eukaryote alpha diversity was significantly lower in rewetted sites than in drained sites in Perco ($p = 0.015$), while no significant difference was observed in Alder and Coast. Alpha diversity of eukaryotes decreased with depth in the rewetted sites. For both pro- and eukaryotes, alpha diversity was negatively correlated with salinity (Fig. S2).

Microbiota composition

Community composition in 16S rRNA and 18S rRNA sequencing analysis of prokaryotes and eukaryotes showed remarkably similar patterns in the non-metric multidimensional scaling (NMDS) plot (Fig. 1), with a significant clustering of microbiota from the same sampling sites (PERMANOVA $p = 0.001$). While the communities of freshwater fens Alder and Perco showed some overlap, the microbiota in Coast were clearly separated along NMDS axis 1, possibly driven by the differences in abiotic factors, such as salinity, total nitrogen (TN), total carbon (TC), C/N ratio and DOM compounds (Fig. 1). Nevertheless, the pro- and eukaryotic microbiota of the dry and wet sites of each fen type were consistently separated along NMDS axis 2, indicating that the water table of the sites influenced the community composition (PERMANOVA $p = 0.001$, $R^2 = 0.267$). Depth showed less impact on community composition (PERMANOVA $p = 0.161$, $R^2 = 0.046$), although some clustering according to sample depth was observed.

Figure 1: NMDS plots showing (a) prokaryotic and (b) eukaryotic community compositions. Symbols

show triplicates of cores and indicate the study sites; the color indicates the depth in cm (see legend).

Environmental vectors with a significance of $p < 0.05$ are shown in black. TN: total nitrogen, hCOM:

humic like carbon, TC: total carbon, PolyC: Carbon polymers, DOM: dissolved organic matter, lCOM:

low molecular organic matter, TS: total Sulphur, C/N: carbon/nitrogen ratio.

Pro- and eukaryotic community compositions were compared between fen types, water conditions

(dry/wet) and sampling depths (Fig. 2a). We detected a higher relative abundance of Acidobacteria in

drained sites, while Betaproteobacteria, Deltaproteobacteria (mainly comprised of Myxobacteria and

sulfate reducers) and Bathyarchaeota showed higher values in all rewetted sites and Perco_dry. SPAM

and Latescibacteria were not detected in Coast, but they occurred in the freshwater fens, whereas

Actinobacteria showed higher values in Coast than in Alder and Perco.

The eukaryotic community varied considerably between fen types and water condition (Fig. 2b).

Among the fungi, ASVs of Ascomycota (which contain saprophytes, yeasts and several plant pathogens)

showed high relative abundance in each site, except Perco_dry. ASV values of Basidiomycota decreased

with depth in Alder and Coast, and showed higher relative abundance in these two site than in Perco.

ASV values of Glomerycota, which form arbuscular mycorrhiza with plants, comprising mostly

Mortirellales were high in Alder_dry and Perco_dry.

Among the protists, ASV values of Apicomplexa were high (up to 40%) in Perco$_{dry}$, but accounted for

less than 5% in other sites. Cercozoa, which contain micro-predators among other life styles, had

higher values in the freshwater sites Alder and Perco. Relative abundance of Lobosa and Ciliophora,

also potential micro-predators, was high in Alder$_{wet}$ and in sites with high water content (rewetted

sites and Perco$_{dry}$) respectively.

Among the Metazoa, Nematoda showed highest relative abundance, followed by Arthropoda. The

latter accounted for 5-10% of total amplicons in Coast and Perco$_{dry}$ and the 5-10 cm layer of Alder$_{dry}$,

while their values were lower in Alder$_{wet}$ and Perco$_{wet}$.

Figure 2: Composition of prokaryotic and eukaryotic microbial communities of drained and rewetted

fens. a) Taxonomic composition of prokaryotic 16S rRNA genes displayed on phylum level, with

Proteobacteria shown on class level. b) Taxonomic composition of eukaryotic 18S rRNA genes, on

phylum level. Bars show triplicate of cores; phyla with < 1% abundance are displayed as "other". The

sample ID on x-axis displays fen type and depth in cm.

A co-occurrence network was constructed to explore the potential interactions between pro- and

eukaryotic species. The network shows two distinct clusters of ASVs (Fig. 3). Color coding of the vertices

according to association with sites shows that ASVs within the same site were more connected than

different sites (Fig. 3b). However, ASVs associated with rewetted sites shows little connections with

those from the drained sites (Fig. 3c), highlighting the significant effect of water conditions on the co-

occurrence patterns. Similarly, the indicator taxa from Coast were mostly connected with each other,

and indicator taxa from Perco and Alder were also distinctly connected in the network (Fig. 3b).

However, there were quite some shared indicator taxa from both Perco and Alder, indicating

similarities between these two habitats compared with Coast, which we already detected in

community composition (Figs. 1 and 2). The degree and closeness centrality of the indicator taxa were

higher in the wet sites than in the dry sites (Fig. 3c). particularly in Coast, followed by Alder, and Perco

(Fig. 3b, c). The transitivity was higher among Coast indicators compared with those from Alder.

The prokaryotic indicator taxa in drained sites mainly comprised ASVs from Acidobacteria and Alpha-

and Gamma-Proteobacteria, whereas the main indicator taxa of rewetted sites were Beta- and Delta-

Proteobacteria. Beta- and Delta-Proteobacteria were also the main phyla in both hydrological status

(Fig. 2). Plants (Tracheophyta) accounted for the largest proportion of the eukaryotic indicator taxa for

both water condition and site. Nematoda ASVs accounted for a larger proportion in the drained than

in the rewetted sites; Ascomycota showed the reverse trend.

In the network with only connections between eukaryotes and prokaryotes (Fig. 3a), eukaryotes

showed higher degrees of connection compared with prokaryotes (Fig. 3). Among the eukaryotes,

fungi (Ascomycota, Basidiomycota and Glomeromycota) possessed the most connections with the

prokaryotes.

Figure 3: Co-occurrence networks. a) Co-occurrence network of prokaryotes and eukaryotes

corresponding to the indicator ASVs of 16S and 18S rRNA amplicon sequencing. ASVs with relative

abundances lower than 0.05% were discarded. b) Co-occurrence network of ASVs associated with (A)

alder carr, (C) coastal fen and (P) percolation fen. c) Co-occurrence of ASVs associated with drained (D)

and rewetted (W) sites. ASVs were associated with sites using the *indicspecies* package (De Cáceres

and Legendre, 2009).. Solid lines indicate significant positive Spearman's rank correlations

(r > 0.7. p < 0.01) between prokaryotic and eukaryotic Indicator ASVs, dashed lines indicate significant

negative correlations (r < -0.7. p < 0.01). Nodes are sized according to the number of correlating

neighbors (degree) and color-coded according to site type or water condition (drained/rewetted). The

relative abundance of prokaryotes and eukaryotes associated with the different site types is shown on

the right hand side of b) and c).

Methanogens

The abundance of methanogens in the samples, assessed by quantitative polymerase chain reaction

(qPCR) of the *mcrA* gene, ranged from 2.1×10^5 to 5.1×10^7 gene copies g^{-1} dry soil (Fig. 4). The *mcrA*

gene abundance was significantly higher in the three rewetted sites compared with the drained sites

(Kruskal-Wallis test, H = 16.5, $p < 0.001$). The difference was largest in the top soils, where *mcrA*

abundances were much higher in rewetted sites. In $Alder_{wet}$ methanogen abundance decreased

significantly with depth, while it was rather constant in $Perco_{wet}$. In contrast, methanogen abundance

in $Alder_{dry}$ was rather constant or increased with depth ($Coast_{dry}$ and $Perco_{dry}$). Positive correlations

were observed between *mcrA* abundance and water, DNA and DOM content (strongest with

biopolymers and lCOM; Fig. S2). Furthermore, TC and TN as well as pH value were positively correlated

with *mcrA* abundance (Fig. S2).

Figure 4: Abundance of methanogens. *mcrA* gene copy numbers per gram dry soil based on qPCR. X-

axis shows mcrA copies per g dry soil, drained sites in white, rewetted sites in black bars, error bars

show standard deviation of triplicate soil cores.

The relative abundance of methanogens in the prokaryotic microbiota was low. Methanogen-affiliated

ASVs in the amplicon sequence datasets were on average below 0.25% of total 16S rRNA amplicons

(Fig. 5). In most of the drained fens, methanogen ASVs were even absent, although they were

detectable by *mcrA* qPCR (see above, Fig. 4). Upper soil samples of Alder$_{wet}$ and Coast$_{wet}$ showed

highest abundances, while methanogens were more abundant in the lower samples of Perco$_{wet}$.

Methanobacteriales and Methanosarcinales (including the families Methanosaetaceae and

Methanosarcinaceae) were the dominant orders in Alder$_{wet}$ and Coast$_{wet}$, while methanogens were

more diverse in Perco$_{wet}$, where Methanomassiliicoccales and Methanomicrobiales were also present.

Despite apparently missing out on some methanogens in the amplicon data, the relative abundance

calculated from *mcrA* gene copies per g dry soil (qPCR) shows a clear positive correlation with the

relative abundance of methanogenic archaea in 16S rRNA amplicon (Spearman correlation, R= 0.578;

p < 0.001).

Figure 5: Relative abundance of functional groups in 16S rRNA gene amplicon datasets. a) Methanogens, b) methanotrophic bacteria and archaea, c) sulfate reducers. Barplots show relative abundance. X-axes represent fen type: alder carr (A), coastal fen (C), percolation fen (P), hydrological state drained (D) and rewetted (W), depth in cm (05, 15, 25). Data are shown as mean values of triplicate soil cores.

Methanotrophs

16S rRNA genes of known methanotrophic prokaryotes were found in all sites (Fig. 5b), but never exceeded 1% of all prokaryotic 16S rRNA genes on average. Their relative abundance and taxon composition differed both between fen types and dry/wet. In drained fens ASVs affiliated with *Ca.* Methylacidiphilum (phylum Verrucomicrobia) were dominant across the soil profile. In contrast, ASVs of Methylococcaceae (class Gammaproteobacteria) were the most abundant methanotrophs in Coast_wet and Perco_wet. Putative anaerobic methanotrophs affiliated with *Ca.* Methanoperedens

(ANME-2d, archaea) and with *Ca*. Methylomirabilis (NC10 phylum), were the dominant methanotroph

guild at the Alder$_{wet}$ site.

Sulfate reducers

ASVs of known sulfate reducers were generally more abundant than methanogens and methanotrophs

(between 0.1% and 5% on average, Fig. 5c). Like methanogens, their relative abundance was higher in

rewetted sites than in drained sites. Highest relative abundance was observed in the Coast$_{wet}$, which is

exposed to recurrent influx of sea water. ASVs of Desulfobacterales were particularly abundant in this

site. Desulfarculales and Desulfosporosinus were only detected in rewetted sites, while

Desulfuromonadales occurred in all sites. Relative abundance of sulfate reducers was positively

correlated to salinity, but not to total sulfur (Fig. S2).

3. Discussion

To our knowledge, this is the first comprehensive study investigating the impact of long-term rewetting

(>> 10 years) on freshwater and coastal fen microbiomes. The parallel study of three different fen

types allowed for the identification of overarching factors next to fen type-specific factors. The NMDS

plots of both prokaryotes and eukaryotes were almost superimposable. It is remarkable that rewetting

caused highly congruent effects on beta-diversity of all three domains of life and it is conceivable that

water table and salinity were the main environmental filters among the physicochemical controlling

factors. Rewetting resulted in more anoxic conditions in the rewetted sites compared with the drained

sites. Increased anoxia is supported by several lines of evidence, including (1) a higher moisture

content and (2) higher proportions of presumably fermentative microorganisms and (3) higher

proportion of taxa with different types of anaerobic respiration, such as methanogenesis and sulfate

reduction.

Environmental filtering effect on pro- and eukaryotic community composition

The network analysis confirmed the conclusions we drew from the community composition: there are

more taxa indicative for each of the different fen types than for water status. Consequently, site

location has a more significant effect on the community assemblages. The higher degree and closeness

centrality of indicators in rewetted sites indicates more connections in these sites. This might be due

to that higher water flow rates in soils create greater homogeneity, and hence weak niche

differentiation, contributing to stronger interactions between soil microbes. Further, it might reflect

the close metabolic interactions in anaerobic microbiomes during plant biomass degradation (Schink,

1997; Tveit *et al.*, 2015). Main taxa of indicator ASVs in rewetted sites were Chloroflexi and Delta- and

Betaproteobacteria, which contained ASVs known from anoxic environments, also highly abundant in

the wet communities (Fig. 2 and Fig. 3) (He *et al.*, 2015). In the drained sites, most indicator taxa were

from Acidobacteria, the predominant phylum, which would correspond with the general acidic site

conditions. Drainage and subsequent oxidation processes in peat soils are associated with acid

production and depending on the buffer capacity of soils pH may go down by one or two units

compared with pre-drained conditions (Lamers, 1998). In contrast, rewetting with base-rich water will

increase alkalinity and pH (Cusell *et al.*, 2013). Especially the two dominant classes Acidobacteria

(subdivision 1) and Solibacteres (subdivision 3) decreased in relative abundance upon rewetting

(Supplementary Fig. S4), which is in line with the obligatory aerobic lifestyle of most characterized

member species (Eichorst *et al.*, 2018). Members of the aerobic Acidobacteria play important roles in

carbohydrate degradation, but some subdivisions also contain facultative or obligate anaerobic species

(Eichorst *et al.*, 2018; Hausmann *et al.*, 2018). Other putative anaerobic phyla such as Ignavibacter and

Bathyarchaeota occurred only in rewetted sites and deep layers of drained sites. Obviously, soil

moisture acted as environmental filter leading to community compositional changes from oxygen-

dependent taxa (drained) to anaerobic and fermentative taxa in the rewetted peat soils (Fig. S3). A

second likely environmental filter in our study is salinity. The higher connectivity and transitivity of

indicators from Coast may result from environmental filtering through higher salinity in these sites,

leading to a change in microbial community with increases of Actinobacteria and sulfate reducers and

decrease of methanogens (Dang et al., 2018, and see below).

Remarkably, the largest proportion of eukaryotic indicator ASVs originated from Tracheophyta

(plants), indicating that the interaction between plants and the soil microbiota is very important.

Similar to the prokaryotes, the interaction was even stronger in rewetted sites than in drained sites. In

Coast with the lowest moisture content, Tracheophyta ASVs showed lower values than in Perco and

Alder. Among the animals, Nematoda indicator ASVs accounted for a larger proportion in drained sites,

likely because they are dependent on oxic conditions.

In the network, we also observed that eukaryotes had higher degrees than prokaryotes. The

prokaryotes mostly connected with fungi, like Ascomycota, Basidiomycota and Glomeromycota,

suggesting that especially fungi may drive the composition of prokaryotic communities. We found

fewer eukaryotic than prokaryotic ASVs, suggesting that one eukaryote was probably connected with

multiple prokaryotes, as implied by higher degrees of connection for eukaryotes. Similarly, the richness

of eukaryotes was far less than the richness of prokaryotes, which is common for soil microbiota, as

shown before in e.g. Tveit et al. (2015). Therefore, our results potentially suggest this as a universal

rule of pro-eukaryote interactions.

The influence of rewetting on methanogens

The important group of methanogenic archaea was hardly included in the analysis of microbial

networks and played no role in the determination of indicator taxa. The reason was the low relative

abundance of methanogens in the amplicon datasets of the microbiota, being less than three per mil

in all samples. Although their relative abundance in the microbiota was low even under rewetted

conditions, their abundance per gram soil was high, especially in the rewetted sites. Wet peatlands are

known to contribute a large share of the global CH_4 emissions (Nazaries *et al.*, 2013; Joosten *et al.*,

2016). The higher *mcrA* abundance after rewetting (Fig. 4) could lead to increased CH_4 emissions from

peatlands, as is the case in rice paddy fields (Reim *et al.*, 2017; Wen *et al.*, 2018). Interestingly, their

abundance was high in the rewetted top soils, at least in $Alder_{wet}$ and $Perco_{wet}$, and not in the subsoils.

This is in contrast to general assumptions about their vertical distribution in the soil profile (reviewed

in Dean et al., 2018) and corroborates recent findings from other ecosystems (Hahn-Schöfl *et al.*, 2011;

Angle *et al.*, 2017). This observation makes peatland management practices such as top soil removal

attractive for mitigation of CH_4 emissions from peatlands (Harpenslager *et al.*, 2015; Zak *et al.*, 2018;

A.Guenther, unpublished). The abundance of methanogens was strongly correlated with soil water

content (Fig. S2), which drives anoxia in the soil. However, many other factors like substrate availability

of DOM, especially lCOM, C and N content of soil, degradation status of the peat and sulfate and nitrate

concentration all correlated with methanogen abundance per gram dry soil as well (Fig. S2). One

significant factor was salinity, resulting from brackish water intrusions from the Baltic Sea in $Coast_{wet}$,

which led to large differences between freshwater fens (Alder and Perco) and Coast. In $Coast_{wet}$ the

abundance of methanogens per gram soil was at least 10x lower than in the freshwater sites $Alder_{wet}$

and $Perco_{wet}$. Thus, this environmental filtering through salinity likely leads to lower CH_4 emissions,

also shown before in several studies(Neubauer *et al.*, 2013; He *et al.*, 2015; van Dijk *et al.*, 2017).

The strong positive correlation of *mcrA* gene abundance in qPCR with the relative abundance of

methanogenic archaea in 16S rRNA gene amplicon datasets proves the validity of the measured results.

However, in most of the drained peats ($Perco_{dry}$, most $Alder_{dry}$ and $Coast_{dry}$), methanogenic taxa were

not present in the dataset, although *mcrA* genes were quantified by qPCR. This can be explained by

the rather shallow average sequencing depth of 20,000 reads per sample, giving a taxon detection limit

of about 0.05‰ in the microbiota, and thus taxa with lower abundances could not be detected. Still,

the distribution of methanogenic taxa led to the assumption that substrate availability was different

in the sites. While hydrogenotrophic Methanobacteriales were dominant in $Coast_{wet}$, $Coast_{dry}$ and

$Alder_{wet}$, $Perco_{wet}$ showed a high variability in methanogenic groups. Here the Methanosarcinales

families of Methanosaetaceae (obligate acetoclastic) and Methanosarcinaceae (flexible substrate

spectrum) occurred, as did the Methanomicrobiales and the Methanomassiliicoccales (that use the H_2

dependent methylotrophic pathway). Methanomassiliicoccales were detected in every site with

Methanomassiliicoccales specific qPCR primers AS1/AS2 (Mihajlovski *et al.*, 2010) (data not shown).

Their high abundance in Perco$_{wet}$ indicates that besides the well reported hydrogenotrophic and

acetotrophic pathways (Lyu *et al.*, 2018), methylotrophic methanogenesis may contribute more to

peatland CH$_4$ emissions than currently assumed (Söllinger *et al.*, 2016; Zalman *et al.*, 2017).

Interactions between methanogens, methanotrophs and sulfate reducers

The second important player in peatland CH$_4$ emissions is the heterogenic group of methanotrophic

Bacteria and Archaea, acting as biological CH$_4$ filter through aerobic or anaerobic oxidation of the

produced CH$_4$ (reviewed in Nazaries et al., 2013 and Knief, 2015). Do the methanogens influence

abundance of methanotrophs? The methanotrophic groups differed with site, and were generally

lower abundant in the drained sites where *Ca.* Methylacidiphilum aerobic CH$_4$ oxidizers were

dominant. These organisms were originally isolated from acidic and volcanic environments, but have

recently also been found in pH neutral environments (van Teeseling *et al.*, 2014; Vaksmaa *et al.*, 2016).

They are characterized by growth on low CH$_4$ concentrations, and are possibly atmospheric CH$_4$

oxidizers, which would make them independent of substrate supply by methanogens. As found in

other wetland studies (e.g. Tveit et al., 2013, Smith et al., 2018) aerobic CH$_4$ oxidizers of

Methylococcaceae dominated in the wet sites. The anaerobic group *Ca.* Methylomirabilis (Vaksmaa *et*

al., 2016), which uses oxygen from nitrite to reduce CH$_4$ appeared in the sites with highest water

content (Alder$_{wet}$, Perco$_{dry}$ and Perco$_{wet}$). The anaerobic CH$_4$ oxidizing ANME-2d (*Ca.*

Methanoperedens) which reduce CH$_4$ by the reverse methanogenic pathway (Welte *et al.*, 2016) were

detected in the water-covered fen Alder$_{wet}$. Although we did not find a clear correlation between

methanogen and methanotroph abundance (see Wen *et al.*, 2018 for similar findings), we did find a

strong division of anaerobic and aerobic methanotrophs depending on water content and therefore

oxygen availability, between sites and depths.

What is the basis of the environmental filter leading to the low methanogen abundance in the coastal

fen? It is known that sulfate reducers can have an impact on methanogens, because methanogens and

sulfate reducers not only share the anaerobic niche, but also compete for the same substrates, such

as hydrogen (Kristjansson and Schönheit, 1983; Thauer *et al.*, 2008; He *et al.*, 2015) and acetate

(Kristjansson *et al.*, 1982). Due to their higher substrate affinity, sulfate reducers can outcompete

methanogens, leading to lower methanogen abundance in the presence of sulfate reducers

(Kristjansson *et al.*, 1982; van Dijk *et al.*, 2017). In fact, Desulfobacterales, known competitors for H_2

as a substrate for methanogenesis, occurred in the coastal site Coast$_{wet}$ with a large proportion, which

was associated with a more than tenfold lower *mcrA* gene abundance than in Alder$_{wet}$ and Perco$_{wet}$.

The influence of salinity in tidal marshes is also known to decrease CH_4 emission compared to

freshwater sites (Poffenbarger *et al.*, 2011; van Dijk *et al.*, 2017). Although no CH_4 emissions were

measured in this study, the higher methanogen abundance in freshwater peatland suggests higher

emission rates (Bartlett *et al.*, 1987) than from the coastal fen. Our study provides microbiome-based

insights into the environmental filtering effects of rewetting of fen ecosystems and suggests that, in

terms of minimizing CH_4 emissions, increased salinity and/or top soil removal might be promising

options. Future analysis of seasonal microbiome dynamics and measurements of CH_4 fluxes will

provide evidence, whether these predictions hold true.

4. Experimental procedures

Sites and sampling

All sampling sites, alder carr, coastal fen and percolation fen, are located in north-eastern Germany

(Fig. S5). The alder carr is part of the Recknitz river valley, was drained at least from 1786 on and has

been used as a managed forest. The sub-site Alder$_{wet}$ (54°13'N 12°49'E) was unintentionally flooded in

1999 due to a blocked drain and has remained wet since then. In the sub-site Alder$_{dry}$ (54°13'N 12°51'E)

long-term deep drainage Alder$_{dry}$ has caused subsidence of the soil surface of about 1 m.

The coastal fen, located on a peninsula, was naturally episodically flooded by brackish seawater coming

from the Bay of Greifswald of the Baltic Sea, typically during winter storm surges. The entire area was

diked in 1850 to allow for pasture and other agricultural use. In 1993, the dykes around the sub site

Coast$_{wet}$ (54°16'N 13°39'E) were removed and, subsequently, the site was episodically flooded again.

The sub-site Coast$_{dry}$ (54°16'N 13°39'E) in contrast remains behind the dyke and, thus, cut off from the

seawater influence. Both, Coast$_{wet}$ and Coast$_{dry}$ are used as cattle pastures.

The percolation fens, representing the third type of peatlands, are located in the catchment areas of

the rivers Trebel and Recknitz. Both sites were deeply drained for intensive land-use and used as deep-

drained grassland in the 20th century. In 1998, the water table at the sub-site Perco$_{wet}$ (54°10'N 12°74'E)

in the Trebel valley was raised again to 5-10 cm above ground level as part of a large rewetting project

(EU LIFE project) (Ministerium für Bau, Landesentwicklung und Umwelt, M-V 1998). In contrast, the

water table of the still drained sub-site Perco$_{dry}$ (54°13'N 12°63'E) in the Recknitz valley remains several

decimeters below the surface and is mown 1-2 times per year.

At all sites peat samples were taken as triplicates at 5-10 cm, 15-20 cm and 25-30 cm depths (Fig. S5)

using a gouge. The samples were stored in 50-ml reaction tubes on ice over night until further

processing.

Soil physico-chemical properties

Soil moisture was measured gravimetrically by drying 2-3 g of soil over night at 90°C until mass

constancy. Soil moisture is expressed as the percentage of lost water weight to wet soil weight. The

potential soil pH was measured at room temperature with a digital pH meter (pH 540 GLP, WTW,

Weilheim, Germany) in 0.01 M CaCl$_2$ solution with a 1: 2.5 soil to solution ratio. Concentrations of total

C, N and S were measured using a CNS analyzer (Vario MICRO cube – Elementar Analysensysteme

GmbH Langenselbold, Germany). DOM was measured in soil extracts derived from mixing 3 g of

previously frozen soil with 30 ml 0.1 M NaCl in 50 ml reaction tubes with subsequent shaking (vortex,

180 rpm) for 30 min. Extracts were filtered through 0.45 µm (pore size) sodium acetate filters, which

were prewashed with 50 µl deionized H$_2$O to remove soluble acetate. The concentrations and the

composition of DOM based on size categories were determined using a size-exclusion chromatography

(SEC) with organic carbon and organic nitrogen detection (LC-OCD-OND analyzer, DOC-Labor Huber,

Karlsruhe, Germany) (Huber *et al.*, 2011). The DOM was classified into three major sub-categories: (i)

biopolymers, (ii) humic or humic-like substances including building blocks, and (iii) low molecular-

weight substances. Time between sample preparation and analysis did never exceed two weeks and

samples were stored at 5°C to avoid significant changes of DOM composition (Heinz and Zak, 2018).

Soluble reactive phosphorus was determined with the ammonium molybdate spectrometric method

(DIN EN 1189 D11) using a Cary 1E Spectrophotometer (Varian). $N-NH_4^+$ and $N-NO_3^-$ were determined

calorimetrically using the photometry CFA method (Skalar SAN, Skalar Analytical B.V., The

Netherlands) according to the guidelines in EN ISO 11732 (DEV-E 23) and EN ISO 13395 (DEV, D 28),

respectively.

DNA extraction and PCR/qPCR

DNA was extracted from 0.25 g soil using the DNeasy PowerSoil Kit (QIAGEN, Hilden, Germany)

according to the manufacturer´s recommendation. The bead beating step was performed using a

FastPrep®-24 5G instrument (MP Biomedicals, Santa Ana, USA), with intensity 5.0 for 45 s. DNA

concentrations were measured with Qubit®2.0 dsDNA High Sensitivity and dsDNA Broad Range

assays (Thermo Fisher, Waltham, USA). DNA size was analyzed using 1% agarose gels.

Quantitative polymerase chain reaction was used to measure the abundances of the *mcrA* gene, coding

for methyl coenzyme M reductase subunit A. Quantification of *mcrA* gene copies in peat soil DNA

extracts was performed on a qTOWER 2.2 (Analytic Jena, Jena, Germany). The primer pair of forward

mlas-mod and reverse *mcrA*- rev (Steinberg and Regan, 2009) was used. For each reaction, 15 µl of

PCR mixture contained 7.5 µl of innuMIX qPCR MasterMix SyGreen (Analytic Jena, Jena, Germany),

0.75 µl of each primer (10 pmol/µl), 5 µl of ddH2O and 1 µl of template DNA. Duplicate measurements

of two concentrations (1 and 2 ng/µl) were performed for each DNA sample. Assay condition was 95°C

for 5 min, 35 cycles of 95°C for 30 s, 55°C for 45 s and 72°C for 45°C, followed by melting curve analysis.

PCR-product specificity was confirmed by melting point analysis. *mcrA* gene copy numbers were

calculated from a standard curve obtained by serially diluting a standard from 10^6 to 10^1 *mcrA* gene

copies per µl. The standard was created with above-mentioned primers by amplifying *mcrA* genes from

cow rumen fluid (Söllinger *et al.*, 2016) and cloning them into the pGEM®-Teasy vector system

(Promega, Mannheim, Germany). Amplicons for *mcrA* standard were generated with vector-specific

primers sp6 and T7 and the resulting PCR product was cleaned with DNA purification kit (Biozym,

Hessisch Oldendorf, Germany), quality-controlled by agarose gel electrophoresis and quantified with

Qubit®2.0 dsDNA High Sensitivity assay. The qPCR assay had the following parameters: slope: 3.41-

3.52, efficiency: 0.92-0.96, R^2: > 0.995.

Amplicon sequencing and bioinformatics

16S and 18S rRNA gene amplicons were prepared and sequenced using 300 bp paired-end read

Illumina MiSeq V3 sequencing by LGC Genomics (Berlin, Germany). This comprised 16S and 18S rRNA

gene PCR amplification, Illumina MiSeq library preparation and sequencing using the primer pair 515YF

(GTG YCA GCM GCC GCG GTA A)/ B806R (GGA CTA CNV GGG TWT CTA AT) (Walters *et al.*, 2015) for

prokaryotes and 1183F (AATTTGACTCAACRCGGG)/ 1443R (GRGCATCACAGACCTG) (Ray *et al.*, 2016)

for eukaryotes.

The data were processed using R version 3.5.1 (The R Core Team, 2018). 16S rRNA and 18S rRNA gene

amplicon sequences were quality filtered, de-replicated and clustered into ASVs using the *dada2*

package (Callahan *et al.*, 2016). The chimeric sequences were de-novo checked and removed with

dada2. Then the representative sequence of each ASV was assigned to taxonomy against a modified

version of the SILVA SSUref_NR_128 data base, containing an updated taxonomy for *Archaea* (Lanzén

et al., 2012), using the programs BLASTn (Altschul *et al.*, 1990) and Megan 5 (Huson *et al.*, 2016). ASVs

that were assigned to Chloroplast or mitochondria were filtered. Due to the low number of sequences

(<500) in one sample (Coast$_{dry}$ 2 15) for both 16S and 18S rRNA genes, this sample was discarded from

further analysis. After filtering, 1,134,953 and 3,990,919 sequences, sequences were retained and

were clustered into 7697 and 5548 ASVs for 16S and 18S rRNA genes, respectively. Plant sequences

which accounted for 288 ASVs were filtered for downstream analysis to increase the resolution of

other eukaryotic taxa.

Alpha diversity (Shannon entropy) of the microbial community was calculated using the package

phyloseq (McMurdie and Holmes, 2013). ASV count table was then normalized using

metagenomeSeq's CSS (cumulative sum scaling) (Paulson *et al.*, 2013). NMDS was performed to

analyze the community composition using the package *vegan* (Oksanen *et al.*, 2016) with defaults.

Pairwise Spearman's rank correlations were conducted among soil properties, DNA concentrations,

qPCR data, NMDS axes and methanogen relative abundances using the package *Hmisc* to detect the

significance of the impact of the factors on microbial communities (Harrell and Dunpont, 2019). The

soil properties with significant impact were fitted into NMDS ordination using the envfit function in

the *vegan* package.

The correlations were visualized by heatmap using *corrplot*. All *P*-values for multiple comparisons were

adjusted by the false discovery rate (FDR) method and the null hypothesis was rejected when *P*-values

were less than 0.05.

A co-occurrence network was constructed to explore the potential interactions between species. Plant

ASVs were included in network analysis because plant residues are a major part of peat and they might

be important in interactions between microbes. To eliminate the influence of rare taxa, ASVs with

relative abundances lower than 0.05% were discarded. The pairwise Spearman's rank correlations

were conducted with the package Hmisc and all *P*-values were adjusted by the FDR. The cut-offs of

adjusted *P*-value and correlation coefficient were 0.01 and 0.7, respectively. The network was

visualized using package *igraph* (Csardi, 2019). A network showing the connections between

eukaryotes and prokaryotes was sub-selected from the original network by keeping the edges

connecting these two groups. The indicators of each group or group combination of a factor (site or

water condition) were identified using *indicspecies* package.

The microbial ecological functional groups were predicted using FAPROTAX (Louca *et al.*, 2016), The

identified groups were verified based on literature knowledge. ANME2-D group within

Methanosarcina known as methanotrophic, using the methanogenic pathway in reverse, was excluded

from methanogens. All plots were created using package *ggplot2* (Wickham, 2016).

Acknowledgement

The authors would like to thank Florian Beyer for providing the maps and arrangement of figure S5.

The European Social Fund (ESF) and the Ministry of Education, Science and Culture of Mecklenburg-

Western Pomerania (Germany) funded this work within the scope of the project WETSCAPES

(ESF/14-BM-A55-0034/16 and ESF/14-BM-A55-0030/16).

References

Altschul, S.F., W. Gish, W. Miller, E.W. Myers, and D.J. Lipman. 1990. Basic local alignment

search tool. J. Mol. Biol. doi: 10.1016/S0022-2836(05)80360-2.

Angle, J.C., T.H. Morin, L.M. Solden, A.B. Narrowe, G.J. Smith, M.A. Borton, C. Rey-Sanchez,

R.A. Daly, G. Mirfenderesgi, D.W. Hoyt, W.J. Riley, C.S. Miller, G. Bohrer, and K.C.

Wrighton. 2017. Methanogenesis in oxygenated soils is a substantial fraction of wetland

methane emissions. Nat. Commun. 8(1): 1567. doi: 10.1038/s41467-017-01753-4.

Bartlett, K.B., D.S. Bartlett, R.C. Harriss, and D.I. Sebacher. 1987. Methane emissions along a

salt marsh salinity gradient. Biogeochemistry 4(3): 183–202. doi: 10.1007/BF02187365.

Callahan, B.J., P.J. McMurdie, M.J. Rosen, A.W. Han, A.J.A. Johnson, and S.P. Holmes. 2016.

DADA2: High-resolution sample inference from Illumina amplicon data. Nat. Methods

13(7): 581–583. doi: 10.1038/nmeth.3869.

Conrad, R. 1996. Soil Microorganisms as Controllers of Atmospheric Trace Gases (H 2 , CO ,

CH 4 , OCS , N 2 O , and NO). 60(4): 609–640.

Couwenberg, J., A. Thiele, F. Tanneberger, J. Augustin, S. Bärisch, D. Dubovik, N.

Liashchynskaya, D. Michaelis, M. Minke, A. Skuratovich, and H. Joosten. 2011. Assessing

greenhouse gas emissions from peatlands using vegetation as a proxy. Hydrobiologia

674(1): 67–89. doi: 10.1007/s10750-011-0729-x.

Dang, C., E.M. Morrissey, W. Virginia, S.C. Neubauer, and R.B. Franklin. 2018. Novel microbial

community composition and carbon biogeochemistry emerge over time following

saltwater intrusion in wetlands. (July). doi: 10.1111/gcb.14486.

Eichorst, S.A., D. Trojan, S. Roux, C. Herbold, T. Rattei, and D. Woebken. 2018. Genomic

insights into the Acidobacteria reveal strategies for their success in terrestrial

environments. Environ. Microbiol. 20(3): 1041–1063. doi: 10.1111/1462-2920.14043.

Estop-Aragonés, C., K. Zajac, and C. Blodau. 2016. Effects of extreme experimental drought

and rewetting on CO_2 and CH_4 exchange in mesocosms of 14 European peatlands with

different nitrogen and sulfur deposition. Glob. Chang. Biol. 22(6): 2285–2300. doi:

10.1111/gcb.13228.

Hahn, J., S. Köhler, S. Glatzel, and G. Jurasinski. 2015. Methane exchange in a coastal fen in

the first year after flooding - A systems shift. PLoS One 10(10). doi:

10.1371/journal.pone.0140657.

Hausmann, B., C. Pelikan, C.W. Herbold, S. Köstlbacher, M. Albertsen, S.A. Eichorst, T.

Glavina Del Rio, M. Huemer, P.H. Nielsen, T. Rattei, U. Stingl, S.G. Tringe, D. Trojan, C.

Wentrup, D. Woebken, M. Pester, and A. Loy. 2018. Peatland Acidobacteria with a

dissimilatory sulfur metabolism. ISME J.: 1–14. doi: 10.1038/s41396-018-0077-1.

He, S., S. a Malfatti, J.W. Mcfarland, F.E. Anderson, A. Pati, M. Huntemann, J. Tremblay, T.

Glavina, M.P. Waldrop, L. Windham-myers, and G. Tringe. 2015. Patterns in Wetland

Microbial Community Composition and Functional Gene Repertoire Associated with

Methane Emissions. Mbio.Asm.Org 6(3): 1–15. doi: 10.1128/mBio.00066-15.Editor.

Heinz, M., and D. Zak. 2018. Storage effects on quantity and composition of dissolved

organic carbon and nitrogen of lake water, leaf leachate and peat soil water. Water Res.

130: 98–104. doi: 10.1016/j.watres.2017.11.053.

Huber, S.A., A. Balz, M. Abert, and W. Pronk. 2011. Characterisation of aquatic humic and

non-humic matter with size-exclusion chromatography - organic carbon detection -

organic nitrogen detection (LC-OCD-OND). Water Res. 45(2): 879–885. doi:

10.1016/j.watres.2010.09.023.

Huson, D.H., S. Beier, I. Flade, A. Górska, M. El-Hadidi, S. Mitra, H.J. Ruscheweyh, and R.

Tappu. 2016. MEGAN Community Edition - Interactive Exploration and Analysis of

Large-Scale Microbiome Sequencing Data. PLoS Comput. Biol. 12(6): 1–12. doi:

10.1371/journal.pcbi.1004957.

Joosten, H., and J. Couwenberg. 2009. Are emission reductions from peatlands measurable,

reportable and verifiable? Wetlands: 14. www.wetlands.org.

Joosten, H., A. Sirin, J. Couwenberg, J. Laine, and P. Smith. 2016. The role of peatlands in

climate regulation. Peatl. Restor. Ecosyst. Serv. Sci. Policy Pract. (May): 63–76. doi:

10.1017/CBO9781139177788.005.

Knief, C. 2015. Diversity and habitat preferences of cultivated and uncultivated aerobic

methanotrophic bacteria evaluated based on pmoA as molecular marker. Front.

Microbiol. 6(DEC). doi: 10.3389/fmicb.2015.01346.

Kristjansson, J.K., and P. Schönheit. 1983. Why do sulfate-reducing bacteria outcompete

methanogenic bacteria for substrates? Oecologia 60(2): 264–266. doi:

10.1007/BF00379530.

Lanzén, A., S.L. Jørgensen, D.H. Huson, M. Gorfer, S.H. Grindhaug, I. Jonassen, L. Øvreås, and

T. Urich. 2012. CREST - Classification Resources for Environmental Sequence Tags. PLoS

One 7(11). doi: 10.1371/journal.pone.0049334.

Louca, S., L. Wegener Parfrey, and M. Doebeli. 2016. Decoupling function and taxonomy in

the global ocean microbiome. Science 353(6305): 1272–1277. doi:

10.1126/science.aaf4507.

Lyu, Z., N. Shao, T. Akinyemi, and W.B. Whitman. 2018. Current Biology. (Figure 1): 727–732.

doi: 10.1016/j.cub.2018.05.021.

Mihajlovski, A., J. Doré, F. Levenez, M. Alric, and J.F. Brugère. 2010. Molecular evaluation of

the human gut methanogenic archaeal microbiota reveals an age-associated increase of

the diversity. Environ. Microbiol. Rep. 2(2): 272–280. doi: 10.1111/j.1758-

2229.2009.00116.x.

Nazaries, L., J.C. Murrell, P. Millard, L. Baggs, and B.K. Singh. 2013. Methane, microbes and

models: Fundamental understanding of the soil methane cycle for future predictions.

Environ. Microbiol. 15(9): 2395–2417. doi: 10.1111/1462-2920.12149.

Neubauer, S.C., R.B. Franklin, and D.J. Berrier. 2013. Saltwater intrusion into tidal freshwater

marshes alters the biogeochemical processing of organic carbon. Biogeosciences

10(12): 8171–8183. doi: 10.5194/bg-10-8171-2013.

Parish, F., A. Sirin, D. Charman, H. Joosten, T. Minayeva, M. Silvius, and L. Stringer. 2008.

Assessment on Peatlands, Biodiversity and Climate Change: Main Report.

Paulson, J.N., O. Colin Stine, H.C. Bravo, and M. Pop. 2013. Differential abundance analysis

for microbial marker-gene surveys. Nat. Methods. doi: 10.1038/nmeth.2658.

Poffenbarger, H.J., B.A. Needelman, and J.P. Megonigal. 2011. Salinity influence on methane

emissions from tidal marshes. Wetlands 31(5): 831–842. doi: 10.1007/s13157-011-

0197-0.

Ray, J.L., J. Althammer, K.S. Skaar, P. Simonelli, A. Larsen, D. Stoecker, A. Sazhin, U.Z. Ijaz, C.

Quince, J.C. Nejstgaard, M. Frischer, G. Pohnert, and C. Troedsson. 2016.

Metabarcoding and metabolome analyses of copepod grazing reveal feeding preference

and linkage to metabolite classes in dynamic microbial plankton communities. Mol.

Ecol. 25(21): 5585–5602. doi: 10.1111/mec.13844.

Reim, A., M. Hernández, M. Klose, A. Chidthaisong, M. Yuttitham, and R. Conrad. 2017.

Response of methanogenic microbial communities to desiccation stress in flooded and

rain-fed paddy soil from Thailand. Front. Microbiol. 8(MAY): 1–17. doi:

10.3389/fmicb.2017.00785.

Schink, B. 1997. Energetics of syntrophic cooperation in methanogenic degradation.

Microbiol. Mol. Biol. Rev. 61(2): 262–80.

http://www.ncbi.nlm.nih.gov/pubmed/9184013.

Singleton, C., C. McCalley, B. Woodcroft, J. Boyd, P. Evans, S. Hodgkins, J. Chanton, S.

Frolking, P. Crill, S. Saleska, V. Rich, and G. Tyson. 2018. Methanotrophy across a natural

permafrost thaw environment. ISME J. 1. doi: 10.1038/s41396-018-0065-5.

Söllinger, A., C. Schwab, T. Weinmaier, A. Loy, A.T. Tveit, C. Schleper, and T. Urich. 2016.

Phylogenetic and genomic analysis of Methanomassiliicoccales in wetlands and animal

intestinal tracts reveals clade-specific habitat. FEMS Microbiol. Ecol. 92(1): 1–12. doi:

10.1093/femsec/fiv149.

Sorokin, D.Y., K.S. Makarova, B. Abbas, M. Ferrer, P.N. Golyshin, E.A. Galinski, S. Ciordia, M.C.

Mena, A.Y. Merkel, Y.I. Wolf, M.C.M. van Loosdrecht, and E. V. Koonin. 2017. Discovery

of extremely halophilic, methyl-reducing euryarchaea provides insights into the

evolutionary origin of methanogenesis. Nat. Microbiol. 2(May): 17081. doi:

10.1038/nmicrobiol.2017.81.

Steinberg, L.M., and J.M. Regan. 2009. mcrA-targeted real-time quantitative PCR method to

examine methanogen communities. Appl. Environ. Microbiol. 75(13): 4435–4442. doi:

10.1128/AEM.02858-08.

van Teeseling, M.C.F., A. Pol, H.R. Harhangi, S. van der Zwart, M.S.M. Jetten, H.J.M. Op den

Camp, and L. van Niftrik. 2014. Expanding the verrucomicrobial methanotrophic world:

Description of three novel species of Methylacidimicrobium gen. nov. Appl. Environ.

Microbiol. 80(21): 6782–6791. doi: 10.1128/AEM.01838-14.

Thauer, R.K., A.-K. Kaster, H. Seedorf, W. Buckel, and R. Hedderich. 2008. Methanogenic

archaea: ecologically relevant differences in energy conservation. Nat. Rev. Microbiol.

6(8): 579–591. doi: 10.1038/nrmicro1931.

The R Core Team. 2018. R: A language and environment for statistical computing.

Tveit, A.T., T. Urich, P. Frenzel, and M.M. Svenning. 2015. Metabolic and trophic interactions

modulate methane production by Arctic peat microbiota in response to warming. Proc.

Natl. Acad. Sci. 112(19): E2507–E2516. doi: 10.1073/pnas.1420797112.

Vaksmaa, A., C. L??ke, T. van Alen, G. Val??, E. Lupotto, M.S.M. Jetten, and K.F. Ettwig.

2016a. Distribution and activity of the anaerobic methanotrophic community in a

nitrogen-fertilized Italian paddy soil. FEMS Microbiol. Ecol. 92(12): 1–11. doi:

10.1093/femsec-fiw181.

Vaksmaa, A., C. Lüke, T. van Alen, G. Valè, E. Lupotto, M.S.M. Jetten, and K.F. Ettwig. 2016b.

Distribution and activity of the anaerobic methanotrophic community in a nitrogen-

fertilized Italian paddy soil. FEMS Microbiol. Ecol. 92(12): 1–11. doi: 10.1093/femsec-

fiw181.

Walters, W., E.R. Hyde, D. Berg-lyons, G. Ackermann, G. Humphrey, A. Parada, J.A. Gilbert,

J.K. Jansson, J.G. Caporaso, J.A. Fuhrman, A. Apprill, and R. Knight. 2015. Improved

Bacterial 16S rRNA GEne(V4 and V4-V5) and Fungal Internal Transcribed Spacer Marker

Gene Primers for Microbial Community Surveys. mSystems 1(1): e0009-15. doi:

10.1128/mSystems.00009-15.Editor.

Welte, C.U., O. Rasigraf, A. Vaksmaa, W. Versantvoort, A. Arshad, H.J.M. Op den Camp,

M.S.M. Jetten, C. Lï¿½ke, and J. Reimann. 2016a. Nitrate- and nitrite-dependent

anaerobic oxidation of methane. Environ. Microbiol. Rep. 8(6): 941–955. doi:

10.1111/1758-2229.12487.

Welte, C.U., O. Rasigraf, A. Vaksmaa, W. Versantvoort, A. Arshad, H.J.M. Op den Camp,

M.S.M. Jetten, C. Lüke, and J. Reimann. 2016b. Nitrate- and nitrite-dependent

anaerobic oxidation of methane. Environ. Microbiol. Rep. 8(6): 941–955. doi:

10.1111/1758-2229.12487.

Wen, X., V. Unger, G. Jurasinski, F. Koebsch, F. Horn, and G. Rehder. 2018a. Predominance of

methanogens over methanotrophs contributes to high methane emissions in rewetted

fens. (April): 1–36.

Wen, X., V. Unger, G. Jurasinski, F. Koebsch, F. Horn, G. Rehder, T. Sachs, D. Zak, G. Lischeid,

K.H. Knorr, M.E. Böttcher, M. Winkel, P.L.E. Bodelier, and S. Liebner. 2018b.

Predominance of methanogens over methanotrophs in rewetted fens characterized by

high methane emissions. Biogeosciences 15(21): 6519–6536. doi: 10.5194/bg-15-6519-

2018.

Zalman, C.A., N. Meade, J. Chanton, J.E. Kostka, S.D. Bridgham, and J.K. Keller. 2017.

Methylotrophic methanogenesis in Sphagnum-dominated peatland soils. Soil Biol.

Biochem. 118(July 2017): 156–160. doi: 10.1016/j.soilbio.2017.11.025.

Ziebarth, M., U. Lenschow, and T. Permien. 2009. Konzept zum Schutz und zur Nutzung der

Moore - Fortschreibung des Konzeptes zur Bestandssicherung und zur Entwicklung der

Moore. : 109.