**Transplacental innate immune training via maternal microbial exposure: the XBP1-ERN1**

**axis in programming dendritic cell precursors**

**Authors:** Kyle T. Mincham[1], Anya C. Jones[1], Marie Bodinier[2], Naomi M. Scott[1], Jean-Francois

Lauzon-Joset[1,3], Philip A. Stumbles[1,4], Anthony Bosco[1], Patrick G. Holt[1,5†], Deborah H.

Strickland[1†*]

**Affiliations:**

[1]Telethon Kids Institute, University of Western Australia, Nedlands, Western Australia,

Australia.

[2]INRA Pays de la Loire, UR 1268 Biopolymers Interactions Assemblies (BIA) Nantes, Nantes,

France.

[3]Centre de recherche de l'Institut de Cardiologie et de Pneumologie de Québec, Université

Laval, Québec, Canada.

[4]College of Science, Health, Engineering and Education, Murdoch University, Perth, Western

Australia, Australia.

[5]Child Health Research Centre, The University of Queensland, Brisbane, Australia.

[†]Joint senior authors.

[*]Corresponding author. Email: Deb.Strickland@telethonkids.org.au Ph: +61 8 6319 1528

Address: Northern Entrance, Perth Children's Hospital, 15 Hospital Avenue, Nedlands,

Western Australia, 6009.

**Conflict of interest statement:** The authors have declared that no conflict of interest exists.

**25** **Abstract**

**26** We recently reported that the offspring of mice treated during pregnancy with the microbial-

**27** derived immunomodulator OM-85 manifest striking resistance postnatally to allergic airways

**28** inflammation, and localised the potential treatment target to the fetal cDC progenitor

**29** compartment which expands to increase the pool of precursors available at birth, enabling

**30** accelerated postnatal seeding of the lung mucosal cDC network required for establishment

**31** of immunological homeostasis in the airways. Here, we profile maternal OM-85 treatment-

**32** associated transcriptomic signatures in fetal bone marrow, and identify a series of

**33** immunometabolic pathways which provide essential metabolites for accelerated

**34** myelopoiesis, that are hallmarks of classical "immune training". In addition, the cDC

**35** progenitor compartment displayed treatment-associated activation of the XBP1-ERN1

**36** signalling axis which has previously been shown to be essential for tissue survival of cDC,

**37** particularly within the lung microenvironment. Our forerunner studies indicate uniquely rapid

**38** turnover of airway mucosal cDCs at baseline, with further large-scale upregulation of

**39** population dynamics during aeroallergen and/or pathogen challenge. XBP1-ERN1 signalling

**40** plays a key role in mitigation of ER stress-associated toxicity which frequently accompanies

**41** DC hyper-activation during intense immunoinflammatory responses, and we suggest that

**42** enhanced capacity for XBP1-ERN1-dependent cDC survival within the airway mucosal tissue

**43** microenvironment may be a crucial element of the OM-85-mediated transplacental "innate

**44** immune training" process which results in enhanced resistance to airway inflammatory

**45** disease during the high-risk early postnatal period.

**46**

**47**

**48**

**Introduction**

The neonatal period represents a time of high risk for infection-related morbidity/mortality resulting from the combined effects of maturational deficiencies in both anti-microbial defense mechanisms that mediate pathogen recognition and elimination, and in the accompanying regulatory mechanisms required for calibration of these responses to minimise inflammatory damage to host tissues[1, 2]. With respect specifically to the lung tissue microenvironment, a crucial factor governing the kinetics of postnatal acquisition of immune competence is the rate of development of the airway mucosal cDC network which controls local immune surveillance[3, 4].

In addition to increased susceptibility to infectious diseases, the seeds for development of a range of non-communicable diseases exemplified by asthma and aero-allergies are also frequently sewn during this early postnatal window period[5], suggesting that maturational deficiencies in immune function(s) may also be risk factors in this context. In this regard, maternal immune perturbations have been acknowledged to significantly influence fetal immune development, but the underlying mechanisms remain poorly characterised[6]. Emerging epidemiological data from studies on traditional farming families in Europe and USA suggesting that benign environmental microbial exposures of mothers during pregnancy can promote prenatal immune maturation within their offspring, leading to reduced susceptibility to postnatal development of respiratory inflammatory diseases[7, 8], have stimulated wide-spread interest in this issue. This capacity for microbial exposures to modulate immune development is consistent with the paradigm of "immune training", whereby exposure to certain classes of microbial stimuli can alter the long-term functional state of innate immune cells, occurring at the progenitor level in the bone marrow (BM)[9-11], leading to optimised

peripheral immune responsiveness to other unrelated microorganisms[12]. With this in mind,

there is growing interest in the concept that immune training can be therapeutically

harnessed[13], particularly during prenatal development, to enhance immunocompetence

within the offspring[14].

We recently reported that oral treatment of pregnant mice with the microbial-derived

immunomodulatory agent OM-85 reduces susceptibility of their offspring to the development

of Th2-driven allergic airways inflammation, and identified myeloid progenitors in the

offspring BM (which supply precursor DC to eventually populate mucosal DC networks) as a

major target for maternal treatment effects[15]. In the study presented here, we employed

transcriptomic profiling to characterise gene networks activated in fetal BM (fBM) as a result

of maternal OM-85 treatment, and identify the principal treatment targets as

immunometabolic pathways supplying cellular cholesterol essential for rapid expansion of

myeloid precursor compartments, and which have previously been recognised as hallmarks

of classical immune training-associated gene signatures. We additionally identify a specific

pathway in the cDC precursor compartment which has previously been associated with

survival-under-stress, especially within the lung mucosal microenvironment.

**Results**

To elucidate the mechanisms-of-action of maternal OM-85 treatment, we examined the *in*

*utero* fetal response at gestation day 18.5 (Figure 1A), 2 days prior to expected natural term

delivery.

***Maternal OM-85 treatment selectively accelerates functional maturation of cDCs in fetal bone marrow***

Phenotypic analysis of the fBM myeloid compartment using multicolour flow cytometry revealed significant expansion of the total dendritic cell (DC) pool in fetuses from OM-85 treated mothers as compared to equivalent fetal samples from untreated mothers (Figure 1B). Further characterisation of the BM DC response demonstrated that this increase was restricted to the conventional DC (cDC) subset as previously described[15], with no parallel changes observed in plasmacytoid DC (pDC) (Figure 1B). These cDC-specific changes in fBM mirror our recent observations of increased cDC yields from BM cultures and peripheral lung from the offspring of OM-85-treated mothers in the early postnatal period[15]. We next turned our attention to fetal DC maturation state as determined by surface I-A/I-E expression. As shown in Figure 1C, maternal OM-85 treatment for the last half of gestation enhanced I-A/I-E expression on cDC in fBM when compared to fBM cDC from untreated mothers. Collectively, these observations suggest that transplacental signals generated at the feto-maternal interface following OM-85 treatment during pregnancy can "train" the developing fetal immune system via accelerating the development of functional competence within the fBM cDC compartment.

***Expansion of fetal bone marrow myeloid progenitor subsets following maternal OM-85 treatment***

Previous studies from our laboratory have additionally identified postnatal expansion of BM myeloid progenitor (MP) cell populations as an effect of maternal treatment with OM-85 during pregnancy[15]. These findings mirror that of recent studies which identified modulation of BM MP as an important component of conventional immune training mediated by both β-

glucan-[10] and Bacillus Calmette-Guérin (BCG)[11]. Based on these findings, we hypothesised

that the enhanced cDC population within fBM following maternal OM-85 treatment would

also be accompanied by concomitant upregulation of upstream MP subsets. Flow cytometric

analysis of fBM (Figure 2A) Lin⁻IL-7Rα⁻c-Kit⁺Sca-1⁻ MP (Figure 2B), Lin⁻IL-7Rα⁻c-Kit⁺Sca-1⁻

CD16/32$^{hi}$CD34$^+$ granulocyte-macrophage progenitor (GMP; Figure 2C) and Lin⁻IL-7Rα⁻c-

Kit⁺Sca-1⁻CD16/32$^{hi}$CD34$^+$CX3CR1$^+$Flt-3$^+$ macrophage-dendritic cell progenitor (MDP; Figure

2D) populations demonstrated a total increase in all 3 progenitor subsets within the BM

compartment following maternal OM-85 treatment, when compared to fBM from untreated

mothers. Consistent with the findings in Figure 1B and C, these data provide further evidence

that maternal OM-85 treatment selectively modulates the offspring BM myeloid lineage *in*

*utero*, beginning at the early-stage myeloid progenitor level through to the terminal cDC

populations which are responsible for seeding peripheral tissues during early postnatal life to

provide local DC-mediated immune surveillance.

***Maternal OM-85 treatment activates key regulators of the UPR pathway in fetal bone***

***marrow***

To gain further insight into the molecular mechanisms underpinning the maternal OM-85-

treatment effects, we employed transcriptomic profiling of fBM cells. Comparison of the

transcriptomic profiles in the treated versus untreated groups indicated that maternal OM-

85 treatment resulted in 152 differentially expressed genes (DEG) in fBM (119 upregulated,

33 downregulated; Figure 3A, Supplementary Table 1). We then interrogated the DEG for

enrichment of biological pathways employing *InnateDB*[16], focusing on the upregulated DEG

response given the limited number of downregulated DEG identified. In fBM, upregulated

DEG were enriched for genes involved in multiple aspects of protein metabolism, the

endoplasmic reticulum (ER) stress response, the unfolded protein response (UPR), cholesterol

biosynthesis and lipid metabolism (Supplementary Table 2).

Upstream regulator analysis was then performed to identify putative molecular drivers of all

observed DEG. The data revealed X-box binding protein 1 (XBP1), a transcription factor central

to the UPR[17] and crucial in the development, survival and function of multiple cell types

including plasma cells[18], eosinophils[19], natural killer (NK) cells[20], T-cell subsets[21] and DCs[22, 23],

as the most strongly activated molecular driver within fBM associated with maternal OM-85

treatment effects ($P$-value = $3.81 \times 10^{-17}$, Z-score = 4.427, Figure 3B; Supplementary Table 3),

and consistent with this, its downstream target, the canonical UPR sensor Activating

Transcription Factor 6 beta (ATF6$\beta$)[24, 25] was upregulated (Supplementary Table 1 and 3).

Additionally, Endoplasmic Reticulum To Nucleus Signalling 1 (ERN1) was identified as an

activated driver gene within the fBM following maternal OM-85 treatment ($P$-value = $3.32 \times 10^{-}$

$^{6}$, Z-score = 2.156; Figure 3B; Supplementary Table 3). Identification of ERN1 is crucial given

that during the ER stress response, this gene encodes the ER stress sensor protein inositol-

requiring enzyme 1 (IRE1$\alpha$), responsible for the unconventional cleavage of a 26 nucleotide

fragment from *Xbp1* mRNA, resulting in the generation of the active spliced variant of XBP1

(XBP1s) and enabling it to function as a potent transcription factor within the UPR signalling

pathway[25]. Collectively, these findings suggest that activation of the UPR pathway may be a

central component of the immune training mechanism induced in fBM as a result of maternal

OM-85 treatment. In addition to UPR pathway drivers, maternal OM-85 treatment also

resulted in the upstream activation of multiple drivers central to immunometabolic pathways

involved in cellular cholesterol homeostasis, including Sterol Regulatory Element Binding

Transcription Factor 1 (SREBF1; $P$-value = $1.95 \times 10^{-6}$, Z-score = 3.056) and 2 (SREBF2; $P$-value

= $2.95 \times 10^{-9}$, Z-score = 2.745) and Sterol Regulatory Element Binding Protein Cleavage-

Activating Protein (SCAP; *P*-value = $1.52 \times 10^{-10}$, Z-score = 2.949; Figure 3B; Supplementary

Table 3).  This was associated with upregulation of their downstream target Low-Density

Lipoprotein Receptor (LDLR; Supplementary Table 1 and 3), while Insulin Inducible Gene 1

(INSIG1) was inhibited in fBM following maternal OM-85 treatment (*P*-value = $5.81 \times 10^{-8}$, Z-

score = -2.931; Figure 3B; Supplementary Table 3).

Additional candidate drivers in fBM following maternal OM-85 treatment included CD38, IL5,

and an array of microRNAs (miR) (Figure 3B; Supplementary Table 3) recognised principally in

the context of cancer-associated functions[26-28]. Of note, miR-149-3p (strongly downregulated

in Figure 3B) has been shown to negatively regulate Toll-like receptor (TLR) 4 expression in

murine monocytic cells *in vitro*[29] and it is possible that other miRs may have similar (but as

yet undefined) innate immune regulatory functions[30]. The relevance of finding TLR4

upregulation in BM-derived myeloid cells in this model merits further investigation. Likewise,

the identification of the T-cell activation-associated markers CD38 and IL-5 suggests possible

contributions from activated T-cells, and these possibilities will also be addressed in follow up

studies.

***Upregulation of XBP1s expression is restricted to fetal bone marrow cDC precursors***

***following maternal OM-85 treatment***

Finally, to obtain evidence confirming that the activated form of XBP1 was upregulated, we

measured expression of the active spliced variant of XBP1 (XBP1s) at the protein level within

fBM. Using multicolour flow cytometry, we identified significant upregulation of fBM

CD11b$^+$CD11c$^+$ pre-cDCs expressing intracellular XBP1s following maternal OM-85 treatment

(Figure 4A, B). While intracellular XBP1s expression was additionally localised within fetal

CD19[+]B220[+] B cells, NKp46[+]CD11b[+]B220[+]CD11c[lo] NK cells and CD3[+] T-cells (Supplementary

Figure 1), as previously described in the literature[18, 20, 21, 31], maternal OM-85 treatment had

no impact on XBP1s expression levels in these cell types.

**Discussion**

In this study, we have characterised the response of the fBM myeloid cell compartment to

maternal treatment with the microbial-derived innate immune modulator OM-85 during

pregnancy. We demonstrate that, consistent with our previous reports in 6-week-old

offspring[15], maternal OM-85 treatment expanded the baseline pool of GMP and MDP in fBM.

Furthermore, we show for the first time the maternal OM-85-induced modulation of

metabolic pathways within fBM responsible for cellular cholesterol homeostasis, with specific

activation of SREBF1, SREBF2, SCAP and LDLR. In this regard, SREBF1 and SREBF2 are

responsible for encoding sterol regulatory element binding proteins (SREBP) which play a

central role in cellular metabolism by controlling the synthesis of cholesterol and other

membrane lipids in the Golgi of mammalian cells, with SREBP2 (encoded by SREBF2) regarded

as the "master regulator" of cellular cholesterol biosynthesis[32]. Furthermore, SCAP is central

to this process by acting as a protein chaperone, mediating the transport of SREBP from the

ER to the Golgi to promote cholesterol biosynthesis[32]. In parallel with this metabolic pathway,

LDLR regulates cellular uptake of low-density lipoprotein (LDL), the upregulation of which

increases downstream cholesterol accumulation within the cell and is required for the

proliferation of hematopoietic stem and progenitor cells[33]. It is now recognised that

myelopoiesis within the BM is heavily reliant upon an increased demand for cellular

cholesterol and enhanced cholesterol biosynthesis[34-36], and we therefore postulate that

upregulation of this immunometabolic pathway following maternal OM-85 treatment is in-

part responsible for the expansion of MP, GMP and MDP observed within fBM. Further

reinforcing the importance of enhanced cholesterol biosynthesis, INSIG1, an ER membrane

protein that prevents trafficking of SCAP/SREBP complexes to the Golgi and thereby restrains

cholesterol biosynthesis[37], was inhibited in fBM following maternal OM-85 treatment.

Together, these findings parallel previous studies demonstrating that enhanced cellular

cholesterol biosynthesis and resultant expansion of BM MP is a hallmark of classical $\beta$-glucan-

mediated central immune training[10]. We further demonstrate that transplacental

mechanisms promoting maternal OM-85-induced immune training within fBM involve a

dynamic process comprising upregulation of immunometabolic pathways that provide key

rate-limiting metabolites required for myelopoiesis and subsequent expansion of MP, GMP

and MDP, and concomitant inhibition of negative feedback loops responsible for arresting

cholesterol biosynthesis.

Downstream of the fBM progenitor response, maternal OM-85 treatment selectively

amplified the overall abundance of fBM cDC, along with enhancing the concomitant

functional maturation of these cDC as demonstrated by upregulated I-A/I-E (MHC Class II)

expression. This BM population is the source of the precursors which subsequently seed the

airway mucosal DC network that progressively develops between birth and weaning[3, 4]. It is

noteworthy that the cDC which initially seed this network postnatally are MHC-II[low] (reflecting

their functionally immature status) relative to the high-level expression seen at later ages[3],

and the findings above in fetal cDC from the treated group may collectively explain the

accelerated postnatal establishment and the enhanced functional maturation of this network

observed in their offspring[15]. This DC network plays an essential "gatekeeper" role in immune

surveillance of airway surfaces, and hence in protection against both allergic and infectious

diseases in the respiratory tract[38, 39], and its relative paucity and reduced functionality during

infancy may be an important contributor to increased susceptibility to these diseases during

this life phase.

Earlier studies from our group also identified unique features of the population dynamics of

this lung cDC network which distinguishes it from comparable populations in other tissues,

notably the exceptionally rapid baseline turnover rate of individual cells within the network,

~85% of which are replaced every 24-36 hours[40], with emigration to draining lymph nodes

(bearing samples of locally acquired antigens) balanced by recruitment of replacements from

BM. Moreover, once development of functional competence is complete[4], this network

develops capability for rapid expansion to up to 5-fold baseline density in the face of acute

challenge with airborne pro-inflammatory irritant, allergenic or microbial stimuli[41], the latter

response exhibiting kinetics that rival neutrophils[42, 43]. These unique population dynamics

suggests that even at baseline, mechanisms that promote lung cDC survival are likely to play

a crucial role in the capacity of the network to perform its immune surveillance functions

which require onward migration to downstream lymph nodes and subsequent interaction

with T-cells as opposed to antigen presentation *in situ*[38]; moreover during prolonged/severe

events exemplified by severe viral infections, the added effects of cDC injury resulting from

inflammation-associated ER stress[44] would place further pressure on survival times.

In this regard, our transcriptomic analyses of fBM from offspring of OM-85-treated mothers

also identified upregulated expression of XBP1, ATF6$\beta$ and ERN1 (encoding IRE1$\alpha$), key

components of the XBP1-ERN1 signalling axis and critical regulators of the UPR pathway[45]

which mitigates the effects of ER stress, and moreover we localised upregulated production

of active XBP1s protein to cDC precursors. Taken together with recent findings on the role for

IRE1$\alpha$-XBP1 signalling and the downstream transcription factor XBP1s in DC development and

function[22, 23], these results suggest a central role for the XBP1-ERN1 signalling axis in this OM-

85-mediated immune training process.

In further support of this suggestion, other studies demonstrate a significant reduction in

CD11c$^+$ cells (mirroring that of our pre-cDC phenotype) within XBP1$^{-/-}$ BM cultures, whilst

forced overexpression of XBP1s in XBP1$^{-/-}$ DC precursors conversely rescues and subsequently

drives expansion of the DC pool *in vitro*[22]. It is also pertinent to note that others have reported

that the cDC population in the lung mucosa is differentially reliant upon XBP1 expression for

survival at baseline relative to cDC from other tissue sites[44], which may be a direct reflection

of the uniquely high turnover rates of cDC within the airway mucosal microenvironment[40].

Collectively, the data presented here indicate that OM-85 likely operates as an immune

training agent, employing cellular and immunometabolic mechanisms previously reported in

independent model systems[10, 11, 46], with the additional capacity to act transplacentally via the

fBM. Furthermore, we go beyond the currently known features of innate immune training to

describe involvement of the XBP1-ERN1 signalling axis. Moreover, classical $\beta$-glucan- and

BCG-mediated immune training has traditionally focused on prototypic innate effector cell

populations (monocytes/macrophages/natural killer cells) resulting in enhanced resistance to

bystander pathogens via the upregulation of pro-inflammatory responses exemplified by

tumour necrosis factor-$\alpha$ (TNF-$\alpha$), interferon-$\gamma$ (IFN-$\gamma$), IL-6 and IL-1$\beta$ production[47-49] and

emergency granulopoiesis-mediated neutrophilic influx[46]. However, studies are now

beginning to recognise that immune training can also occur in DC populations, resulting *inter*

*alia* in epigenetic reprograming of pro-inflammatory cytokine responses[50]. We have recently

extended these observations to include OM-85-mediated training effects on key

immunoregulatory functions in both pregnant mice and their offspring, including effects on

both cDC and pDC populations and downstream T-regulatory cells, which are collectively

associated with enhanced resistance of both mothers and offspring to the pro-inflammatory

effects of bacterial, viral and allergenic stimulation[15, 51]. Of note, similar transplacental

immune training-like effects targeting immunoregulatory mechanisms in offspring have been

reported in relation to pregnant maternal exposure to extracts from *Acinetobacter lwoffii*[52]

and *Helicobacter pylori*[53], suggesting that the phenomenon reported here may be

generalisable, and may point towards a novel approach to mitigation of disease risk in the age

group that is in greatest need of protection.

**Methods**

***Animals***

Specific pathogen-free BALB/c mice were purchased from the Animal Resource Centre

(Murdoch, Western Australia, Australia). All mice were housed under specific pathogen-free

conditions at the Telethon Kids Institute Bioresources Centre.

***Time-mated pregnancies***

Female BALB/c mice 8-12 weeks of age were time-mated with male BALB/c studs 8-26 weeks

of age. Male studs were housed individually with 1-2 females overnight. The detection of a

vaginal plug the following morning was designated gestation day (GD) 0.5.

***Maternal OM-85 treatment***

OM-85 (OM Pharma) is an endotoxin-low lyophilised extract containing a cocktail of TLR

ligands derived from 8 major respiratory tract bacterial pathogens (*Haemophilus influenzae,*

*Streptococcus pneumoniae, Streptococcus pyogenes, Streptococcus viridians, Klebsiella*

*pneumoniae, Klebsiella ozaenae, Staphylococcus aureus* and *Neisseria catarrhalis*)[54, 55]. Based

on previously optimised dosing concentrations[51, 56], time-mated pregnant BALB/c mice

selected at random received daily oral feeding of lyophilised OM-85 reconstituted in

phosphate-buffered saline (PBS) to a concentration of 400mg/kg body weight for the second

half of gestation (GD9.5 – 17.5). Control pregnant mice were left untreated for the duration

of the study. All maternal treatment was performed with a single batch of OM-85 (batch

1812162).

***Tissue collection***

Pregnant BALB/c mice were sacrificed 24 hours after the final OM-85 dose at GD18.5. Both

horns of the uterus were removed and fetuses sacrificed by decapitation. Fetal hind legs

(cleaned of excess tissue) were removed. Fetal hind leg samples for flow cytometry were

collected into cold PBS + 0.1% bovine serum albumin (BSA) and stored on ice. Fetal hind leg

samples for transcriptomic analysis were collected into RNAlater® stabilisation solution

(Ambion). Samples collected into RNAlater® were stored overnight at 4°C, then transferred

to 1.5ml Eppendorf tubes (Eppendorf) and frozen at -80°C for future transcriptome profiling.

Dead fetuses were excluded from the study.

*Single-cell suspension preparation*

Fetal hind legs were prepared by mincing with a scalpel followed by enzymatic digestion, as

previously detailed[15]. Briefly, minced tissue was resuspended in 10ml GKN + 10% fetal calf

serum (FCS; Serana) with collagenase IV (Worthington Biochemical Corp.) and DNase (Sigma-

Aldrich) at 37°C under gentle agitation for 60 minutes. Digested cells were filtered through

sterile cotton wool columns, centrifuged and resuspended in cold PBS for total cell counts.

*Flow cytometry*

Fetal bone marrow single-cell suspensions were used for all immunostaining. Panels of

monoclonal antibodies were developed to enable phenotypic characterisation of committed

myeloid cells: CD3-FITC (BD Bioscience; clone 17A2), CD11b-BV510 (BD Bioscience; clone

M1/70), CD11c-BV711 (BD Bioscience; clone HL3), CD19-APC-H7 (BD Bioscience; clone 1D3),

Gr-1-Biotin (BD Bioscience; clone RB6-6B2), CD45R/B220-PerCP-Cy5.5 (BD Bioscience; clone

RA3-6B2) NKp46-PE-Cy7 (BioLegend; clone 29A1.4), SIRP-$\alpha$-APC (BioLegend; clone P84), I-A/I-

E-BV421 (BioLegend; clone M5/114.15.2), F4/80-BV785 (BioLegend; clone BM8), Viability-

AF700 (BD Bioscience), Streptavidin-BV605 (BD Bioscience); hematopoietic stem and

progenitor cells: CD2-Biotin (BD Bioscience; clone RM2-5), CD3-Biotin (BD Bioscience; clone

145-2C11), CD4-Biotin (BD Bioscience; clone GK1.5), CD5-Biotin (BD Bioscience; clone 53-7.3),

CD8$\alpha$-Biotin (BD Bioscience; clone 53-6.7), CD19-Biotin (BD Bioscience; clone 1D3),

CD45R/B220-Biotin (BD Bioscience; RA3-6B2), Gr-1-Biotin (BD Bioscience; clone RB6-6B2),

Ter119-Biotin (BD Bioscience; clone TER-119), CD16/32-PerCP-Cy5.5 (BD Bioscience; clone

2.4G2), CD34-FITC (BD Bioscience; clone RAM34), IL-7R$\alpha$-PE-Cy7 (BD Bioscience; clone

SB/199), Flt-3-PE (BD Bioscience; clone A2F10.1), c-Kit-APC-Cy7 (BD Bioscience; clone 2B8),

Sca-1-BV510 (BD Bioscience; clone D7), CX3CR1-APC (BioLegend; clone SA011F11), NKG2D-

BV711 (BD Bioscience; CX5), Viability-AF700 (BD Bioscience), Streptavidin-BV605 (BD

Bioscience) and XBP1s-expressing bone marrow cells: CD3-FITC (BD Bioscience; clone 17A2),

CD11b-BV510 (BD Bioscience; clone M1/70), CD11c-AF700 (BD Bioscience; clone HL3), CD19-

APC-H7 (BD Bioscience; clone 1D3), I-A/I-E-AF647 (BD Bioscience; clone M5/114.15.2),

CD45R/B220-PE-CF594 (BD Bioscience; RA3-6B2), Gr-1-Biotin (BD Bioscience; clone RB6-6B2),

NKp46-PE-Cy7 (BioLegend; clone 29A1.4), F4/80-BV785 (BioLegend; clone BM8), XBP1s-

BV421 (BD Bioscience; clone Q3-695), Streptavidin-BV605 (BD Bioscience). Intracellular

staining for XBP1s was performed using an intracellular staining buffer kit (eBioscience). Data

acquisition was performed on a 4-laser LSR Fortessa (BD Bioscience). All samples were kept

as individuals and not pooled. Immune cell phenotypic characterisation was performed using

FlowJo software (version 10.1, Tree Star). Fluorescence minus one staining controls were used

for all panels.

***Flow cytometric statistical analyses***

Statistical analysis and graphing was performed using GraphPad Prism (GraphPad software;

version 7.0a). Statistical significance of $p < 0.05$ was considered significant. Unpaired, two-

tailed Student's *t*-test or Mann Whitney *U* test were used based on distribution of the data as

determined by D'Agostino-Pearson omnibus normality test. Part of the cDC flow cytometry

data presented in Figure 1 and MDP data presented in Figure 2 has been published in a

forerunner manuscript[15].

***Transcriptome profiling (RNA-Seq)***

*Tissue preparation, RNA extraction and transcriptome profiling.* Fetal bone marrow samples

were homogenised using a rotor-star homogeniser (Qiagen) and total RNA extracted via

TRIzol (Ambion), followed by clean-up using RNeasy MinElute (Qiagen). RNA integrity was

determined using an Agilent 2100 Bioanalyzer (Agilent Technologies; RIN: $10 \pm 0$ (mean $\pm$ SD)).

1μg total RNA (*n* = 32) was shipped on dry ice to the Australia Genome Research Facility

(AGRF) for library preparation (TruSeq Stranded mRNA Library Prep Kit, Illumina) and

sequencing (Illumina HiSeq2500, 50-bp singe-end reads, v4 chemistry). The raw sequencing

data are available from GEO (accession; GSE140143).

***RNA-Seq data analysis***

*Pre-processing and exploratory data analysis:* RNA-Seq data was analysed in the R

environment for statistical computing. Sequencing data quality control (QC) was performed

with the Bioconductor package Rqc[57]. Sequencing reads were aligned to the reference murine

genome (mm10) using Subread and summarised at the gene level using featureCounts[58].

Genes with <500 total counts across the data were removed from the analysis. Sample QC

was performed by analysing the distribution of the raw read counts to check for sample

outliers using boxplots, relative log-transformed expression (RLE) plots and principal

component analysis (PCA), before and after global-scale median normalisation. *Differential*

*expression analysis:* Differentially expressed genes (DEG) were identified employing the

*DESeq2* package[59]. *DESeq2* utilises a negative binomial distribution model, with a False

Discovery Rate adjusted *P*-value for multiple comparisons. Genes were deemed significant

with an adjusted *P*-value < 0.1. *Pathways analysis:* Pathways enrichment analysis was

performed using the InnateDB database[16] with Benjamini & Hochberg adjusted *P*-value $\leq 0.05$

deemed significant. *Upstream regulator analysis:* Ingenuity Systems Upstream Regulator

Analysis[60] was employed to identify putative molecular drives of the DEG patterns.

Significance was determined by activation Z-score $\geq 2$ and *P*-value of overlap $\leq 0.05$.

*Study approval*

All animal experiments were formally approved by the Telethon Kids Institute Animal Ethics

Committee, operating under the guidelines developed by the National Health and Medical

Research Council of Australia for the care and use of animals in scientific research.

**Acknowledgements**

The authors wish to acknowledge the animal technicians at the Telethon Kids Institute

Bioresources Centre. This study was funded by the National Health and Medical Research

Council of Australia. OM Pharma provided the OM-85 at no cost.

**Author contributions**

KTM, PGH and DHS designed the study. KTM, MB, NMS and JFLJ performed the experiments.

KTM, ACJ, MB and DHS analysed the data. PAS and AB contributed to the project design and

discussions on data interpretation. KTM, PAS, PGH and DHS wrote the manuscript. All authors

reviewed the final version of the manuscript.

**Figure 1. Maternal OM-85 treatment during pregnancy selectively modulates the fetal bone marrow conventional DC subset. (A)** Kinetics of maternal OM-85 treatment beginning at gestation day (GD) 9.5 with daily oral treatment until GD 17.5 and autopsy 24 hours post final treatment at GD 18.5. **(B)** Absolute numbers of dendritic cells (DC), CD11b$^+$B220$^-$CD11c$^+$Gr-1$^-$SIRP-$\alpha^+$I-A/I-E$^+$ conventional dendritic cells (cDC) and CD11b$^-$B220$^+$CD11c$^+$Gr-1$^+$I-A/I-E$^+$ plasmacytoid dendritic cells (pDC) in BM of fetuses from OM-85-treated and untreated mothers. **(C)** Mean fluorescence intensity (MFI) of I-A/I-E expression on cDCs and pDCs in fBM. Data are presented from individual animals comparing fetuses from OM-85-treated and untreated mothers and displayed as bar graphs showing mean $\pm$ SEM of *n* = 8 independent experiments. Statistical significance was determined using Mann-Whitney *U* test (B) or Student's *t* test (C) based on distribution of the data as determined by D'Agostino-Pearson omnibus normality test.

**Figure 2. Treatment of mothers with OM-85 during pregnancy boosts myeloid progenitor subsets within fetal bone marrow. (A)** Representative flow cytometry plots for the identification of Lin⁻IL-7Rαc-Kit⁺Sca-1⁻ myeloid progenitors (MP), Lin⁻IL-7Rαc-Kit⁺Sca-1⁻CD16/32ʰⁱCD34⁺ granulocyte-macrophage progenitors (GMP) and Lin⁻IL-7Rαc-Kit⁺Sca-1⁻CD16/32ʰⁱCD34⁺CX3CR1⁺Flt-3⁺ macrophage-dendritic cell progenitors (MDP) within fBM. Absolute numbers of **(B)** MP, **(C)** GMP and **(D)** MDP in BM of fetuses from OM-85-treated and

445 untreated mothers. Data are presented from individual animals comparing fetuses from OM-

446 85-treated and untreated mothers and displayed as bar graphs showing mean $\pm$ SEM of $n = 8$

447 independent experiments. Statistical significance was determined using Student's $t$ test (C) or

448 Mann-Whitney $U$ test (B, D) based on distribution of the data as determined by D'Agostino-

449 Pearson omnibus normality test.

**Figure 3. Maternal OM-85-induced changes in fetal bone marrow gene expression profiles.**

**(A)** Differentially expressed genes (DEG) within fBM comparing offspring from OM-85-treated and untreated mothers. DEG are summarised as a volcano plot showing data along axes of statistical significance (-$\log_{10}$ adjusted *P*-value) and differential expression magnitude ($\log_2$ fold change) for *n* = 16 individual animals per group collected from *n* = 7 independent experiments. Dashed horizontal lines indicate a False Discovery Rate (FDR) adjusted *P*-value < 0.1. Genes shown in red were upregulated and those shown in blue were downregulated. **(B)** activated (red) and inhibited (blue) molecular drivers of the differential expression patterns were identified using Upstream Regulator Analysis.

**Figure 4. XBP1s expression in pre-cDCs within fetal bone marrow. (A)** Representative flow

cytometry plots demonstrating intracellular XBP1s expression in CD11b$^+$B220$^-$CD11c$^+$Gr-1$^-$I-

A/I-E$^-$XBP1s$^+$ pre-conventional dendritic cells (pre-cDCs) within fBM. **(B)** Absolute numbers of

XBP1s$^+$ pre-cDCs in BM of fetuses from OM-85-treated and untreated mothers. Data are

presented from individual animals comparing fetuses from OM-85-treated and untreated

mothers and displayed as bar graphs showing mean $\pm$ SEM of $n$ = 4 independent experiments.

Statistical significance was determined using Mann-Whitney $U$ test.

**References**

1.    Levy O. Innate immunity of the newborn: basic mechanisms and clinical correlates.

Nat Rev Immunol. 2007;7(5):379-90.

2.    Adkins B, Leclerc C, Marshall-Clarke S. Neonatal adaptive immunity comes of age. Nat

Rev Immunol. 2004;4(7):553-64.

3.    Nelson DJ, McMenamin C, McWilliam AS, Brenan M, Holt PG. Development of the

airway intraepithelial dendritic cell network in the rat from class II major

histocompatibility (Ia)-negative precursors: differential regulation of Ia expression at

different levels of the respiratory tract. J Exp Med. 1994;179(1):203-12.

4.    Nelson DJ, Holt PG. Defective regional immunity in the respiratory tract of neonates is

attributable to hyporesponsiveness of local dendritic cells to activation signals. J

Immunol. 1995;155(7):3517-24.

5.    Gollwitzer ES, Marsland BJ. Impact of Early-Life Exposures on Immune Maturation and

Susceptibility to Disease. Trends Immunol. 2015;36(11):684-96.

6.    Apostol AC, Jensen KDC, Beaudin AE. Training the Fetal Immune System Through

Maternal Inflammation-A Layered Hygiene Hypothesis. Front Immunol. 2020;11:123-

36.

7.    von Mutius E, Vercelli D. Farm living: effects on childhood asthma and allergy. Nat Rev

Immunol. 2010;10(12):861-8.

8.    Ober C, Sperling AI, von Mutius E, Vercelli D. Immune development and environment:

lessons from Amish and Hutterite children. Curr Opin Immunol. 2017;48:51-60.

9.    Kleinnijenhuis J, et al. Long-lasting effects of BCG vaccination on both heterologous

Th1/Th17 responses and innate trained immunity. J Innate Immun. 2014;6(2):152-8.

10.    Mitroulis I, et al. Modulation of Myelopoiesis Progenitors Is an Integral Component of

Trained Immunity. Cell. 2018;172(1-2):147-61.

11.    Kaufmann E, et al. BCG Educates Hematopoietic Stem Cells to Generate Protective

Innate Immunity against Tuberculosis. Cell. 2018;172(1):176-90.e19.

12.    Netea MG, et al. Defining trained immunity and its role in health and disease. Nat Rev

Immunol. 2020.

13.    Mulder WJM, Ochando J, Joosten LAB, Fayad ZA, Netea MG. Therapeutic targeting of

trained immunity. Nat Rev Drug Discov. 2019;18(7):553-66.

14.    Holt PG, Strickland DH, Custovic A. Targeting maternal immune function during

pregnancy for asthma prevention in offspring: harnessing the "farm effect"? J Allergy

Clin Immunol. 2020;*In-press*.

15.    Mincham KT, et al. Transplacental immune modulation with a bacterial-derived agent

protects against allergic airway inflammation. J Clin Invest. 2018;128(11):4856-69.

16.    Breuer K, et al. InnateDB: systems biology of innate immunity and beyond-recent

updates and continuing curation. Nucleic Acids Res. 2013;41(Database issue):D1228-

33.

17.    Iwakoshi NN, et al. Plasma cell differentiation and the unfolded protein response

intersect at the transcription factor XBP-1. Nat Immunol. 2003;4(4):321-9.

18.    Reimold AM, et al. Plasma cell differentiation requires the transcription factor XBP-1.

Nature. 2001;412(6844):300-7.

19.    Bettigole SE, et al. The transcription factor XBP1 is selectively required for eosinophil

differentiation. Nat Immunol. 2015;16(8):829-37.

20.    Dong H, et al. The IRE1 endoplasmic reticulum stress sensor activates natural killer cell

immunity in part by regulating c-Myc. Nat Immunol. 2019;20(7):865-78.

21.    Pramanik J, et al. Genome-wide analyses reveal the IRE1a-XBP1 pathway promotes T

helper cell differentiation by resolving secretory stress and accelerating proliferation.

Genome Med. 2018;10(1):76.

22.    Iwakoshi NN, Pypaert M, Glimcher LH. The transcription factor XBP-1 is essential for

the development and survival of dendritic cells. J Exp Med. 2007;204(10):2267-75.

23.    Osorio F, et al. The unfolded-protein-response sensor IRE-1alpha regulates the

function of CD8alpha+ dendritic cells. Nat Immunol. 2014;15(3):248-57.

24.    Yoshida H, et al. Endoplasmic reticulum stress-induced formation of transcription

factor complex ERSF including NF-Y (CBF) and activating transcription factors 6alpha

and 6beta that activates the mammalian unfolded protein response. Mol Cell Biol.

2001;21(4):1239-48.

25.    Yoshida H, Matsui T, Yamamoto A, Okada T, Mori K. XBP1 mRNA is induced by ATF6

and spliced by IRE1 in response to ER stress to produce a highly active transcription

factor. Cell. 2001;107(7):881-91.

26.    Xia B, et al. Upregulation of miR-874-3p and miR-874-5p inhibits epithelial ovarian

cancer malignancy via SIK2. J Biochem Mol Toxicol. 2018;32(8):e22168.

27.    Stark MS, et al. The "melanoma-enriched" microRNA miR-4731-5p acts as a tumour

suppressor. Oncotarget. 2016;7(31):49677-87.

28.    Yang D, Du G, Xu A, Xi X, Li D. Expression of miR-149-3p inhibits proliferation,

migration, and invasion of bladder cancer by targeting S100A4. Am J Cancer Res.

2017;7(11):2209-19.

29.    Shen W, et al. Repression of Toll-like receptor-4 by microRNA-149-3p is associated

with smoking-related COPD. Int J Chron Obstruct Pulmon Dis. 2017;12:705-15.

30.    Kirchner B, Pfaffl MW, Dumpler J, von Mutius E, Ege MJ. microRNA in native and

543           processed cow's milk and its implication for the farm milk effect on asthma. J Allergy

544           Clin Immunol. 2016;137(6):1893-5.e13.

545    31.    Todd DJ, et al. XBP1 governs late events in plasma cell differentiation and is not

546           required for antigen-specific memory B cell development. J Exp Med.

547           2009;206(10):2151-9.

548    32.    Goldstein JL, DeBose-Boyd RA, Brown MS. Protein sensors for membrane sterols. Cell.

549           2006;124(1):35-46.

550    33.    Tolani S, et al. Hypercholesterolemia and reduced HDL-C promote hematopoietic stem

551           cell proliferation and monocytosis: studies in mice and FH children. Atherosclerosis.

552           2013;229(1):79-85.

553    34.    Yvan-Charvet L, et al. ATP-binding cassette transporters and HDL suppress

554           hematopoietic stem cell proliferation. Science (New York, NY). 2010;328(5986):1689-

555           93.

556    35.    Murphy AJ, et al. ApoE regulates hematopoietic stem cell proliferation, monocytosis,

557           and monocyte accumulation in atherosclerotic lesions in mice. J Clin Invest.

558           2011;121(10):4138-49.

559    36.    Morgan PK, Fang L, Lancaster GI, Murphy AJ. Hematopoiesis is regulated by

560           cholesterol efflux pathways and lipid rafts: connections with cardiovascular diseases.

561           J Lipid Res. 2020;61(5):667-75.

562    37.    Yang T, et al. Crucial step in cholesterol homeostasis: sterols promote binding of SCAP

563           to INSIG-1, a membrane protein that facilitates retention of SREBPs in ER. Cell.

564           2002;110(4):489-500.

38.  Holt PG, Strickland DH, Wikstrom ME, Jahnsen FL. Regulation of immunological

homeostasis in the respiratory tract. Nat Rev Immunol. 2008;8(2):142-52.

39.  Lambrecht BN, Hammad H. Lung dendritic cells in respiratory viral infection and

asthma: from protection to immunopathology. Annu Rev Immunol. 2012;30:243-70.

40.  Holt PG, Haining S, Nelson DJ, Sedgwick JD. Origin and steady-state turnover of class

II MHC-bearing dendritic cells in the epithelium of the conducting airways. J Immunol.

1994;153(1):256-61.

41.  McWilliam AS, et al. Dendritic Cells Are Recruited into the Airway Epithelium during

the Inflammatory Response to a Broad Spectrum of Stimuli. J Exp Med.

1996;184(6):2429-32.

42.  Jahnsen FL, et al. Accelerated antigen sampling and transport by airway mucosal

dendritic cells following inhalation of a bacterial stimulus. J Immunol.

2006;177(9):5861-7.

43.  McWilliam AS, Nelson D, Thomas JA, Holt PG. Rapid dendritic cell recruitment is a

hallmark of the acute inflammatory response at mucosal surfaces. J Exp Med.

1994;179(4):1331-6.

44.  Tavernier SJ, et al. Regulated IRE1-dependent mRNA decay sets the threshold for

dendritic cell survival. Nat Cell Biol. 2017;19(6):698-710.

45.  Grootjans J, Kaser A, Kaufman RJ, Blumberg RS. The unfolded protein response in

immunity and inflammation. Nat Rev Immunol. 2016;16(8):469-84.

46.  Brook B, et al. BCG vaccination–induced emergency granulopoiesis provides rapid

protection from neonatal sepsis. Sci Transl Med. 2020;12(542):eaax4517.

47.  Quintin J, et al. Candida albicans infection affords protection against reinfection via

functional reprogramming of monocytes. Cell Host Microbe. 2012;12(2):223-32.

48.    Kleinnijenhuis J, et al. Bacille Calmette-Guerin induces NOD2-dependent nonspecific

protection from reinfection via epigenetic reprogramming of monocytes. Proc Natl

Acad Sci U S A. 2012;109(43):17537-42.

49.    Sun JC, Beilke JN, Lanier LL. Adaptive Immune Features of Natural Killer Cells. Nature.

2009;457(7229):557-61.

50.    Hole CR, et al. Induction of memory-like dendritic cell responses in vivo. Nat Commun.

2019;10(1):2955.

51.    Scott NM, et al. Protection against maternal infection-associated fetal growth

restriction: proof-of-concept with a microbial-derived immunomodulator. Mucosal

Immunol. 2017;10(3):789-801.

52.    Conrad ML, et al. Maternal TLR signaling is required for prenatal asthma protection by

the nonpathogenic microbe Acinetobacter lwoffii F78. J Exp Med. 2009;206(13):2869-

77.

53.    Kyburz A, et al. Transmaternal Helicobacter pylori exposure reduces allergic airway

inflammation in offspring through regulatory T cells. J Allergy Clin Immunol.

2019;143(4):1496-512.e11.

54.    Razi CH, et al. The immunostimulant OM-85 BV prevents wheezing attacks in

preschool children. J Allergy Clin Immunol. 2010;126(4):763-9.

55.    Rozy A, Chorostowska-Wynimko J. Bacterial immunostimulants--mechanism of action

and clinical application in respiratory diseases. Pneumonol Alergol Pol.

2008;76(5):353-9.

56.    Strickland DH, et al. Boosting airway T-regulatory cells by gastrointestinal stimulation

as a strategy for asthma control. Mucosal Immunol. 2011;4(1):43-52.

57.     de Souza W, Carvalho BdS, Lopes-Cendes I. Rqc: A Bioconductor Package for Quality

613           Control of High-Throughput Sequencing Data. Journal of Statistical Software, Code

614           Snippets. 2018;87(2):1-14.

615     58.     Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable read mapping

616           by seed-and-vote. Nucleic Acids Res. 2013;41(10):e108.

617     59.     Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

618           RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.

619     60.     Krämer A, Green J, Pollard J, Jr., Tugendreich S. Causal analysis approaches in Ingenuity

620           Pathway Analysis. Bioinformatics. 2014;30(4):523-30.