The nose is mightier than the tooth: larger male proboscis monkeys have smaller canines

Ikki Matsuda[1,2,3,4*], Danica J. Stark[5], Diana A. Ramirez Saldivar[5,6], Augustine Tuuga[6], Senthilvel K. S. S. Nathan[6], Benoit Goossens[5,6,7,8], Carel P. van Schaik[9], Hiroki Koda[10*]

[1]Chubu University Academy of Emerging Sciences, 1200, Matsumoto-cho, Kasugai-shi, Aichi 487-8501, Japan

[2]Wildlife Research Center of Kyoto University, Kyoto, Japan

[3]Japan Monkey Centre, Inuyama, Japan

[4]Institute for Tropical Biology & Conservation, Universiti Malaysia, Sabah, Malaysia

[5]Danau Girang Field Centre, c/o Sabah Wildlife Department, Sabah, Malaysia

[6]Sabah Wildlife Department, Sabah, Malaysia

[7]Sustainable Places Research Institute, Cardiff University, Cardiff, UK

[8] Organisms and Environment Division, Cardiff School of Biosciences, Cardiff University, Cardiff, UK

[9]Anthropological Institute and Museum, University of Zurich, Zurich, Switzerland

[10]Primate Research Institute, Kyoto University, Inuyama, Aichi 484-8506, Japan

*Correspondence: ikki-matsuda@isc.chubu.ac.jp or koda.hiroki.7a@kyoto-u.ac.jp

The uniquely enlarged noses of male proboscis monkeys (*Nasalis larvatus*) are prominent adornments, and a sexually selected male masculine trait, as recently revealed by the significant correlations among nose, body, and testis sizes and the clear association between nose size and the number of females in a male's harem. However, till date, the analyses of other common male traits, i.e., canines, are lacking. Whereas male nose size had a positive correlation with body size, we unexpectedly found a negative correlation between body and canine sizes. Similar relationship between nose and body size was observed in females, whereas only a weak correlation was noted between canine and body size. We explain this by an interaction between sexual and natural selection. Larger noses in males may interfere with the use of canines, thereby reducing their effectiveness as weapons. However, longer canines are opposed by natural selection because they impose a larger gape upon its bearer and reduce foraging efficiency, particularly in folivores. The difference between the sexes regarding the regression intercept for body size vs. canine size was significant, although the difference of the slopes was insignificant, consistent with the hypothesis that the nose interferes with the effectiveness of canines as weapons in males but not in females. Moreover, a mathematical simulation supported our prediction. This unique case of decoupling of body and canine size reveals that large canines carry an ecological cost. It also raises interesting questions regarding the developmental mechanism that produces a negative correlation among adult body, nose size, and canine size.

Introduction

Darwinian evolutionary theory explains male-specific characteristics as a consequence of sexual selection [1]. Large canines in some mammals, including primates, represent a typical example of a weapon used in contests with conspecific male rivals [2]. By contrast, ornaments independently evolve for a function in combat, reflecting their function as a partner attraction signal (ultimately determined by female choice). However, some male-specific traits are neither weapons nor ornaments, and they serve as a badge of male social status, advertising competitive potential with respect to other males [1, 3], particularly as observed for social rank-dependent masculine traits in some primate species [4]. Proboscis monkeys (*Nasalis larvatus*) represent a typical sexually dimorphic primate with male-specific enlarged noses that are prominent adornments, which have been linked with their sexually competitive social system of one-male groups. Recently, we reported significant positive correlations among nose, body, and testis size as well as a clear association between nose size and both nasalized acoustic signals and the number of females in one-

male groups; this supports the hypothesis that larger noses audiovisually advertise a
male's fighting ability with conspecifics [5], and thus serve as status badges rather than
ornaments.

Till date, an analysis of another common sexually selected male trait, i.e., canine
size, is lacking. Sexually dimorphic male traits (e.g., body mass, nose, testis, and canines)
develop after sexual maturation, primarily in response to endocrine changes in the levels
of androgens such as testosterone. Comparisons of these male traits would enable us to
study the relationships among weapons, badges, ornaments, and testes. In this study, we
compared canine size among previously investigated wild proboscis monkeys [5]. Notably,
proboscis monkey one-male groups (and all-male groups) aggregate on a regular basis at
their sleeping sites (suggestive of a "multilevel social system" [6]), and female dispersal
among the groups is typical [7], creating the potential for both female choice and male-male
competition for females. Therefore, male canines are expected to serve as important
weapons for mate competition or defense, especially because canine size generally
reflects the strength of intra-sexual selection more directly than body size [2, 8].

Material and methods
Between July 2011 and December 2016, we captured 18 free-ranging adult male
proboscis monkeys in the Lower Kinabatangan Floodplain, Sabah, Borneo, Malaysia
(5°18′N to 5°42′N and 117°54′E to 18°33′E), as described in detail in the methods and
ethical statement of Stark *et al.*[9]. In this study, we added the data for maxillary canine
length to those on body mass and nose size (length × width) for the same males obtained
by Koda *et al.* [5] as well as data for female canine length, body mass, and nose size. The
maxillary canine was selected because of its larger size and greater importance in
behavioral displays and weaponry [10]. A linear model was used to establish whether body
mass was related to other physical properties such as nose and canine size. To obtain a
linear dimension comparable to the canine dimension, body mass and nose size were cube
root- and square root-transformed, respectively, to generate response and explanatory
variables. The variance inflation factors were 1.00 for both nose size and body mass,
indicating that collinearity among independent factors would not affect the results [11]. We
examined a set of models with all possible combinations of the explanatory variables and
ranked them according to the corrected version of the Akaike information criterion (AIC)
for small sample sizes, called AICc [12]. Following guidelines published for wildlife
research, we selected models with ΔAICc $\leq$ 2, where ΔAICc = AICc − minimum AICc
within the candidate model set [12]. Analysis of covariance served to compare the slopes
and intercepts of regression lines between the sexes. These analyses were performed using

R ver. 3.3.2 [13].

Our motivation for the model simulation was to clarify the mechanism by which
the termination time of canine development t_1 determines the final results of body mass
and canine size. We simulated the growth pattern based on several deterministic
parameters, considering the developmental evidence for proboscis monkeys. Additionally,
we tried to apply the model for both males and females within the same developmental
frameworks, considering the sex differences of the developmental parameters. More
details of the mathematical models and parameter proposals are described in the
Supplementary   Information   or   code   repositories
(https://github.com/hkoda/nasalis_canine_bodymass_model).

Results and discussion
The best-fitting model included both male nose and canine size. Whereas nose size was
positively correlated with body mass, the model with canine size was unexpectedly
negative (Fig. 1), although the fit of the model including only canine size was almost as
good (i.e., ΔAICc < 2.0: Table 1). Conversely, for females, the best-fitting model only
included nose size, which had a positive effect on body mass. The second-best model was
the null model (Table 1). The difference in the regression slopes and intercepts for body
size vs. nose size between the sexes was not significant (slope: $t = -1.41$, $P = 0.17$; $t =$
0.28; intercept: $P = 0.78$). However, the difference in the regression intercept for body
size vs. canine size was significant ($t = 8.28$, $P < 0.0001$), although the slopes were not
different ($t = -0.33$, $P = 0.74$). Previous results illustrated that secondary sexual male
traits reflect both pre- and post-copulatory competition; specifically, larger noses serve as
both "female attraction" and "badge of status," and simultaneously, larger testes reflect
sperm competition [5]. Body size also contributes to both male-male competition and
female attraction.

These results can be interpreted in two ways. The minimal interpretation is that
canine size does not contribute to the fighting ability of harem-holding males, unlike the
findings in other primate species forming harem societies [2]. Battles featuring fatal
physical attacks using canines are commonly observed in typical harem species such as
hanuman langurs (*Semnopithecus entellus*) [14] or howler monkeys (*Alouatta seniculus*) [15].
Nonetheless, the benefit of a larger body size in proboscis monkey males is clear. Indeed,
branch shaking is commonly used to keep groups apart [6], and in chases without physical
contact, males repeatedly exhibit such displays accompanied by leaps between trees (IM.
pers obs). Additionally, among proboscis monkey males, nose size, which is highly
correlated with body size, determines the formant frequencies of loud vocalizations,

which may decide contests without any direct physical competition, or at least direct

biting [5, 7, 16]. Thus, a larger body, which contributes to male athleticism, would allow more

effective displays of fighting ability, whereas nose size is the best indicator of fighting

ability because it is easier to assess via vocalization opposed to visual examination.

Consequently, males might invest their physiological resources to develop nose, body,

and testis size more than canine weapons, in line with the trade-off phenomenon recently

reported in other harem primates e.g., [17].

Contrarily, whereas females do not compete through sound as do males, they

unexpectedly exhibited the same nose size/body size correlation, suggesting that female

nose size is linked to the effect on males. Female noses are probably sufficiently small to

avoid interfering with their activity, and thus, selection toward sex-differentiated

expression does not occur.

However, the significantly negative correlation between male canine and body

size supports an alternative interpretation; namely, it suggests an ecological cost of large

canines, as suggested for the taxonomic variation in canine size observed across primate

lineages [18, 19]. The prominent nose may interfere with effective canine bites, thereby

reducing their effectiveness as weapons. Moreover, given their reduced effectiveness,

natural selection may actually favor smaller canines because larger canines increase gape,

which reduces bite force and thus mastication efficiency [18]. This foraging cost may be

especially severe in large-sized folivores, such as proboscis monkeys and gorillas [19]. Thus,

a negative correlation might evolve in the rate of post-adolescent growth of the size of

the body frame and that of canines. We therefore predict that in proboscis monkeys, body-

size dependent displays are more effective at greater distances than the facial displays

commonly observed in most other primates [20].

Contrary to the significant negative slope for the male canine size-body size

correlation, we only found a weak correlation for the female slope. Although the

difference of the slopes between the sexes was not significant, possibly because of the

limited sample size, the significant sex difference in the intercept for body size vs. canine

size suggests that canine size has a different relationship with body mass between males

and females. The difference in the sex-differentiated pattern between noses and canines

would confirm the hypothesis that large canines interfere with effective biting in males,

making them less effective as weapons and thus leading to negative selection on canine

size in the largest males.

Concerning the mechanism underlying the trade-off among body mass, nose size,

and canine size, we predict that canine growth would actually exhibit a negative

correlation with testosterone levels, opposite to the common and ancestral situation [17]. To

test this "costly canine hypothesis", we conducted preliminary mathematical simulations
considering body mass development with a growth constrained by canine size (for a
model description, see Supplementary Information). Echoing our empirical findings, the
negative relationship between body and canine size was observed only in males when we
assumed a large cost of canine development (Fig. 2). Future detailed studies focusing on
the behavior and development system of proboscis monkeys are therefore required.
172

Acknowledgements
We thank the Sabah Biodiversity Centre and the Sabah Wildlife Department for granting
permission to carry out this research, our research assistants for support in the field, and
the Japan Monkey Centre (Dr. N. Kimura) for providing the information used in our
mathematical simulations. We appreciate Prof. I. Tsuda to support verifying the
mathematical simulations. The authors would like to thank Enago for the English
language review. This study was partially funded by Japan Science and Technology
Agency Core Research for Evolutional Science and Technology 17941861
(#JPMJCR17A4), Ministry of Education, Culture, Sports, Science and Technology Grant-
in-aid for Scientific Research on Innovative Areas #4903 (Evolinguistic), 17H06380, and
Yayasan Sime Darby (to BG). This paper is dedicated to the memory of Dr. Diana Angeles
Ramirez Saldivar.

Author contributions
IM, BG, CV and HK conceptualized the initial idea, IM, DJS, DARS, SKSSN performed
the wild morphology sampling, AT BG arranged the sampling in the wild, IM, HK
performed and interpreted the statistical analyses and mathematical simulation, IM, CV,
HK drafted the manuscript. All authors contributed to the final version of the manuscript.

Competing interests
The authors declare that they have no competing interests.

Table and figure legends
Table 1 Summary of model selection. Linear models were used to investigate whether
body mass was related to other body traits such as nose and canine size (only the models
with ΔAICc ≤ 2 are shown).

Figure 1 Relationships of nose size (left) and canine size and (right) with body mass.

202 Figure 2 Relationships of normalized canine size with body mass in the mathematical
203 simulation. We observed a negative correlation between body mass and canine size for
204 both male statuses, but no correlation was noted for females, consistent with our
205 observations in free-ranging specimens (see Supplementary Information for details of the
206 simulation).
207
208

References

1. Darwin C. *The descent of man, and selection in relation to sex*. John Murray (1871).

2. Plavcan JM, van Schaik CP. Intrasexual competition and canine dimorphism in anthropoid primates. *Am J Phys Anthropol* **87**, 461-477 (1992).

3. Anderson M. *Sexual Selection*. Princeton University Press (1994).

4. Grueter CC, Isler K, Dixson BJ. Are badges of status adaptive in large complex primate groups? *Evolution and Human Behavior*， (2015).

5. Koda H*, et al.* Nasalization by *Nasalis larvatus*: Larger noses audiovisually advertise conspecifics in proboscis monkeys. *Science advances* **4**, eaaq0250 (2018).

6. Yeager CP. Proboscis monkey (*Nasalis larvatus*) social organization: Intergroup patterns of association. *Am J Primatol* **23**, 73-86 (1991).

7. Matsuda I, Tuuga A, Bernard H, Furuichi T. Inter-individual relationships in proboscis monkeys: a preliminary comparison with other non-human primates. *Primates* **53**, 13-23 (2012).

8. Plavcan JM. Sexual selection, measures of sexual selection, and sexual dimorphism in primates. In: *Sexual Selection in Primates: New and Comparative Perspectives* (eds Kappeler PM, van Schaik CP). Cambridge University Press (2004).

9. Stark DJ, Vaughan IP, Ramirez Saldivar DA, Nathan SK, Goossens B. Evaluating methods for estimating home ranges using GPS collars: A comparison using proboscis monkeys (*Nasalis larvatus*). *PLOS One* **12**, e0174891 (2017).

10. Leutenegger W, Kelly JT. Relationship of sexual dimorphism in canine size and body size to social, behavioral, and ecological correlates in anthropoid primates. *Primates* **18**, 117–136 (1977).

11. Quinn GP, Keough MJ. *Experimental design and data analysis for biologists*. Cambridge University Press (2002).

12. Burnham KP, Anderson DR. *Model selection and multimodel inference: a practical information-theoretic approach*, 2nd edn. Springer (2002).

13. R-Core-Development-Team. R: a language and environment for statistical computing.). Foundation for Statistical Computing (2019).

14. Sugiyama Y. On the social change of hanuman langurs (*Presbytis entellus*) in their natural condition. *Primates* **6**, 381-418 (1965).

15. Crockett CM, Pope T. Inferring patterns of aggression from red howler monkey injuries. *Am J Primatol* **15**, 289-308 (1988).

16. Matsuda I, Kubo T, Tuuga A, Higashi S. A Bayesian analysis of the temporal change of local density of proboscis monkeys: implications for environmental effects on a multilevel society. *Am J Phys Anthropol* **142**, 235-245 (2010).

17. Dunn JC*, et al.* Evolutionary trade-off between vocal tract and testes dimensions in howler monkeys. *Curr Biol* **25**, 2839-2844 (2015).

18. Hylander WL. Functional links between canine height and jaw gape in catarrhines with special reference to early hominins. *Am J Phys Anthropol* **150**, 247-259 (2013).

265   19.   Greenfield LO. Origin of the human canine: A new solution to an old enigma. *Am J Phys Anthropol*
266   **35**, 153-185 (1992).
267
268   20.   van Hooff JARAM. The Facial Displays of the Catarrhine Monkeys and Apes. In: *Primate*
269   *ethology* (ed Morris D). AldineTransaction (1967).
270
271

272 Table 1. Summary of model selection. Linear models were used to investigate whether
273 body mass was related to other body traits such as nose and canine size (only the models
274 with ΔAICc ≤ 2 are shown).
275

Intercept	Nose size	Canine size	df	Log-likelihood	AICc	Δ-AICc	AICc weight
Male							
42.3	0.11	-0.01	4	25.0	-39.0	0	0.41
33.0		-0.01	3	23.2	-38.7	0.34	0.34
Female							
1.387	0.2724		3	12.3	-14.6	0	0.77

278   Figure 1. Relationships of nose size (left) and canine size and (right) with body mass.

Cost parameter: 0.99

280
281 Figure 2. Relationships of normalized canine size with body mass in the mathematical
282 simulation. We observed a negative correlation between body mass and canine size for
283 both male statuses, but no correlation was noted for females, consistent with our
284 observations in free-ranging specimens (see Supplementary Information for details of the
285 simulation).