FUS gene is dual-coding with both proteins united in FUS-

mediated toxicity

Marie A. Brunet[1, 2,*], Jean-Francois Jacques[1,2], Sonya Nassari[3], Giulia E. Tyzack[4,5], Philip

McGoldrick[6], Lorne Zinman[7], Steve Jean[3], Janice Robertson[6], Rickie Patani[4,5], Xavier Roucou[1, 2,*]

[1] Department of Biochemistry and Functional Genomics, Université de Sherbrooke, Sherbrooke,

Québec, Canada;

[2] PROTEO, Quebec Network for Research on Protein Function, Structure, and Engineering;

[3] Immunology and Cell Biology Department, Université de Sherbrooke, Sherbrooke, Québec,

Canada;

[4] The Francis Crick Institute, 1 Midland Road, London, UK;

[5] Department of Neuromuscular Diseases, UCL Queen Square Institute of Neurology, Queen

Square, London, UK;

[6] Tanz Centre for Research in Neurodegenerative Diseases, University of Toronto, Toronto,

Canada;

[7] Division of Neurology, Department of Medicine, Sunnybrook Health Sciences Centre, University

of Toronto, Toronto, Canada.

*Corresponding authors:

Xavier Roucou, Department of Biochemistry and Functional Genomics, Université de Sherbrooke,

3201 Jean Mignault, Sherbrooke, Québec J1E 4K8, Canada, Tel. (819) 821-8000x72240; E-Mail:

xavier.roucou@usherbrooke.ca

Marie A. Brunet, Department of Biochemistry and Functional Genomics, Université de

Sherbrooke, 3201 Jean Mignault, Sherbrooke, Québec J1E 4K8, Canada, Tel. (819) 821-

8000x72189; E-Mail: marie.brunet@usherbrooke.ca

ABSTRACT:

Novel functional coding sequences (altORFs) are camouflaged within annotated ones (CDS) in a

different reading frame. We discovered an altORF nested in the FUS CDS encoding a conserved

169 amino acid protein, altFUS. AltFUS is endogenously expressed in human tissues, notably in

the motor cortex and motor neurons. Overexpression of wild-type FUS and/or amyotrophic lateral

sclerosis-linked FUS mutants is known to trigger toxic mechanisms in different models. These

include an inhibition of autophagy, loss of mitochondrial potential, and accumulation of

cytoplasmic aggregates. We show here that altFUS, not FUS, is responsible for the inhibition of

autophagy. AltFUS is also pivotal in the mechanisms leading to the mitochondrial potential loss

and accumulation of cytoplasmic aggregates. Suppression of altFUS expression in a *Drosophila*

model of *FUS*-related toxicity protects against neurodegeneration. Some mutations found in ALS

patients are overlooked because of their synonymous effect on the FUS protein. Yet we showed

they exert a deleterious effect via their missense consequence on the overlapping altFUS protein.

These findings demonstrate that *FUS* is a bicistronic gene and suggest that both proteins, FUS and

altFUS, cooperate in toxic mechanisms.

 MAIN

FUS is a nuclear RNA-binding protein, with a C-terminal nuclear localization signal (NLS)[1,2]. The

protein is involved in RNA processing, DNA repair and cellular proliferation, although its functions

are not precisely elucidated[1]. Mutations in *FUS* gene associate with Amyotrophic Lateral Sclerosis

(ALS), frontotemporal lobar dementia (FUS-FTLD) and essential tremor, all characterized by FUS

cytoplasmic inclusions in neurons and glial cells[1]. Such cytoplasmic FUS aggregates are

pathological features in patients with *FUS* mutations or sporadic disease.

Recently, mutations in *FUS* 3'UTR were described in ALS patients and linked to an increased level

of *FUS* mRNA and protein[3–5]. Surprisingly, over-expression of wild-type FUS provokes an

aggressive ALS phenotype in mice and fruit flies, in accordance with findings in yeast and

mammalian cells[6–10]. The mechanism of the wild-type or ALS-linked mutated FUS toxicity remains

unclear[1,11,12].

With currently non-annotated proteins being increasingly reported[13–16], we hypothesized that the

toxicity resulting from wild-type FUS over-expression may come from another, unseen, actor[16].

These novel proteins are coded by alternative open reading frames (altORFs) that are located

within "non-coding" RNAs (ncRNA), within the 5' or 3' "untranslated" regions (UTR) of mRNAs, or

overlapping a known coding sequence (CDS) within a different frame of an mRNA[15–17].

Serendipitous discoveries and ribosome profiling have recently highlighted the distribution of

altORFs throughout the human genome, and the consequences of their absence from current

databases[16]. For example, mass spectrometry-based proteomics has become the gold standard

for protein identification and has been extensively used in ALS studies[18,19]. However, if a protein

is not annotated, it is not included in the protein database (e.g. UniProtKB), and thus cannot be

detected by mass spectrometry. An estimated 50 % of mass spectra from a proteomics

experiment are unmatched at the end of the analysis[16,20].

Genome annotations must avoid spurious ORF annotations. Thus, unless functional

characterization has been published, they rely upon 2 arbitrary criteria: a minimum length of 100

codons and a single ORF per transcript. Several groups developed tools to challenge such criteria,

such as the sORFs repository[21] and the OpenProt database[17], which offer a data-driven broader

view of eukaryotic proteomes. The OpenProt database is based on a polycistronic model of ORF

annotation[17] and reports any ORF longer than 30 codons within any frame of an mRNA or ncRNA.

It contains currently annotated proteins (RefProts), novel Isoforms, and novel alternative proteins

(altProts). Here, we used the OpenProt database (www.openprot.org) to ask whether *FUS* may

encode additional proteins that could explain the toxicity of the wild-type protein over-

expression. In support of this hypothesis, FUS displays an N-terminal prion-like domain. These

low-complexity domains are known to harbour overlapping ORFs[22,23].

 ALTFUS IS A NOVEL 170 AMINO ACID PROTEIN, ENDOGENOUSLY EXPRESSED IN CELL LINES AND TISSUES

We began by querying OpenProt[17] predictions for *FUS* canonical mRNA (*ENST00000254108* or

NM_004960), which led to 8 predicted altORFs, either overlapping the coding sequence (CDS) or

within the 3'UTR (**Appendix Table 1**). Amongst these, IP_243680 or altFUS, a 170 codon altORF

overlapping FUS N-terminal prion-like domain, presents convincing experimental evidence of

expression (OpenProt v1.3). AltFUS overlaps the FUS CDS in an open reading frame shifted by one

nucleotide (**Fig 1A**, **Fig EV1A**). *FUS* is a complex gene with 13 annotated transcripts resulting from

alternative splicing. Based on the GTEx expression data in brain tissues and nerves, five transcripts

are more abundant and represent 85 % of all transcripts (**Fig EV1B**). Three of them (*FUS-206*, *FUS-*

211, *FUS-203*) are non-coding according to Ensembl, while only two (*FUS-211*, *FUS-203*) are non-

coding according to OpenProt (**Fig 1B, Fig EV1C**). Ensembl[24] annotates two transcripts as coding

(*FUS-201* and *FUS-202*), either for the 526 amino acid FUS protein or its 525 amino acid isoform

(**Fig EV1D**). From OpenProt prediction, these two transcripts also encode altFUS (IP_243680), or

its 169 amino acid isoform (IP_243691) respectively (**Fig EV1E**). Moreover, the second most

abundant transcript in brain tissues and nerves (*FUS-206*), representing about 20 % of all

transcripts, is non-coding according to Ensembl, but OpenProt predicts it contains the altFUS CDS

(**Fig 1B**, **Fig EV1C**). Thus, of the five most abundant transcripts in brain tissues and nerves, two

code for both FUS and altFUS proteins, one codes for altFUS alone, and the remaining two are

non-coding.

We retrieved nucleotide conservation scores (PhyloP) for *FUS* transcripts over 100 vertebrates.

PhyloP scores range from -10 (highly variable) to 10 (highly conserved). Scores over the FUS CDS

are under a constraint at the altFUS CDS locus (average score of 2.6 instead of 4 elsewhere on the

FUS CDS), which is consistent with a selection pressure across 2 overlapping frames (**Fig 1C**)[25]. We

then retrieved altFUS protein sequences over 84 species and observed a strong protein

conservation across mammals, and primates notably (75 to 99.4 % of sequence identity -

Appendix Table 2, **Fig EV1F and Appendix Source Data 1**). Thus, AltFUS is well conserved, with

domains showing little to no sequence variations (**Fig 1D**). No functional domain nor clear

secondary structures could be inferred from bioinformatics predictions (data not shown).

Published RIBO-seq data in Human, retrieved from the Gwips portal[26], revealed an accumulation

of initiating ribosomes around the altFUS initiating methionine, in association with an increase in

the density of elongating ribosomes over altFUS CDS (**Fig 1E**). These results suggest that altFUS is

translated. Similar results were observed in Mouse (**Fig EV2A**).

Based on the OpenProt database, AltFUS was identified in multiple proteomics datasets, with up

to 7 confident peptides (**Fig 1F, Appendix Table 1**). These peptides are unique to altFUS, or shared

with its isoform (IP_243691), and represent a 41 % sequence coverage (**Fig EV2B**). Furthermore,

we used a peptide-centric approach to query the TCGA datasets for altFUS expression with the

PepQuery algorithm. This approach allowed us to identify 28 peptides unique to altFUS, or shared

with its isoform (IP_243691), confidently mapped to mass spectra that could not be better

explained by any known protein (hg38_Ensembl database) (**Appendix Table 3**). These peptides

span through the entire altFUS sequence, representing a full sequence coverage (**Fig EV2C**). Out

of these, 20 peptides were confidently mapped to mass spectra that could not be better explained

by any known protein with any post-translational modification and/or chemical artefact (**Fig 1G**,

Fig EV2C).

To further validate altFUS protein expression, we developed a custom antibody targeting two

unique altFUS peptides (**Fig EV3A**) and tested it using three constructs: FUS, altFUS and FUS$^{(\emptyset)}$.

The latter is a monocistronic *FUS* version, where all altFUS methionines are mutated for

threonines in a manner synonymous for FUS (**Fig EV3B-D**). Thus, the FUS protein sequence is

unchanged, but the altFUS sequence does not contain any methionines. Transfection of HEK293

cells revealed expression of both proteins, FUS and altFUS, from the FUS nucleotide sequence (**Fig**

2A). As expected, altFUS expression was lost with the monocistronic FUS$^{(\emptyset)}$ construct. HEK293 cells

transfected with a siRNA targeting *FUS* mRNA showed a significant knockdown of both proteins,

FUS and altFUS; whereas altFUS endogenous expression was visible in scrambled control siRNA

and mock transfected cells (**Fig 2B**). These results validate the specificity of the custom antibody

for altFUS protein detection by western blot and demonstrate altFUS endogenous expression in

HEK293 cultured cells.

AltFUS endogenous expression was visible in control human tissues, HEK293 and HeLa cell lines

(**Fig 2C**). In order to test altFUS expression in pathological tissues, we retrieved motor cortex

lysates from 3 ALS patients with a *C9orf72* mutation (most common genetic cause) and 3 sporadic

ALS patients (most common aetiology). AltFUS endogenous expression was detected in all cases

(**Fig 2D**). Furthermore, as ALS is a motor neuron disease, we derived functional ventral spinal

motor neurons from induced pluripotent stem cells (iPSCs)[27,28] from healthy controls and ALS

patients carrying valosin-containing protein mutations (3 lines per group). AltFUS endogenous

expression was detected in all samples (**Fig 2E**). We noticed that brain (**Fig 2C**) and motor cortex

lysates (**Fig 2D**), as well as iPSCs-derived motor neurons (**Fig 2E**), from healthy controls and ALS

patients, presented a higher band detected with the custom altFUS antibody. This band is not

present in cultured cell lines or other tissues. It could come from a non-specific signal or a post-

translational modification on altFUS that is specific to the motor cortex and spinal cord motor

neurons (**Fig 2C-E**). Deep learning predictions of post-translational modifications on altFUS

revealed an extensive propensity to phosphorylation and O-GlcNAcylation (O-linked β-N-

acetylglucosamine glycosylation), with up to 19 and 18 sites respectively under a high stringency

model (**Appendix Table 4**). These two post-translational modifications are abundant in the

eukaryotic brain, and known to be dysregulated in neurodegenerative diseases or aging[29–32]. We

could also observe a lower band in the line 2 of controls motor neurons, which may correspond

to a degradation product or an initiation at a downstream methionine in altFUS sequence. This

band has never been observed in other samples so far. We demonstrate that the *FUS* gene

encodes two proteins, FUS and altFUS, both endogenously expressed in human tissues, iPSCs and

cell lines.

ALTFUS IS A MITOCHONDRIAL PROTEIN

FLAG-tagged altFUS (altFUS-FLAG) displayed a strong co-localization with a common mitochondrial

marker, TOMM20 (**Fig 2F**). Additionally, mitochondrial extracts showed an enrichment in altFUS-

FLAG (**Fig 2G**). Cellular fractionation of cells over-expressing untagged altFUS further validated

altFUS mitochondrial localization (**Fig 2H**). The endogenous altFUS protein was found in the

mitochondrial fraction, although it displayed a weak cytoplasmic signal as well (**Fig 2I**), consistent

with the immunofluorescence data (**Fig 2F**). Furthermore, cells over-expressing altFUS showed an

altered mitochondrial network, with a significant increase in fragmented mitochondria (globular)

compared to mock cells that displayed more tubular structures (**Fig 2J-K**, **Fig EV4A**).

Mitochondrial fragmentation is observed in models of overexpression of FUS mutants[33–35], and

we reproduced here a similar effect when over-expressing altFUS alone. Thus, we wondered

whether altFUS played a role in other mitochondrial dysfunctions observed in *FUS*-linked toxicity

models. To this end, we reproduced an ALS-associated FUS mutant: FUS-R495x[34,35]. This mutant

leads to a premature stop codon before FUS NLS and is linked to severe fALS and sALS cases[1]. In

this construct, altFUS is still present and not affected by the mutation (**Fig EV4B**). Similar to FUS$^{(\emptyset)}$,

we also generated the monocistronic construct FUS$^{(\emptyset)}$-R495x, which contains synonymous

mutations for FUS-R495x, but prevents altFUS expression. V5-FUS$^{(\emptyset\text{-FLAG})}$ and V5-FUS$^{(\emptyset\text{-FLAG})}$-R495x

did not express altFUS, but only the FUS protein, wild-type or ALS-linked mutant R495x

respectively (**Fig EV4C**). We first investigated the effect of altFUS on the mitochondrial membrane

potential using the potential sensitive dye TMRE (**Fig 3A-B, Fig EV4D**). As previously described[34],

over-expression of bicistronic FUS or FUS-R495x led to a decrease in mitochondrial membrane

potential. The mitochondrial membrane potential remained normal when over-expressing

monocistronic FUS$^{(\emptyset)}$ or FUS$^{(\emptyset)}$-R495x, underlining the role of altFUS. However, over-expression of

altFUS alone did not alter the mitochondrial membrane potential, which suggests both proteins

cooperate for this FUS-associated toxicity hallmark.

To further characterize altFUS, we investigated its protein interactors. Using stimulated emission

depletion microscopy (STED), we observed that altFUS localized in puncta following a cristae-like

pattern inside the mitochondria, delimited using an outer-membrane mitochondrial marker,

TOMM20 (**Fig 3C-D**). We then used size-exclusion chromatography on mitochondrial extracts to

isolate altFUS-FLAG macromolecular complexes (**Fig EV5A-B**). Following a FLAG affinity purification

and mass spectrometry (AP-MS) analysis, using a no-bait control for quantitative comparison, we

confidently identified 12 interacting proteins (**Fig 3E**, **Fig EV5C-D**, **Appendix Table 5**). These

proteins were identified with a minimum of 2 unique peptides and displayed over a 2-fold

enrichment to the control. A gene enrichment analysis to the Human proteome on subcellular

localization showed that the identified interactors are in majority proteins known to localize at

the mitochondria (**Fig 3F**). Amongst the identified interactors is the heat shock protein HSPA9, a

chaperone crucial in the mitochondrial iron-sulfur cluster biogenesis[36] (**Fig EV5D, Appendix Table**

5). Although HSPA9 was identified with 6 unique peptides and a fold-change of 6.53 to the control,

this protein is commonly detected in AP-MS datasets and thus likely non specific nor functionally

relevant[37]. Several altFUS interacting proteins are known to interact together, such as PHB,

ATAD3A, ERLIN2, RANBP2 and EMD (**Fig EV5D**). A refined gene enrichment analysis to Human

mitochondrial proteome identified three significantly enriched biological processes: autophagy-

related pathways, mitochondrial metabolism and cellular response to stress (**Fig 3G**). Disruptions

within these pathways are pathological hallmarks of *FUS*-linked toxicity and ALS[10,11,35,38].

ALTFUS INHIBITS AUTOPHAGY AND DRIVES THE ACCUMULATION OF FUS- AND TDP43-POSITIVE CYTOPLASMIC

AGGREGATES

Following on these results, we hypothesized that the inhibition of autophagy observed with ALS-

associated FUS mutants may instead be attributed to altFUS. We used the mCherry-GFP-LC3

reporter to track the autophagic flux by confocal microscopy (**Fig EV6A**). Under basal conditions,

cells displayed red and yellow foci as expected (**Fig 4A**). An accumulation of yellow foci was

observed when cells were treated with bafilomycin, an inhibitor of autophagy. Similarly, cells

over-expressing altFUS displayed a significant accumulation of yellow foci (**Fig 4A**). Furthermore,

our results were consistent with previously published data[39] as cells transfected with FUS or FUS-

R495x displayed a decreased autophagic flux (**Fig 4A, Fig EV6B**). This accumulation of yellow foci

was absent in cells that express monocistronic FUS constructs, thus lacking altFUS expression

(FUS$^{(\emptyset)}$ or FUS$^{(\emptyset)}$-R495x). We used bafilomycin followed by LC3 probing to further validate the

impact of altFUS on autophagy (**Fig 4B, Fig EV6C**). Similarly, an inhibition of autophagy was

observed only in cells over-expressing altFUS. Furthermore, in cells over-expressing monocistronic

FUS$^{(\emptyset)}$-R495x, the inhibition of autophagy could be restored by co-transfecting altFUS (**Fig 4B, Fig**

EV6C). These results establish altFUS, rather than FUS, as the protein responsible of the inhibition

of autophagy.

Furthermore, altFUS interactome analysis suggested a role in the cellular stress response, which

is known to be altered in ALS with a TDP-43 cytoplasmic accumulation in 98 % of patients[40,41]. In

FUS-linked ALS and some sALS cases, FUS cytoplasmic aggregates or mislocalization are also

observed[42–44]. In our hands, cells over-expressing FUS-R495x displayed cytoplasmic aggregates

that were positive for both FUS-R495x and TDP-43 (**Fig 4C**). Although TDP-43 aggregates are not

a common observation with FUS-associated mutants, co-aggregation has already been reported

in patients, animal models and cultured cell lines across multiple studies[42,44–49]. In cells over-

expressing the monocistronic FUS$^{(\emptyset)}$-R495x construct, thus lacking altFUS expression, FUS-R495x

displayed a more diffuse cytoplasmic localization, and TDP-43 remained in the nucleus (**Fig 4C**).

FUS cytoplasmic aggregates were significantly more numerous and larger when altFUS was co-

expressed (**Fig 4D-E**). Accumulation of FUS-R495x and TDP-43 in cytoplasmic aggregates could be

reconstituted by co-transfecting altFUS and the monocistronic FUS$^{(\emptyset)}$-R495x construct (**Fig 4C**).

These observations were repeated across all 7 ALS-associated FUS mutations tested (**Fig EV6D-F**).

The cytoplasmic aggregates were also TIAR-positive as observed with ALS-linked FUS mutants in

previous work[40] (**Fig EV7**). These results suggest that altFUS enhances the assembly of cytoplasmic

FUS mutants aggregates and is responsible for the recruitment of TDP-43 in these aggregates.

ALTFUS PROTECTS AGAINST NEURODEGENERATION IN FUS-ASSOCIATED DROSOPHILA MODELS

In order to investigate the role of altFUS in an already established *in vivo* model of FUS-related

neurodegeneration, we generated *Drosophila* models expressing either the bicistronic, FUS and

FUS-R495x constructs, or the monocistronic, FUS$^{(\emptyset)}$ and FUS$^{(\emptyset)}$-R495x, constructs. We used the

Elav-GeneSwitch-GAL4 Driver strain, as previously described[50], as it allows for an inducible over-

expression in motor neurons and avoids lethality at the larval stage from FUS over-expression in

the central nervous system[50,51]. First, we generated flies containing the sequences for UASt-

altFUS, UASt-FUS, UASt-FUS$^{(\emptyset)}$, UASt-FUS-R495x or UASt-FUS$^{(\emptyset)}$-R495x. These flies were then

crossed with the Elav-GeneSwitch-GAL4 driver strain (**Fig 5A**). UASt-mCherry flies were used as

controls. Selected F1 individuals were then divided into 2 groups with equal proportions of

males/females. The first group received standard food, while the other received RU-486 treated

food. The treatment induces a conformational change in the Elav-GeneSwitch driver, which allows

activation of the UAS promoter and thus expression of the target protein. We retrieved flies at

selected time points to validate protein expression in the RU-486 treated population through

time, while the controls showed no expression (**Fig 5B**).

The motor neuron degeneration linked to ALS provokes a progressive locomotion loss measurable

with a well-described climbing assay[52]. The control populations did not show any significant

locomotion loss at day 1, 10 nor 20 (**Fig 5C-E**). Similarly to the RU-486 treated control group did

not show a significant effect, although a decrease in climbing ability could be observed as

previously reported[53,54] (mCherry transgenic flies – **Fig 5C**). AltFUS flies did not show any

significant locomotion loss through time (**Fig 5C**). This result is consistent with the *in cellulo* data

showing altFUS alone is not sufficient to provoke pathological hallmarks. As previously shown with

this model[8], the bicistronic FUS flies, which express both FUS and altFUS proteins, displayed a

significant locomotion loss (**Fig 5D**). Bicistronic ALS-linked FUS-R495x flies showed an even greater

motor neuron degeneration through time compared to FUS (**Fig 5E**). Monocistronic FUS$^{(\emptyset)}$ (**Fig 5D**)

and FUS$^{(\emptyset)}$-R495x (**Fig 5E**) flies, which do not express altFUS, displayed both a delay in the onset

of motor neuron degeneration and a reduced drop in climbing success at 20 days post-induction

(40% vs. 60% for FUS$^{(\emptyset)}$, 30% vs. 70% for FUS$^{(\emptyset)}$-R495x). These results in *Drosophila* confirm a role

for altFUS in *FUS*-related neurodegeneration *in vivo*, are consistent with our *in cellulo*

observations and highlight the toxic cooperation between FUS and altFUS.

ALS-ASSOCIATED MUTATIONS, SYNONYMOUS FOR FUS, ALTER ALTFUS AND LEAD TO TDP-43 CYTOPLASMIC

AGGREGATES

As of today, over 50 mutations in the *FUS* gene have been associated with ALS[1]. However, most

of these locate at the carboxyl end of the protein and as such have no effect on altFUS (**Fig 1A**).

We wondered whether mutations altering altFUS might have been overlooked as non-

consequential in the FUS reading frame. We retrieved FUS synonymous mutations found in ALS

patients, with an allelic frequency below 0.01 %, from previous studies and the ALS Variant Server

(http://als.umassmed.edu/ - **Appendix Table 6**). The retrieved mutations clustered on the altFUS

locus (**Fig 6A**), with 60 % of FUS synonymous mutations found in sALS patients and 50 % of FUS

synonymous mutations found in fALS patients, which is significantly higher than expected by

chance (34 %) (**Appendix Table 6**). We selected 4 mutations for further analysis based on the

residue conservation: altFUS-P31L, altFUS-A38V, altFUS-A46V and altFUS-R64P. We generated

them in GFP-FUS constructs: GFP-FUS[(P31L-FLAG)]-S44=; GFP-FUS[(A38V-FLAG)]-G51=; GFP-FUS[(A46V-FLAG)]-G59=

and GFP-FUS[(R64P-FLAG)]-S77=. All altFUS mutants still localized to the mitochondria (**Fig EV8A**). To

investigate whether these mutations may provoke an ALS-like phenotype, we quantified the

number of cells presenting TDP-43 aggregates. All 4 altFUS mutants displayed clear TDP-43

aggregates, and showed a 1.8 to 2.4 fold increase compared to wild-type altFUS (**Fig 6B-C, Fig**

EV8B). This result indicates that altFUS mutations potentiate TDP-43 cytoplasmic aggregation, a

pathological hallmark in 98 % of ALS cases. Hence, some *FUS* mutations, synonymous for the FUS

protein, exert a deleterious effect through their missense consequence on the altFUS protein.

CONCLUSIONS: FUS AND ALTFUS, ENCODED BY THE SAME GENE, COOPERATE IN FUS-RELATED TOXICITY

Despite considerable advances in the field, current genome annotations still uphold arbitrary

assumptions, such as the monocistronic nature of eukaryotic genes[16]. Here, we demonstrate *FUS*

is a bicistronic gene. We discovered *FUS* CDS contains a second protein-coding sequence in a

shifted frame overlapping its prion-like intrinsically disordered domain, regions known to host

dual-coding events[22,23]. This novel protein, named altFUS, is not an isoform but an entirely new

sequence of 170 amino acids. AltFUS is endogenously expressed in human tissues and cultured

cell lines, as demonstrated by ribosome profiling, mass spectrometry and with a custom antibody.

AltFUS is notably expressed in the motor cortex and iPSCs-derived motor neurons of healthy

controls and ALS patients. Because altFUS is embedded within the *FUS* CDS, this discovery is of

crucial importance to the field. Indeed, over-expression studies on *FUS* actually implicate two

proteins: FUS and altFUS. Similarly FUS knockdown or knockout studies actually inhibit expression

of both proteins[55]. Moreover, previous work has shown that gene editing techniques targeting a

specific CDS do not necessarily result in knockout of the gene in case of dual-coding gene[56]. Our

discovery thus suggests that *FUS* edited cells or models, notably targeting its last exons[57–59], might

only impair FUS protein expression but not altFUS, thus not resulting in a true *FUS* knockout. Our

work provides a more accurate view of *FUS* coding potential to better understand its physiological

function and models of FUS-related neurodegeneration.

Following the discovery of altFUS, we developed tools in order to differentiate the specific roles

and phenotypes of FUS and altFUS. Our study demonstrates that altFUS is necessary for three

toxic molecular hallmarks previously attributed to FUS: mitochondrial fragmentation and loss of

mitochondrial membrane potential, inhibition of autophagy, and cytoplasmic aggregation of FUS.

The inhibition of autophagy was observed when over-expressing altFUS alone, absent when over-

expressing FUS alone (wild-type or ALS-associated R495x mutant) and reconstituted when co-

expressing FUS and altFUS. This demonstrates that the inhibition of autophagy, previously

described in *FUS*-ALS[10,39], has been incorrectly associated to the FUS protein. AltFUS inhibits

autophagy. Moreover, altFUS is necessary but not sufficient for the mitochondrial membrane

potential loss and cytoplasmic aggregation of FUS and TDP-43. Although TDP-43 aggregates are

not commonly seen with FUS mutations[44], it has already been reported[42,44–49] and this study lends

support for altFUS conspiring with FUS and TDP-43 to lead to this molecular hallmark. Our data

suggests the stoichiometry between FUS and altFUS may be important for the development of

cytoplasmic aggregates. Both proteins are required to observe the phenotype, highlighting a

functional alliance between FUS and altFUS. This pathological synergy was also observed in the

Drosophila model.

The physiological function of altFUS is still unclear, although our work provides evidence for its

role in mitochondrial dynamics and the cellular response to stress. One mechanism put forward

in ALS is that the disease originates from a sub-optimal resolution of cellular stresses, which can

come from environmental sources or mutated proteins[38,60]. We have shown that altFUS, not FUS,

inhibits autophagy, most likely via its interaction partners (**Appendix Table 5**). Since an inhibition

of autophagy inhibits the dissociation of stress granules[38], we suggest altFUS potentiates stress

granule accumulation under stress conditions, and that FUS phase separation properties then lead

to the formation of solid and toxic aggregates[61]. Further work is needed to fully understand the

role of altFUS in physiological and pathological conditions, yet our study shows this novel protein

plays a crucial role in *FUS*-linked gain-of-toxic dysfunctions in models of *FUS*-related

neurodegeneration.

Recent studies have addressed the toxicity resulting from over-expression of wild-type FUS.

Bogaert and colleagues used FUS domain truncation mutants to investigate wild-type FUS

toxicity[51]. A FUS mutant lacking its N-terminal intrinsically disordered domain, thus lacking altFUS,

displayed reduced toxicity. This study corroborates our findings that the absence of altFUS

reduces the toxicity. Furthermore, Bogaert and colleagues concluded that FUS N-terminal

synergizes with the C-terminal domain to mediate toxicity in *Drosophila*[51]. Here, we showed that

altFUS synergizes with FUS to mediate ALS-like toxic features in cultured cells and toxicity in

Drosophila. Despite different laboratories and different techniques, our data are in agreement

with theirs. Yet, we point to an alternative (not necessarily mutually exclusive) explanation

whereby altFUS, not the FUS N-terminal domain, synergizes for toxicity. Additionally, this

discussion shows that not being aware of overlapping CDSs, especially in deletion studies,

precludes alternative interpretations.

Current genome annotations guide the interpretation of data and the design of studies, however

they also affect the way we screen for pathological mutations[16]. Most of the ALS-linked *FUS*

mutations affect the carboxyl end of the protein, as shown with the example used in this study,

the R495x mutation[62]. These mutations do not alter the altFUS protein, which is embedded at the

beginning of the FUS CDS and span on exons 3 to 6. How could altFUS be important for the disease

if most pathological mutations do not alter it? To answer this question, one needs to grasp how

much genome annotations shape today's research. For example, when screening for pathological

mutations, those that are synonymous for FUS are discarded early in analyses as insignificant[63].

Yet, a synonymous mutation for FUS may not be for altFUS. Our work shows that FUS synonymous

mutations found in patients cluster on altFUS genomic locus. We tested 4 of these mutations,

synonymous for the FUS protein and missense for the altFUS protein, and we found that each of

them potentiated TDP-43 cytoplasmic aggregation, a pathological hallmark of ALS[41].

Overall, we have shown that *FUS* is a bicistronic gene and our results indicate that altFUS

interferes with mitochondrial homeostasis and autophagy in cell culture and induces motor

neuron toxicity in *Drosophila* models. It will be important to further characterize the function of

altFUS and determine if it has a role in ALS and/or FTLD.

METHODS

FUS constructs

FUS and altFUS sequences were obtained from Bio Basic Gene Synthesis service. All FUS constructs

were subcloned into pcDNA3.1- (Invitrogen) using Gibson assembly (New England Biolabs,

E26115). FUS and altFUS wild-type sequences correspond to that of the Human FUS canonical

transcript (*ENST00000254108* or *NM_004960*). FUS and altFUS proteins were tagged with V5

(GKPIPNPLLGLDST) and 2 Flag (DYKDDDDKDYKDDDDK) respectively. FUS was tagged on the N-

terminal, altFUS was tagged on the C-terminal. For immunofluorescence assays, N-terminal GFP-

tagged FUS was also cloned into pcDNA3.1- by Gibson assembly. The necessary gBlocks were

purchased from IDT. The monocistronic constructs $FUS^{(\emptyset)}$ and $FUS^{(\emptyset)}$-R495x were generated by

mutating all altFUS methionines (ATG) to threonines (ACG). These mutations are synonymous in

the FUS CDS (TAT > TAC, both coding for tyrosine). The altFUS mutated sequence was obtained

from Bio Basic Gene Synthesis service, and then subcloned in FUS sequences in pcDNA3.1- using

Gibson assembly. The bicistronic constructs are named as follows throughout the article: FUS,

FUS-R495x, or $FUS^{(Flag)}$ and $FUS^{(Flag)}$-R495x when altFUS is Flag-tagged in the +2 reading frame. The

monocistronic constructs are named as follows throughout the article: $FUS^{(\emptyset)}$ or $FUS^{(\emptyset)}$-R495x to

indicate altFUS absence.

Cell culture, transfections, western blots and immunofluorescence

HEK293 and HeLa cells cultures tested negative for mycoplasma contamination (ATCC 30–1012K).

Transfections, immunofluorescence, confocal analyses and western blots were carried out as

previously described[64]. For FUS knock-down, 150 000 HEK293 cells in a 6-well plate were

transfected with 25 nM FUS SMARTpool: siGENOME siRNA (Dharmacon, Canada, L-009497-00-

0005) or ON-TARGET plus Nontargeting pool siRNAs (Dharmacon, D-001810-10-05) with

DharmaFECT one transfection reagent (Dharmacon, T-2001–02) according to the manufacturer's

protocol. Cell media was changed every 24 hrs and cells were processed 72 hrs after transfection.

For immunofluorescence, primary antibodies were diluted as follows: anti-Flag (Sigma, F1804) 1/

1000, anti-TOMM20 (Abcam, ab186734) 1/500, anti-V5 (Cell Signalling Technologies, #13202)

1/1000, anti-TDP-43 (ProteinTech, 10782-2-AP) 1/500, and anti-TIAR (Cell Signalling Technologies,

#8611) 1/1600. For western blots, primary antibodies were diluted as follows: anti-Flag (Sigma,

F1804) 1/8000, anti-V5 (Sigma, V8012) 1/8000, anti-actin (Sigma, A5441) 1/40000, anti-FUS

(Abcam, ab84078) 1/500, anti-altFUS (Abcam, custom antibody) 1/3000, anti-LC3 (Cell Signalling

Technologies, #2775) 1/1000, anti-Hsp70 (Thermo Fisher Scientific, MA3-028) 1/1000, anti-

Tubulin (Thermo Fisher Scientific, a11126) 1/2000 and anti-VDAC (Abcam, ab15895) 1/10000. The

altFUS antibody was generated by injection two rabbits, each with 2 unique altFUS peptide. The

purified antibody from rabbit 2 was used in this study at a 1/2000 dilution. Mitochondrial

morphology was evaluated using the microP tool[65]. A minimum of 100 cells per replicate were

counted across 3 independent experiments (n = 3, i.e. minimum 300 cells for each experimental

condition). Colocalization analyses were performed using the JACoP plugin (Just Another

Colocalization Plugin) implemented in Image J software, as previously described[13]. When

specified, images obtained by confocal microscopy on the Leica TCS SP8 STED 3X were

deconvolved using the Huygens software (Scientific Volume Imaging B.V., Hilversum,

Netherlands). The software uses a signal reassignment algorithm for deconvolution, identical

deconvolution parameters were applied to all images. The default parameters were used,

including the Classic Maximum Likelihood Estimation (CMLE) algorithm, signal to noise ratio, and

background estimation radius. The maximum iteration number was set at 30. Human tissue

lysates for altFUS endogenous expression were purchased from Zyagen Laboratories (San Diego,

California, USA).

Rɪʙᴏ-seq data and conservation analyses

Global aggregate reads for initiating ribosomes and elongating ribosomes footprints across all

available studies were downloaded from the Gwips portal (https://gwips.ucc.ie/), for *Homo*

sapiens and for *Mus musculus*. For altFUS protein conservation analysis, all FUS mRNAs with at

least EST evidence were retrieved across all available species from NCBI RefSeq. We performed

an *in silico* 3-frame translation and retrieved the best matching protein sequence per species that

displayed a minimum of 20 % sequence identity with the Human altFUS sequence over 25 % of

Human altFUS length. AltFUS homologous sequences were found in 83 species, and we manually

added that of *Drosophila melanogaster* that displayed a 37.5 % sequence identity over 19 % of

the Human altFUS length. All retrieved altFUS sequences were then aligned using Clustalω with

default parameters.

Peptide-centric analysis of proteomics datasets

The stand alone PepQuery tool (v1.0) was downloaded from the PepQuery website

(http://www.pepquery.org/). The tool was run on the following datasets from the TCGA

consortium: colon cancer (COCA) proteome, ovarian cancer (OVCA) proteome, phosphoproteome

and glycoproteome, and the breast cancer (BRCA) proteome and phosphoproteome. The

reference database was set to the Ensembl datababase (hg38_Ensembl_20190910). The following

parameters were set for all runs unless specified: carbamidomethylation of cysteine as fixed

modification (as well as iTRAQ 4-plex of K, iTRAQ 4-plex of peptide N-term for BRCA and OVCA);

oxidation of methionine as variable modification (as well as iTRAQ 4-plex of Y for BRCA and OVCA);

a maximum of 3 modifications per peptides; trypsin digestion with maximum of 1 miscleavage;

precursor tolerance of 10 ppm (20 ppm for COCA); fragment mass tolerance of 0.05 Da (0.6 Da

for COCA); the hyperscore was used as a scoring metric, and 10 000 randoms. For

phosphoproteomes, phosphorylation of Y, T and S were added as variable modifications. For the

glycoproteome, deamidation of Q and N were added as variable modifications. PepQuery was run

in the protein mode, with altFUS (IP_243680) whole sequence as input. Spectra were visualised

and drawn using an in-house python script.

Human induced pluripotent stem cell differentiation into motor neurons

Directed differentiation to human iPSC-motor neurons was performed as previously reported[27].

Briefly, iPSCs were maintained on Geltrex (Life Technologies) with Essential 8 Medium media (Life

Technologies), and passaged using EDTA (Life Technologies, 0.5mM). All cell cultures were

maintained at 37°C and 5% carbon dioxide. For motor neuron differentiation, iPSCs were

differentiated to neuroepithelium by plating to 100% confluency in chemically defined medium

consisting of DMEM/F12 Glutamax, Neurobasal, L-Glutamine, N2 supplement, non-essential

amino acids, B27 supplement, β-mercaptoethanol (Life Technologies) and insulin (Sigma).

Treatment with the following small molecules from day 0-7: 1µM Dorsomorphin (Millipore), 2µM

SB431542 (Tocris Bioscience), and 3.3µM CHIR99021 (Miltenyi Biotec). At day 8, cells patterned

for 7 days with 0.5µM retinoic acid and 1µM Purmorphamine. At day 14 spinal cord motor neuron

precursors were treated with 0.1µM Purmorphamine for a further 4 days before being terminally

differentiated for >10 days in 0.1 µM Compound E (Enzo Life Sciences) to promote cell cycle exit.

Throughout the neural conversion and patterning phase (D0-18) the neuroepithelial layer was

enzymatically dissociated twice (at D4-5 and D10-12) using dispase (GIBCO, 1 mg ml-1).

Preparation of tissue lysates of the motor cortex of ALS patients

Approximately 100mg of motor cortex from 4 sporadic ALS and 4 C9orf72-ALS cases was lysed in

10x RIPA (50mM Tris HCl pH7.8, 150mM NaCl, 0.5% sodium deoxycholate, 1% NP40;

supplemented with protease inhibitors and EDTA) volume using TissueLyzer equipment (Qiagen).

Lysates were incubated on ice 20 minutes followed by centrifugation at 20,000xg for 20 minutes

at 4°C. Supernatant was taken as 'RIPA fraction' and pellets resuspended in RIPA and SDS (final

concentration of 2%). 3 sporadic ALS and 3 C9orf72-ALS samples were subsequently used as they

were sufficiently concentrated to load 100 ug of proteins onto SDS-page gels.

Mitochondrial extracts and cellular fractionation

Mitochondrial extracts were prepared as previously described[56]. Briefly, HEK293 cells grown up

to 80 % confluence, were rinsed twice with PBS and gathered using a cell scraper. Cells were

pelleted by centrifugation at 500 x g for 10 mins at 4 °C. Supernatant was discarded and cells

suspended in mitochondrial buffer (210 mM mannitol, 70 mM sucrose, 1 mM EDTA, 10 mM

HEPES-NaOH, pH 7.5, 0.5 mM PMSF and EDTA-free protease inhibitor (Thermo Fisher Scientific)).

Cells were disrupted by 15 consecutive passages through a 25G1 0.5 x 25 needle syringe on ice,

followed by a 3 min centrifugation at 2 000 x g at 4 °C. Supernatant was collected and the pellet

suspended in mitochondrial buffer. The cell disruption was repeated four times and all retrieved

supernatants containing mitochondria were again passed through syringe needle in mitochondrial

buffer and cleared by centrifugation for 3 mins at 2000 × g at 4 °C. Supernatants were pooled and

centrifuged for 10 mins at 13 000 x g at 4 °C to pellet mitochondria. The pellet was suspended in

200 µL of mitochondrial buffer until further processing. Cellular fractionation was performed

using the Cell Fractionation Kit (#9038S, Cell Signaling Technology). Briefly, HEK293 cells were

grown up to 80 % confluence, washed twice with PBS and gathered using a cell scraper. Cells were

spun at 350 x g for 5 mins at 4 °C and 2.5×10^6 cells were suspended in 500 µL of ice-cold PBS. An

aliquot of 100 µL was spun at 350 x g for 5 mins at 4 °C and resuspended in SDS buffer (4 % SDS,

Tris-HCl 100 mM pH 7.6) and kept as WCL (Whole Cell Lysate). The rest of the collected cells

(remaining 400 µL) were spun at 500 x g for 5 mins at 4 °C. Supernatant was discarded and pellet

resuspended in 500 µL of CIB (Cytoplasmic Isolation Buffer) from the kit, vortexed for 5 secs and

incubated on ice for 5 mins. After centrifugation at 500 x g for 5 mins at 4 °C, the supernatant was

collected as the cytosplasmic fraction. The pellet was resuspended in 500 µL of MIB buffer

(Membrane Isolation Buffer) from the kit, vortexed for 15 secs and incubated on ice for 5 mins.

After centrifugation at 8 000 x g for 5 mins at 4 °C, the supernatant was collected as the membrane

and organelles fraction. To each 100 µL of fraction was added 60 µL of loading buffer 1 x (from

ColdSpring Laemmli sample buffer: 50 mM Tris pH 6.8, 2 % SDS, 10 % Glycerol, 5 % β-

mercaptoethanol) before processing for western blot.

Mitochondrial membrane potential measurements

Mitochondrial membrane potential was measured by flow cytometry in HEK293 cells using TMRE

(Tetramethylrhodamine, Ethyl ester, Abcam, ab113852). FCCP was used as a positive control to

validate each independent experiment. Cells were grown up to 80 % confluence and washed twice

with PBS. The cells were then incubated for 5 mins at 37 °C, 5 % CO_2 with PBS/A (0.2 % BSA in PBS)

solution (experimental) or 3 µM FCCP in PBS/A solution (positive control). Then, 100 nM of TMRE

was added and cells were incubated 15 mins at 37 °C, 5 % CO_2. After incubation, cells were

trypsinized and centrifuged at 800 x g for 5 mins at 4 °C and resuspended in 500 µL of PBS and

kept on ice. Cells were immediately analysed by flow cytometry. A gate for living cells was set, as

well as a second gate to filter out cell doublets. TMRE fluorescence (PE-A) was recorded over a

minimum of 50 000 gated cells for each experimental condition. The mean TMRE fluorescence

intensity was measured over 3 independent experiments for each experimental condition.

Stimulated Emission Depletion (STED) microscopy

Samples were prepared as described above for confocal microscopy. A Leica TCS SP8 STED 3X was

used with a 100x objective lens and immersion oil for dual-color STED images. Images were

obtained by sequential scanning of a given area. The combination of Alexa Fluor 488 (Thermo

Fisher Scientific, A-11017) and Alexa Fluor 568 (Thermo Fisher Scientific, A-21069) dyes was

chosen for STED imaging. Alexa Fluor 488 dye was excited with a white light laser (WLL) at 488 nm

and was depleted using the 660 nm STED laser. Alexa Fluor 568 dye was excited with a WLL at

561 nm and was depleted using the 660 nm STED laser. The STED laser (660 nm) was applied at

80 % of maximum power.

Fast Protein Liquid Chromatography (FPLC) and affinity-purification mass spectrometry (AP-MS)

Mitochondrial extracts of HEK293 cells were centrifuged at 13 000 x g for 10 mins at 4 °C to

remove the supernatant and were resuspended in FPLC buffer (50 mM Tris-HCl, 1 mM EDTA, 150

mM NaCl, 1 % Triton X-100, pH 7.5, filtered with 0.2 µm filters) at 2 mg/mL for a total of 4 mg of

mitochondrial proteins. Samples were incubated on ice for 15 mins and then centrifuged at 10 000

x g for 5 mins at 4 °C and the supernatant was loaded in the injection syringe without disrupting

the pellet. The FPLC was performed on a HiLoad 16/60 Superdex 200 pg column (GE Healthcare,

Chicago, USA) at 4 °C. The column was pre-equilibrated with the FPLC buffer for up to 0.2 CV

(column volume) and the sample was applied at a flow rate of 0.5 mL/min with a pressure alarm

set at 0.5 MPa. The elution was performed over 72 fractions of 1.5 mL for a maximum of 1.1 CV.

For altFUS probing by western blot, proteins were precipitated from 150 µL of each 4 fractions in

technical duplicates. First, 600 µL of methanol was added to each tube and mixed gently, before

adding 150 µL of chloroform. Tubes were gently inverted 10 times before adding 450 µL of milliQ

H_2O and vortexing briefly. After centrifugation at 12 000 x g for 3 mins, the upper phase was

discarded, and 400 µL of methanol was added. Tubes are centrifuged at 16 000 x g for 4 mins and

the pellet was resuspended in loading buffer. For interactome analysis by mass spectrometry,

fractions of interest (8 to 14) were pooled together and incubated at 4 °C overnight with magnetic

FLAG beads (Sigma, M8823) pre-conditioned with FPLC buffer. The beads were then washed 3

times with 5 mL of FPLC buffer, and 5 times with 5 mL of 20 mM NH_4HCO_3 (ABC). Proteins were

eluted and reduced from the beads using 10 mM DTT (15mins at 55 °C), and then treated with 20

mM IAA (1 hour at room temperature in the dark). Proteins were digested overnight by adding 1

µg of trypsin (Promega, Madison, Wisconsin) in 100 µL ABC at 37 °C overnight. Digestion was

quenched using 1 % formic acid and supernatant was collected. Beads were washed once with

acetonitrile/water/formic acid (1/1/0.01 v/v) and pooled with supernatant. Peptides were dried

with a speedvac, desalted using a C18 Zip-Tip (Millipore Sigma, Etobicoke, Ontario, Canada) and

resuspended into 30 µl of 1% formic acid in water prior to MS analysis.

Mass-spectrometry analysis

Peptides were separated in a PepMap C18 nano column (75 µm × 50 cm, Thermo Fisher Scientific).

The setup used a 0–35% gradient (0–215 min) of 90% acetonitrile, 0.1% formic acid at a flow rate

of 200 nL/min followed by acetonitrile wash and column re-equilibration for a total gradient

duration of 4 h with a RSLC Ultimate 3000 (Thermo Fisher Scientific, Dionex). Peptides were

sprayed using an EASYSpray source (Thermo Fisher Scientific) at 2 kV coupled to a quadrupole-

Orbitrap (QExactive, Thermo Fisher Scientific) mass spectrometer. Full-MS spectra within a m/z

350–1600 mass range at 70,000 resolution were acquired with an automatic gain control (AGC)

target of 1e6 and a maximum accumulation time (maximum IT) of 20 ms. Fragmentation (MS/MS)

of the top ten ions detected in the Full-MS scan at 17,500 resolution, AGC target of 5e5, a

maximum IT of 60 ms with a fixed first mass of 50 within a 3 m/z isolation window at a normalized

collision energy (NCE) of 25. Dynamic exclusion was set to 40 s. Mass spectrometry RAW files were

searched with Andromeda search engine implemented in MaxQuant 1.5.5.1. The digestion mode

was set at Trypsin/P with a maximum of two missed cleavages per peptides. Oxidation of

methionine and acetylation of N-terminal were set as variable modifications, and

carbamidomethylation of cysteine was set as fixed modification. Precursor and fragment

tolerances were set at 4.5 and 20 ppm respectively. Files were searched using a target-decoy

approach against UniprotKB (*Homo sapiens* 03/2017 release) with the addition of altFUS

sequence for a total of 92,949 entries. The false discovery rate (FDR) was set at 1% for peptide-

spectrum-match, peptide and protein levels. Protein interactions were then scored using the

SAINT algorithm, with Mock cells as control and the magnetic FLAG beads in HEK293 cells

crapome[66]. Proteins with a SAINT score above 0.99 were considered, as well as those presenting

a SAINT score above 0.88 with a minimum of two unique peptides.

Biological processes and cellular compartment enrichment analysis

Proteins identified in altFUS interactome were screened for cellular compartment and biological

processes enrichment using Gene Ontology (GO) enrichment. Proteins were queried against the

whole Human Proteome for cellular compartment and against the Human mitochondrial

proteome (MitoCarta 2.0) for biological processes. The statistical analysis used a Fisher's Exact

test with a FDR set at 1 %.

Autophagic flux measurements

The mCherry-GFP-LC3 was used to evaluate the autophagic vesicles within HeLa cells by confocal

microscopy. Before fusion with the lysosome, the LC3 molecules on the autophagosome display

a yellow fluorescence (combined mCherry and GFP fluorescence). After fusion, the GFP

fluorescence is quenched by the lysosomal pH, and as such the LC3 molecules display a red signal

(mCherry alone). This allows a visual representation of the autophagic flux in a given cell. Cells

treated with 50 nM Bafilomycin for 4 hours were used as a positive control to validate each

independent experiment. Observations were made across 2 technical duplicates for each

biological condition, across 3 independent experiments (n=3). Alternatively, the autophagic flux

was also evaluated by LC3 probing before and after bafilomycin treatment (50 nM for 4 hours).

The quantification corresponds to the treated / untreated ratio of LC3-II abundance.

Cytoplasmic aggregates measurements

Images of HeLa cells were taken by confocal microscopy and then processed using the Image J 3D

Objects Counter plugin. FUS cytoplasmic aggregates were then quantified in number and size

(μm^2) for each cell. A total of 100 cells across two technical replicates were taken for each

independent experiment (n=3, i.e. a minimum of 300 cells per biological conditions).

Transgenic *Drosophila* and climbing assay

The bicistronic constructs, FUS and FUS-R495x, and the monocistronic constructs, altFUS, FUS$^{(\emptyset)}$

and FUS$^{(\emptyset)}$-R495x, were subcloned in the pUASTattB expression vector for site specific insertion

into attP2 on chromosome 3. Transgenic flies were generated by Best Gene (Best Gene Inc.,

California, USA). The Elav-GeneSwitch-GAL4 driver (stock number: 43642, genotype: y[1] w[*];

P{w[+mC]=elav-Switch.O}GSG301) and the UAS-mCherry flies (stock number: 35787, genotype:

y[1] sc[*] v[1]; P{y[+t7.7] v[+t1.8]=UAS-mCherry.VALIUM10}attP2) was purchased from

Bloomington (Bloomington Drosophila Stock Center, Indiana, USA). All stocks were in a *w^{1118}*

background and were cultured on standard medium at 25°C or room temperature. Transgenic

flies were crossed with the Elav-GeneSwitch-GAL4 driver strain. The F1 was equally divided in two

groups with equal proportion of males and females: one group will feed on standard food

supplemented with ethanol (0.2 % - control flies), the other on standard food supplemented with

RU-486 at 10 μM diluted in ethanol (induced flies). The climbing assay was performed as

previously described[52]. Briefly, flies were transferred into an empty vial and tapped to the bottom.

After 18 s, the number of flies at the top of the tube were considered successful. The assay was

done at day 1, 10 and 20 post-induction, across 4 independent F1. Five flies were taken at day 1,

10 and 20 post-induction to validate expression of the proteins of interest.

Statistical analyses and representation

Unless otherwise stated, the statistical analysis carried was a two-way ANOVA with Tukey's

multiple comparison correction. The box plots represent the mean with the 5 to 95 % percentile.

The bar graphs represent the mean, and error bars correspond to the standard deviation. When

using parametric tests, normality of data distribution was verified beforehand using the Shapiro-

Wilk test.

DATA AVAILABILITY STATEMENT

The OpenProt database is available at www.openprot.org. The GTEx portal is available via

www.gtexportal.org. The Gwips portal is available at www.gwips.ucc.ie. The proteomics data are

available on the PRIDE repository with the accession PXD------. Any other relevant data are

available from the corresponding authors upon reasonable request.

ACKNOWLEDGEMENTS

This research was supported by CIHR grants MOP-137056 and MOP-136962, by an ALS Canada

Project Grant, and by a Canada Research Chair in Functional Proteomics and Discovery of Novel

Proteins to X.R. X.R and S.J are members of the Fonds de Recherche du Québec Santé (FRQS)-

supported Centre de Recherche du Centre Hospitalier Universitaire de Sherbrooke. P.M. is

supported by the ALS Double Play Christopher Chiu Postdoctoral Fellowship. We thank J. Ule, M-

J. Boucher and D. Hunting for helpful discussions. The autopsy program is supported by the James

Hunter and Family ALS Initiative. The Genotype-Tissue Expression (GTEx) Project was supported

by the Common Fund of the Office of the Director of the National Institutes of Health, and by NCI,

NHGRI, NHLBI, NIDA, NIMH, and NINDS. The data used for the analyses described in this

manuscript were obtained from the GTEx Portal on April 2019.

CONTRIBUTIONS

MA.B and X.R designed and wrote the study. MA.B and JF.J did the experiments, and MA.B did

the analyses and figures. S.N and S.J assisted with the Drosophila experiments. P.M, L.Z and J.R

REFERENCES

1. Deng, H., Gao, K. & Jankovic, J. The role of FUS gene variants in neurodegenerative diseases.

Nat. Rev. Neurol. **10**, 337–348 (2014).

2. Vance, C. *et al.* Mutations in FUS, an RNA Processing Protein, Cause Familial Amyotrophic

Lateral Sclerosis Type 6. *Science* **323**, 1208–1211 (2009).

3. Zou, Z.-Y. *et al.* Screening of the FUS gene in familial and sporadic amyotrophic lateral

sclerosis patients of Chinese origin. *Eur. J. Neurol.* **19**, 977–983 (2012).

4. Dini Modigliani, S., Morlando, M., Errichelli, L., Sabatelli, M. & Bozzoni, I. An ALS-associated

mutation in the FUS 3'-UTR disrupts a microRNA–FUS regulatory circuitry. *Nat. Commun.* **5**,

4335 (2014).

5. Sabatelli, M. *et al.* Mutations in the 3' untranslated region of FUS causing FUS overexpression

are associated with amyotrophic lateral sclerosis. *Hum. Mol. Genet.* **22**, 4748–4755 (2013).

6. Ajmone-Cat, M. A. *et al.* Increased FUS levels in astrocytes leads to astrocyte and microglia

activation and neuronal death. *Sci. Rep.* **9**, 4572 (2019).

7. Ju, S. *et al.* A Yeast Model of FUS/TLS-Dependent Cytotoxicity. *PLoS Biol.* **9**, (2011).

8. Chen, Y. *et al.* Expression of human FUS protein in Drosophila leads to progressive

neurodegeneration. *Protein Cell* **2**, 477–486 (2011).

9. Miguel, L. *et al.* Accumulation of insoluble forms of FUS protein correlates with toxicity in

Drosophila. *Neurobiol. Aging* **33**, 1008.e1–15 (2012).

10. Ling, S.-C. *et al.* Overriding FUS autoregulation in mice triggers gain-of-toxic dysfunctions

in RNA metabolism and autophagy-lysosome axis. *eLife* **8**, e40811 (2019).

11. Taylor, J. P., Brown Jr, R. H. & Cleveland, D. W. Decoding ALS: from genes to mechanism.

Nature **539**, 197–206 (2016).

12.   Nolan, M., Talbot, K. & Ansorge, O. Pathogenesis of FUS-associated ALS and FTD: insights

from rodent models. *Acta Neuropathol. Commun.* **4**, (2016).

13.   Samandi, S. *et al.* Deep transcriptome annotation enables the discovery and functional

characterization of cryptic small proteins. *eLife* **6**, e27860 (2017).

14.   Saghatelian, A. & Couso, J. P. Discovery and characterization of smORF-encoded

bioactive polypeptides. *Nat. Chem. Biol.* **11**, 909–916 (2015).

15.   Delcourt, V., Staskevicius, A., Salzet, M., Fournier, I. & Roucou, X. Small Proteins Encoded

by Unannotated ORFs are Rising Stars of the Proteome, Confirming Shortcomings in Genome

Annotations and Current Vision of an mRNA. *Proteomics* (2017)

doi:10.1002/pmic.201700058.

16.   Brunet, M. A., Levesque, S. A., Hunting, D. J., Cohen, A. A. & Roucou, X. Recognition of

the polycistronic nature of human genes is critical to understanding the genotype-phenotype

relationship. *Genome Res.* (2018) doi:10.1101/gr.230938.117.

17.   Brunet, M. A. *et al.* OpenProt: a more comprehensive guide to explore eukaryotic coding

potential and proteomes. *Nucleic Acids Res.* **47**, D403–D410 (2019).

18.   Umoh, M. E. *et al.* A proteomic network approach across the ALS-FTD disease spectrum

resolves clinical phenotypes and genetic vulnerability in human brain. *EMBO Mol. Med.*

e201708202 (2017) doi:10.15252/emmm.201708202.

19.   Collins, M. A., An, J., Hood, B. L., Conrads, T. P. & Bowser, R. P. Label-Free LC–MS/MS

Proteomic Analysis of Cerebrospinal Fluid Identifies Protein/Pathway Alterations and

Candidate Biomarkers for Amyotrophic Lateral Sclerosis. *J. Proteome Res.* **14**, 4486–4501

(2015).

20.   Chick, J. M. *et al.* A mass-tolerant database search identifies a large proportion of

unassigned spectra in shotgun proteomics as modified peptides. *Nat. Biotechnol.* **33**, 743–

749 (2015).

21.   Olexiouk, V., Van Criekinge, W. & Menschaert, G. An update on sORFs.org: a repository

of small ORFs identified by ribosome profiling. *Nucleic Acids Res.* **46**, D497–D502 (2018).

22.   Kovacs, E., Tompa, P., Liliom, K. & Kalmar, L. Dual coding in alternative reading frames

correlates with intrinsic protein disorder. *Proc. Natl. Acad. Sci.* **107**, 5429–5434 (2010).

23.   Pancsa, R. & Tompa, P. Coding Regions of Intrinsic Disorder Accommodate Parallel

Functions. *Trends Biochem. Sci.* **41**, 898–906 (2016).

24.   Zerbino, D. R. *et al.* Ensembl 2018. *Nucleic Acids Res.* **46**, D754–D761 (2018).

25.   Pavesi, A. Asymmetric evolution in viral overlapping genes is a source of selective

protein adaptation. *Virology* **532**, 39–47 (2019).

26.   Michel, A. M., Kiniry, S. J., O'Connor, P. B. F., Mullan, J. P. & Baranov, P. V. GWIPS-viz:

2018 update. *Nucleic Acids Res.* doi:10.1093/nar/gkx790.

27.   Hall, C. E. *et al.* Progressive Motor Neuron Pathology and the Role of Astrocytes in a

Human Stem Cell Model of VCP-Related ALS. *Cell Rep.* **19**, 1739–1749 (2017).

28.   Luisier, R. *et al.* Intron retention and nuclear loss of SFPQ are molecular hallmarks of

ALS. *Nat. Commun.* **9**, 1–15 (2018).

29.   Hart, G. W. & Akimoto, Y. The O-GlcNAc Modification. in *Essentials of Glycobiology* (eds.

Varki, A. et al.) (Cold Spring Harbor Laboratory Press, 2009).

30.   Thompson, J. W., Sorum, A. W. & Hsieh-Wilson, L. C. Deciphering the Functions of O-

GlcNAc Glycosylation in the Brain: The Role of Site-Specific Quantitative O-GlcNAcomics.

Biochemistry **57**, 4010–4018 (2018).

31. Didonna, A. & Benetti, F. Post-translational modifications in neurodegeneration.

Biophys. 2016 Vol 3 Pages 27-49 (2015) doi:10.3934/biophy.2016.1.27.

32. Santos, A. L. & Lindner, A. B. Protein Posttranslational Modifications: Roles in Aging and

Age-Related Disease. *Oxid. Med. Cell. Longev.* **2017**, (2017).

33. Carrì, M. T., D'Ambrosi, N. & Cozzolino, M. Pathways to mitochondrial dysfunction in ALS

pathogenesis. *Biochem. Biophys. Res. Commun.* **483**, 1187–1193 (2017).

34. Deng, J. *et al.* FUS Interacts with HSP60 to Promote Mitochondrial Damage. *PLOS Genet.*

11, e1005357 (2015).

35. Nakaya, T. & Maragkakis, M. Amyotrophic Lateral Sclerosis associated FUS mutation

shortens mitochondria and induces neurotoxicity. *Sci. Rep.* **8**, 15575 (2018).

36. Shan, Y. & Cortopassi, G. Mitochondrial Hspa9/Mortalin regulates erythroid

differentiation via iron-sulfur cluster assembly. *Mitochondrion* **26**, 94–103 (2016).

37. Mellacheruvu, D. *et al.* The CRAPome: a contaminant repository for affinity purification-

mass spectrometry data. *Nat. Methods* **10**, 730–736 (2013).

38. Monahan, Z., Shewmaker, F. & Pandey, U. B. Stress granules at the intersection of

autophagy and ALS. *Brain Res.* **1649**, 189–200 (2016).

39. Marrone, L. *et al.* FUS pathology in ALS is linked to alterations in multiple ALS-associated

proteins and rescued by drugs stimulating autophagy. *Acta Neuropathol. (Berl.)* (2019)

doi:10.1007/s00401-019-01998-x.

40. Aulas, A. & Vande Velde, C. Alterations in stress granule dynamics driven by TDP-43 and

FUS: a link to pathological inclusions in ALS? *Front. Cell. Neurosci.* **9**, (2015).

41. Hergesheimer, R. C. *et al.* The debated toxic role of aggregated TDP-43 in amyotrophic

lateral sclerosis: a resolution in sight? *Brain* **142**, 1176–1194 (2019).

42.   Deng, H.-X. *et al.* FUS-immunoreactive inclusions are a common feature in sporadic and

non-SOD1 familial amyotrophic lateral sclerosis. *Ann. Neurol.* **67**, 739–748 (2010).

43.   Tyzack, G. E. *et al.* Widespread FUS mislocalization is a molecular hallmark of

amyotrophic lateral sclerosis. *Brain* doi:10.1093/brain/awz217.

44.   Farrawell, N. E. *et al.* Distinct partitioning of ALS associated TDP-43, FUS and SOD1

mutants into cellular inclusions. *Sci. Rep.* **5**, 13416 (2015).

45.   Wiesner, D. *et al.* Reversible induction of TDP-43 granules in cortical neurons after

traumatic injury. *Exp. Neurol.* **299**, 15–25 (2018).

46.   Cohen, N. R., Hammans, S. R., Macpherson, J. & Nicoll, J. A. R. New neuropathological

findings in Unverricht–Lundborg disease: neuronal intranuclear and cytoplasmic inclusions.

Acta Neuropathol. (Berl.) **121**, 421–427 (2011).

47.   Shan, X., Chiang, P.-M., Price, D. L. & Wong, P. C. Altered distributions of Gemini of

coiled bodies and mitochondria in motor neurons of TDP-43 transgenic mice. *Proc. Natl.*

Acad. Sci. U. S. A. **107**, 16325–16330 (2010).

48.   Keller, B. A. *et al.* Co-aggregation of RNA binding proteins in ALS spinal motor neurons:

evidence of a common pathogenic mechanism. *Acta Neuropathol. (Berl.)* **124**, 733–747

(2012).

49.   Kryndushkin, D., Wickner, R. B. & Shewmaker, F. FUS/TLS forms cytoplasmic aggregates,

inhibits cell growth and interacts with TDP-43 in a yeast model of amyotrophic lateral

sclerosis. *Protein Cell* **2**, 223–236 (2011).

50.   Lanson, N. A. *et al.* A Drosophila model of FUS-related neurodegeneration reveals

genetic interaction between FUS and TDP-43. *Hum. Mol. Genet.* **20**, 2510–2523 (2011).

51.   Bogaert, E. *et al.* Molecular Dissection of FUS Points at Synergistic Effect of Low-

Complexity Domains in Toxicity. *Cell Rep.* **24**, 529-537.e4 (2018).

52.   Chambers, R. P. *et al.* Nicotine increases lifespan and rescues olfactory and motor

deficits in a Drosophila model of Parkinson's disease. *Behav. Brain Res.* **253**, 95–102 (2013).

53.   Mawhinney, R. M. S. & Staveley, B. E. Expression of GFP can influence aging and climbing

ability in Drosophila. *Genet. Mol. Res. GMR* **10**, 494–505 (2011).

54.   Robles-Murguia, M., Hunt, L. C., Finkelstein, D., Fan, Y. & Demontis, F. Tissue-specific

alteration of gene expression and function by RU486 and the GeneSwitch system. *Npj Aging*

Mech. Dis. **5**, 1–5 (2019).

55.   Scekic-Zahirovic, J. *et al.* Toxic gain of function from mutant FUS protein is crucial to

trigger cell autonomous motor neuron loss. *EMBO J.* **35**, 1077–1097 (2016).

56.   Delcourt, V. *et al.* The Protein Coded by a Short Open Reading Frame, Not by the

Annotated Coding Sequence, Is the Main Gene Product of the Dual-Coding Gene MIEF1. *Mol.*

Cell. Proteomics MCP **17**, 2402–2411 (2018).

57.   Kino, Y. *et al.* FUS/TLS deficiency causes behavioral and pathological abnormalities

distinct from amyotrophic lateral sclerosis. *Acta Neuropathol. Commun.* **3**, (2015).

58.   Hicks, G. G. *et al.* Fus deficiency in mice results in defective B-lymphocyte development

and activation, high levels of chromosomal instability and perinatal death. *Nat. Genet.* **24**,

175–179 (2000).

59.   An, H. *et al.* ALS-linked FUS mutations confer loss and gain of function in the nucleus by

promoting excessive formation of dysfunctional paraspeckles. *Acta Neuropathol. Commun.* **7**,

7 (2019).

60.   Al-Chalabi, A. & Hardiman, O. The epidemiology of ALS: a conspiracy of genes,

environment and time. *Nat. Rev. Neurol.* **9**, nrneurol.2013.203 (2013).

61.   Qamar, S. *et al.* FUS Phase Separation Is Modulated by a Molecular Chaperone and

Methylation of Arginine Cation-π Interactions. *Cell* **173**, 720-734.e15 (2018).

62. Naumann, M. *et al.* Impaired DNA damage response signaling by FUS-NLS mutations

leads to neurodegeneration and FUS aggregate formation. *Nat. Commun.* **9**, 335 (2018).

63. Richards, S. *et al.* Standards and guidelines for the interpretation of sequence variants: a

joint consensus recommendation of the American College of Medical Genetics and Genomics

and the Association for Molecular Pathology. *Genet. Med. Off. J. Am. Coll. Med. Genet.* **17**,

405–424 (2015).

64. Vanderperre, B. *et al.* An overlapping reading frame in the PRNP gene encodes a novel

polypeptide distinct from the prion protein. *FASEB J.* **25**, 2373–2386 (2011).

65. Peng, J.-Y. *et al.* Automatic Morphological Subtyping Reveals New Roles of Caspases in

Mitochondrial Dynamics. *PLOS Comput. Biol.* **7**, e1002212 (2011).

66. Choi, H. *et al.* SAINT: Probabilistic Scoring of Affinity Purification - Mass Spectrometry

Data. *Nat. Methods* **8**, 70–73 (2011).

    LEGENDS

    Main Figures

Figure 1: *FUS* is a bicistronic gene

A, FUS gene bicistronic annotation, with the canonical FUS CDS (in blue, +1 frame) and altFUS CDS
(in green, +2 frame) represented on the FUS canonical transcript (*ENST00000254108* or
NM_004960). Sequence length proportions are respected, the scale bar corresponds to 300
nucleotides. **B**, Genome browser view of FUS gene. The five most abundant transcripts in the brain
are shown in the 'Transcript' track. Transcripts predicted coding by the OpenProt resource are
coloured in blue, and in grey if predicted non-coding. The 'Protein' track contains all predicted
protein products. The known FUS protein (ENSP00000254108) and its isoform
(ENSP00000369594) are coloured in green. The novel predicted altFUS protein (IP_243680) and
its isoform (IP_243691) are coloured in red. **C**, PhyloP nucleotidic conservation scores are
represented in grey across the *FUS* mRNA (ENST00000254108). The noise reduction after FFT (Fast
Fourrier Transformation) is outlined in blue. The average PhyloP score on the bicistronic and the
monocistronic region are represented as dotted red lines. The position of the FUS CDS is
represented by a blue rectangle, and that of altFUS CDS by a green rectangle. **D**, Alignment
(Clustalω) of altFUS protein sequences in Human (*Homo sapiens*), Chimpanzee (Chimp. - *Pan
troglodytes*), Rat (*Rattus norvegicus*), Mouse (*Mus musculus*) and Dog (*Canis lupus familiaris*).
Residues are coloured based on their identity across species, from white (not conserved) to red
(conserved in all species). **E**, Rɪʙo-seq data over the *FUS* gene from the Gwips portal. Initiating
ribosome reads are indicated by blue bars, and elongating ribosomes footprints are indicated by
the blue curve. The graph capture the beginning of the FUS gene with FUS and altFUS methionines
indicated by blue arrows. The genomic positions are indicated relative to the start of exon 1. **F-G**,
Genome browser view of *FUS* gene, centered on altFUS. The 'Transcript' track contains the
beginning of the canonical FUS transcript (*ENST00000254108*) in blue. The 'Protein' track contains
the beginning of the FUS protein (green) and the whole altFUS protein (red). In **F**, the 'Peptide'
track contains all the peptides identified by the OpenProt resource using the classical spectrum-
centric approach. The peptides sequences are indicated and are unique to altFUS or its isoform
(see **Fig EV2B** for an example spectra). In **G**, the 'Peptide' track contains all the peptides identified
by our peptide-centric approach. The peptides indicated matched better to at least one spectra
than any known protein, and are coloured in yellow if they matched better than any known
protein with any PTM (see **Fig EV2C** for an example spectra). The peptides sequences are indicated
and are unique to altFUS or its isoform.

Figure 2: altFUS, a novel endogenous mitochondrial protein

A, Expression of both FUS and altFUS proteins from transfection of the *FUS* cDNA in HEK293 cells
by western blot, and expression of FUS with the monocistronic construct FUS$^{(\emptyset)}$ (representative
image from n=3). The western blot corresponds to a low exposition so that the endogenous FUS

and altFUS are non visible. **B**, AltFUS endogenous expression in HEK293 cells using a siRNA
targeting *FUS* mRNA as negative control and over-expression of altFUS CDS as positive control
(representative image from n=3). **C**, AltFUS (arrow) endogenous expression in Human tissues
(brain, muscles and kidney – 100 ug), in HEK293 and HeLa cultured cells (100 ug) and using the
over-expression of altFUS CDS in HEK293 cells (50 ug) as positive control (representative image
from n=3). The asterisk indicates a protein species detected with the anti-altFUS antibody
specifically in the brain. **D-E**, AltFUS (arrow) endogenous expression in the motor cortex of three
C9orf72 and three sporadic ALS patients (**D**) or in iPSC-derived motor neurons of three lines from
controls and from ALS patients (**E**) (representative image from n=3). The asterisk indicates a
protein species detected with the anti-altFUS antibody specifically in the brain. **F**, Images by
confocal microscopy of altFUS-Flag (green) in HeLa cells, using Tomm20 (red) as a mitochondrial
marker (representative image from n=3, Pearson's correlation r = 0.92). The white scale bar
corresponds to 10 μm. **G**, AltFUS-Flag enrichment in mitochondrial extracts from transfected
HEK293 cells (representative image from n=3) with Hsp70 used as a mitochondrial marker (WCL
= Whole Cell Lysate, Mito = Mitochondrial extract). **H**, AltFUS mitochondrial expression in
transfected HEK293 cells following fractionation (representative image from n=3), with Tubulin as
a cytosolic fraction marker and VDAC as a mitochondrial fraction marker (WCL = Whole Cell Lysate,
Cyto = Cytosol fraction, Mito = mitochondrial fraction). **I**, Endogenous altFUS mitochondrial
expression in HEK293 cells following fractionation (representative image from n=3), with Tubulin
as a cytosolic fraction marker and VDAC as a mitochondrial fraction marker. We used si*FUS*
transfected cells as a negative control and altFUS transfected cells as a positive control for altFUS
expression (WCL = Whole Cell Lysate, Cyto = Cytosol fraction, Mito = mitochondrial fraction). **J**,
Representative images of the mitochondrial network (Tomm20 in red) in mock and altFUS-Flag
(green) transfected HeLa cells (n=3). The white scale bar corresponds to 10μm. **K**, Proportion of
tubules and globules in the mitochondrial network of mock and altFUS-Flag transfected HeLa cells
(see **Fig EV4A**). Quantification was done over a minimum of 100 cells across a technical duplicate
per independent experiments (n=3, i.e. a minimum of 300 cells per biological conditions, p-value
< 0.001, Mann-Whitney U test).

Figure 3: altFUS is involved in mitochondrial dysfunction and autophagy processes

A, Representative traces of TMRE fluorescence measured by flow cytometry in mock transfected
cells and cells overexpressing the bicistronic FUS-R495x or the monocistronic FUS$^{(\emptyset)}$-R495x
constructs (n=4, minimum of 50 000 live cells per independent replicates). Mean fluorescence
intensity of mock transfected cells treated with a decoupling agent, FCCP, is indicated by a grey
dotted line. **B**, Mean TMRE fluorescence intensity measures in mock transfected cells, cells over-
expressing altFUS, FUS, FUS$^{(\emptyset)}$, FUS-R495x or FUS$^{(\emptyset)}$-R495x, or mock transfected cells treated with
FCCP across 3 independent experiments (n=4, also see **Fig EV4D**). Statistical significance is relative
to the mock condition unless otherwise indicated (*** = p value < 0.001, **** = p value < 0.0001,
n.s. = non-significant). **C**, Representative image by stimulated emission depletion microscopy
(STED) of altFUS-Flag (green) localization within mitochondria (Tomm20 marker in red). The white
bar across the mitochondria represents the region of interest quantified in panel **D**. The white
scale bar corresponds to 3μm. **D**, Relative fluorescence histogram for altFUS-Flag and Tomm20
across the region of interest highlighted by a white line on panel **C**. **E**, Scatter plot of the proteins
identified by AP-MS (see **Fig EV5**) indicating their enrichment (Fold-change over control) and their
SAINT probability score. Proteins above the 0.8 threshold are indicated in black, others in grey.
AltFUS is indicated in red (bait) and preys known to regulate the autophagy or the cellular stress

response are indicated in green. **F**, Subcellular localizations of proteins identified by AP-MS from
(see **Fig EV5**). **G**, Enrichment of biological processes in altFUS interacting proteins compared to
the Human mitochondrial proteome (n=2, Fisher's Exact test with FDR < 0.1 %). The number of
proteins identified in each GO term is indicated next to the corresponding bar.

Figure 4: AltFUS is necessary for FUS-associated inhibition of autophagy and accumulation of
FUS/TDP-43 cytoplasmic aggregates

A, Images by confocal microscopy of mCherry-GFP-LC3 signal in HeLa cells across biological
conditions: untreated mock, bafilomycin treated mock, altFUS, FUS, FUS$^{(\emptyset)}$, FUS-R495x and FUS$^{(\emptyset)}$-
R495x (representative images of n=3). The white scale bar corresponds to 10 µm and the zoomed
in region (right panels) is delimited by a white box. **B**, LC3-II accumulation after bafilomycin
treatment from mock, altFUS, FUS, FUS$^{(\emptyset)}$, FUS-R495x, FUS$^{(\emptyset)}$-R495x transfected cells and FUS$^{(\emptyset)}$-
R495x and altFUS co-transfected cells across 3 independent experiments (n=3). The quantification
corresponds to the treated/untreated ratio of LC3-II abundance. Statistical significance is relative
to the mock condition unless otherwise indicated (**** = p value < 0.0001, *** = p value < 0.001,
** = p value < 0.01, n.s. = non-significant). **C**, Images by confocal microscopy of altFUS (FLAG
tagged- white), FUS (GFP tagged - green) and TDP-43 (red) signals in HeLa cells transfected with
the bicistronic GFP-FUS$^{(FLAG)}$-R495x or the monocistronic GFP-FUS$^{(\emptyset-FLAG)}$-R4955x constructs, or co-
transfected with the monocistronic GFP-FUS$^{(\emptyset-FLAG)}$-R495x and altFUS-FLAG constructs
(representative images from n=3). The white scale bar corresponds to 10 µm. **D-E**, Quantification
of FUS cytoplasmic granules, number (**D**) and area (µm^2) (**E**) in cells over-expressing the bicistronic
(+) or monocistronic (-) construct for FUS, FUS-G156E, FUS-R495x, FUS-K510E, FUS-Q519x, FUS-
Q519I-fs527x, FUS-R521C, and FUS-P525L. Statistical comparisons are made between bicistronic
and monocistronic versions of each construct (**** = p value < 0.0001, *** = p value < 0.001, **
= p value < 0.01, * = p value < 0.05, n.s. = non-significant, one-way ANOVA test with Sidak's
multiple comparison).

Figure 5: AltFUS expression is necessary for the full *FUS*-linked ALS phenotype in *Drosophila*

A, Cross-breeding strategy for *Drosophila* generation using the Elav-GeneSwitch-GAL driver as an
inducible expression system specific to the motor neurons. **B**, FUS and altFUS expression in
mCherry (control), altFUS, FUS, FUS$^{(\emptyset)}$, FUS-R495x or FUS$^{(\emptyset)}$-R495x expressing *Drosophila* from the
control F1 and the RU-486 treated F1 (see panel **A**) at 1, 10 or 20 days post-induction
(representative image from n=2). **C-E**, Locomotion assay using the percentage of climbing success
in control and RU-486-treated transgenic *Drosophila* expressing mCherry or altFUS (**C**), the
bicistronic FUS or the monocistronic FUS$^{(\emptyset)}$ (**D**), and the bicistronic FUS-R495x or the
monocistronic FUS$^{(\emptyset)}$-R495x (**E**) at day 1, 10 and 20 post-induction. Statistical comparison were
made between each population (n=4). Indicated significance are between the monocistronic and
the bicistronic transgenic flies of the RU-486-treated population (n.s. = non-significant, * = p value
< 0.05, *** = p value < 0.001).

Figure 6: FUS mutations, synonymous for FUS but missense for altFUS, potentiate TDP-43
cytoplasmic aggregation

A, Graphical representation of FUS synonymous mutations (yellow) found in ALS patients. The
canonical FUS mRNA is represented in dark blue (*ENST00000254108* or *NM_004960*). The FUS
protein coding sequence is indicated in light blue, and the altFUS protein coding sequence is
indicated in green. **B**, Images by confocal microscopy of TDP-43 (yellow) and altFUS (FLAG tagged
- red) in HeLa cells over-expressing GFP-FUS(FLAG) or GFP-FUS(P31L-FLAG)-S44= (representative images
from n=3). The white scale bar corresponds to 10 μm. **C**, Quantification of cells with TDP-43
aggregates in HeLa cells (see panel **B** and **Fig EV8B**). The data are represented as the fold-change
compared to the GFP-FUS(FLAG) expressing cells. Statistical significance is indicated above the bars
(n=3, **** = p value < 0.0001).

909   ***<u>Extended View Figures</u>***

Figure EV1: *FUS* gene protein diversity

A, AltFUS is encoded within an alternative open reading frame (ORF) overlapping the FUS
canonical CDS. When read in the +1 frame, the *FUS* mRNA (here *ENST00000254108*) codes for a
526 amino acid protein (highlighted in blue), named FUS. In the +2 frame, the *FUS* mRNA contains
a second ORF (highlighted in green) that codes for a novel 170 amino acid protein, named altFUS.
The two proteins are not isoforms. **B**, GTEx portal data on *FUS* mRNA expression in the brain and
nerves are shown in blue colored scale (TPM = Transcripts Per Million). Transcripts are identified
with Ensembl accessions (the number after the dot corresponds to the version used in the
analysis) and are quantified across 14 tissues. The five transcripts framed in red share 85 % of all
mRNA expression level. **C**, Table compiling the protein information relayed by Ensembl and
OpenProt resources for the five transcripts highlighted in panel **B**. The FUS protein is highlighted
in blue. The altFUS protein is highlighted in green. D-E, Alignment (Clustalω) of protein sequences
for FUS (ENSP00000254108) and its isoform (ENSP00000369594) in panel **D** (blue) and altFUS
(IP_243680) and its isoform (IP_243691) in panel **E** (green). The residues are coloured based on
their degree of identity. **F**, Heatmap of altFUS protein sequences identity across 84 species (see
Appendix Table 2 and **Source Data 1**). Primates, Rodents and Mammals in general display strong
protein conservation. The sequence identity is colored from blue (0 %) to red (100 %).

Figure EV2: AltFUS is endogenously expressed

A, RIBO-seq data over the mouse *FUS* gene from the Gwips portal (*Mus musculus*). Initiating
ribosome reads are indicated by green bars, and elongating ribosomes footprints are indicated by
the green curve. The graph capture the beginning of the *FUS* gene with FUS and altFUS
methionines indicated by green arrows. The genomic positions are indicated relative to the start
of exon 1. **B**, MS/MS spectra confidently mapped to a peptide unique to altFUS by OpenProt
resource. Peaks are represented by their mass over charge ratios (m/z) and their intensity relative
to the highest (relative intensity). The y ions are coloured in blue, the b ions in red and the
unannotated peaks appear in grey. The charge state and m/z error of the matched peptide are

indicated on the top right corner of the spectra. The original study, file and spectra number are

indicated below the graph. **C**, Comparison of one MS/MS spectra confidently mapped to a peptide

unique to altFUS (left spectra) and its best possible annotation with any known protein with any

post-translational modification (PTM - right spectra). Peaks are represented by their mass over

charge ratios (m/z) and their intensity relative to the highest (relative intensity). The y ions are

coloured in blue, the b ions in red and the unannotated peaks appear in grey. The charge state

and m/z error for each of the matched peptides are indicated on the top right corner of each

panel. The original study, file and spectra number are indicated below the graph. On the top of

each panel is indicated the matched spectra with its corresponding protein and PTMs.

Figure EV3: Tools to explore altFUS expression and roles

A, AltFUS custom antibody strategy, with epitopes highlighted in bold on altFUS sequence (green),

and ELISA test results on both bleeds from each rabbits. B, Nucleotidic mutations on *FUS* mRNA

(*ENST00000254108*) are highlighted in red and mutate all altFUS methionines to threonines

(altFUS_Ø). **C-D**, Protein alignment of altFUS (**C**) and FUS (**D**) proteins from FUS bicistronic

construct and the monocistronic (FUS$^{(\emptyset)}$) construct. The residues are coloured based on sequence

identity (dark blue for identical residues).

Figure EV4: AltFUS is involved in mitochondrial functions

A, MicroP processing of confocal images of mock and altFUS transfected HeLa cells. The nucleus

is coloured in blue, altFUS in green and the mitochondrial network in red (TOMM20). Tubules are

coloured in purple, globules are coloured in green. **B**, Graphical representation of bicistronic

constructs FUS and FUS-R495x (early stop codon highlighted in red), and their monocistronic

equivalent FUS$^{(\emptyset)}$ and FUS$^{(\emptyset)}$-R495x. AltFUS CDS is highlighted in green when present, FUS CDS is

highlighted in blue. **C**, V5-FUS, V5-FUS-R495x and altFUS-FLAG expression from bicistronic or

monocistronic constructs in HEK293 cells (representative image from n=3). **D**, Representative

traces of TMRE fluorescence measured by flow cytometry in mock cells and cells overexpressing

the bicistronic constructs FUS or FUS-R495x (in red) in the top panels, and the monocistronic

constructs, FUS$^{(\emptyset)}$ or FUS$^{(\emptyset)}$-R495x (in blue) in the bottom panels (n=3, minimum of 50 000 live

cells per independent replicates).

Figure EV5: AltFUS mitochondrial interactome

A, Workflow of mitochondrial extraction and size exclusion chromatography (FPLC) on mock or

altFUS-FLAG transfected HEK293 cells. **B**, AltFUS expression in FPLC fractions on transfected

HEK293 cells mitochondrial extracts (representative image from n=2). Every 4 fractions were

loaded, the size scale through the fractions is indicated in red. **C**, Workflow of magnetic beads

FLAG trap on FPLC fractions 8 to 14, from mock or altFUS-FLAG transfected HEK293 cells. **D**, Protein-

protein interaction network of altFUS (bait, in green). Identified proteins (preys) are indicated in

blue. The network was expanded using String database, with a maximum of 2 additional nodes

(grey nodes and grey dotted edges). Interactions (edges) identified by our AP-MS are in black.
Interactions between preys were added using the String database (blue edges).

Figure EV6: AltFUS inhibits the autophagy and potentiate FUS/TDP-43 cytoplasmic aggregates

A, Graphical representation of the mCherry-GFP-LC3 construct to study the autophagic flux. **B**,
Images by confocal microscopy of mCherry-GFP-LC3 and altFUS (FLAG, in blue) signal in HeLa cells
across biological conditions: untreated mock, bafilomycin treated mock, altFUS, FUS, FUS$^{(\emptyset\text{-FLAG})}$,
FUS-R495x and FUS$^{(\emptyset\text{-FLAG})}$-R495x (representative images of n=3). The white scale bar corresponds
to 10 μm and the zoomed in region (right panel) is delimited as a white box (related to **Fig4A**). **C**,
LC3-II expression from untreated (-) or bafilomycin treated (+) cells transfected with mock, altFUS,
FUS, FUS$^{(\emptyset)}$, FUS-R495x, FUS$^{(\emptyset)}$-R495x, or co-transfected with FUS$^{(\emptyset)}$-R495x and altFUS. **D-F**, Images
by confocal microscopy of FLAG (white), GFP (green) and TDP-43 (red) signals in HeLa cells
transfected with the bicistronic constructs (**D**), monocistronic constructs (**E**) or co-transfected
with altFUS-FLAG and the monocistronic constructs (**F**) of 6 ALS-associated mutants: FUS-G156E,
FUS-K510E, FUS-Q519x, FUS-Q519I-fs527x, FUS-R521C, and FUS-P525L (representative images
from n=3). The white scale bar corresponds to 10 μm.

Figure EV7: AltFUS potentiate FUS recruitment to stress granules

A, Images by confocal microscopy of FLAG (white), GFP (green) and TIA-1 (red) signals in HeLa cells
transfected with the bicistronic GFP-FUS$^{(\text{FLAG})}$-R495x or the monocistronic GFP-FUS$^{(\emptyset\text{-FLAG})}$-R495x
constructs, or co-transfected with the monocistronic GFP-FUS$^{(\emptyset\text{-FLAG})}$-R495x and altFUS-FLAG
constructs (representative images from n=3). The white scale bar corresponds to 10 μm. **B-D**,
Images by confocal microscopy of FLAG (white), GFP (green) and TIA-1 (red) signals in HeLa cells
transfected with the bicistronic constructs (**B**), monocistronic constructs (**C**) or co-transfected
with altFUS-FLAG and the monocistronic constructs (**D**) of 6 ALS-associated mutants: FUS-G156E,
FUS-K510E, FUS-Q519x, FUS-Q519I-fs527x, FUS-R521C, and FUS-P525L (representative images
from n=3). The white scale bar corresponds to 10 μm.

1004

Figure EV8: Mitochondrial altFUS mutants potentiates TDP-43 cytoplasmic aggregates

A, Images by confocal microscopy of altFUS (FLAG tagged - green) and mitochondria (TOMM20
marker, red) in HeLa cells over-expressing altFUS-FLAG, altFUS-P31L-FLAG, altFUS-A38V-FLAG,
altFUS-A46V-FLAG or altFUS-R64P-FLAG constructs (representative images from n=3). The white
scale bar corresponds to 10 μm. Deconvolution over a maximum of 30 iterations on the green and
red channels was performed for the zoomed in pictures. The white scale bar corresponds to 3 μm.
B, Images by confocal microscopy of TDP-43 (yellow) and altFUS (FLAG tagged - red) in HeLa cells
over-expressing GFP-FUS$^{(\text{A38V-FLAG})}$-G51=, GFP-FUS$^{(\text{A46V-FLAG})}$-G59= or GFP-FUS$^{(\text{R64P-FLAG})}$-S77=
constructs (representative images from n=3). The white scale bar corresponds to 10 μm.

1014

1015 ***Appendix Tables***

1016

Appendix Table 1: *FUS* gene annotation on OpenProt

OpenProt search results for FUS transcripts *ENST00000254108* and *NM_004960.3*. The OpenProt
version 1.3 was used. Each protein prediction is identified with a unique protein accession
(starting with II_ for novel predicted isoform, and IP_ for predicted alternative proteins). All
predicted alternative proteins are coloured in light blue. The protein localization is relative to the
canonical FUS coding sequence (CDS). The furthered studied protein, named altFUS (IP_243680)
is highlighted in dark blue. The workbook also contains the list of peptides identified in re-analysis
of mass spectrometry experiments using the OpenProt database (version 1.3).

1025

Appendix Table 2: altFUS protein sequences across species

1027 AltFUS conservation across vertebrates. The workbook contains the list of *FUS* transcripts
1028 retrieved from NCBI RefSeq database (transcript accession and nucleotidic sequences); the list of
1029 transcripts containing an altFUS sequence (transcript accession, species, and altFUS protein
1030 sequence); and the matrix of altFUS protein sequence identity between each species.

1031

Appendix Table 3: Peptide-Centric proteomic results for altFUS endogenous expression

1033 Peptide-spectrum matches confidently identified using PepQuery (v1.0) on the TCGA datasets.
1034 The first sheet of the workbook contains the legends for the summary and results tables. The
1035 summary table lists the number of peptides and PSMs in each dataset. The details of results for
1036 each dataset queried is also listed.

1037

Appendix Table 4: Post-translational modification predictions on altFUS sequence

1039 Predictions for post-translational modifications (PTMs) on altFUS sequence. The summary sheets
1040 contains the name of the modification predicted, the target residues, the tool used and the
1041 number of predicted sites with their confidence score. Results for the 2 more abundant PTMs
1042 predicted (phosphorylation and O-Glc-NAcylation) are detailed.

1043

Appendix Table 5: AltFUS interacting proteins

Proteins identified by mass spectrometry following size exclusion chromatography (FPLC) and
FLAG affinity purification on mitochondrial extracts from HEK293 cells over-expressing altFUS-
FLAG. The proteins are listed by their UniProtKB accession alongside their annotated function, the
number of unique peptides supporting the detection and the number of peptide spectrum
matches (PSMs) within each experimental condition. Confidence of the interaction was scored
using the SAINT algorithm. The SAINT score is also reported. Proteins involved in autophagy
related processes are indicated in red. In light greys are proteins removed from the analysis
because lacking protein coverage (unique peptides < 2).

1053

1054 **Appendix Table 6: FUS synonymous mutations in ALS patients and their consequence on altFUS**

1055 *FUS* mutations, synonymous for FUS but missense for altFUS. The first sheet contains mutations
1056 identified from a manually curated literature review. The second sheet retrieves variants in sALS
1057 cases from the ALS Variant Server (http://als.umassmed.edu/). The third sheet retrieves variants
1058 in fALS cases from the ALS Variant Server.

1059

1060 ***Source Data***

1061

1062 **Source Data 1: altFUS protein alignment across species**

1063 Alignment of altFUS protein sequences across all 82 species. The alignment was done using
1064 Clustalω, and coloured based on conservation from white (highly variable residue) to dark blue
1065 (highly conserved residue).

1066

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

FUS = synonymous

5'   3'

AltFUS = missense

B

	AltFUS	TDP-43	Merged
WT			
P31L			

C

Cells with TDP-43 aggregates
(Fold-change to WT)

AltFUS mutations in GFP-FUS$^{(FLAG)}$ backbone

P31L   A38V   A46V   R64P

Figure 6

A

B

C

D

E

F

Figure EV1

Figure EV2

a

MGPT**PPSPGRAIPSRAVSPTDSR**VTVVIASPRTLQAMARAAILLMARARTQAMELSQLPRDMARLAAMAVA**RAPNRLTGSSPPTLA**MASSQLPAAPREVTVAVLRA
AAMGSPRVGATASSLAMVDSSKAMDSSKAIIPLRAMDSRTSTTAAVVVEVEVEVEVTMAKINPP

b

>Mutated_sequence
ACGGGGCCTACCCCACCCAGCCCGGGCAGGGCTATTCCCAGCAGAGCAGTCAGCCCTACGGACAGCCAGAGTTACAGTGGTTATAGC
CAGTCCACGGACACTTCAGGCT**ACG**GCCAGAGCAGCTATTCTTCTT**ACG**GCCAGAGCCAGAACACAGGCT**ACG**GAACTCAGTCAAC
TCCCCAGGGAT**ACG**GCTCGACTGGCGGCT**ACG**GCAGTAGCCAGAGCTCCCAATCGTCTTACGGGCAGCAGTCCTCCTACCCTGGCT**A
CG**GCCAGCAGCCAGCTCCCAGCAGCACCTCGGGAAGTTACGGTAGCAGTTCTCAGAGCAGCAGCT**ACG**GGCAGCCCCAGAGTGGG
AGCTACAGCCAGCAGCCTAGCT**ACG**GTGGACAGCAGCAAAGCT**ACG**GACAGCAGCAAAGCTATAATCCCCCTCAGGGCT**ACG**GACA
GCAGAACCAGTACAACAGCAGCAGTGGTGGTGGAGGTGGAGGTGGAGGTGGAGGTAACT**ACG**GCCAAGATCAATCCTCCATGA

>altFUS_Ø
TGPTPPSPGRAIPSRAVSPTDSRVTVVIASPRTLQA**T**ARAAILL**T**ARARTQA**T**ELSQLPRD**T**ARLAA**T**AVARAPNRLTGSSPPTLA**T**ASSQLP
AAPREVTVAVLRAAA**T**GSPRVGATASSLA**T**VDSSKA**T**DSSKAIIPLRA**T**DSRTSTTAAVVVEVEVEVEVT**T**AKINPP

c

AltFUS	1	MGPTPPSPGRAIPSRAVSPTDSRVTVVIASPRTLQAMARAAILLMARARTQAMELSQLPR	60
AltFUS-Ø	1	TGPTPPSPGRAIPSRAVSPTDSRVTVVIASPRTLQATARAAILLTARARTQATELSQLPR	60
AltFUS	61	DMARLAAMAVARAPNRLTGSSPPTLAMASSQLPAAPREVTVAVLRAAAMGSPRVGATASS	120
AltFUS-Ø	61	DTARLAATAVARAPNRLTGSSPPTLATASSQLPAAPREVTVAVLRAAATGSPRVGATASS	120
AltFUS	121	LAMVDSSKAMDSSKAIIPLRAMDSRTSTTAAVVVEVEVEVEVTMAKINPP	170
AltFUS-Ø	121	LATVDSSKATDSSKAIIPLRATDSRTSTTAAVVVEVEVEVEVTTAKINPP	170

d

FUS	1	MASNDYTQQATQSYGAYPTQPGQGYSQQSSQPYGQQSYSGYSQSTDTSGYGQSSYSSYGQ	60
FUS(Ø)	1	MASNDYTQQATQSYGAYPTQPGQGYSQQSSQPYGQQSYSGYSQSTDTSGYGQSSYSSYGQ	60
FUS	61	SQNTGYGTQSTPQGYGSTGGYGSSQSSQSSYGQQSSYPGYGQQPAPSSTSGSYGSSSQSS	120
FUS(Ø)	61	SQNTGYGTQSTPQGYGSTGGYGSSQSSQSSYGQQSSYPGYGQQPAPSSTSGSYGSSSQSS	120
FUS	121	SYGQPQSGSYSQQPSYGGQQQSYGQQQSYNPPQGYGQQNQYNSSSGGGGGGGGGGNYGQD	180
FUS(Ø)	121	SYGQPQSGSYSQQPSYGGQQQSYGQQQSYNPPQGYGQQNQYNSSSGGGGGGGGGGNYGQD	180
FUS	181	QSSMSSGGGSGGGYGNQDQSGGGSGGYGQQDRGGRGRGGSGGGGGGGGGGGYNRSSGGYE	240
FUS(Ø)	181	QSSMSSGGGSGGGYGNQDQSGGGSGGYGQQDRGGRGRGGSGGGGGGGGGGGYNRSSGGYE	240
FUS	241	PRGRGGGRGGRGGMGGSDRGGFNKFGGPRDQGSRHDSEQDNSDNNTIFVQGLGENVTIES	300
FUS(Ø)	241	PRGRGGGRGGRGGMGGSDRGGFNKFGGPRDQGSRHDSEQDNSDNNTIFVQGLGENVTIES	300
FUS	301	VADYFKQIGIIKTNKKTGQPMINLYTDRETGKLKGEATVSFDDPPSAKAAIDWFDGKEFS	360
FUS(Ø)	301	VADYFKQIGIIKTNKKTGQPMINLYTDRETGKLKGEATVSFDDPPSAKAAIDWFDGKEFS	360
FUS	361	GNPIKVSFATRRADFNRGGGNGRGGRGRGGPMGRGGYGGGGSGGGGRGGFPSGGGGGGGQ	420
FUS(Ø)	361	GNPIKVSFATRRADFNRGGGNGRGGRGRGGPMGRGGYGGGGSGGGGRGGFPSGGGGGGGQ	420
FUS	421	QRAGDWKCPNPTCENMNFSWRNECNQCKAPKPDGPGGGPGGSHMGGNYGDDRRGGRGGYD	480
FUS(Ø)	421	QRAGDWKCPNPTCENMNFSWRNECNQCKAPKPDGPGGGPGGSHMGGNYGDDRRGGRGGYD	480
FUS	481	RGGYRGRGGDRGGFRGGRGGGDRGGFGPGKMDSRGEHRQDRRERPY	526
FUS(Ø)	481	RGGYRGRGGDRGGFRGGRGGGDRGGFGPGKMDSRGEHRQDRRERPY	526

Figure EV3

Figure EV4

Figure EV5

Figure EV6

Figure EV7

Figure EV8

```
                                            10        20        30        40        50        60        70
                                   .    :    .    :    .    :    .    :    .    :    .    :    .    :
Homo sapiens                       MGP---------T---PPS---PG--RA-IPSRAVSPTDSRVTVVIASPRTL-----QAM-ARAAI-L-LMAR
Pan paniscus                       MGP---------T---PPS---PG--RA-IPSRAVSPTDSRVTVVIASPRTL-----QAM-ARAAI-L-LMAR
Gorilla gorilla gorilla            MGP---------T---PPS---PG--RA-IPSRAVSPTDSRVTVVIASPRTL-----QAM-ARAAI-L-LMAR
Pan troglodytes                    MGP---------T---PPS---PG--RA-IPSRAVSPTDSRVTVVIASPRTL-----QAM-ARAAI-L-LMAR
Nomascus leucogenys                MGP---------T---PPS---PG--RA-IPSRAVSPTDSRVTAVIASPRTL-----QAM-ARAAI---LMAR
Pongo abelii                       MGP---------T---PPS---PG--RA-IPSRAVSPTDSRVTALIASPRTL-----QAM-ARAAI-L-LTAR
Chlorocebus sabaeus                MGP---------T---PPS---PG--RA-IPSRAVSPTDSRVIVVTASPRT------QAM-ARAAI-L-LTAR
Rhinopithecus roxellana            MGP---------T---PPS---PG--RA-IPSRAVSPTDSRVTAVTASPRTL-----QAM-ARAAI-L-LTAR
Rhinopithecus bieti                MGP---------T---PPS---PG--RA-IPSRAVSPTDSRVTAVTASPRTL-----QAM-ARAAI-L-LTAR
Macaca mulatta                     MGP---------T---PPS---LG--RA-IPSRAVSPTDSRVTAVTASPRTL-----QAM-ARAAI-L-LTAR
Colobusangolensispalliatus         MGP---------T---PPS---PV--RA-IPSRAVSPTDSRVTAVTASPRTL-----QAM-ARAAI-L-LTAR
Macaca nemestrina                  MGP---------T---PPS---LG--RA-IPSRAVSPTDSRVTAVTASPRTL-----QAM-ARAAI-L-LTAR
Macaca fascicularis                MGP---------T---PPS---LG--RA-FPSRAVSPTDSRVTAVTASPRTL-----QAM-ARAAI-L-LTAR
Mandrillus leucophaeus             MGP---------T---PPS---LG--RA-IPSRAVSPTDSRVTAVTASPRTL-----QAM-ARAAI-L-LTAR
Cercocebus atys                    MGP---------T---PPS---LG--RA-IPSRAVSPTDSRVTAVTASPRTL-----QAM-ARAAI-L-LTAR
Papio anubis                       MGP---------T---PPS---LG--RA-IPSRAVSPTDSRVTAVTASPRTL-----QAM-ARAAI-L-LTAR
Callithrix jacchus                 ----------------------------------------------------------------------
Saimiri boliviensis boliviensis    MGP---------T---PPS---PG--RA-IPSRAVSPTDSRVTAVTASPQTL-----QAM-ARAAMVL-LTAR
Cebus capucinus imitator           MGP---------T---PPS---PG--RA-IPSRAVSPTDSRVTVVMGSLQIL-----QAM-ARAAMVL-LTAR
Sus scrofa                         ---------------------------------------------------------M-ARAAMAV-LMDR
Physeter catodon                   MGP---------T---QPS---LG--RA-IPSRAISPMDSRVTVAMASQRTL-----QAM-ARAAMVV-LMDR
Tursiops truncatus                 MGP---------T---RPS---LG--RA-IPSRAISPMDSRVTVATASQRML-----QAM-ARAAMVV-LTDR
Orcinus orca                       MGP---------T---RPS---LG--RA-IPSRAISPMDSRVTVATASQRML-----QAM-ARAAMVV-LTDR
Balaenoptera acutorostratas        MGP---------T---QPS---LG--RA-IPSRAISPMDSRVTVATASQRTL-----QAM-ARAAMVV-LMDR
Sorex araneus                      ----------------------------------------------------------------------
Aotus nancymaae                    MGP---------T---PRS---PG--RA-IPSRAVSPMDSRVTAVMASPQIL-----QAM-AKAAMVL-LTAR
Trichechus manatus latirostris     -------------------------------------------------------M-ARAAMVL-LMDK
Togninia minima                    MTL--------L--QAR---H---------------------------------------------------
Ceratotherium simum simum          MGP---------T---PPS---LG--RA-TPSRAISPMDSRVTVVTVSQQTL-----QAM-ARAAMVL-LMDR
Condylura cristata                 MGP---------T---PPS---LG--RA-IPSRAISPTDSRVTVVMVV-LMDR----QAM-ARAAMVV-LMDR
Galeopterus variegatus             MGP---------T---PPS---LG--RA-TPSRAISPTDSRVTVVTASQQTL-----QAM-ARAAIVL-LMDR
Odobenus rosmarus divergens        MGP---------T---PPS---LG--RA-IPSRAVSPMDSRVTVVMASQQTL-----QAM-ARAAMVL-LTDR
Tarsius syrichta                   MGP---------T---LPS---LG--RA-TPSRAVSPMGSRVTVVMASQRTL-----QAM-ARAAIVL-LMDR
Microcebus murinus                 MGP---------T---PPS---RG--RA-TRSRAVSPMDSRVTVVTAS-QTL-----QAM-VRAATVV-LMDR
Bison bison bison                  -------------------------------------------------------M-ARAATVV-LMDR
Propithecus coquereli              MGP---------T---PPS---RG--RA-THSRAVSPMDSRVTVVTAS-QTL-----QAM-ARAATVV-LMDR
Bos mutus                          -------------------------------------------------------M-ARAATVV-LMDR
Pteropus vampyrus                  MGL---------T---PPS---LD--RA-IPSRAVSPMDSRVTVVTASQRTL-----LAM-ARAAM-V-LMDR
Pteropus alecto                    MGL---------T---PPS---LD--RA-IPSRAVSPMDSRVTVVTASQRTL-----LAM-ARAAM-V-LMDR
Vicugna pacos                      MGP---------T---LPS---LG--RA-IPSRAISPTDSRVTVVMASHRTL-----QAM-ARAAIVV-LMDR
Rousettus aegyptiacus              MGL---------T---PPS---LD--RA-IPSRAVSPMDSRVTAVTASHRTP-----QAM-ARAAM-V-LMDR
Bos taurus                         -------------------------------------------------------M-ARAATVV-LMDR
Equus caballus                     MGP---------T---PPS---LG--RA-IPSRAINPMDSRVTAVTVSQQTL-----QAM-ARAAMVL-LMDR
Miniopterus natalensis             MGP---------T---PPS---LG--RV-IPSRAISPMDNRVMVATASQGTL-----QAM-ARAPM-V-LMDR
Canis lupus familiaris             MGP---------T---PPS---LG--RA-IPSRAISPMDSRVTVVTVSQQTL-----QAT-ARAAMVL-LMDR
Camelus ferus                      MGP---------T---LPS---LG--RA-IPSRAISPMDSRVTVVMANHRTL-----QAM-ARAAMVV-LMDR
Camelus bactrianus                 MGP---------T---LPS---LG--RA-IPSRAISPMDSRVTVVMANHRTL-----QAM-ARAAMVV-LMDR
Equus przewalskii                  MGP---------T---PPS---LG--RA-IPSRAINPMDSRVTAVTVSQQTL-----QAM-ARAAMVL-LMDR
Equus asinus                       MGP---------T---PPS---LG--RA-IPSRAINPMDSRVTAVTVSQQTL-----QAM-ARAAMVL-LMDR
Eptesicus fuscus                   MGP---------T---PPS---LG--RV-IPSRATSPMDSRVTVATASLGTL-----QAM-ARAPT-V-LTGR
Leptonychotes weddellii            MGP---------T---PPS---LG--RA-IPSRAISPMDSRVTVVTASQQTL-----QAM-ARAAMVL-LTDR
Octodon degus                      MGP---------T---PPS---LG--RA-TPSRAISPMDSRVTVAMASQQTL-----QAM-ARAAMVL-LMDR
Otolemur garnettii                 MGP---------T---PPS---LG--RA-TPNRAISPMDSRVTVVMASQ-IL-----QAM-ARAAMV--LMDR
Myotis davidii                     MGP---------T---PPS---LG--RA-IPSRAASPMDSRVTVATASLGTP-----QAM-ARAPM-V-PTDR
Dipodomys ordii                    MGH---------T---LLS---LG--RA-TPSRAASPMDSRATVVMASQQTL-----QAM-VRAAMVL-LMDR
Jaculus jaculus                    MGP---------I---PHS---LG--RA-TLSRAISLMDSRVTVVMASQQIL-----QDM-ARAAMVL-LMDR
Ochotona princeps                 ---------------------------MASRAMAATGSRTL-----PAM-ARAAMAP-RTDR
Microtus ochrogaster               MGP---------I---LPS---LG--RA-TPNRAINPMASRVTVVMANQLTL-----QDT-ARAAMAL-LMDR
Mus musculus                      -----------------------------MANRVTVVMASQLTL-----QDT-ARAAMVL-LMDR
Peromyscus maniculatus bairdii     MGL---------I---LPS---LG--RD-TPSRAINPMASRVTVVMASQLTL-----RDM-ARAAMGL-LMDR
Erinaceus europaeus                MGR---------T---PPS---LG--RA-IPSRATSPTDSRVTEVTASQRTR-----QAM-ARAAMAL-PMDK
Nannospalax_galili                 MGP---------T---LPS---LG--RV-TPNRAISPMDSRVTVDMASQLTL-----QGM-ARAAMVL-LMDR
Pantholops_hodgsonii               MGP---------T---QLS---LG--KA-TPSRAVPTGSRVTVATASLRTL-----RAM-ARAATVV-LTDR
Drosophila_melanogaster_fusilli    A-----------------------------------MFL-----TGMRACCSLRS-RMDG
Rattus_norvegicus                  MGP---------I---LPS---LG--RA-IPNRAINPMASRVTVVMANQLTL-----QDT-ARAAMVL-LMDR
Capra_hircus                       MGP---------T---QLS---LG--KA-THSRAVNPTGSRVTVATASLRTL-----RAM-ARAATVV-LTDR
Brachypodium_distachyon            MAT--------R--EGT---------------------------------------------------
Ovis_aries                         MGP---------T---QLS---LG--KA-THSRAVNPTGSRVTVATASLQTL-----RAM-ARAATVV-LTDR
Ovis_aries_musimon                 MGP---------T---QLS---LG--KA-THSRAVNPTGSRVTVATASLQTL-----RAM-ARAATVV-LTDR
Oreochromis_niloticus              MVE---------V--A-V---LA--RA-TVSRLARVTASRATEATTRART------------------
Haplochromis_burtoni               MVE---------V--A-V---LA--RA-TVSRLARVTASRATEATTRART------------------
Perkinsus_marinus                  MEV-----------VGRVPGS----------------NSRVVVAV-----------------------
Eucalyptus_grandis                 MASPAAVAPPGTTT---SPA---TT-AA-IPSRRTARIGTPTTPDLELVQVLYLLLVIGM-QNAAIV------
Gallus_gallus                      ----------------------------------------------------------------------
Sarcophilus_harrisii               MGL---------T---LLS---LD--KV-IPNRAVSPMANRVIVDMASQLIP-----QVM-VRAAMAP-PTDK
Ictalurus_punctatus                MVD---------M--EVS---PV--RA-TASRVTEDTARVRTAVPGLTVTD-----TAP-ATARVDT-VPSR
Anolis_carolinensis                MGP---------T---QLH---PH--RV-ILSRLVSRMGSRAMVDMDSQQIL-----QDT-AKVVTAL-LMDR
Trypanosoma                        MQI---------F--VKT---L---------------TGKTI-----ALE-VESSDTI--EN
Xenopus tropicalis                 MGH---------T---PPN---QA--RV-MGSRAASPTASKVTVVTASLQVA-----PRM-DRVATAL-HMVK
Danio rerio                        MIM---------A--KPQAMAMGVMED-SPVKATANLLPRTTVSRAMEDTT-----KAQ-RAAP-LLIIRED
Latimeria chalumnae                MV------------LQ--LG--RV-TLSKPLRRTASRDMVTTASQQTL-----RHM-DRVATLH-PMDK
Xenopus_laevis                     MGH---------I---PPS---QA--RVMVSSRAARPTASKVTVATANLQVA-----PRM-DRVATAL-RMVK
```

```
                                        80          90         100         110         120         130         140
Homo sapiens                   ART - - - - - QAMELSQLPRD-MARLAAMAVAR - - - - - - - - - - - - - - - APNRL - - - - - - - - - - - - - TGSSPP-T
Pan paniscus                   ART - - - - - QAMELSQLPRD-MARLAAMAVAR - - - - - - - - - - - - - - - APNRL - - - - - - - - - - - - - TGSSPP-T
Gorilla gorilla gorilla        ART - - - - - QAMELSQLPRD-MARLAAMAVAR - - - - - - - - - - - - - - - APNRL - - - - - - - - - - - - - TGSSPP-T
Pan troglodytes                ART - - - - - QAMELSQLPRD-MARLAAMAVAR - - - - - - - - - - - - - - - APNRL - - - - - - - - - - - - - TGSSPP-T
Nomascus leucogenys            ART - - - - - QAMELSQLPRD-MARLAAIAVAR - - - - - - - - - - - - - - - APNRL - - - - - - - - - - - - - TGSSPP-T
Pongo abelii                   ART - - - - - QAMELSQLPRD-MARLAAMAVAR - - - - - - - - - - - - - - - APNRL - - - - - - - - - - - - - TGSSPP-T
Chlorocebus sabaeus            ART - - - - - QAMELSQLPRD-MARLAAMAVAR - - - - - - - - - - - - - - - APNLL - - - - - - - - - - - - - MGSSPP-T
Rhinopithecus roxellana        ART - - - - - QAMELSQLPRD-MAQLAAMAVAR - - - - - - - - - - - - - - - APNLL - - - - - - - - - - - - - TGSSPP-T
Rhinopithecus bieti            ARTVSLSQQAMELSQLPRD-MAQLAAMAVAR - - - - - - - - - - - - - - - APNLL - - - - - - - - - - - - - TGSSPP-T
Macaca mulatta                 ARA - - - - - QAMELSQLPRD-MARLAAMAVAR - - - - - - - - - - - - - - - APNLL - - - - - - - - - - - - - MGSSPP-T
Colobusangolensispalliatus     ART - - - - - QAMELSQLPRD-MARLAAMAVAR - - - - - - - - - - - - - - - APNLL - - - - - - - - - - - - - TGSSPP-T
Macaca nemestrina              ARA - - - - - QAMELSQLPRD-MARLAAMAVAR - - - - - - - - - - - - - - - APNLL - - - - - - - - - - - - - MGSSPP-T
Macaca fascicularis            ARA - - - - - QAMELSQLPRD-MARLAAMAVAR - - - - - - - - - - - - - - - APNLL - - - - - - - - - - - - - MGSSPP-T
Mandrillus leucophaeus         ARA - - - - - QAMELSQLPRD-MARLAAMAVAR - - - - - - - - - - - - - - - APNLL - - - - - - - - - - - - - MGSSPP-T
Cercocebus atys                ARA - - - - - QAMELSQLPRG-MARLAAMAVAR - - - - - - - - - - - - - - - APNLL - - - - - - - - - - - - - MGSSPP-T
Papio anubis                   VRAVSLSQQAMELSQLPRD-MARLAAMAVAR - - - - - - - - - - - - - - - APSLL - - - - - - - - - - - - - MGSSPP-T
Callithrix jacchus             - - - - - - - - MALSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - APNHL - - - - - - - - - - - - - TGSSPP-T
Saimiri boliviensis boliviensis PRT - - - - - QAMALSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - APNHL - - - - - - - - - - - - - TGSSPL-T
Cebus capucinus imitator       PRT - - - - - QAMALSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - APNHL - - - - - - - - - - - - - TGSSPL-T
Sus scrofa                     PRT - - - - - QAMAHSQLPRD-MAQLVAMVVAR - - - - - - - - - - - - - - - VLNRL - - - - - - - - - - - - - TGSSLH-T
Physeter catodon               PRT - - - - - QAMAHSQLPRD-MAQLVAMAVAK - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - MGSSPH-T
Tursiops truncatus             PRT - - - - - - AMAHSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - TGSSPH-T
Orcinus orca                   PRT - - - - - QAMAHSQLPRD-MAQLVAMAVAK - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - TGSSPH-T
Balaenoptera acutorostratas    PRT - - - - - QAMAHSQLPRD-MAQLVAMAVAK - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - TGSSPH-T
Sorex araneus                  - - - - - - - - MALSPLPRD-TAQLVAMAAVR - - - - - - - - - - - - - - - VLSRP - - - - - - - - - - - - - MDSSPP-T
Aotus nancymaae                PRT - - - - - QAMALSQLPRD-MAQLVAIVVAR - - - - - - - - - - - - - - - APNHL - - - - - - - - - - - - - TGSSPP-T
Trichechus manatus latirostris PRT - - - - - QVMAPSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - VPNHL - - - - - - - - - - - - - TGSSLR-I
Togninia minima                PRT - - - - - A - PEMA - PPRR - TAR - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - -
Ceratotherium simum simum      PRT - - - - - QAMELSQLPKD-MAQLAAMAVAR - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - TGSSPP-T
Condylura cristata             PRT - - - - - QVMALSQLPRD-MAQLAAMAVAR - - - - - - - - - - - - - - - VLNRL - - - - - - - - - - - - - TGSSPP-T
Galeopterus variegatus         PRT - - - - - QAMALNQLPRD-MAQPVAMAVVR - - - - - - - - - - - - - - - VLNRL - - - - - - - - - - - - - TGSSPP-T
Odobenus rosmarus divergens    PRT - - - - - QAMALSQVPRD-MAQPVAMAVAR - - - - - - - - - - - - - - - VLSHL - - - - - - - - - - - - - MGNSPL-I
Tarsius syrichta               PRT - - - - - QVIALSQLHRD-MAQ-VAMAVAR - - - - - - - - - - - - - - - VPSRL - - - - - - - - - - - - - MGSSPP-T
Microcebus murinus             LRT - - - - - QAIVLSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - VPNRL - - - - - - - - - - - - - MGSSPL-T
Bison bison bison              PRT - - - - - QATAHSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - VLSHL - - - - - - - - - - - - - MGNSPR-I
Propithecus coquereli          PRT - - - - - QAIALRQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - VPSRL - - - - - - - - - - - - - MGSSPP-I
Bos mutus                      PRT - - - - - QATAHSQLPRD-IAQLVAMAVAR - - - - - - - - - - - - - - - VLSHL - - - - - - - - - - - - - MGNSPR-I
Pteropus vampyrus              PRT - - - - - QAITLSQLPRD-MAQQVAMA-AR - - - - - - - - - - - - - - - VLNLL - - - - - - - - - - - - - MGSSPP-T
Pteropus alecto                PRT - - - - - QAITLSQLPRD-MAQRVAMA-AR - - - - - - - - - - - - - - - VLNLL - - - - - - - - - - - - - MGSSPP-T
Vicugna pacos                  LRT - - - - - QAMAHSQLPRV-MAQLVAMAAAK - - - - - - - - - - - - - - - VRSRL - - - - - - - - - - - - - MGSSPH-T
Rousettus aegyptiacus          PRT - - - - - QAIALSQLPRD-MAQRVAMA-AR - - - - - - - - - - - - - - - VLNLL - - - - - - - - - - - - - MGSSPP-T
Bos taurus                     PRT - - - - - - ATAHSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - VLSHL - - - - - - - - - - - - - MGNSPR-T
Equus caballus                 PRT - - - - - QAMALSQLPRD-MAQLVDMAVAR - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - TGSSPP-T
Miniopterus natalensis         PRT - - - - - QAMALSQLPRD-MAQLVAMAVDR - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - MGSSP - -T
Canis lupus familiaris         PRT - - - - - QAMALSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - VLSHL - - - - - - - - - - - - - MGNSPL-I
Camelus ferus                  PRT - - - - - QAMAHSQLPRV-MAQLVATAAAK - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - MGSSPH-T
Camelus bactrianus             PRT - - - - - QAMAHSQLPRV-MAQLVATAAAK - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - MGSSPH-T
Equus przewalskii              PRT - - - - - QAMALSQLPRD-MAQLVDMAVAR - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - TGSSPP-T
Equus asinus                   LRT - - - - - QAMALSQLPRD-MAQLVDMAVAR - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - TGSSPP-T
Eptesicus fuscus               PRT - - - - - QAMAPSQLPRD-MAQPVAMVVAR - - - - - - - - - - - - - - - VLSHL - - - - - - - - - - - - - MGSSP - -T
Leptonychotes weddellii        PRT - - - - - QAMALSQVPRD-MAQPVAMAVAR - - - - - - - - - - - - - - - VLSHL - - - - - - - - - - - - - MGNSPL-I
Octodon degus                  LRT - - - - - QVMAHSQLPRV-MAPLVAMAVAR - - - - - - - - - - - - - - - VPSHL - - - - - - - - - - - - - MVSSPL-I
Otolemur garnettii             PRT - - - - - QVIALSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - VLSHL - - - - - - - - - - - - - MGSSPL-I
Myotis davidii                 PRT - - - - - QAMAPSQLPRV-MAQPVAMVVAR - - - - - - - - - - - - - - - VLSHL - - - - - - - - - - - - - MGSSP - -T
Dipodomys ordii                LRT - - - - - QVMELSQLPRG-TAPLVGMAAAR - - - - - - - - - - - - - - - VPSRL - - - - - - - - - - - - - TGSSLP-T
Jaculus jaculus                PRT - - - - - QVMALSQPPRD-MVPLVDMAVAR - - - - - - - - - - - - - - - VPN-L - - - - - - - - - - - - - MGSSPH-T
Ochotona princeps              HRT - - - - - QAMALSQLPRV-MARLAAMVAAR - - - - - - - - - - - - - - - VPSRL - - - - - - - - - - - - - TDNSPP-T
Microtus ochrogaster           PKT - - - - - QAMELSQLPRD-MVPLEAMAAAR - - - - - - - - - - - - - - - VPNLL - - - - - - - - - - - - - MGSSPP-T
Mus musculus                   PKT - - - - - QAMELSQLPRD-MVPLEAMAAAK - - - - - - - - - - - - - - - VHNLL - - - - - - - - - - - - - MGSSPP-T
Peromyscus maniculatus bairdii PKT - - - - - QAMALSQLPRD-MVPLEAMAAAK - - - - - - - - - - - - - - - VPSLL - - - - - - - - - - - - - MGSSPP-T
Erinaceus europaeus            PRT - - - - - QAMVLSQPPRA-TAPPAAMAAVR - - - - - - - - - - - - - - - VLS-L - - - - - - - - - - - - - TGSSRP-T
Nannospalax_galili             PRT - - - - - QAMALSRLPRD-MVPLEAMAVAK - - - - - - - - - - - - - - - VPNLL - - - - - - - - - - - - - MDNSPP-I
Pantholops_hodgsonii           PRI - - - - - QATAHSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - TGNSLR-T
Drosophila_melanogaster_fusilli PRV - - - - - MPSCYSQTRRM-R - - - - - - - - - - - - - - - - - - - - - PRRW - - - - - - - - - - - - - AGTVNPLG
Rattus_norvegicus              PKT - - - - - QAMELSQHPRD-MVPLEAMAAAK - - - - - - - - - - - - - - - VPSLL - - - - - - - - - - - - - MGSSPP-T
Capra_hircus                   PRI - - - - - QATARSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - TGNSLR-T
Brachypodium_distachyon        - - - - - - - - - - - - - VVQVLEMALYL - - - - - - - - - - - - - - - VVE - - - - - - - - - - - - - - - - - - - - - -
Ovis_aries                     PRI - - - - - QATARSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - TGNSLR-T
Ovis_aries_musimon             PRI - - - - - QATARSQLPRD-MAQLVAMAVAR - - - - - - - - - - - - - - - VLSRL - - - - - - - - - - - - - TGNSLR-T
Oreochromis_niloticus          - - - - - - - - - - - AAPAP - - - TIREAIAATAI - - - - - - - - - - - - - - - LSQVD - - - - - - - - - - - - - TGLHPP-T
Haplochromis_burtoni           - - - - - - - - - - - AAPAP - - - TVREAIAATAI - - - - - - - - - - - - - - - LSQVD - - - - - - - - - - - - - TGLHPP-T
Perkinsus_marinus              - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - -
Eucalyptus_grandis             - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - -
Gallus_gallus                  - - - - - - - - - - - - - - - MAAAEGTAVGR - - - - - - - - - - - - - - - AHSHP - - - - - - - - - - - - - MGRPPP-T
Sarcophilus_harrisii           PRT - - - - - QAMVPNRLPKD-MVQPVAMEVAR - - - - - - - - - - - - - - - VPNLL - - - - - - - - - - - - - MVSSHP-I
Ictalurus_punctatus            PHS - - - - - PT - - - - - ARVVLTA-AAAVSLL - - - - - - - - - - - - - - - RAE - - - - - - - - - - - - - QRTISRPP-T
Anolis_carolinensis            LKT - - - - - LAMALSPHHRP-MAHQQDMAAAK - - - - - - - - - - - - - - - LLRHP - - - - - - - - - - - - - MDSSHH-I
Trypanosoma                    VKA - - - - - KIQDKEGIPPD-QQRLIFAGKQL - - - - - - - - - - - - - - - E - - - - - - - - - - - - - - - - - DGR-T
Xenopus tropicalis             TRV - - - - - LDIVLSQLHKD-MD-LVVMETPN - - - - - - - - - - - - - - - HLSPLMAAGSHSSSPSPLTPATPS-S
Danio rerio                    TAP - - - - - TMDSPSQEDMD-LKPLLKATVSPVSRIALVATVTPASLHQLKA - - - - - - - - - - - EATASSPL-T
Latimeria chalumnae            HKL - - - - - LGMEHSPRPRV-MAP-VAMEALK - - - - - - - - - - - - - - - HHR - - - - - - - - - - - - - TPMDNNLL-T
Xenopus_laevis                 IRA - - - - - LGIVLSQLHNK-GMDLEVMETPN - - - - - - - - - - - - - - - HLSPLMAA - - - - - GS - - -HSSRPL-I
```

```
                                        150       160       170       180       190       200       210       220
                                         .    |    .    |    .    |    .    |    .    |    .    |    .    |    .    |

Homo sapiens                        - - - - - - - LAMASSQLPAAPR-EV-T-VAVLRA- - - - - - - - -AA- MGS-PRVGATAS-SLAMV- - -DSSK- - - - -
Pan paniscus                        - - - - - - - LVMASSQLPAAPR-EV-T-VAVLRA- - - - - - - - -AA- MGS-PRVGATAS-SLAMV- - -DSSK- - - - -
Gorilla gorilla gorilla             - - - - - - - LAMASSQLPAAPR-EV-T-VAVLRA- - - - - - - - -AA- MGS-PRVGATAS-SLAMV- - -DSSL- - - - -
Pan troglodytes                     - - - - - - - LVMASSQLPAAPR-EV-T-VAVLRA- - - - - - - - -AA- MGS-PRVGATAS-SLAMV- - -DSSK- - - - -
Nomascus leucogenys                 - - - - - - - LAMASSHLPAAPR-EV-T-VAVLRP- - - - - - - - -AA- MGS-PRVGAMAS-SLAMV- - -DSSQ- - - - -
Pongo abelii                        - - - - - - - LAMASSQLPAAPR-EV-T-VAVLRP- - - - - - - - -AA- MGS-PRVGATAS-SLAMV- - -DSSK- - - - -
Chlorocebus sabaeus                 - - - - - - - LAMASSQLPAAPR-EV-T-VAVLRP- - - - - - - - -AA- MGS-PRVGATAS-SLAMV- - -DSSK- - - - -
Rhinopithecus roxellana             - - - - - - - LVMASSQLPAAPR-EV-T-VAVLRP- - - - - - - - -AA- MGS-PQVGATAS-SLAMV- - -DSSK- - - - -
Rhinopithecus bieti                 - - - - - - - LVMASSQLPAAPR-EV-T-VAVLRP- - - - - - - - -AA- MGS-PQVGATAS-SLAMV- - -DSSK- - - - -
Macaca mulatta                      - - - - - - - LAMASSQLPAAPR-EV-T-VAVLRP- - - - - - - - -AA- MGS-PRVGATAS-SLAMV- - -DSSK- - - - -
Colobusangolensispalliatus          - - - - - - - LVMASSQLPAAPR-EV-T-VAVLRP- - - - - - - - -AA- MGS-PQVGATAS-SLAMV- - -DSSK- - - - -
Macaca nemestrina                   - - - - - - - LAMASSQLPAAPR-EV-T-VAVLRP- - - - - - - - -AA- MGS-PRVGATAS-SLAMV- - -DSSK- - - - -
Macaca fascicularis                 - - - - - - - LAMASSQLPAAPR-EV-T-VAVLRP- - - - - - - - -AA- MGS-PRVGATAS-SLAMV- - -DSSK- - - - -
Mandrillus leucophaeus              - - - - - - - LAMASSQLPAAPR-EV-T-VAVLRP- - - - - - - - -AA- MGS-RRVGATAS-SLAMV- - -DSSK- - - - -
Cercocebus atys                     - - - - - - - LAMASSQLPAAPR-EV-T-VAVLRP- - - - - - - - -AA- MGS-PRVGATAS-SLAMV- - -DSSK- - - - -
Papio anubis                        - - - - - - - LAMASSQLPAAPR-EV-T-VAVLRP- - - - - - - - -AA- MGS-PQVGATAS-SLAMV- - -DSSK- - - - -
Callithrix jacchus                  - - - - - - - LAMVSSQLPAAPR-EV-T-GAVLRA- - - - - - - - -AA- MGS-PRVAATAS-NLAMV- - -DSSK- - - - -
Saimiri boliviensis boliviensis     - - - - - - - LAMASSQLPAAPR-EV-M-GAVLRA- - - - - - - - -AA-TGSPPRVAATAS-SLAMV- - -DSSK- - - - -
Cebus capucinus imitator            - - - - - - - LAMASSQLPAAPR-EV-T-GAVLRA- - - - - - - - -AA- MGSPPRVAAMAS-SLAMV- - -DSRK- - - - -
Sus scrofa                          - - - - - - - LAMASSQLLAAPR-EV-M-VLVRRA- - - - - - - - -VA- MGS-PRVGAMAS-NLAMV- - -DSSK- - - - -
Physeter catodon                    - - - - - - - LAMASSQLLAAPR-EV-T-VAVLRA- - - - - - - - -AA- MGS-PRVGAMAS-SLAMV- - -DSSK- - - - -
Tursiops truncatus                  - - - - - - - LAMASSQLLAAPR-EV-T-VAVLRA- - - - - - - - -AA- MGS-PRVGAMAS-SLAMV- - -DSSK- - - - -
Orcinus orca                        - - - - - - - LAMASSQLLAAPR-EV-T-VAVLRA- - - - - - - - -AA- MGS-PRVGAMAS-SLAMV- - -DSSK- - - - -
Balaenoptera acutorostratas         - - - - - - - LAMASSQLLAAPR-EV-T-VAVLRA- - - - - - - - -AA- MGS-PRVGAMAS-SLAMV- - -DSLK- - - - -
Sorex araneus                       - - - - - - - LATASSQPRAVPR-EV-M-AAVLRA- - - - - - - - -AA- MGS-PRVAAMAS-SLAMV- - -GSSK- - - - -
Aotus nancymaae                     - - - - - - - LAMASSQLRAAPR-EV-M-GAVLRA- - - - - - - - -AA- MGSPPRVAATAS-SLAMV- - -YSSK- - - - -
Trichechus manatus latirostris      - - - - - - - LAMASSQLPAAPR-EA-M-EVVLRA- - - - - - - - -AA- MGS-PRVEAMAN-SLDMV- - -DSSQ- - - - -
Togninia minima                     - - - - - - - - - -PTPSL- - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - -
Ceratotherium simum simum           - - - - - - - LAMASNRLLAAPR-EV-T-AAVPRA- - - - - - - - -AA- MGS-PRVVAMS-SLAMV- - -DSSQ- - - - -
Condylura cristata                  - - - - - - - LAMASSQLLAAPR-EV-M-GAVLRA- - - - - - - - -AA-TDS-PRVEAMAS-SLATV- - -DSSK- - - - -
Galeopterus variegatus              - - - - - - - LAMVNSQLPAVPR-EV-M-VAVLRA- - - - - - - - -AA- MGS-PQVGAMAN-SLAMV- - -DSSK- - - - -
Odobenus rosmarus divergens         - - - - - - - LAMASSQLLAALR-EV-T-VVVLRA- - - - - - - - -AA-TGS-PRVGAMAS-SLPIV- - -DSSQ- - - - -
Tarsius syrichta                    - - - - - - - LAMVSNQLPAAPR-EV-T-VAVLRA- - - - - - - - -AA- MGS-PRVGAMAN-SLAMV- - -DSRK- - - - -
Microcebus murinus                  - - - - - - - LAMASSRLPAAPR-EV-T-VAVLRA- - - - - - - - -AA-TGS-PRVGAMAN-SLAMV- - -DSNK- - - - -
Bison bison bison                   - - - - - - - LAMASSQLLAAPQ-EV-T-VAVLKA- - - - - - - - -AA- MGS-PRVEATAS-SLAMV- - -GSSK- - - - -
Propithecus coquereli               - - - - - - - LAMANSRLPAAPR-EV-T-VAVPRA- - - - - - - - -AA- MDS-PQAGAMAS-SLAMV- - -DSNK- - - - -
Bos mutus                           - - - - - - - LAMASSQLLAAPQ-EV-T-VAVLKA- - - - - - - - -AA- MGS-PRVEATAS-SLAMV- - -GSSK- - - - -
Pteropus vampyrus                   - - - - - - - LAMASSQPLAAPR-EV-M-VAVLRA- - - - - - - -AATMGS-PRVGVMVS-SLAMV- - -DSSK- - - - -
Pteropus alecto                     - - - - - - - LAMASSQPLAAPR-EV-M-VAVLRA- - - - - - - -AATMGS-PRVGVMVS-SLAMV- - -DSSK- - - - -
Vicugna pacos                       - - - - - - - LAMASSQLLVAPR-EA-T-VVVLRA- - - - - - - - -AA-MAS-PRAGATDS-SLAMV- - -DSSR- - - - -
Rousettus aegyptiacus               - - - - - - - LAMASSQPLAAPR-EV-M-VAVLRA- - - - - - - -AATMGS-PRVGVMVS-SLAMV- - -DNSK- - - - -
Bos taurus                          - - - - - - - LAMASSQLLAAPQ-EV-T-VAVLKA- - - - - - - - -AA- MGS-PRVEATAS-SLAMV- - -GSSK- - - - -
Equus caballus                      - - - - - - - LGMVSSQLLAAPR-EV-M-VAVLRA- - - - - - - - -VA- MDS-PKVEAMAS-SLAMV- - -DSSQ- - - - -
Miniopterus natalensis              - - - - - - - LAMASSQLLAAPL-EV-T-VAVLRA- - - - - - - - -AT- MGS-PRVGAMVS-SLAMV- - -DSSL- - - - -
Canis lupus familiaris              - - - - - - - LAMASSQLLAAPR-EV-M-VVVLRA- - - - - - - - -AA-TGS-PRVGAMAN-SLAIV- - -DSSK- - - - -
Camelus ferus                       - - - - - - - LAMASSQLLVAPR-EA-T-VAVLRA- - - - - - - - -AA-MAS-PRAGATAS-SLAMV- - -DSSR- - - - -
Camelus bactrianus                  - - - - - - - LAMASSQLLVAPR-EA-T-VAVLRA- - - - - - - - -AA-MAS-PRAGATAS-SLAMV- - -DSSR- - - - -
Equus przewalskii                   - - - - - - - LGMVSSQLLAAPR-EV-M-VAVLRA- - - - - - - - -VA- MDS-PKVEAMAS-SLAMV- - -DSSQ- - - - -
Equus asinus                        - - - - - - - LGMVSSQLLAAPR-EV-M-VAVLRA- - - - - - - - -VA- MDS-PKVEAMAS-SLAMV- - -DSSQ- - - - -
Eptesicus fuscus                    - - - - - - - LAMASSQLLAAPR-EV-T-VAVLRA- - - - - - - - -AT- MGS-PRVGAMVS-SLAMV- - -DSSL- - - - -
Leptonychotes weddellii             - - - - - - - LAMASSQLLAAPR-EV-M-VLLRA- - - - - - - - -AA-TGS-PRVGAMAS-SRPIV- - -DSSK- - - - -
Octodon degus                       - - - - - - - LAMVSSQLPAAPR-EV-M-VVVLRA- - - - - - - - -AA- MGS-PRVGAMAN-SLAMV- - -DSNR- - - - -
Otolemur garnettii                  - - - - - - - LAMASSQLLAAPR-EV-T-AAVLRA- - - - - - - - -AA- MDS-PQVGAMAN-SPAMA- - -DSSR- - - - -
Myotis davidii                      - - - - - - - PAMASSPLLAARR-EV-T-VAVLRA- - - - - - - - -AT- MGS-PRVGAMVS-SLAM- - - -DSSL- - - - -
Dipodomys ordii                     - - - - - - - LAMASSQLPAAPL-EV-M-VAVPRA- - - - - - - - -AA- MGS-LKVGATAN-SLAME- - -DSNR- - - - -
Jaculus jaculus                     - - - - - - - LAMASSQLLAAPQ-EV-T-VAVLRA- - - - - - - - -AA- MGS-LRVGAMAN-SLGMA- - -DSNK- - - - -
Ochotona princeps                   - - - - - - - LAMGSSQLPAAPQ-EV-T-AAVLRA- - - - - - - - -AA-TGS-PRAGAMAS-NLAMA- - -DSNK- - - - -
Microtus ochrogaster                - - - - - - - LAMANSQLLAAPQ-EV-M-VAVLRA- - - - - - - - -AA- MGS-PRVEAMAN-SPAMV- - -DSNK- - - - -
Mus musculus                        - - - - - - - LAMANSQLLAAPQ-EV-M-VAVLRA- - - - - - - - -AA- MGN-PRVEVMVN-SLAMV- - -DSNK- - - - -
Peromyscus maniculatus bairdii      - - - - - - - LAMANSQLLAAPR-EV-T-VAVLRA- - - - - - - - -AA- MGS-PRVEAMAN-SQAMV- - -DSNK- - - - -
Erinaceus europaeus                 - - - - - - - PAMASSRLLATPL-EV-T-VAALRA- - - - - - - - -AA- MGS-PRAGATGS-SRATA- - -DSSK- - - - -
Nannospalax_galili                  - - - - - - - LAMANNQLLAALQ-EV-T-VGALRA- - - - - - - - -AA- MGS-PRVEAMAN-SQAMV- - -DSNR- - - - -
Pantholops_hodgsonii                - - - - - - - LAMVSSQLLAAPQ-EV-T-VAVLKA- - - - - - - - -AA- MGS-PRVEAMAS-SLAMV- - -GSSK- - - - -
Drosophila_melanogaster_fusilli     SGTLSYSV- - - - - - -PRPPR-C- - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - -
Rattus_norvegicus                   - - - - - - - PVMANSQLLVAPQ-EV-M-VEVLRA- - - - - - - - -AA- MGS-PRVEAMVN-SLAMV- - -DSNK- - - - -
Capra_hircus                        - - - - - - - LAMASSQLLAAPQ-EV-T-VAVLKA- - - - - - - - -AA- MGS-PRVEAMAS-SLAMV- - -GSSK- - - - -
Brachypodium_distachyon             - - - - - - - - - -MGTGREAPAPH-SV-E-VEV- - - - - - - - - - - - - - -GAVATETVSMV- - -MTQR- - - - -
Ovis_aries                          - - - - - - - LAMASSQLLAAPQ-EV-T-VAVLKA- - - - - - - - -AA- MGS-PRVEAMAS-SLAMV- - -GSSK- - - - -
Ovis_aries_musimon                  - - - - - - - LAMASSQLLAAPQ-EV-T-VAVLKA- - - - - - - - -AA- MGS-PRVEAMAS-SLAMV- - -GSSK- - - - -
Oreochromis_niloticus               - - - - - - - -RAV- - - - - - - - - - - - - - -A-V- - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - -
Haplochromis_burtoni                - - - - - - - -RVV- - - - - - - - - - - - - - - -VVV- - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - -
Perkinsus_marinus                   - - - - - - - - - - - - - - - -R- - - -G-IAIVRA- - - - - - - - -AI- MGV-PRTTTTRV-GVALGA- -VTG- - - -V
Eucalyptus_grandis                  - - - - - - - - - - - - - - - - - - - - - -Q-ILIVRR- - - - - - - - -V- AVNF-LAEGSLVG-S- - - - - - - - - - - - -
Gallus_gallus                       - - - - - - - PATTSRPPPAPPP-EA-M-GAAPRA- - - - - - - -PA-TGS-RPAPATAS-SPHTA- -ASSP- - - - -
Sarcophilus_harrisii                - - - - - - - LATANSQLLAALQ-EA-M-VAVPRA- - - - - - - -VA- MDS-PRVGAMVR-SLVMV- - -DNKV- - - - -
Ictalurus_punctatus                 - - - - - - - RPTASSSRRLLPQLQA-S-PAALG- - - - - - - -HQA-TGS-REEAG- - - - - - - - - - - - - - - - - -
Anolis_carolinensis                 - - - - - - - PATPSSQ-PVPLL-EV-M-APALKP- - - - - - - -PA- MGN-PRVEVMAS-NQAMA- -ASSR- - - - -
Trypanosoma                         - - - - - - - LADYNIQ- - - - -K-ES-TLHLVLRLR- - - - - - -GG-V-ME-PTLEALAK-KYNW- - - -EKK- - - - -
Xenopus_tropicalis                  - - - - - - - LQPVATQEVAAPS-HP-A-TSSSRV- - - - - - -AB- MGS-NQVGVMEV-SRAAMEVNSNP- - - -V
Danio_rerio                         - - - - - - - LAITS-LLLHLPL-VAIA-AALSR- - - - - - - -LA-TVSSSSRVEV- - -G-MEAVV- -VSLEDMGAV
Latimeria_chalumnae                 - - - - - - - LVTASRPPLRPPQ-PA-T-EEVAPS- - - - - - - - -LPA-TGS-PRVEPMDR-ILLTA- -SSSR- - - - -
Xenopus_laevis                      - - - - - - - LATRSSLQPVATQ-EV-M-VGAPSP-LPTSSNKVVD- MDS-RQVGVMEA-SRAAMEVNNNP- - - -V
```

Multiple sequence alignment (positions ~225–293)

Species	Sequence
Homo sapiens	AM-DSS--KA-IIPLRAMDSRTSTTA ... AV-VV
Pan paniscus	AM-DSS--KA-IIPLRAMDSRTSTTA ... AV--V
Gorilla gorilla gorilla	AMVDSS--KA-IIPLRAMDSRTSTTA ... AV--V
Pan troglodytes	AM-DSS--KA-IIPLRAMDSRTSTTA ... AV--V
Nomascus leucogenys	AM-DSS--KA-IIPLRAMDSRTSTTA ... AA--V
Pongo abelii	AM-DSS--KA-IIPLRAMDSRTSTTT ... AA-AV
Chlorocebus sabaeus	AM-DSS--KA-IIPLRAMDSRTSTTA ... AA-VV
Rhinopithecus roxellana	AM-DSS--KA-IIPLRAMDSRTSTTA ... AA-VV
Rhinopithecus bieti	AM-DSS--KA-IIPLRAMDSRTSTTA ... AA-VV
Macaca mulatta	AM-DSS--KA-IIPLRAMDSRTSTTA ... AA-VV
Colobusangolensispalliatus	AM-DSS--KA-IIPLRAMDSRTSTTA ... AA-VV
Macaca nemestrina	AM-DSS--KA-IIPLRVMDSRTSTTA ... AA-VV
Macaca fascicularis	AM-DSS--KA-IIPLRAMDSRTSTTA ... AA-VV
Mandrillus leucophaeus	AM-DSS--KA-IIPLRVMDSRISTTA ... AA-VV
Cercocebus atys	AM-DSS--RA-IIPLRAMDSRISTTA ... AA-VV
Papio anubis	AM-DSS--KA-IIPLRAMDNRIISTTA ... AA-VV
Callithrix jacchus	AT-DSS--KAPIIRLRAMDSRTSTTA ... AA-AAA
Saimiri boliviensis boliviensis	PT-DSS--KAPIIRLRAMDSRTSTTA ... AA-AAA
Cebus capucinus imitator	PT-DSS--KAPIIRLRAMDSRTSTTA ... AA-AAA
Sus scrofa	AM-DSS--KA-IIPLRAMDSRTSTTV ... AV-AV
Physeter catodon	AMVDSS--KA-IIPLRAMGSRTSTTA ... AV-E
Tursiops truncatus	AM-DSS--KA-IIPLRAMGSRTSTTA ... AV-E
Orcinus orca	AM-DSS--KA-IIPLRAMGSRTSTTA ... AV-E
Balaenoptera acutorostrata	AM-DSS--KA-IIPLRATGSRTSTTV ... AV-E
Sorex araneus	AM-DSN--KAPIIHLKVMDSRTSTTA ... AV-EV
Aotus nancymaae	PT-DSS--KAPIIRLRAMDSRTSTTA ... AA-AAA
Trichechus manatus latirostris	AT-DSS--KAPIIPLRAMDSRTSTTV ... AV-GA
Togninia minima	----------------LRTPATL ... AV-----
Ceratotherium simum simum	AM-DSS--QAPIIPLRAMDSRTSTTV ... AV-EV
Condylura cristata	AM-GSS--KAPITPLKAMDSRTSTTV ... AV-EV
Galeopterus variegatus	AT-DSS--KAPIIPLRAMDSRASTTV ... AA-EV
Odobenus rosmarus divergens	AM-DSS--KAPIIPLKAMDSRTSTTV ... AV-
Tarsius syrichta	AT-DSS--KAPITPLRAMDSRTSTTV ... AV-EV
Microcebus murinus	TT-GSS--KAPIIPLRAMDSRASTTV ... AV-EV
Bison bison bison	AM-DSS--RA-ITRLRAMGSRTSTTV ... AV-A
Propithecus coquereli	AT-GSS--KAPIIPLRAMDSRASTTV ... AV--V
Bos mutus	AM-DSS--RA-ITRLRAMGSRTSTTV ... AV-A
Pteropus vampyrus	AM-DSS--KAPIILLRAMDSRTSTTV ... AV-EV
Pteropus alecto	AM-DSS--KAPIILLRAMDSRTSTTL ... AV-EV
Vicugna pacos	AM-DSS--RA-ITPLRAMDSRTSTTA ... AV-
Rousettus aegyptiacus	AM-DSS--KAPIILLRAMDSRTSTTV ... AV-EV
Bos taurus	AM-GSS--RA-ITRLRAMGSRASTTV ... AV-A
Equus caballus	AM-DSS--QAPIILLRAMDSRTSTTA ... AV-EV
Miniopterus natalensis	AM-DSS--KAPITLLRAMDSRTSTTV ... AV-EV
Canis lupus familiaris	AM-DSS--KAPIIPLKAMDSRTSTTV ... AV-EV
Camelus ferus	AM-DSS--RA-ITPLRAMDSRTSTTA ... AV-
Camelus bactrianus	AM-DSS--RA-ITPLRAMDSRTSTTA ... AV-
Equus przewalskii	AM-DSS--QAPIILLRSMDSRTSTTA ... AV-EV
Equus asinus	AM-DSS--QAPIILLRAMDSRTSTTA ... VV-EV
Eptesicus fuscus	AM-DSS--RAPIILLRAMDSRTSTTV ... AV-EV
Leptonychotes weddellii	AM-DSS--KAPIIPLKAMDSRTSTTV ... AV-EV
Octodon degus	AT-DSS--KAPTIPLRAMDSRASTTA ... AV-A
Otolemur garnettii	AT-GNS--KA-IIPLRAMDNRTSTTV ... AV-EV
Myotis davidii	AM-DSS--KVPIILLRAMDSRTSTTV ... AV-EV
Dipodomys ordii	AM-DSS--KALIIPLRAMDNRTSITV ... AV-AE
Jaculus jaculus	AM-DSS--KVPITPPRAMDSRTSTTV ... AV-EV
Ochotona princeps	AM-GSS--KAPITPLRATGSRTSTTA ... AA-EVA
Microtus ochrogaster	AM-DNN--KAPITHLRVMDNRTSTTA ... AV-EV
Mus musculus	AT-DNN--KAPITHLRVMDNRTSTTV ... AV-EV
Peromyscus maniculatus bairdii	AM-DNS--KAPITHLRVMDNRTSTTA ... AV-EV
Erinaceus europaeus	AT-DSS--KALTTPRRATGSRTSTTA ... AA-EVA
Nannospalax_galili	AM-DSN--RAPIIPLRAMDSRTSITV ... AV-EV
Pantholops_hodgsonii	AM-ASS--RA-ITRLRATGSRTSVVV ... VV-AA
Drosophila_melanogaster_fusilli	----------------SRCSTGR ... WT
Rattus_norvegicus	AT-DSN--KAPITHLRVMDNRTSTTA ... AV-EV
Capra_hircus	AM-GSS--RA-ITRLRATGSRTSVVV ... VV-AA
Brachypodium_distachyon	---------------GHLGEIMSATV ... VL
Ovis_aries	AM-GSS--RA-ITRLRATGSRTSTAV ... VV-AA
Ovis_aries_musimon	AM-GSS--RA-ITRLRATGSRTSTAV ... VV-AA
Oreochromis_niloticus	---------------VGMVTQVSPTA ... PE
Haplochromis_burtoni	---------------VGMVTQVSPTA ... PV
Perkinsus_marinus	IT---------EGA---TISL-GINMIIV-VTTTMTTTRTPTTR ... TALTML
Eucalyptus_grandis	---------------ET-LLP ... I-MV
Gallus_gallus	PT-ASS--PH-TARRRATASRASTAT ... AA-P
Sarcophilus_harrisii	PM-DSS--KAPTIPLKDMASRTSITA ... AV-GA
Ictalurus_punctatus	----------------------
Anolis_carolinensis	AP-MGS--SLPTTHLRAMASKVSMAV ... AV-AV
Trypanosoma	VC-RRC--YA-RLPVRASNCRKKAC
Xenopus tropicalis	MGASRVRVA--MAVSS---QALM-DSSSNQA-TASPRAMASSKHSIVVELVDIIRTLLPWAAEAMV-----
Danio rerio	EARAVDLGVAEANTSPLSLEEGLTAH-LLT--TA-PLHHKAMGNKVSTDR
Latimeria chalumnae	----------PS--RART-GSS--RA-TTLSRAMDSSLNMAA ... VA-MVRIHPHL
Xenopus_laevis	MGASRARVATAVSSSS-----QALM-DSSN-QA-TTSLRAIASSKHNIV ... AV-ELV-----

Multiple sequence alignment (positions ~295–365)

Ruler: 300 · 310 · 320 · 330 · 340 · 350 · 360 ·

Species	Sequence (visible residues)
Homo sapiens	----------E------------------VE - - - V - E - V----------------
Pan paniscus	---------------------------AE - - - - - V----------------
Gorilla gorilla gorilla	---------------------------AE - - - - - V----------------
Pan troglodytes	---------------------------AE - - - - - V----------------
Nomascus leucogenys	----------------------------E - - - - - V----------------
Pongo abelii	-------------V-------------VE - - - V - E - V----------------
Chlorocebus sabaeus	---------------------------VE - - - - - V----------------
Rhinopithecus roxellana	---------------------------VE - - - - - V----------------
Rhinopithecus bieti	---------------------------VE - - - - - V----------------
Macaca mulatta	---------------------------VE - - - - - V----------------
Colobusangolensispalliatus	---------------------------VE - - - - - V----------------
Macaca nemestrina	---------------------------VE - - - - - V----------------
Macaca fascicularis	---------------------------VE - - - - - V----------------
Mandrillus leucophaeus	---------------------------VE - - - - - V----------------
Cercocebus atys	---------------------------VE - - - - - V----------------
Papio anubis	---------------------------VE - - - - - - ----------------
Callithrix jacchus	---------------AVG---------VE - - G - G - V----------------
Saimiri boliviensis boliviensis	----------------V----------VE - - - V - G - V----------------
Cebus capucinus imitator	--------------VVV----------EE - - - V - G - V----------------
Sus scrofa	---------------------------VE - - G - V - A----------------
Physeter catodon	---------------------------VE - - G - V - V----------------
Tursiops truncatus	---------------------------VE - - G - V - V----------------
Orcinus orca	---------------------------VE - - G - V - V----------------
Balaenoptera acutorostratas	---------------------------VE - - G - V - V----------------
Sorex araneus	---------------------------VA - - G - V - V----------------
Aotus nancymaae	----------------VVV--------VE - - - V - G - V----------------
Trichechus manatus latirostris	---------------------------AE - - E - V V E----------------
Togninia minima	------------------------------------ - - - V - V----------------
Ceratotherium simum simum	---------------------------AE - - G - V - V----------------
Condylura cristata	---------------------------AE - - G - V - V----------------
Galeopterus variegatus	---------------------------AE - - G - V - V----------------
Odobenus rosmarus divergens	--
Tarsius syrichta	---------------------------AE - - G - A - V----------------
Microcebus murinus	---------------------------AE - - G - A - V----------------
Bison bison bison	---------------------------AE - - G - V - A----------------
Propithecus coquereli	---------------------------AE - - G - V - V----------------
Bos mutus	---------------------------AE - - G - V - A----------------
Pteropus vampyrus	---------------------------AE - - G - V - V----------------
Pteropus alecto	---------------------------AE - - G - V - V----------------
Vicugna pacos	---------------------------VE - - G - V - V----------------
Rousettus aegyptiacus	---------------------------AE - - G - V - V----------------
Bos taurus	---------------------------AE - - G - V - A----------------
Equus caballus	---------------------------AE - - G - V - V----------------
Miniopterus natalensis	---------------------------AE - - G - V - V----------------
Canis lupus familiaris	----------------V----------AE - - G - V - V----------------
Camelus ferus	---------------------------VE - - G - V - V----------------
Camelus bactrianus	---------------------------VE - - G - V - V----------------
Equus przewalskii	---------------------------AE - - G - V - V----------------
Equus asinus	---------------------------AE - - G - V - V----------------
Eptesicus fuscus	---------------------------AE - - G - V - E----------------
Leptonychotes weddellii	----------------V----------AE - - G - V - V----------------
Octodon degus	---------------------------VE - - G - V - V----------------
Otolemur garnettii	----------------V----------VE - - G - V - V----------------
Myotis davidii	----------------------------E - - G - V - E----------------
Dipodomys ordii	---------------------------AE - - G - V - V----------------
Jaculus jaculus	---------------------------VE - - G - V - V----------------
Ochotona princeps	----------------------------E - - G - V - A----------------
Microtus ochrogaster	---------------------------VE - - G - V - V----------------
Mus musculus	---------------------------VE - - G - V - V----------------
Peromyscus maniculatus bairdii	---------------------------VE - - G - V - V----------------
Erinaceus europaeus	----------------------------E - - G - V - A----------------
Nannospalax_galili	----------------------------E - - G - - - V----------------
Pantholops_hodgsonii	---------------------------AE - - G - V - A----------------
Drosophila_melanogaster_fusilli	---------------------------P - - - - R - TM----------------
Rattus_norvegicus	---------------------------VE - - G - V - V----------------
Capra_hircus	---------------------------AE - - G - V - A----------------
Brachypodium_distachyon	---------------------------AE - - - - - ----------------
Ovis_aries	---------------------------AE - - G - V - A----------------
Ovis_aries_musimon	---------------------------AE - - G - V - A----------------
Oreochromis_niloticus	--
Haplochromis_burtoni	--
Perkinsus_marinus	---------------EM------------------ - - I - LV IDS - S----------------
Eucalyptus_grandis	----------------------------EE - - G - V - V----------------
Gallus_gallus	--
Sarcophilus_harrisii	---------------------------VE - - G - V - V----------------
Ictalurus_punctatus	--
Anolis_carolinensis	----------------------------EE - - AAV - VD - - - - - - - - - - -
Trypanosoma	------------------------------------ G - H - C----------------
Xenopus_tropicalis	----------EG-----------------------------L - I RVAMEAK I EEGDVEDLVVVV
Danio_rerio	--
Latimeria chalumnae	VVGMVAKSRVALGV- - - - - - - - - - - - - - SR ITA - - - - - - - - - AE - - - G - E - VDLAVE -DL IE - - - - - - - - -
Xenopus_laevis	----------DI IRTLLPWAAEAMVVDL IRVAMEGKTAEAAGDAEDLVG - E - EEAAL - TAV - - - - - - - - - - -

Multiple sequence alignment (positions ~380–440)

Species	Sequence
Homo sapiens	E V T M
Pan paniscus	E V T M
Gorilla gorilla gorilla	E V T M
Pan troglodytes	E V T M
Nomascus leucogenys	E V T M
Pongo abelii	E V T M
Chlorocebus sabaeus	E V A M
Rhinopithecus roxellana	E V T M
Rhinopithecus bieti	E V T M
Macaca mulatta	E V A M
Colobusangolensispalliatus	E V T M
Macaca nemestrina	E V A M
Macaca fascicularis	E V A M
Mandrillus leucophaeus	E V A M
Cercocebus atys	E V A M
Papio anubis	E V A M
Callithrix jacchus	E V A M
Saimiri boliviensis boliviensis	E V A T
Cebus capucinus imitator	E V T M
Sus scrofa	E V V M
Physeter catodon	E V T M
Tursiops truncatus	E V T M
Orcinus orca	E V T M
Balaenoptera acutorostratas	E V T M
Sorex araneus	E V T M
Aotus nancymaae	E V A M
Trichechus manatus latirostris	E V T M
Togninia minima	P L V P
Ceratotherium simum simum	E V A M
Condylura cristata	E V A M
Galeopterus variegatus	E V A M
Odobenus rosmarus divergens	E V T M
Tarsius syrichta	E V V M
Microcebus murinus	E V T M
Bison bison bison	E V A M
Propithecus coquereli	E V T M
Bos mutus	E V A M
Pteropus vampyrus	E V T T
Pteropus alecto	E V T T
Vicugna pacos	E V A M
Rousettus aegyptiacus	E V T T
Bos taurus	E V A T
Equus caballus	E V A M
Miniopterus natalensis	E V T M
Canis lupus familiaris	E V T M
Camelus ferus	E V A M
Camelus bactrianus	E V A M
Equus przewalskii	E V A M
Equus asinus	E V A M
Eptesicus fuscus	E V T M
Leptonychotes weddellii	E V T M
Octodon degus	E A I T
Otolemur garnettii	E V T M
Myotis davidii	E V A M
Dipodomys ordii	E A T M
Jaculus jaculus	E A T M
Ochotona princeps	E G A M
Microtus ochrogaster	E A T M
Mus musculus	E A T M
Peromyscus maniculatus bairdii	E A T M
Erinaceus europaeus	E V T M
Nannospalax_galili	E A V M
Pantholops_hodgsonii	E V A M
Drosophila_melanogaster_fusilli	S R A V
Rattus_norvegicus	E A T M
Capra_hircus	E V A M
Brachypodium_distachyon	- - - -
Ovis_aries	E V A M
Ovis_aries_musimon	E V A M
Oreochromis_niloticus	A I T A
Haplochromis_burtoni	A I T A
Perkinsus_marinus	N S S S
Eucalyptus_grandis	V G P M
Gallus_gallus	P A A M
Sarcophilus_harrisii	E V A M
Ictalurus_punctatus	- - - -
Anolis_carolinensis	L V A M
Trypanosoma	S N L R
Xenopus tropicalis	V V V E E E Q V A L T A A E G A A V V A E A A W A V V N G E D S V N L V D H V K A A D P P D M K W P N K T T L T T T Q S S C K D L E R M L Q L S Q L
Danio rerio	E - V T
Latimeria chalumnae	- V E E E A - - V G D E E A - - - A V A V A W A V V N V V - - - - - - A S V N L V D P G
Xenopus_laevis	G G A P A - - - V A E E A W A V V N G E D S V N L V D H M K V A D P P E M K W L N K T T L T T T Q S L C K D L E R M L Q L S Q

Multiple sequence alignment (positions ~450–470)

Species	Sequence
Homo sapiens	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Pan paniscus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Gorilla gorilla gorilla	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Pan troglodytes	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Nomascus leucogenys	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Pongo abelii	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Chlorocebus sabaeus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Rhinopithecus roxellana	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Rhinopithecus bieti	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Macaca mulatta	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Colobusangolensispalliatus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Macaca nemestrina	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Macaca fascicularis	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Mandrillus leucophaeus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Cercocebus atys	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Papio anubis	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Callithrix jacchus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Saimiri boliviensis boliviensis	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Cebus capucinus imitator	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Sus scrofa	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Physeter catodon	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Tursiops truncatus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Orcinus orca	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Balaenoptera acutorostratas	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Sorex araneus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Aotus nancymaae	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Trichechus manatus latirostris	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Togninia minima	- - - - - - - - - - - T M M - - - - - - - - - - - - - - M P P
Ceratotherium simum simum	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Condylura cristata	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N P P
Galeopterus variegatus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Odobenus rosmarus divergens	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Tarsius syrichta	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S L P
Microcebus murinus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Bison bison bison	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Propithecus coquereli	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Bos mutus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Pteropus vampyrus	- - - - - - - - - - - V K I - - - - - - - - - - - - - - N P Q
Pteropus alecto	- - - - - - - - - - - V K I - - - - - - - - - - - - - - N P Q
Vicugna pacos	- - - - - - - - - - - A K T - - - - - - - - - - - - - - S P P
Rousettus aegyptiacus	- - - - - - - - - - - V K I - - - - - - - - - - - - - - N P P
Bos taurus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Equus caballus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Miniopterus natalensis	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Canis lupus familiaris	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Camelus ferus	- - - - - - - - - - - A K T - - - - - - - - - - - - - - S P P
Camelus bactrianus	- - - - - - - - - - - A K T - - - - - - - - - - - - - - S P P
Equus przewalskii	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Equus asinus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Eptesicus fuscus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Leptonychotes weddellii	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Octodon degus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Otolemur garnettii	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Myotis davidii	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Dipodomys ordii	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Jaculus jaculus	- - - - - - - - - - - V K I - - - - - - - - - - - - - - S P P
Ochotona princeps	- - - - - - - - - - - A K T - - - - - - - - - - - - - - S P L
Microtus ochrogaster	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Mus musculus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P L
Peromyscus maniculatus bairdii	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Erinaceus europaeus	- - - - - - - - - - - A K T - - - - - - - - - - - - - - S P P
Nannospalax_galili	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Pantholops_hodgsonii	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Drosophila_melanogaster_fusilli	- - - - - - - - - - - G T A - - - - - - - - - - - - - - S R R
Rattus_norvegicus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P L
Capra_hircus	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Brachypodium_distachyon	- A L G
Ovis_aries	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Ovis_aries_musimon	- - - - - - - - - - - A K I - - - - - - - - - - - - - - S P P
Oreochromis_niloticus	- - - - - - - - - - - A T S - - - - - - - - - - - - - - R P A
Haplochromis_burtoni	- - - - - - - - - - - A T S - - - - - - - - - - - - - - R P A
Perkinsus_marinus	- - - - - - - - - - - R T T - - - - - - - - - - - - - - I T P
Eucalyptus_grandis	- - - - - - - - - - - V - - A G V L M V L D W W L G H S E V N P
Gallus_gallus	- - - - - - - - - - - G R S - - - - - - - - - - - - - - R P R
Sarcophilus_harrisii	- - - - - - - - - - - A K I - - - - - - - - - - - - - - N H L
Ictalurus_punctatus	- T A E
Anolis_carolinensis	- - - - - - - - - - - A R I - - - - - - - - - - - - - - S P L
Trypanosoma	- - - - - - - - - - - M K K - - - - - - - - - - - - - - K L R
Xenopus tropicalis	L T T S N K L E S L - R - L T R R L - - - - - - - - - - - G S L
Danio rerio	- - - - - - - - - - - T K I - - - - - - - - - - - - - - A H Q
Latimeria chalumnae	- - - - - - - - - - I K E D Q G - - - - - - - - - - - - M M Q
Xenopus_laevis	- - - - - - - - L R T T S N R L A - - - - - - - - - - - S S R