Head Stabilization During Standing in People with Persisting Symptoms

after Mild Traumatic Brain Injury

Peter C. Fino[1], Tiphanie E Raffegeau[1], Lucy Parrington[2], Robert J Peterka[2,3], Laurie A. King[2,3]

[1] University of Utah, Department of Health, Kinesiology, and Recreation, Salt Lake City, UT

[2] Oregon Health Sciences University, Department of Neurology, Portland, OR

[3] National Center for Rehabilitative Auditory Research, VA Portland Health Care System,

Portland, OR

Corresponding Author:

Peter C. Fino, PhD

250 S 1850 E, Rm 257, Salt Lake City, UT, 84112, USA

peter.fino@utah.edu

801-213-1363

ABSTRACT

Increased postural sway is often observed in people with mild traumatic brain injury

(mTBI), but our understanding of how individuals with mTBI control their head during

stance is limited. The purpose of this study was to determine if people with mTBI exhibit

increased sway at the head compared with healthy controls. People with persisting

symptoms after mTBI ($n = 59$, 41 women) and control participants ($n = 63$, 38 women)

stood quietly for one minute in four conditions: eyes open on a firm surface (EO-firm),

eyes closed on a firm surface (EC-firm), eyes open on a foam pad (EO-foam), and eyes

closed on foam (EC-foam). Inertial sensors at the head, sternum, and lumbar region

collected tri-axial accelerations. Root-mean-square (RMS) accelerations in

anteroposterior (AP) and mediolateral (ML) directions. Sway ratios between the head

and sternum, head and lumbar, and sternum and lumbar region, were compared

between groups. Temporal coupling of anti-phase motion between the upper and lower

body angular accelerations was assessed with magnitude squared coherence and

cross-spectral phase angles. People with mTBI demonstrated greater sway than

controls across conditions and directions. During foam-surface conditions, the control

group, but not the mTBI group, reduced ML sway at their head and trunk relative to their

lumbar by increasing the expression of an anti-phase hip strategy within the frontal

plane. These results are consistent with suggestions of inflexible or inappropriate

postural control in people with mTBI.

KEY TERMS: Posture, Balance, Sway, Sensory Integration, Concussion, Stability

INTRODUCTION

Considerable work has demonstrated the importance of head stabilization in space (i.e., maintaining static head position within a global reference system) for whole-body postural control during stance (Lund and Broberg, 1983;Di Fabio and Emasithi, 1997;Paloski et al., 2006;Hansson et al., 2010) and locomotion (Grossman et al., 1988;Pozzo et al., 1990;Pozzo et al., 1991;Farkhatdinov et al., 2019). Objective assessments of postural sway in people with mild traumatic brain injury (mTBI) often reveal increased sway of the center of mass (CoM) or center of pressure (CoP) compared with healthy controls (Guskiewicz, 2011;Haran et al., 2016;King et al., 2017;Gera et al., 2018), but these results do not inform us about how individuals may stabilize the head in space. While there is some evidence that head stabilization may be impaired while walking in those with mTBI (Sessoms et al., 2015), we are unaware of studies that specifically examine the sway of the head in individuals with mTBI during standing.

The lack of knowledge about head stability during standing in those with mTBI likely stems from two sources: 1) the predominant use of force plates or sensors at the waist to quantify postural sway; and 2) the reliance on single-link inverted pendulum mechanics to understand postural control. Force plates are ideal for quantifying the center-of-pressure (CoP) and reflective of CoM sway measured with inertial sensors at the waist (Mancini et al., 2012). However, the motion of the CoP and CoM does not match the motion of the head (Sakaguchi et al., 1995) because upright postural control is not strictly determined by single-link inverted pendulum mechanics. While a single-link inverted pendulum representation can account for some aspects of body motion

during stance in controlled and unperturbed conditions (Gage et al., 2004), a multi-link

representation of body mechanics provides a larger feasible control manifold (Kuo and

Zajac, 1993;Hsu et al., 2007). Multi-link postural control can also facilitate adaptations to

situational demands (Horak and Nashner, 1986) and sensory integration (Mergner and

Rosemeier, 1998;Stoffregen et al., 2000) through in-phase and anti-phase motion

between the upper and lower body (Creath et al., 2005). Torque generated at the hip

joint can stabilize superior body segments such as the head to optimize the sensitivity of

sensory inputs to vestibular and visual signals (Pozzo et al., 1995;Farkhatdinov et al.,

2019). In addition to the ankle and hip joint, multi-link postural control models can

include a joint at the neck (Nicholas et al., 1998). Neck afferent signals are involved in

perceptual functions of self-motion (Pettorossi and Schieppati, 2014) and reflex

responses such as the cervico-spinal and cervico-ocular reflex, suggesting

somatosensory cervical input converges with vestibular input to mediate multisensory

control of orientation, gaze, and posture (Brandt and Bronstein, 2001). Head

stabilization could therefore be achieved by anti-phase strategies at the hip, neck, or

both.

While numerous studies have investigated postural stability during standing in

people with mTBI (Guskiewicz, 2011;Sosnoff et al., 2011;Buckley et al., 2016;Fino et

al., 2016;Gera et al., 2018), the control of head stability after mTBI remains unclear.

Thus, we conducted an ancillary analysis of a larger study (Fino et al., 2017) to address

two pertinent questions: 1) Do individuals with persisting symptoms after mTBI exhibit

greater sway at the head compared with healthy controls when standing under various

sensory conditions?, and 2) Does the relationship between sway at the lumbar spine

level (approximate CoM), sternum, and head differ between those with persisting

symptoms after mTBI and healthy controls as sensory conditions change? In

accordance with previous reports of larger CoP and CoM sway after mTBI, we

hypothesized that individuals with mTBI would sway more at the head compared with

controls. We did not have a direction-specific *a priori* hypothesis for the second

exploratory question, but we anticipated that segmental sway patterns in people with

mTBI would differ from controls under more challenging sensory conditions. For

example, if normalized to anthropometry, smaller sway at the head relative to the

sternum would indicate more active anti-phase neck control, and smaller sway at the

sternum relative to the lumbar would indicate more active anti-phase hip control.

MATERIALS and METHODS

<u>Participants</u>

Fifty-nine individuals with chronic (>3 months) balance complaints after a

clinically diagnosed mTBI and 63 healthy controls (Table 1) were recruited as part of a

larger study (ClinicalTrials.gov Identifier NCT02748109) in people with chronic mTBI

(detailed elsewhere; see (Fino et al., 2017)). Briefly, participants were recruited through

posters in athletic facilities, physical therapy clinics, hospitals, concussion clinics,

community notice boards and cafes in and around the Portland, Oregon metropolitan

area. Participants were included if they were: >3 months post-mTBI and reporting a

nonzero symptom score on the Sport Concussion Assessment Tool 3

(SCAT3)(Guskiewicz et al., 2013) balance problems symptom, or had no history of brain

injury in the past year for the control group; zero to minimal cognitive deficits as

determined by the Short Blessed Test (Katzman et al., 1983) (score ≤8); and were

between the ages of 18 and 60 years. For the purposes of this study, mTBI was defined

and classified using the criteria from the United States Department of Defense: no CT

scan, or a normal CT scan if obtained; loss of consciousness not exceeding 30 min;

alteration of consciousness/mental state up to 24 h; and post-traumatic amnesia not

exceeding one day (Woodson, 2015). The mechanism of injury was not restricted.

Exclusion criteria for both groups consisted of: musculoskeletal injury in the previous

year that could have seriously impacted gait or balance; current moderate or severe

substance abuse; any past peripheral vestibular or oculomotor pathology from before

their reported mTBI; or refusal to abstain from medications that could impact their

mobility for the duration of testing. Participants were asked to abstain from medications

that could impact their mobility starting 24 hours prior to their first testing date. These

medications included sedatives, benzodiazepines, narcotic pain medications and

alcohol. All recruitment procedures were approved by the Oregon Health & Science

University and Veterans Affairs Portland Health Care System joint institutional review

board and participants provided written informed consent prior to commencing the

study.

127   Table 1

Procedures

Subjects completed demographic and symptom-related questionnaires

(Neurobehavioral Symptom Inventory, NSI(Cicerone and Kalmar, 1995)) and an

assessment of postural control for 60 seconds standing under four different sensory

conditions: (1) firm ground with eyes open (EO-Firm), (2) firm ground with eyes closed

(EC-Firm), (3) foam surface (Airex Balance-pad Elite, Airex AG, Sins, Switzerland) with

eyes open (EO-Foam), (4) and foam surface with eyes closed (EC-Foam)(Shumway-

Cook and Horak, 1986;Fino et al., 2017;Freeman et al., 2018;Gera et al., 2018). For all

conditions, participants were instructed to stand with their feet together with their hands

on their hips, and to hold that position for 60 seconds. The duration of 60 seconds was

chosen to ensure a long enough period to capture reliable sway data after removing

transient effects (Scoppa et al., 2013). If the participant lost balance or deviated from

the starting position (e.g., opening eyes during an eyes closed condition) before the

completion of the 60 seconds, the trial was stopped. Trials that were stopped early were

not repeated, and these trials were excluded from future analysis. Trials were always

presented in the same order: EO-Firm, EC-Firm, EO-Foam, EC-Foam, and rest breaks

were provided between each condition as needed.

<u>Data Analysis</u>

Data were collected at 128 Hz using wearable inertial sensors (Opal, APDM Inc.)

affixed over the lumbar spine region (approximate CoM), sternum, forehead, and

bilaterally on the dorsum of the feet using elastic straps. Each sensor provided tri-axial

acceleration, angular velocity, and magnetometer data. For this ancillary analysis, only

data from the head, sternum, and lumbar sensors were analyzed.

For each condition, the sensors' axes were rotated to align with the global

coordinate system (Moe-Nilssen, 1998). Acceleration data were low-pass filtered using

a phaseless 4th order, 10 Hz low pass Butterworth filter. After filtering, the first and last

10 seconds of each trial were removed to reduce the influence of transient effects from

the start of the trial, movement coincident with the end of the trial, or end distortion from

filtering. The middle 40 seconds of each trial were retained for analysis. For each

condition, accelerations in the transverse plane were extracted for each sensor (Figure

1), and the root mean square (RMS) of AP and ML accelerations were calculated for the

head, sternum, and lumbar sensors. To examine the sway at the head relative to the

sway at the lumbar, the ratio of head sway over lumbar sway was calculated in each

direction for all trials. To account for the effect of pendulum length on linear

accelerations, acceleration sway ratios were normalized based on the height of the

sensor. For example, the head-to-lumbar sway ratio was multiplied by the ratio of the

height of the lumbar sensor over the height of the head sensor,

$$SwayRatio_{Head2Lumbar} = \frac{RMS_{Head}}{RMS_{Lumbar}} \times \frac{h_{Lumbar}}{h_{Head}} \tag{1}$$

where h_{Head} *and* h_{Lumbar} *are* the height of the head and lumbar sensors, respectively and

RMS is the RMS of acceleration at each segment. Similarly, the ratio of head-to-

sternum and sternum-to-lumbar sway was also calculated to determine if the neck

played a significant role in attenuating acceleration between the lumbar and head. As

the height of each sensor was not initially recorded, but the anatomical placement was

consistent across subjects, the height of each sensor was estimated based on the

percentage of total height based on standard anthropometric data: h_{Head} = 0.96, $h_{Sternum}$

= 0.76, h_{Lumbar} = 0.59 (de Leva, 1996). Sway ratios equal to one indicate single-link

sway about the ankle, sway ratios less than 1 indicate anti-phase multi-link sway where

the superior segment is stabilized relative to the inferior segment, and sway ratios

greater than one indicate in-phase multi-link sway where the superior segment sway is

amplified relative to the inferior segment.

177   Figure 1

To examine the temporal coupling between segmental sway, measurements of

AP and ML accelerations at head and lumbar levels were used to calculate angular

accelerations of the upper-body trunk/head segment and the lower-body segment within

the sagittal and frontal planes, respectively. Cross-spectral analysis between these two

segmental angular motions was used to investigate the presence of anti-phase motion

coupling between the segments (i.e. 180° out-of-phase motion) and the strength

(coherence) of the coupling. Comparisons of segmental angular accelerations to assess

temporal coupling between segments is advantageous over the direct use of linear

acceleration measures; a perfectly stabilized head would have zero linear acceleration

thus preventing any meaningful conclusion about the temporal relation between lumbar

and head accelerations, while comparison of segmental angular accelerations would

reveal anti-phase temporal coupling.

The angular acceleration at head and lumbar levels with respect to a rotation axis

at ground level was calculated by dividing the measured transverse accelerations at

these two levels by the respective heights above the ground: $\alpha_{head} = a_{head}/h_{head}$, $\alpha_L =$

a_{lumbar}/h_{lumbar}. The upper body angular acceleration α_{UB} is given by the difference

between α_{head} and α_{lumbar}: $\alpha_{UB} = \alpha_{head} - \alpha_{lumbar}$ and the lower body angular

acceleration $\alpha_{LB} = \alpha_{lumbar}$.

To evaluate the temporal coupling between segments, the magnitude-squared

coherence and phase of the cross power spectral density (CSD) of $\boldsymbol{\alpha}_{UB}$ and $\boldsymbol{\alpha}_{LB}$ was

computed for both AP and ML directions. The coherence and CSD were implemented in

MATLAB using the functions 'mscohere' and 'cpsd', respectively, using a 10 second

Hamming window with 50% overlap.

<u>Statistical Analysis</u>

To examine whether individuals with mTBI and healthy controls had similar rates

of failure (e.g., loss of balance before 60 seconds), Chi-squared proportions tests

compared the number of failed trials in each condition using a 0.05 significance level.

To assess differences between groups across conditions, linear mixed effects

regression models were fit for RMS sway at each body segment in each direction.

Outcomes were first assessed for normality; all segments and directions exhibited

skewed distributions for RMS sway and were therefore log-transformed. Each linear

mixed effects regression model contained fixed effects of group, condition, the

group×condition interaction, and random intercepts to account for the within-subject

correlations. Condition was modeled as a categorical variable with EO-Firm serving as

the reference condition. The control group served as the reference condition for group.

Post-hoc pairwise comparisons were performed using independent sample t-tests and

Cohen's *d* effect sizes to further investigate group×condition interactions for any sway

ratio outcome. As coherence and cross-spectral phase were obtained as

complementary secondary outcomes, descriptive statistics were extracted and

presented, but no statistical comparisons were performed on these outcomes. All

significance values were corrected for multiple comparisons using a false discovery rate

(FDR) correction (Benjamini and Hochberg, 1995) and a significance level of 0.05. All

statistical analysis was performed in MATLAB (r2018a, The MathWorks Inc.).

RESULTS

<u>Task Completion</u>

No participants in the control group failed to complete the 60 seconds during any

of the trials. Comparatively, mTBI subjects failed to complete a total of 23 trials (EO-

firm: $n = 2$ (3%), $p = 0.447$, EC-firm: $n = 8$ (14%), $p = 0.008$, EO-foam: $n = 3$ (5%), $p =$

0.220, EC-foam: $n = 10$ (17%), $p = 0.002$). All subjects who failed the EO-firm trial failed

all other trials. Subjects who failed the EC-firm trial or the EO-foam trial also failed the

EC-foam trial. Compared with controls, more mTBI subjects failed the EC trials, but did

not have significantly higher failure rates during EO trials.

<u>Head Sway</u>

Participants with mTBI swayed more at the head for all conditions and directions

relative to controls (Tables 2,3). Relative to the baseline EO-Firm condition, RMS head

sway increased in both AP and ML directions for every condition in control participants

(main condition effects, Table 3). Compared with the between-condition changes in the

control group, the mTBI group exhibited larger increases in head sway in both directions

and all conditions except for EO-Foam in the ML direction (group×condition interactions,

Table 3).

238   Table 2

239   Table 3

<u>Sternum Sway</u>

Participants with mTBI exhibited greater RMS sway at the sternum for all

conditions in the ML direction only (Tables 2,3). Relative to the baseline EO-Firm

condition, RMS sternum sway increased in both AP and ML directions for all other

conditions in control participants except for EC-Firm in the AP direction (condition

effects, Table 3). Compared with the between-condition changes in the control group,

the mTBI group had larger increases in AP sternum sway in all conditions and larger

changes in ML sternum sway in the EC-Foam condition (group×condition interactions,

Table 3).

Lumbar Sway

Similar to sway at the head, participants with mTBI exhibited greater RMS sway

at the lumbar for all conditions and directions (Tables 2,3). Relative to the baseline EO-

Firm condition, RMS lumbar sway increased in both directions for all other conditions in

control participants except for EC-Firm in the AP direction (condition effects, Table 3).

Compared to the between-condition changes relative to the baseline EO-Firm condition

in the control group, the mTBI group had larger increases in AP lumbar sway in EC-Firm

and EC-Foam conditions (group×condition interactions, Table 3).

Head to Lumbar Sway Ratio

In foam conditions, control participants decreased the ML head-to-lumbar sway

ratio relative to the baseline EO-Firm condition (condition effects, Table 4, Figure 2F).

Comparatively, the mTBI group exhibited little change in the ML head-to-lumbar sway

ratio in foam conditions (EO-Foam, EC-Foam) relative to the EO-Firm condition

(group×condition interactions, Table 4, Figure 2F). Post-hoc t-tests revealed the mTBI

group had significantly greater sway ratios than the control group during foam

conditions (EO-Foam $d = 0.74$, *adjusted p* < 0.001; EC-Foam $d = 0.91$, *adjusted p* <

0.001). No effects of group, condition, or interactions were detected for the head-to-

lumbar sway ratio in the AP direction.

267   Table 4

268   Head to Sternum Sway Ratio

269   The head-to-sternum sway ratio did not differ by group or condition in the AP

270   direction. In the ML direction, the head-to-sternum sway ratio decreased in the EC-

271   Foam condition only (condition effects, Table 4); no other group, condition, or

272   group×condition effects were found.

273   Sternum to Lumbar Sway Ratio

274   In foam conditions, control participants decreased the ML sternum-to-lumbar

sway ratio relative to the baseline EO-Firm condition (condition effects, Table 4).

Comparatively, the mTBI group exhibited increased ML sternum-to-lumbar sway ratio in

foam conditions (EO-Foam, EC-Foam) (group×condition interactions, Table 4). Post-hoc

t-tests revealed the mTBI group had significantly greater sway ratios than the control

group during foam conditions (EO-Foam $d = 1.05$, *adjusted p* < 0.001; EC-Foam $d =$

1.15, *adjusted p* < 0.001). No effects of group, condition, or interactions were found for

the sternum-to-lumbar sway ratio in the AP direction.

282   Figure 2

283   Temporal Coupling of Segments

284   The coherence and cross-spectral phase between α_{UB} and α_{LB} in the frequency

band most associated with postural sway (<1 Hz) were consistent with the sway ratio

resultss. In foam conditions relative to firm conditions, control participants exhibited

stronger anti-phase control of the upper and lower body segments in the frontal plane,

characterized by greater coherence between frontal plane α_{UB} and α_{LB} and a cross-

spectral phase angle near 180° (Figure 3). Despite having similar coherence and cross-

spectral phase to controls in firm-ground conditions, the mTBI group exhibited a weaker

anti-phase relationship in foam conditions, characterized by lower coherence and cross-

spectral phase angles that were farther from 180°. In both groups, a peak in coherence

was observed between 0-1 Hz, centered between 0.35-0.50 Hz, during the foam

conditions. Consistent with the sway ratio results, no group differences were observed

in coherence or cross-spectral phase within the sagittal plane (Figure 4).

296   Figure 3

297   Figure 4

298   **DISCUSSION**

299   Our primary objective was to compare postural sway at the head between

individuals with mTBI and healthy control subjects under varying sensory conditions.

Additionally, we sought to examine the relationship between stabilization of the head

and stabilization of inferior segments (sternum and lumbar) to understand potential

postural control strategies. We found individuals with mTBI swayed more at the head,

sternum, and lumbar spine, extending previous results indicating postural sway of the

CoM or CoP is greater in people with mTBI (Guskiewicz, 2011;King et al., 2017;Gera et

al., 2018). Between-group differences in RMS sway, however, were greatest at the

head location. Additionally, the largest between-group differences were observed in the

ML direction, agreeing with previous studies suggesting the importance of

accelerometry-derived ML sway in differentiating acute, symptomatic mTBI from healthy

controls (King et al., 2017).

We found healthy control subjects tended to stabilize the head in space when

standing on foam, while individuals with mTBI did not exhibit the same degree of head

stabilization. Fujisawa et al. (2005) represented body mechanics as a double inverted

pendulum that allowed joint rotations at the ankles and hip while assuming a rigid head-

on-trunk coupling and showed that balance control shifted toward greater use of hip

joint torques compared to ankle joint torques when the surface was narrowed. With

double pendulum body mechanics, proprioceptors at the ankle determine the ankle

angle, proprioceptors at the hip determines the hip angle, and the visual and vestibular

sensors determine the angle of the head and trunk relative to vertical (Figure 5). At any

point in time, the horizontal position of the CoM, x_{CoM} can then be described using one

of three equations

$$x_{CoM} = h_{Hip} \sin(\theta) + \left(h_{CoM} - h_{Hip}\right) \sin(\theta + \beta) \qquad (3)$$

322   where h_{Hip} is the height of the hip, h_{CoM} is the height of the CoM, θ is the ankle

323   angle, and β is the hip angle,

$$x_{CoM} = h_{Hip} \sin(\theta) + \left(h_{CoM} - h_{Hip}\right) \sin(\gamma) \qquad (4)$$

324   where γ is the angle of the upper body relative to vertical, or

$$x_{CoM} = h_{Hip} \sin(\beta + \gamma) + \left(h_{CoM} - h_{Hip}\right) \sin(\gamma) . \qquad (5)$$

325   Adding a noise term onto each angle estimate to reflect sensory noise common

326   and dividing by h_{CoM} yields

$$\sin \Phi = H \sin(\theta + \epsilon_\theta) + (1 - H) \sin(\theta + \epsilon_\theta + \beta + \epsilon_\beta) \tag{6}$$

$$\sin \Phi = H \sin(\theta + \epsilon_\theta) + (1 - H) \sin(\gamma + \epsilon_\gamma) \tag{7}$$

$$\sin \Phi = H \sin(\gamma + \epsilon_\gamma - \beta + \epsilon_\beta) + (1 - H) \sin(\gamma + \epsilon_\gamma) \tag{8}$$

where Φ is the angle with respect to vertical of a single inverted pendulum connecting

the CoM to the ankle joint and H is the ratio $\frac{h_{Hip}}{h_{CoM}}$. Since H is routinely close to 0.9, the

above equations are largely dominated by the first term. Therefore, the state estimation

of the position of the center of mass will be influenced by the sensory error at the ankle

ϵ_α in Eq.6 and 7, or the combined sensory error at the hip and head $\epsilon_\beta + \epsilon_\gamma$ in Eq. 8.

The head-in-space stabilization demonstrated by controls may have been achieved by a

strategy that places greater reliance on hip proprioceptors and visual/vestibular

information at the head to estimate body position (Eq. 8) when ankle proprioceptive

cues are unreliable in the foam surface condition. The reduced head stabilization in the

mTBI group may be due to an inflexible postural control strategy that was unable to shift

toward increased use of hip and head sensory information to reduce the use of

unreliable ankle proprioceptive cues. This interpretation agrees with studies using

entropic measures of sway; individuals with mTBI, at various times since injury, have

greater regularity and less complexity in their postural sway that are typically indicative

of inflexible postural control (Cavanaugh et al., 2006;Sosnoff et al., 2011;Buckley et al.,

2016;Fino et al., 2016;Schmidt et al., 2018). However, the postural control strategy

exhibited by the mTBI group may not be inflexible, but rather inappropriate given the

task demands; a perception of instability or fear of falling related to self-reported

symptoms may increase muscle co-contraction at the hips thereby decreasing head

stabilization through anti-phase motion about the hip. Future electromyography studies

focused on the muscles that span the hip joint are needed to directly answer this

question.

Our results also suggest the head-in-space stabilization exhibited by control

subjects during foam conditions was achieved primarily through an anti-phase

relationship between the angular acceleration of the upper and lower body due to

control at the hip. While ankle and hip strategies are traditionally defined in the sagittal

plane (Horak and Nashner, 1986), similar control mechanisms consisting of in-phase

(i.e., ankle strategy) and anti-phase (i.e., hip strategy) motion of the upper and lower

body exist in the frontal plane (Zhang et al., 2007;Goodworth and Peterka, 2010) - we

only detected this head stabilization and increased hip strategy during foam conditions

in the ML direction. The ML sternum-to-lumbar sway ratio was less than one for all

conditions in healthy controls, with lower sway ratios in foam conditions (see Figure 2E).

Additionally, the coherence between upper and lower body frontal-plane angular

accelerations increased in the foam conditions while maintaining a cross-spectral phase

angle near 180°, indicating a stronger anti-phase relationship in the dominant frequency

band of postural sway (Zatsiorsky and Duarte, 1999). These results are consistent with

the idea that a continuum exists between ankle and hip strategies (Creath et al., 2005).

Specific to the narrow stance width employed here, ML balance control can shift from

an in-phase ankle strategy to an anti-phase hip strategy based on the amplitude of

surface perturbations (Goodworth and Peterka, 2010). Thus, the increased expression

of an anti-phase hip strategy in the ML direction when standing on the foam surface in

healthy control subjects may be specific to the narrow stance width used here.

The ML head-to-sternum ratios were greater than or equal to one for all

conditions, indicating a lack of a strong anti-phase relationship in both groups (see

Figure 2D). These results suggest that the stiffness of the neck joint increased in the

EC-Foam trial, but no active stabilization caused by anti-phase motion was observed at

the neck. Pozzo et al. noted head stabilization in the frontal plane during complex

balance tasks tended to occur at the neck, but a rigid head-trunk unit with minimal

actuation at the neck compensated for small oscillations (Pozzo et al., 1995). Our

results suggesting both groups increased the head-to-trunk coupling during EC-Foam

conditions are consistent with the small oscillation condition reported by Pozzo et al.

(1995). It is possible that complex balance tasks with larger frontal plane angular

displacements than those examined here may elicit compensation utilizing head-on-

trunk stabilization that reveals differences between individuals with mTBI and healthy

controls. Nevertheless, the lack of a between-group difference in the head-to-sternum

sway ratio suggests that neck problems in mTBI subjects, such as neck pain or

whiplash, did not influence our results.

Several limitations should be acknowledged and considered. First, participants

were tested with their shoes on. Traditionally, postural sway would be assessed

barefoot to remove any confounding factor of footwear. However, out of concern for

safety and other aspects of the larger study, balance assessments were performed with

shoes. Therefore, some heterogeneity in our data may stem from differences in

footwear. Second, the actual height of the sensors was not recorded at the time of data

collection. As the anatomical placement of the sensors was consistent across subjects,

we relied on anthropometric ratios to estimate the height of the sensors. However, as

anthropometry varies across individuals, the use of ratios likely introduced some degree

of random error into our results. Nevertheless, our findings were robust and it is unlikely

this error was biased based on group. Finally, we excluded trials in which individuals

lost balance before the end of the trial, removing 17% of our mTBI subjects from the

EC-Foam condition and potentially leading to a bias in our sample. However, excluding

incomplete trials was a conservative approach. A loss of balance creates extremely

large RMS values. Since these trials only occurred in our mTBI groups, removing these

trials with a loss of balance likely led to smaller sway in the mTBI group and may have

biased our results toward smaller between-group effects. Our results should therefore

be interpreted as the conservative estimate of the difference in head stability between

mTBI and controls.

In conclusion, we found that people with persisting balance complaints following

mTBI exhibited greater sway, quantified with inertial sensors on the head, sternum, and

lumbar, compared with healthy control subjects. Further, control subjects reduced the

sway at the head relative to sway at the lumbar when standing on foam by increasing

the strength of the anti-phase relationship of upper body and lower body angular

acceleration. Conversely, those with mTBI did not change postural control strategies

between conditions. The attenuation of sway predominantly occurred over the trunk

segment, suggestive of a shift towards increasing expression of an anti-phase hip

strategy that was supported by coherence and cross-spectral phase results.

Speculatively, these results suggest healthy control subjects are more capable, or more

willing, to shift control into head-centric postural control using hip torque, while people

with persisting symptoms after mTBI may continue to use the same strategy regardless

of sensory information.

ACKNOWLEDGEMENTS

The authors would like to acknowledge the research assistants who contributed

to the subject recruitment, enrollment, and data collection throughout this process. This

work was supported by the Assistant Secretary of Defense for Health Affairs under

Award No. W81XWH-15-1-0620. Additionally, PCF was supported by the Eunice

Kennedy Shiver National Institute of Child Health & Human Development of the National

Institutes of Health under Award Number K12HD073945. Opinions, interpretations,

conclusions and recommendations are those of the authors and are not necessarily

endorsed by the Department of Defense or the official views of the National Institutes of

Health.

References:

Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate - a Practical and
Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B-*
Statistical Methodology 57**,** 289-300.
Brandt, T., and Bronstein, A.M. (2001). Cervical vertigo. *J Neurol Neurosurg Psychiatry* 71**,** 8-
12.
Buckley, T.A., Oldham, J.R., and Caccese, J.B. (2016). Postural control deficits identify
lingering post-concussion neurological deficits. *J Sport Health Sci* 5**,** 61-69.
Cavanaugh, J.T., Guskiewicz, K.M., Giuliani, C., Marshall, S., Mercer, V.S., and Stergiou, N.
(2006). Recovery of postural control after cerebral concussion: new insights using
approximate entropy. *J Athl Train* 41**,** 305-313.
Cicerone, K.D., and Kalmar, K. (1995). Persistent postconcussion syndrome: The structure of
subjective complaints after mild traumatic brain injury. *The Journal of Head Trauma*
Rehabilitation 10**,** 1-17.
Creath, R., Kiemel, T., Horak, F., Peterka, R., and Jeka, J. (2005). A unified view of quiet and
perturbed stance: simultaneous co-existing excitable modes. *Neurosci Lett* 377**,** 75-80.
De Leva, P. (1996). Adjustments to Zatsiorsky-Seluyanov's segment inertia parameters. *J*
Biomech 29**,** 1223-1230.
Di Fabio, R.P., and Emasithi, A. (1997). Aging and the mechanisms underlying head and
postural control during voluntary motion. *Phys Ther* 77**,** 458-475.
Farkhatdinov, I., Michalska, H., Berthoz, A., and Hayward, V. (2019). Gravito-inertial ambiguity
resolved through head stabilization. *Proc Math Phys Eng Sci* 475**,** 20180010.
Fino, P.C., Nussbaum, M.A., and Brolinson, P.G. (2016). Decreased high-frequency center-of-
pressure complexity in recently concussed asymptomatic athletes. *Gait Posture* 50**,** 69-
74.
Fino, P.C., Peterka, R.J., Hullar, T.E., Murchison, C., Horak, F.B., Chesnutt, J.C., and King, L.A.
(2017). Assessment and rehabilitation of central sensory impairments for balance in
mTBI using auditory biofeedback: a randomized clinical trial. *BMC Neurol* 17**,** 41.
Freeman, L., Gera, G., Horak, F.B., Blackinton, M.T., Besch, M., and King, L. (2018).
Instrumented Test of Sensory Integration for Balance: A Validation Study. *J Geriatr*
Phys Ther 41**,** 77-84.
Fujisawa, N., Masuda, T., Inaoka, Y., Fukuoka, H., Ishida, A., and Minamitani, H. (2005).
Human standing posture control system depending on adopted strategies. *Med Biol Eng*
Comput 43**,** 107-114.
Gage, W.H., Winter, D.A., Frank, J.S., and Adkin, A.L. (2004). Kinematic and kinetic validity of
the inverted pendulum model in quiet standing. *Gait Posture* 19**,** 124-132.
Gera, G., Chesnutt, J., Mancini, M., Horak, F.B., and King, L.A. (2018). Inertial Sensor-Based
Assessment of Central Sensory Integration for Balance After Mild Traumatic Brain
Injury. *Mil Med* 183**,** 327-332.
Goodworth, A.D., and Peterka, R.J. (2010). Influence of stance width on frontal plane postural
dynamics and coordination in human balance control. *J Neurophysiol* 104**,** 1103-1118.
Grossman, G.E., Leigh, R.J., Abel, L.A., Lanska, D.J., and Thurston, S.E. (1988). Frequency and
velocity of rotational head perturbations during locomotion. *Exp Brain Res* 70**,** 470-476.
Guskiewicz, K.M. (2011). Balance assessment in the management of sport-related concussion.
Clin Sports Med 30**,** 89-102, ix.

Guskiewicz, K.M., Register-Mihalik, J., Mccrory, P., Mccrea, M., Johnston, K., Makdissi, M.,
Dvorak, J., Davis, G., and Meeuwisse, W. (2013). Evidence-based approach to revising
the SCAT2: introducing the SCAT3. *Br J Sports Med* 47**,** 289-293.
Hansson, E.E., Beckman, A., and Hakansson, A. (2010). Effect of vision, proprioception, and the
position of the vestibular organ on postural sway. *Acta Otolaryngol* 130**,** 1358-1363.
Haran, F.J., Slaboda, J.C., King, L.A., Wright, W.G., Houlihan, D., and Norris, J.N. (2016).
Sensitivity of the Balance Error Scoring System and the Sensory Organization Test in the
Combat Environment. *J Neurotrauma* 33**,** 705-711.
Horak, F.B., and Nashner, L.M. (1986). Central programming of postural movements: adaptation
to altered support-surface configurations. *J Neurophysiol* 55**,** 1369-1381.
Hsu, W.L., Scholz, J.P., Schoner, G., Jeka, J.J., and Kiemel, T. (2007). Control and estimation of
posture during quiet stance depends on multijoint coordination. *J Neurophysiol* 97**,** 3024-
3035.
Katzman, R., Brown, T., Fuld, P., Peck, A., Schechter, R., and Schimmel, H. (1983). Validation
of a short Orientation-Memory-Concentration Test of cognitive impairment. *Am J
Psychiatry* 140**,** 734-739.
King, L.A., Mancini, M., Fino, P.C., Chesnutt, J., Swanson, C.W., Markwardt, S., and Chapman,
J.C. (2017). Sensor-Based Balance Measures Outperform Modified Balance Error
Scoring System in Identifying Acute Concussion. *Ann Biomed Eng* 45**,** 2135-2145.
Kuo, A.D., and Zajac, F.E. (1993). Human standing posture: multi-joint movement strategies
based on biomechanical constraints. *Prog Brain Res* 97**,** 349-358.
Lund, S., and Broberg, C. (1983). Effects of different head positions on postural sway in man
induced by a reproducible vestibular error signal. *Acta Physiol Scand* 117**,** 307-309.
Mancini, M., Salarian, A., Carlson-Kuhta, P., Zampieri, C., King, L., Chiari, L., and Horak, F.B.
(2012). ISway: a sensitive, valid and reliable measure of postural control. *J Neuroeng
Rehabil* 9**,** 59.
Mergner, T., and Rosemeier, T. (1998). Interaction of vestibular, somatosensory and visual
signals for postural control and motion perception under terrestrial and microgravity
conditions--a conceptual model. *Brain Res Brain Res Rev* 28**,** 118-135.
Moe-Nilssen, R. (1998). A new method for evaluating motor control in gait under real-life
environmental conditions. Part 1: The instrument. *Clin Biomech (Bristol, Avon)* 13**,** 320-
327.
Nicholas, S.C., Doxey-Gasway, D.D., and Paloski, W.H. (1998). A link-segment model of
upright human posture for analysis of head-trunk coordination. *J Vestib Res* 8**,** 187-200.
Paloski, W.H., Wood, S.J., Feiveson, A.H., Black, F.O., Hwang, E.Y., and Reschke, M.F.
(2006). Destabilization of human balance control by static and dynamic head tilts. *Gait
Posture* 23**,** 315-323.
Pettorossi, V.E., and Schieppati, M. (2014). Neck proprioception shapes body orientation and
perception of motion. *Front Hum Neurosci* 8**,** 895.
Pozzo, T., Berthoz, A., and Lefort, L. (1990). Head stabilization during various locomotor tasks
in humans. I. Normal subjects. *Exp Brain Res* 82**,** 97-106.
Pozzo, T., Berthoz, A., Lefort, L., and Vitte, E. (1991). Head stabilization during various
locomotor tasks in humans. II. Patients with bilateral peripheral vestibular deficits. *Exp
Brain Res* 85**,** 208-217.
Pozzo, T., Levik, Y., and Berthoz, A. (1995). Head and trunk movements in the frontal plane
during complex dynamic equilibrium tasks in humans. *Exp Brain Res* 106**,** 327-338.

Sakaguchi, M., Taguchi, K., Ishiyama, T., Netsu, K., and Sato, K. (1995). Relationship between
head sway and center of foot pressure sway. *Auris Nasus Larynx* 22**,** 151-157.
Schmidt, J.D., Terry, D.P., Ko, J., Newell, K.M., and Miller, L.S. (2018). Balance Regularity
Among Former High School Football Players With or Without a History of Concussion. *J*
Athl Train 53**,** 109-114.
Scoppa, F., Capra, R., Gallamini, M., and Shiffer, R. (2013). Clinical stabilometry
standardization: basic definitions--acquisition interval--sampling frequency. *Gait Posture*
37**,** 290-292.
Sessoms, P.H., Gottshall, K.R., Sturdy, J., and Viirre, E. (2015). Head Stabilization
Measurements as a Potential Evaluation Tool for Comparison of Persons With TBI and
Vestibular Dysfunction With Healthy Controls. *Military Medicine* 180**,** 135-142.
Shumway-Cook, A., and Horak, F.B. (1986). Assessing the influence of sensory interaction of
balance. Suggestion from the field. *Phys Ther* 66**,** 1548-1550.
Sosnoff, J.J., Broglio, S.P., Shin, S., and Ferrara, M.S. (2011). Previous mild traumatic brain
injury and postural-control dynamics. *J Athl Train* 46**,** 85-91.
Stoffregen, T.A., Hettinger, L.J., Haas, M.W., Roe, M.M., and Smart, L.J. (2000). Postural
instability and motion sickness in a fixed-based flight simulator. *Hum Factors* 42**,** 458-
469.
Woodson, J. (2015). "Traumatic Brain Injury: Updated Definition and Reporting", (ed.) D.O.
Defense. (Washington, DC).
Zatsiorsky, V.M., and Duarte, M. (1999). Instant equilibrium point and its migration in standing
tasks: rambling and trembling components of the stabilogram. *Motor control* 3**,** 28-38.
Zhang, Y., Kiemel, T., and Jeka, J. (2007). The influence of sensory information on two-
component coordination during quiet stance. *Gait Posture* 26**,** 263-271.

Table 1. Demographic information for each group of participants. Data are presented as
mean (SD) unless otherwise noted.

	mTBI	Control
N	59	63
Age (yrs)	39.3 (10.9)	37.7 (12.8)
Gender (F / M)	41/18	38/25
Height (cm)	170.8 (9.4)	171.5 (9.5)
Mass (kg)	79.8 (19.7)	75.2 (19.0)
NSI Symptom Score*	37.0 (23.0)	3.0 (5.0)
One or more lifetime mTBI (n)	59	5
Time Since Most Recent mTBI (yrs)*	1.12 (1.95)	13.28 (11.10)

* Shown as median (interquartile range)
NSI: Neurobehavioral Symptom Inventory

554 **Table 2.** Descriptive statistics of median and inter-quartile range (IQR) for RMS sway at
555 the head, trunk, and lumbar spine in mTBI and control groups.

Location	Head				Trunk				Lumbar			
Group	mTBI		Control		mTBI		Control		mTBI		Control	
Statistic	*Median*	*IQR*	*Median*	*IQR*	*Median*	*IQR*	*Median*	*IQR*	*Median*	*IQR*	*Median*	*IQR*
AP direction												
EO-Firm	0.18	0.12	0.12	0.08	0.13	0.06	0.11	0.06	0.11	0.05	0.08	0.05
EC-Firm	0.24	0.22	0.14	0.08	0.15	0.12	0.11	0.04	0.13	0.10	0.09	0.05
EO-Foam	0.24	0.17	0.15	0.07	0.16	0.10	0.13	0.06	0.14	0.08	0.11	0.03
EC-Foam	0.40	0.23	0.22	0.12	0.27	0.23	0.17	0.07	0.22	0.16	0.13	0.06
ML direction												
EO-Firm	0.12	0.08	0.08	0.03	0.08	0.06	0.05	0.02	0.06	0.04	0.05	0.02
EC-Firm	0.15	0.13	0.08	0.03	0.09	0.10	0.06	0.02	0.08	0.06	0.05	0.02
EO-Foam	0.18	0.19	0.12	0.05	0.14	0.12	0.09	0.04	0.12	0.08	0.10	0.03
EC-Foam	0.32	0.22	0.17	0.05	0.25	0.21	0.14	0.05	0.20	0.12	0.15	0.06

556 *Note: All values given in units of m/s^2*
557

558 **Table 3.** Mixed effect regression model fit and parameters for head, sternum, and lumbar accelerations.

Fixed-Effects	Head β	SE	Corrected p	Sternum β	SE	Corrected p	Lumbar β	SE	Corrected p
AP direction									
Intercept	-2.07	0.06	**< 0.001**	-2.26	0.06	**< 0.001**	-2.49	0.05	**< 0.001**
Condition: EO-Firm	**Ref.**	--	--	**Ref.**	--	--	**Ref.**	--	--
Condition: EC-Firm	0.15	0.06	**0.019**	0.09	0.06	0.191	0.04	0.05	0.463
Condition: EO-Foam	0.17	0.06	**0.006**	0.19	0.06	**0.003**	0.26	0.05	**< 0.001**
Condition: EC-Foam	0.59	0.06	**< 0.001**	0.53	0.06	**< 0.001**	0.54	0.05	**< 0.001**
Group: Control	**Ref.**	--	--	**Ref.**	--	--	**Ref.**	--	--
Group: mTBI	0.29	0.09	**0.002**	0.19	0.09	0.051	0.25	0.07	**0.002**
Condition: EO-Firm × Group	**Ref.**	--	--	**Ref.**	--	--	**Ref.**	--	--
Condition: EC-Firm × Group	0.25	0.08	**0.008**	0.19	0.08	**0.041**	0.25	0.08	**0.004**
Condition: EO-Foam × Group	0.21	0.08	**0.019**	0.18	0.08	**0.049**	0.11	0.08	0.244
Condition: EC-Foam × Group	0.27	0.08	**0.004**	0.30	0.08	**0.001**	0.21	0.08	**0.018**
ML direction									
Intercept	-2.55	0.05	**< 0.001**	-2.99	0.06	**< 0.001**	-3.04	0.05	**< 0.001**
Condition: EO-Firm	**Ref.**	--	--	**Ref.**	--	--	**Ref.**	--	--
Condition: EC-Firm	0.08	0.04	**0.048**	0.18	0.05	**0.003**	0.14	0.04	**0.003**
Condition: EO-Foam	0.44	0.04	**< 0.001**	0.54	0.05	**< 0.001**	0.67	0.04	**< 0.001**
Condition: EC-Foam	0.81	0.04	**< 0.001**	1.02	0.05	**< 0.001**	1.15	0.04	**< 0.001**
Group: Control	**Ref.**	--	--	**Ref.**	--	--	**Ref.**	--	--
Group: mTBI	0.43	0.08	**< 0.001**	0.48	0.09	**< 0.001**	0.38	0.07	**< 0.001**
Condition: EO-Firm × Group	**Ref.**	--	--	**Ref.**	--	--	**Ref.**	--	--
Condition: EC-Firm × Group	0.17	0.06	**0.016**	0.13	0.08	0.189	0.08	0.06	0.302
Condition: EO-Foam × Group	0.09	0.06	0.259	0.12	0.08	0.199	-0.09	0.06	0.220
Condition: EC-Foam × Group	0.16	0.07	**0.025**	0.19	0.08	**0.033**	-0.03	0.07	0.722

559 *Note:* Degrees of freedom = 456 for all effects. Significance denoted by bolded *p*-value.
560 Parameter estimates: slope estimate (*β*), standard error (*SE*), and corrected *p*-value (*p*).

Table 4. Mixed effect regression model fit and parameters for head to lumbar, head to sternum, and sternum to lumbar sway ratios.

Fixed-Effects	Head to Lumbar			Head to Sternum			Sternum to Lumbar		
	β	SE	Corrected p	β	SE	Corrected p	β	SE	Corrected p
AP direction									
Intercept	-0.07	0.05	0.299	-0.04	0.05	0.414	-0.02	0.04	0.653
Condition: EO-Firm	**Ref.**	--	--	**Ref.**	--	--	**Ref.**	--	--
Condition: EC-Firm	0.10	0.06	0.168	0.06	0.05	0.354	0.04	0.05	0.475
Condition: EO-Foam	-0.09	0.06	0.253	-0.01	0.05	0.775	-0.07	0.05	0.262
Condition: EC-Foam	0.05	0.06	0.434	0.06	0.05	0.325	-0.01	0.05	0.921
Group: Control	**Ref.**	--	--	**Ref.**	--	--	**Ref.**	--	--
Group: mTBI	0.05	0.08	0.617	0.11	0.07	0.144	-0.07	0.06	0.392
Condition: EO-Firm × Group	**Ref.**	--	--	**Ref.**	--	--	**Ref.**	--	--
Condition: EC-Firm × Group	-0.01	0.09	0.949	0.05	0.08	0.597	-0.05	0.08	0.519
Condition: EO-Foam × Group	0.10	0.09	0.339	0.03	0.08	0.761	0.07	0.07	0.394
Condition: EC-Foam × Group	0.05	0.09	0.615	-0.04	0.08	0.648	0.10	0.08	0.298
ML direction									
Intercept	0.01	0.04	0.911	0.21	0.03	**< 0.001**	-0.20	0.04	**< 0.001**
Condition: EO-Firm	**Ref.**	--	--	**Ref.**	--	--	**Ref.**	--	--
Condition: EC-Firm	-0.05	0.04	0.336	-0.08	0.03	0.081	0.03	0.04	0.461
Condition: EO-Foam	-0.23	0.04	**< 0.001**	-0.10	0.04	0.019	-0.13	0.04	**0.003**
Condition: EC-Foam	-0.34	0.04	**< 0.001**	-0.21	0.04	**< 0.001**	-0.13	0.04	**0.003**
Group: Control	**Ref.**	--	--	**Ref.**	--	--	**Ref.**	--	--
Group: mTBI	0.05	0.06	0.475	-0.05	0.05	0.368	0.10	0.05	0.092
Condition: EO-Firm × Group	**Ref.**	--	--	**Ref.**	--	--	**Ref.**	--	--
Condition: EC-Firm × Group	0.09	0.06	0.179	0.05	0.06	0.463	0.04	0.06	0.479
Condition: EO-Foam × Group	0.18	0.06	**0.005**	-0.03	0.06	0.633	0.21	0.06	**< 0.001**
Condition: EC-Foam × Group	0.19	0.06	**0.004**	-0.03	0.06	0.696	0.22	0.06	**< 0.001**

Note: Degrees of freedom = 456 for all effects. Significance denoted by bolded *p*-value.
Parameter estimates: slope estimate (*β*), standard error (*SE*), and corrected *p*-value (*p*).

Figure 1. Representation of acceleration sway within the transverse plane at the head, sternum, and lumbar sensor locations for a single subject during EO-Firm (left) and EC-Foam (right) conditions.

Figure 2. Sway ratios for each direction and condition. A-C) Sway ratios for the head-to-sternum, sternum-to-lumbar, and head-to-lumbar sway ratios in the anteroposterior direction. D-F) Sway ratios for the head-to-sternum, sternum-to-lumbar, and head-to-lumbar sway ratios in the mediolateral direction. Across all figures, the horizontal green line indicates a sway ratio equal to one. In E and F, * indicates significant between-group differences based on post-hoc pairwise t-tests.

Figure 3. Mean and standard deviation envelopes (±one SD) of magnitude-squared coherence (top) and cross-spectral phase (bottom) between frontal plane α_{UB} and α_{LB} for both mTBI (red) and Control (black) groups in each condition. In the foam conditions, the Control group exhibited greater coherence between 0-1 Hz, and maintained a stronger anti-phase relationship between α_{UB} and α_{LB} compared to the mTBI group. The stronger anti-phase relationship are consistent with increased head-in-space stabilization in the foam conditions.

Figure 4. Mean and standard deviation envelopes (±one SD) of magnitude-squared coherence (top) and cross-spectral phase (bottom) between sagittal plane α_{UB} and α_{LB} for both mTBI (red) and Control (black) groups in each condition. No group differences are noted.

Figure 5. Schematic of two-link postural control state estimation. Motion in the sagittal or frontal plane can be determined from the combined height of the hip joint, h_{hip}, height of the center of mass, h_{CoM}, and two of the three angles: the angle of the ankle with respect to vertical, θ, encoded through ankle proprioceptors; the angle of the hip with respect to the lower body, β, encoded through hip proprioceptors; and the angle of the upper body with respect to vertical, γ, encoded through visual and vestibular sensory systems. Using Equations 6, 7, or 8, the horizontal position of the CoM and the angle of a single-link inverted pendulum model, Φ, (shown in red) can be determined. All angles are defined as positive for a counterclockwise rotation with respect to the reference.