Bacterial quorum sensing allows graded and bimodal cellular responses to variations in population density

Jennifer B. Rattray[1,2], Stephen A. Thomas[1,2,3], Yifei Wang[1,2,4], Evgeniya Molotkova[1], James Gurney[1,2], John J. Varga[1,2], Sam P. Brown[1,2]

[1] School of Biological Sciences, Georgia Institute of Technology, Atlanta, 30332 GA, USA
[2] Center for Microbial Dynamics and Infection, Georgia Institute of Technology, Atlanta, 30332 GA, USA
[3] Graduate Program in Quantitative Biosciences (QBioS), Georgia Institute of Technology, Atlanta, 30332 GA, USA
[4] The Institute for Data Engineering and Science (IDEaS), Georgia Institute of Technology, Atlanta, 30332 GA, USA

Corresponding author: Sam P. Brown

Email: sam.brown@biology.gatech.edu

ORCIDs. Jennifer B. Rattray https://orcid.org/0000-0003-3478-2242; Stephen A. Thomas http://orcid.org/0000-0003-0773-0236; Yifei Wang http://orcid.org/0000-0002-7510-7959, James Gurney http://orcid.org/0000-0002-8364-3380, John J. Varga https://orcid.org/0000-0002-4868-1971

Keywords

Bacterial communication, quorum sensing, sociomicrobiology, reaction norm

Author Contributions

J.B.R and S.P.B. conceived the idea and designed the experiment; J.B.R., E.M., J.G., and J.J.V. performed the experiments; J.B.R analyzed the data; and J.B.R., S.A.T, Y.W., J.G., J.J.V., and S.P.B. wrote the paper.

Abstract

Quorum sensing (QS) is a mechanism of cell–cell communication that connects gene expression

to environmental conditions (e.g. density) in many bacterial species, mediated by diffusible signal

molecules. Current functional studies focus on a dichotomy of QS on/off (or, quorate / sub-

quorate) states, overlooking the potential for intermediate, graded responses to shifts in the

environment. Here, we track QS regulated protease (*lasB*) expression and show that

Pseudomonas aeruginosa can deliver a graded behavioral response to fine-scale variation in

population density, on both the population and single-cell scales. On the population scale, we see

a graded response to variation in environmental population density. On the single-cell scale, we

see significant bimodality at higher densities, with separate OFF and ON sub-populations that

respond differentially to changes in density; static OFF cells and increasing intensity of

expression among ON cells. While the QS-controlled behavioral output is graded, the underlying

multi-signal dynamics display a threshold shift in signal concentration with increasing density,

reflecting the onset of positive signal auto-regulation at intermediate densities. Together these

results indicate that QS can tune gene expression to graded environmental change, with no

critical cell mass or 'quorum' at which behavioral responses are activated on either the individual

cell or population scale. In an infection context, our results indicate there is not a hard threshold

separating sub-quorate 'stealth' mode and a quorate 'attack' mode.

Main Text

Introduction

Many species of bacteria are capable of a form of cell-cell communication via diffusible signal

molecules, generally referred to as quorum sensing (QS). The study of QS has largely focused on

the intracellular gene regulatory scale, leading to a detailed understanding of the regulatory

mechanisms shaping the production of and response to signal molecules in model organisms

such as *Vibrio cholerae*, *Bacillus cereus* and *Pseudomonas aeruginosa* (1–3). We now

understand that QS is mediated by multiple diffusible signals that together control a diverse array

of responses, including swarming, luminescence, competence and the production of diverse

secreted factors (4, 5)

While the molecular mechanisms of QS have been described for model organisms in remarkable

detail, the functional and evolutionary context of QS continues to be disputed. In other words,

while we now have a better understanding of *how* QS works, we still have limited understanding

of *why* bacteria use this system to control behavior. What are the functions of QS? How do these

QS functions help bacteria to survive and grow? The standard answer is that bacteria use QS to

sense when they are at sufficient density ('quorate') to efficiently turn on cooperative behaviors

such as secretion of toxins and enzymes in order to collectively modify their environment (6).

Other researchers have argued that QS is an asocial sensing apparatus, where individual cells

produce and monitor signal levels in order to infer their physical environment (am I in an open or

enclosed space?) (7). More recently, integration of molecular and evolutionary approaches has

increased the menu of potential functions to include sensing multiple aspects of both the social

and physical environment (6, 8–10) and coordinating complex social strategies that limit the

profitability of non-cooperating 'cheat' strains (11–18).

A critical step in assessing the various adaptive hypotheses is establishing the functional

capacities and limits of QS. Previous studies have demonstrated 'density sensing' functions –

populations can use QS to sense when they exceed a density threshold (6, 19, 20). In addition,

Darch et al. (2012) demonstrated that responding with increased QS controlled cooperative

activity at high density can provide a fitness benefit (6). Other studies have demonstrated

'diffusion sensing' functions (7) – QS systems can functionally respond to variation in physical

containment, so that even a single cell can become 'quorate' (turn on a QS controlled reporter

gene) if isolated in a sufficiently small contained space (9). More recently, some studies have

demonstrated 'genotype sensing' functions – QS can respond to variation in the genotypic

composition of a population, restricting QS-controlled responses to populations that are enriched

with wildtypes (11, 14, 21, 22).

The functional studies outlined above largely focus on a dichotomy of QS on/off (or, quorate /

sub-quorate) states, overlooking the potential for intermediate, graded responses (Fig 1A). The

threshold quorate/non-quorate concept is ingrained in the QS literature following the use of the

legal 'quorum' analogy (20), and is also supported by mathematical models of QS signal

dynamics that highlight how sufficiently strong positive feedback control of signal production can

produce a sharp threshold response to changes in environmental parameters such as density or

diffusion (23, 24). However, these same mathematical models indicate that graded responses are

also possible, dependent on the model parameterization. More generally, Fig 1A highlights that

the phenotypic response of QS bacteria to differing environmental conditions can be viewed as a

'reaction norm' (25–28) that can in principle take differing shapes. Reaction norms describe

phenotypic responses of a single genotype (y-axis, Fig 1A) to varying environmental inputs (x-

axis, Fig 1A). Incorporating a reaction norm framework provides a menu of quantitative metrics to

define QS responses to environmental variation (e.g., slope, intercept, and variances). With this

reaction norm framework, it is important to emphasize that in our study the x-axis is not time, but

instead captures a gradient of environmental conditions. Whether responses are graded or

thresholded during the growth towards high density is a separate line of inquiry (29). Describing

the reaction-norms of QS cells and populations to contrasting environments is an important step

towards understanding the capacities of QS systems to differentially respond to novel

environments.

Figure 1. Schematic of potential population and single cell responses to variation in cell density. A)

Population response (y-axis) across discrete carrying capacity environments (N, x-axis), given a threshold

(left) or graded response (right). In (B) and (C) we outline alternative cell-scale responses (intensity of green

cells) that are consistent with discrete population scale behaviors (yellow arrows). (B) threshold (ON/OFF)

cellular responses can produce a threshold or graded responses on population scale. (C) graded individual

responses can produce threshold or graded responses on a population scale.

Whether the population scale reaction norm to environmental variation is threshold-like or graded

(Fig 1A), a separate question is how collective population-level responses are constructed out of

individual cellular contributions (Fig 1B,C). Studies of QS on a single-cell scale have revealed

substantial heterogeneity in response to QS signals (9, 30–36), highlighting that cell-cell

communication does not necessarily result in tight synchronization of individual cell activity (Fig

1B,C). In some systems, heterogeneity can be quenched by the addition of extra signal (31, 33),

implying a lack of receptor saturation. However, this is not a universal result (30), indicating that

other molecular processes can drive cellular variation in response. Regardless of the molecular

details, we currently lack a behavioral understanding of how individual cellular responses vary

with changes in the environment.

In the current study we address the canonical 'density sensing' function of QS, using the

environmental generalist and opportunistic pathogen *Pseudomonas aeruginosa*, and an

unprecedented scale of environmental resolution (13 discrete limiting carbon levels conducted in
triplicate, generating 39 density environments). Our first challenge is to map the population scale
resolving power of QS to quantitatively discriminate graded differences in population density (Fig
1A). Does *P. aeruginosa* respond in a purely threshold manner, collapsing quantitative
differences in population density into a simple low / high qualitative output, or can QS allow *P.*
aeruginosa to deliver a graded response to distinct environmental densities? Our second
challenge is to understand how collective responses are partitioned across individual cells. Are
changes in collective responses governed primarily by changes in the proportion of cells in an on
state (Fig 1B) or changes in the individual cell intensity of response (Fig 1C), or both?
105
106
Results
Collective level of response to density is graded and linear. Our first challenge is to map out
the population scale reaction norm of the collective QS-controlled protease (*lasB*) response to
variation in population densities. To provide a detailed picture of the QS response reaction norm
to varying density, we grew a QS reporter strain (PAO1 pMHLAS containing the *PlasB::gfp(ASV)*
reporter construct for QS regulated protease expression (37)) under 13 conditions of carbon
limitation in triplicate and measured average fluorescence output per cell as the populations
reach carrying capacity (Fig 2). Dead cells with compromised membranes were identified with a
propidium iodide stain and excluded from analysis. The range of cell densities generated from
this method is from 1×10^8 cells/ml to 2×10^9 cells/ml. Figure 2 shows that QS response is linear
with increasing culture density, providing intermediate levels of average per-capita response to
intermediate densities. To confirm the lack of threshold behavior we assessed alternate statistical
models including threshold functions, and found that a linear fit model supports the data better
than a step-function fit (AIC linear: 89, AIC step-function: 190; relative likelihood that the linear
model is the best fit compared to step-function $> 10^9$, see (38)), supporting a graded population
response as outlined in Figure 1. This agrees with literature that QS induction at lower population
densities is possible (6, 9, 19), but differs in that there is no observable population density at
which populations 'switch', or reach quorum, into a responsive state.

Figure 2. Population response to increasing cell density is linear and graded. 13 distinct culture
carrying capacities were generated by manipulating the concentration of casein as the limiting resource (Fig
S1). Cells were grown to carrying capacity in triplicate and immediately assayed for QS response via
fluorescence microscopy imaging. Response is determined by a fusion of the quorum sensing controlled
lasB promoter and an unstable green fluorescent protein (PAO1 pMHLAS containing *PlasB::gfp(ASV)*).
Individual cell pixel intensity is a measure of cellular quorum sensing response and average pixel intensity is
calculated across all cells in the population as a proxy for total population expression. Microscopy averages
are congruent with population scale plate reader results (Figure S2). A quorum sensing signal knockout
(Δ*lasI*Δ*rhlI*), yellow star, shows background response with no signal in the environment. Average population
investment in QS increases as culture density increases with no observable density threshold (AIC linear:
89, AIC step-function: 190).

Individual response to density is bimodal at high densities. Figure 2 establishes that on a
collective population scale, the response to environmental variation (in density) is smoothly
graded. Next, we ask how this collective response is built from individual cell contributions. Is the
graded increase due to more cells turning on at higher densities (Figure 1B), cells turning on to a
greater extent (Figure 1C), or both? To address this question, we take the same data presented

143 in Figure 2 and now present the distribution of individual cellular responses rather than simply the
144 mean response (Figure 3).
145

146
Figure 3. Individual response is heterogenous and bimodal at higher densities. Ridgeline density plot
(bandwidth = 0.435) of single-cell *lasB* reporter response data showing the distribution of individual cell QS
expression across the population. For brevity and plotting purposes, carrying capacities were averaged
across 3 replicates for each of the 13 carbon environments before plotting. A full plot of each independent
replicate environment can be found in Figure S3. Each line summarizes 18,000 to 30,000 individual cell
measurements, scaled to a unit height. Asterisks indicate significant bimodality (Hartingan's Dip Test (39),
Figure S4). The quorum sensing signal knockout (Δ*lasl*Δ*rhll*) is denoted with yellow boxes. A total of
345,000 individual cell measurements were analyzed.
155
156 As expected from prior studies in other QS systems (9, 30–36), plotting all individual responses
157 within a population shows cell-to-cell variation in QS response within a single population despite
158 isogenic and homogenous culture conditions (Figure 3). In addition, at higher densities we see
159 significant bimodality (defined by Hartigan's Dip Test, Figure S4), with the population segregating
160 into an unresponsive, sub-quorate, OFF state and a responsive, quorate, ON state.
161

162 In light of this bimodality, we fit a two-component finite mixture model to the data (Figure 4A, see
163 Figures S10-S13 and Table S2 for extended analysis), which allows us to define the average
164 intensity of the ON state (Figure 4B) and the proportion of cells in the OFF or ON states (Figure
165 4C).

166
Figure 4. Proportion of cells responding and level of response varies with density. In light of the
bimodal responses in Figure 3, we course-grain the single-cell *lasB* response data into discrete ON/OFF
states. A) Method summary. We quantify distinct ON/OFF states by fitting a two-component finite mixture
model at each measured optical density, where the OFF state is fixed to the OFF state of the highest density
environment. The histogram shows the distribution of cellular expression levels at a single density treatment
(0.76 OD_{600}), the grey line is the fitted OFF state and the green dashed line is the fitted ON state. B) The
mean intensity of the ON (green circle) and OFF (grey triangle) states is determined from the means of
mixture model component fits (green and grey lines in panel A). The mean intensity of the ON state
distribution increases as culture density increases, while the mean of the OFF state remains constant. C)
The proportion of cells ON in the population is determined from the relative mass of cells in the model
component fits. The proportion ON increases with culture density but does not reach 100%.
178
Figure 4B illustrates a graded linear increase in the intensity of the ON state with increasing
environmental density, and a density-invariant off state. Figure 4C illustrates that the proportion of
cells that are ON plateaus at around 85% at densities with consistent support for bimodality
(above 0.36 OD_{600}). At lower densities, the intensity of the ON state (Figure 4B) declines to a
point where the OFF and ON states are no longer significantly different and the dip test fails to
reject uni-modality (Figure S4). In supplemental materials, we present alternate statistical

analyses of this data, and of other related datasets. Across other experiments, we find consistent

support for the graded and bimodal response pattern on the single-cell scale across multiple

assay time-points (Figure S5) and across two reporter strain constructs (Figure S6) and support

for the graded and linear response pattern on the population scale across fluorescent and lux

reporters (Figure S2, Figure S7). We find further support for the graded reaction norm on the

population scale across two additional QS controlled genes (*pqsA, rhlI;* Figure S7).

As quorum sensing is a signal mediated behavior, we sought to connect these behavioral results

with the underlying signaling dynamics in the environment. QS in *Pseudomonas aeruginosa* is

heavily studied in a high-density context, revealing a complex mechanism of multi-signal control

(40–42). The *P. aeruginosa* QS system is dominated by the acyl-homoserone lactone (AHL)

signaling systems LasI/R and RhlI/R (40, 43). Each of these signaling systems codes for a signal

synthase (LasI, RhlI), which guide the production of a diffusible AHL signal molecule (3-oxo-C12

HSL, C4-HSL) at an initially basal level. Binding of each signal to its cognate receptor (LasR,

RhlR) results in an active transcriptional factor which can up-regulate cognate synthase activity

(signal auto-induction) along with other genes in the QS regulon. For more details on the

complexities of *P. aeruginosa* QS wiring, see (44–47).

In light of this established mechanistic understanding of high-density behavior, we outline three

alternate hypotheses for the reaction norms of multi-signal concentrations on density: (1) First,

under a null model of no autoinduction, we predict a linear increase in signal concentration,

reflecting a constant (baseline) per-capita signal production. (2) Second, under a threshold model

for the onset of autoinduction we predict a piecewise linear reaction norm, with a steeper slope in

the higher density environment (reflecting higher per-capita signal production following a

threshold onset of auto-induction). (3) Third, under a graded onset model, we predict a smoothly

accelerating reaction norm, with the slope at a given density reflecting the graded degree of onset

of autoinduction.

To test these hypotheses, we measured the environmental concentration of 3-oxo-C12 HSL and

C4-HSL from previous experiments (at the time point of QS response assays, Figure 3) using *E.*

coli biosensors (48) (Figure 5). To discriminate among the three explicit models, we fit multiple

alternate models to the data and compared their goodness of fit using information criteria. (Figure

5; Tables S3-S5 and Figures S14-S15). A linear fit, representing no positive signal auto-

regulation, is the worst of the models considered for both 3-oxo-C12 HSL and C4 HSL data.

Further statistical tests provide additional evidence for a non-linear relationship (Table S5). In

Figure 5, we show model fits for the best fitting continuously accelerating functions alongside a

threshold model fit. Evaluating these two models via AICc values (corrected AIC to account for

smaller datasets (38)) we find that the threshold model is the best supported model for both the 3-
oxo-C12 HSL and C4 HSL data (Figure 5, inset). More specifically, we can assess the strength of
support for the threshold model via the relative likelihood versus the next-best exponential model
(38). For the C4 HSL data, we find the threshold model has 14-fold greater support, and for the 3-
oxo-C12 HSL data, the threshold model has 8-fold greater support (Table S5). This analysis
indicates there is a critical intermediate density that triggers an abrupt shift in the extent of signal
auto-regulation, separating a basal signal regime from a higher density auto-regulated regime.
229

230
231 **Figure 5. 3-oxo-C12 HSL and C4 HSL concentration increases in a threshold (piecewise linear)**
232 **manner with increasing density.** Signal environment was characterized using filtered culture supernatant
233 (extracted at the same timepoints as for the gene expression data in Figure 3) and *E. coli* biosensors (48).
234 Model fits are based on nonlinear least squares estimates. Model performance was assessed via AICc (see
235 figure insets and Table S5). See Tables S3-S5 and Figures S14-S15 for statistical analysis details.
236

Discussion

Our results show that populations of *P. aeruginosa* can respond in a smoothly graded manner to
variation in environmental density (Figure 2), and that populations exhibit significant bimodality at
higher densities (Figure 3), and that this population scale graded response can be described by
the number of responsive 'ON' cells and the intensity of the 'ON' state (Figure 4). Turning to the
underlying signal mechanics, we further illustrate a threshold onset of signal auto-induction at

intermediate densities (Figure 5). The ability to achieve a graded population scale response

implies in principle that *P. aeruginosa* can tune collective responses (such as the secreted

elastase virulence factor produced by our *lasB* reporter gene) to graded environmental changes,

rather than simply course-graining into a simple 'high / low' dichotomy. A similar population scale

graded response to continuous environmental variation is visible in the data from Allen et al.,

which looked at variation in the genotypic composition of mixed populations grown to the same

total density (11). As the proportion of wildtype (PAO1 versus Δ*lasR* 'cheats') increased, the

wildtype per-capita investment in cooperative LasB secretions also increased, providing a simple

behavioral mechanism to protect cooperative investments from exploitation by cheats (11, 22).

The existence of graded population scale responses across two continuously varying

environmental inputs (density, genotypic composition) raises the question of why use a graded

response? Is there an evolutionary rationale for a graded response, or is a graded increase

simply the 'best approximation' of a threshold response, given a simple system working under

genetic constraints? Existing evolutionary theory suggest that graded investment reaction norms

can be adaptive, under a range of distinct scenarios (49, 50) (34, 35). In the specific context of

quorum-sensing bacteria, evolutionary theory suggests that population scale responses to

increasing density should depend critically on the shape of the cost and benefit functions of

increasing cooperative investments. Specifically, a graded response is predicted to be the optimal

strategy if the benefit function is decelerating and costs are linear with increasing investment (51).

To further consider the functional context of the graded reaction norms, we turn to the single cell

scale data, which reveals how the graded population response is built from the contributions of

individual cells. In agreement with previous work in multiple quorum-sensing organisms (9, 30–

36, 52, 53), we find cell-scale heterogeneity. In addition, our results illustrate how cellular

heterogeneity changes with the environment, demonstrating the onset of ON/OFF bimodality at

intermediate densities, with both the proportion of cells ON and the intensity of the cellular ON

states increasing with increases in culture carrying capacity (Figures 3 & 4).

The presence of a bimodal QS response is in contrast with the common view of QS as a

mechanism of cell synchronization yet can be viewed as a striking example of widely observed

cellular heterogeneity under QS control. Indeed, bimodal responses are implicit in some of the

previous single-cell QS literature (31, 32, 53), for example Darch et al. (2018) report distinct

populations of QS-responsive and non-responsive cells within single experimental runs (53). The

degree of heterogeneity in any cellular trait can be interpreted as the interplay of biochemical

properties of molecules and the architecture of gene-regulatory networks (54). Given that

regulatory networks are subject to mutation and selection, this implies that the degree of

heterogeneity is an evolvable trait (55). In the context of QS, positive feedback loops (signal auto-
regulation (56)) and the presence of cooperative transcription factor binding (57) provides
recognized regulatory ingredients for bimodal expression (58). Recently, the presence of
heterogeneous QS response at the single-cell scale has been ascribed to a potential bet-hedge
against mis-directed QS induction (52), suggesting that our OFF cells are poised to more quickly
resume growth in the event of a rapid return to a growth-friendly environment.
286
We made a number of specific observational choices in order to conduct our experiment that
could have shaped our results in ways that are not generalizable to other contexts. In the
supplementary we detail a number of additional experiments (and alternate statistical analysis
approaches) that collectively illustrate the robustness of our findings. In brief, we found that our
single cell results are not sensitive to the time the population was sampled (Figure S5), the
presence of a potentially leaky P*lac::lasR* on the pMHLAS construct (Figure S6), or the plasmid
nature of the pMHLAS construct (Figure S6). Additionally, we recognize that *lasB* is only one
gene out of hundreds that are controlled by QS (3), and is often co-regulated by other factors
(59–61). We chose to initially focus on *lasB* as it is a traditionally studied QS-controlled trait (62–
64) that has clinical significance as a virulence factor (65, 66). To begin to address the generality
of our results across genes in *P. aeruginosa*, we show that two other QS regulated genes with
complex promoters, *pqsA* and *rhlI*, also support a graded population response (Figure S7). It
remains to be seen whether the graded responses we report here are consistent across all QS
controlled genes in *P. aeruginosa*, and across QS systems in other species,
301
A recent transcriptomic analysis of clinical versus *in vitro* gene expression in *P. aeruginosa* called
into question the clinical relevance of *in vitro* models of QS, reporting that QS activity (including
lasB expression) was systematically higher in *in vitro* models (67). Our results provide a simple
interpretation of this difference: *in vitro* models are conducted under higher experimental
densities, resulting in higher levels of average QS gene expression (Figure 2). Consistent with
this graded response interpretation, Cornforth et al. (2018) also reported higher levels of relative
expression in *in vitro* biofilm models (close-packed cells, the highest local density achievable)
compared to *in vitro* planktonic models.
310
In summary, our results provide a finely resolved mapping of the QS reaction norm to
environmental density in PAO1, on both the collective and single-cell scale. On the population
scale we see a graded linear response across a range of cellular densities (1×10^{8} cells/ml to
2×10^{9} cells/ml) and significant individual-scale bimodality at higher densities. We further resolve
this linear population response (Figure 2) into a combination of the likelihood of being responsive
and the intensity of response (Figure 4). The underlying signal dynamics support a threshold

onset of signal auto-induction at intermediate densities, leading to increased levels of QS signal

production (Figure 5). In an infection context, our results indicate that there is no hard threshold

separating sub-quorate 'stealth' mode and a quorate 'attack' mode (68). One implication is that

attempts to control virulence and biofilm expression in medicine and industry via QS inhibition

could have impacts across a wider spectrum of population densities. In this applied context, it is

important to assess the generality of our results and ask, how do QS reaction-norms vary across

strains and species of QS bacteria? How do they vary across environments? More broadly, our

work undermines the threshold concept of a 'quorum', instead placing QS bacteria in the graded

world of reaction norms.

Materials and Methods

Bacterial Strains and Growth Conditions. The two main bacterial strains used in this study are

P. aeruginosa NPAO1 (Nottingham-PAO1) containing the *PlasB::gfp(ASV)* quorum sensing

reporter pMHLAS (37) and a double signal synthase mutant incapable of producing QS signal

molecules, *P. aeruginosa* NPAO1 *ΔlasI/ΔrhlI* containing the same *PlasB::gfp(ASV)* quorum

sensing reporter pMHLAS. A complete table of strains used in the main text and supplemental

figures can be found in Supplemental Table 1. Overnight cultures were grown in lysogeny broth

(LB), supplemented with 50 ug/ml gentamicin to maintain the pMHLAS plasmid, with shaking at

37 °C. Experiments were conducted in lightly buffered (50 mM MOPS) M9 minimal defined media

composed of an autoclaved basal salts solution (Na_2HPO_4, 6.8 gL^{-1}; KH_2PO_4, 3.0 gL^{-1}; NaCl, 0.5

gL^{-1}), and filter-sterilized 1 mM $MgSO_4$, 100 uM $CaCl_2$, and 1X Hutner's Trace Elements with

casein (CAA) as the sole carbon source.

Controlling Culture Carrying Capacity. We manipulated density by controlling the limiting

resource in the media, carbon, allowing us to tune the carrying capacity of each treatment (Figure

S1). To cover a variety of densities, we generated a CAA range between 0.05% and 0.25% via

dilutions of a 0.5% CAA minimal media stock for a total of 13 different carrying capacities with

three replicates each. This produced a range of densities environments from 1.18×10^8 cells/ml to

2.02×10^9 cells/ml. Overnight cultures were grown in LB gentamicin 50 ug/ml and centrifuged at

8,500 x g for 2 minutes. The cells were then washed twice with carbonless minimal media and

then each carbon treatment was adjusted to OD_{600} = 0.05. Then, 200 uL of each sample was

added to a 96-well microplate. Plates were incubated with continuous shaking at 37 °C in a

Cytation/BioSpa plate reader and growth curves were generated by absorbance readings taken

at 30-min intervals.

Measuring Population QS Response. To measure population response, we performed growth-curve experiments as previously described using PAO1 *PlasB::gfp(ASV)*, additionally taking fluorescence readings at 30-min intervals. Fluorescence, population level response, was recorded when populations reached the end of their exponential growth phase, before they entered stationary phase. Controls for background fluorescence of the reporter were done with the QS signal deficient mutant PAO1 *ΔlasI/ΔrhlI PlasB::gfp(ASV)*. The population microplate data (Figure S2) and averaged microscope data (Figure 2) agreed, so the latter is provided in the primary text.

Measuring Individual QS Response. To measure individual response, we performed growth-curve experiments as previously described, but removed samples for microscopy once cells reached end exponential phase. Since we control carrying capacity with the amount of carbon, the exact time that cells reach the end of exponential growth differs across treatments by 2-3 hours. To robustly sample cultures at this specific point, the slope of the two most recent time points on the growth curve was monitored and samples were taken as the slope approached 0. Replicate wells were kept growing to confirm that the treatment entered stationary phase right after the sampling time point. We also determined that our results are generalizable even when sampling at a pre-determined hour across concentrations (Figure S5). Samples were stained with propidium iodide to differentiate between life and dead cells and a small aliquot (5 ul) was added to a 0.01% poly-l-lysine coated slide to immobilize cells and immediately imaged to avoid changes in expression between sample acquisition and imaging in the dark on a Nikon Eclipse TI inverted microscope at 20x magnification. Live cell fluorescence microscopy was used for this study as fluorophores can be sensitive to fixation/permeabilization. These techniques can result in a decrease in fluorescence and therefore decrease in the observable dynamic range. Bright field, green fluorescence (20% Lumencor light engine power, 200ms exposure, and 64x gain-sufficient for imaging of low fluorescent cells without saturating pixel intensity), and red fluorescence (20% Lumencor light engine power, 800ms exposure, and 64x gain) channels were captured. Between 5,000 and 15,000 individual cells were captured for each sample. Aliquots were diluted immediately before imaging with carbonless minimal media when required to ensure an even distribution of cells.

Single cell image analysis. A custom macro in ImageJ was written to analyze the image, outlined in Figure S8. The macro uses ImageJ's "analyze particles" command to identify single cells on the bright field image. This then generates a ROI (region of interest) for each individual cell and these ROIs were then overlaid onto the corresponding fluorescent image. The red fluorescence channel was used to identify dead cells with compromised membranes, which were excluded from further analysis. The green fluorescence channel reflected the QS reporter and pixel intensity was measured as a proxy for level of QS response. This tabulated live cell

expression data was then analyzed using *Stata Statistical Software: Release 17* from StataCorp

LLC. In order to improve the fit of the mixed models, the lowest pixel intensity measurement in the

highest carbon PAO1 *ΔlasI/ΔrhlI PlasB::gfp(ASV)* treatment was subtracted from all pixel

intensities so that expression started at 0.

Statistical analysis summary. The analysis was done using *Stata Statistical Software: Release*

17 from StataCorp LLC and the additional third party resources: (69–73). Each of the 39

populations was fit to a finite mixture model of two Gamma distributions. The latent classes in the

mixture model correspond to OFF and ON cells. Gamma distributions are preferred to Normal

distributions as gene expression is strictly non-negative and necessarily right-skewed. The

models provide maximum likelihood estimates of the proportion of cells in each latent class and

the shape and scale parameters of the component Gamma distributions. Mean expression level

for each distribution is the product of shape and scale parameters. Information criteria for

aggregate mean expression level was also calculated using Stata.

Quantifying Signal Concentration. AHL signal concentration was estimated using S17-

1 *Escherichia coli* containing either the p56536 or pSB1142 plasmids (74), which luminesce in

response to short and long chain AHLs, respectively. Filtered culture supernatant was diluted

1/100 in LB broth and mixed 1:1 with exponentially growing bioreporter strains at an OD_{600} of 0.1

in LB broth. A calibration curve was generated by exposing the bioreporters to synthetic signal at

various concentrations. Signal bioreporters were grown with diluted supernatant for 3 hours at

37 °C taking reads of optical density and luminescence every 30 min. Using the peak

luminesce, a calibration curve was then fitted to calculate signal concentrations in experimental

samples.

Acknowledgments

We thank Steve Diggle, Marvin Whitely, Kathleen O'Connor and members of the Center for

Microbial Dynamics and Infection (CMDI) for valuable comments and discussion on this work. We

thank Dr. Kasper Norskov Kragh for the *PlasB::gfp(ASV)* reporter construct. This research was

supported by the National Science Foundation Graduate Research Fellowship Program under

Grant No. DGE-1650044 and by The Simons Foundation Grant No. 396001.

References

1. S. T. Rutherford, B. L. Bassler, Bacterial quorum sensing: Its role in virulence and
possibilities for its control. *Cold Spring Harb. Perspect. Med.* **2** (2012).

2. R. G. Abisado, S. Benomar, J. R. Klaus, A. A. Dandekar, J. R. Chandler, Bacterial
Quorum Sensing and Microbial Community Interactions. *MBio* **9**, e02331-17 (2018).

3.   M. Whiteley, S. P. Diggle, E. P. Greenberg, Progress in and promise of bacterial quorum
sensing research. *Nature* **551**, 313–320 (2017).

4.   M. B. Miller, B. L. Bassler, Quorum Sensing in Bacteria. *Annu. Rev. Microbiol.* **55**, 165–
199 (2001).

5.   R. Popat, D. M. Cornforth, L. McNally, S. P. Brown, Collective sensing and collective
responses in quorum-sensing bacteria. *J. R. Soc. Interface* **12** (2015).

6.   S. E. Darch, S. A. West, K. Winzer, S. P. Diggle, Density-dependent fitness benefits in
quorum-sensing bacterial populations. *Proc. Natl. Acad. Sci.* **109**, 8259–8263 (2012).

7.   R. J. Redfield, Is quorum sensing a side effect of diffusion sensing? *Trends Microbiol.* **10**,
365–70 (2002).

8.   B. A. Hense, *et al.*, Does efficiency sensing unify diffusion and quorum sensing? *Nat. Rev.*
Microbiol. **5**, 230–239 (2007).

9.   J. Q. Boedicker, M. E. Vincent, R. F. Ismagilov, Microfluidic Confinement of Single Cells of
Bacteria in Small Volumes Initiates High-Density Behavior of Quorum Sensing and Growth
and Reveals Its Variability. *Angew. Chemie Int. Ed.* **48**, 5908–5911 (2009).

10. D. M. Cornforth, *et al.*, Combinatorial quorum sensing allows bacteria to resolve their
social and physical environment. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 4280–4284 (2014).

11. R. C. Allen, L. McNally, R. Popat, S. P. Brown, Quorum sensing protects bacterial co-
operation from exploitation by cheats. *ISME J.* **10**, 1706–1716 (2016).

12. A. Eldar, Social conflict drives the evolutionary divergence of quorum sensing. *Proc. Natl.*
Acad. Sci. U. S. A. **108**, 13635–13640 (2011).

13. N. E. Smalley, D. An, M. R. Parsek, J. R. Chandler, A. A. Dandekar, Quorum Sensing
Protects Pseudomonas aeruginosa against Cheating by Other Species in a Laboratory
Coculture Model. *J. Bacteriol.* **197**, 3154 LP – 3159 (2015).

14. E. Even-Tov, *et al.*, Social Evolution Selects for Redundancy in Bacterial Quorum
Sensing. *PLOS Biol.* **14**, e1002386 (2016).

15. E. T. Granato, R. Kümmerli, The path to re-evolve cooperation is constrained in
Pseudomonas aeruginosa. *BMC Evol. Biol.* **17**, 214 (2017).

16. Ö. Özkaya, R. Balbontín, I. Gordo, K. B. Xavier, Cheating on Cheaters Stabilizes
Cooperation in Pseudomonas aeruginosa. *Curr. Biol.* **28**, 2070-2080.e6 (2018).

17. R. Chen, E. Déziel, M. C. Groleau, A. L. Schaefer, E. P. Greenberg, Social cheating in a
Pseudomonas aeruginosa quorum-sensing variant. *Proc. Natl. Acad. Sci. U. S. A.* **116**,
7021–7026 (2019).

18. J. Gurney, S. Azimi, S. P. Brown, S. P. Diggle, Combinatorial quorum sensing in
Pseudomonas aeruginosa allows for novel cheating strategies. *Microbiology*, micro000941
(2020).

19. J. L. Connell, *et al.*, Probing prokaryotic social behaviors with bacterial "lobster traps."
MBio **1** (2010).

20. W. C. Fuqua, S. C. Winans, E. P. Greenberg, Quorum sensing in bacteria: The LuxR-LuxI
family of cell density- responsive transcriptional regulators. *J. Bacteriol.* **176**, 269–275
(1994).

21. J. Schluter, A. P. Schoech, K. R. Foster, S. Mitri, The Evolution of Quorum Sensing as a
Mechanism to Infer Kinship. *PLOS Comput. Biol.* **12**, e1004848 (2016).

22. Y. Wang, J. B. Rattray, S. A. Thomas, J. Gurney, S. P. Brown, In silico bacteria evolve
robust cooperaion via complex quorum-sensing strategies. *Sci. Rep.* **10**, 1–10 (2020).

23. A. B. Goryachev, Understanding Bacterial Cell–Cell Communication with Computational
Modeling. *Chem. Rev.* **111**, 238–250 (2011).

24. S. James, P. Nilsson, G. James, S. Kjelleberg, T. Fagerström, Luminescence control in
the marine bacterium Vibrio fischeri: An analysis of the dynamics of lux regulation. *J. Mol.*
Biol. **296**, 1127–1137 (2000).

25. Woltereck, Weitere experimentelle Untersuchungen uber Artveranderung, speziel uber
das Wesen quantitativer Artuntershiede bei Daphniden. *Science (80-.).* **32**, 344–345
(1910).

26. C. H. Waddington, Canalization of development and the inheritance of acquired
characters. *Nature* **150**, 563–565 (1942).

27. C. Schlichting, M. Pigliucci, Phenotypic Evolution: A Reaction Norm Perspective (1998).

28. A. B. Paaby, N. D. Testa, "Developmental Plasticity and Evolution" in *Evolutionary*
Developmental Biology, (Springer International Publishing, 2018), pp. 1–14.

29. R. L. Scholz, E. Peter Greenberg, Positive autoregulation of an Acyl- homoserine lactone
quorum-sensing circuit synchronizes the population response. *MBio* **8** (2017).

30. B. B. Pradhan, S. Chatterjee, Reversible non-genetic phenotypic heterogeneity in bacterial
quorum sensing. *Mol. Microbiol.* **92**, 557–569 (2014).

31. C. Anetzberger, T. Pirch, K. Jung, Heterogeneity in quorum sensing-regulated
bioluminescence of *Vibrio harveyi*. *Mol. Microbiol.* **73**, 267–277 (2009).

32. P. D. Pérez, S. J. Hagen, Heterogeneous response to a quorum-sensing signal in the
luminescence of individual vibrio fischeri. *PLoS One* **5** (2010).

33. J. Grote, D. Krysciak, W. R. Streit, Phenotypic heterogeneity, a phenomenon that may
explain why quorum sensing does not always result in truly homogenous cell behavior.
Appl. Environ. Microbiol. **81**, 5280–5289 (2015).

34. D. Garmyn, *et al.*, Evidence of autoinduction heterogeneity via expression of the agr
system of Listeria monocytogenes at the single-cell level. *Appl. Environ. Microbiol.* **77**,
6286–6289 (2011).

35. E. L. Haseltine, F. H. Arnold, Implications of rewiring bacterial quorum sensing. *Appl.*
Environ. Microbiol. **74**, 437–445 (2008).

36. L. Plener, *et al.*, The Phosphorylation Flow of the Vibrio harveyi Quorum-Sensing
Cascade Determines Levels of Phenotypic Heterogeneity in the Population. *J. Bacteriol.*
197, 1747 (2015).

37. M. Hentzer, *et al.*, Inhibition of quorum sensing in Pseudomonas aeruginosa biofilm
bacteria by a halogenated furanone compound. *Microbiology* **148**, 87–102 (2002).

38. K. P. Burnham, D. R. Anderson, K. P. Huyvaert, AIC model selection and multimodel
inference in behavioral ecology: some background, observations, and comparisons.
Behav. Ecol. Sociobiol. 2010 651 **65**, 23–35 (2010).

39.   J. A. Hartigan, P. M. Hartigan, The Dip Test of Unimodality.
507   *https://doi.org/10.1214/aos/1176346577* **13**, 70–84 (1985).

508   40.   S. PC, P. L, I. BH, Activation of the Pseudomonas aeruginosa lasI gene by LasR and the
509   Pseudomonas autoinducer PAI: an autoinduction regulatory hierarchy. *J. Bacteriol.* **177**,
510   654–659 (1995).

511   41.   M. LM, W. M, Membrane vesicles traffic signals and facilitate group activities in a
512   prokaryote. *Nature* **437**, 422–425 (2005).

513   42.   K. Papenfort, B. L. Bassler, Quorum sensing signal–response systems in Gram-negative
514   bacteria. *Nat. Rev. Microbiol. 2016 149* **14**, 576–588 (2016).

515   43.   L. J, Z. L, The hierarchy quorum sensing network in Pseudomonas aeruginosa. *Protein
516   Cell* **6**, 26–41 (2015).

517   44.   D. V, D. E, Revisiting the quorum-sensing hierarchy in Pseudomonas aeruginosa: the
518   transcriptional regulator RhlR regulates LasR-specific factors. *Microbiology* **155**, 712–723
519   (2009).

520   45.   J. B. Feltner, *et al.*, LasR variant cystic fibrosis isolates reveal an adaptable quorum-
521   sensing hierarchy in Pseudomonas aeruginosa. *MBio* **7** (2016).

522   46.   S.-A. MP, *et al.*, Inactivation of the quorum-sensing transcriptional regulators LasR or
523   RhlR does not suppress the expression of virulence factors and the virulence of
524   Pseudomonas aeruginosa PAO1. *Microbiology* **165**, 425–432 (2019).

525   47.   C. RL, *et al.*, RhlR-Regulated Acyl-Homoserine Lactone Quorum Sensing in a Cystic
526   Fibrosis Isolate of Pseudomonas aeruginosa. *MBio* **11** (2020).

527   48.   W. MK, *et al.*, Construction and analysis of luxCDABE-based plasmid sensors for
528   investigating N-acyl homoserine lactone-mediated quorum sensing. *FEMS Microbiol. Lett.*
529   **163**, 185–192 (1998).

530   49.   , The evolutionary consequences of plasticity in host–pathogen interactions -
531   ScienceDirect (November 10, 2019).

532   50.   S. Via, *et al.*, Adaptive phenotypic plasticity: consensus and controversy. *Trends Ecol.
533   Evol.* **10**, 212–217 (1995).

534   51.   S. Heilmann, S. Krishna, B. Kerr, Why do bacteria regulate public goods by quorum
535   sensing?-How the shapes of cost and benefit functions determine the form of optimal
536   regulation. *Front. Microbiol.* **6** (2015).

537   52.   P. Jayakumar, S. A. Thomas, S. P. Brown, R. Kümmerli, Pseudomonas aeruginosa
538   reaches collective decisions via transient segregation of quorum sensing activities across
539   cells. *bioRxiv*, 2021.03.22.436499 (2021).

540   53.   S. E. Darch, *et al.*, Spatial determinants of quorum signaling in a Pseudomonas
541   aeruginosa infection model. *Proc. Natl. Acad. Sci.* **115**, 4779–4784 (2018).

542   54.   K. Fujimoto, S. Sawai, A Design Principle of Group-level Decision Making in Cell
543   Populations. *PLOS Comput. Biol.* **9**, e1003110 (2013).

544   55.   M. Ackermann, A functional perspective on phenotypic heterogeneity in microorganisms.
545   *Nat. Rev. Microbiol. 2015 138* **13**, 497–508 (2015).

546   56.   J. L. Cherry, F. R. Adler, How to make a Biological Switch. *J. Theor. Biol.* **203**, 117–133

547   (2000).

548   57.   M. Schuster, M. L. Urbanowski, E. P. Greenberg, Promoter specificity in Pseudomonas
549   aeruginosa quorum sensing revealed by DNA binding of purified LasR. *Proc. Natl. Acad.*
550   *Sci.* **101**, 15833–15839 (2004).

551   58.   A. Ochab-Marcinek, M. Tabaka, Bimodal gene expression in noncooperative regulatory
552   systems. *Proc. Natl. Acad. Sci.* **107**, 22096–22101 (2010).

553   59.   B. Mellbye, M. Schuster, Physiological framework for the regulation of quorum sensing-
554   dependent public goods in Pseudomonas aeruginosa. *J. Bacteriol.* **196**, 1155–1164
555   (2014).

556   60.   S. P. Diggle, K. Winzer, A. Lazdunski, P. Williams, M. Cámara, Advancing the quorum in
557   Pseudomonas aeruginosa: MvaT and the regulation of N-acylhomoserine lactone
558   production and virulence gene expression. *J. Bacteriol.* **184**, 2576–2586 (2002).

559   61.   S. A. Chugani, *et al.*, QscR, a modulator of quorum-sensing signal synthesis and virulence
560   in Pseudomonas aeruginosa. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 2752–2757 (2001).

561   62.   M. J. Gambello, B. H. Iglewski, Cloning and characterization of the Pseudomonas
562   aeruginosa lasR gene, a transcriptional activator of elastase expression. *J. Bacteriol.* **173**,
563   3000 (1991).

564   63.   J. P. Pearson, L. Passador, B. H. Iglewski, E. P. Greenberg, A second N-acylhomoserine
565   lactone signal produced by Pseudomonas aeruginosa. *Proc. Natl. Acad. Sci.* **92**, 1490–
566   1494 (1995).

567   64.   B. JM, O. DE, Synthesis of multiple exoproducts in Pseudomonas aeruginosa is under the
568   control of RhlR-RhlI, another set of regulators in strain PAO1 with homology to the
569   autoinducer-responsive LuxR-LuxI family. *J. Bacteriol.* **177**, 7155–7163 (1995).

570   65.   C. F, *et al.*, The LasB Elastase of Pseudomonas aeruginosa Acts in Concert with Alkaline
571   Protease AprA To Prevent Flagellin-Mediated Immune Recognition. *Infect. Immun.* **84**,
572   162–171 (2015).

573   66.   C. Cigana, *et al.*, Pseudomonas aeruginosa Elastase Contributes to the Establishment of
574   Chronic Lung Colonization and Modulates the Immune Response in a Murine Model.
575   *Front. Microbiol.* **0**, 3443 (2021).

576   67.   D. M. Cornforth, *et al.*, Pseudomonas aeruginosa transcriptome during human infection.
577   *Proc. Natl. Acad. Sci.* **115**, E5125–E5134 (2018).

578   68.   K. Winzer, P. Williams, Quorum sensing and the regulation of virulence gene expression
579   in pathogenic bacteria. *Int. J. Med. Microbiol.* **291**, 131–143 (2001).

580   69.   N. J. Cox, DIPTEST: Stata module to compute dip statistic to test for unimodality. *Stat.*
581   *Softw. Components* (2016) (November 8, 2021).

582   70.   N. J. Cox, S. P. Jenkins, GAMMAFIT: Stata module to fit a two-parameter gamma
583   distribution. *Stat. Softw. Components* (2011) (November 8, 2021).

584   71.   P. Deb, FMM: Stata module to estimate finite mixture models. *Stat. Softw. Components*
585   (2012) (November 8, 2021).

586   72.   B. Jann, PALETTES: Stata module to provide color palettes, symbol palettes, and line
587   pattern palettes. *Stat. Softw. Components* (2020) (November 8, 2021).

73. B. Jann, COLRSPACE: Stata module providing a class-based color management system
in Mata. *Stat. Softw. Components* (2020) (November 8, 2021).

74. M. K. Winson, *et al.*, Construction and analysis of luxCDABE -based plasmid sensors for
investigating N -acyl homoserine lactone-mediated quorum sensing . *FEMS Microbiol.*
Lett. **163**, 185–192 (1998).

75. J. Meisner, J. B. Goldberg, The Escherichia coli rhaSR-PrhaBAD inducible promoter
system allows tightly controlled gene expression over a wide range in Pseudomonas
aeruginosa. *Appl. Environ. Microbiol.* **82**, 6715–6727 (2016).

76. F. MP, *et al.*, A dual biosensor for 2-alkyl-4-quinolone quorum-sensing signal molecules.
Environ. Microbiol. **9**, 2683–2693 (2007).

77. H. TT, K. AJ, B. A, S. HP, Integration-proficient plasmids for Pseudomonas aeruginosa:
site-specific integration and use for engineering of reporter and expression strains.
Plasmid **43**, 59–72 (2000).

78. N. Friedman, L. Cai, X. S. Xie, Linking Stochastic Dynamics to Population Distribution: An
Analytical Framework of Gene Expression. *Phys. Rev. Lett.* **97**, 168302 (2006).

79. A. P. Dempster, N. M. Laird, D. B. Rubin, Maximum Likelihood from Incomplete Data via
the EM Algorithm. *J. R. Stat. Soc. Ser. B* **39**, 1–38 (1977).

80. T. L. Bailey, C. Elkan, Fitting a mixture model by expectation maximization to discover
motifs in biopolymers. *Proceedings. Int. Conf. Intell. Syst. Mol. Biol.* **2**, 28–36 (1994).

81. J. Abrevaya, W. Jiang, A Nonparametric Approach to Measuring and Testing Curvature. *J.*
Bus. Econ. Stat. **23**, 1–19 (2005).

82. G. C. Chow, Tests of Equality Between Sets of Coefficients in Two Linear Regressions.
Econometrica **28**, 591–605 (1960).