A high-quality genome assembly of the North American Song Sparrow, *Melospiza melodia*.

Swarnali Louha[1,*], David A. Ray[2], Kevin Winker[3], Travis Glenn[1,4]

[1]Institute of Bioinformatics, University of Georgia, Athens, GA, USA

[2]Department of Biological Science, Texas Tech University, Lubbock, TX, USA

[3]University of Alaska Museum, Fairbanks, AK, USA

[4]Department of Environmental Health Science, College of Public Health, University of Georgia, Athens, GA, USA

*Correspondence: Swarnali Louha, Institute of Bioinformatics, University of Georgia, Athens, GA 30602; E-mail: sl50708@uga.edu

Keywords: *Melospiza melodia*, Dovetail genomics, Passeriforms, whole genome sequencing, *de novo* assembly

Abstract

The song sparrow, *Melospiza melodia*, is one of the most widely distributed species of

songbirds found in North America. It has been used in a wide range of behavioral and ecological

studies. This species' pronounced morphological and behavioral diversity across populations

makes it a favorable candidate in several areas of biomedical research. We have generated a

high-quality *de novo* genome assembly of *M. melodia* using Illumina short read sequences from

genomic and *in vitro* proximity-ligation libraries. The assembled genome is 978.3 Mb, with a

coverage of 24.9×, N50 scaffold size of 5.6 Mb and N50 contig size of 31.7 Kb. Genes within

our genome assembly are largely complete, with 87.5% full-length genes present out of a set of

4,915 universal single-copy orthologs present in most avian genomes. We annotated our genome

assembly and constructed 15,086 gene models, a majority of which have high homology to

related birds, *Taeniopygia guttata* and *Junco hyemalis*. In total, 83% of the annotated genes are

assigned with putative functions. Furthermore, only ~7% of the genome is found to be repetitive;

these regions and other non-coding functional regions are also identified. The high-quality *M.*

melodia genome assembly and annotations we report will serve as a valuable resource for

facilitating studies on genome structure and evolution that can contribute to biomedical research

and serve as a reference in population genomic and comparative genomic studies of closely

related species.

20   **INTRODUCTION**

The oscine passerines (Order Passeriformes) are songbirds having specialized vocal

learning capabilities. Many species of songbirds have been widely used by neuroscientists to

study the processes underlying memory and learning and social interactions (Doupe et al. 1999,

White 2010). The song sparrow (*Melospiza melodia*) is one of the most morphologically diverse

songbirds found in North America, with 26 recognized subspecies (Pruett et al. 2008). It has

been recognized as a model vertebrate species for field studies of birds and has been the subject

of extensive research integrating behavioral and ecological studies over the last 70 years (Arcese

et al. 2002). Though several species of songbirds have been sequenced and studied (Warren et al.

2010, Jarvis et al. 2014), few offer the plethora of biomedical research potential presented by the

song sparrow. For example, differences in neural growth and song-center brain development

among different populations of song sparrows could play an instrumental role in basic research

involving human brain neurogenesis (NIH 2001). The molecular processes underlying the

regeneration of "hair" cells in the auditory system of song sparrows might also provide therapies

useful in human hearing loss (Hawkins et al. 2003, Hawkins and Lovett 2004). Several other

areas of biomedical research in which the song sparrow might serve as a model system are

hepatic lipogenesis (Gosler 1996, Schubert et al. 2007), craniofacial development (Brugmann et

al. 2010, Powder et al. 2012), and variations in body size (Sutter et al. 2007, Allen et al. 2010).

The last of these is a polygenic trait, and elucidation of the underlying gene network affecting

different metabolic pathways can help clarify several biological phenomena, including human

diseases. Given its significant biomedical potential and experimental tractability in the field and

aviary, the song sparrow will continue to be used for answering research questions related to

mechanisms causing variation in behavior, morphology, and demographics across populations

(Arcese et al. 2002, Nietlisbach et al. 2015).

Prior work on population genetics of song sparrows in Alaska has shown how sequential

bottlenecking of populations in the Aleutian Archipelago has given rise to a naturally inbred

population with large body size (Pruett and Winker 2005). Previous work has also been done on

the song sparrow transcriptome, developing genomic markers to screen at population levels

(Srivastava et al. 2012). A high-quality genome assembly of *M. melodia* furthers the

development of genomic markers to screen loci associated with phenotypic traits of interest. An

ever-growing number of songbirds have sequenced genomes, but relatively few have been

published so far, including the American crow (*Corvus brachyrhynchos*), golden-collared

manakin (*Manacus vitellinus*; Jarvis et al. 2014), Zebra finch (*Taeniopygia guttata*; Warren et al.

2010), medium ground finch (*Geospiza fortis*; Parker et al. 2012) and the dark-eyed junco (*Junco*

hyemalis; Friis et al. 2018). In this study, we provide the genome assembly of *Melospiza*

melodia, a member of the family Passerellidae; this will facilitate genomic and phylogenetic

comparisons with other songbirds, and will also serve as a reference genome for comparative

genomic studies.

Our high-quality draft genome assembly of *M. melodia* was created by combining both

traditional Illumina paired-end libraries and a *de novo* proximity-ligation Chicago library. The

Chicago library method together with Dovetail Genomics' HiRise software pipeline is designed

to significantly reduce gaps in alignment arising from repetitive elements in the genome (Putnam

et al. 2016) and increases assembly contiguity. The draft genome was annotated using

transcribed RNA and protein sequences from *M. melodia* and related songbird species, *Junco*

hyemalis and *Taeniopygia guttata*. Genomic features of interest other than coding sequences,

such as microsatellites, repeat elements, transposable elements, and non-coding RNA, were also

annotated and the genome assembly was evaluated for quality by comparing it to related avian

species.

69   **METHODS**

Library preparation and *de novo* shotgun assembly

The *de novo* assembly of the song sparrow genome was constructed using Illumina paired end libraries. Genomic DNA was isolated from a blood sample of a single male song sparrow, obtained from the wild in the Aleutian Islands of Alaska on 16 Sep 2003 and archived as a voucher specimen at the University of Alaska Museum (http://arctos.database.museum/guid/UAM:Bird:31500). We sheared the genomic DNA using the Covaris S2 (Covaris, Woburn, MA, USA) targeting a 600bp average fragment size. The sheared DNA was end-repaired, adenylated, and ligated to TruSeq LT adapters using a TruSeq DNA PCR-Free Library Preparation Kit (Illumina, San Diego, CA, USA). We purified the ligation reaction using a Qiaquick Gel Extraction Kit (Qiagen, Venlo, The Netherlands) from a 2% agarose gel. We sequenced the library on an Illumina HiSeq 2500 at the HudsonAlpha Institute for Biotechnology (Huntsville, AL, USA) to obtain paired-end (PE) ~100 bp reads. The sequence data consisted of 276 million read pairs sequenced from a total of 41.3 Gbp of paired-end libraries (~50× coverage). Reads were trimmed for quality, sequencing adapters, and mate pair adapters using Trimmomatic (Bolger et al. 2015). The reads were assembled at Dovetail Genomics (Santa Cruz, CA, USA) using Meraculous 2.0.4 (Chapman et al. 2011) with a *k-mer* size of 29. This yielded a 972.4 Mbp assembly with a contig N50 of 22.5 Kbp and a scaffold N50 of 33 Kbp.

Chicago library preparation and scaffolding the draft genome

To improve the *de novo* assembly, a Chicago library was prepared at Dovetail Genomics using previously described methods (Putnam et al. 2016). Around 500 ng of high-molecular-weight genomic DNA (mean fragment length = 50 kbp) was used for chromatin reconstitution *in vitro* and fixed with formaldehyde. Fixed chromatin was digested with *Dpn*II, the 5' overhangs

filled in with biotinylated nucleotides, and free blunt ends were ligated together. After ligation,

crosslinks were reversed and DNA was purified from protein. Purified DNA was treated to

remove biotin that was not internal to ligated fragments. Next, DNA was sheared to ~350 bp

mean fragment size and sequencing libraries were generated using NEBNext Ultra enzymes

(New England Biolabs, Ipswich, MA, USA) and Illumina-compatible adapters. Biotin-containing

fragments were isolated using streptavidin beads before PCR enrichment of the library. The

Chicago library was sequenced on an Illumina HiSeq 2500 to produce 47 million 150 bp paired

end reads (1-50 kb pairs).

Dovetail Genomics' HiRise scaffolding software pipeline (Putnam et al. 2016) was used

to map the shotgun and Chicago library sequences to the draft *de novo* assembly using a

modified SNAP read mapper (http://snap.cs.berkeley.edu). The separations of Chicago read pairs

mapped within draft scaffolds were analyzed by HiRise to produce a likelihood model for

genomic distance between read pairs, and the model was used to identify and break putative

misjoins, to score prospective joins, and make joins above a threshold. After scaffolding, shotgun

sequences were used to close gaps between contigs.

Identification of microsatellites and transposable elements

Transposable elements (TEs) in the song sparrow genome were identified using a

combination of *de novo* and homology-based TE identification methods, in addition to a manual

curation step (Platt et al. 2016). First, we used RepeatModeler v1.0.11 (Smit and Hubley 2008-

2015) with default parameters (File S1) to generate a custom repeat library consisting of 672

consensus repeat sequences. RepeatModeler uses two *de novo* repeat identification programs,

RECON v1.08 (Bao and Eddy 2002) and RepeatScout v1.0.6 (Pevzner et al. 2005), for

identifying repetitive elements from sequence data. To ensure accurate and complete

representation of putative TEs, the RepeatModeler derived consensus sequences were filtered for

size (>100 bp), and then subjected to iterative homology-based searches against the genome,

followed by manual curation (Platt et al. 2016). The final set of manually curated TEs was

queried against CENSOR (Kohany et al. 2006) and TEclass (Abrusan et al. 2009) for

classification. TEs not identifiable in CENSOR were also searched against the NCBI nucleotide

and protein databases using BLASTN and BLASTX respectively. Finally, a custom *de novo*

repeat library comprising song sparrow-specific TEs and existing simple and low complexity

repeats in birds was used to screen for repeats in the song sparrow genome assembly with

RepeatMasker v4.0.9.

Microsatellites in the song sparrow genome were identified and described with GMATA

v2.01 (Wang et al. 2016) with sequence motifs ranging in length from 2-20 bp, and each motif

repeated at least 5 times (File S2).

De novo **gene annotation and function prediction**

Genes were predicted in the song sparrow genome with the MAKER v2.31.9

genome annotation pipeline (Campbell et al. 2014). A custom repeat library of 900 repeat

sequences consisting of TEs identified in the song sparrow genome and other existing avian

repeat elements was used to soft mask the genome. Transcriptome evidence sets for MAKER

included the assembled song sparrow transcriptome (Srivastava et al. 2012) and Trinity (v2.4.0)

mRNA-seq assemblies from multiple tissues of *Junco hyemalis* (Peterson et al. 2012, NCBI

BioProject Accession: PRJNA256328). Protein evidence sets used by MAKER included

annotated proteins for song sparrow, *Junco hyemalis*, and *Taeniopygia guttata* from the NCBI

Protein database. The MAKER pipeline consisted of the following steps: 1) Transcriptomic and

protein evidence sets were used to make initial evidence-based annotations with MAKER; 2) the

initial annotations were used to train two *ab initio* gene predicters: Augustus (Stanke et al. 2006),

which was trained once, and SNAP (Korf 2004), which was iteratively trained twice; and 3) the

trained gene prediction tools SNAP and Augustus were used to generate the final set of gene

annotations (File S3-S8).

Functional annotations of the predicted genes were obtained by making homology-based

searches with BLASTP against the Uniprot/Swiss-Prot protein database (Pundir et al. 2016, File

S9). InterProScan v5.29 (Zdobnov et al. 2001) was used to find protein domains associated with

the genes. The putative functions and protein domains were added to the gene annotations using

scripts provided with MAKER (File S9).

To quantitatively assess the completeness of the song sparrow genome assembly and

annotated gene set, we ran BUSCO (Benchmarking Universal Single-Copy Orthologs) v3.0.2

(Waterhouse et al. 2017) with 4,915 single-copy orthologous genes in the Aves lineage group

(Aves_odb9; https://busco.ezlab.org/), using "chicken" as the Augustus reference species (File

S10). The 4,915 orthologous genes are present in at least 90% of the 40 species included within

the Aves lineage group, and thus are likely to be found in the genome of related species. We also

used the JupiterPlot pipeline (https://github.com/JustinChu/JupiterPlot) to visually compare the

zebra finch (*T. guttata*) genome assembly (Warren et al. 2010) to our assembly in a Circos plot,

using the largest scaffolds making up 95% of our genome assembly, and all scaffolds greater

than 100 kbp in the Zebra finch genome (File S11). The zebra finch was selected because it was

the first fully sequenced songbird genome, it is one of the most complete bird genomes, and it is

often used for comparative genomic studies in birds.

Non-coding RNA prediction

Transfer RNAs (tRNAs) were predicted in the song sparrow genome with tRNAscan-SE

v2.0 (Lowe and Chan 2016, File S12). A training set comprising eukaryotic tRNAs was used to

train the covariance models employed by tRNAscan-SE, and tRNAs were searched against the

genome with Infernal v1.1.2 (Nawrocki et al. 2014). tRNAscan-SE also provides functional

classification of tRNAs based on a comparative analysis using a suite of isotype-specific tRNA

covariance models. A random sample of 10 predicted tRNAs were selected and searched against

the tRNA databases GtRNAdb (Chan and Lowe 2016) and tRNAdb (Juhling et al.2009).

Identification of miRNAs (microRNAs), snoRNAs (small nucleolar RNAs), snRNAs

(small nuclear RNAs), rRNAs (ribosomal RNAs), and lncRNAs (long non-coding RNAs) was

achieved by using a homology-based prediction method. Structural homologs to eukaryotic

ncRNA covariance models from the Rfam database v14.1 (Gardner et al. 2009) were searched

against the song sparrow genome using Infernal's "cmscan" program (File S13). All low-scoring

overlapping hits and hits with an E-value greater than 10_{-5} were discarded, and the remaining

ncRNAs were grouped into different classes.

Lastly, we compared the predicted classes of different ncRNAs in the song sparrow

genome to those reported in the genomes of related birds, *Taeniopygia guttata* and *Ficedula*

albicollis (collared flycatcher).

179   **DATA AVAILABILITY**

Raw reads have been deposited in the NCBI Sequence Read Archive (SRR10491484 and

SRR10451714 for the Meraculous assembly, and SRR10424475 for the Chicago HiRise

assembly). Supplemental files available at FigShare. Supplemental File S1 contains submission

script for RepeatModeler. Supplemental File S2 contains primary configuration file used to run

GMATA (default_cfg.txt). Supplemental File S3 contains submission script for MAKER.

Supplemental File S4 contains MAKER executable file (maker_exe.ctl). Supplemental File S5

contains specifications for downstream filtering of BLAST and Exonerate alignments

(maker_bopts.ctl). Supplemental File S6 contains primary configuration of MAKER specific

options (maker_opts.ctl). Supplemental File S7 contains scripts for training SNAP. Supplemental

File S8 contains scripts for training Augustus. Supplemental File S9 contains scripts for running

BLASTP and InterProScan for functional annotation of predicted genes; and scripts for adding

the functional annotations to gene annotation files. Supplemental File S10 contains submission

script for BUSCO. Supplemental File S11 contains submission scripts for JupiterPlot pipeline.

Supplemental File S12 contains submission script for tRNAscan-SE. Supplemental File S13

contains submission script for Infernal. Supplemental File S14 contains classification of

predicted transposable elements. Supplemental File S15 contains annotation of microsatellites

with their genomic locations. Supplemental File S16 contains percentage of different

microsatellites present in the genome. Supplemental File S17 contains frequency of occurrence

of microsatellites in each scaffold of the genome. Supplemental File S18 contains the distribution

of the length of microsatellites. Supplemental File S19 contains predicted function of annotated

genes by BLASTP. Supplemental File S20 contains prediction of protein domains, GO

annotations and pathway annotations of predicted genes by InterProScan. Supplemental File S21

contains sequence and structure of tRNAs identified in the Song Sparrow genome. Supplemental

File S22 contains classification of predicted tRNAs. Supplemental File S23 contains

classification of different ncRNAs predicted in the genome with Infernal. Supplemental File S24

contains the M. melodia Chicago HiRise genome sequence (Mmel_1.0). Supplemental File S25

contains annotations for the M. melodia genome assembly. Supplemental Table S1 contains

genome sizes of birds related to M. melodia. Supplemental Figure S1 contains the distribution of

the percentage of annotated genes with their corresponding AED scores.

210   **RESULTS AND DISCUSSION**

Assembly

We produced the *de novo* genome assembly of song sparrow, with a total length of 978.3

Mb, using a Chicago library and the HiRise assembly pipeline. The N50 scaffold size was 5.6

Mb and contig size was 31.7 Kb. This assembly showed significant improvement over the initial

shotgun assembly, with a 169-fold increase in scaffold N50 and a 60-fold increase in scaffold

N90 (Table 1). These increases in scaffold size were also accompanied by an increase in

assembly contiguity, with the total number of scaffolds decreasing from 74,832 to 13,785 (Fig 1,

Table 1).

Microsatellites and Transposable Elements

In total, 89 as yet unnamed TEs were identified in the song sparrow genome. Fifty-five of

these did not have any significant matches in CENSOR (Kohany et al. 2006) and are considered

novel (File S14). A TE was considered to have a significant match to a known element in

CENSOR only when it had a length of at least 80 bp and 80% identity to the known element over

80% of its length, the 80-80-80 rule (Schulman et al. 2007). The predicted TEs were classified

into DNA transposons and retrotransposons (i.e. LINEs, LTRs, and SINEs) using CENSOR and

TEclass (File S14). Approximately 7.4% of the genome comprises repeats with the majority of

that consisting of TEs (~ 48%). Among the different TEs, LTRs (~ 40%) and LINEs (~ 49%)

were found to be most abundant (Table 2). The song sparrow genome assembly was found to be

less repetitive when compared to sequenced genomes of related songbirds, primarily due to the

lower content of LTRs and LINEs than other songbirds (Fig 2).

Overall, 112,419 microsatellites with motifs ranging in size from 2-20 bp were found in

the song sparrow genome (File S15 contains all microsatellites with their genomic locations).

The majority of the microsatellites were made up of 2-, 3-, 4-, and 5-mers, with 2-mers making

up about 71% of all microsatellites identified (Fig 3, File S16). The distribution of the top base-

pair composition of microsatellite motifs present in the genome is shown in Fig 4. The frequency

of occurrence of microsatellites in every scaffold and a distribution of their lengths are provided

in Files S17 and S18, respectively.

Gene annotation and function prediction

The MAKER genome annotation pipeline predicted 15,086 genes and 139 pseudogenes

in the song sparrow genome, fewer than *T. guttata, F. albicollis*, and *M. vitellinus*, but higher

than *G. fortis* (Table 3). The average gene length, exon length, intron length, and the total

number of exons and introns predicted are also less compared to closely related species (Table

3). Out of the 15,086 predicted genes, 12,541 genes were assigned putative functions with

BLASTP (File S19). InterProScan assigned functional domains to 11,298 (74.9%) predicted

genes (File S20). A total of 7,010 genes obtained GO annotations. Pathway annotations were

assigned to 2,716 genes.

Annotated genes were assigned annotation edit distance (AED) scores with values

ranging from 0 to 1. AED is a distance metric score that signifies how closely gene models

match transcript and protein evidence. Gene models with AED scores closer to 0 have better

alignment with the evidence provided in the MAKER pipeline. A distribution of the percentage

of genes with their corresponding AED scores shows close similarity of the annotated genes with

the transcript and protein evidence provided in the MAKER pipeline (Fig S1).

The song sparrow genome assembly contained 4,318 complete universal single-copy

orthologs (BUSCOs; 87.9%) from a total of 4,915 BUSCO groups searched. Among all complete

BUSCOs, 99.4% were present as single-copy genes and 0.6% were duplicated. About 7.4%

(356) of the orthologous gene models were partially recovered, and 4.9% (241) had no

significant matches. The incomplete and missing gene models could either be partially present or

missing, or could indicate genes that are too divergent or have very complex structures, making

their prediction difficult. Incomplete and missing gene models could also suggest problems

associated with the genome assembly and gene annotation. The results from the BUSCO analysis

are in agreement with the Circos plot (Fig 5), in which few scaffolds in the zebra finch genome

assembly are not represented in our assembly and very few small translocations exist between

the two genome assemblies.

Non-coding RNA prediction and identification

A total of 267 tRNAs were detected in the song sparrow genome by tRNAscan-SE (see

File S21 for sequence and structure of tRNAs), out of which 129 were found coding for the

standard twenty amino acids. The predicted output from tRNAscan-SE (File S22) contained 114

tRNAs with low Infernal as well as Isotype scores; these were characterized as pseudogenes

lacking tRNA-like secondary structures (Lowe and Chan 2016). Two tRNAs had undetermined

isotypes and 22 were chimeric, with mismatch isotypes. Chimeric tRNAs contain point

mutations in their anticodon sequence, rendering different predicted isotypes than those predicted

by structure-specific tRNAscan-SE covariance models. Among all predicted tRNAs, 11

contained introns within their sequences. No suppressor tRNAs and tRNAs coding for

selenocysteine were predicted. The subset of 10 randomly selected tRNAs was also predicted in

many other species in both GtRNAdb and tRNAdb databases.

Infernal searches predicted a total of 364 ncRNAs in the song sparrow genome,

comprising 166 miRNAs, 8 rRNAs, 154 snoRNAs, 16 snRNAs, and 20 lncRNAs (File S23).

Compared to the genomes of related avian species (*T. guttata* and *F. albicollis*), the song sparrow

genome has the highest number of predicted tRNAs, but fewer other ncRNAs (Table 4).

281   **CONCLUSION**

The Chicago and shotgun sequencing libraries along with the HiRise assembly software

enabled accurate and highly contiguous *de novo* assembly of the song sparrow genome. The

genome assembly is 978.3 Mb, with 48 scaffolds (L50) making up half the genome size. A

previous estimate of genome size of *M. melodia* from densitometry analysis provided a C-value

of 1.43 pg (1,398.54 Mb) (Gregory et al. 2009). Our own *k-mer* based estimate of genome size

from paired reads in the shotgun and Chicago libraries using Kmergenie v1.7044 (Chikhi and

Medvedev 2013) yielded an estimated size of 1,127.25 Mb. Both these genome size estimates

and the genome sizes of related birds (Table S1) are slightly higher than our genome assembly

(978.3 Mb); this could be due to the low TE content observed in the song sparrow. Vertebrate

genome sizes are typically correlated with TE content and birds are no exception (Kidwell 2002,

Gregory and Elliott 2015). Indeed, birds, along with bats, are proposed to have been subject to

extensive DNA loss via an 'accordion model' of genome evolution (Kapusta et al. 2017). Our

data suggest that, among birds, the song sparrow may have been particularly susceptible to the

segmental deletions that are critical to this model. The small genome size of the song sparrow

can also be attributed to the compression of repetitive regions, which is generally observed in

assemblies generated from short-read sequencing data. This is also consistent with the fact that

our genome contains fewer repeats when compared to related songbirds (Fig 2). Although short

reads limit our ability to characterize all genomic repeats, we have been able to resolve the vast

majority of repeats into LINEs, SINEs, LTRs, and DNA retrotransposons (Fig 2, Table 2).

Our genome is highly complete, with 87.5% full-length genes present out of 4,915

universal orthologous genes in avian species. A large set of genes (15,086) with known

homology to related birds was annotated in our study. A majority of these genes (83%) were

assigned with putative functions. The improved scaffold lengths and gene model annotations will

facilitate studies to identify genes responsible for multiple phenotypic traits of interest.

Additionally, longer scaffolds in the Chicago HiRise assembly will help detect regions under

selection, including SNPs and structural variants such as insertions/deletions or copy number

variations which are potentially responsible for the phenotypic diversity observed in this species.

Though we report fewer miRNAs, snRNAs, snoRNAs, rRNAs, and lncRNAs in this

genome than in related songbirds, we have high confidence in the predicted ncRNAs because we

used conservative cutoffs to reduce false positives. Pending the availability of long-read data,

this genome assembly provides an excellent reference for a range of genetic, ecological,

functional, and comparative genomic studies in song sparrows and other songbirds.

ACKNOWLEDGEMENTS

This research was supported in part by an Institutional Development Award (IDeA) from the

National Institute of General Medical Sciences of the NIH (P20GM103395). The content is

solely the responsibility of the authors and does not necessarily reflect the official views of the

NIH. We also thank David Sonneborn and Jack Withrow for their roles in making a vouchered

specimen available for our work. We thank the staff of Dovetail Genomics for help in preparing

and processing the Chicago library and HiRise assemblies. The high performance computing

cluster at Georgia Advanced Computing Resource Center (GACRC) at the University of Georgia

provided computational infrastructure and technical support throughout the work.

References

References

Abrusan G., Grundmann N., DeMeester L., Makalowski W., 2009 TEclass: a tool for automated

classification of unknown eukaryotic transposable elements. Bioinformatics. 25:1329-1330

Andrews C.B., Mackenzie S.A., Gregory T.R., 2009 Genome size and wing parameters in

passerine birds. Proc Biol Sci. 276(1654):55-61. doi:10.1098/rspb.2008.1012

Arcese P., Sogge M.K., Marr A.B., Patten M.A., 2002 Song Sparrow (*Melospiza melodia*) In:

Poole A., Gill F., editors. The Birds of North America. Philadelphia, PA: The Birds of North

America Inc. No. 704

Bolger A.M., Lohse M., Usadel B., 2014 Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics. 30: 2114–2120. doi:10.1093/bioinformatics/btu170

Brugmann S.A., Powder K.E., Young N.M., Goodnough L.H., Hahn S.M., James A.W., Helms

J.A., Lovett M., 2010 Comparative gene expression analysis of avian embryonic facial structures

reveals new candidates for human craniofacial disorders. Human Molecular genetics. 19: 920-

930

Campbell M. S., Holt C., Moore B., Yandell M., 2014 Genome annotation and curation using

MAKER and MAKER-P. Current Protocols in Bioinformatics. 48:4.11.1–39.

doi:10.1002/0471250953.bi0411s48

Chan P. P., Lowe T. M., 2016 GtRNAdb 2.0: An expanded database of transfer RNA genes

identified in complete and draft genomes. Nucleic Acids Res. 44: D184–D189.

doi:10.1093/nar/gkv1309

Chapman J.A., Ho I., Sunkara S., Luo S., Schroth G.P., Rokhsar D.S., 2011 Meraculous: de novo

genome assembly with short paired-end reads. PLoS One. 6(8):e23501.

doi:10.1371/journal.pone.0023501

Chikhi R, Medvedev P., 2014 Informed and automated k-mer size selection for genome

assembly. Bioinformatics. 30(1):31-37. doi: 10.1093/bioinformatics/btt310

Doupe A.J., Kuhl P.K., 1999 Birdsong and Human Speech: Common Themes and Mechanisms.

Annual Review of Neuroscience. 22(1) 567-631. DOI: 10.1146/annurev.neuro.22.1.567

Elliott T.A., Gregory T.R., 2015 What's in a genome? The C-value enigma and the evolution of

eukaryotic genome content. Philos Trans R Soc Lond B Biol Sci. 370(1678):20140331. doi:

10.1098/rstb.2014.0331

Friis G., Fandos G., Zellmer A.J., McCormack J.E.,4, Faircloth B.C., Milá B., 2018 Genome-

wide signals of drift and local adaptation during rapid lineage divergence in a songbird. Mol

Ecol. 27(24):5137-5153. doi:10.1111/mec.14946

Gardner P.P., Daub J., Tate J., Moore B.L., Osuch I.H., Griffiths-Jones S., Finn R.D., Nawrocki

E.P., Kolbe D.L., Eddy S.R., Bateman A., 2011 Rfam: Wikipedia, clans and the 'decimal'

release. Nucleic Acids Res. 39(Database issue):D141-5. doi:10.1093/nar/gkq1129

Gosler A.G., 1996 Environmental and social determinants of winter fat storage in the Great Tit

Parus major. J. Animal Ecology. 65:1-17

Hawkins R.D., Bashiardes S., Helms C.A., Hu L., Saccnone N.L., Warchol M.E., Lovett M.,

2003 Gene expression differences in quiescent versus regenerating hair cells of avian sensory

epithelia: implications for human hearing and balance disorders. Human Molecular Genetics. 12:

1261-1272

Hawkins R.D., Lovett M., 2004 The developmental genetics of auditory hair cells. Human

Molecular Genetics. 13: R289-R296

Jarvis E.D., Mirarab S., Aberer A.J., Li B., Houde P., Li C., Ho S.Y., Faircloth B.C., et al., 2014

Whole-genome analyses resolve early branches in the tree of life of modern birds. Science.

12;346(6215):1320-31. doi: 10.1126/science.1253451

Juhling F., Morl M., Hartmann R.K., Sprinzl M., Stadler P.F., Pütz J., 2009 tRNAdb 2009:

Compilation of tRNA sequences and tRNA genes. Nucleic Acids Res. 37: D159–D162.

doi:10.1093/nar/gkn772

Kapusta A., Suh A., Feschotte C., 2017 Dynamics of genome size evolution in birds and

mammals. Proc Natl Acad Sci USA. 114(8):E1460-E1469. doi:10.1073/pnas.1616702114

Kidwell M.G., 2002 Transposable elements and the evolution of genome size in eukaryotes.

Genetica. 115(1):49-63

Kohany O., Gentles A.J., Hankus L., Jurka J., 2006 Annotation, submission and screening of

repetitive elements in Repbase: RepbaseSubmitter and Censor. BMC Bioinformatics. 25;7:474

Korf I., 2004 Gene finding in novel genomes. BMC Bioinformatics 5:59. doi:10.1186/1471-

2105-5-59

Lango A.H., Estrada K., Lettre G., Berndt S.I., Weedon M.N., et al., 2010 Hundreds of variants

clustered in genomic loci and biological pathways affect human height. Nature. 467:832-838

Lowe T.M., Chan P.P., 2016 tRNAscan-SE On-line: Search and Contextual Analysis of Transfer

RNA Genes. Nucleic Acids Res. 44:W54-57

Nawrocki E.P., 2014 Annotating functional RNAs in genomes using Infernal. Methods Mol Biol.

1097:163–97. doi: 10.1007/978-1-62703-709-9_9

Nietlisbach P., Camenisch G., Bucher T., Slate J., Keller L.F., Postma E., 2015 A microsatellite-

based linkage map for song sparrows (*Melospiza melodia*). Molecular Ecology Resources

15:1486-1496

NIH (2001) What we learned from songbirds: The adult brain can generate new nerve cells. NIH

Publication No. 01-4602

Parker P., Li B., Li H., Wang J., 2012 The genome of Darwin's Finch (Geospiza fortis).

GigaScience. http://dx.doi.org/10.5524/100040

Peterson M.P., Whittaker D.J., Ambreth S., Sureshchandra S., Buechlein A., Podicheti R., Choi

J.H., Lai Z., Mockatis K., Colbourne J., Tang H., Ketterson E.D., 2012 De novo transcriptome

sequencing in a songbird, the dark-eyed junco (*Junco hyemalis*): genomic tools for an ecological

model system. BMC Genomics. 13:305. doi: 10.1186/1471-2164-13-305

Platt R.N. 2nd, Blanco-Berdugo L., Ray D.A., 2016 Accurate Transposable Element Annotation

Is Vital When Analyzing New Genome Assemblies. Genome Biology and Evolution. 8(2):403-

10. doi: 10.1093/gbe/evw009

Powder K.E., Ku Y.C., Brugmann S.A., Veile R.A., Renaud N.A., Helms J.A., Lovett M., 2012

A cross-species analysis of microRNAs in the developing avian face. PLoS One. 7(4):e35111.

doi: 10.1371/journal.pone.0035111

Price A.L., Jones N.C., Pevzner P.A., 2005 De novo identification of repeat families in large

genomes. Bioinformatics. Suppl 1:i351-8

Pruett C.L., Arcese P., Chan Y.L., Wilson A.G., Patten M.A., Keller L.F., Winker K., 2008

Concordant and discordant signals between genetic data and described subspecies of Pacific

Coast Song Sparrows. The Condor. 110(2) 359-364. DOI: 10.1525/cond.2008.8475

Pruett C.L., Winker K., 2005 Northwestern Song Sparrow populations show genetic effects of

sequential colonization. Molecular Ecology. 14:1421-1434

Pundir S., Martin M. J., O'Donovan C., 2016 UniProt Tools. Curr. Protoc. Bioinformatics.

53:1.29.1-15. doi: 10.1002/0471250953.bi0129s53

Putnam N.H., O'Connell B.L., Stites J.C., Rice B.J., Blanchette M., Calef R., Troll C.J., Fields

A., Hartley P.D., Sugnet C.W., Haussler D., Rokhsar D.S., Green R.E., 2016 Chromosome-scale

shotgun assembly using an in vitro method for long-range linkage. Genome Research. 26:342-

350

Schubert K.A., Mennill D.J., Ramsay S.M., Otter K.A., Boag P.T., Ratcliffe L.M., 2007

Variation in social rank acquisition influences lifetime reproductive success in black-capped

chickadees. Biol. J. Linn. Soc. 90:85-95

Smit A.F.A., Hubley R., RepeatModeler. Open-1.0.11, 2008–2015

(http://www.repeatmasker.org)

Srivastava A., Winker K., Shaw T.I., Jones K.L., Glenn T.C., 2012 Transcriptome analysis of a

North American songbird, Melospiza melodia. DNA Research. 19:325-333

Stanke M., Tzvetkova A., Morgenstern B., 2006 AUGUSTUS at EGASP: using EST, protein

and genomic alignments for improved gene prediction in the human genome. BMC Genome

Biology. 7(Suppl 1):S11. 1-8

Sutter N.B., Bustamante C.D., Chase K., Gray M.M., Zhao K., Zhu L., Padhukasahasram B.,

Karlins E., Davis S., Jones P.G., Quignon P., Johnson G.S., Parker H.G., Fretwell N., Mosher

D.S., Lawler D.F., Satyaraj E., Nordborg M., Lark K.G., Wayne R.K., Ostrander E.A., 2007 A

single IGF1 allele is a major determinant of small size in dogs. Science. 316(5821):112-115

Wang X., Wang L., 2016 GMATA: An Integrated Software Package for Genome-Scale SSR

Mining, Marker Development and Viewing. Front Plant Sci. 13;7:1350. doi:

10.3389/fpls.2016.01350. eCollection 2016

Warren W.C., Clayton D.F., Ellegren H., Arnold A.P., Hillier L.W., et al., 2010 The genome of a

songbird. Nature. 464(7289):757-62. doi: 10.1038/nature08819

Waterhouse R.M., Seppey M., Simão F.A., Manni M., Ioannidis P., Klioutchnikov G.,

Kriventseva E.V., Zdobnov E.M., 2017 BUSCO applications from quality assessments to gene

prediction and phylogenomics. Mol Biol Evol. doi: 10.1093/molbev/msx319

White S.A., 2010 Genes and vocal learning, Brain and Language. 115(1) 21-28.

DOI:10.1016/j.bandl.2009.10.002

Wicker T., Sabot F., Hua-Van A., Bennetzen J.L., Capy P., Chalhoub B., Flavell A., Leroy P.,

Morgante M., Panaud O., Paux E., SanMiguel P., Schulman A.H., 2007 A unified classification

system for eukaryotic transposable elements. Nat Rev Genet. 8(12):973-82

Z. Bao, S. R. Eddy., 2002 Automated de Novo Identification of Repeat Sequence Families in

Sequenced Genomes. Genome Research. 12:1269-1276

Zdobnov E.M., Apweiler R., 2001 InterProScan-an integration platform for the signature-

recognition methods in InterPro. Bioinformatics. 17:847-848

Zhang G., Li C., Li Q., Li B., Larkin D.M., Lee C., Storz J.F., Antunes A., et al., 2014

Comparative genomics reveals insights into avian genome evolution and adaptation. Science.

346(6215):1311-20. doi:10.1126/science.1251385

 Tables and Figures

516 Table 1: A comparison of assembly quality statistics from the initial shotgun sequencing

517 assembled by Meraculous and the final HiRise assembly.

	Meraculous Assembly	**Chicago HiRise Assembly**
Total length	972.4 Mb	978.3 Mb
Scaffold N50	33 kb	5.58 Mb
Scaffold N90	5 kb	303 kb
Scaffold L50	7,552 scaffolds	48 scaffolds
Scaffold L90	35,731 scaffolds	324 scaffolds
Longest scaffold	366,149	26,942,064
Number of scaffolds	74,832	13,785
Number of scaffolds > 1kb	74,806	13,768
Contig N50	22.5 kb	31.7 kb
Number of gaps	53,577	95,490
Percent of genome in gaps	1.427%	1.847%

520 Table 2: Number and percentage of repeats in the *M. melodia* genome assembly.

Classification	Number of copies	Percentage of assembly
LINEs	104,032	3.01
LTRs	85,276	2.83
SINEs	6,695	0.08
DNA Transposons	13,521	0.21

Unclassified	4,884	0.12
Total transposable elements	**214,408**	**6.25**
Satellites	569	0.00
Low complexity repeats	38,561	0.20
Microsatellites	192,996	0.90
Total	**446,534**	**7.35**

Table 3: Numbers of genes predicted in the *M. melodia* genome compared to *Taeniopygia guttata* (zebra finch) and *Ficedulla albicollis* (collared flycatcher), *Manacus vitellinus* (golden-collared manakin) and *Geospiza fortis* (medium ground finch).

	M. melodia	*T. guttata*[1]	*F. albicollis*[2]	*M. vitellinus*[3]	*G. fortis*[4]
Number of genes	15,086	17,561	16,763	18,976	14,388
Mean gene length (bp)	14,457	26,458	31,394	27,847	30,164
Mean CDS length (bp)	1,325	1,677	1,942	1,929	1,766
Number of exons	131,940	171,767	189,043	190,390	164,721
Mean exon length (bp)	153	225	253	264	195
Mean number of exons/gene	8.67	10.25	12.22	11.51	11.41
Number of introns	116,724	153,909	171,236	171,089	149,563
Mean intron length (bp)	1,695	2,930	3,257	3,294	2,813

[1]https://www.ncbi.nlm.nih.gov/genome/annotation_euk/Taeniopygia_guttata/103/

[2]https://www.ncbi.nlm.nih.gov/genome/annotation_euk/Ficedula_albicollis/101/

[3]https://www.ncbi.nlm.nih.gov/genome/annotation_euk/Manacus_vitellinus/102/

[4]https://www.ncbi.nlm.nih.gov/genome/annotation_euk/Geospiza_fortis/101/

Table 4: Numbers of ncRNAs predicted in the *Melospiza melodia* genome compared to

Taeniopygia guttata (zebra finch) and *Ficedulla albicollis* (collared flycatcher).

	M. melodia	*T. guttata*[1,2]	*F. albicollis*[1,3]
tRNA	267	184	179
miRNA	166	302	510
snRNA	16	44	32
snoRNA	154	241	199
rRNA	8	100	22
lncRNA	20	908	1473

[1]http://useast.ensembl.org/info/data/ftp/index.html

[2]https://www.ncbi.nlm.nih.gov/genome/annotation_euk/Taeniopygia_guttata/103/

[3]https://www.ncbi.nlm.nih.gov/genome/annotation_euk/Ficedula_albicollis/101/

Figure 1: Comparison of assembly contiguity.

Figure 2: Comparison of percentages of transposable elements (TEs) among related songbirds.

∗Zhang et al. Science. 2014 Dec 12;346(6215):1311-20. DOI: 10.1126/science.1251385.

Figure 3: Distribution of microsatellite repeat motif size classes in the *M. melodia* genome

assembly (details are given in Supplemental File S15).

Figure 4: Distribution of the top base-pair composition of microsatellite motifs in the *M. melodia*

genome

Figure 5: Jupiter plot correlating zebra finch and song sparrow genome assemblies, considering

scaffolds greater than 100 kbp in the reference zebra finch genome and the largest scaffolds

representing 95% of the query song sparrow genome.