Family History of Depression is Associated with Alterations in Task-Dependent Connectivity between the Cerebellum and Ventromedial Prefrontal Cortex

Lindsey J. Tepfer, Lauren B. Alloy & David V. Smith[*]

Department of Psychology, Temple University, Philadelphia, PA, USA

*Corresponding Author:

David V. Smith, Ph.D.
Assistant Professor of Psychology
Temple University
Weiss Hall, Room 825
1701 North 13th Street Philadelphia, PA 19122
Office Phone: 215-204-1552
Email: david.v.smith@temple.edu

Acknowledgments

This work was supported, in part, by grants from the National Institutes of Health (R21-MH113917 and R03-DA046733 to DVS) and the American Association for University Women (Career Development Grant to LJT). DVS was a Research Fellow of the Public Policy Lab at Temple University during the preparation of this manuscript (2019-2020 academic year). We thank Ka-Yi Chat for helpful feedback on the manuscript.

Abstract

Background A family history of major depressive disorder (MDD) increases the likelihood of a future depressive episode, which itself poses a significant risk for disruptions in reward processing and social cognition. However, it is unclear whether a family history of MDD is associated with alterations in the neural circuitry underlying reward processing and social cognition.

Methods We subdivided 279 participants from the Human Connectome Project into three groups: 71 with a lifetime history of MDD, 103 with a family history of MDD (FH), and 105 healthy controls (HC). We then evaluated task-based fMRI data on a social cognition and a reward processing task and found a region of the ventromedial prefrontal cortex (vmPFC) that responded to both tasks, independent of group. To investigate whether the vmPFC shows alterations in functional connectivity between groups, we conducted psychophysiological interaction (PPI) analyses using the vmPFC as a seed region.

Results We found that FH (relative to HC) was associated with increased sadness scores, and MDD (relative to both FH and HC) was associated with increased sadness and MDD symptoms. Additionally, the FH group had increased vmPFC functional connectivity within the nucleus accumbens, left dorsolateral PFC, and subregions of the cerebellum relative to HC during the social cognition task.

Conclusions These findings suggest that aberrant neural mechanisms among those with a familial risk of MDD may underlie vulnerability to altered social cognition.

Keywords: Depression, Family History, Functional Connectivity, fMRI

Introduction

Major Depressive Disorder (MDD) substantially interferes with the ability to carry out a normal life. As a leading contributor to global disability (Ferrari et al., 2013; World Health Organization, 2017), MDD carries profoundly disruptive symptoms that range from hopelessness (Abramson et al., 1989; Iacoviello et al., 2010) to diminished motivation (Sherdell et al., 2012; Treadway et al., 2012) as a part of anhedonia (American Psychiatric Association, 2013; Pizzagalli, 2014), and is a prominent predictor of suicidality (Klonsky et al., 2016; Nock et al., 2009). Symptom recurrence is likely (Kovacs et al., 2016; Solomon et al., 2000), and each subsequent episode increases the chance that more will follow (Colman et al., 2011; Hoertel et al., 2017). Depression is highly comorbid with substance use disorders (Swensden & Merikangas, 2000), including alcohol (Fergusson et al., 2009; Pedrelli et al., 2016; L. E. Sullivan et al., 2005) and tobacco (Covey et al., 1998; Weinberger et al., 2017), and there is evidence that the relationship between MDD and substance use may impart similar patterns in the children of those who suffer from depression (Luthar, Merikangas, Rounsaville et al., 1993) as both are highly familial (Merikangas &Swendsen, 1997). Importantly, the probability that an individual will experience MDD during their lifetime increases for those with a family history (FH) of depression (Klein et al., 2005, 2013; Monroe et al., 2014; Zimmermann et al., 2008). Indeed, those with depressed first-degree relatives are nearly three times as likely to develop the disorder later in life (Sullivan et al., 2000). Upon the onset of MDD, the systems that support social cognition also are affected (Cusi et al., 2012; Derntl et al., 2011), which limits the capacity to maintain supportive connections with others and negatively impacts recovery (George et al., 1989; Santini et al., 2015). Understanding the abnormalities among the neural mechanisms underlying social cognition and reward sensitivity in MDD, and how they are vulnerable in those with a familial risk of MDD,

stands to elucidate the relationship between early pathophysiology in the brain and subsequently altered cognition.

A wealth of research suggests that many of the symptoms associated with MDD involve abnormalities in specific neural circuits. For example, studies have found grey matter volume reductions in frontotemporal areas (Cai et al., 2014; Serra-Blasco et al., 2013), the hippocampus (Arnone et al., 2013; Zhao et al., 2014), the orbitofrontal cortex (OFC; (Bremner et al., 2002; Lacerda et al., 2004), the ventromedial prefrontal cortex (vmPFC; (Wise et al., 2017), the dorsomedial prefrontal cortex (dmPFC), the anterior cingulate cortex (ACC), and the bilateral insula (Caetano et al., 2006; Salvadore et al., 2011). Recent reports also indicate a putative role for cerebellar grey matter volume decrease in depression (Wise et al., 2017). Indeed, abnormal structural circuitry in the cerebellum appears to underlie many common mental disorders (Romer et al., 2018) with a handful of studies indicating subregional specificity, such as the vermis and posterior cerebellum. For example, lesioning in the posterior and vermis subregions often is associated with emotion dysregulation and other affective symptoms (Schmahmann, 2004; Schmahmann et al., 2007), and increased anterior vermis volume has been found in MDD patients (Yucel et al., 2013). Likewise, white matter fractional anisotropy investigations also show disruptions across the lifespan among individuals with depression (Alexopoulos et al., 2008; Bae et al., 2006; Cruwys et al., 2014; Cullen et al., 2010; Steingard et al., 2002; Taylor et al., 2004; Yang et al., 2007). In conjunction with structural changes, MDD is associated with functional aberrations in systems associated with reward processing. Evidence of differences in activation and connectivity in healthy individuals relative to those with MDD typically implicate the OFC (Cheng et al., 2018; Ng et al., 2019), the ventral striatum (VS; (Furman et al., 2011;

Kumar et al., 2018; Robinson et al., 2012), and the vmPFC (Koenigs et al., 2008; Myers-Schulz & Koenigs, 2012). Similarly, alterations in function also have been found in the cerebellum among depressed patients, including increased blood flow to the cerebellar vermis (Bench et al., 1992; Dolan et al., 1992), increasing vermis-PCC functional connectivity with symptom severity (Alalade et al., 2011), left-posterior cerebellar activation during rewarded sustained attention (Chantiluke et al., 2012), as well as decreases in vmPFC-bilateral lobule VIIb functional connectivity (Liu et al., 2012). Together, this work suggests that we may be able to understand some of the canonical symptoms seen in MDD symptomatology by taking a closer look at the neural correlates that underlie them.

Tied to the functional and structural impairments seen among reward processing systems (e.g., the striatum), a hallmark symptom of MDD involves the marked loss in the ability to experience pleasure (Cooper et al., 2018). Individuals with MDD often endure significant decreases in motivation (Hershenberg et al., 2016), diminishment in emotional responses to things that are pleasant (Alloy et al., 2016), and no longer pursue the interests they once found enjoyable (Fried & Nesse, 2014). Moreover, research implicates that social rewards share the neural circuitry represented by non-social rewards (Bhanji & Delgado, 2014; Fareri & Delgado, 2014). Thus, abnormalities in reward processing often overlap with social cognitive deficits, as reward system activation is critical for facilitating positive social exchanges (Carta et al., 2019; Smith & Delgado, 2015; Wang et al., 2016). Deficits in social cognition persist throughout recovery from MDD and are associated with later relapse (Inoue et al., 2004, 2006).

Individuals with a FH of depression also appear to be at risk for similar degrees of deviation in reward and social processing early on (Monk et al., 2008; Weinberg et al., 2015). For example, individuals with a familial risk of depression show abnormal responses to reward and punishment (McCabe et al., 2012), have reduced activity in the VS and ACC in response to social reward during adolescence (Olino et al., 2015), and decreased VS and anterior insula activity in response to non-social rewards during childhood (Luking et al., 2016). Furthermore, those with a familial history of MDD begin to show alterations in theory of mind (ToM) processing - the ability to infer the mental states of others (Harkness et al., 2011).

However, despite the work underscoring the relationship between reward, social cognition, and depression, it remains unclear whether individuals with a familial history of MDD demonstrate neural alterations associated with processing social stimuli in the absence of reward. Investigating how the neural and behavioral alterations observed in MDD emerge in association with a FH is key to understanding the mechanisms that drive these changes. Moreover, evaluating cases with a predisposition to such alterations may inform current treatment practices that can help provide early intervention upon the identification of at-risk populations. Thus, in the present study, we investigate whether a FH of depression, in the absence of a personal diagnosis, is associated with altered social cognition and reward processing in the brain. Our analyses focused on three main hypotheses. First, we expected to see increased uncertainty during ToM processing, as well as increased depressive symptoms, reports of sadness, alcohol, and tobacco use among individuals with a personal history of MDD, relative to those with a FH of MDD or HCs. Second, we predicted that the MDD group would show blunted temporoparietal junction (TPJ) response and increased vermis and posterior cerebellar activity during a ToM

task, and decreased striatal activation in response to reward during a gambling task. Our third hypothesis was that the MDD group would show decreased connectivity between the cerebellum's bilateral lobule VIIb and the vmPFC during the social task. For all of these hypotheses, we anticipated a dose-dependent type of effect across the three groups, in which the MDD group would show the highest magnitude of difference relative to the FH and HC groups, with the FH group resembling any effect seen in the MDD group, albeit at a smaller magnitude. For example, should the MDD group demonstrate hypothesized blunted TPJ activity during the ToM task, the FH group would similarly show decreases in TPJ activity relative to HCs, but not decreased to the same extent as the MDD group.

Materials and Methods

Participants. The sample included 279 participants (males, 120; females, 159; ages, 22-36; mean ± SD, 28.45 ± 3.75 years) selected from the Human Connectome Project (HCP) dataset, a large collection of primarily healthy young adults (N=1206) from the 1200 Subjects Release (WU-Minn HCP consortium S1200 Data). The study protocol was performed in compliance with the Code of Ethics of the World Medical Association (Declaration of Helsinki). Participants provided written informed consent, and procedures were approved by the ethics committee in accordance with guidelines of WU-Minn HCP. We subdivided participants into three groups. Individuals in the MDD group (N=71) were selected if they had a lifetime history of a DSM-IV Major Depressive Episode diagnosis. Individuals in the FH group (N=103) were selected if they had not received a DSM-IV Major Depressive Episode diagnosis in their lifetime, but either their mother or father had. We note that the 71 MDD and 103 FH comprised all participants who met criteria for these groups from the full 1206 HCP sample. HCs (N=105) consisted of participants who had not

received a DSM-IV Major Depressive Episode diagnosis in their lifetime, and had no FH of psychiatric or neurologic disorders. The number of HCs that qualified based on these criteria was large (N=573); to facilitate comparison, we randomly selected HC participants to match our largest experimental group (N=105).

Exclusion Criteria Participants were excluded for head motion if their average relative head motion (calculated using a root-mean-squared approach; Jenkinson et al., 2002) exceeded the 75th percentile plus the value of 150% of the interquartile range of average relative head motion for all subjects (i.e., a standard boxplot threshold; (Smith et al., 2014). Additionally, participants who met DSM-IV criteria for alcohol abuse were excluded from the current study.

Task Paradigms

All 279 participants in the current study performed 7 tasks in the scanner across two sessions, with two runs of each task. Here, we analyze data from only 2 tasks: social cognition and reward processing. Each task involved alternating blocks of stimuli. The descriptions of the two tasks of interest here are adaptations of the details provided elsewhere (Barch et al., 2013). The social cognition task required participants to indicate whether or not they interpreted a social interaction among brief animations of geometric shapes, a robust and reliable measurement of theory of mind processing. The reward processing task prompted participants to guess whether a 'mystery card' held a value of higher or lower than 5, and subsequently received feedback on the outcome of their choice. For more details regarding the block timing and composition of each task, see **Supplemental Information**.

Image Acquisition

All 279 participants in the current study underwent T1 and T2-weighted structural scans, resting-state and task-based fMRI scans, and diffusion-weighted MRI scans, all via a 3T Siemens scanner with a standard Siemens 32 channel RF head coil. Identical multi-band EPI sequence parameters leveraged high spatial resolution at 2mm isotropic using the following parameters: repetition time (TR) = 720 ms; echo time (TE) = 33.1 ms, 72 slices, 2mm isotropic voxels, using a multi-band acceleration factor of 8 (Feinberg et al., 2010; Moeller et al., 2010; Setsompop et al., 2012; Xu et al., 2013). The current investigation used minimally preprocessed BOLD fMRI data to increase ease of use and mitigate the possibility of complex issues relating to imaging data quality; the general preprocessing steps are presented in full detail in previous reports (Glasser et al., 2013; Smith et al., 2013; Uğurbil et al., 2013; Van Essen et al., 2013). Tasks were performed twice, with one run in a left-to-right direction, and then reversed in the second run. In addition to the minimal preprocessing already featured in the HCP dataset, we identified and removed artifacts sourced from excess head-motion using the ICA-AROMA Software Package. Preceding ICA-AROMA, we first smoothed the data with a spatial smoothing of 6mm and the entire 4D dataset received grand-mean intensity normalization using a single multiplicative factor. This grand-mean intensity normalization step was applied to ensure scaling is not shifted following denoising and the normalization (mean = 10,000) is maintained. For convenience, the current study worked solely in volumetric MNI-space, although future work would benefit from transitioning to a CIFTI space (Coalson et al., 2018; Glasser, Coalson, et al., 2016; Glasser, Smith, et al., 2016).

fMRI Analysis

We analyzed task-based fMRI data using FEAT (FMRI Expert Analysis Tool) Version 6.00, part of FSL (FMRIB's Software Library; (Jenkinson et al., 2012)). Specifically, we used FILM (FMRIB's Improved Linear Model) with local autocorrelation (M. W. Woolrich et al., 2001) correction to assess task-evoked changes in neural responses to social and reward stimuli employing two main models. Both models employed a general linear model (GLM) approach, convolved using the canonical hemodynamic response function. Each first level GLM included two regressors to account for the block structure in a task: in the gambling task, these regressors corresponded to reward and loss, and in the social task, mental trials and random trials. The reward regressor was modeled from the onset of the first trial of the block, which consisted of mostly reward until the offset of that block; similarly, the punishment regressor was elicited the same way, but drawn from the mostly-loss blocks. The mental regressor was modeled from the onset of the first block of the run that consisted of mostly mental videos until the offset of that run; the random regressor was similarly derived from the run that consisted of mostly random trials.

To investigate task-evoked changes in brain connectivity, we utilized a psychophysiological interaction (PPI) analysis (Friston et al., 1997). Our PPI models focused on cerebellar coupling with vmPFC and were later expanded in an exploratory whole brain analysis, using the vmPFC as the seed region. The vmPFC region of interest (ROI) was defined based on a preliminary analysis using FLAME. For this analysis, we thresholded and corrected for multiple comparisons within a priori ROI (http://aspredicted.org/blind.php?x=8qw2h3) at the voxel level. ROIs were identified via a Cerebellar Atlas in MNI152 space after normalization with FNIRT and the

Harvard-Oxford cortical and subcortical atlases in FSL. In addition to the same regressors used in the activation models, our PPI models included regressors to represent the activation time course in connectivity between the vmPFC seed region and cerebellar target regions. Similar to our analysis of activation, we evaluated two distinct PPI models for each task: the social cognition and reward processing tasks.

To combine data across runs for our second-level analysis, we used a fixed effects model, with random effects variance forced to zero in FLAME (Beckmann et al., 2003; M. Woolrich, 2008; M. W. Woolrich et al., 2004). Next, we combined data across participants using FLAME (FMRIB's Local Analysis of Mixed Effects) stage 1 (Beckmann et al., 2003; M. Woolrich, 2008; M. W. Woolrich et al., 2004). To detect group differences in neural activity for the social cognition and reward processing tasks, and to control for other factors that influence our results, we used a GLM consisting of 11 regressors. The 11 regressors contained data describing participant head movement during the task, gender, depressive symptoms, sadness scores, impulsivity, alcohol and tobacco use. Gaussianized z-statistic images were thresholded with voxel-level GRF-theory-based maximum height thresholding with a corrected significance threshold of $p=0.05$ (Worsley, 2001). Our model also included F-tests to directly evaluate our ability to reject the null hypothesis that MDD = FH = HC with regard to each of our hypotheses. We controlled for family structure in the data (Winkler et al., 2015) using PALM (Winkler et al., 2014) to evaluate pre-registered hypotheses regarding group differences in behavior and neural responses tied to activation and connectivity. We note, however, that the seed for our connectivity analysis (vmPFC) was identified through conventional parametric statistics since the goal of that analysis was simply to identify regions that respond similarly to social and

reward, irrespective of group. Our application of PALM was one-tailed, and we controlled for the number of comparisons across contrasts using the -corrcon flag, which includes FWER-correction of p-values. We also enabled threshold-free cluster enhancement (TFCE) inference for univariate tests, and was permuted 5000 times for all behavioral, activation, and PPI analyses.

In addition, we performed t-tests between MDD, FH, and HCs to determine whether the groups differed by head motion variables in both the gambling and social cognition task. We found no difference among groups on the gambling task between MDD and FH (t(172)=-1.28, p=0.20), MDD and HC (t(174)=-0.13, p=0.88), FH and HC (t(206)=1.26, p=0.20), nor any difference among groups on the social cognition task between MDD and FH (t(172)=-0.99, p=0.31), MDD and HC (t(174)=0.01, p=0.99), FH and HC (t(206)=1.15, p=0.24).

Results

Behavioral and Clinical Analyses. Our analyses began by testing whether a FH of depression was associated with higher depression symptoms and sadness experienced across one's lifetime. To compare depression symptoms, we evaluated the average number of lifetime depressive symptoms reported by each participant who met DSM-IV criteria. We found that individuals with a FH reported more lifetime depressive symptoms across their lives compared to HCs (t(206)=2.54, p=0.011). Individuals in the depression group had higher depressive symptom rates than both FH (t(172)= 19.22, p<0.01) and HC (t(174)=27.00, p<0.01; see Figure 1A) groups. Similarly, we used unadjusted sadness scores as measured by the NIH Toolbox Sadness Survey to compare self-reports of sadness across groups. The NIH Toolbox Sadness Survey is a subcomponent of a larger battery, the NIH Toolbox Emotion Battery, a standardized form of measuring emotional health across the lifespan (www.nihtoolbox.org). We found that unlike the

depressive symptoms, sadness scores in participants with a FH of depression did not differ significantly from HCs ($t(206)=1.88$, $p=0.06$). However, as expected, the levels of sadness reported by the depression group surpassed both FH ($t(172)=4.39$, $p<0.01$) and HC groups ($t(174)=6.09$, $p<0.01$, See Figure 1b). Additionally, we tested for differences in alcohol and tobacco use (in the past 7 days) between groups, as previous work has found a relationship between tobacco use (Edwards et al., 2011) or occasional bouts of heavy drinking (Manninen et al., 2006) and depression. However, we did not find significant differences in either alcohol or tobacco use across groups (see Table 1).

We hypothesized that the depression group would be more unsure about the social nature of the stimuli relative to FH or HC groups. To test this hypothesis, we compared the selections each group made while engaging in the ToM task in the scanner. Contrary to our expectations, we initially found that the depression group had a higher percentage of 'social' selections relative to FH ($t(172)=5.81$ $p<0.01$) and HC ($t(174)=5.66$, $p<0.01$) groups. The depression group also initially demonstrated lower percentages of 'random' selections relative to FH ($t(172)=-3.89$ $p<0.01$) and HC ($t(174)=-3.87$, $p<0.01$) groups. Depression and FH ($t(172)=-1.81$ $p=0.07$), depression and HCs ($t(174)=-0.57$, $p=0.57$), FH and HCs ($t(206)=1.41$, $p=0.16$) did not differ significantly with respect to percentage of 'unsure' selections throughout the task (see Figure 1c). However, after controlling for family structure in the data, we found no differences in performance between groups on the social cognition task (all $-\log_{10}(p)$ values < 1.3010, where p is 0.05, corrected for multiple comparisons across voxels and contrasts).

Figure 1: Self-reported depression symptoms and levels of sadness across depression, family history, and healthy control groups. ***A***, Depression (MDD) group shows higher average depression symptoms compared to both individuals with a family history of depression (FH; $t(172)$= 19.22, $p<0.01$, d=2.82) and healthy controls (HCs; $t(174)$=27.00, $p<0.01$. d=24.00). Individuals in the FH group show higher depression symptoms compared to HCs ($t(206)$=2.54, p=0.011, d=0.35). ***B***, Unadjusted sadness scores are significantly higher in MDD when comparing to both FH ($t(172)$=4.39, $p<0.01$, d=0.68) and HC groups ($t(174)$=6.11, $p<0.01$, d=0.93), but did not reach our significance threshold between FH and HCs ($t(206)$=1.88, p=0.06, d=0.26). ***C***, Average social, random, and unsure percent responses to a Theory of Mind (ToM) social cognition task presented to participants during the fMRI scanning paradigm. Initially, MDD showed a higher percentage of 'social' selections relative to FH ($t(172)$=5.81 $p<0.01$, d=0.86) and HC ($t(174)$=5.66, $p<0.01$, d=0.83) groups. MDD also initially demonstrated lower percentages of 'random' selections relative to FH ($t(172)$=-3.89 $p<0.01$, d=-0.59) and HCs ($t(174)$=-3.87, $p<0.01$, d=-0.59). MDD and FH ($t(172)$=-1.81 p=0.07), MDD and HCs ($t(174)$=-0.57, p=0.57, d=-0.08), FH and HCs ($t(206)$=1.41, p=0.16, d=0.19) did not differ significantly with respect to percentage of 'unsure' selections throughout the task. However, after controlling for family structure in the data, any differences in performance between groups on the social cognition task were eliminated (all $-\log_{10}(p)$ values < 1.3010, where p=0.05, corrected for multiple comparisons across voxels and contrasts).

Group	Participants (N)	Gender	Alcohol Use (drinks/7 days)	Tobacco Use (cigarettes/7 days)
Depression (MDD)	71	61% F; 40% M	4.661	5.450
Family history (FH)	103	62% F; 38% M	4.165	6.533
Healthy controls (HC)	105	50% F; 50% M	4.295	5.123

Table 1: Percentage of male and female participants, as well as average tobacco (measured in number of cigarettes) and alcohol (measured in number of drinks) used in the past 7 days across each group. Groups did not differ significantly in their use of alcohol over the past 7 days. Alcohol use in MDD and FH ($t(172)$=0.47, p=0.63, d=0.07), MDD and HCs ($t(174)$=0.34, p=0.73, d=0.05), FH and HCs ($t(206)$=-0.15, p=0.88, d=-0.02). Similarly, there were no differences in tobacco use between MDD and FH ($t(172)$= -0.36, p=0.71, d=-0.05), MDD and HCs ($t(174)$= 0.12, p=0.90, d=0.01), and FH and HCs ($t(206)$=0.51, p=0.61, d=0.07).

ROI Analyses

Next, we evaluated whether the social cognition and reward processing tasks activate canonical mentalizing and reward processing regions respectively, and if activity overlaps. We found that independent of group, the social cognition task activated the TPJ on the mental > random trial contrast, and the reward processing task activated the nucleus accumbens on the reward > punishment trial contrast. A subsequent conjunction analysis revealed that a region of the vmPFC responded to both tasks, independent of groups (Figure 2). We then used this region of the vmPFC as a seed for psychophysiological interaction (PPI) analyses to evaluate vmPFC connectivity patterns with the cerebellum and later, across the entire brain.

Figure 2: Activation and Conjunction Analyses. *A*, Independent of group, the social cognition task evoked activity in the temporoparietal junction specific to Mental > Random trials contrasts. *B*, Similarly, the reward processing task evoked striatal activation on the Reward > Punishment trial contrasts. *C*, Both tasks also demonstrate shared activation in a region of the ventromedial prefrontal cortex (vmPFC). Gaussianized z-statistic images were thresholded with voxel-level GRF-theory-based maximum height thresholding with a corrected significance threshold of p=0.05.

We then evaluated the presence of dose-dependent responsivity among the groups during the social cognition task. We predicted that during this task, MDD would demonstrate blunted TPJ activation, but increased cerebellar activity, particularly in the vermis and posterior hemispheres relative to HC and FH groups. Our dose-dependency hypotheses also anticipated that while the

MDD group would show the largest responses (either by way of increase or decrease), the FH group would trend towards the pattern exhibited by the depression group, with blunted TPJ activation and increased cerebellar activity relative to HCs. Additionally, we expected to see that the MDD group would have blunted striatal activation in response to the reward processing task, relative to HC and FH groups, while the FH group also would show blunted striatal activation relative to HCs. Despite these hypotheses, we found no differences in TPJ or cerebellar activation within the social task, nor striatal activation differences during the reward processing task across the groups.

Table 2: vmPFC Seed Functional Connectivity by task, condition, and group comparison							
vmPFC-Cerebellar PPI analyses							
Task	Condition	Comparison	Probabilistic Anatomical label	x	y	z	$-\log_{10}(p)$
Gambling	Rew>Pun	MDD>FH	Right Crus I (15%), Right Crus II (5%)	9	-84	-20	1.6307
Gambling	Pun>Rew	FH>MDD	Right Crus I (15%), Right Crus II (5%)	9	-84	-20	1.6307
Social	Mental	MDD>HC	Right VIIa (45%), Right VIIb (45%)	14	-72	-54	1.6497
Social	Mental	FH>HC	Left Crus I (100%)	-47	-68	-36	1.6055
Social	Random	FH>HC	Left Crus I (71%), Left VI (28%)	-36	-61	-28	1.6126
vmPFC-whole brain PPI analyses							
Social	Mental	FH>HC	Insular Cortex (39%)	-30	13	6	1.4294
			Left Putamen (98%)	-17	9	-7	
			Frontal Pole (78%)	-36	42	25	
Social	Random	FH>MDD	Frontal Pole (90%)	-42	53	-3	1.3851

Table 2: Regions that demonstrate significant activation above the $-\log_{10}(p)$ value cutoff of 1.3010 (where p is 0.05, corrected for multiple comparisons across voxels and contrasts) among Reward (Rew), Punishment (Pun), Mental, and Random conditions for the Gambling and Social tasks, respectively. Sites of functional connectivity activation marked with probabilistic anatomical labels correspond to anatomical coordinates (x,y,z) for each group comparison: major depressive disorder (MDD), family history (FH) and healthy controls (HC).

We also expected to see decreased vmPFC-cerebellar (specifically with respect to the bilateral lobule VIIb) connectivity during the social cognition task for the MDD group relative to FH and HC groups. As with our activation hypotheses, we expected that the FH group also would show decreased connectivity between the vmPFC and cerebellar subregions, albeit relative to HCs. Our results do not show any differences in vmPFC-cerebellar functional connectivity between MDD and FH groups, but are seen when comparing MDD and HC groups, such that MDD groups show

a small region of increased vmPFC-cerebellar connectivity during the mental condition (see Table 2). Nevertheless, we find increased vmPFC-cerebellar functional connectivity in the FH group relative to HCs. Specifically, relative to HCs, the FH group shows increases in connectivity between the vmPFC and the left crus I of the cerebellum during the mentalizing trials. Random trials, on the other hand, also show increased connectivity between the vmPFC and left crus I, as well as the left VIIb region of the cerebellum, with the FH group again showing greater connectivity than the HCs in particular (see Figure 3).

Figure 3: Psychophysiological (PPI) analysis using vmPFC as the seed region. *A*, Relative to healthy controls, the family history group showed increases in task-based functional connectivity between the vmPFC and the left crus I of the cerebellum during the "Mentalizing" trials. Areas of activation passed an a priori small volume correction (cerebellum) using threshold-free-cluster enhancement (TFCE) and permutation-based testing using PALM. *B*, Interrogation of the left crus I region of the cerebellum revealed increased connectivity relative to the decrease in activation seen in healthy controls during the mental trials. *C*, Increased connectivity between the vmPFC and left crus I and left VIIb regions of the cerebellum was selective for the random stimuli in the social task, with the family history

group showing a greater response than the healthy controls. Areas of activation passed an a priori small volume correction (cerebellum) using threshold-free-cluster enhancement (TFCE) and permutation-based testing using PALM. **D**, Interrogation of the left crus I and left VIIb regions of the cerebellum showed the family history group had a larger increase in connectivity relative to healthy controls during the mental trials.

Exploratory Analyses

We expanded our search beyond the pre-registered hypotheses and a priori cerebellar target to include a whole-brain analysis of vmPFC connectivity. We again find functional connectivity differences primarily for the FH and HC groups during the social cognition task. Specifically, individuals with a familial risk of depression, relative to healthy controls, demonstrated increased vmPFC-nucleus accumbens and vmPFC-dorsolateral PFC functional connectivity during the mentalizing trials (see Figure 4).

Figure 4: Exploratory whole-brain analysis. A, Relative to healthy controls, the family history group showed greater task-based functional connectivity between the vmPFC and the nucleus accumbens (NAcc) during the mental trials of the social cognition task. Areas of activation passed a whole brain correction using threshold-free-cluster enhancement (TFCE) and permutation-based testing. B, Interrogation of the NAcc revealed increased connectivity in the family history group, whereas the healthy control group demonstrated a relative decrease in connectivity during the mental trials.

Discussion

A deeper understanding of the neural mechanisms underlying social cognition and reward processing can elucidate how these systems typically function, and the conditions under which they are susceptible to change. Mapping task-based neural activity among those with a well-established risk factor for MDD allows us to probe this susceptibility to change in reward and social systems. Moreover, capturing the state of the brain before MDD onset in an at-risk group is especially relevant to clinical efforts in treatment and intervention. Thus, the goal of the current study centered on evaluating task-based neural processing of social and reward systems in those with a FH of major depressive disorder, using data from the Human Connectome Project. In line with our expectations, we found that relative to HCs, individuals with a FH of MDD demonstrate increases in depressive symptoms, and the MDD group showed increased sadness scores and depressive symptoms relative to both FH and HC groups. Yet, we did not find significant neural differences between HCs and those with a personal history of MDD on either social or reward processing tasks as expected. Nevertheless, we confirmed our initial hypothesis that the FH group would show neural alterations relative to HCs during the social cognition task. Specifically, we found that relative to HCs, individuals with a FH of MDD demonstrated increased task-dependent functional connectivity between the vmPFC and the nucleus accumbens, left dorsolateral PFC, and subregions of the cerebellum during a mentalizing task. Altogether, these results reveal that in addition to upticks in pre-clinical reports of depressive symptoms, individuals with a FH of depression may be at risk for neural abnormalities underlying ToM processes, with altered cerebellar-vmPFC functional connectivity contributing to this difference.

A host of recent studies have begun to reveal the functional role of the cerebellum in social, cognitive, and affective neuroscience (Van Overwalle et al., 2014, 2015). For example, the

cerebellum has been linked to social reward (Carta et al., 2019), social preference (Badura et al., 2018), and emotional processing (Baumann & Mattingley, 2012). Of particular relevance to our current results, the cerebellum has been linked to mentalizing processes by way of connectivity with the cortex, including the TPJ (Van Overwalle et al., 2019) and dorsomedial PFC (Van Overwalle et al., 2015). Cerebellar connectivity also has been tied to the default mode network (Buckner et al., 2011), which participates in social inference (Buckner & DiNicola, 2019). In light of this past work, the present study dovetails with the existing literature by providing additional evidence demonstrating involvement of cerebellar connectivity during mentalizing. Specifically, our findings implicating task-based vmPFC-cerebellar connectivity differences in a ToM task appears to fall in line with previous suggestions that the cerebellum is involved in social reasoning (Van Overwalle et al., 2019).

Unlike prior studies of the cerebellum, our work highlights a novel role for the region's involvement in the vulnerabilities inherited through a FH of depression. Whereas investigations of autism (Becker & Stoodley, 2013), schizophrenia (Andreasen & Pierson, 2008), and major depressive disorder (Liu et al., 2012) have shown alterations in the cerebellum related to social and affective processing, these data demarcate populations that already have developed a disorder. As such, relatively little is known about the neural signatures involved in a FH of depression as it relates to social cognition. Through the identification of altered task-based cerebellar functional connectivity with the vmPFC, we shed new light on possible mechanisms at play that may lead to the early stages of MDD. In addition to building a deeper understanding of the cerebellum's role in social cognition, this work may inform interventions geared towards mitigating later emergence of MDD. Indeed, considering the cerebellum's influence on social and cognitive development, the region previously has been suggested as a potential site for treatment (S. S. H. Wang et al., 2014).

Moreover, altered functional connectivity between the cortex and cerebellum reliably can predict the presence of depression (Ma et al., 2013). The results of the present study indicate that consideration of the cerebellum as a therapeutic target may be promising early on, before the onset of MDD.

Although our study demonstrates that task-dependent connectivity with the cerebellum is altered in those with a FH of depression, we note that our findings are accompanied by some limitations. First, we note that the current work uses a block design and our analyses focus on differences between blocks, consistent with other work using the HCP data (e.g., Barch et al., 2013). Yet, some studies using the HCP data have suggested that the reward task also could be analyzed based on individual events, which may help separate processes tied to valence and salience (Zhang et al., 2017). Future studies may be able to build on our findings by examining salience and valence and by considering other datasets that are optimized for such questions (e.g., ABCD). Additionally, our results that vmPFC-cerebellar connectivity was enhanced in the FH group relative to HCs was not specific to the mental or random conditions in the social task. We speculate that the lack of differences between the two conditions in the social task could be due to the nature of the task. Indeed, although the task used by the HCP is a robust probe of TPJ activation, it may not evoke similar levels of activation in other regions that process more subtle differences in how participants process and interpret a range of social interactions. Future work building on these findings would do well to involve measures of social cognition that characterize mental representations of others in a variety of ways. For instance, it may be particularly relevant to examine social interactions as seen in the virtual school paradigm (Jarcho et al., 2016), as this may better probe how the cerebellum processes the mental states of characters in a more ecologically valid setting.

In addition to limitations with the social task, we also note that our study may not be able to speak directly to the mechanisms underlying risk for developing MDD, given the cross-sectional nature of the Human Connectome Project. Although our original hypothesis—that aberrant patterns of connectivity would deviate further from HCs when examining people with a personal history of depression—was not supported by our findings, we note that this is an area where longitudinal data would be necessary for understanding whether and how altered connectivity is associated with the development of MDD. To address this question, future research could leverage other open datasets that include longitudinal assessments of cerebellar connectivity, such as the Adolescent Brain Cognitive Development study (ABCD; (Casey et al., 2018). Thus, in the absence of longitudinal data, it is unclear whether neural differences between FH and HC groups can predict a subsequent MDD diagnosis.

Despite these limitations, our study demonstrates that relative to HCs, individuals with a FH of MDD exhibit altered task-dependent vmPFC-cerebellar activity during a social cognition task. Previous research shows that the cerebellum plays a key role in processing social, affective and rewarding stimuli and is a region of interest in psychiatric illnesses, including MDD (Konarski et al., 2005). Evidence for neural alterations underlying social cognition in advance of a personal episode of MDD implies the possibility of early intervention and support that may mitigate later emergence of the disorder. Abnormal neural connectivity has been documented in subclinical depression in another study using the HCP dataset (Ely et al., 2016), as well as other disorders (Young et al., 2015) and may play a central role in the development of psychiatric illnesses (Zhang et al., 2015) and maladaptive behaviors (Diehl et al., 2018) such as substance use (Cheng et al., 2019). From a clinical lens, the identification of neural alterations and their ability to establish a diagnosis or predict how an individual will respond to treatment remains unclear

(Savitz et al., 2013). Yet, as research accumulates data detailing the mechanisms driving neural alterations, future work may more reasonably employ emergent technology—such as brain stimulation—to intervene (Diehl et al., 2018). Considering the limits MDD places on social cognition (Knight et al., 2018), addressing abnormalities in the underlying neural mechanisms of social cognition in particular lends itself to the development of treatments that facilitate social support and connection, a leading protective factor against future depressive episodes (Cruwys et al., 2014). Additional work must continue to investigate how the cerebellum connects with other regions across a variety of affective disorders to understand the full spectrum of the site's role in healthy and abnormal social cognition alike.

Data Availability Statement

Data were provided [in part] by the Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at Washington University. This study was pre-registered, and can be found at http://aspredicted.org/blind.php?x=8qw2h3. Study materials and code is available on Open Science Framework (DOI 10.17605/OSF.IO/JU32V). Images are available on the NeuroVault repository at https://identifiers.org/neurovault.collection:6130.

References

Abramson, L. Y., Metalsky, G. I., & Alloy, L. B. (1989). Hopelessness Depression: A Theory-Based Subtype of Depression. *Psychological Review*. https://doi.org/10.1037/0033-295X.96.2.358

Alalade, E., Denny, K., Potter, G., Steffens, D., & Wang, L. (2011). Altered cerebellar-cerebral

functional connectivity in geriatric depression. *PLoS ONE*, *6*(5).

https://doi.org/10.1371/journal.pone.0020035

Alexopoulos, G. S., Murphy, C. F., Gunning-Dixon, F. M., Latoussakis, V., Kanellopoulos, D.,

Klimstra, S., Lim, K. O., & Hoptman, M. J. (2008). Microstructural white matter

abnormalities and remission of geriatric depression. *American Journal of Psychiatry*.

https://doi.org/10.1176/appi.ajp.2007.07050744

Alloy, L. B., Olino, T., Freed, R. D., & Nusslock, R. (2016). Role of Reward Sensitivity and

Processing in Major Depressive and Bipolar Spectrum Disorders. In *Behavior Therapy*.

https://doi.org/10.1016/j.beth.2016.02.014

American Psychiatric Association. (2013). American Psychiatric Association, 2013. Diagnostic

and statistical manual of mental disorders (5th ed.). In *American Journal of Psychiatry*.

https://doi.org/10.1176/appi.books.9780890425596.744053

Andreasen, N. C., & Pierson, R. (2008). The Role of the Cerebellum in Schizophrenia. In

Biological Psychiatry. https://doi.org/10.1016/j.biopsych.2008.01.003

Arnone, D., Mckie, S., Elliott, R., Juhasz, G., Thomas, E. J., Downey, D., Williams, S., Deakin, J. F.

W., & Anderson, I. M. (2013). State-dependent changes in hippocampal grey matter in

depression. *Molecular Psychiatry*. https://doi.org/10.1038/mp.2012.150

Badura, A., Verpeut, J. L., Metzger, J. W., Pereira, T. D., Pisano, T. J., Deverett, B., Bakshinskaya,

D. E., & Wang, S. S. H. (2018). Normal cognitive and social development require

posterior cerebellar activity. *ELife*. https://doi.org/10.7554/eLife.36401

Bae, J. N., MacFall, J. R., Krishnan, K. R. R., Payne, M. E., Steffens, D. C., & Taylor, W. D. (2006).

Dorsolateral Prefrontal Cortex and Anterior Cingulate Cortex White Matter Alterations

in Late-Life Depression. *Biological Psychiatry*.

https://doi.org/10.1016/j.biopsych.2006.03.052

Barch, D. M., Burgess, G. C., Harms, M. P., Petersen, S. E., Schlaggar, B. L., Corbetta, M., Glasser,

M. F., Curtiss, S., Dixit, S., Feldt, C., Nolan, D., Bryant, E., Hartley, T., Footer, O., Bjork, J.

M., Poldrack, R., Smith, S., Johansen-Berg, H., Snyder, A. Z., & Van Essen, D. C. (2013).

Function in the human connectome: Task-fMRI and individual differences in behavior.

NeuroImage. https://doi.org/10.1016/j.neuroimage.2013.05.033

Baumann, O., & Mattingley, J. B. (2012). Functional topography of primary emotion processing

in the human cerebellum. *NeuroImage*.

https://doi.org/10.1016/j.neuroimage.2012.03.044

Becker, E. B. E., & Stoodley, C. J. (2013). Autism spectrum disorder and the cerebellum. In

International Review of Neurobiology. https://doi.org/10.1016/B978-0-12-418700-

9.00001-0

Beckmann, C. F., Jenkinson, M., & Smith, S. M. (2003). General multilevel linear modeling for

group analysis in FMRI. *NeuroImage*. https://doi.org/10.1016/S1053-8119(03)00435-X

Bench, C. J., Friston, K. J., Brown, R. G., Scott, L. C., Frackowiak, R. S. J., & Dolan, R. J. (1992). The

anatomy of melancholia—Focal abnormalities of cerebral blood flow in major

depression. *Psychological Medicine*. https://doi.org/10.1017/S003329170003806X

Bhanji, J. P., & Delgado, M. R. (2014). The social brain and reward: Social information processing

in the human striatum. In *Wiley Interdisciplinary Reviews: Cognitive Science*.

https://doi.org/10.1002/wcs.1266

Bremner, J. D., Vythilingam, M., Vermetten, E., Nazeer, A., Adil, J., Khan, S., Staib, L. H., &

Charney, D. S. (2002). Reduced volume of orbitofrontal cortex in major depression.

Biological Psychiatry. https://doi.org/10.1016/S0006-3223(01)01336-1

Buckner, R. L., & DiNicola, L. M. (2019). The brain's default network: Updated anatomy,

physiology and evolving insights. In *Nature reviews. Neuroscience*.

https://doi.org/10.1038/s41583-019-0212-7

Buckner, R. L., Krienen, F. M., Castellanos, A., Diaz, J. C., & Thomas Yeo, B. T. (2011). The

organization of the human cerebellum estimated by intrinsic functional connectivity.

Journal of Neurophysiology. https://doi.org/10.1152/jn.00339.2011

Caetano, S. C., Kaur, S., Brambilla, P., Nicoletti, M., Hatch, J. P., Sassi, R. B., Mallinger, A. G.,

Keshavan, M. S., Kupfer, D. J., Frank, E., & Soares, J. C. (2006). Smaller cingulate volumes

in unipolar depressed patients. *Biological Psychiatry*.

https://doi.org/10.1016/j.biopsych.2005.10.011

Cai, Y., Liu, J., Zhang, L., Liao, M., Zhang, Y., Wang, L., Peng, H., He, Z., Li, Z., Li, W., Lu, S., Ding,

Y., & Li, L. (2014). Grey matter volume abnormalities in patients with bipolar I depressive

disorder and unipolar depressive disorder: A voxel-based morphometry study.

Neuroscience Bulletin. https://doi.org/10.1007/s12264-014-1485-5

Carta, I., Chen, C. H., Schott, A. L., Dorizan, S., & Khodakhah, K. (2019). Cerebellar modulation of

the reward circuitry and social behavior. *Science*.

https://doi.org/10.1126/science.aav0581

Casey, B. J., Cannonier, T., Conley, M. I., Cohen, A. O., Barch, D. M., Heitzeg, M. M., Soules, M.

E., Teslovich, T., Dellarco, D. V., Garavan, H., Orr, C. A., Wager, T. D., Banich, M. T.,

Speer, N. K., Sutherland, M. T., Riedel, M. C., Dick, A. S., Bjork, J. M., Thomas, K. M., …

Dale, A. M. (2018). The Adolescent Brain Cognitive Development (ABCD) study: Imaging

acquisition across 21 sites. In *Developmental Cognitive Neuroscience*.

https://doi.org/10.1016/j.dcn.2018.03.001

Chantiluke, K., Halari, R., Simic, M., Pariante, C. M., Papadopoulos, A., Giampietro, V., & Rubia,

K. (2012). Fronto-striato-cerebellar dysregulation in adolescents with depression during

motivated attention. *Biological Psychiatry*.

https://doi.org/10.1016/j.biopsych.2011.09.005

Cheng, W., Rolls, E. T., Qiu, J., Xie, X., Wei, D., Huang, C. C., Yang, A. C., Tsai, S. J., Li, Q., Meng, J.,

Lin, C. P., Xie, P., & Feng, J. (2018). Increased functional connectivity of the posterior

cingulate cortex with the lateral orbitofrontal cortex in depression. *Translational

Psychiatry*. https://doi.org/10.1038/s41398-018-0139-1

Cheng, W., Rolls, E. T., Robbins, T. W., Gong, W., Liu, Z., Lv, W., Du, J., Wen, H., Ma, L., Quinlan,

E. B., Garavan, H., Artiges, E., Papadopoulos Orfanos, D., Smolka, M. N., Schumann, G.,

Kendrick, K., & Feng, J. (2019). Decreased brain connectivity in smoking contrasts with

increased connectivity in drinking. *ELife*. https://doi.org/10.7554/eLife.40765

Coalson, T. S., Essen, D. C. V., & Glasser, M. F. (2018). The impact of traditional neuroimaging methods on the spatial localization of cortical areas. *Proceedings of the National Academy of Sciences*, *115*(27), E6356–E6365. https://doi.org/10.1073/pnas.1801582115

Colman, I., Naicker, K., Zeng, Y., Ataullahjan, A., Senthilselvan, A., & Patten, S. B. (2011). Predictors of long-term prognosis of depression. *CMAJ*. https://doi.org/10.1503/cmaj.110676

Cooper, J. A., Arulpragasam, A. R., & Treadway, M. T. (2018). Anhedonia in depression: Biological mechanisms and computational models. In *Current Opinion in Behavioral Sciences*. https://doi.org/10.1016/j.cobeha.2018.01.024

Covey, L. S., Glassman, A. H., Stetner, F., & Covey, S. (1998). Cigarette Smoking and Major Depression. *Journal of Addictive Diseases*, *17*(1), 35–46. https://doi.org/10.1300/J069v17n01_04

Cruwys, T., Haslam, S. A., Dingle, G. A., Haslam, C., & Jetten, J. (2014). Depression and Social Identity: An Integrative Review. *Personality and Social Psychology Review*. https://doi.org/10.1177/1088868314523839

Cullen, K. R., Klimes-Dougan, B., Muetzel, R., Mueller, B. A., Camchong, J., Houri, A., Kurma, S., & Lim, K. O. (2010). Altered White Matter Microstructure in Adolescents With Major Depression: A Preliminary Study. *Journal of the American Academy of Child and Adolescent Psychiatry*. https://doi.org/10.1097/00004583-201002000-00011

Cusi, A. M., Nazarov, A., Holshausen, K., MacQueen, G. M., & McKinnon, M. C. (2012). Systematic review of the neural basis of social cognition in patients with mood

disorders. In *Journal of Psychiatry and Neuroscience*.

https://doi.org/10.1503/jpn.100179

Derntl, B., Seidel, E. M., Eickhoff, S. B., Kellermann, T., Gur, R. C., Schneider, F., & Habel, U.

(2011). Neural correlates of social approach and withdrawal in patients with major

depression. *Social Neuroscience*. https://doi.org/10.1080/17470919.2011.579800

Diehl, M. M., Lempert, K. M., Parr, A. C., Ballard, I., Steele, V. R., & Smith, D. V. (2018). Toward

an integrative perspective on the neural mechanisms underlying persistent maladaptive

behaviors. In *European Journal of Neuroscience*. https://doi.org/10.1111/ejn.14083

Dolan, R. J., Bench, C. J., Scott, L. C., Friston, K. J., Frackowiak, R. S. J., Brown, R. G., Dolan, R. J.,

& Frackowiak, R. S. J. (1992). Regional cerebral blood flow abnormalities in depressed

patients with cognitive impairment. *Journal of Neurology, Neurosurgery and Psychiatry*,

55(9), 768–773. https://doi.org/10.1136/jnnp.55.9.768

Edwards, A. C., Maes, H. H., Pedersen, N. L., & Kendler, K. S. (2011). A population-based twin

study of the genetic and environmental relationship of major depression, regular

tobacco use and nicotine dependence. *Psychological Medicine*.

https://doi.org/10.1017/S0033291710000589

Ely, B. A., Xu, J., Goodman, W. K., Lapidus, K. A., Gabbay, V., & Stern, E. R. (2016). Resting-state

functional connectivity of the human habenula in healthy individuals: Associations with

subclinical depression. *Human Brain Mapping*. https://doi.org/10.1002/hbm.23179

Fareri, D. S., & Delgado, M. R. (2014). Social rewards and social networks in the human brain. In

Neuroscientist. https://doi.org/10.1177/1073858414521869

Feinberg, D. A., Moeller, S., Smith, S. M., Auerbach, E., Ramanna, S., Glasser, M. F., Miller, K. L., Ugurbil, K., & Yacoub, E. (2010). Multiplexed echo planar imaging for sub-second whole brain fmri and fast diffusion imaging. *PLoS ONE*. https://doi.org/10.1371/journal.pone.0015710

Fergusson, D. M., Boden, J. M., & Horwood, L. J. (2009). Tests of causal links between alcohol abuse or dependence and major depression. *Archives of General Psychiatry*, *66*(3), 260–266. https://doi.org/10.1001/archgenpsychiatry.2008.543

Ferrari, A. J., Charlson, F. J., Norman, R. E., Patten, S. B., Freedman, G., Murray, C. J. L., Vos, T., & Whiteford, H. A. (2013). Burden of Depressive Disorders by Country, Sex, Age, and Year: Findings from the Global Burden of Disease Study 2010. *PLoS Medicine*. https://doi.org/10.1371/journal.pmed.1001547

Fried, E. I., & Nesse, R. M. (2014). The impact of individual depressive symptoms on impairment of psychosocial functioning. *PLoS ONE*. https://doi.org/10.1371/journal.pone.0090311

Friston, K. J., Buechel, C., Fink, G. R., Morris, J., Rolls, E., & Dolan, R. J. (1997). Psychophysiological and modulatory interactions in neuroimaging. *NeuroImage*. https://doi.org/10.1006/nimg.1997.0291

Furman, D. J., Hamilton, J. P., & Gotlib, I. H. (2011). Frontostriatal functional connectivity in major depressive disorder. *Biology of Mood & Anxiety Disorders*. https://doi.org/10.1186/2045-5380-1-11

George, L. K., Blazer, D. G., Hughes, D. C., & Fowler, N. (1989). Social support and the outcome of major depression. *British Journal of Psychiatry*. https://doi.org/10.1192/bjp.154.4.478

Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E., Ugurbil, K.,

Andersson, J., Beckmann, C. F., Jenkinson, M., Smith, S. M., & Van Essen, D. C. (2016). A

multi-modal parcellation of human cerebral cortex. *Nature*, *536*(7615), 171–178.

https://doi.org/10.1038/nature18933

Glasser, M. F., Smith, S. M., Marcus, D. S., Andersson, J. L. R., Auerbach, E. J., Behrens, T. E. J.,

Coalson, T. S., Harms, M. P., Jenkinson, M., Moeller, S., Robinson, E. C., Sotiropoulos, S.

N., Xu, J., Yacoub, E., Ugurbil, K., & Van Essen, D. C. (2016). The Human Connectome

Project's neuroimaging approach. *Nature Neuroscience*, *19*(9), 1175–1187.

https://doi.org/10.1038/nn.4361

Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L., Xu, J.,

Jbabdi, S., Webster, M., Polimeni, J. R., Van Essen, D. C., & Jenkinson, M. (2013). The

minimal preprocessing pipelines for the Human Connectome Project. *NeuroImage*.

https://doi.org/10.1016/j.neuroimage.2013.04.127

Harkness, K. L., Washburn, D., Theriault, J. E., Lee, L., & Sabbagh, M. A. (2011). Maternal history

of depression is associated with enhanced theory of mind in depressed and

nondepressed adult women. *Psychiatry Research*.

https://doi.org/10.1016/j.psychres.2011.06.007

Hershenberg, R., Satterthwaite, T. D., Daldal, A., Katchmar, N., Moore, T. M., Kable, J. W., &

Wolf, D. H. (2016). Diminished effort on a progressive ratio task in both unipolar and

bipolar depression. *Journal of Affective Disorders*.

https://doi.org/10.1016/j.jad.2016.02.003

Hoertel, N., Blanco, C., Oquendo, M. A., Wall, M. M., Olfson, M., Falissard, B., Franco, S., Peyre, H., Lemogne, C., & Limosin, F. (2017). A comprehensive model of predictors of persistence and recurrence in adults with major depression: Results from a national 3-year prospective study. *Journal of Psychiatric Research*. https://doi.org/10.1016/j.jpsychires.2017.07.022

Iacoviello, B. M., Alloy, L. B., Abramson, L. Y., & Choi, J. Y. (2010). The early course of depression: A longitudinal investigation of prodromal symptoms and their relation to the symptomatic course of depressive episodes. *Journal of Abnormal Psychology*. https://doi.org/10.1037/a0020114

Inoue, Y., Tonooka, Y., Yamada, K., & Kanba, S. (2004). Deficiency of theory of mind in patients with remitted mood disorder. *Journal of Affective Disorders*. https://doi.org/10.1016/j.jad.2004.04.004

Inoue, Y., Yamada, K., & Kanba, S. (2006). Deficit in theory of mind is a risk for relapse of major depression. *Journal of Affective Disorders*. https://doi.org/10.1016/j.jad.2006.04.018

Jarcho, J. M., Davis, M. M., Shechner, T., Degnan, K. A., Henderson, H. A., Stoddard, J., Fox, N. A., Leibenluft, E., Pine, D. S., & Nelson, E. E. (2016). Early-Childhood Social Reticence Predicts Brain Function in Preadolescent Youths During Distinct Forms of Peer Evaluation. *Psychological Science*. https://doi.org/10.1177/0956797616638319

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved Optimization for the Robust and Accurate Linear Registration and Motion Correction of Brain Images. *NeuroImage*, *17*(2), 825–841. https://doi.org/10.1006/nimg.2002.1132

Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W., & Smith, S. M. (2012). FSL -

    Review. *NeuroImage*. https://doi.org/10.1016/j.neuroimage.2011.09.015

Klein, D. N., Glenn, C. R., Kosty, D. B., Seeley, J. R., Rohde, P., & Lewinsohn, P. M. (2013).

    Predictors of first lifetime onset of major depressive disorder in young adulthood.

    Journal of Abnormal Psychology. https://doi.org/10.1037/a0029567

Klein, D. N., Lewinsohn, P. M., Rohde, P., Seeley, J. R., & Olino, T. M. (2005). Psychopathology in

    the adolescent and young adult offspring of a community sample of mothers and fathers

    with major depression. *Psychological Medicine*.

    https://doi.org/10.1017/S0033291704003587

Klonsky, E. D., May, A. M., & Saffer, B. Y. (2016). Suicide, Suicide Attempts, and Suicidal

    Ideation. *Annual Review of Clinical Psychology*. https://doi.org/10.1146/annurev-

    clinpsy-021815-093204

Knight, M. J., Air, T., & Baune, B. T. (2018). The role of cognitive impairment in psychosocial

    functioning in remitted depression. *Journal of Affective Disorders*.

    https://doi.org/10.1016/j.jad.2018.04.051

Koenigs, M., Huey, E. D., Calamia, M., Raymont, V., Tranel, D., & Grafman, J. (2008). Distinct

    regions of prefrontal cortex mediate resistance and vulnerability to depression. *Journal

    of Neuroscience*. https://doi.org/10.1523/JNEUROSCI.2324-08.2008

Konarski, J. Z., McIntyre, R. S., Grupp, L. A., & Kennedy, S. H. (2005). Is the cerebellum relevant

    in the circuitry of neuropsychiatric disorders? In *Journal of Psychiatry and Neuroscience*.

Kovacs, M., Obrosky, S., & George, C. (2016). The course of major depressive disorder from

childhood to young adulthood: Recovery and recurrence in a longitudinal observational

study. *Journal of Affective Disorders*. https://doi.org/10.1016/j.jad.2016.05.042

Kumar, P., Goer, F., Murray, L., Dillon, D. G., Beltzer, M. L., Cohen, A. L., Brooks, N. H., &

Pizzagalli, D. A. (2018). Impaired reward prediction error encoding and striatal-midbrain

connectivity in depression. *Neuropsychopharmacology*.

https://doi.org/10.1038/s41386-018-0032-x

Lacerda, A. L. T., Keshavan, M. S., Hardan, A. Y., Yorbik, O., Brambilla, P., Sassi, R. B., Nicoletti,

M., Mallinger, A. G., Frank, E., Kupfer, D. J., & Soares, J. C. (2004). Anatomic evaluation

of the orbitofrontal cortex in major depressive disorder. *Biological Psychiatry*.

https://doi.org/10.1016/j.biopsych.2003.08.021

Liu, L., Zeng, L. L., Li, Y., Ma, Q., Li, B., Shen, H., & Hu, D. (2012). Altered cerebellar functional

connectivity with intrinsic connectivity networks in adults with major depressive

disorder. *PLoS ONE*. https://doi.org/10.1371/journal.pone.0039516

Luking, K. R., Pagliaccio, D., Luby, J. L., & Barch, D. M. (2016). Depression Risk Predicts Blunted

Neural Responses to Gains and Enhanced Responses to Losses in Healthy Children.

Journal of the American Academy of Child and Adolescent Psychiatry.

https://doi.org/10.1016/j.jaac.2016.01.007

Ma, Q., Zeng, L. L., Shen, H., Liu, L., & Hu, D. (2013). Altered cerebellar-cerebral resting-state

functional connectivity reliably identifies major depressive disorder. *Brain Research*.

https://doi.org/10.1016/j.brainres.2012.12.002

Manninen, L., Poikolainen, K., Vartianen, E., & Laatikainen, T. (2006). Heavy drinking occasions

and depressions. *Alcohol and Alcoholism*. https://doi.org/10.1093/alcalc/agh246

McCabe, C., Woffindale, C., Harmer, C. J., & Cowen, P. J. (2012). Neural processing of reward

and punishment in young people at increased familial risk of depression. *Biological*

Psychiatry. https://doi.org/10.1016/j.biopsych.2012.04.034

Moeller, S., Yacoub, E., Olman, C. A., Auerbach, E., Strupp, J., Harel, N., & Uğurbil, K. (2010).

Multiband multislice GE-EPI at 7 tesla, with 16-fold acceleration using partial parallel

imaging with application to high spatial and temporal whole-brain FMRI. *Magnetic*

Resonance in Medicine. https://doi.org/10.1002/mrm.22361

Monk, C. S., Klein, R. G., Telzer, E. H., Schroth, E. A., Mannuzza, S., Moulton, J. L., Guardino, M.,

Masten, C. L., McClure-Tone, E. B., Fromm, S., Blair, R. J., Pine, D. S., & Ernst, M. (2008).

Amygdala and nucleus accumbens activation to emotional facial expressions in children

and adolescents at risk for major depression. *American Journal of Psychiatry*.

https://doi.org/10.1176/appi.ajp.2007.06111917

Monroe, S. M., Slavich, G. M., & Gotlib, I. H. (2014). Life stress and family history for depression:

The moderating role ofpast depressive episodes. *Journal of Psychiatric Research*.

https://doi.org/10.1016/j.jpsychires.2013.11.005

Myers-Schulz, B., & Koenigs, M. (2012). Functional anatomy of ventromedial prefrontal cortex:

Implications for mood and anxiety disorders. In *Molecular Psychiatry*.

https://doi.org/10.1038/mp.2011.88

Ng, T. H., Alloy, L. B., & Smith, D. V. (2019). Meta-analysis of reward processing in major

depressive disorder reveals distinct abnormalities within the reward circuit.

Translational Psychiatry. https://doi.org/10.1038/s41398-019-0644-x

Nock, M. K., Hwang, I., Sampson, N., Kessler, R. C., Angermeyer, M., Beautrais, A., Borges, G.,

Bromet, E., Bruffaerts, R., De Girolamo, G., De Graaf, R., Florescu, S., Gureje, O., Haro, J.

M., Hu, C., Huang, Y., Karam, E. G., Kawakami, N., Kovess, V., … Williams, D. R. (2009).

Cross-national analysis of the associations among mental disorders and suicidal

behavior: Findings from the WHO World Mental Health Surveys. *PLoS Medicine*.

https://doi.org/10.1371/journal.pmed.1000123

Olino, T. M., Silk, J. S., Osterritter, C., & Forbes, E. E. (2015). Social reward in youth at risk for

depression: A preliminary investigation of subjective and neural differences. *Journal of

Child and Adolescent Psychopharmacology*. https://doi.org/10.1089/cap.2014.0165

Pedrelli, P., Borsari, B., Ketchen Lipson, S., Heinze, J. E., & Eisenberg, D. (2016). Gender

Differences in the Relationships Among Major Depressive Disorder, Heavy Alcohol Use,

and Mental Health Treatment Engagement Among College Students A PPROXIMATELY

40% OF COLLEGE STUDENTS report engaging in heavy episodic drinking. In *JOURNAL OF

STUDIES ON ALCOHOL AND DRUGS*.

Pizzagalli, D. A. (2014). Depression, Stress, and Anhedonia: Toward a Synthesis and Integrated

Model. *Annual Review of Clinical Psychology*. https://doi.org/10.1146/annurev-clinpsy-

050212-185606

Robinson, O. J., Cools, R., Carlisi, C. O., Sahakian, B. J., & Drevets, W. C. (2012). Ventral striatum

response during reward and punishment reversal learning in unmedicated major

depressive disorder. *American Journal of Psychiatry*.

https://doi.org/10.1176/appi.ajp.2011.11010137

Romer, A. L., Knodt, A. R., Houts, R., Brigidi, B. D., Moffitt, T. E., Caspi, A., & Hariri, A. R. (2017).

Structural alterations within cerebellar circuitry are associated with general liability for

common mental disorders. *Molecular Psychiatry*. https://doi.org/10.1038/mp.2017.57

Salvadore, G., Nugent, A. C., Lemaitre, H., Luckenbaugh, D. A., Tinsley, R., Cannon, D. M.,

Neumeister, A., Zarate, C. A., & Drevets, W. C. (2011). Prefrontal cortical abnormalities

in currently depressed versus currently remitted patients with major depressive

disorder. *NeuroImage*. https://doi.org/10.1016/j.neuroimage.2010.11.011

Santini, Z. I., Koyanagi, A., Tyrovolas, S., Mason, C., & Haro, J. M. (2015). The association

between social relationships and depression: A systematic review. In *Journal of Affective

Disorders*. https://doi.org/10.1016/j.jad.2014.12.049

Savitz, J. B., Rauch, S. L., & Drevets, W. C. (2013). Clinical application of brain imaging for the

diagnosis of mood disorders: The current state of play. In *Molecular Psychiatry*.

https://doi.org/10.1038/mp.2013.25

Schmahmann, J. D. (2004). Disorders of the cerebellum: Ataxia, dysmetria of thought, and the

cerebellar cognitive affective syndrome. In *Journal of Neuropsychiatry and Clinical

Neurosciences*. https://doi.org/10.1176/jnp.16.3.367

Schmahmann, J. D., Weilburg, J. B., & Sherman, J. C. (2007). The neuropsychiatry of the

cerebellum—Insights from the clinic. In *Cerebellum*.

https://doi.org/10.1080/14734220701490995

Serra-Blasco, M., Portella, M. J., Gómez-Ansón, B., De Diego-Adelinõ, J., Vives-Gilabert, Y.,

Puigdemont, D., Granell, E., Santos, A., Álvarez, E., & Pérez, V. (2013). Effects of illness

duration and treatment resistance on grey matter abnormalities in major depression.

British Journal of Psychiatry. https://doi.org/10.1192/bjp.bp.112.116228

Setsompop, K., Gagoski, B. A., Polimeni, J. R., Witzel, T., Wedeen, V. J., & Wald, L. L. (2012).

Blipped-controlled aliasing in parallel imaging for simultaneous multislice echo planar

imaging with reduced g-factor penalty. *Magnetic Resonance in Medicine*.

https://doi.org/10.1002/mrm.23097

Sherdell, L., Waugh, C. E., & Gotlib, I. H. (2012). Anticipatory pleasure predicts motivation for

reward in major depression. *Journal of Abnormal Psychology*.

https://doi.org/10.1037/a0024945

Smith, S. M., Beckmann, C. F., Andersson, J., Auerbach, E. J., Bijsterbosch, J., Douaud, G., Duff,

E., Feinberg, D. A., Griffanti, L., Harms, M. P., Kelly, M., Laumann, T., Miller, K. L.,

Moeller, S., Petersen, S., Power, J., Salimi-Khorshidi, G., Snyder, A. Z., Vu, A. T., …

Glasser, M. F. (2013). Resting-state fMRI in the Human Connectome Project.

NeuroImage. https://doi.org/10.1016/j.neuroimage.2013.05.039

Smith, D. V., & Delgado, M. R. (2015). Reward Processing. In *Brain Mapping: An Encyclopedic

Reference*. https://doi.org/10.1016/B978-0-12-397025-1.00255-4

Smith, David V., Utevsky, A. V., Bland, A. R., Clement, N., Clithero, J. A., Harsch, A. E. W., McKell

Carter, R., & Huettel, S. A. (2014). Characterizing individual differences in functional

connectivity using dual-regression and seed-based approaches. *NeuroImage*.

https://doi.org/10.1016/j.neuroimage.2014.03.042

Solomon, D. A., Keller, M. B., Leon, A. C., Mueller, T. I., Lavori, P. W., Shea, M. T., Coryell, W., Warshaw, M., Turvey, C., Maser, J. D., & Endicott, J. (2000). Multiple recurrences of major depressive disorder. *American Journal of Psychiatry*. https://doi.org/10.1176/appi.ajp.157.2.229

Steingard, R. J., Renshaw, P. F., Hennen, J., Lenox, M., Cintron, C. B., Young, A. D., Connor, D. F., Au, T. H., & Yurgelun-Todd, D. A. (2002). Smaller frontal lobe white matter volumes in depressed adolescents. *Biological Psychiatry*. https://doi.org/10.1016/S0006-3223(02)01393-8

Sullivan, L. E., Fiellin, D. A., & O'Connor, P. G. (2005). The prevalence and impact of alcohol problems in major depression: A systematic review. In *American Journal of Medicine* (Vol. 118, Issue 4, pp. 330–341). Elsevier Inc. https://doi.org/10.1016/j.amjmed.2005.01.007

Sullivan, P. F., Neale, M. C., & Kendler, K. S. (2000). Genetic epidemiology of major depression: Review and meta-analysis. In *American Journal of Psychiatry*. https://doi.org/10.1176/appi.ajp.157.10.1552

Taylor, W. D., MacFall, J. R., Payne, M. E., McQuoid, D. R., Provenzale, J. M., Steffens, D. C., & Krishnan, R. R. (2004). Late-life depression and microstructural abnormalities in dorsolateral prefrontal cortex white matter. *American Journal of Psychiatry*. https://doi.org/10.1176/appi.ajp.161.7.1293

Treadway, M. T., Bossaller, N. A., Shelton, R. C., & Zald, D. H. (2012). Effort-based decision-making in major depressive disorder: A translational model of motivational anhedonia. *Journal of Abnormal Psychology*. https://doi.org/10.1037/a0028813

Uğurbil, K., Xu, J., Auerbach, E. J., Moeller, S., Vu, A. T., Duarte-Carvajalino, J. M., Lenglet, C.,

    Wu, X., Schmitter, S., Van de Moortele, P. F., Strupp, J., Sapiro, G., De Martino, F., Wang,

    D., Harel, N., Garwood, M., Chen, L., Feinberg, D. A., Smith, S. M., … Yacoub, E. (2013).

    Pushing spatial and temporal resolution for functional and diffusion MRI in the Human

    Connectome Project. *NeuroImage*. https://doi.org/10.1016/j.neuroimage.2013.05.012

Van Essen, D. C., Smith, S. M., Barch, D. M., Behrens, T. E. J., Yacoub, E., & Ugurbil, K. (2013).

    The WU-Minn Human Connectome Project: An overview. *NeuroImage*.

    https://doi.org/10.1016/j.neuroimage.2013.05.041

Van Overwalle, F., Baetens, K., Mariën, P., & Vandekerckhove, M. (2014). Social cognition and

    the cerebellum: A meta-analysis of over 350 fMRI studies. In *NeuroImage*.

    https://doi.org/10.1016/j.neuroimage.2013.09.033

Van Overwalle, F., D'aes, T., & Mariën, P. (2015). Social cognition and the cerebellum: A meta-

    analytic connectivity analysis. *Human Brain Mapping*.

    https://doi.org/10.1002/hbm.23002

Van Overwalle, F., De Coninck, S., Heleven, E., Perrotta, G., Taib, N. O. Ben, Manto, M., &

    Mariën, P. (2019). The role of the cerebellum in reconstructing social action sequences:

    A pilot study. *Social Cognitive and Affective Neuroscience*.

    https://doi.org/10.1093/scan/nsz032

Wang, K. S., Smith, D. V., & Delgado, M. R. (2016). Using fMRI to study reward processing in

    humans: Past, present, and future. In *Journal of Neurophysiology*.

    https://doi.org/10.1152/jn.00333.2015

Wang, S. S. H., Kloth, A. D., & Badura, A. (2014). The Cerebellum, Sensitive Periods, and Autism. In *Neuron*. https://doi.org/10.1016/j.neuron.2014.07.016

Weinberg, A., Liu, H., Hajcak, G., & Shankman, S. A. (2015). Blunted neural response to rewards as a vulnerability factor for depression: Results from a family study. *Journal of Abnormal Psychology*. https://doi.org/10.1037/abn0000081

Weinberger, A. H., Kashan, R. S., Shpigel, D. M., Esan, H., Taha, F., Lee, C. J., Funk, A. P., & Goodwin, R. D. (2017). Depression and cigarette smoking behavior: A critical review of population-based studies. *The American Journal of Drug and Alcohol Abuse*, *43*(4), 416–431. https://doi.org/10.3109/00952990.2016.1171327

Winkler, A. M., Ridgway, G. R., Webster, M. A., Smith, S. M., & Nichols, T. E. (2014). Permutation inference for the general linear model. *NeuroImage*. https://doi.org/10.1016/j.neuroimage.2014.01.060

Winkler, A. M., Webster, M. A., Vidaurre, D., Nichols, T. E., & Smith, S. M. (2015). Multi-level block permutation. *NeuroImage*. https://doi.org/10.1016/j.neuroimage.2015.05.092

Wise, T., Radua, J., Via, E., Cardoner, N., Abe, O., Adams, T. M., Amico, F., Cheng, Y., Cole, J. H., Périco, C. de A. M., Dickstein, D. P., Farrow, T. F. D., Frodl, T., Wagner, G., Gotlib, I. H., Gruber, O., Ham, B. J., Job, D. E., Kempton, M. J., … Arnone, D. (2017). Common and distinct patterns of grey-matter volume alteration in major depression and bipolar disorder: Evidence from voxel-based meta-analysis. *Molecular Psychiatry*. https://doi.org/10.1038/mp.2016.72

Woolrich, M. (2008). Robust group analysis using outlier inference. *NeuroImage*. https://doi.org/10.1016/j.neuroimage.2008.02.042

Woolrich, M. W., Behrens, T. E. J., Beckmann, C. F., Jenkinson, M., & Smith, S. M. (2004).

Multilevel linear modelling for FMRI group analysis using Bayesian inference.

NeuroImage. https://doi.org/10.1016/j.neuroimage.2003.12.023

Woolrich, M. W., Ripley, B. D., Brady, M., & Smith, S. M. (2001). Temporal autocorrelation in

univariate linear modeling of FMRI data. *NeuroImage*.

https://doi.org/10.1006/nimg.2001.0931

World Health Organization. (2017). Depression and other common mental disorders: Global

health estimates. *World Health Organization*. https://doi.org/CC BY-NC-SA 3.0 IGO

Worsley, K. J. (2001). Testing for signals with unknown location and scale in a χ2 random field,

with an application to fMRI. *Advances in Applied Probability*.

https://doi.org/10.1239/aap/1011994029

Xu, J., Moeller, S., Auerbach, E. J., Strupp, J., Smith, S. M., Feinberg, D. A., Yacoub, E., & Uğurbil,

K. (2013). Evaluation of slice accelerations using multiband echo planar imaging at 3T.

NeuroImage. https://doi.org/10.1016/j.neuroimage.2013.07.055

Yang, Q., Huang, X., Hong, N., & Yu, X. (2007). White matter microstructural abnormalities in

late-life depression. *International Psychogeriatrics*.

https://doi.org/10.1017/S1041610207004875

Young, J. S., Smith, D. V., Coutlee, C. G., & Huettel, S. A. (2015). Synchrony between sensory and

cognitive networks is associated with subclinical variation in autistic traits. *Frontiers in

Human Neuroscience*. https://doi.org/10.3389/fnhum.2015.00146

Yucel, K., Nazarov, A., Taylor, V. H., Macdonald, K., Hall, G. B., & MacQueen, G. M. (2013). Cerebellar vermis volume in major depressive disorder. *Brain Structure and Function*, *218*(4), 851–858. https://doi.org/10.1007/s00429-012-0433-2

Zhang, J., Kendrick, K. M., Lu, G., & Feng, J. (2015). The fault lies on the other side: Altered brain functional connectivity in psychiatric disorders is mainly caused by counterpart regions in the opposite hemisphere. *Cerebral Cortex*. https://doi.org/10.1093/cercor/bhu173

Zhao, Y. J., Du, M. Y., Huang, X. Q., Lui, S., Chen, Z. Q., Liu, J., Luo, Y., Wang, X. L., Kemp, G. J., & Gong, Q. Y. (2014). Brain grey matter abnormalities in medication-free patients with major depressive disorder: A meta-analysis. *Psychological Medicine*. https://doi.org/10.1017/S0033291714000518

Zimmermann, P., Brückl, T., Lieb, R., Nocon, A., Ising, M., Beesdo, K., & Wittchen, H. U. (2008). The Interplay of Familial Depression Liability and Adverse Events in Predicting the First Onset of Depression During a 10-Year Follow-up. *Biological Psychiatry*. https://doi.org/10.1016/j.biopsych.2007.05.020