Bacterial lipopolysaccharide induces settlement and metamorphosis in a marine larva.

Marnie L. Freckelton[1], Brian T. Nedved[1], You-Sheng Cai[2,3], Shugeng Cao[2], Helen Turano[4], Rosanna A. Alegado[4,5] and Michael G. Hadfield[1*]

[1] Kewalo Marine Laboratory, University of Hawai□i, Honolulu, Hawai□i, United States, 96813.
[2] Department of Pharmaceutical Sciences, Daniel K Inouye College of Pharmacy, University of Hawai□i at Hilo, 200 W. Kawili Street, Hilo, Hawai□i 96720
[3] Institute of TCM and Natural Products, School of Pharmaceutical Sciences, Wuhan University, 185 Donghu Road, Wuhan, Hubei 430071, People's Republic of China
[4] Department of Oceanography, University of Hawai□i Mānoa, Honolulu, Hawai□i, United States, 96813.
[5] Sea Grant College Program, University of Hawai□i Mānoa, Honolulu Hawai□i, United States, 96813.

*Correspondence: Michael G. Hadfield

Email: hadfield@hawaii.edu

Author Contributions: Conceptualization, M.L.F, M.G.H., S.C., R.A.A., B.T.N.; Methodology, M.L.F., M.G.H., B.T.N., H.T., R.A.A.; Investigation, M.L.F, B.T.N., Y.C., H.T.; Resources, M.G.H., S.C., R.A.A.; Writing – Original Draft, M.L.F. and M.G.H with support from all the co-authors; Supervision and Funding Acquisition, M.G.H., B.T.N., S.C., R.A.A.

Competing Interest Statement: The authors declare no competing interests.

Classification

Major: Biological Sciences
Minor: Ecology

Keywords*: Hydroides elegans*, metamorphosis, lipopolysaccharide, *Cellulophaga lytica*, biofilms

This PDF file includes:

Main Text

Figures 1 to 5

Table 1

Abstract

How larvae of the many phyla of marine invertebrates find places appropriate for settlement,

metamorphosis, growth and reproduction is an enduring question in marine science. Biofilm

induced metamorphosis has been observed in marine invertebrate larvae from nearly every major

marine phylum. Despite the widespread nature of this phenomenon the mechanism of induction

remains poorly understood. The serpulid polychaete *Hydroides elegans* is a well-established

model for investigating bacteria-induced larval development. A broad range of biofilm bacterial

species elicit larval metamorphosis in *H. elegans* via at least two mechanisms, including outer

membrane vesicles and phage-tail bacteriocins. We investigated the interaction between larvae

of *H. elegans* and the inductive bacterium *Cellulophaga lytica*, which produces an abundance of

OMVs but not phage-tail bacteriocins. We asked whether the OMVs of *C. lytica* induce larval

settlement due to cell membrane components or through delivery of specific cargo. Employing a

biochemical structure-function approach with a strong ecological focus, the cells and outer

membrane vesicles produced by *C. lytica* were interrogated to determine the nature of the

inductive molecule. Here we report that the cue produced by *C. lytica* that induces larvae of

H. elegans to metamorphose is lipopolysaccharide (LPS). The widespread prevalence of LPS

and its associated taxonomic and structural variability suggest it may be a broadly employed cue

for bacterially induced larval settlement of marine invertebrates.

Significance Statement

New surfaces in the sea are quickly populated by dense communities of invertebrate animals,

whose establishment and maintenance require site-specific settlement of larvae from the

plankton. Although it is recognized that larvae selectively settle in sites where they can

metamorphose and thrive, and that the biofilm bacteria residing on these surfaces supply

inductive cues, the nature of the cues used to identify 'right places' has remained enigmatic. In

this paper, we reveal that lipopolysaccharide (LPS) from the outer membrane of a marine Gram-

negative bacterium cue metamorphosis for a marine worm and discuss the likelihood that LPS

provides the variation necessary to explain settlement site selectivity for many of the bottom-living

invertebrate animals that metamorphose in response to bacterial biofilms.

Main Text

Introduction

More than two-thirds of the earth's surface is covered by the oceans. The solid surface beneath
these salt waters is greater than that of all the jungles, rain forests, boreal forests, prairies,
desserts, tundras, and mountains combined. Yet, we know more about the dynamics of the
biological world living above the ocean's shores than we do about the enormously diverse benthic
communities that extend from the higher reaches of the intertidal to the greatest depths of the
seas. With global climate change causing massive melting of the earth's glaciers, sea levels are
rising, and the amount of land covered by the seas is increasing dramatically. The establishment
and maintenance of benthic marine animal communities depends heavily on recruitment of
minute larvae from the plankton. Larval recruitment is an essential process not only for all marine
benthic communities, but also for creating harvestable populations of mariculture species such as
clams, oysters, and shrimp. It is also the initial process in the establishment of costly biofouling
communities on the hulls of ships and in the pipes that bring cooling seawater to coastal electrical
generators (1). A critical element of this process that remains poorly understood is how larvae
select habitats suitable not just for attachment and metamorphosis, but long-term survival, growth
and reproduction (2, 3).

In recent decades we have learned that the cues most invertebrate larvae employ to select right
places to metamorphose are almost universally biological, the majority originating from microbes
residing in surface biofilms (4, 5). Biofilm- or bacterial-induced settlement has been shown for
larvae of sponges (6–8), cnidarians (9, 10), bryozoans (11, 12), molluscs (13–16), annelids (17),
echinoderms (18, 19), crustaceans (20, 21) and urochordates (22), i.e., all of the major
invertebrate phyla. Although the number of larval types recorded to settle and metamorphose in
response to bacterial films is now so great as to suggest a nearly universal mechanism for both
settlement induction and larval response, we remain largely ignorant about the diversity of
bacteria that stimulate larvae to settle, the chemical nature of the bacterial cues, and the
mechanisms involved in larval recognition of bacteria.

Larvae of the serpulid polychaete *Hydroides elegans* provide an excellent model for investigating
bacteria-stimulated settlement (17, 23). *H. elegans* is a cosmopolitan member of the benthic and
biofouling communities in warm-water bays and ports throughout the world (23). As is the case
for a vast majority of marine invertebrates, an external stimulant induces an extremely rapid
series of morphogenetic events that quickly transform the swimming larva of *H. elegans* into a

bottom-living, tube-dwelling juvenile that will grow rapidly into a reproductive adult (23). Larvae of

H. elegans typically do not settle in the absence of a biofilm, and this response follows direct

contact with surface biofilms (17, 23–28). Investigations of the complex biofilm communities that

induce settlement and metamorphosis in *H. elegans* and the cues that they produce must take

place in the context of this ecology. Inductive cues must be naturally entrained within the biofilm

and fast acting, because even in a low turnover environment such as Pearl Harbor, Hawai□i,

current speeds and turbulence at the biofilm interface would otherwise rapidly disperse and

remove dissolved compounds (29–31).

Evidence has accumulated that it is particular bacterial species residing in the biofilms that induce

the metamorphic response in the larvae of *H. elegans* (5, 28, 32). Inductive bacteria from a broad

range of phyla, including both Gram-negative and Gram-positive strains, are inductive when

cultured in mono-species biofilms (24, 28, 32–35). However, most biofilm-bacterial species do not

induce settlement in *H. elegans* (28, 36). To date, we have identified at least two different

induction mechanisms from two different Gram-negative bacteria, *Pseudoalteromonas*

luteoviolacea and *Cellulophaga lytica.* For *P. luteoviolacea* strain HI1, a highly inductive biofilm

species isolated from Pearl Harbor, Hawai□i and genetically characterized in our lab, we

identified specific structural elements derived from phage-tail proteins that are in some manner

involved in settlement induction (37–39). However, examination of other inductive bacterial

species revealed that induction of larval settlement cannot alone be explained by phage-tail

bacteriocins. *Cellulophaga lytica,* another biofilm bacterium, induces larvae of *H. elegans* to settle

by an entirely different mechanism. Although cell-free preparations from broth cultures of *C. lytica*

induce the tubeworm larvae to metamorphose, analysis of the genome of *C. lytica* (40) yielded

none of the genes that transcribe the phage-tail elements of bacteriocins (34). Additionally,

transmission electron microscopy (TEM) of inductive supernatants from *C. lytica* revealed an

abundance of outer membrane vesicles (OMVs) and none of the bacteriocins found in similar

preparations from *P. luteoviolacea* (34).

We have previously suggested that such membrane vesicles should be considered for their

potential to be a common mechanism of interaction between biofilm bacteria and invertebrate

larvae (35). Gram-negative bacteria ubiquitously produce OMVs (41, 42), but their structures and

contents are variable per species and strain. Interactions between microbes and eukaryotes are

mediated largely by a suite of conserved molecules knowns as Microbe-Associated Molecular

Patterns (MAMPs) (43, 44). Two major MAMP classes, lipopolysaccharide and peptidoglycan,

comprise important components of the envelope of both whole cells and OMVs. Additionally,

OMVs may enclose genetic material, virulence factors, proteins, small bioactive metabolites and

lipids (41). Consequently, OMVs provide the mechanism for multiple ecological roles fulfilled by

Gram-negative bacteria, including cell-to-cell signaling and transfer of DNA, proteins and small

signaling molecules between cells (41, 42, 45–53). The ubiquity of OMVs means that the

response to their detection by a eukaryote is entirely context dependent. In the present study, we

asked whether the OMVs of *C. lytica* induce larval settlement because they are accessible pieces

of bacterial cell membranes or because they deliver a specific cargo.

To gain key insight into the molecular structure of the larval settlement inducer produced by

C. lytica and to determine if it is a protein, nucleic acid, or lipid, we subjected the OMV fractions to

a battery of broad-spectrum enzymatic and chemical treatments and then tested the treated

OMVs for loss of larval settlement-inducing activity. Subsequently, because OMVs are composed

of pieces of the bacterial outer membrane and because of the need for larger quantities of

compounds for treatments and assays, the components of the cell envelope (whole membrane,

peptidoglycan and lipopolysaccharide) were assayed directly by extraction from whole cells.

These approaches, including *in silico* analyses of the bacterial genome, HPLC fractionation of

methanol extracts, and enzymatic destructions of OMVs and isolated cell membrane

components, combined with larval settlement assays, allowed us to reject all but bacterial LPS as

cues for larval settlement. As the outermost element of Gram-negative bacterial cells, membrane

LPS provides the surface-contact stimulus indicated by behavioral studies of larval settlement in

H. elegans (26). We conclude LPS is a key element in selective larval settlement by *H. elegans*,

and we predict this will hold true for many other benthic marine invertebrate animals.

Results

<u>OMV production, size, and metamorphic bioactivity as a function of growth phase in *C. lytica.*</u>

OMVs isolated from broth cultures of *C. lytica* induced metamorphosis in larvae of *H. elegans* at

all growth phases (mid-log, late-log and stationary phase) with the highest average response to

OMVs isolated during stationary phase and lowest from mid-log phase OMVs (Figure 1A).

Notably, although all size classes (<3 kDa, 3-30 kDa, 30-100 kDa) induced metamorphosis in

larvae of *H. elegans* (Figure 1B), the highest abundance of vesicles was also observed during

stationary phase (Figure 1C).

<u>Enzymatic interrogations of OMVs for larval metamorphic cues: Nucleic acids and proteins.</u> The

enzymatic interrogations of OMVs from *C. lytica* with nucleases and proteases had no impact on

the metamorphosis-inducing capacity of the OMVs (Figure 2A). The absence of nucleic acids in

the OMV samples was confirmed by electrophoresis and UV absorbance, and the enzymatic

destruction of proteins was confirmed by SDS-PAGE. Interestingly, both proteases had to be

removed from the treatment by ultracentrifugation prior to larval assay, because the residual

enzymes themselves induced metamorphosis (Fig S2).

<u>Enzymatic interrogations of OMVs for larval metamorphic cues: Lipids.</u>

The enzymatic interrogation of OMVs from *C. lytica* with lipases and phospholipases significantly

decreased OMV induction of metamorphosis (Figure 2B). Metamorphosis after OMV treatment

with Phospholipase A_1, while still significantly higher than the treatments with all other lipases,

was significantly lower than the untreated OMV and positive controls ($p<0.05$; Figure 2B). No

molecular structure-activity relationship for the metamorphic cue can be predicted from the

difference in reduction of metamorphosis between the phospholipase A_1 and A_2 treatments

(Figure 2B). Phospholipase A_2 is composed mostly of linear sheets and consequently will be

subject to less steric hindrance than the barrel shaped phospholipase A_1 (54). These results

indicate the involvement of a bioactive lipid.

<u>Enzymatic interrogations of OMVs for larval metamorphic cues: Peptidoglycan.</u> Enzymatic

interrogation of OMVs from *C. lytica* with lysozyme (to break down peptidoglycan) had no impact

on metamorphosis of larvae of *H. elegans* (Figure 2C).

<u>Bioassay guided fractionation of *C. lytica*.</u>

None of the HPLC fractions or purified compounds were as active as the parent samples. The

compounds isolated through the bioassay-guided fractionation of *C. lytica* were all extracted from

moderately active HPLC fractions of the cells of *C. lytica* (Figures S2-6). The amino acids

(phenylalanine and leucine), as well as, niacin, thymidine, adenosine, and a 2,5-diketopiperazine

of phenylalanine and proline were isolated and had minor levels of bioactivity (Figure 3).

Bioactivity of isolated compounds was consistent with their commercial counterparts (Figures S5-

6). It should be noted that induction of larval metamorphosis did not begin until after more than

6 h of exposure to any of the fractions, which is exceptionally slow.

Analysis of the genome of *C. lytica* with AntiSMASH (55, 56) identified one potential region of

secondary metabolite significance from 445 027–465 863 bp. A protein -BLAST of this contig,

identified it as a phytoene synthase (Table S1-2). Analysis of the genome of *C. lytica* with

NaPDos (57) identified three potential ketosynthase domains, (Table S3), however, closer

examination showed that these sequences are involved in fatty acid biosynthesis and are

unrelated to the production of bioactive secondary metabolites.

<u>Analysis of cell envelope components for larval metamorphic cues:</u> Isolated peptidoglycan at first

appeared to induce settlement in larvae of *H. elegans*, but the activity disappeared upon

treatment with lipase suggesting the isolated peptidoglycan was contaminated with a lipid. This

activity was unaffected by treatment with lysozyme (Figure 4). Commercial muramyl dipeptides

and muramyl tripeptides, tested across a concentration range of 1–100 µM, resulted in no

metamorphosis of larvae of *H. elegans* when tested separately or in combination.

Because lipases significantly reduced settlement induction by OMV samples and no significantly

inductive lipophilic small molecules were found in the natural products isolations, our focus

returned to the remaining major lipophilic component of the bacterial outer membrane,

lipopolysaccharide.

Extracted and purified LPS successfully induced metamorphosis in the larvae of *H. elegans*

(62%; Figure 5). Importantly, metamorphosis in larvae of H. elegans began less than one hour

after exposure to LPS samples. A dose-response curve generated for this LPS fraction, yielded a

peak of metamorphosis at 5 µg·ml^{-1}. Metamorphosis was not induced when the concentration of

the fraction dropped below 0.5 µg·ml^{-1}, and toxicity and death were observed in larvae exposed to

concentrations of LPS at or above 50 µg·ml^{-1} (Figure 5). Enzymatic treatment of the LPS sample

was consistent with the results gained from the OMV experiments; only lipase reduced the

induction of metamorphosis in larvae of *H.* elegans (Figure 5). The absence of peptidoglycan and

proteins from this preparation was confirmed using SDS-PAGE, comparing Coomassie blue

staining with silver staining (Figure 5). Additionally, the absence of proteins was indicated by a

lack of absorption of the sample at 260 nm (supplementary information, part E). The absence of

other contaminating glycolipids was confirmed with SDS-PAGE gels stained with the lipophilic

stain Pro-Q Emerald 300 (Figure 5) as well as TLC developed with 5-hydroxy-1-tetralone. The

Pro-Q Emerald 300 stain suggests that there could be at least 2 LPS molecules within the purified

LPS sample.

Discussion

The broadly distributed marine polychaete *Hydroides elegans* serves as a useful proxy for

undoubtedly thousands of benthic marine invertebrate species whose larvae settle and

metamorphose in response to specific biofilm-dwelling bacteria (6, 12, 23, 24, 34, 58-61).

However, it can be difficult to know whether bioactive chemicals identified in the laboratory are

the same as those that act *in situ* or simply stimulate processes downstream of the receptor (62–

66). The practice of isolating and identifying compounds involves the exposure of the larva to

substances with which it may not naturally interact (62). While experimentally valuable, artificial

induction of larval metamorphosis does not enable us to learn more of the true induction of
metamorphosis that is so essential to benthic marine organisms. Fortunately, the wealth of
knowledge accumulated for larval settlement conditions provides sound criteria for the
acceptance or rejection of the ecological relevance of any isolated cue. Our conditions for
accepting an ecologically relevant cue to induce metamorphosis in *H. elegans* require it to be
bacterial in origin, fast acting and entrained to the biofilm (26). It was with this focus that we
asked the question: what chemical component of marine biofilm bacterium *Cellulophaga lytica*
cues the larva of *H. elegans* to settle and metamorphose?
216
OMVs, a key component for bacterial interactions with the environment, can be considered a
secretion or a delivery system (42, 67). Previous studies have shown that OMVs produced during
different culture growth stages or of different sizes can vary in composition and biological activity
(42, 67, 68). Our data, however, demonstrated bioactivity in all growth stages, with a steady but
non-significant increase over time, suggesting that the bioactive compound produced by *C. lytica*
may be either constitutively produced or has a long half-life and may proportionately increase in
cultures over time. To determine whether OMVs induce metamorphosis through the delivery of
"cargo", we subjected OMV fractions from *C. lytica* to a battery of enzymatic treatments and
bioassay guided fractionation to identify components that impact larval settlement. Enzymatic
interrogations of the OMVs strongly indicated that induction of metamorphosis was not due to a
nucleic acid or protein cargo. Lipases, however, did inhibit induction of metamorphosis by the
OMVs, triggering an investigation for the presence of a small bioactive lipid with bioassay-guided
fractionation. Methanol and DCM extractions of OMV preparations followed by HPLC separation
and bioassays failed to identify any bioactive lipids.
231
The bioassay-guided fractionation did identify a few low-activity common small molecules such as
amino acids, but these were judged to not have ecological relevance. First, it is known that
induction of metamorphosis can be stimulated in larvae of *H. elegans* by artificial inducers
(25,64). Previous studies into the induction of metamorphosis in *H. elegans* by dissolved free
amino acids (DFAAs) was rejected due to delayed onset of metamorphosis, low levels of
metamorphosis, non-specificity of DFAAs, undetectability of DFAAs in field studies and the
presence of contaminating bacterial biofilms in the initial experimental set up (69–71). Second,
the turbulent flow present at the field site, Pearl Harbor, would quickly dissipate FAAs and keep
them well below inductive levels (29–31). Our results agree with these conclusions; when we
tested amino acids, metamorphosis was rare and delayed. Additionally, all the extracts, HPLC
fractions, and purified compounds were less active than their corresponding parent samples. For
these reasons, we concluded that none of the low-activity metabolites from the bioassay-guided

fractionations were involved in natural metamorphic induction. The *in-silico* searches further

confirmed the results from bioassay-guided fractionation approach; no gene clusters or domains

from known secondary metabolite biosynthetic pathways were identified that could transcribe

compounds likely to induce complex larval metamorphosis.

The results of the enzymatic interrogations of the OMVs, in combination with the discovery that, in

culture, OMVs were active settlement inducers regardless of their size or during which bacterial

growth phase they were collected, suggested that OMVs induced metamorphosis because they

were small pieces of cell membranes, rather than because they were acting as a delivery system.

For this reason, we next focused our investigations on studies of the bacterial cell membranes.

Cell membranes can contain a few larger lipid classes that may have been missed by the

bioassay guided fractionation, including phospholipids, glycolipids, and glycosphingolipids.

Phospholipids, although considered a major source for activation-dependent signaling molecules

(72), can be almost immediately dismissed from the list of potential settlement inducers. TLC

chromatography of the active LPS sample showed a complete absence of phospholipid.

Furthermore, not only is phospholipid signaling overwhelmingly intracellular (72, 73), but its

activity also arises from secondary messenger molecules derived from the membrane

phospholipids. These secondary messenger molecules are produced when phospholipases

selectively hydrolyze phospholipids (72, 73). Consequently, the phospholipases should have

increased production of the secondary messenger molecules and resulted in higher

metamorphosis and not the significant reduction in metamorphosis that was observed (Fig. 2).

Alternatively, the reduced activity of OMVs treated with phospholipase could reflect the

requirement for an intact or negatively charged membrane (73). Selective externalization of

phospholipids in the outer membrane can alter the biophysical properties of the membrane

affecting their binding properties (72). Additionally, intact membranes may hold active cue

molecules in particular conformations that would not occur if their components were in a free

solution. If that conformation is required to activate the larval receptor, removal of the compound

from the membrane would result in a similar loss of activity. This explanation, however, was

quickly rejected by the discovery of inductive activity among the remaining classes of compounds

obtained by extraction of the cell envelope. Rather, the inhibition of induction by lipase treatments

is most likely the result of non-specific enzyme action.

The remaining classes of compounds included the important MAMP groups, peptidoglycan and

lipopolysaccharide. Of these two groups, peptidoglycan has been shown to be an active

component of non-pathogenic bacterial signaling in other systems (74, 75), and, for this reason,

was chosen as the next component examined for its capacity to induce larval settlement. Results
of larval assays utilizing isolated peptidoglycan, commercial peptidoglycan monomers or OMVs
subjected to lysozyme treatments were all negative. These results are not entirely unexpected
since the peptidoglycan layer resides beneath the outer membrane and would not be the first
point of interaction between a larva and a bacterium or an OMV.

LPS, which could be described as a "prototypical MAMP", was found to induce metamorphosis in
larvae of *H. elegans* when provided at a concentration that reflects the concentration of cells in an
active biofilm of *C. lytica* (24)*.* In addition to being a major MAMP, LPS molecules comprise the
vast outer portion of the outer membrane of Gram-negative bacteria, and, as such, would be the
first bacterial molecules to interact with a larva upon contact, which is in line with how the process
naturally occurs (26). Similarly, LPS forms the outer portion of the outer membrane of OMVs and
is thus consistent with the observed activity of OMVs from *C. lytica*. It is important to note here
that, although isolations of LPS can include residual traces of both peptidoglycan and other
membrane lipids (phospholipids, glycosphingolipids etc.) (76), purity checks using both SDS-
PAGE and TLC revealed none of these classes of compounds in the active LPS extract.
Additionally, lipase and phospholipase treatments of the isolated LPS inhibited the inductive
activity, consistent with the results from the OMV samples and suggesting that non-specific
enzyme activity occurred. In other words, the discovery that the inductive cue for settlement and
metamorphosis of larvae of *H. elegans* is comprised of bacterial LPS is consistent with all of the
observations made previously and in this study.

While LPS may largely explain the widespread occurrence of inductive activity across Gram-
negative biofilm bacteria, it cannot account for the same activity found in Gram-positive bacteria
(34) which lack both an outer membrane and its LPS element. Instead, the cell envelope of
Gram-positive bacteria consists of a thickened cell wall of peptidoglycan that can be populated by
teichoic and lipoteichoic acids (LTA) (77,78). LTA is a major MAMP of Gram-positive bacteria; it is
a component of nearly all Gram-positive membranes, *as well as a functional equivalent to LPS*
(79). Thus, we predict that LTA from the inductive Gram-positive bacteria will be the active moiety
in induction of metamorphosis. However, given the great predominance of Gram-negative
bacteria in marine biofilms (80) may render this possibility irrelevant, i.e., settlement induction in
nature may always be by Gram-negative bacteria.

The strong immunogenicity of LPS, combined with its ubiquitous expression by Gram-negative
bacteria and inherent structural variability, serve to make it both a likely target of evolutionarily
conserved innate-immune receptors in larvae as well as suggest a possibly wide-spread

mechanism for bacterial induction of marine-invertebrate metamorphosis. LPS is a highly variable
macromolecule with this variability tied closely to bacterial taxonomic lines and growth conditions
(81-85). Indeed, our purified LPS sample appears to contain at least 3 different LPS molecules,
which is not unusual (86). LPS molecules are composed of three main components: lipid A
(endotoxin); an inner core oligosaccharide (kDO); and an outer core oligosaccharide (O-antigen),
with variation occurring primarily in the lipid A and O-antigen components (86). While beyond the
scope of this paper, determining the location of the inductive portion of the macromolecule or
whether it is a specific moiety of a specific LPS molecule produced by *C. lytica* will comprise our
next focus of study. If variations in the structure of LPS are responsible for the observed inductive
activity of different species and strains of bacteria, it can explain why many – but definitely not all
– marine Gram-negative bacteria induce settlement and metamorphosis in the larvae of
H. elegans (17, 23, 28) and other invertebrate species. It may also explain why not all strains of
the same bacterium induce larvae of a single invertebrate species to settle (37). Most importantly,
the inherent variation of LPS molecules could explain the differential settlement of different
invertebrates in different habitats.
331
Materials and Methods
Larval culture and bioassays: Dioecious adult *H. elegans were* collected from docks in Pearl
Harbor and maintained in the laboratory in continuously flowing, unfiltered seawater for several
weeks without a noticeable decrease in fecundity. Gametes were obtained and larval cultured
according to Nedved and Hadfield (23). Briefly, worms were induced to spawn by removal from
their tubes. Gametes were mixed. Fertilization and development proceeded rapidly to a feeding
trochophore stage in one day. Larvae were fed the single-celled alga *Isochrysis galbana* Tahitian
Strain at a concentration of approximately 60,000 cells·ml^{-1}. The larvae developed to the
metamorphically competent nectochaete stage by day five and were utilized in metamorphosis
experiments for 1-2 days. Because larvae of *H. elegans* will not metamorphose without exposure
to bacterial biofilms or their products, wild, complex natural films developed on glass slides
readily served as positive larval controls, and maintaining the larvae in double-filtered, autoclaved
seawater (DFASW) served as negative larval controls. Assays also included a biofilm of *C. lytica*
for reference. Assays were carried out in 24-well plates, and the percent of larvae that
metamorphosed was determined at 20–24 hours. Significant differences (p<0.05) were calculated
using Kruskal-Wallis followed by pairwise comparisons with false detection rate (FDR) correction
(87).
349
Bacterial culture and OMV isolation: *Cellulophaga lytica* were grown as previously described by
Freckelton et al. (34) on ½ FSWt media (88). Briefly, cryogen bacterial stocks were streaked onto

½ FSWt agar plates and incubated for 24-48 h at 25°C. Single colonies were then used to

inoculate 10 ml cultures of ½ FSWt broth and incubated for 12 h at 25°C. An aliquot of this starter

culture was then used to inoculate 200 ml cultures and incubated for a further 14 h at 25°C. Cells

were harvested by low-speed centrifugation (4000 g, 25 min 4°C). The cell-free broth was then

filtered (0.2 µm, Millipore) to remove any remaining cells and then ultra-centrifuged (200,000 g,

2 h, 4°C) (34, 89) to isolate OMVs. All samples were examined by TEM using negatively stained

preparations to confirm the presence of OMVs as previously described (34).

<u>Determination of relative concentration of OMVs:</u> OMV production is highly dependent on growth

conditions, and pelagic and biofilm bacterial populations can produce widely disparate amounts of

OMVs (41, 42). The amount of OMVs produced by a bacterium can likewise be impacted by the

growth phase of the population or the presence of other species. This variability results in

difficulties in predicting an ecologically relevant concentration for testing. Consequently, isolated

OMV suspensions were diluted and tested across three orders of magnitude. The lower (0.001X)

dilution reflects that of the number of cells in 1 h biofilm (10^5 cells/mm^2) (24) and the highest

concentration (1X) reflect the concentration used to inoculate those biofilms (10^8 cells·ml^{-1}). For

clarity of presentations all figures present the data for the lowest dilution only, unless otherwise

stated. Additional OMV dilutions are presented in the supplementary information part A.

To maintain consistent relative concentration of OMVs between culture batches an aliquot of

each sample was stained with the fluorescent lipophilic styryl dye FM-4-64 (Molecular Probes; T-

3166) following the protocol described by Schooling and Beverage (46). Briefly, samples were

suspended in Hank's buffered saline solution (HBSS) and stained with 20 µM FM-4-64 in the dark

for 20 min at 25°C. Fluorescence was read using a Molecular Devices SpectraMax M5

spectrophotometer (EX: 544 EM:680). Cholesterol was used to calibrate the dye's response.

<u>Analysis of OMVs from different growth phases of cultures of *C. lytica*:</u> Cultures of *C. lytica* were

monitored for growth rate by measuring the OD$_{600}$ of an aliquot hourly to determine the growth

phase. This process was repeated with triplicate cultures, and samples were collected at early

log, late log and stationary phases. OMVs were isolated from each sample time point as

described above and diluted to reflect the concentration of a culture at 10^5 cells·ml^{-1} (as above) in

keeping with their FM-46-4 readings.

<u>Analysis of OMV size classes:</u> Cell-free, filtered (0.2 µm), conditioned media was used to isolate

OMVs. OMVs from *C. lytica* were separated into size classes by ultrafiltration (Millipore, Centricon

Plus-70, 3 kDa, 30 kDa, 100 kDa). Aliquots (2 x 70 ml) of conditioned media were applied to each

ultrafilter and then centrifuged (4 000 g x 30 min x 4°C). Aliquots of the retentate and filtrate were

taken for metamorphosis assays and TEM analysis. The remaining filtrate was then applied to the

next ultrafilter (100 kDa > 30 kDa > 3 kDa).

<u>Enzyme treatment of OMVs:</u> Isolated OMVs were combined with enzymes (nucleases, proteases,

lipases, or lysozyme) and incubated in a water bath at the optimal temperatures and time periods

described in Table 1. Lysozyme is an enzyme that catalyzes the breakdown of 1,4-beta-linkages

between N-acetylmuramic acid and N-acetyl-D-glucosamine residues in peptidoglycan (90).

Enzyme action was halted by heat treatment alone, except for the proteases which had to be

removed from the treatment by ultracentrifugation prior to larval assay, because the residual

enzymes themselves induced metamorphosis (supplementary information part B, Fig S1). For

each enzyme or set of enzymes, a seawater control was also performed to ensure that the

responses observed were not a result of the enzymes or heat treatments alone. Successes of

nuclease treatments were assessed by spectrometry and electrophoresis. Successes of protease

treatments were assessed by spectrometry and SDS-PAGE.

<u>Bioassay-guided fractionation of *C. lytica*:</u> Cell pellets (CLP), cell-free supernates (CLS) and

OMVs were extracted with organic solvents directly to identify metamorphosis-inducing

molecules. CLP and OMV samples were extracted with methanol (MeOH) followed by a

dichloromethane-methanol mixture (DCM: MeOH, 1:1). Methanol was used as the first extracting

solvent to both quickly quench the cells and to remove any residual water. As this aqueous

methanol extract would have primarily targeted polar molecules, this extraction was followed by a

DCM: MeOH to recover any non-polar molecules. Some non-polar molecules will be extracted by

both solvent systems but should exist in higher concentration in the DCM: MeOH extract. CLS

(3.5 L) was extracted by liquid: liquid partition with ethyl acetate (EtOAc), with nonpolar

compounds extracted into the EtOAc layer.

The initial OMV extracts were subjected to metamorphosis bioassays first at 10 $\mu g \cdot ml^{-1}$. Any

samples that produced a toxic response in the larvae were re-assayed at 1 $\mu g \cdot ml^{-1}$ and 0.1 $\mu g \cdot ml^{-1}$

. Extracts were added to wells by dissolving in MeOH at the concentration required for the

addition of no more than 50 μL of MeOH to sample wells. MeOH was allowed to evaporate before

drying overnight in a lyophilizer to remove any residual MeOH. A MeOH solvent control was

added to the assay to ensure any observed induction of metamorphosis was not due to the

residual presence of MeOH.

Additionally, each extract (CLP, CLS, OMV) was subjected to reverse phase HPLC (Thermo

Scientific 3000), and all fractions were subjected to metamorphosis bioassay with larvae of

H. elegans. CLP HPLC fractions were generated for bioactivity testing through an acetonitrile

(ACN) gradient (10-25%, 20 mins; 25-40%, 10 mins; 40-100%, 20 mins; 100%, 10 mins). Active

fractions were further purified using HPLC (Phenomenex, Luna, 5 μm, C18, 100 Å, 250 x

100 mm, MeOH 10%, 30 mins). For CLS, extracts were fractionated using a phenyl-hexyl column

(Phenomenex, Luna, 5 μm, Phenyl-Hexyl, 250 x 100 mm) and a water: acetonitrile gradient.

^{1}H NMR and LC-ESI-MS data were collected on all active fractions using a Bruker (400 MHz) and

an Agilent 1260 HPLC system coupled to a 6120 quadrupole LC-MS spectrometer, respectively.

Active fractions underwent ^{1}H NMR and HR-MS analysis to assess purity, and, if fractions were

still mixtures of compounds, they underwent additional fractionation. This process was repeated

until pure fractions were obtained. Specific purification details of isolated compounds are provided

in the supplementary information part C.

<u>*In silico* exploration for the presence of secondary metabolite gene clusters in the genome of</u>

<u>*C. lytica*</u>: AntiSMASH and NaPDos are bioinformatic platforms that allow genome mining for

secondary metabolite biosynthesis gene clusters in bacterial genomes (55, 56) and allow rapid

genome-wide identification of secondary metabolite genes, C- or KS- domains or amino acids. *In*

silico analyses of the genome of *C. lytica* (CP009239.1) using both AntiSMASH 5.0 (56, 57) and

NaPDos (57) were run with default, relaxed parameters to determine the potential for *C. lytica* to

produce secondary metabolites (supplementary information part D, Tables S3-5).

<u>Extraction of cell-envelope peptidoglycan of *C. lytica*</u>: Peptidoglycans were obtained from cultures

of *C. lytica* with a protocol adapted from Leulier et al. (76). Culture growth was halted by chilling

cultures on ice to 4°C. Cells were collected by centrifugation, washed once with a cold 3.5% NaCl

solution and centrifuged again. Cells were then suspended in hot (95°C) aqueous SDS solution

(4%) for 30 min and then cooled to room temperature (1 h). The SDS was removed by

ultracentrifugation (20000 *xg;* 40 mins) and the pellet was resuspended in water and washed

again for a total of 5 washes. The pellet was then resuspended in water and incubated with both

DNase I, RNase, Proteinase K, Pronase E and Trypsin (Table 1) to remove any contaminating

proteins or nucleic acids. Enzymes were removed following the same ultracentrifuge protocol

described above. Metamorphic activity in the peptidoglycan extract was further interrogated with

either a lysozyme or lipase treatment. Induction of metamorphosis in larvae of *H. elegans was*

conducted before and after lipase and lysozyme treatments.

<u>Bioactivity testing of commercially available cell-envelope components,</u> Peptidoglycan monomers

in the form of muramyl dipeptides and muramyl tripeptides (Sigma Aldrich) were tested for

induction of metamorphosis in larvae of *H. elegans*, individually and across a range of

concentrations (0.1–10 µM), by dissolving them in water. Additionally, a solution containing 10 µM

concentrations of both compounds was also screened for metamorphic induction. LPS from

Escherichia coli 055:B5*, Salmonella enterica* serotype enteritidis and *Pseudomonas aeruginosa*

10 (Sigma Aldrich) were tested for induction of metamorphosis in larvae of *H. elegans*,

individually across a range of concentrations (0.1–10 µM), by dissolving them in DMSO adding

them to the sample wells before freezing and lyophilizing overnight to remove any residual

solvent. A solvent control was included in the assay to ensure any observed induction of

metamorphosis was not due to the presence of residual solvent. All commercial LPSs were

checked for contaminants with thin layer chromatography and ultraviolet spectroscopy.

<u>Extraction of cell-envelope lipopolysaccharide of *C. lytica*</u>: Lipopolysaccharides (LPS) were

extracted using the Westphal hot phenol method (91) as updated by Apicella et al. (92, 93). The

LPS fraction was assessed for metamorphic induction in larvae of *H. elegans* across a range of

concentrations determined from the inoculation density of metamorphically inductive biofilms of *C.*

lytica (10^5 cells·mm^{-2}) and estimates of the amount of LPS present per cell range from 25-

62 fg·cell^{-1} (94-96). This calculation estimates that active biofilms of *C. lytica* contain between 2

and 6 µg LPS per well in a metamorphosis assay. Consequently, a range from 0.5 µg·ml^{-1} to

10 µg·ml^{-1} LPS was utilized to develop a dose-response series for assessment of induction of

metamorphosis in larvae of *H. elegans*.

To reduce issues with solubility of the LPS extract and to determine a weight-for-volume

concentration, an aliquot of the concentrated LPS sample was dried completely using a speed-

vac. This mass was then used to calculate the concentration of the remaining solution. The LPS

sample was also subjected to enzymatic interrogation with lipase, lysozyme and pronase E as

described above.

<u>Analysis of LPS extract purity</u>: Aqueous LPS solutions were checked for contaminating proteins

by measuring the absorption at 260 nm. Electrophoretic behavior and additional protein

contamination checks were performed by SDS-PAGE (Coomassie Blue and Silver staining) (76,

97). The presence of glycolipids was identified with additional SDS-PAGE gels stained with

Invitrogen Pro-Q™ Emerald 300 Lipopolysaccharide Gel Stain Kit according to manufacturer's

instructions. The presence of contaminating lipid classes in the extracted LPS was checked by

utilizing the rapid fluorometric micro-detection system of Watanabe and Mizuta (98). Briefly, the

LPS extract was run on an alumina backed silica TLC plate (AAdvance Instruments, 60Å F254)

alongside LPS and glucose standards, as well as the phospholipid standard, phosphatidyl

ethanolamine (Sigma Aldritch). The TLC was developed by heating with acidified 5-hydroxy-1-

tetralone (98). Under UV light (365 nm) hexoses, including oligosaccharides and polysaccharides

containing hexoses appear as green spots; acidic and neutral glycosphingolipids appear as a

yellow spot and phospholipids appear as blue spots.

<u>Examination of bacterial preparations with transmission electron microscopy (TEM):</u> Bacterial

preparations and cell-free preparations were applied to glow-discharged formvar grids, negatively

stained with 2% uranyl acetate aqueous solution and air dried. Samples were imaged using a

120 kV Hitachi HT7700 electron microscope with an AMT XR-41 camera.

Acknowledgments

This research was funded in part by the Gordon and Betty Moore Foundation through Grant

GBMF5009 and the Office of Naval Research grant no. N00014-08-1-2658 to MGH. The authors

especially thank Prof. Michael Apicella for his invaluable advice regarding all things pertaining to

LPS and Mrs. Tina M. Carvalho at the Biological Electron Microscopy Facility, University of

Hawai☐i at Mānoa for her technical advice. Previous lab members, Nidhi Vijayan, Audrey

Asahina and Shaun Hennings assisted us in many ways, which we greatly appreciate.

References

1.   M. P. Schultz, J. A. Bendick, E. R. Holm, W. M. Hertel, Economic impact of biofouling on a
naval surface ship. *Biofouling* **27**, 87–98 (2011).

2.   M. G. Hadfield, Settlement and recruitment of marine invertebrates: a perspective and
some proposals. *B. Mar. Sci.* **39**, 418–425 (1986).

3.   D. J. Crisp, "Factors influencing the settlement of marine invertebrate larvae" in
Chemoreception in Marine Organisms, P. T. Grant, A. M. Mackie, Eds. (Academic Press,
1974), pp. 177–265.

4.   S. Dobretsov, D. Rittschof, Love at first taste: induction of larval settlement by marine
microbes. *Int. J. Mol. Sci.* **21**, 731 (2020).

5.   M. G. Hadfield, Biofilms and marine invertebrate larvae: What bacteria produce that larvae
use to choose settlement sites. *Annu. Rev. Mar. Sci.* **3**, 453–470 (2011).

6.   S. Whalan, N. S. Webster, Sponge larval settlement cues: the role of microbial biofilms in
a warming ocean. *Sci. Rep.* **4**, 4072 (2014).

7.   S. Whalan, P. Ettinger-Epstein, C. Battershill, R. de Nys, Larval vertical migration and
hierarchical selectivity of settlement in a brooding marine sponge. *Mar. Ecol. Prog. Ser.*
368, 145–54 (2008).

8.   M. A. A. Wahab, R. de Nys, N. Webster, S. Whalan, Larval behaviours and their
contribution to the distribution of the intertidal coral reef sponge *Carteriospongia*
foliascens. PLoS ONE **9**, 5 (2014).

9.   N. S. Webster, *et al.*, Metamorphosis of a scleractinian coral in response to microbial
biofilms. *Appl. Environ. Microbiol.* **70**, 1213–1221 (2004).

10.   N. S. Webster, R. Soo, R. Cobb, A. P. Negri, Elevated seawater temperature causes a
microbial shift on crustose coralline algae with implications for the recruitment of coral
larvae. *ISME J.* **5**, 759 (2011).

11.   J. F. Bertrand, R. M. Woollacott, G protein-linked receptors and induction of
metamorphosis in *Bugula stolonifera* (Bryozoa). *Invertebr. Biol.* **122**, 380–385 (2003).

12.   S. Dobretsov, P. Y. Qian, Facilitation and inhibition of larval attachment of the bryozoan
Bugula neritina in association with mono-species and multi-species biofilms. *J. Exp. Mar.*
Biol. Ecol. **333**, 263–274 (2006).

13.   W. Y. Bao, C. G. Satuito, J. L. Yang, H. Kitamura, Larval settlement and metamorphosis
of the mussel *Mytilus galloprovincialis* in response to biofilms. *Mar. Biol.* **150**, 565–574
(2007).

14.   M. N. Tamburri, M. W. Luckenbach, D. L. Breitburg, S. M. Bonniwell, Settlement of
Crassostrea ariakensis larvae: effects of substrate, biofilms, sediment and adult chemical
cues. *J. Shellfish Res.* **27**, 601–609 (2008).

15.   P. E. Gribben, J. T. Wright, W. A. O'Connor, P. Steinberg, Larval settlement preference of
a native bivalve: the influence of an invasive alga versus native substrata. *Aquat. Biol.* **7**,
217–227 (2009).

16.　A. C. Alfaro, T. Young, A. M. Ganesan, Regulatory effects of mussel (*Aulacomya*
maoriana Iredale 1915) larval settlement by neuroactive compounds, amino acids and
bacterial biofilms. *Aquaculture* **322**, 158–168 (2011).

17.　M. G. Hadfield, C. C. Unabia, C. M. Smith, T. M. Michael, "Settlement preferences of the
ubiquitous fouler Hydroides elegans." in *Recent Developments in Biofouling Control*,
(1994), pp. 65–74.

18.　M. J. Huggett, J. E. Williamson, R. de Nys, S. Kjelleberg, P. D. Steinberg, Larval
settlement of the common Australian sea urchin *Heliocidaris erythrogramma* in response
to bacteria from the surface of coralline algae. *Oecologia* **149**, 604–619 (2006).

19.　S. A. Dworjanyn, I. Pirozzi, Induction of settlement in the sea urchin *Tripneustes gratilla* by
macroalgae, biofilms and conspecifics: A role for bacteria? *Aquaculture* **274**, 268–274
(2008).

20.　J. A. Anderson, C. E. Epifanio, Induction of metamorphosis in the Asian shore crab
Hemigrapsus sanguineus: characterization of the cue associated with biofilm from adult
habitat. *J. Exp. Mar. Biol. Ecol.* **382**, 34–39 (2009).

21.　L. Khandeparker, A. C. Anil, S. Raghukumar, Relevance of biofilm bacteria in modulating
the larval metamorphosis of *Balanus amphitrite. FEMS Microbiol. Ecol.* **58**, 425–438
(2006).

22.　B. Roberts, *et al.*, A complement response may activate metamorphosis in the ascidian
Boltenia villosa. Dev. Genes Evol. **217**, 449-458. (2007).

23.　B. T. Nedved, M. G. Hadfield, "A Model for Biofouling Research" in *Marine and Industrial*
Biofouling, (Springer-Verlag, 2008), pp. 203–217.

24.　S. Huang, M. G. Hadfield, Composition and density of bacterial biofilms determine larval
settlement of the polychaete Hydroides elegans. *Mar. Ecol. Prog. Ser.* **260**, 161–172
(2003).

25.　E. Carpizo-Ituarte, M. G. Hadfield, Stimulation of metamorphosis in the polychaete
Hydroides elegans Haswell (Serpulidae). *Biol. Bull.* **194**, 14–24 (1998).

26.　M. G. Hadfield, B. T. Nedved, S. Wilbur, M. A. R. Koehl, Biofilm cue for larval settlement in
Hydroides elegans (Polychaeta): Is contact necessary? *Mar. Biol.* **161**, 2577–2587 (2014).

27.　E. R. Holm, B. T. Nedved, E. Carpizo-Ituarte, M. G. Hadfield, Metamorphic-signal
transduction in *Hydroides elegans* (Polychaeta: Serpulidae) is not mediated by a G
protein. *Biol. Bull.* **195**, 21–29 (1998).

28.　C. R. C. Unabia, M. G. Hadfield, Role of bacteria in larval settlement and metamorphosis
of the polychaete Hydroides elegans. *Mar. Biol.* **133**, 55–64 (1999).

29.　M. A. R. Koehl, Hydrodynamics of larval settlement into fouling communities. *Biofouling*
23, 357–368 (2007).

30.　M. A. R. Koehl, M. G. Hadfield, Hydrodynamics of larval settlement from a larva's point of
view. *Integr. Comp. Biol.* **50**, 539–551 (2010).

31.　M. A. R. Koehl, J. P. Crimaldi, D. E. Dombroski, Wind chop and ship wakes determine
hydrodynamic stresses on larvae settling on different microhabitats in fouling
communities. *Mar. Ecol. Prog. Ser.* **479**, 47–62 (2013).

32. S. C. K. Lau, P. Y. Qian, Larval settlement in the serpulid polychaete *Hydroides elegans* in
response to bacterial films: An investigation of the nature of putative larval settlement cue.
Mar. Biol. **138**, 321–328 (2001).

33. P. Y. Qian, S. C. Lau, H. U. Dahms, S. Dobretsov, T. Harder, Marine biofilms as mediators
of colonization by marine macroorganisms: implications for antifouling and aquaculture.
Mar. Biotechnol. **9**, 399–410 (2007).

34. M. L. Freckelton, B. T. Nedved, M. G. Hadfield, Induction of invertebrate larval settlement:
Different bacteria, different mechanisms? *Sci. Rep.* **7** (2017).

35. S. C. Lau, K. K. Mak, F. Chen, P. Y. Qian, Bioactivity of bacterial strains isolated from
marine biofilms in Hong Kong waters for the induction of larval settlement in the marine
polychaete *Hydroides elegans*. *Mar. Ecol. Prog. Ser.* **226**, 301–310 (2002).

36. N. Vijayan, M. G. Hadfield, Bacteria known to induce settlement of larvae of *Hydroides
elegans* are rare in natural inductive biofilm. *Aquat. Microb. Ecol.* **84**, 31-42 (2020).

37. Y. Huang, S. Callahan, M. G. Hadfield, Recruitment in the sea: Bacterial genes required
for inducing larval settlement in a polychaete worm. *Sci. Rep.* **2**, 228 (2012).

38. N. J. Shikuma, *et al.*, Marine tubeworm metamorphosis induced by arrays of bacterial
phage tail-like structures. *Science* **343**, 529–533 (2014).

39. C. F. Ericson, *et al.*, A contractile injection system stimulates tubeworm metamorphosis by
translocating a proteinaceous effector. *eLife, .* **8** (2019).

40. A. Y. Asahina, M. G. Hadfield, Complete genome sequence of *Cellulophaga lytica* HI1
using PacBio single-molecule real-time sequencing. *Genome Announc.* **2**, e01148-14
(2014).

41. B. L. Deatherage, B. T. Cookson, Membrane vesicle release in bacteria, eukaryotes, and
archaea: a conserved yet underappreciated aspect of microbial life. *Infect. Immun.* **80**,
1948–1957 (2012).

42. J. B. Lynch, R. A. Alegado, Spheres of hope, packets of doom: the good and bad of outer
membrane vesicles in interspecies and ecological dynamics. *J. Bacteriol.* **199**, e00012-17
(2017).

43. T. A. Koropatnick, Microbial factor-mediated development in a host-bacterial mutualism.
Science **306**, 1186–1188 (2004).

44. N. E. Madala, M. R. Leone, A. Molinaro, I. A. Dubery, Deciphering the structural and
biological properties of the lipid A moiety of lipopolysaccharides from *Burkholderia cepacia*
strain ASP B 2D, in *Arabidopsis thaliana*. *Glycobiology* **21**, 184–194 (2011).

45. L. Brown, J. M. Wolf, R. Prados-Rosales, A. Casadevall, Through the wall: extracellular
vesicles in Gram-positive bacteria, mycobacteria and fungi. *Nat. Rev. Microbiol.* **13**, 620–
630 (2015).

46. S. R. Schooling, T. J. Beveridge, Membrane vesicles: An overlooked component of the
matrices of biofilms. *J. Bacteriol.* **188**, 5945–5957 (2006).

47. A. J. Manning, M. J. Kuehn, Contribution of bacterial outer membrane vesicles to innate
bacterial defense. *BMC Microbiol.* **11**, 258 (2011).

48.   J. Berleman, M. Auer, The role of bacterial outer membrane vesicles for intra- and
interspecies delivery. *Environ. Microbiol.* **15**, 347–354 (2013).

49.   G. Raposo, W. Stoorvogel, Extracellular vesicles: Exosomes, microvesicles, and friends.
J. Cell Biol. **200**, 373–383 (2013).

50.   S. J. Biller, *et al.*, Bacterial vesicles in marine ecosystems. *Science* **343**, 183–186 (2014).

51.   M. Renelli, V. Matias, R. Y. Lo, T. J. Beveridge, DNA-containing membrane vesicles of
Pseudomonas aeruginosa PAO1 and their genetic transformation potential. *Microbiology*
150, 2161–2169 (2004).

52.   T. Baumgarten, *et al.*, Membrane vesicle formation as a multiple-stress response
mechanism enhances *Pseudomonas putida* DOT-T1E cell surface hydrophobicity and
biofilm formation. *Appl. Environ. Microbiol.* **78**, 6217–6224 (2012).

53.   T. N. Ellis, M. J. Kuehn, Virulence and immunomodulatory roles of bacterial outer
membrane vesicles. *Microbiol. Mol. Biol. Rev.* **74**, 81–94 (2010).

54.   L. de Maria, J. Vind, K. M. Oxenbøll, A. Svendsen, S. Patkar, Phospholipases and their
industrial applications. *Appl. Microbiol. Biotechnol.* **74**, 290–300 (2007).

55.   K. Blin, *et al.*, AntiSMASH 2.0-a versatile platform for genome mining of secondary
metabolite producers. *Nucleic acids Res.* **41**, w204–w212 (2013).

56.   M. H. Medema, *et al.*, AntiSMASH: Rapid identification, annotation and analysis of
secondary metabolite biosynthesis gene clusters in bacterial and fungal genome
sequences. *Nucleic acids Res.* **39**, W339–W346 (2011).

57.   N. Ziemert, *et al.*, The natural product domain seeker NaPDoS: A phylogeny based
bioinformatic tool to classify secondary metabolite gene diversity. *PLoS ONE* **7**, e34064
(2012).

58.   M. A. Wahab, R. de Nys, S. Whalan, Larval behaviour and settlement cues of a brooding
coral reef sponge. *Coral Reefs* **30**, 451–460 (2011).

59.   C. Tran, M. G. Hadfield, Larvae of *Pocillopora damicornis* (Anthozoa) settle and
metamorphose in response to surface-biofilm bacteria. *Mar. Ecol. Prog. Ser.* **433**, 85–96
(2011).

60.   J. Tebben, C. Motti, D. Tapiolas, P. Thomas-Hall, T. Harder, A coralline algal-associated
bacterium, *Pseudoalteromonas* strain J010, yields five new korormicins and a
bromopyrrole. *Mar. Drugs* **12**, 2802–2815 (2014).

61.   J. Tebben, *et al.*, Induction of larval metamorphosis of the coral *Acropora millepora* by
tetrabromopyrrole isolated from a *Pseudoalteromonas* bacterium. *PLoS ONE* **6**, e19802
(2011).

62.   R. A. Jensen, D. E. Morse, R. L. Petty, N. Hooker, Artificial induction of larval
metamorphosis by free fatty acids. *Mar. Ecol. Prog. Ser.* **67**, 55–71 (1990).

63.   R. W. Woollacott, M. G. Hadfield, Induction of metamorphosis in larvae of a sponge.
Invertebr. Biol. **115**, 257–262 (1996).

64.   A. J. Yool, *et al.*, Excess potassium induces larval metamorphosis in four marine
invertebrate species. *Biol. Bull.* **170**, 255–266 (1986).

65. C. M. Pearce, R. E. Scheibling, Induction of metamorphosis of larval echinoids
(*Strongylocentrotus droebachiensis* and *Echinarachnius parma*) by potassium chloride
(KCI). *Invertebr. Reprod. Dev.* **26**, 213–220 (1994).

66. B. M. Degnan, D. Souter, S. M. Degnan, S. C. Long, Induction of metamorphosis with
potassium ions requires development of competence and an anterior signalling centre in
the ascidian *Herdmania momus*. *Dev. Genes Evol.* **206**, 370–376 (1997).

67. A. Kulp, M. J. Kuehn, Biological Functions and Biogenesis of Secreted Bacterial Outer
Membrane Vesicles. *Annu. Rev. Microbiol.* **64**, 163–184 (2010).

68. C. Schwechheimer, A. Kulp, M. J. Kuehn, Modulation of bacterial outer membrane vesicle
production by envelope structure and content. *BMC Microbiol.* **14**, 324 (2014).

69. T. Jin, P.-Y. Qian, Effect of amino acids on larval metamorphosis of the polychaete
Hydroides elegans. *Mar. Ecol. Prog. Ser.* **267**, 209–218 (2004).

70. T. Harder, P.-Y. Qian, Induction of larval attachment and metamorphosis in the serpulid
polychaete *Hydroides elegans* by dissolved free amino acids: isolation and identification.
Mar. Ecol. Prog. Ser. **179**, 259–271 (1999).

71. M. Beckmann, T. Harder, P.-Y. Qian, Induction of larval attachment and metamorphosis in
the serpulid polychaete *Hydroides elegans* by dissolved free amino acids: mode of action
in laboratory bioassays. *Mar. Ecol. Prog. Ser.* **190**, 167–178 (1999).

72. V. B. O'Donnell, J. Rossjohn, M. J. O. Wakelam, Phospholipid signaling in innate immune
cells. *J. Clin. Investig.* **128**, 2670–2679 (2018).

73. J. H. Hurley, Y. Tsujishita, M. A. Pearson, Floundering about at cell membranes: a
structural view of phospholipid signaling. *Curr. Opin. Struct. Biol.* **10**, 737–743 (2000).

74. K. A. Cloud-Hansen, *et al.*, Breaching the great wall: peptidoglycan and microbial
interactions. *Nat. Rev. Microbiol.* **4**, 710–716 (2006).

75. S. V. Nyholm, M. McFall-Ngai, The winnowing: establishing the squid–vibrio symbiosis.
Nat. Rev. Microbiol. **2**, 632–642 (2004).

76. F. Leulier, *et al.*, The *Drosophila* immune system detects bacteria through specific
peptidoglycan recognition. *Nat. Immunol.* **4**, 478–484 (2003).

77. T. J. Silhavy, D. Kahne, S. Walker, The bacterial cell envelope. *Cold Spring Harb.*
Perspect. Biol. **2**, a000414 (2010).

78. S. Morath, A. Geyer, T. Hartung, Structure–function relationship of cytokine induction by
lipoteichoic acid from *Staphylococcus aureus*. *J. Exp. Med.* **193**, 393–398 (2001).

79. L. Brown, A. Kessler, P. Cabezas-Sanchez, J. L. Luque-Garcia, A. Casadevall,
Extracellular vesicles produced by the Gram-positive bacterium *Bacillus subtilis* are
disrupted by the lipopeptide surfactin. *Mol. Microbiol.* **93**, 183–198 (2014).

80. K. A. Lema, F. Constancias, S. A. Rice, M. G. Hadfield, High bacterial diversity in
nearshore and oceanic biofilms and their influence on larval settlement by Hydroides
elegans (Polychaeta). Environmental Microbiology 21, 3472–3488 (2019).

81. C. R. H. Raetz, C. M. Reynolds, M. S. Trent, R. E. Bishop, Lipid A modification systems in
Gram-negative bacteria. *Annu. Rev. Biochem.* **76**, 295–329 (2007).

82. Y. Zhang, C. Arias, C. Shoemaker, P. Klesius, Comparison of lipopolysaccharide and
protein profiles between *Flavobacterium columnare* strains from different genomovars. *J.*
Fish Dis. **29**, 657–663 (2006).

83. J. Schletter, H. Heine, A. J. Ulmer, E. T. Rietschel, Molecular mechanisms of endotoxin
activity. *Arch. Microbiol.* **164**, 383–389 (1995).

84. H. M. Aucken, T. L. Pitt, Lipopolysaccharide profile typing as a technique for comparative
typing of Gram-negative bacteria. *J. Clin. Microbiol.* **31**, 1286–1289 (1993).

85. K. Leker, I. Lozano-Pope, K. Bandyopadhyay, B. P. Choudhury, M. Obonyo, Comparison
of lipopolysaccharides composition of two different strains of *Helicobacter pylori. BMC*
Microbiol. **17**, 1–9 (2017).

86. J. D. King, D. Kocíncová, E. L. Westman, J. S. Lam, Lipopolysaccharide biosynthesis in
Pseudomonas aeruginosa. Inn. Immun. **15**, 261–312 (2009).

87. Y. Benjamini, A. Krieger, D. Yekutieli, Adaptive linear step-up procedures that control the
false discovery rate. *Biometrika* **93**, 491–507 (2006).

88. K. J. Boettcher, E. G. Ruby, Depressed light emission by symbiotic *Vibrio fischeri* of the
sepiolid squid *Euprymna scolopes. J. Bacteriol.* **172**, 3701–3706 (1990).

89. S. J. Gould, R. G, As we wait: coping with an imperfect nomenclature for extracellular
vesicles. *J Extracell. Vesicles* **2**, 3–5 (2013).

90. V. Tullio, S. Roberta, P. Manuela, Lysozymes in the animal kingdom. *Human and*
Mosquito Lysozymes: Old Molecules for New Approaches Against Malaria **35**, 45–57
(2015).

91. O. Westphal, Bacterial lipopolysaccharides extraction with phenol-water and further
applications of the procedure. *Meth. Carbohydr. Chem.* **5**, 83–91 (1965).

92. M. A. Apicella, "Isolation and characterization of lipopolysaccharides" in *Bacterial*
Pathogenesis: Methods and Protocols, F. R. DeLeo, M. Otto, Eds. (Humana Press, 2008),
pp. 3–13.

93. M. A. Apicella, J. Griffiss, H. Schneider, Isolation and characterization of
lipopolysaccharides, lipooligosaccharides, and lipid A. *Meth. Enzymol.* **235**, 242 (1994).

94. J. C. Hurley, Endotoxemia: Methods of detection and clinical correlates. *Clin. Microbiol.*
Rev. **8**, 268–292 (1995).

95. S. W. Watson, T. J. Novitsky, H. L. Quinby, F. W. Valois, Determination of bacterial
number and biomass in the marine environment. *Appl. Environ. Microbiol.* **33**, 940–946
(1977).

96. P. Yagupsky, N. FS, Quantitative aspects of septicemia. *Clin. Microbiol. Rev.* **3**, 269–279
(1990).

97. C.-M. Tsai, C. E. Frasch, A sensitive silver stain for detecting lipopolysaccharides in
polyacrylamide gels. *Anal. Biochem.* **119**, 115–119 (1982).

98. K. Watanabe, M. Mizuta, Fluorometric detection of glycosphingolipids on thin layer
chromatographic plates. *J. Lipid Res.* **36**, 1848–1855 (1995).

99.   D. Suck, DNA recognition by DNase I. *J. Mol. Recognit.* **7**, 65–70 (1994).

100.   K. D. Dyer, H. F. Rosenberg, The RNase A superfamily: Generation of diversity and innate
754   host defense. *Mol. Divers.* **10**, 585–597 (2006).

755   101.   P. J. Sweeney, J. M. Walker, "Proteinase K (EC 3.4.21.14)" in *Enzymes of Molecular*
756   *Biology*, (Humana Press), pp. 305–312.

757   102.   P. Bermejo-Barrera, S. Fernández-Nocelo, A. Moreda-Piñeiro, A. Bermejo-Barrera,
758   Usefulness of enzymatic hydrolysis procedures based on the use of pronase E as sample
759   pre-treatment for multi-element determination in biological materials. *J. Anal. At.*
760   *Spectrom.* **14**, 1893–1900 (1999).

761   103.   E. Altindis, Y. Fu, J. J. Mekalanos, Proteomic analysis of *Vibrio cholerae* outer membrane
762   vesicles. *PNAS* **111**, E1548–E1556 (2014).

763   104.   T. Nakae, H. Nikaido, Outer membrane as a diffusion barrier in *Salmonella typhimurium*.
764   Penetration of oligo- and polysaccharides into isolated outer membrane vesicles and cells
765   with degraded peptidoglycan layer. *J. Biol. Chem.* **250**, 7359–7365 (1975).

766   105.   V. K. Sidhu, F.-J. Vorhölter, K. Niehaus, S. A. Watt, Analysis of outer membrane vesicle
767   associated proteins isolated from the plant pathogenic bacterium *Xanthomonas*
768   *campestris* pv. *campestris. BMC Microbiol.* **8**, 87 (2008).

106.   G. S. Richmond, T. K. Smith, Phospholipases A1. *Int. J. Mol. Sci.* **12**, 588–612 (2011).

107.   M. Murakami, I. Kudo, Phospholipase A2. *J. Biochem.* **131**, 285–292 (2002).

Figures and Tables

Figure 1. Metamorphosis of larvae of *H. elegans* induced by outer membrane vesicles from

C. lytica. (A) Metamorphosis of larvae of *H. elegans* when exposed to OMVs from different size

classes. (B) Metamorphosis of larvae of *H. elegans* when exposed to OMVs from each growth

phase (mid log, late log and stationary). Metamorphosis was counted after 24 h exposure, with

filtered seawater serving as a negative control, and a multispecies biofilm serving as a positive

control. Letters show significant differences. (C) Negatively stained TEM image of OMVs from the

mid log and stationary stage of a culture of *C. lytica*. Scale bar 200 nm.

Figure 2. Induction of metamorphosis in larvae of *H. elegans* from enzyme treated outer membrane vesicles (OMVs) from *C. lytica*. OMVs and seawater (SW) controls were exposed to either (A) Nucleases (DNase (200 U) or RNase (200 U) and Proteases (trypsin (20 µg·ml^{-1}) followed by either Proteinase K or Pronase E (100 µg·ml^{-1}); (B) lipase (100 U), phospholipase A$_1$ (PLA1; 100 U), phospholipase A$_2$ (PLA2; 100 U) and phospholipase C (PLC; 100 U); (C) lysozyme (500 U). Metamorphosis was counted after 24 h exposure. Filtered seawater (SW) served as a negative control, and a multispecies biofilm, a *C. lytica* biofilm and an untreated *C. lytica* OMV sample were used as positive controls. Letters indicate significant differences.

Figure 3. Metamorphosis induction in larvae of *H. elegans* was consistently low from methanol extracts (10 μg·ml^{-1}) of A) isolated OMV fraction; B) cell-free culture supernatant containing outer membrane vesicles (OMVs) of *C. lytica;* C) cell pellet. Extracts were solubilized in ethanol, transferred to wells, evaporated to remove solvent and sterile seawater and larvae were added. Solvent control: ethanol, negative control: sterile seawater (SW), positive controls: Wild type (WT) biofilm and monospecific *C. lytica* biofilm.

800
801

Figure 4. A lipid that coextracted with peptidoglycan in the SDS extraction of cells of *C. lytica* induced metamorphosis in the larvae of *H. elegans*. Metamorphosis was counted 24 h after exposure to the SDS extract (10 μg·ml^{-1}), SDS extract that had been treated with lysozyme (500U) or SDS extract treated with lipase (100U). Negative control: sterile seawater (SW); positive controls a wild type (WT) biofilm, a *C. lytica* biofilm and *C. lytica* OMV sample.

Figure 5. LPS induces metamorphosis in larvae of *Hydroides elegans.* (A) Induction of

metamorphosis by LPS was not significantly affected by treatment with lysozyme or lipase

(lysozyme (500 U), lipase (100 U); negative control: sterile filtered seawater (SW); positive

control: wild-type (WT) biofilm and biofilm of *C. lytica*. Letters indicate significant differences). (B)

Metamorphosis dose response curve of LPS isolated from *C. lytica*. (C) Coomassie Blue stained

SDS-PAGE gel showing standard protein ladder, the presence of proteins in the OMVs isolated

from *C. lytica*, and the absence of proteins in the extracted LPS sample. (D) SDS-PAGE gel

stained with ProQ Emerald 300 showing *E. coli* LPS standard followed by boiled cells of *C. lytica*,

OMVs from *C. lytica* and isolated LPS from *C. lytica*.

Table 1: Enzyme treatments of OMVs from *C. lytica*

Cell component targeted	Enzyme	Incubation			Thermal Inactivation	
		Conc	Time	°C	°C	Time
Nucleic acid	DNAse (99)	200 U	2.5 h	37°C	75°C	10 min
	RNAse (100)	200 U	2.5 h	37°C	75°C	10 min
Proteins	Proteinase K (101)	100 µg·ml^{-1}	2.5 h	55°C	100°C	10 min
	Pronase E (102)	100 µg·ml^{-1}	2.5 h	55°C	100°C	10 min
	Trypsin (103)	20 µg·ml^{-1}	2.5 h	37°C	100°C	10 min
Peptidoglycan	Lysozyme (104)	500 U	2.5 h	37°C	100°C	10 min
Lipid	Lipase (105)	100 U	2.5 h	55°C	100°C	10 min
	Phospholipase A$_1$ (106)	100 U	2.5 h	55°C	100°C	10 min
	Phospholipase A$_2$ (107)	100 U	2.5 h	55°C	100°C	10 min
	Phospholipase C	100 U	2.5 h	55°C	100°C	10 min

PNAS

www.pnas.org

Supplementary Information for

Bacterial lipopolysaccharide induces settlement and metamorphosis in a marine larva.

Marnie L Freckelton, Brian T. Nedved, You-Sheng Cai, Shugeng Cao, Helen Turano,
Rosanna A. Alegado and Michael G. Hadfield[*]

*correspondence: Michael G. Hadfield

Email: hadfield@hawaii.edu

This PDF file includes:

Supplementary text

Figures S1 to S5

Tables S1 to S3

SI References

844
845

Supplementary Information Text

<u>Part A: Growth phase analysis of OMVs</u>

Figure. S1. Metamorphosis of larvae of *H. elegans* induced by outer membrane
vesicles from a culture of *C. lytica* at different growth phases. Metamorphosis of
larvae of *H. elegans* when exposed to OMVs from each growth phase (mid log, late log
and stationary). 1X is equivalent to the concentration of OMVs found in an inductive 10^8
cells·ml^{-1} biofilm. 0.01X is a 1/100th dilution. Metamorphosis was counted after 24 h
exposure, with filtered seawater serving as a negative control, and a multispecies biofilm
serving as a positive control. Letters show significant differences.

<u>Part B. Enzyme treatment of OMVs:</u>

Figure. S2. Proteases induce metamorphosis in larvae of *C. lytica*. Larvae of *H. elegans*
were exposed to either Proteinase K (5 U) or Pronase E (5 U). Metamorphosis was
counted after 24 h exposure. Filtered seawater (SW) served as a negative control, and a
multispecies biofilm and untreated OMVs were used as positive controls.

Part C: Bioassay-guided fractionation of *C. lytica*

1. Small molecule analysis of cell pellets from *C. lytica*:

Cell pellets of *C. lytica* were extracted with MeOH twice followed by repeated extraction

with MeOH: DCM (1:1). The MeOH: DCM extract was subjected to semi-preparative

reverse chromatography HPLC (C18, ACN: 10-25%, 20 mins; 25-40%, 10 mins; 40-

100%, 20 mins; 100% 10 mins). Fractions were collected every minute, dried and

resuspended at 20 mg/ml filtered seawater and tested for settlement-induction activity at

a concentration of 10 µg/ml. Activity was detected in fraction 7 and fraction 29; fraction

CLP-29 did not contain enough material or any indicative absorbance to pursue further.

Semi-preparative reverse chromatography of CLP7 (C18; MeOH 10%, 30 mins) yielded

activity in the 2nd and 4th fractions (Fig S3A). LC-Mass spectrometry and ^{1}H NMR

spectroscopy were consistent with the presence of phenylalanine and adenosine

(Fig S3B-D). To confirm the activity and further elucidate the stereochemistry, both

isolated and commercial versions of adenosine (Sigma-Aldritch) and phenylalanine (D

and L; Sigma-Aldritch) were tested for metamorphic induction (Fig S4). Additional

commercially available compounds (thymidine, leucine (D and L) and niacin) were also

tested (Fig S5).

2. Small molecule analysis of cell-free supernatant from *C. lytica* containing

OMVs.

Conditioned media of *C. lytica* cultures (3.7 L) were extracted with ethyl acetate (EtOAc)

and dried (363 mg). The EtOAc extract was subjected to semi-preparative reverse

chromatography HPLC (C18, ACN: 10-25%, 20 mins; 25-40%, 10 mins; 40-100%,

20 mins; 100% 10 mins). Fractions were collected every minute, dried and resuspended

at 20 mg/ml ethanol and tested for metamorphic activity at a concentration of (10 µg/ml).

An additional solvent (ethanol) control was included in bioassay. Activity was identified in

the 25[th] and 26[th] fractions which were combined. LC-MS revealed a compound

consistent with the presence of a diketopiperazine of phenylalanine and proline (Cyclo(-

Phe-Pro)). Isolated and commercial versions of this compound were subjected to

metamorphosis assay as previously described (Fig S6).

896
Figure. S3. Spectral confirmation of isolation of adenosine and phenylalanine from most
active cell pellet fraction (CLP7). (A) UV-Vis absorbance chromatogram B) Total Ion
Chromatogram (TIC) positive mode, Ions were detected within a mass range m/z 100-
1100. (C) Extracted ion chromatogram of the active peak at 2.9 min yielded m/z of
268.1043.(D) Extracted ion chromatogram of the active peak at 4.9 min yielded m/z of
123.0459.

Figure. S4. Metamorphosis of larvae of *H. elegans* when exposed to isolated (A) and
commercial (C) adenosine and isolated (B) and commercial (D) phenylalanine from
Fraction 7 of the cell pellet for 24 hrs. Positive Control: Wild Type (WT) Biofilm. Negative
Control: Sterile Seawater (SW).

Figure. S5. Metamorphosis of larvae of *H. elegans* when exposed to commercial
versions of compounds detected in fraction 7 for 24 hrs. A) Adenosine; B) DL-
Phenylalanine; C) L-Phenylalanine; D) D-Phenylalanine; E) Niacin; F) Thymidine; G)
L=Leucine; H) D-Leucine. Positive Control: Wild Type (WT) Biofilm. Negative Control:
Sterile Seawater (SW).

Figure. S6. Metamorphosis of larvae of *H. elegans* when exposed to commercial version
of the diketopiperizine of phenylalanine and proline (Cyclo(-Phe-Pro) detected in fraction
25 of the supernatant from a *C. lytica* culture for 24 hrs. Positive Control: Wild Type (WT)
Biofilm. Negative Control: Sterile Seawater (SW).

<u>Part D. Genome Mining for Secondary Metabolites</u>

Analysis of the genome of *C. lytica* for the presence of secondary metabolite

biosynthesis gene clusters revealed a single contig *(ctg1_392)* located between 455,027

- 455,863. A protein blast of this contig, identified it as a phytoene synthase (Table S1).

The Minimum Information about a Biosynthetic Gene cluster (MIBiG) failed to match any

enzymes with greater than 50% sequence identity (Table S2). Similar low matches were

generated using NaPDos (Table S3).

Table S1. NCBI BlastP on ctg1_392 from genome of *C. lytica* generated by AntiSMASH
(1)

Accession	Description	Max Score	Total Score	Query Cover	E value	% Ident
WP_013620042.1	phytoene synthase [*Cellulophaga lytica*]	563	563	100%	0.0	100.00%

934   **Table S2.** Minimum Information about a Biosynthetic Gene cluster (MIBiG) search
935   results from genome of *C. lytica* generated by AntiSMASH (1)

MIBiG Protein	Description	MIBiG Cluster	MiBiG Product	% ID	% Coverage	BLAST Score	E-value
ABB88949.1	CrtB	BGC0000650	terpene	50.0	99.6	255.0	1.2e-67
AAK64298.1	Phytoene synthase	BGC0000637	terpene	37.0	100.4	168.0	1.5e-41
ABD24399.1	Phytoene synthase	BGC0000644	terpene	33.0	97.1	155.0	9.9e-38
AAF65581.1	Phytoene synthase	BGC0000636	terpene	34.0	102.5	134.0	2.4e-31
ABP56869.1	Phytoene synthase	BGC0001087	saccharide-terpene	25.0	89.9	80.0	5.3e-15

936
937

938 **Table S3:** Polyketide Synthase (KS) domains within the genome of *C. lytica* generated
939 by NaPDos (2). Search was run with relaxed parameters.

	Query ID	Database match ID	% ident.	Align length	e-val	Pathway product
KS2	CP009239.1_3_157_15_425	FabF_Bacillus_FAS	55	409	5e^{-112}	fatty acid synthesis
KS1	CP009239.1_4_290_11_391	FabF_Bacillus_FAS	26	414	8e^{-24}	fatty acid synthesis
KS3	CP009239.1_6_112_8_221	Myca_YP880565_1KSB	30	131	2e^{-06}	mycocerosic acid synthase

941   <u>Part E: LPS purity checks</u>

944   **Figure S7.** UV-VIS absorbance spectrum of LPS sample shows no absorbance above
945   215 nm

SI References

1. M. H. Medema, *et al.*, AntiSMASH: Rapid identification, annotation and analysis of secondary metabolite biosynthesis gene clusters in bacterial and fungal genome sequences. *Nucleic Acids Res.* **39**, W339–W346 (2011).

2. N. Ziemert, *et al.*, The natural product domain seeker NaPDoS: A phylogeny based bioinformatic tool to classify secondary metabolite gene diversity. *PLoS One* **7**, e34064 (2012).