Title: ETV7 limits antiviral gene expression and control of SARS-CoV-2 and influenza viruses

Authors: Heather M. Froggatt[1], Alfred T. Harding[1], Brook E. Heaton[1], and Nicholas S. Heaton[1,*]

[1]Department of Molecular Genetics and Microbiology
Duke University School of Medicine
Durham, NC 27710, USA

*To whom correspondence should be addressed:
Nicholas S. Heaton, PhD
Assistant Professor
Department of Molecular Genetics and Microbiology (MGM)
Duke University Medical Center
213 Research Drive, 426 CARL Building, Box 3054
Durham, NC 27710
Tel: 919-684-1351
Fax: 919-684-2790
Email: nicholas.heaton@duke.edu

One Sentence Summary: ETV7 is an interferon-induced, repressive transcription factor that negatively regulates antiviral interferon-stimulated genes essential for controlling influenza virus and SARS-CoV-2 infections.

Abstract

The type I interferon (IFN) response is an important component of the innate immune response to viral infection. Precise control of interferon responses is critical; insufficient levels of interferon-stimulated genes (ISGs) can lead to a failure to restrict viral spread while excessive ISG activation can result in interferon-related pathologies. While both positive and negative regulatory factors control the magnitude and duration of IFN signaling, it is also appreciated that a number of ISGs regulate aspects of the interferon response themselves. However, the mechanisms underlying these ISG regulatory networks remain incompletely defined. In this study, we performed a CRISPR activation screen to identify new regulators of the type I IFN response. We identified ETS variant transcription factor 7 (ETV7), a strongly induced ISG, as a protein that acts as a negative regulator of the type I IFN response; however, ETV7 did not uniformly suppress ISG transcription. Instead, ETV7 preferentially targeted a subset of known antiviral ISGs. Further, we showed the subset of ETV7-modulated ISGs was particularly important for IFN-mediated control of some viruses including influenza viruses and SARS-CoV-2. Together, our data assign a function for ETV7 as an IFN response regulator and also identify ETV7 as a therapeutic target to increase innate responses and potentiate the efficacy of interferon-based antiviral therapies.

Introduction

The type I interferon (IFN) response is a transient innate immune defense system that, upon activation by viral infection or therapeutic IFN treatment, induces the transcription of hundreds of interferon-stimulated genes (ISGs) (*1*). Many ISGs have characterized antiviral roles that restrict viral replication by either interfering with viral processes directly or altering cellular pathways required for viral replication (*2*). However, because replication mechanisms and points of interaction with the cell differ between viruses, individual ISGs have varying potencies against different viruses (*3–5*). As a result, unique combinations of ISGs are thought to mediate successful antiviral responses against distinct viruses (*1*, *6*).

The canonical activation pathway of the type I IFN signaling pathway induced by viral infection, or after therapeutic administration (*7*), is well understood (*8*, *9*). Extracellular IFN is bound by its cognate plasma membrane-localized receptor (IFNAR1/2). Downstream effectors (JAK proteins) are phosphorylated to then activate formation of the interferon-stimulated gene factor 3 (ISGF3) complex. Finally, the ISGF3 complex of STAT1, STAT2, and IRF9 translocates to the nucleus (*8*) and binds the interferon sensitive response element (ISRE), with the consensus DNA motif GAAANNGAAA, to activate transcription of ISGs (*10*).

As infection is cleared and virally-derived innate immune activators become scarce, interferon production is reduced and the interferon-stimulated gene response is downregulated. To facilitate this return to cell homeostasis, IFN induced negative regulators, such as SOCS1 (*11*) and USP18 (*12*), act at multiple levels in the signaling pathway (*13*). Thus, negative regulators of IFN responses are an important group of IFN-stimulated genes that control the duration of ISG induction and activity. Antagonism of interferon response negative regulators has been proposed as a mechanism to enhance host antiviral responses to clear infection, both alone and in conjunction

with IFN treatment (*14–16*), and most recently during the COVID-19 pandemic (*17*). However,

regulators working upstream of transcription impact ISGs indiscriminately, including suppression

of the pro-inflammatory effectors, that when overabundant, can induce a cytokine storm; for

example, agonists of an IFNAR-downregulating protein, S1PR1 (*18*), are proposed for use against

pandemic influenza viruses (*19*) and SARS-CoV-2 (*20*) to instead limit excessive immune

responses associated with interferon signaling. An ideal negative regulator of the IFN response for

antiviral therapeutic targeting would enhance virus restricting ISGs specifically, without affecting

pro-immune cytokines.

In addition to the upstream regulators that broadly activate or suppress IFN responses, there are

interferon-induced transcriptional regulators that enhance, limit, or fine-tune ISG activity (*21*).

Many ISGs themselves participate in innate immune signaling to amplify IFN and other pro-

immune responses (*22*). Activators also add complexity by inducing non-canonical IFN response

pathways or specific groups of ISGs. Interferon responsive factors (IRFs) 1 and 7 are ISGs and

transcription factors that activate subsets of ISGs (*23*, *24*). Further, recent work has shown that

ELF1 (E74-like ETS transcription factor) is induced by IFN and facilitates the expression of a

group of genes not otherwise activated by the IFN response (*25*). Additional regulatory steps for

ISGs, post-JAK/STAT signaling, likely exist to allow the cell to fine-tune its antiviral activity for

an effective and appropriate response. While interferon-induced positive regulators of the IFN

response are known to shape the complexity of ISG activation, reports of analogous roles for

negative regulators remain conspicuously absent.

To address this gap in knowledge and identify genes able to shape the IFN response through

negative regulation, we performed a CRISPR activation (CRISPRa) screen that selected for factors

sufficient to prevent expression of an ISRE-containing IFN response reporter. We identified ETV7

(ETS variant transcription factor 7) as a negative regulator of the type I IFN response with a role

in controlling the expression of specific ISGs. We further showed the ETV7-modulated ISGs are

important for control of some IFN-sensitive respiratory viruses. Together, these data demonstrate

ETV7 is a suppressive component of the complex ISG regulatory network that could be targeted

to enhance specific antiviral responses against influenza viruses and SARS-CoV-2 (*1*, *26*).

Results

A CRISPR activation screen identifies ETV7 as a negative regulator of the type I IFN response.

In order to identify negative regulators of the type I IFN response, we developed a type I IFN

response reporter that included seven copies of the consensus interferon sensitive response element

(ISRE) ahead of a minimal CMV promoter controlling expression of sfGFP (**Fig. 1A**). To make

our reporter temporally specific, sfGFP was fused to a mouse ornithine decarboxylase (MODC)

protein degradation domain to decrease its half-life (*27*). We stably introduced this construct into

the A549 lung epithelial cell line along with a dCAS9-VP64 fusion protein and a MS2-p65-HSF1

activator complex required for the SAM CRISPR activation (CRISPRa) system (*28*). After clonal

selection, 99.8% of the A549-SAM-IFN response cells expressed GFP in response to type I IFN

treatment (**Fig. 1B and C**).

To perform the screen, we took the A549-SAM-IFN response cell line and introduced a lentivirus

library containing sgRNAs designed to activate every putative ORF in the human genome (*28*)

(**Fig. 1D**). After 48 hours, half of the cells were collected to determine the transduction efficiency

and the remaining cells were re-plated for IFN stimulation. At 72 hours post-sgRNA introduction,

the cells were treated with IFN-α and collected for fluorescence-activated cell sorting (FACS).

During sorting, we eliminated reporter-positive cells and collected only cells that were

nonresponsive to IFN as this population should theoretically be overexpressing a negative

Figure 1

Fig. 1. A CRISPR activation screen identifies ETV7 as a negative regulator of the type I interferon response. (**A**) Diagram of the IFN response reporter used to identify cells responding to IFN. ISRE = interferon sensitive response element, MODC = protein degradation domain. (**B**) Flow cytometry histogram and (**C**) bar graph of A549-SAM-IFN response cells before and after IFN-α treatment (1000 U/mL, 6 h) (data shown as mean ± SD, n=4, statistical analysis relative to

Fig. 1. (continued) untreated control). (**D**) Diagram of CRISPRa screen workflow to identify negative regulators of the type I IFN response. (**E**) Flow cytometry plots of 293T cells transfected with the IFN response reporter and overexpression plasmids for the indicated screen hits and then treated with IFN-α (100 U/mL, 9 h). (**F**) Quantification of **E** showing brightness of cells expressing GFP compared to the mCherry-expressing control (data shown as mean ± SD, n=4). (**G**) Brightness of cells expressing GFP from a constitutively expressing plasmid in cells overexpressing the indicated genes (positive control = EIF2AK1/HRI, shuts off translation) compared to control (data shown as mean ± SD, n=4). (**H**) Western blot showing ETV7 protein levels in 293T cells transfected with the ETV7 overexpression plasmid. Stain-free gel imaging was used to confirm equal loading. (**I**) Endogenous ISG mRNA expression levels measured using RT-qPCR after IFN-α treatment (100 U/mL, 9 h) (data shown as mean ± SD, n=4). (**J**) Western blot comparing IFIT1 protein levels in control and ETV7 overexpressing cells after IFN-α treatment (500 U/mL, 18 h). Stain-free gel imaging was used to confirm equal loading. For all panels: P-values calculated using unpaired, two-tailed Student's t-tests (*p<0.05, **p<0.001) compared to mCherry-expressing control samples unless otherwise noted.

regulator of the IFN response. We performed two independent biological replicates of the screen

and sequenced the sgRNA-containing amplicons derived from our input DNA, unselected

transduced cells, and cells that were nonresponsive to type I IFN. Raw sequencing data was

aligned, mapped, and subsequently analyzed using the MAGeCK pipeline (*29*) to generate z-score

values for each gene. Genes were defined as "hits" if their z-scores exceeded two standard

deviations from the mean, resulting in an overlap of 10 high-confidence genes between the two

screen replicates (**Fig. S1A, Table S1, Data files S1 and S2**). We were seeking to identify

regulators of the IFN response that are regulated by IFN themselves; therefore, we selected hits

for validation previously reported to have at least a two-fold induction after IFN stimulation in the

Interferome database (*30*). This analysis identified three hits (C1GALT1, ETV7, and NUP153) as

potential negative regulators of the type I IFN response (**Table S1**).

To validate our three hits, and to eliminate any potential off-target effects of CRISPRa, we cloned

the three ORFs and validated overexpression of the genes in 293T cells (**Fig. S1B**). Co-transfection

of the overexpression plasmids and IFN response plasmid, followed by stimulation with IFN-α,

resulted in lower reporter expression compared to a control mCherry-expressing plasmid (**Fig. 1E**

and F, Fig. S1C). To verify this repressive activity was specific to the IFN response and the hits

were not general inhibitors of transcription or translation, we transfected the overexpression

plasmids along with a constitutively active GFP-expressing plasmid. We included a positive

control (EIF2AK1/HRI), which is known to suppress host translation when overexpressed (*31*).

None of the potential hits significantly reduced expression of the GFP-expressing plasmid (**Fig.**

1G, Fig. S1D). We therefore chose ETV7 for further characterization because: 1) NUP153 has

previously been shown to control the distribution of STAT1 in the cell (*32*), 2) ETV7 had not been

reported to play a role in the IFN response, and 3) it had the strongest inhibitory phenotype against

the IFN response reporter.

After confirming overexpression of ETV7 at the protein level (**Fig. 1H**), we verified that the

inhibitory effects of ETV7 were not restricted to the IFN response reporter plasmid. We collected

mRNA and protein from IFN-α stimulated ETV7 overexpression cells to quantify effects on the

expression of endogenous ISGs. ETV7 overexpression significantly repressed the induction of

three prototypical ISGs (IFIT1, MX1, and ISG15) at the RNA level (**Fig. 1I**). The reduction of

ISG expression during ETV7 overexpression was also demonstrated at the protein level for IFIT1

(**Fig. 1J**). These data together demonstrate that overexpression of ETV7 alone is sufficient to

repress ISG induction by type I IFN.

ETV7 acts as a transcription factor to repress IFN-induced expression.

ETV7 is known to be a repressive transcription factor (*33, 34*), although a role in repressing type

I IFN responses has never been reported. To determine whether ETV7 acts as a transcription factor

in this context, we generated a previously validated mutant of ETV7, called ETV7(KALK), which

is unable to bind DNA (**Fig. 2A and B**) (*35*). Overexpression of ETV7(KALK) and stimulation

Figure 2

Fig. 2. ETV7 acts as a transcription factor to negatively regulate the type I IFN response. (**A**) Diagram showing the ETV7 protein domains and amino acid changes made to generate the DNA binding mutant, ETV7(KALK). (**B**) Western blot showing ETV7 protein levels in 293T cells transfected with WT or DNA binding mutant (KALK) ETV7 expression plasmids. Stain-free gel imaging was used to confirm equal loading. (**C**) Percent of 293T cells expressing GFP from the IFN response reporter with overexpression of WT or DNA binding mutant (KALK) ETV7 after IFN-α treatment (100 U/mL, 9 h) compared to control (data shown as mean ± SD, n=4). (**D**) ETV7's DNA binding position weighted matrix (PWM) generated using enoLOGOS (*99*) with data from Wei et al. (*36*). (**E**) Diagrams of the IFN response reporters containing (+ETS) and not containing (-ETS) potential ETV7 binding sites (ETS site, highlighted in yellow). (**F**) Normalized percent of 293T cells expressing GFP from IFN response reporters either containing or not

Fig. 2. (continued) containing ETS sites after overexpression of ETV7 and IFN-α treatment (100 U/mL, 6 h) compared to mCherry-expressing control (data shown as mean ± SD, n=4). (**G**) Diagram of the DNA pull-down experiment using biotinylated DNA and streptavidin-coated magnetic beads to show binding of a transcription factor to a specific DNA sequence. Sequences of biotinylated DNA bait used to show binding of ETV7 to the WT ISG15 ISRE (*93*) and loss of binding with mutation of the ETS binding site (-ETS). (**H**) Western blot of DNA pull-down using biotinylated oligos containing the ISG15 ISRE sequence in its wild-type form (WT) or with a single nucleotide mutation (-ETS) to eliminate the ETS site, incubated with nuclear lysate from cells expressing ETV7 with IFN treatment (100 U/ml, 9 h). (**I**) Heat map displaying RNA levels of genes upregulated at least 2.5-fold following IFN-α treatment (100 U/mL, 9 h) in control (mCherry) cells. Yellow = upregulated, blue = downregulated. For all panels: P-values calculated using unpaired, two-tailed Student's t-tests (*p<0.05, **p<0.001) relative to IFN-stimulated, mCherry-expressing control samples.

with IFN-α had no measurable effect on expression of the IFN response reporter, in contrast to

WT ETV7 overexpression (**Fig. 2C, Fig. S2A**).

ETV7 has been reported to bind the canonical ETS family DNA motif, GGAA (*36*), known as an

"ETS" site (**Fig. 2D**). The original IFN response reporter design contained multiple ETS sites (**Fig.**

2E), which likely explains why it is negatively impacted by ETV7 in our screen. To test the

requirement of ETS sites for ETV7 repressive activity against our reporter, we generated an IFN

response reporter containing seven consensus ISREs from canonical ISGs that all lack ETS sites

(ISRE -ETS) (**Fig. 2E**). We transfected the two reporter plasmids (ISRE +ETS and ISRE -ETS)

independently into 293T cells and stimulated with IFN-α. As expected, both reporter plasmids

responded to IFN treatment and were repressed by overexpression of known negative regulators

of the type I IFN response (SOCS1, USP18) that function upstream of transcription (**Fig. S2B**).

When ETV7 was transfected into cells with the IFN response reporters and stimulated with IFN

however, the repressive activity of ETV7 was restricted to the reporter plasmid containing ETS

motifs (**Fig. 2F, Fig. S2C**).

To provide evidence of ETV7 directly binding ISRE motifs containing ETS sites, we performed a

DNA oligo-based pull-down experiment using the ISG15 ISRE sequence (**Fig. 2G**). We chose this

ISRE because 1) it contains an ETS binding site, 2) it was included in our initial IFN response

reporter, and 3) ETV7 was shown to impact ISG15 induction (**Fig. 1I**). In addition to the wild-

type ISG15 ISRE biotinylated oligos, we tested biotinylated oligos with a single nucleotide

mutation that eliminates the ETS site in the ISG15 ISRE (-ETS). We generated nuclear lysates

from 293T cells overexpressing ETV7 and treated with IFN-α, incubated the lysate with either the

biotinylated WT or -ETS ISG15 ISRE oligos, and pulled down the biotinylated DNA using

streptavidin-coated magnetic beads. Western blotting for ETV7 protein revealed the transcription

factor bound the WT ISG15 ISRE containing the ETS site, but not the mutated version of the

promoter lacking the ETS site (**Fig. 2H**). Together, these experiments demonstrate that the

repressive activity of ETV7 requires both its ability to bind DNA and the presence of ETS sites in

the target ISG promoters.

ETV7 non-uniformly represses IFN-stimulated gene expression

Since consensus ISREs can either contain or lack a GGAA motif (**Table S2**), we hypothesized

ETV7 could differentially act on specific ISGs based on the presence or absence of ETS sites in

their ISRE sequences and promoters. To perform an unbiased examination of the effect of ETV7

upregulation, we performed RNA sequencing in cells with or without ETV7 overexpression and

IFN stimulation (**Data file S3 and Fig. S2E).** To identify the types of genes, processes, and

functions impacted by overexpression of ETV7, we generated lists of the most downregulated

genes during ETV7 overexpression, both before and after IFN treatment, and performed gene set

enrichment analyses. This analysis technique takes a gene list and returns overrepresented terms

from gene and protein databases (*37*). In the absence of IFN treatment, the most significantly

enriched terms were not biological pathways, but rather ETS family transcription factor binding

motifs (**Table S3**); this is expected, as ETV7 is an ETS family member and the entire family binds

the same core motif, GGAA. In contrast, the genes specifically downregulated by ETV7 during

IFN treatment were highly enriched for processes and pathways associated with responses to

interferon and viral infection (**Table S4**). To visualize the effect of ETV7 upregulation specifically

on ISGs, we generated a heat map of the genes induced at least 2.5-fold upon IFN treatment in our

RNA sequencing experiment (**Fig. 2I**). Consistent with our hypothesis, we found some genes were

suppressed by ETV7 overexpression more than others (**Fig. 2I**).

Although we had previously immunoprecipitated ETV7 with ISRE-containing oligos (**Fig. 2H**),

we next wanted to demonstrate ETV7 occupancy on native ISG promoters for genes targeted by

ETV7 via chromatin immune-precipitation followed by quantitative PCR (ChIP-qPCR). To enable

these experiments, we first generated a FLAG-tagged ETV7 (**Fig. S2E**) and demonstrated that the

tag does not disrupt its ability to repress the IFN response reporter (**Fig. S2F**). We next selected

the promoter of IFI44L for ChIP analysis as it was both highly affected by ETV7 and also harbors

many ETS sites within ISRE-like sequences in its promoter (**Fig. 2I and Fig. S2G and H**). To

perform ChIP-qPCR, we generated sheared chromatin after cross-linking cells transfected with

FLAG-tagged ETV7 and treated with IFN-α. We then performed immunoprecipitation using either

nonspecific IgG or anti-FLAG antibodies raised in mice or rabbits. After qPCR analysis, we found

no enrichment for a negative control region (the gene desert on chromosome 12) between the IgG

and anti-FLAG samples (**Fig. S2I**). In contrast, DNA corresponding to the IFI44L promoter was

significantly enriched with both the rabbit and mouse anti-FLAG samples compared to the

nonspecific IgG samples (**Fig. S2I**). These experiments demonstrate that ETV7 acts as a

transcription factor to bind and suppress ISG promoters that containing GGAA motifs.

ETV7 loss enhances antiviral IFN-stimulated gene expression.

Our experiments up to this point had primarily utilized overexpression of ETV7. One major disadvantage of this approach is the non-physiological kinetics and magnitude of ETV7 expression relative to what is observed after IFN stimulation in cells (**Fig. 3A**). To define the physiological effects of ETV7 induction during the IFN response, we performed a series of loss-of-function experiments. Our expectation was that the knockout of ETV7 would have the reciprocal effect on IFN responses as protein overexpression (*38*). We transduced A549-IFN response reporter cells (the original reporter with ISRE +ETS sites) with Cas9 and separately, two independent sgRNAs targeting ETV7, KO1(sg16667) and KO2(sg16668), generated pooled lines using antibiotic selection, and then stimulated with IFN-α. Both guides resulted in significantly more IFN-induced sfGFP expression compared to a control sgRNA (**Fig. 3B, Fig. S3A**).

We next generated clonal ETV7 knockout A549 lung epithelial cell lines (KO1-1, KO1-2) lacking the reporter and sequenced the resulting DNA lesions to confirm ETV7 knockout. Since ETV7 is normally only expressed after IFN stimulation, we treated with IFN-α and verified a reduction in ETV7 expression at the RNA level, presumably via nonsense mediated RNA decay (**Fig. 3C**). A representative housekeeping gene (ACTB) displayed no significant increases in transcription in any of the ETV7 KO clones (**Fig. 3D**). We then selected twelve ISGs (IFITM1, RSAD2, OAS1-3, IFIT1-3, IFI44L, ISG15, BST2, and IFI44) that were highly impacted by ETV7 in our RNA sequencing results for RT-qPCR analysis after IFN treatment. Compared to clonal lines containing a non-targeting guide, the ETV7 KO lines showed significant increases in induction of each of the twelve ISGs (**Fig. 3E-K, Fig. S3B-F**). Thus, the physiological induction of ETV7 after IFN stimulation affects the expression of ISGs.

Figure 3

Fig. 3. Loss of ETV7 enhances expression of ISGs. (**A**) ETV7 and other ISG mRNA levels in A549 lung epithelial cells after IFN-α treatment (100 U/mL, 6 h) (data shown as mean ± SD, n=4). (**B**) Percentage of cells expressing GFP from the IFN response reporter in two A549 ETV7 KO cell lines (pooled, 2 different guides) after IFN-α treatment (1000 U/mL, 6 h) compared to non-targeting control cells (data shown as mean ± SD, n=3). (**C**) mRNA levels of ETV7 in non-targeting control and ETV7 KO A549 clonal cell lines after IFN-α treatment (1000 U/mL, 6 h) (data shown as mean ± SD, n=4). (**D-K**) mRNA levels of a housekeeping gene (**D**) and ISGs (**E-K**) in control and ETV7 KO A549 clonal cell lines after IFN-α treatment (100 U/mL, 9 h) (data shown as mean ± SD, n=4). For all panels: P-values calculated using unpaired, two-tailed Student's t-tests (*p<0.05, **p<0.001) compared to IFN-stimulated, non-targeting sgRNA control samples.

Suppression of ETV7 enhances IFN-mediated control of influenza viruses and SARS-CoV-2.

In trying to predict the physiological significance of excessive induction of these ETV7-regulated

ISGs during the type I IFN response, we noted that many have recognized antiviral functions (*2,*

39, 40). To determine whether the effects of ETV7 suppression of ISG expression were relevant

in the context of a viral infection, we wanted to identify a virus restricted by the ISGs most affected

by ETV7 (*41*). Many of these genes with well-recognized antiviral functions (IFITM1, IFIT1-3,

OAS1-3, BST2, RSAD2) have been reported to play important roles in the restriction of influenza

viruses (*42*). IFITM1 has been shown to prevent viral entry (*43*), OAS proteins activate RNase L

to degrade viral RNA (*44*), IFITs bind viral RNA and promote antiviral signaling (*45*), and

BST2/Tetherin and RSAD2/Viperin restrict viral budding and egress (*46, 47*). Therefore, we

hypothesized that ETV7 dysregulation would affect influenza virus replication and spread.

We first infected our ETV7 KO A549 cells with a laboratory-adapted H1N1 influenza A virus

(IAV), A/Puerto Rico/8/1934 (PR8). Using a hemagglutination (HA) assay to measure the number

of viral particles released over time, we observed reduced virus production in our ETV7 KO cells

compared to control cells (**Fig. 4A**). This was the anticipated outcome because loss of a negative

regulator (i.e. ETV7) is expected to enhance expression of antiviral ISGs. We also measured

infectious viral titers and found a significant reduction in our ETV7 KO cells compared to control

cells (**Fig. S4A**). Using a fluorescent reporter strain of PR8 (PR8-mNeon) (*48*), we next visualized

infection and spread. As expected, we observed fewer cells expressing mNeon in ETV7 KO cells

using both microscopy (**Fig. 4B**) and flow cytometry readouts (**Fig. 4C**). We also tested whether

this phenotype would extend to a more contemporary H1N1 IAV strain, A/California/07/2009

(Cal/09), as well as an unrelated Victoria lineage influenza B virus strain, B/Malaysia/2506/2004

(Mal/04) (*49*). Using fluorescent reporter strains of these viruses, we observed significant

decreases in the number of Cal/09- and Mal/04-infected cells when comparing ETV7 KO cells to

control cells (**Fig. 4D and E**). In order to rule out non-IFN-related effects of ETV7 on inhibition

of influenza viral replication, we also performed these experiments with a non-influenza virus. We

selected Sendai virus (SeV) because, although it is known to induce a strong IFN response,

Fig. 4. Loss of ETV7 leads to restricted growth of multiple strains of influenza virus and SARS-CoV-2 with IFN treatment. (**A**) Hemagglutination (HA) assay of virus collected at indicated time points from non-targeting control and ETV7 KO A549 clonal cell lines after infection with WT PR8 virus (MOI=0.05, multicycle infection) (data shown as mean ± SD, n=3). (**B**) Control and ETV7 KO A549 clonal cell lines after mock or PR8-mNeon reporter virus infection (24 h, MOI=0.1, multicycle infection). Green = mNeon, blue = nuclei. Scale bar, 200 μm. (**C**) Flow cytometry quantification of control and ETV7 KO A549 clonal cell lines after infection with PR8-mNeon reporter virus (24 h, MOI=0.1, multicycle infection) (data shown as mean ± SD, n=4). (**D, E**) Percentage of infected (reporter+) cells from ETV7 KO A549 clonal cell lines compared to a control cell line after infection with (**D**) Cal/09-sfGFP or (**E**) Mal/04-mNeon

Fig. 4. (continued) reporter viruses (24 h, multicycle infection) (data shown as mean ± SD, n=4). (**F**) Percentage of infected (reporter+) cells from ETV7 KO A549 clonal cell lines compared to a control cell line after infection with Sendai-GFP reporter virus (24 h) (data shown as mean ± SD, n=4). (**G**) Percentage of infected (reporter+) cells from ETV7 KO A549 clonal cell lines compared to a control cell line after pre-treatment with IFN-α (100 U/ml, 6 h) then infection with PR8-mNeon reporter virus (24 h, MOI=0.1, multicycle) (data shown as mean ± SD, n=4). (**H**) qPCR detecting SARS-CoV-2 N vRNA in infected cells from ETV7 KO A549 clonal cell lines compared to a control cell line after pre-treatment with IFN-α (1000 U/ml, 6 h) then infection with SARS-CoV-2 virus (24 h, MOI=0.1) (data shown as mean ± SD, n=4). For all panels: P-values calculated using unpaired, two-tailed Student's t-tests (*p<0.05, **p<0.001) compared to infected, non-targeting sgRNA control samples.

previous reports (*50*) show SeV replication is relatively unaffected by type I IFN. Furthermore,

many of the ISGs most impacted by ETV7 have little to no effect against SeV, including IFITM1

(*51*), IFIT1 (*52*), and BST2/Tetherin (*53*). Using of a fluorescent reporter SeV, we observed no

significant change in the number of infected cells when comparing ETV7 KO cells to control cells

(**Fig. 4F**).

In addition to physiological induction of IFN after infection, exogenous IFN treatment is

frequently utilized or proposed as an antiviral therapy (*7*), either when there is difficulty

developing targeted antivirals (e.g. hepatitis C virus (*54*)) or there is an unexpected emergence of

a viral pathogen (e.g. swine or highly pathogenic avian influenza viruses (*55*), SARS-CoV (*56*),

Ebola virus (*57*), SARS-CoV-2 (*58–60*)). Therefore, we wanted to assess whether suppressing

ETV7 could be an effective strategy to enhance the antiviral effects of therapeutic IFN

administration. In support of this idea, we found treatment of ETV7 KO cells with IFN-α prior to

infection with PR8-mNeon led to an additional 5-fold enhancement of viral restriction compared

to IFN-treated WT control cells (**Fig. 4G**). Finally, since recent reports indicate that although

SARS-CoV-2 infections fail to induce a strong IFN responses (*61–63*), the virus is susceptible to

IFN treatment (*64–66*), we wanted to test if loss of ETV7 would potentiate IFN-mediated

suppression of SARS-CoV-2. With the same experimental design as the influenza virus

experiment, we found that pre-treatment of ETV7 KO cells with IFN-α resulted in a further 10-

fold reduction in SARS-CoV-2 vRNA compared to IFN-treated control cells (**Fig. 4H**). This

enhanced control was likely mediated via the ISG LY6E, which potently restricts SARS-CoV-2

(*67*, *68*) and we found to be regulated by ETV7 (**Fig. S4B**). Together, these experiments

demonstrate the potential of targeting ETV7 to enhance either the physiological or therapeutic

effects of IFN to control viral infection.

Discussion

In this study, we performed a CRISPR activation screen to identify negative regulators of the type

I IFN response. Specifically, we were interested in negative regulators that contribute to the types

of differentiated ISG profiles that are essential for effective control of viral infections. From this

screen, we identified ETV7 as a negative IFN regulator and subsequently showed it acts as a

transcription factor to repress subsets of ISGs through its recognized DNA binding motif. We also

demonstrated that the regulatory activity of ETV7 impacts the replication and spread of multiple

strains of influenza viruses. ETV7 loss also increased the antiviral effects of exogenous interferon

treatment against an H1N1 influenza virus and SARS-CoV-2. These findings demonstrate the

importance of ETV7 in fine-tuning the IFN response through specificity and transcriptional

repression to regulate antiviral ISG targets, and its potential as a target to enhance antiviral IFN-

based therapeutics.

ETV7 is a member of the ETS family of transcription factors. This family performs diverse

functions despite recognizing the same core DNA sequence, GGAA. Because these factors share

the same core motif, specificities are gained in other ways such as expression patterns (cell type,

basal expression, or immune pathway induction), active or repressive activity, and potential

binding partners (*69*, *70*). ETV7 is induced as an ISG and is repressive, whereas most ETS

transcription factors are activators, including ETS transcription factors recognized to assist in the

induction of ISG expression (e.g. ELF1 (*25*), PU.1 (*71*)). We also found repeatedly in our work

that the reporters and genes impacted by ETV7 contained ETS sites within ISRE sequences (**Fig.**

2E-H, S2G-I, S4B). This is not unexpected because it is recognized that many ETS family

members bind extended motifs similar to ISREs called ETS-IRF composite elements (EICE) (*72*).

EICE-associated activity is reported to require an IRF binding partner to direct ETS transcription

factor activity (*71*, *73*); therefore, it is likely ETV7 has an IRF binding partner. If ETV7 does

require a binding partner, this protein's induction and distribution likely contribute to the timing,

gene targets, and activity of ETV7 during the IFN response. It is known that IRFs can be basally

expressed (IRF2, IRF3), IFN-induced (IRF1, IRF7), or IFN-responsive (IRF9) (*72*), and the

availability of a binding partner could dramatically affect the timing and magnitude of effects on

EICE-controlled ISGs. Future work will define if ETV7 has specific binding partners and how

those interactions may contribute to the nonuniform, repressive activity of ETV7 during the type

I IFN response reported in this study.

IFN-induced regulators control the magnitude and duration of IFN responses in addition to the

temporal regulation of specific waves of ISGs (*74*). These coordinated waves of ISG induction can

peak early or late during the IFN response and are thought to correspond to specific stages of virus

replication or immune processes (*1*, *6*). We compared the induction of ETV7 with IFI44L, IFIT1,

ISG15 and RSAD2, and we observed ETV7 is upregulated at earlier time points than these

prototypical ISGs (**Fig. 3A**). We expanded our analysis to published datasets of human gene

expression during respiratory infections and concluded that ETV7 is generally induced earlier than

many ISGs (*75*). Although not the focus of our study, ETV7's early induction pattern suggests it

may be a key regulator of the first stages of IFN-mediated gene induction. We favor a model

wherein early ETV7 expression is responsible for reducing the accumulation, or delaying the

expression, of ISGs controlled by ISREs and promoters containing ETS motifs.

ETV7 is induced during infections across many vertebrate species (*76, 77*), indicating a potential

conserved, relevant role in the immune response; however, ETV7 has been lost in mice and closely

related rodents (*78*). Since mice and rodents have an intact interferon response pathway, a natural

question is: how are the activities of ETV7 being accounted for in these animals? While we have

no clear answer from the data in this study, it is well-recognized that IFN responses contain many

redundancies (*41*). Accordingly, we believe other ETS family members, potentially the closely

related ETV6 (which is also induced by IFNs), may perform the role of ETV7 in mice (*79*). Future

studies will be required to test the hypothesis that mice induce an ETV7-related alternative during

the type I IFN response.

Another important question is why the IFN-induced activity of ETV7 has been maintained

throughout evolution. In this report, we provide evidence that ETV7's activity reduces a cell's

ability to restrict virus infection; this seems counterintuitive to ETV7 benefitting the host. We

hypothesize that regulators like ETV7 are important to prevent excessive inflammatory signaling.

It is appreciated that negative regulators of the IFN response are required to prevent extreme and

prolonged immune responses, which are associated with poor disease outcomes after infection

(*80–82*). ETV7 potentially contributes to the cumulative activities of negative IFN regulators to

limit IFN responses during pathogen clearance. Additionally, it stands to reason that individual

ISGs have different toxic effects on the cell. It is tempting to speculate that ETV7 suppresses ISGs

whose accumulation is particularly harmful to cell viability and host recovery after infection.

Additionally, the relevance of controlled IFN responses goes beyond infectious disease; patients

with dysfunctional USP18, a negative regulator of the IFN response, develop a type I

interferonopathy that results in a severe pseudo-TORCH (Toxoplasmosis, Other agents, Rubella,

Cytomegalovirus, and Herpes simplex) syndrome (*83*). Mouse knockouts for other negative

regulators of the IFN response (SOCS1, SOCS3, USP18) also develop non-pathogen associated,

chronic inflammatory diseases (*84–87*). ETV7's lack of a murine homolog eliminates an easily

generated animal knockout model to experimentally show ETV7's relevance as a general innate

immune repressor. However, genome-wide association studies (GWAS) have linked ETV7 to

autoimmune diseases including rheumatoid arthritis and multiple sclerosis (*88*, *89*); both of these

autoimmune diseases have evidence of enhanced ISG expression (*90*, *91*). Thus, although the

specific contributions of ETV7 activity to IFN regulation are currently undefined, its potential role

is not limited to viral infections.

In conclusion, this study identified ETV7 as a negative regulator of the type I IFN response.

Previously, ETV7 was appreciated to be an ISG; however, a specific function during the IFN

response was unknown. We determined that ETV7 acts as a transcription factor to target specific

ISGs for repression, potentially contributing to the complex ISG transcriptional landscape.

Additionally, many of the ETV7-regulated ISGs restrict respiratory viruses (*42*, *67*, *68*), and we

showed that loss of ETV7 can further enhance the ability of type I IFN to control influenza virus

and SARS-CoV-2 replication. Further work is required to understand the complexity of IFN

regulation, while therapeutic targeting of factors like ETV7 could lead to the development of a

new class of host-directed antivirals that can enhance or tailor ISG responses to specific viruses.

Materials and Methods

Cloning

To generate reporters sensitive to IFN, we designed gBlocks (IDT) containing ISREs to be cloned into the pTRIP vector ahead of a minimal CMV promoter controlling expression of sfGFP. To clone and express the open reading frames (ORFs) of our screen hits, we designed primers for cloning into the pLEX-MCS vector using Gibson Assembly (NEB). To amplify ETV7, NUP153 and USP18 we used cDNA templates from Transomic Technologies. To amplify C1GALT1 and EIF2AK1 we used RNA from IFN-stimulated A549 cells. The DNA binding mutant ETV7, ETV7(KALK) (*35*), and codon-optimized SOCS1 expression plasmids were generated using a gBlock (IDT). FLAG-tagged ETV7, ETV7(FLAG), was generated using a primer including the FLAG tag. Non-targeting and ETV7-targeting CRISPR KO sgRNAs were cloned by annealing oligos encoding the desired sgRNA sequence and ligating them directly into the lentiCRISPRv2 vector (Addgene). DNA was transformed into NEB 5-alpha high efficiency competent cells. Insert size was verified with PCR and purified plasmids were sequenced using Sanger sequencing.

Cells

All cells were obtained from ATCC and grown at 37°C in 5% CO_2. A549 and 293T cells were grown in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 5% fetal bovine serum, GlutaMAX, and penicillin-streptomycin. Madin-Darby canine kidney (MDCK) cells were grown in minimal essential media (MEM) supplemented with 5% fetal bovine serum, HEPES, $NaHCO_3$, GlutaMAX, and penicillin-streptomycin. The A549 CRISPR-SAM cells were previously validated (*92*) and transduced with the IFN response reporter three times before being clonally selected. The A549 CRISPR KO cells were transduced (ETV7 KO1 = sg16667, ETV7 KO2 = sg16668, nontargeting control) and then selected using puromycin (10 μg/mL). A549-IFN response reporter ETV7 KO lines were selected with puromycin and used as pooled lines. A549

ETV7 KO lines were selected with puromycin and subsequently plated at a dilution to isolate

single cells, which were grown until colonies of an appropriate size allowed for collection. These

clonally selected colonies were grown up and verified to be KO lines (control, KO1-1, KO1-2,

KO2-1, KO2-2) by sequencing the DNA lesions generated as a result of Cas9 endonuclease

activity, results shown below. Guide sequences = underlined, insertions = bolded, deletions =

dashes, exonic sequences = uppercase, intronic sequences = lowercase.

ETV7 KO guide 1 (sg16667)
WT   5' CTG CCA TGC ACC G<u>CG GAG CAC GGG TTC GAG ATG</u> AAC GGA CGC GCC 3'
403   protein length = 342aa
404
405 KO1-1 allele 1 5' CTG CCA TGC ACC GCG GA**A** GCA CGG GTT CGA GAT GAA CGG ACG CGC 3'
406   predicted protein length = 104aa
407 KO1-1 allele 2 5' CTG CCA TGC ACC GCG GA**A** GCA CGG GTT CGA GAT GAA CGG ACG CGC 3'
408   predicted protein length = 104aa
409
410 KO1-2 allele 1 5' CTG CCA TGC ACC GCG GA**A** GCA CGG GTT CGA GAT GAA CGG ACG CGC 3'
411   predicted protein length = 104aa
412 KO1-2 allele 2 5' CTG CCA TGC ACC GCG GA**A** GCA CGG GTT CGA GAT GAA CGG ACG CGC 3'
413   predicted protein length = 104aa
414
415 ETV7 KO guide 2 (sg16668)
416 WT   5' GCG ATG CCG CAG GCC <u>CCC ATT GAC GGC AGG ATC GCT</u> Ggtgagtgggagg 3'
417   protein length = 342aa
418
419 KO2-1 allele 1 5' GCG ATG CCG CAG GCC CCC ATT GAC G-- --- --- -CT Ggtgagtgggagg 3'
420   predicted protein length = 333aa
421 KO2-1 allele 2 5' GCG ATG CCG CAG GCC CCC ATT GAC GGC AGG --- --- --tgagtgggagg 3'
422   predicted protein length = unknown, loss of splice site
423
424 KO2-2 allele 1 5' GCG ATG CCG CAG GCC CCC ATT GAC G-- --- --- -CT Ggtgagtgggagg 3'
425   predicted protein length = 333aa
426 KO2-2 allele 2 5' GCG ATG CCG CAG GCC CCC ATT GAC GGC AGG -T**GA** CGC TGgtgagtgggagg 3'
427   predicted protein length = 220aa
428

Flow Cytometry

Cells were trypsinized and analyzed on a Fortessa X-20 (BD) machine with standard laser and

filter combinations. Data was visualized and processed with FlowJo software.

CRISPR Activation Screen

The sgRNA library was packaged into lentivirus as previously described (*92*). After packaging

and titering the lentivirus, $2x10^8$ A549-CRISPR-SAM-IFN response reporter cells were seeded

onto 15 cm plates (10 plates total). The next day they were transduced with the packaged sgRNA

library (MOI=0.5). After 48 h, the transduced cells were split and half were collected as a

transduction control, while the remaining half were plated back onto 15 cm plates. The next day,

cells were treated with IFN-α ($4x10^3$ U/mL) for 6 h. Cells were then collected and sorted on a

Beckman Coulter Astrios cell sorter. Specifically, gates were set to sort GFP-negative cells as the

population of interest, as well as GFP-positive cells as a control population of cells still capable of

signaling. This screen was performed in duplicate. Genomic DNA was extracted from sorted cells

using the Zymo Quick gDNA micro prep kit. PCR was subsequently performed using barcoded

primers as previously described using the NEB Next High Fidelity 2x PCR master mix (*92*). PCR

bands were gel purified using the Thermo GeneJET gel extraction kit. Samples were then

sequenced via next-generation Illumina MiSeq using paired-end 150 bp reads.

Screen Analysis

Raw MiSeq read files were aligned to the CRISPR SAM sgRNA library and raw reads for each

sgRNA were counted using the MAGeCK pipeline (*29*). sgRNA enrichment was determined using

the generated count files and the MAGeCK-MLE analysis pipeline. Genes were sorted based on

z-score and determined to be significantly enriched if their z-score was at least two standard

deviations above the average z-score of the entire sorted population.

Western Blotting

Cells were trypsinized and $1x10^6$ cells were pelleted at 800 x g for 5 min. Equal amounts of cellular

material were loaded into 4-20% acrylamide gels (Bio-Rad) and imaged using a ChemiDoc

Imaging System (Bio-Rad). Protein was transferred to a nitrocellulose membrane at 60V for 60

min. PBS with 5% (w/v) non-fat dried milk and 0.1% Tween-20 were used to block for 1 h at 4°C.

Primary antibodies were then incubated with the membrane overnight at 4°C. Antibodies used

were rabbit anti-ETV7 (Sigma, HPA029033), rabbit anti-IFIT1 (Cell Signaling, D2X9Z), mouse

anti-FLAG (Sigma, F3165) and mouse anti-αtubulin (Sigma, T5168). Membranes were washed

five times in PBS with 0.1% Tween-20 and then an anti-rabbit-HRP (Thermo, A16104) or anti-

mouse-HRP (Thermo, A16072) secondary antibody was added for 1 h. The membrane was then

washed five times and Clarity or Clarity Max ECL substrate (Bio-Rad) was added before being

exposed to film and developed.

RT-qPCR

For all experiments except SARS-CoV-2 infection experiments, total RNA was collected and

prepared using Monarch Total RNA Miniprep Kits (NEB). For SARS-CoV-2 experiments, cells

were collected in TRIzol (Invitrogen) and RNA was isolated using Phasemaker tubes (Invitrogen).

One-step RT-qPCR was performed with commercial TaqMan assays from Thermo for ETV7

(Hs00903229_m1), C1GALT1 (Hs00863329_g1), NUP153 (Hs01018919_m1), ISG15

(Hs00196051_m1), MX1 (Hs00895608_m1), IFIT1 (Hs00356631_g1), IFI44L

(Hs00915292_m1), RSAD2 (Hs00895608_m1), IFITM1 (Hs00705137_s1), OAS1

(Hs00973635_m1), IFIT2 (Hs00533665_m1), IFIT3 (Hs01922752_s1), IFI44 (Hs00197427_m1),

BST2 (Hs01561315_m1), and ACTB (Hs01060665_g1) and primers/probe from BEI targeting the

SARS-CoV-2 N region using the EXPRESS One-Step Superscript qRT-PCR Kit on an Applied

Biosystems StepOnePlus or QuantStudio 3 instrument. RNA was normalized using an endogenous

18S rRNA primer/probe set (Applied Biosystems).

DNA pull-down assay

483 Nuclear extract from 293T cells expressing ETV7 and treated with IFN-α (100 U/mL, 9 h) was

484 generated using NE-PER extraction reagents (Thermo Scientific). This nuclear extract was

485 incubated with poly(dI-dC) as nonspecific competitor DNA, 10x ligase buffer (Invitrogen), 200

486 mM EDTA, biotinylated DNA bait (either WT or -ETS ISG15 ISRE (*93*)), and 20x excess

487 nonbiotinylated DNA competitor for 30 min at room temperature. This mixture was then incubated

488 with streptavidin-coated magnetic beads (Invitrogen), followed by four washes with TTBS.

489 Proteins bound to the biotinylated DNA bait captured by the streptavidin beads were eluted with

490 protein sample buffer and then detected by Western blot. Input (nuclear extract) was diluted 1:200

491 for loading. No bait (control) contained no biotinylated DNA.

493 *RNA sequencing*

494 293T cells were transfected with ETV7- or control-expressing plasmids and selected using

495 puromycin (20 µg/mL) for 24 h before treatment with IFN-α (100 U/mL). Total RNA was

496 collected at 9 h post-IFN treatment using Monarch Total RNA Miniprep Kits (NEB). RNA was

497 prepped for RNA sequencing submission using the NEBNext Poly(A) mRNA Magnetic Isolation

498 Module (NEB), NEBNext Ultra II RNA Library Prep Kit for Illumina (NEB), and NEBNext

499 Multiplex Oligos for Illumina (NEB). Samples were analyzed on one lane of an Illumina HiSeq

500 4000 using 50 bp single strand reads. Mapping of the raw reads to the human hg19 reference

501 genome was accomplished using a custom application on the Illumina BaseSpace Sequence Hub

502 (*94*). After data normalization, average read values were compared across samples. For

503 comparisons in which some samples had zero reads detected for a specific gene, one read was

504 added to all values in the sample to complete analyses that required non-zero values. The heat map

505 shows genes upregulated 2.5-fold (after normalization) with IFN treatment in the control samples.

506 Values shown are z-scores.

ChIP-qPCR

293T cells were transfected with a FLAG-tagged ETV7-expressing plasmid, ETV7(FLAG), 48 h before treatment with IFN-α (100 U/mL, 9 h). Chromatin immunoprecipitation was performed using the ChIP-IT Express Enzymatic kit according to the manufacturer's directions (Active Motif). DNA was enzymatically sheared for 10 min. ChIP antibodies included mouse IgG (Active Motif), rabbit anti-FLAG (Cell Signaling, 14793S), and mouse anti-FLAG (Sigma, F3165). Quantitative PCR was performed using SYBR Green (Bio-Rad) and primers targeting the Chr12 gene desert (Active Motif) or IFI44L promoter (forward primer = 5' TTTCATGCCTGCCTACATAC 3', reverse primer = 5' ATGCCAACTGCCACTAAC 3') in a region containing two potential ISRE sequences overlapping with ETS sites, and analyzed using the ChIP-IT qPCR Analysis kit (Active Motif).

Viruses

PR8-mNeon was generated via insertion of the mNeon fluorescent gene (*95*) into segment 4 of the virus (*48*). Mal/04-mNeon was generated by inserting the mNeon fluorescent gene (*95*) into segment 4 of the Mal/04 genome (*49*). Cal/09-sfGFP was generated via insertion of the sfGFP gene (*96*) into segment 8 of the virus using the same scheme previously used to insert Cre recombinase (*97*). Sendai-GFP was a gift from Benhur Lee (*98*). For influenza virus infections, cells were either mock- or virus-infected for 1 h and then cultured in OptiMEM supplemented with bovine serum albumin (BSA), penicillin-streptomycin, and 0.2 μg/mL TPCK-treated trypsin protease (Sigma). For Sendai infections, cells were infected for 1 h and then cultured in DMEM. For SARS-CoV-2 infections, cells were washed with PBS before infection with SARS-CoV-2 isolate USA-WA1/2020 from BEI Resources in 2% FBS DMEM infection media for 1 hour. PR8 WT, PR8-mNeon, Cal/09-sfGFP, Sendai-GFP, and SARS-CoV-2 were incubated at 37°C, Mal/04-mNeon was incubated at 33°C.

Viral Growth Assays

Hemagglutination (HA) assays to measure the number of viral particles were performed by diluting influenza infected cell supernatants collected at the indicated time points in cold PBS. An equal amount of chicken blood diluted 1:40 in PBS was mixed with serially diluted virus and incubated at 4°C for 2-3 h before scoring. Infectious viral titers were determined using standard plaque assay procedures on MDCK cells. Infected cell supernatants were collected at 18 h, serially diluted, and used to infect confluent 6-well plates for 1 h before removing the virus and adding the agar overlay. Cells were then incubated at 37°C for 48 h before being fixed in 4% PFA overnight. The 4% PFA was then aspirated, and the agar layer was removed before washing cells with PBS. Serum from WT PR8 infected mice was diluted 1:2,000 in antibody dilution buffer (5% (w/v) non-fat dried milk and 0.05% Tween-20 in PBS) and incubated on cells at 4°C overnight. Cells were then washed twice with PBS and incubated for 1 h with anti-mouse IgG horseradish peroxidase (HRP)-conjugated sheep antibody (GE Healthcare) diluted 1:4,000 in antibody dilution buffer. Assays were then washed twice with PBS and exposed to 0.5 mL of TrueBlue peroxidase substrate (KPL) for 20 min. Plates were then washed with water and dried before plaques were counted.

549
550

551 **Supplementary Materials**
552
553 Fig. S1. CRISPR activation screen hit identification and validation.
554
555 Fig. S2. ETV7 represses IFN-stimulated expression and directly binds an ISG promoter.
556
557 Fig. S3. ETV7 loss increases IFN-stimulated expression.
558
559 Fig. S4. ETV7 loss results in decreased viral titers and increased antiviral gene expression.
560
561 Table S1. Hits from CRISPRa screen for negative regulators of the type I IFN response.
562
563 Table S2. ISRE sequences identified in the literature.
564
565 Table S3. Gene set enrichment analysis of 200 most downregulated genes in ETV7-
566 expressing cells without IFN treatment.
567
568 Table S4. Gene set enrichment analysis of 200 most downregulated genes in ETV7-
569 expressing cells after 9hr IFN treatment.
570
571 Data file S1. CRISPRa screen for negative regulators of the type I IFN response
572 sgRNA sequences, transduction read counts, and screen read counts.
573
574 Data file S2. CRISPRa screen for negative regulators of the type I IFN response
575 MAGeCK z-scores and potential hit cutoffs.
576
577 Data file S3. RNA sequencing with ETV7 overexpression and IFN treatment
578 reads and results.
579

References

1.   W. M. Schneider, M. D. Chevillotte, C. M. Rice, Interferon-Stimulated Genes: A
Complex Web of Host Defenses. *Annu. Rev. Immunol.* **32**, 513–545 (2014).

2.   J. W. Schoggins, Interferon-Stimulated Genes: What Do They All Do? *Annu. Rev. Virol.*
6, 567–584 (2019).

3.   J. W. Schoggins, S. J. Wilson, M. Panis, M. Y. Murphy, C. T. Jones, P. Bieniasz, C. M.
Rice, A diverse range of gene products are effectors of the type I interferon antiviral
response. *Nature.* **472**, 481–485 (2011).

4.   J. W. Schoggins, D. A. MacDuff, N. Imanaka, M. D. Gainey, B. Shrestha, J. L. Eitson, K.
B. Mar, R. B. Richardson, A. V. Ratushny, V. Litvak, R. Dabelic, B. Manicassamy, J. D.
Aitchison, A. Aderem, R. M. Elliott, A. García-Sastre, V. Racaniello, E. J. Snijder, W. M.
Yokoyama, M. S. Diamond, H. W. Virgin, C. M. Rice, Pan-viral specificity of IFN-
induced genes reveals new roles for cGAS in innate immunity. *Nature.* **505**, 691–695
(2014).

5.   M. Kane, T. M. Zang, S. J. Rihn, C. M. Rice, S. J. Wilson, P. D. Bieniasz, Identification
of Interferon-Stimulated Genes with Antiretroviral Activity. *Cell Host Microbe.* **20**, 392–
405 (2016).

6.   E. V. Mesev, R. A. LeDesma, A. Ploss, Decoding type I and III interferon signalling
during viral infection. *Nat. Microbiol.* **4**, 914–924 (2019).

7.   B. X. Wang, E. N. Fish, The yin and yang of viruses and interferons. *Trends Immunol.* **33**
(2012), doi:10.1016/j.it.2012.01.004.

8.   G. R. Stark, J. E. Darnell, The JAK-STAT Pathway at Twenty. *Immunity.* **36**, 503–514
(2012).

9.   K. Honda, H. Yanai, A. Takaoka, T. Taniguchi, Regulation of the type I IFN induction: a
current view. *Int. Immunol.* **17**, 1367–1378 (2005).

10.   D. E. Levy, D. S. Kessler, R. Pine, N. Reich, J. E. Darnell, Interferon-induced nuclear
factors that bind a shared promoter element correlate with positive and negative
transcriptional control. *Genes Dev.* **2**, 383–93 (1988).

11.   N. P. D. Liau, A. Laktyushin, I. S. Lucet, J. M. Murphy, S. Yao, E. Whitlock, K.
Callaghan, N. A. Nicola, N. J. Kershaw, J. J. Babon, The molecular basis of JAK/STAT
inhibition by SOCS1. *Nat. Commun.* **9**, 1558 (2018).

12.   M. P. Malakhov, O. A. Malakhova, K. Il Kim, K. J. Ritchie, D.-E. Zhang, UBP43
(USP18) specifically removes ISG15 from conjugated proteins. *J. Biol. Chem.* **277**, 9976–
81 (2002).

13.   K.-I. Arimoto, S. Miyauchi, S. A. Stoner, J.-B. Fan, D.-E. Zhang, Negative regulation of
type I IFN signaling. *J. Leukoc. Biol.* **103**, 1099–1116 (2018).

14.   G. Randall, L. Chen, M. Panis, A. K. Fischer, B. D. Lindenbach, J. Sun, J. Heathcote, C.
M. Rice, A. M. Edwards, I. D. McGilvray, Silencing of USP18 Potentiates the Antiviral
Activity of Interferon Against Hepatitis C Virus Infection. *Gastroenterology.* **131**, 1584–
1591 (2006).

15.   K. G. Frey, C. M. I. Ahmed, R. Dabelic, L. D. Jager, E. N. Noon-Song, S. M. Haider, H.
M. Johnson, N. J. Bigley, HSV-1-induced SOCS-1 expression in keratinocytes: use of a
SOCS-1 antagonist to block a novel mechanism of viral immune evasion. *J. Immunol.*
183, 1253–62 (2009).

16.   C. J. Carbone, H. Zheng, S. Bhattacharya, J. R. Lewis, A. M. Reiter, P. Henthorn, Z.-Y.
Zhang, D. P. Baker, R. Ukkiramapandian, K. K. Bence, S. Y. Fuchs, Protein tyrosine
phosphatase 1B is a key regulator of IFNAR1 endocytosis and a target for antiviral
therapies. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 19226–31 (2012).

17. M. Sa Ribero, N. Jouvenet, M. Dreux, S. Nisole, Interplay between SARS-CoV-2 and the
type I interferon response. *PLOS Pathog.* **16**, e1008737 (2020).

18. J. R. Teijaro, S. Studer, N. Leaf, W. B. Kiosses, N. Nguyen, K. Matsuki, H. Negishi, T.
Taniguchi, M. B. A. Oldstone, H. Rosen, S1PR1-mediated IFNAR1 degradation
modulates plasmacytoid dendritic cell interferon-α autoamplification. *Proc. Natl. Acad.*
Sci. U. S. A. **113**, 1351–6 (2016).

19. K. B. Walsh, J. R. Teijaro, P. R. Wilker, A. Jatzek, D. M. Fremgen, S. C. Das, T.
Watanabe, M. Hatta, K. Shinya, M. Suresh, Y. Kawaoka, H. Rosen, M. B. A. Oldstone,
Suppression of cytokine storm with a sphingosine analog provides protection against
pathogenic influenza virus. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 12018–23 (2011).

20. F. Naz, M. Arish, Battling COVID-19 Pandemic: Sphingosine-1-Phosphate Analogs as an
Adjunctive Therapy? *Front. Immunol.* **11**, 1102 (2020).

21. P. J. Hertzog, B. R. G. Williams, Fine tuning type I interferon responses. *Cytokine Growth*
Factor Rev. **24**, 217–225 (2013).

22. J. P. B. Viola, R. P. Donnelly, H. M. Johnson, H. A. R. Bluyssen, A. Michalska, K.
Blaszczyk, J. Wesoly, A Positive Feedback Amplifier Circuit That Regulates Interferon
(IFN)-Stimulated Gene Expression and Controls Type I and Type II IFN Responses. **9**, 28
(2018).

23. N. Tanaka, T. Kawakami, T. Taniguchi, Recognition DNA sequences of interferon
regulatory factor 1 (IRF-1) and IRF-2, regulators of cell growth and the interferon system.
Mol. Cell. Biol. **13**, 4531–8 (1993).

24. M. Sato, N. Hata, M. Asagiri, T. Nakaya, T. Taniguchi, N. Tanaka, Positive feedback
regulation of type I *IFN* genes by the IFN-inducible transcription factor IRF-7. *FEBS Lett.*
441, 106–110 (1998).

25. L. L. Seifert, C. Si, D. Saha, M. Sadic, M. de Vries, S. Ballentine, A. Briley, G. Wang, A.
M. Valero-Jimenez, A. Mohamed, U. Schaefer, H. M. Moulton, A. García-Sastre, S.
Tripathi, B. R. Rosenberg, M. Dittmann, The ETS transcription factor ELF1 regulates a
broadly antiviral program distinct from the type I interferon response. *PLOS Pathog.* **15**,
e1007634 (2019).

26. P. Hubel, C. Urban, V. Bergant, W. M. Schneider, B. Knauer, A. Stukalov, P. Scaturro, A.
Mann, L. Brunotte, H. H. Hoffmann, J. W. Schoggins, M. Schwemmle, M. Mann, C. M.
Rice, A. Pichlmair, A protein-interaction network of interferon-stimulated genes extends
the innate immune system landscape. *Nat. Immunol.* **20**, 493–502 (2019).

27. X. Li, X. Zhao, Y. Fang, X. Jiang, T. Duong, C. Fan, C.-C. Huang, S. R. Kain, Generation
of Destabilized Green Fluorescent Protein as a Transcription Reporter. *J. Biol. Chem.* **273**,
34970–34975 (1998).

28. S. Konermann, M. D. Brigham, A. E. Trevino, J. Joung, O. O. Abudayyeh, C. Barcena, P.
D. Hsu, N. Habib, J. S. Gootenberg, H. Nishimasu, O. Nureki, F. Zhang, Genome-scale
transcriptional activation by an engineered CRISPR-Cas9 complex. *Nature.* **517**, 583–8
(2015).

29. W. Li, H. Xu, T. Xiao, L. Cong, M. I. Love, F. Zhang, R. A. Irizarry, J. S. Liu, M. Brown,
X. S. Liu, MAGeCK enables robust identification of essential genes from genome-scale
CRISPR/Cas9 knockout screens. *Genome Biol.* **15**, 554 (2014).

30. I. Rusinova, S. Forster, S. Yu, A. Kannan, M. Masse, H. Cumming, R. Chapman, P. J.
Hertzog, INTERFEROME v2.0: an updated database of annotated interferon-regulated
genes. *Nucleic Acids Res.* **41**, D1040–D1046 (2012).

31. A.-P. Han, C. Yu, L. Lu, Y. Fujiwara, C. Browne, G. Chin, M. Fleming, P. Leboulch, S.
H. Orkin, J. Chen, Heme-regulated eIF2alpha kinase (HRI) is required for translational
regulation and survival of erythroid precursors in iron deficiency. *EMBO J.* **20**, 6909–

6918 (2001).

32.   A. Marg, Y. Shan, T. Meyer, T. Meissner, M. Brandenburg, U. Vinkemeier,
Nucleocytoplasmic shuttling by nucleoporins Nup153 and Nup214 and CRM1-dependent
nuclear export control the subcellular distribution of latent Stat1. *J. Cell Biol.* **165**, 823–33
(2004).

33.   M. D. Potter, A. Buijs, B. Kreider, L. van Rompaey, G. C. Grosveld, Identification and
characterization of a new human ETS-family transcription factor, TEL2, that is expressed
in hematopoietic tissues and can associate with TEL1/ETV6. *Blood.* **95**, 3341–8 (2000).

34.   H. Poirel, R. G. Lopez, V. Lacronique, V. Della Valle, M. Mauchauffé, R. Berger, J.
Ghysdael, O. A. Bernard, Characterization of a novel ETS gene, TELB, encoding a
protein structurally and functionaly related to TEL. *Oncogene.* **19**, 4802–4806 (2000).

35.   F. C. Harwood, R. I. Klein Geltink, B. P. O'Hara, M. Cardone, L. Janke, D. Finkelstein, I.
Entin, L. Paul, P. J. Houghton, G. C. Grosveld, ETV7 is an essential component of a
rapamycin-insensitive mTOR complex in cancer. *Sci. Adv.* **4**, eaar3938 (2018).

36.   H. Wei, G. Badis, M. F. Berger, T. Kivioja, K. Palin, M. Enge, M. Bonke, A. Jolma, M.
Varjosalo, A. R. Gehrke, J. Yan, S. Talukder, M. Turunen, M. Taipale, H. G.
Stunnenberg, E. Ukkonen, T. R. Hughes, M. L. Bulyk, J. Taipale, Genome-wide analysis
of ETS-family DNA-binding in vitro and in vivo. *EMBO J.* **29**, 2147–2160 (2010).

37.   J. Reimand, T. Arak, P. Adler, L. Kolberg, S. Reisberg, H. Peterson, J. Vilo, g:Profiler-a
web server for functional interpretation of gene lists (2016 update). *Nucleic Acids Res.* **44**,
W83–W89 (2016).

38.   N. E. Sanjana, O. Shalem, F. Zhang, Improved vectors and genome-wide libraries for
CRISPR screening. *Nat. Methods.* **11**, 783–784 (2014).

39.   D. C. Busse, D. Habgood-Coote, S. Clare, C. Brandt, I. Bassano, M. Kaforou, M. Levin,
J.-F. Eleouet, P. Kellam, J. S. Tregoning, Interferon-induced Protein-44 and Interferon-
induced Protein 44-like restrict 1 replication of Respiratory Syncytial Virus 2 Downloaded
from. *J. Virol* (2020), doi:10.1128/JVI.00297-20.

40.   D. J. Morales, D. J. Lenschow, The antiviral activities of ISG15. *J. Mol. Biol.* **425**, 4995–
5008 (2013).

41.   J. W. Schoggins, Recent advances in antiviral interferon-stimulated gene biology.
F1000Research. **7**, 309 (2018).

42.   A. García-Sastre, Induction and evasion of type I interferon responses by influenza
viruses. *Virus Res.* **162**, 12–18 (2011).

43.   A. L. Brass, I.-C. Huang, Y. Benita, S. P. John, M. N. Krishnan, E. M. Feeley, B. J. Ryan,
J. L. Weyer, L. van der Weyden, E. Fikrig, D. J. Adams, R. J. Xavier, M. Farzan, S. J.
Elledge, The IFITM Proteins Mediate Cellular Resistance to Influenza A H1N1 Virus,
West Nile Virus, and Dengue Virus. *Cell.* **139**, 1243–1254 (2009).

44.   Y. Li, S. Banerjee, Y. Wang, S. A. Goldstein, B. Dong, C. Gaughan, R. H. Silverman, S.
R. Weiss, Activation of RNase L is dependent on OAS3 expression during infection with
diverse human viruses. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 2241–6 (2016).

45.   A. Pichlmair, C. Lassnig, C.-A. Eberle, M. W. Górna, C. L. Baumann, T. R. Burkard, T.
Bürckstümmer, A. Stefanovic, S. Krieger, K. L. Bennett, T. Rülicke, F. Weber, J. Colinge,
M. Müller, G. Superti-Furga, IFIT1 is an antiviral protein that recognizes 5′-triphosphate
RNA. *Nat. Immunol.* **12**, 624–630 (2011).

46.   M. A. Yondola, F. Fernandes, A. Belicha-Villanueva, M. Uccelini, Q. Gao, C. Carter, P.
Palese, Budding capability of the influenza virus neuraminidase can be modulated by
tetherin. *J. Virol.* **85**, 2480–91 (2011).

47.   X. Wang, E. R. Hinson, P. Cresswell, The Interferon-Inducible Protein Viperin Inhibits
Influenza Virus Release by Perturbing Lipid Rafts. *Cell Host Microbe.* **2**, 96–105 (2007).

48.   A. T. Harding, B. E. Heaton, R. E. Dumm, N. S. Heaton, Rationally Designed Influenza
Virus Vaccines That Are Antigenically Stable during Growth in Eggs. *MBio*. **8** (2017),
doi:10.1128/mBio.00669-17.

49.   R. E. Dumm, J. K. Fiege, B. M. Waring, C. T. Kuo, R. A. Langlois, N. S. Heaton, Non-
lytic clearance of influenza B virus from infected cells preserves epithelial barrier
function. *Nat. Commun.* **10**, 779 (2019).

50.   J. R. Bedsaul, L. A. Zaritsky, K. C. Zoon, Type I interferon-mediated induction of
antiviral genes and proteins fails to protect cells from the cytopathic effects of sendai virus
infection. *J. Interf. Cytokine Res.* **36**, 652–665 (2016).

51.   J. M. Perreira, C. R. Chin, E. M. Feeley, A. L. Brass, IFITMs restrict the replication of
multiple pathogenic viruses. *J. Mol. Biol.* **425** (2013), pp. 4937–4955.

52.   D. F. Young, J. Andrejeva, X. Li, F. Inesta-Vaquera, C. Dong, V. H. Cowling, S.
Goodbourn, R. E. Randall, Human IFIT1 Inhibits mRNA Translation of Rubulaviruses but
Not Other Members of the Paramyxoviridae Family. *J. Virol.* **90**, 9446–9456 (2016).

53.   C. Bampi, L. Rasga, L. Roux, Antagonism to human BST-2/tetherin by Sendai virus
glycoproteins. *J. Gen. Virol.* **94**, 1211–9 (2013).

54.   M. P. Manns, H. Wedemeyer, M. Cornberg, Treating viral hepatitis C: efficacy, side
effects, and complications. *Gut*. **55**, 1350–9 (2006).

55.   E. Haasbach, K. Droebner, A. B. Vogel, O. Planz, Low-dose interferon Type I treatment is
effective against H5N1 and swine-origin H1N1 influenza A viruses in vitro and in vivo. *J.*
Interferon Cytokine Res. **31**, 515–25 (2011).

56.   M. R. Loutfy, L. M. Blatt, K. A. Siminovitch, S. Ward, B. Wolff, H. Lho, D. H. Pham, H.
Deif, E. A. LaMere, M. Chang, K. C. Kain, G. A. Farcas, P. Ferguson, M. Latchford, G.
Levy, J. W. Dennis, E. K. Y. Lai, E. N. Fish, Interferon Alfacon-1 Plus Corticosteroids in
Severe Acute Respiratory Syndrome. *JAMA*. **290**, 3222 (2003).

57.   M. K. Konde, D. P. Baker, F. A. Traore, M. S. Sow, A. Camara, A. A. Barry, D. Mara, A.
Barry, M. Cone, I. Kaba, A. A. Richard, A. H. Beavogui, S. Günther, on behalf of E. M.
L. European Mobile Laboratory Consortium, M. Pintilie, E. N. Fish, Interferon β-1a for
the treatment of Ebola virus disease: A historically controlled, single-arm proof-of-
concept trial. *PLoS One*. **12**, e0169255 (2017).

58.   Q. Zhou, V. Chen, C. P. Shannon, X.-S. Wei, X. Xiang, X. Wang, Z.-H. Wang, S. J.
Tebbutt, T. R. Kollmann, E. N. Fish, Interferon-α2b Treatment for COVID-19. *Front.*
Immunol. **11**, 1061 (2020).

59.   E. Davoudi-Monfared, H. Rahmani, H. Khalili, M. Hajiabdolbaghi, M. Salehi, L.
Abbasian, H. Kazemzadeh, M. S. Yekaninejad, Efficacy and safety of interferon β-1a in
treatment of severe COVID-19: A randomized clinical trial. *Antimicrob. Agents*
Chemother. (2020), doi:10.1128/AAC.01061-20.

60.   I. F.-N. Hung, K.-C. Lung, E. Y.-K. Tso, R. Liu, T. W.-H. Chung, M.-Y. Chu, Y.-Y. Ng,
J. Lo, J. Chan, A. R. Tam, H.-P. Shum, V. Chan, A. K.-L. Wu, K.-M. Sin, W.-S. Leung,
W.-L. Law, D. C. Lung, S. Sin, P. Yeung, C. C.-Y. Yip, R. R. Zhang, A. Y.-F. Fung, E.
Y.-W. Yan, K.-H. Leung, J. D. Ip, A. W.-H. Chu, W.-M. Chan, A. C.-K. Ng, R. Lee, K.
Fung, A. Yeung, T.-C. Wu, J. W.-M. Chan, W.-W. Yan, W.-M. Chan, J. F.-W. Chan, A.
K.-W. Lie, O. T.-Y. Tsang, V. C.-C. Cheng, T.-L. Que, C.-S. Lau, K.-H. Chan, K. K.-W.
To, K.-Y. Yuen, Triple combination of interferon beta-1b, lopinavir-ritonavir, and
ribavirin in the treatment of patients admitted to hospital with COVID-19: an open-label,
randomised, phase 2 trial. *Lancet (London, England)*. **395**, 1695–1704 (2020).

61.   D. Blanco-Melo, B. E. Nilsson-Payant, W. C. Liu, S. Uhl, D. Hoagland, R. Møller, T. X.
Jordan, K. Oishi, M. Panis, D. Sachs, T. T. Wang, R. E. Schwartz, J. K. Lim, R. A.
Albrecht, B. R. tenOever, Imbalanced Host Response to SARS-CoV-2 Drives

774   Development of COVID-19. *Cell*. **181**, 1036-1045.e9 (2020).

775 62. H. Chu, J. F.-W. Chan, Y. Wang, T. T.-T. Yuen, Y. Chai, Y. Hou, H. Shuai, D. Yang, B.
776   Hu, X. Huang, X. Zhang, J.-P. Cai, J. Zhou, S. Yuan, K.-H. Kok, K. K.-W. To, I. H.-Y.
777   Chan, A. J. Zhang, K.-Y. Sit, W.-K. Au, K.-Y. Yuen, Comparative Replication and
778   Immune Activation Profiles of SARS-CoV-2 and SARS-CoV in Human Lungs: An Ex
779   Vivo Study With Implications for the Pathogenesis of COVID-19. *Clin. Infect. Dis.*
780   (2020), doi:10.1093/cid/ciaa410.

781 63. J. Hadjadj, N. Yatim, L. Barnabei, A. Corneau, J. Boussier, N. Smith, H. Péré, B. Charbit,
782   V. Bondet, C. Chenevier-Gobeaux, P. Breillat, N. Carlier, R. Gauzit, C. Morbieu, F. Pène,
783   N. Marin, N. Roche, T.-A. Szwebel, S. H. Merkling, J.-M. Treluyer, D. Veyer, L.
784   Mouthon, C. Blanc, P.-L. Tharaux, F. Rozenberg, A. Fischer, D. Duffy, F. Rieux-Laucat,
785   S. Kernéis, B. Terrier, Impaired type I interferon activity and inflammatory responses in
786   severe COVID-19 patients. *Science* (2020), doi:10.1126/science.abc6027.

787 64. A. Vanderheiden, P. Ralfs, T. Chirkova, A. A. Upadhyay, M. G. Zimmerman, S. Bedoya,
788   H. Aoued, G. M. Tharp, K. L. Pellegrini, C. Manfredi, E. Sorscher, B. Mainou, J. E.
789   Kohlmeier, A. C. Lowen, P.-Y. Shi, V. D. Menachery, L. J. Anderson, A. Grakoui, S. E.
790   Bosinger, M. S. Suthar, Type I and Type III IFN Restrict SARS-CoV-2 Infection of
791   Human Airway Epithelial 1 Cultures 2 3 Downloaded from. *J. Virol* (2020),
792   doi:10.1128/JVI.00985-20.

793 65. K. G. Lokugamage, A. Hage, M. de Vries, A. M. Valero-Jimenez, C. Schindewolf, M.
794   Dittmann, R. Rajsbaum, V. D. Menachery, *bioRxiv*, in press,
795   doi:10.1101/2020.03.07.982264.

796 66. E. Mantlo, N. Bukreyeva, J. Maruyama, S. Paessler, C. Huang, Antiviral activities of type
797   I interferons to SARS-CoV-2 infection. *Antiviral Res.* **179**, 104811 (2020).

798 67. S. Pfaender, K. B. Mar, E. Michailidis, A. Kratzel, I. N. Boys, P. V'kovski, W. Fan, J. N.
799   Kelly, D. Hirt, N. Ebert, H. Stalder, H. Kleine-Weber, M. Hoffmann, H. Heinrich
800   Hoffmann, M. Saeed, R. Dijkman, E. Steinmann, M. Wight-Carter, M. B. McDougal, N.
801   W. Hanners, S. Pöhlmann, T. Gallagher, D. Todt, G. Zimmer, C. M. Rice, J. W.
802   Schoggins, V. Thiel, LY6E impairs coronavirus fusion and confers immune control of
803   viral disease. *Nat. Microbiol.*, 1–10 (2020).

804 68. X. Zhao, S. Zheng, D. Chen, M. Zheng, X. Li, G. Li, H. Lin, J. Chang, H. Zeng, J.-T. Guo,
805   LY6E Restricts the Entry of Human Coronaviruses, Including the Currently Pandemic
806   SARS-CoV-2. *J. Virol.* (2020), doi:10.1128/JVI.00562-20.

807 69. R. Li, H. Pei, D. K. Watson, Regulation of Ets function by protein–protein interactions.
808   *Oncogene*. **19**, 6514–6523 (2000).

809 70. A. D. Sharrocks, The ETS-domain transcription factor family. *Nat. Rev. Mol. Cell Biol.* **2**,
810   827–837 (2001).

811 71. D. Meraro, M. Gleit-Kielmanowicz, H. Hauser, B.-Z. Levi, IFN-Stimulated Gene 15 Is
812   Synergistically Activated Through Interactions Between the Myelocyte/Lymphocyte-
813   Specific Transcription Factors, PU.1, IFN Regulatory Factor-8/IFN Consensus Sequence
814   Binding Protein, and IFN Regulatory Factor-4: Characterization o. *J. Immunol.* **168**,
815   6224–6231 (2002).

816 72. T. Taniguchi, K. Ogasawara, A. Takaoka, N. Tanaka, IRF Family of Transcription Factors
817   as Regulators of Host Defense. *Annu. Rev. Immunol.* **19**, 623–655 (2001).

818 73. A. L. Brass, E. Kehrli, C. F. Eisenbeis, U. Storb, H. Singh, Pip, a lymphoid-restricted IRF,
819   contains a regulatory domain that is important for autoinhibition and ternary complex
820   formation with the Ets factor PU.1. *Genes Dev.* **10**, 2335–2347 (1996).

821 74. W. Wang, L. Xu, J. Su, M. P. Peppelenbosch, Q. Pan, Transcriptional Regulation of
822   Antiviral Interferon-Stimulated Genes. *Trends Microbiol.* **25**, 573–584 (2017).

75. T.-Y. Liu, T. Burke, L. P. Park, C. W. Woods, A. K. Zaas, G. S. Ginsburg, A. O. Hero, An
individualized predictor of health and disease using paired reference and target samples.
BMC Bioinformatics. **17**, 47 (2016).

76. P. C. De La Cruz-Rivera, M. Kanchwala, H. Liang, A. Kumar, L.-F. Wang, C. Xing, J. W.
Schoggins, The IFN Response in Bats Displays Distinctive IFN-Stimulated Gene
Expression Kinetics with Atypical RNASEL Induction. *J. Immunol.* **200**, 209–217 (2018).

77. K. Haq, J. T. Brisbin, N. Thanthrige-Don, M. Heidari, S. Sharif, Transcriptome and
proteome profiling of host responses to Marek's disease virus in chickens. *Vet. Immunol.*
Immunopathol. **138**, 292–302 (2010).

78. H. Kawagoe, M. Potter, J. Ellis, G. C. Grosveld, TEL2, an ETS factor expressed in human
leukemia, regulates monocytic differentiation of U937 cells and blocks the inhibitory
effect of TEL1 on Ras-induced cellular transformation. *Cancer Res.* **64**, 6091–6100
(2004).

79. P. Rasighaemi, A. C. Ward, ETV6 and ETV7: Siblings in hematopoiesis and its disruption
in disease. *Crit. Rev. Oncol. Hematol.* **116**, 106–115 (2017).

80. J. A. Nick, S. M. Caceres, J. E. Kret, K. R. Poch, M. Strand, A. V. Faino, D. P. Nichols,
M. T. Saavedra, J. L. Taylor-Cousar, M. W. Geraci, E. L. Burnham, M. B. Fessler, B. T.
Suratt, E. Abraham, M. Moss, K. C. Malcolm, Extremes of Interferon-Stimulated Gene
Expression Associate with Worse Outcomes in the Acute Respiratory Distress Syndrome.
PLoS One. **11**, e0162490 (2016).

81. Y. Muramoto, J. E. Shoemaker, M. Q. Le, Y. Itoh, D. Tamura, Y. Sakai-Tagawa, H. Imai,
R. Uraki, R. Takano, E. Kawakami, M. Ito, K. Okamoto, H. Ishigaki, H. Mimuro, C.
Sasakawa, Y. Matsuoka, T. Noda, S. Fukuyama, K. Ogasawara, H. Kitano, Y. Kawaoka,
Disease severity is associated with differential gene expression at the early and late phases
of infection in nonhuman primates infected with different H5N1 highly pathogenic avian
influenza viruses. *J. Virol.* **88**, 8981–97 (2014).

82. S. Davidson, M. K. Maini, A. Wack, Disease-promoting effects of type I interferons in
viral, bacterial, and coinfections. *J. Interferon Cytokine Res.* **35**, 252–64 (2015).

83. M. E. C. Meuwissen, R. Schot, S. Buta, G. Oudesluijs, S. Tinschert, S. D. Speer, Z. Li, L.
van Unen, D. Heijsman, T. Goldmann, M. H. Lequin, J. M. Kros, W. Stam, M. Hermann,
R. Willemsen, R. W. W. Brouwer, W. F. J. Van IJcken, M. Martin-Fernandez, I. de Coo,
J. Dudink, F. A. T. de Vries, A. Bertoli Avella, M. Prinz, Y. J. Crow, F. W. Verheijen, S.
Pellegrini, D. Bogunovic, G. M. S. Mancini, Human USP18 deficiency underlies type 1
interferonopathy leading to severe pseudo-TORCH syndrome. *J. Exp. Med.* **213**, 1163–74
(2016).

84. J.-C. Marine, D. J. Topham, C. McKay, D. Wang, E. Parganas, D. Stravopodis, A.
Yoshimura, J. N. Ihle, SOCS1 Deficiency Causes a Lymphocyte-Dependent Perinatal
Lethality. *Cell*. **98**, 609–616 (1999).

85. W. S. Alexander, R. Starr, J. E. Fenner, C. L. Scott, E. Handman, N. S. Sprigg, J. E.
Corbin, A. L. Cornish, R. Darwiche, C. M. Owczarek, T. W. . Kay, N. A. Nicola, P. J.
Hertzog, D. Metcalf, D. J. Hilton, SOCS1 Is a Critical Inhibitor of Interferon γ Signaling
and Prevents the Potentially Fatal Neonatal Actions of this Cytokine. *Cell*. **98**, 597–608
(1999).

86. B. A. Croker, D. L. Krebs, J.-G. Zhang, S. Wormald, T. A. Willson, E. G. Stanley, L.
Robb, C. J. Greenhalgh, I. Förster, B. E. Clausen, N. A. Nicola, D. Metcalf, D. J. Hilton,
A. W. Roberts, W. S. Alexander, SOCS3 negatively regulates IL-6 signaling in vivo. *Nat.*
Immunol. **4**, 540–545 (2003).

87. K.-P. Knobeloch, O. Utermöhlen, A. Kisser, M. Prinz, I. Horak, Reexamination of the
Role of Ubiquitin-Like Modifier ISG15 in the Phenotype of UBP43-Deficient Mice

Downloaded from. *Mol. Cell. Biol.* **25**, 11030–11034 (2005).

88.   Y. Okada, D. Wu, G. Trynka, T. Raj, C. Terao, K. Ikari, Y. Kochi, K. Ohmura, A. Suzuki,
S. Yoshida, R. R. Graham, A. Manoharan, W. Ortmann, T. Bhangale, J. C. Denny, R. J.
Carroll, A. E. Eyler, J. D. Greenberg, J. M. Kremer, D. A. Pappas, L. Jiang, J. Yin, L. Ye,
D.-F. Su, J. Yang, G. Xie, E. Keystone, H.-J. Westra, T. Esko, A. Metspalu, X. Zhou, N.
Gupta, D. Mirel, E. A. Stahl, D. Diogo, J. Cui, K. Liao, M. H. Guo, K. Myouzen, T.
Kawaguchi, M. J. H. Coenen, P. L. C. M. van Riel, M. A. F. J. van de Laar, H.-J.
Guchelaar, T. W. J. Huizinga, P. Dieudé, X. Mariette, S. Louis Bridges Jr, A. Zhernakova,
R. E. M. Toes, P. P. Tak, C. Miceli-Richard, S.-Y. Bang, H.-S. Lee, J. Martin, M. A.
Gonzalez-Gay, L. Rodriguez-Rodriguez, S. Rantapää-Dahlqvist, L. Ärlestig, H. K. Choi,
Y. Kamatani, P. Galan, M. Lathrop, S. Eyre, J. Bowes, A. Barton, N. de Vries, L. W.
Moreland, L. A. Criswell, E. W. Karlson, A. Taniguchi, R. Yamada, M. Kubo, J. S. Liu,
S.-C. Bae, J. Worthington, L. Padyukov, L. Klareskog, P. K. Gregersen, S. Raychaudhuri,
B. E. Stranger, P. L. De Jager, L. Franke, P. M. Visscher, M. A. Brown, H. Yamanaka, T.
Mimori, A. Takahashi, H. Xu, T. W. Behrens, K. A. Siminovitch, S. Momohara, F.
Matsuda, K. Yamamoto, R. M. Plenge, K. Yamamoto, R. M. Plenge, Genetics of
rheumatoid arthritis contributes to biology and drug discovery. *Nature*. **506**, 376–381
(2014).

89.   T. James, M. Lindén, H. Morikawa, S. J. Fernandes, S. Ruhrmann, M. Huss, M. Brandi, F.
Piehl, M. Jagodic, J. Tegnér, M. Khademi, T. Olsson, D. Gomez-Cabrero, I. Kockum,
Impact of genetic risk loci for multiple sclerosis on expression of proximal genes in
patients. *Hum. Mol. Genet.* **27**, 912–928 (2018).

90.   J. E. Castañeda-Delgado, Y. Bastián-Hernandez, N. Macias-Segura, D. Santiago-Algarra,
J. D. Castillo-Ortiz, A. L. Alemán-Navarro, P. Martínez-Tejada, L. Enciso-Moreno, Y.
Garcia-De Lira, D. Olguín-Calderón, L. A. Trouw, C. Ramos-Remus, J. A. Enciso-
Moreno, Type I Interferon Gene Response Is Increased in Early and Established
Rheumatoid Arthritis and Correlates with Autoantibody Production. *Front. Immunol.* **8**,
285 (2017).

91.   A. Hundeshagen, M. Hecker, B. K. Paap, C. Angerstein, O. Kandulski, C. Fatum, C.
Hartmann, D. Koczan, H.-J. Thiesen, U. Klaus Zettl, Elevated type I interferon-like
activity in a subset of multiple sclerosis patients: molecular basis and clinical relevance. *J.*
Neuroinflammation. **9**, 574 (2012).

92.   B. E. Heaton, E. M. Kennedy, R. E. Dumm, A. T. Harding, M. T. Sacco, D. Sachs, N. S.
Heaton, A CRISPR Activation Screen Identifies a Pan-avian Influenza Virus Inhibitory
Host Factor. *Cell Rep.* **20**, 1503–1512 (2017).

93.   S. Schmid, M. Mordstein, G. Kochs, A. García-Sastre, B. R. TenOever, Transcription
factor redundancy ensures induction of the antiviral state. *J. Biol. Chem.* **285**, 42013–
42022 (2010).

94.   N. S. Heaton, R. A. Langlois, D. Sachs, J. K. Lim, P. Palese, R. Benjamin, Long-term
survival of influenza virus infected club cells drives immunopathology. **211**, 1707–1714
(2014).

95.   N. C. Shaner, G. G. Lambert, A. Chammas, Y. Ni, P. J. Cranfill, M. A. Baird, B. R. Sell,
J. R. Allen, R. N. Day, M. Israelsson, M. W. Davidson, J. Wang, A bright monomeric
green fluorescent protein derived from Branchiostoma lanceolatum. *Nat. Methods*. **10**,
407–9 (2013).

96.   J.-D. Pédelacq, S. Cabantous, T. Tran, T. C. Terwilliger, G. S. Waldo, Engineering and
characterization of a superfolder green fluorescent protein. *Nat. Biotechnol.* **24**, 79–88
(2006).

97.   B. S. Chambers, B. E. Heaton, K. Rausch, R. E. Dumm, J. R. Hamilton, S. Cherry, N. S.

Heaton, DNA mismatch repair is required for the host innate response and controls
cellular fate after influenza virus infection. *Nat. Microbiol.* **4**, 1964–1977 (2019).

98.   S. M. Beaty, A. Park, S. T. Won, P. Hong, M. Lyons, F. Vigant, A. N. Freiberg, B. R.
tenOever, W. P. Duprex, B. Lee, Efficient and Robust Paramyxoviridae Reverse Genetics
Systems. *mSphere.* **2** (2017), doi:10.1128/mSphere.00376-16.

99.   C. T. Workman, Y. Yin, D. L. Corcoran, T. Ideker, G. D. Stormo, P. V. Benos,
enoLOGOS: a versatile web tool for energy normalized sequence logos. *Nucleic Acids*
Res. **33**, W389–W392 (2005).

100. S. Babicki, D. Arndt, A. Marcu, Y. Liang, J. R. Grant, A. Maciejewski, D. S. Wishart,
Heatmapper: web-enabled heat mapping for all. *Nucleic Acids Res.* **44**, W147–W153
(2016).

101. H. A. R. Bluyssen, R. J. Vlietstra, P. W. Faber, E. M. E. Smit, A. Hagemeijer, J. Trapman,
Structure, Chromosome Localization, and Regulation of Expression of the Interferon-
Regulated Mouse Ifi54/Ifi56 Gene Family. *Genomics.* **24**, 137–148 (1994).

102. T. Ohtomo, Y. Sugamata, Y. Ozaki, K. Ono, Y. Yoshimura, S. Kawai, Y. Koishihara, S.
Ozaki, M. Kosaka, T. Hirano, M. Tsuchiya, Molecular Cloning and Characterization of a
Surface Antigen Preferentially Overexpressed on Multiple Myeloma Cells. *Biochem.*
Biophys. Res. Commun. **258**, 583–591 (1999).

103. W. Yang, J. Tan, R. Liu, X. Cui, Q. Ma, Y. Geng, W. Qiao, Interferon-γ Upregulates
Expression of IFP35 Gene in HeLa Cells via Interferon Regulatory Factor-1. *PLoS One.* **7**
(2012), doi:10.1371/JOURNAL.PONE.0050932.

104. M. G. Wathelet, C. H. Lin, B. S. Parekh, L. V Ronco, P. M. Howley, T. Maniatis, Virus
infection induces the assembly of coordinately activated transcription factors on the IFN-
beta enhancer in vivo. *Mol. Cell.* **1**, 507–18 (1998).

105. B. Testoni, V. Schinzari, F. Guerrieri, S. Gerbal-Chaloin, G. Blandino, M. Levrero, p53-
paralog DNp73 oncogene is repressed by IFNα/STAT2 through the recruitment of the
Ezh2 polycomb group transcriptional repressor. *Oncogene.* **30**, 2670–8 (2011).

106. M. N. Rutherford, G. E. Hannigan, B. R. Williams, Interferon-induced binding of nuclear
factors to promoter elements of the 2-5A synthetase gene. *EMBO J.* **7**, 751–9 (1988).

107. Q. Wang, G. Floyd-Smith, The p69/71 2-5A synthetase promoter contains multiple
regulatory elements required for interferon-α-induced expression. *DNA Cell Biol.* **16**,
1385–1394 (1997).

108. D. Rebouillat, A. Hovnanian, G. David, A. G. Hovanessian, B. R. G. Williams,
Characterization of the Gene Encoding the 100-kDa Form of Human 2′,5′Oligoadenylate
Synthetase. *Genomics.* **70**, 232–240 (2000).

109. N. Wang, Q. Dong, J. Li, R. K. Jangra, M. Fan, A. R. Brasier, S. M. Lemon, L. M. Pfeffer,
K. Li, Viral induction of the zinc finger antiviral protein is IRF3-dependent but NF-
kappaB-independent. *J. Biol. Chem.* **285**, 6080–90 (2010).

959
960

Acknowledgements: We acknowledge assistance from Mike Cook and the Duke Cancer Institute Flow Cytometry Core. We thank Robert Lefkowitz and his laboratory for assistance with and use of the ChemiDoc Imaging System. We would like to thank Dr. Clare Smith for help establishing SARS-CoV-2 viral infection assays at BSL3. We also thank Ephraim Tsalik, Micah McClain, and Bandita Gershon for helpful discussions and Ben Chambers, Stacy Webb, and other members of the Heaton lab for critical reading of the manuscript. **Funding:** Biocontainment work was performed in the Duke Regional Biocontainment Laboratory under the direction of Greg Sempowski, which received partial support for construction from the National Institutes of Health, National Institute of Allergy and Infectious Diseases (UC6-AI058607). H.M.F. and A.T.H. were supported by NIH training grant T32-CA009111. N.S.H. is supported by R01-HL142985, R01-AI137031, and funding from the Defense Advanced Research Projects Agency's (DARPA) PReemptive Expression of Protective Alleles and Response Elements (PREPARE) program (Cooperative agreement #HR00111920008). The views, opinions and/or findings expressed are those of the author and should not be interpreted as representing the official views or policies of the U.S. Government. **Author contributions:** H.M.F., A.T.H. and N.S.H. designed the study. H.M.F., A.T.H., and B.E.H. performed experiments. H.M.F., A.T.H. and N.S.H. analyzed data. H.M.F. and N.S.H. wrote the paper. **Competing interests:** The authors declare no competing interests. **Data and materials availability:** All next generation sequencing data are available at NCBI GEO under accession number GSE140718. The following reagent was deposited by the Centers for Disease Control and Prevention and obtained through BEI Resources, NIAID, NIH: SARS-Related Coronavirus 2, Isolate USA-WA1/2020, NR-52281.