ETV7 represses a subset of interferon-stimulated genes that restrict influenza viruses

Heather M. Froggatt[1], Alfred T. Harding[1], and Nicholas S. Heaton[1,*]

[1]Department of Molecular Genetics and Microbiology
Duke University School of Medicine
Durham, NC 27710

Keywords:

*To whom correspondence should be addressed:
Nicholas S. Heaton, PhD
Assistant Professor
Department of Molecular Genetics and Microbiology (MGM)
Duke University Medical Center
213 Research Drive, 426 CARL Building, Box 3054
Durham, NC 27710
Tel: 919-684-1351
Fax: 919-684-2790
Email: nicholas.heaton@duke.edu

Abstract

The type I interferon (IFN) response is an important component of the innate immune response to viral infection. Precise control of interferon responses is critical, as insufficient levels of interferon-stimulated genes (ISGs) can lead to a failure to restrict viral spread, while excessive ISG activation results in interferon-related pathologies. While both positive and negative regulatory factors can control the magnitude and duration of IFN signaling, it is also appreciated that a number of ISGs regulate aspects of the interferon response themselves. However, the mechanisms underlying complex ISG regulatory networks remain incompletely defined. In this study, we performed a CRISPR activation screen to identify new regulators of type I IFN responses. We identified ETS variant transcription factor 7 (ETV7), a strongly induced ISG, as a protein that acts as a negative regulator of the type I IFN response; however, ETV7 did not uniformly suppress ISG transcription. Instead, ETV7 specifically targeted a subset of ISGs for regulation based on their promoter sequences. We further showed the subset of ETV7-modulated ISGs is particularly important for control of influenza viruses. Together, our data demonstrate that ETV7 is a component of the complex ISG regulatory network by controlling the expression of a subset of ISGs with a potential role in directing the interferon response against specific viruses.

Significance

Interferons (IFNs) were first described in 1957 and are now known to be critical for restriction of viruses. Still, our understanding of the complex web of interactions that underlie IFN responses remains incomplete. In particular, negative regulation of interferon responses has received disproportionately less study. In this work, we performed a genome-wide overexpression screen for factors capable of suppressing IFN response signaling. We identified a DNA binding transcription factor (ETV7) that, after induction by interferon, acts to suppress a subset of IFN-stimulated genes required for control of influenza viruses. Our work highlights the importance of understanding negative IFN signaling not only with respect to the magnitude and duration of the response, but also the specificity of its antiviral effects.

Introduction

The type I interferon (IFN) response is a transient innate immune defense system that, upon activation by viral infection, induces the transcription of hundreds of interferon-stimulated genes (ISGs) (1). Many ISGs have characterized antiviral roles that restrict viral replication by either interfering with viral processes directly or regulating the cellular pathways required for viral replication (2). However, because replication mechanisms and points of interaction with the cell differ between viruses, individual ISGs have varying potencies against different viruses (3–5). As a result, unique combinations of ISGs are thought to mediate successful antiviral responses against distinct viruses (1, 6).

The activation of the type I IFN signaling pathway in response to viral infection is well understood (7, 8). Extracellular IFN, which is released after a cell recognizes virus-derived nucleic acid, is bound by its cognate plasma membrane-localized receptor (IFNARs). Downstream effectors (JAK proteins) are phosphorylated to then activate interferon-stimulated gene factor 3 (ISGF3) complex formation. Finally, the ISGF3 complex of STAT1, STAT2, and IRF9 translocates to the nucleus (7). There, ISGF3 binds the interferon sensitive response element (ISRE), with the consensus DNA motif GAAANNGAAA, to activate transcription of ISGs (9).

As infection is cleared and virally derived innate immune activators become scarce, interferon production is reduced and the interferon-stimulated gene response is downregulated. To facilitate this return to cell homeostasis, negative regulators are induced and act at multiple levels in the signaling pathway (10). For example, PKD2 is an

ISG that recruits ubiquitin to the IFN receptor, IFNAR1, resulting in its degradation (11).

SOCS1 and SOCS3 are upregulated during, and act to limit, the IFN response through

direct interactions with JAK proteins, while SOCS1 also ubiquitinates other pathway

components (12). USP18 is induced by IFN to help return the cell to homeostasis by

removing the ubiquitin-like ISG15 from target proteins (13). Thus, negative regulators of

IFN responses are an important group of IFN-stimulated genes that control the duration

of ISG induction and activity.

In addition to activating or suppressing IFN responses, there are a number of interferon-

induced regulators that enhance, limit, or fine-tune antiviral activity (14). Many ISGs

themselves participate in innate immune signaling to amplify IFN, and other pro-immune,

responses. For example, IFN signaling increases the levels of STAT1/2 and IRF9, thus

forming a positive feedback loop that enhances further ISG expression (15). Activators

also add complexity by inducing non-canonical IFN response pathways or specific groups

of ISGs. Interferon responsive factors (IRFs) 1 and 7 are ISGs and transcription factors

that activate subsets of ISGs (16, 17). Recent work has shown ELF1 (E74-like ETS

transcription factor) is induced by IFN, resulting in the expression of a group of genes not

otherwise activated by the IFN response (18). These differential ISG profiles are thought

to allow the cell to fine-tune its antiviral activity for an effective and appropriate response.

While interferon-induced positive regulators of the IFN response are known to shape the

complexity of ISG activation, reports of analogous roles for negative regulators remain

conspicuously absent.

To address this gap in knowledge and identify genes able to shape the IFN response

through negative regulation, we performed a CRISPR activation (CRISPRa) screen that

selected for factors sufficient to prevent expression of an ISRE-containing IFN response

reporter. We identified ETV7 (ETS variant transcription factor 7) as a negative regulator

of the type I IFN response with a role in controlling the expression of a subset of ISGs.

We further showed the ETV7-modulated ISGs are important for control of influenza

viruses. Together, these data demonstrate ETV7 is a suppressive component of the

complex ISG regulatory network that could be targeted to enhance specific antiviral

responses against influenza viruses (1, 19).

Results

A CRISPR activation screen identifies potential negative regulators of the type I IFN

response.

In order to identify negative regulators of the type I IFN response, we developed a type I

IFN response reporter (IFNrsp) that included seven copies of the consensus interferon

sensitive response element (ISRE) ahead of a minimal CMV promoter controlling

expression of sfGFP (**Fig. 1A**). To make our reporter temporally specific, sfGFP was

fused to a mouse ornithine decarboxylase (MODC) protein degradation domain to

decrease its half-life (20). We stably introduced this construct into the A549 lung epithelial

cell line along with a dCAS9-VP64 fusion protein and a MS2-p65-HSF1 activator complex

required for the SAM CRISPR activation system (21). After clonal selection, 99.8% of the

A549-SAM-IFNrsp cells expressed GFP in response to type I IFN treatment (**Fig. 1B and**

C). To perform the screen, we took the A549-SAM-IFNrsp cell line and transduced 2×10^8

Fig. 1. A CRISPR activation screen to identify negative regulators of the type I interferon response. A) Diagram of the IFN response reporter (IFNrsp) used to identify cells responding to IFN. ISRE = interferon sensitive response element, MODC = protein degradation domain. B) Flow cytometry histogram and C) bar graph of A549-SAM-IFNrsp cells before and after IFN-α treatment (1000 U/mL, 6 h) (data shown as mean ± SD, n=4). Data shown are representative of two independent experiments. P-values calculated using unpaired, two-tailed Student's t-tests, *p<0.05, **p<0.001. D) Diagram of CRISPRa screen workflow to identify negative regulators of the type I IFN response. E) Results of the two independent CRISPRa screens. Z-score values from the replicate screens with a cutoff of 2 standard deviations from the mean were used to identify top "hits". F) Venn diagram indicating overlapping hits from the replicate screens and genes upregulated by interferon at least two-fold, according to the Interferome database (23).

cells at a multiplicity of infection (MOI) of 0.5 with a lentivirus library containing sgRNAs

designed to activate every putative ORF in the human genome (21) (**Fig. 1D**). After 48

hours, half of the cells were collected to determine the transduction efficiency and the

remaining cells were re-plated for IFN stimulation. At 72 hours post-sgRNA introduction,

the cells were treated with 4,000 U/mL IFN-α for 6 hours and collected for fluorescence-

activated cell sorting. During sorting, we eliminated reporter-positive cells and collected

only cells that were nonresponsive to IFN, because this population should theoretically

be overexpressing a negative regulator of the IFN response. We performed two

independent biological replicates of the screen and sequenced the sgRNA-containing

amplicons derived from our input DNA, unselected transduced cells, and cells that were

nonresponsive to type I IFN. Raw sequencing data was aligned and mapped and

subsequently analyzed using the MAGeCK pipeline (22) to generate z-score values for

each gene. Genes were defined as "hits" if their z-scores exceeded two standard

deviations from the mean, resulting in an overlap of 10 genes between the two screen

replicates (**Fig. 1E, Supplementary Data 1 and 2, and Supplementary Table 1**). We

were seeking to identify regulators of the IFN response that are regulated by IFN

themselves; therefore, we selected hits for validation previously reported to have at least

a two-fold induction after IFN stimulation in the Interferome database (23). This analysis

identified three hits (C1GALT1, ETV7, and NUP153) as potential negative regulators of

the type I IFN response (**Fig. 1F and Supplementary Table 1**).

<u>Overexpression of ETV7 is sufficient to negatively regulate the type I IFN response.</u>

To validate our three hits, and to avoid potential false positive results as the result of off-

target effects of CRISPRa, we cloned the three ORFs and validated overexpression of

the genes in 293T cells (**Supplementary Figure 1**). Co-transfection of the

overexpression plasmids and IFNrsp plasmid, followed by stimulation with IFN-α, resulted

in significantly fewer GFP-expressing cells compared to a control mCherry-expressing

plasmid (**Fig. 2A and B**). To verify this repressive activity was specific to the IFN response

and the hits were not general inhibitors of transcription or translation, we transfected the

overexpression plasmids along with a constitutively active GFP-expressing plasmid (**Fig.**

2C). We included a positive control (EIF2AK1/HRI), which is known to shut off translation

when overexpressed (24). C1GALT1 overexpression significantly downregulated GFP

expression, indicating the repressive activity of C1GALT1 is not completely specific to the

IFN response. While the overexpression of either ETV7 or NUP153 specifically affected

the IFNrsp plasmid, NUP153 has previously been shown to control the distribution of

STAT1 in the cell (25). We therefore chose ETV7 for further characterization because: 1)

ETV7 had not been previously reported to play a role in the IFN response, and 2) it had

the strongest inhibitory phenotype against the IFNrsp reporter.

After confirming overexpression of ETV7 at the protein level (**Fig. 2D**), we verified the

inhibitory effects of ETV7 were not restricted to the reporter plasmid. We collected mRNA

and protein from IFN-α stimulated ETV7 overexpression cells to quantify effects on the

expression of endogenous ISGs. ETV7 overexpression significantly repressed the

induction of three prototypical ISGs (IFIT1, MX1, and ISG15) at the RNA level (**Fig. 2E**).

The reduction of ISG expression during ETV7 overexpression was also demonstrated at

the protein level for IFIT1 (**Fig. 2F**). These experiments show that overexpression of ETV7

is sufficient to repress ISG induction by type I IFN.

Fig. 2. ETV7 overexpression suppresses ISG expression. A) Flow cytometry plots of 293T cells transfected with the IFNrsp reporter and overexpression plasmids for the indicated screen hits then treated with IFN-α (100 U/mL, 6 h). B) Quantification of A showing normalized percentage of cells expressing GFP compared to the mCherry-expressing control (data shown as mean ± SD, n=4). C) Normalized percentage of cells expressing GFP from a constitutively expressing plasmid in cells overexpressing the indicated genes (positive control = EIF2AK1/HRI, shuts off translation) compared to control (data shown as mean ± SD, n=4). D) Western blot showing ETV7 protein levels in 293T cells transfected with the ETV7 overexpression plasmid. Stain-free gel imaging was used to confirm equal loading. E) Endogenous ISG mRNA expression levels measured using RT-qPCR after IFN-α treatment (100 U/mL, 9 h) (data shown as mean ± SD, n=4). F) Western blot comparing IFIT1 protein levels in control and ETV7 overexpressing cells after IFN-α treatment (500 U/mL, 18 h). ns = nonspecific band. Stain-free gel imaging was used to confirm equal loading. For all panels: Data shown are representative of two independent experiments. P-values calculated using unpaired, two-tailed Student's t-tests (*p<0.05, **p<0.001) compared to IFN-stimulated, mCherry-expressing control samples.

ETV7 acts as a transcription factor to repress the type I IFN response.

ETV7 is known to be a repressive transcription factor (26, 27), although a role in

repressing type I IFN responses has never been reported. To determine whether ETV7

acts as a transcription factor in this context, we generated a previously validated mutant

of ETV7, called ETV7(KALK), which is unable to bind DNA (**Fig. 3A and B**) (28).

Overexpression of ETV7(KALK) and stimulation with IFN-α had no measurable effect on

expression of the IFNrsp reporter, in contrast to WT ETV7 overexpression (**Fig. 3C**).

ETV7 has been reported to bind the canonical ETS family DNA motif, GGAA (29), known

as an "ETS" site (**Fig. 3D**). Since consensus ISREs can either contain or lack a GGAA

motif (**Supplementary Table 2**), we hypothesized ETV7 could act on specific ISGs based

on the presence of ETS sites in their promoters. The original IFN response reporter design

contained multiple ETS sites (**Fig. 3E**), which potentially explains why it is negatively

impacted by ETV7. To test the requirements of ETS sites for ETV7 repressive activity, we

generated an IFN response reporter containing seven consensus ISREs from canonical

ISGs that all lack ETS sites (ISRE -ETS) (**Fig. 3E**). We transfected the two reporter

plasmids (ISRE +ETS and ISRE -ETS) independently into 293T cells and stimulated with

IFN-α. As expected, both reporter plasmids responded to IFN treatment, but the

repressive activity of ETV7 was restricted to the reporter plasmid containing ETS motifs

(**Fig. 3F**). These experiments together demonstrate that ETV7's repressive activity

requires both its ability to bind DNA and the presence of ETS sites in ISG promoters.

Fig. 3. ETV7 acts as a transcription factor to negatively regulate the type I IFN response. A) Diagram showing the ETV7 protein domains and amino acid changes made to generate the DNA binding mutant, ETV7(KALK). B) Western blot showing ETV7 protein levels in 293T cells transfected with WT (1μg) or DNA binding mutant (KALK) ETV7 expression plasmids. Stain-free gel imaging was used to confirm equal loading. C) Normalized percentage of 293T cells expressing GFP from the IFNrsp reporter with overexpression of WT or DNA binding mutant (KALK) ETV7 after IFN-α treatment (100 U/mL, 6 h) compared to control (data shown as mean ± SD, n=4, statistical analysis relative to IFN-stimulated, mCherry-expressing control samples). D) ETV7's DNA binding position weighted matrix (PWM) generated using enoLOGOS (69) with data from Wei et al. (70) and the conserved ETS family binding site, GGAA, highlighted in yellow. E) Diagrams of the IFNrsp reporters containing (+ETS) and not containing (-ETS) potential ETV7 binding sites (ETS site, highlighted in yellow). F) Normalized percentage of 293T cells expressing GFP from IFNrsp reporters either containing or not containing ETS sites after overexpression of ETV7 and IFN-α treatment (100 U/mL, 6 h) compared to mCherry-expressing control (data shown as mean ± SD, n=4). For all panels: Data shown are representative of two independent experiments. P-values calculated using unpaired, two-tailed Student's t-tests (*p<0.05, **p<0.001).

<u>ETV7 differentially regulates genes based on their ISRE sequence.</u>

Our data suggested ETV7 likely does not affect all ISG promoters. To perform an

unbiased examination of ETV7's repressive activity against ISGs with a variety of

potential regulatory sites, we performed RNA sequencing in cells with or without ETV7

overexpression and IFN stimulation (**Supplementary Data 3**). We then generated a

dendrogram using the 2,000 most differentially expressed genes after IFN treatment to

compare the four conditions: overexpression of control protein (mCherry) or ETV7, and

with or without IFN-α treatment. When comparing the impact of IFN treatment on control

and ETV7-overexpressing cells, we observed a larger divergence in the transcriptional

profile of control cells compared to ETV7-overexpressing cells after IFN treatment (**Fig.**

4A). This difference demonstrates that ETV7 generally "dampens" the transcriptional

impact of IFN treatment. Using a heat map to observe patterns in genes that increased

at least two-fold upon IFN treatment, we found some genes are more suppressed during

ETV7 overexpression than others (**Fig. 4B**). We divided these genes into three groups

(from I = most affected to III = least affected) depending on their response to ETV7

overexpression and we examined their promoters to identify motifs associated with ETS

transcription factors and IFN regulation. Unexpectedly, comparing the number of ETS

binding sites (GGAA) across these three groups revealed no significant difference

between the differentially affected groups (**Fig. 4C**). However, it is known that ETS sites

sometimes occur as a part of combined motif related to ISREs, known as ETS-IRF

combined elements (EICEs) with the consensus sequence GGAANN(N)GAAA (30, 31).

We therefore tested the hypothesis that ETV7 negatively regulates ISGs with EICE sites.

The number of EICE sites was significantly different between the most and least ETV7-

affected groups (**Fig. 4D**), indicating ETV7 impacts the expression of specific ISGs by

targeting an extended DNA binding motif.

Fig. 4. ETV7 differentially regulates ISGs during the type I IFN response based on ISRE-related regulatory elements. A) Dendrogram of genes most differentially expressed in cells overexpressing either a control protein (mCherry) or ETV7 before and after IFN-α treatment (100 U/mL, 9 h) as measured using RNA sequencing. Three independent, biological replicates per condition. Red box highlights control samples, yellow box highlights ETV7-expressing samples, shading indicates IFN-stimulated samples. The box width indicates the linkage distance between samples before and after IFN, indicating control cells' transcriptional profile is more diverged after IFN treatment compared to ETV7-expressing cells. B) Heat map displaying RNA levels of genes upregulated at least two-fold following IFN-α treatment (100 U/mL, 9 h) in control cells. Expression was normalized to control cells after IFN treatment (averaged across replicates). Yellow = downregulated, blue = upregulated. C,D) Motif counts in promoter regions (-1000 bp, +500 bp) for the genes most and least affected by ETV7 overexpression in the RNA sequencing results. ETS sites (GGAA) highlighted in yellow. P-values calculated using unpaired, two-tailed Mann-Whitney U tests (*p<0.05, **p<0.001).

<u>ETV7 is required to negatively regulate specific ISGs.</u>

Our experiments to this point used an overexpression system to demonstrate that ETV7

is sufficient to suppress ISG expression. However, this approach leads to constitutive

ETV7 expression at high levels relative to the physiological magnitude and IFN-induced

expression of ETV7 (**Fig. 5A**). To determine the importance of ETV7 induction during the

IFN response, we performed a series of loss of function experiments that we expected

would have the reciprocal effect on IFN responses (32). We transduced A549-IFNrsp

reporter cells (the original reporter with ISRE +ETS sites) with Cas9 and one of two

different sgRNAs targeting ETV7 (ETV7 KO1, ETV7 KO2), selected for edited cells, and

then stimulated with IFN-α. Both guides resulted in significantly more IFN-induced sfGFP

expression compared to a control sgRNA (**Fig. 5B**). We next generated clonal ETV7

knockout A549 lung epithelial cell lines and sequenced the resulting DNA lesions to

confirm ETV7 knockout (**Supplementary Figure 2**). Since ETV7 is normally only

expressed after IFN stimulation, we treated with IFN-α and verified a reduction in ETV7

expression at the RNA level (**Fig. 5C**). We then selected five ISGs for RT-qPCR analysis.

Three (IFI44L, RSAD2/Viperin, IFIT3) were from the group most affected by ETV7 (Group

I) that contained multiple EICEs in their promoters. Two (IFIT5, IRF9) were chosen from

the less affected groups (Groups II and III). The EICE-rich genes (Group I) showed

significantly higher levels of RNA expression in the ETV7 knockout cells (**Fig. 5D**), while

genes with few EICEs were not significantly impacted by the loss of ETV7 (**Fig. 5E**). Thus,

the physiological level of ETV7 induction after IFN stimulation affects the expression of a

subset of ISGs.

Fig. 5. ETV7 loss enhances expression of specific ISGs. A) ISG15 and ETV7 mRNA levels in A549 lung epithelial cells after IFN-α treatment (1000 U/mL, 6 h) (data shown as mean ± SD, n=4). B) Percentage of A549 cells expressing GFP from IFNrsp reporter after knockout of ETV7 and IFN-α treatment (1000 U/mL, 6 h) (data shown as mean ± SD, n=3, statistical analysis represents p-values for both of the two ETV7 KO sgRNAs compared to a non-targeting control). C) mRNA levels of ETV7 in non-targeting control and ETV7 KO A549 cells after IFN-α treatment (1000 U/mL, 6 h) (data shown as mean ± SD, n=4). D,E) Representative genes were chosen from the groups D) most affected by ETV7 (Group I) and E) least affected (Groups II and III) in the RNA sequencing analysis (**Fig. 4B**). Each gene's potential ISRE sequences (ETS sites highlighted in yellow) are shown, along with its mRNA levels in control and ETV7 KO cells after IFN-α treatment (1000 U/mL, 6 h) (data shown as mean ± SD, n=4). For all panels: Data shown are representative of two independent experiments. P-values calculated using unpaired, two-tailed Student's t-tests (*p<0.05, **p<0.001) compared to IFN-stimulated, non-targeting sgRNA control samples unless otherwise indicated.

<u>Loss of ETV7 restricts influenza viral replication.</u>

For a successful antiviral response, individual ISGs are thought to work together to restrict

multiple parts of the virus replication cycle (1). To determine whether the effects of ETV7

suppression of ISG expression were relevant in the context of a viral infection, we wanted

to identify a virus restricted by the genes regulated by ETV7 (i.e. Group I genes) (33).

Considering the Group I genes with well recognized antiviral functions (IFITM1, IFIT1-3,

OAS1-3, BST2, RSAD2), we found each had been reported to play important roles in the

restriction of influenza viruses (34). IFITM1 has been shown to prevent viral entry (35),

OAS proteins activate RNase L to degrade viral RNA (36), IFITs bind viral RNA and

promote antiviral signaling (37), and BST2/Tetherin and RSAD2/Viperin restrict viral

budding and egress (38, 39).

To determine whether ETV7 regulation affected influenza virus infection, we first infected

our ETV7 knockout A549 cells with a laboratory-adapted H1N1 influenza A virus (IAV),

A/Puerto Rico/8/1934 (PR8). Using a hemagglutination (HA) assay to measure the

number of viral particles released over time, we observed reduced virus production in our

ETV7 KO cells compared to control cells (**Fig. 6A**). This was the anticipated outcome

because loss of a negative regulator (i.e. ETV7) is expected to enhance expression of

antiviral ISGs. We also measured infectious viral titers and found a significant reduction

in our ETV7 KO cells compared to control cells (**Fig. 6B**). Using a fluorescent reporter

strain of PR8 (PR8-mNeon) (40), we next visualized infection and spread. As expected,

we observed fewer cells expressing mNeon in ETV7 KO cells using both microscopy (**Fig.**

6C) and flow cytometry readouts (**Fig. 6D**). Next, we tested whether ETV7's impact on

influenza virus infection and spread would extend to a more contemporary H1N1 IAV

strain, A/California/07/2009 (Cal/09), as well as an unrelated Victoria lineage influenza B

virus strain, B/Malaysia/2506/2004 (Mal/04) (41). Using these fluorescent reporter

Fig. 6. Loss of ETV7 limits replication of multiple influenza viruses. A) Hemagglutination (HA) assay of virus collected at indicated time points from non-targeting control and ETV7 KO A549 cells after infection with WT PR8 virus (MOI=0.05, multicycle infection) (data shown as mean ± SD, n=3). B) Titer of virus collected from control and ETV7 KO A549 cells after infection with WT PR8 virus (18 h, MOI=0.05, multicycle infection) (data shown as mean ± SD, n=4). C) Control or ETV7 KO A549 cells after mock or PR8-mNeon reporter virus infection (24 h, MOI=0.1, multicycle infection). Green = mNeon, blue = nuclei. Scale bar, 200 μm. D) Flow cytometry quantification of control or ETV7 KO A549 cells after infection with PR8-mNeon reporter virus (24 h, MOI=0.01, multicycle infection) (data shown as mean ± SD, n=3). E,F) Normalized percentage of infected (reporter+) cells in ETV7 KO A549 cells compared to control cells after infection with E) Cal/09-sfGFP or F) Mal/04-mNeon reporter viruses (24 h, multicycle infection) (data shown as mean ± SD, n=4). For all panels: Data shown are representative of two independent experiments. P-values calculated using unpaired, two-tailed Student's t-tests (*p<0.05, **p<0.001) compared to influenza infected, non-targeting sgRNA control samples.

289   viruses, we observed significant decreases in the number of Cal/09- and Mal/04-infected

290   cells when comparing ETV7 KO cells to control cells (**Fig. 6E and F**). These experiments

291   demonstrate that loss of ETV7 leads to decreased viral replication across multiple,

292   unrelated influenza viruses.

Discussion

In this study, we performed a CRISPR activation screen to identify negative regulators of the type I IFN response. Specifically, we were interested in negative regulators that contribute to the types of differentiated ISG profiles IFN-induced activators are reported to produce. From this screen, we identified ETV7 as a negative IFN regulator and showed it acts as a transcription factor to repress subsets of ISGs dependent on a motif related to the ISRE, the EICE. We also showed ETV7's regulatory activity impacts the replication and spread of multiple strains of influenza viruses. These findings demonstrate the importance of ETV7 in fine-tuning the IFN response through specificity and transcriptional repression to regulate particular, antiviral ISG targets.

ETV7 is a member of the ETS family of transcription factors. This family performs diverse functions despite recognizing the same core DNA sequence, GGAA, by acting on extended motifs requiring binding partners (42, 43). In our work we identified the EICE, a recognized ETS transcription factor-associated motif, as a regulatory element related to ISREs that ETV7 uses to discriminate genes for regulation. The EICE has previously been reported to require an IRF binding partner to direct ETS transcription factor activity (30, 31); therefore, it is likely ETV7 has an IRF binding partner. If ETV7 does require a binding partner, this protein's induction and distribution likely contribute to the timing, gene targets, and activity of ETV7 during the IFN response. It is known that IRFs can be basally expressed (IRF2, IRF3) or IFN-induced (IRF1, IRF7) (44) and the availability of a binding partner could dramatically affect the timing and magnitude of effects on EICE-controlled ISGs. Future work will define if ETV7 has specific binding partners and how those

interactions may contribute to the nonuniform, repressive activity of ETV7 during the type

I IFN response reported in this study.

IFN-induced regulators control the magnitude and duration of IFN responses in addition

to the temporal regulation of specific waves of ISGs (45). These coordinated waves of

ISG induction can peak early or late during the IFN response and are thought to

correspond to specific stages of virus replication or immune processes (1, 6). We

compared the induction of ETV7 and ISG15 and observed ETV7 is both upregulated and

downregulated at earlier time points than this prototypical ISG (**Fig. 5A**). We expanded

our analysis to published datasets of human gene expression during respiratory infections

and concluded that ETV7 is generally induced earlier than many ISGs (46). Although not

the focus of our study, ETV7's early and short induction pattern suggests it may be a key

regulator of the first stages of IFN-mediated gene induction. We favor a model wherein

early ETV7 expression is responsible for reducing the accumulation, or delaying the

expression, of ISGs controlled by EICE motifs (**Supplementary Fig. 3**).

ETV7 is induced during infections across many vertebrate species (47, 48), indicating a

potential conserved, relevant role in the immune response; however, ETV7 has been lost

in mice and closely related rodents (49). Since mice and rodents have an intact interferon

response pathway, a natural question is: how are the activities of ETV7 being accounted

for in these animals? While we have no clear answer from the data in this study, it is well-

recognized that IFN responses contains many redundancies (33). Accordingly, we believe

other ETS family members, potentially the closely related ETV6 (which is also induced by

IFNs), may perform the role of ETV7 in mice (50). Future studies will be required to test

the hypothesis that mice induce an ETV7-related alternative during the type I IFN

response.

Another important question is why ETV7's IFN-induced activity has been maintained

throughout evolution. In this report, we provide evidence that ETV7's activity reduces a

cell's ability to restrict influenza virus infection; this seems counterintuitive to ETV7

benefitting the host. We hypothesize that regulators like ETV7 are important to prevent

excessive inflammatory signaling. It is appreciated that negative regulators of the IFN

response are required to prevent extreme and prolonged immune responses, which are

associated with poor disease outcomes after infection (51–53). ETV7 potentially

contributes to the cumulative activities of negative IFN regulators to limit IFN responses

during pathogen clearance. Additionally, it stands to reason that different individual ISGs

have differing toxic effects on the cell. It is tempting to speculate that ETV7 suppresses

ISGs whose accumulation is particularly harmful to cell viability and host recovery after

infection.

Additionally, the relevance of controlled IFN responses goes beyond infectious disease;

patients with dysfunctional USP18, a negative regulator of the IFN response, develop a

type I interferonopathy that results in a severe pseudo-TORCH syndrome (54). Mouse

knockouts for other negative regulators of the IFN response (SOCS1, SOCS3, USP18)

also develop non-pathogen associated, chronic inflammatory diseases (55–58). ETV7's

lack of murine homolog eliminates an easily generated animal-knockout model to

experimentally show ETV7's relevance as a general innate immune repressor. However,

genome wide association studies (GWAS) have linked ETV7 to autoimmune diseases

including rheumatoid arthritis and multiple sclerosis (59, 60); both of these autoimmune

diseases have evidence of enhanced ISG expression (61, 62). Thus, although the specific

contributions of ETV7 activity to IFN regulation are currently undefined, its potential role

is not limited to viral infections.

In conclusion, we identified ETV7 as a negative regulator of the type I IFN response.

Previously, ETV7 was appreciated to be an ISG; however, a specific function during the

IFN response was unknown. We determined that ETV7 acts as a transcription factor to

target specific ISGs for repression, potentially contributing to the complex ISG

transcriptional landscape. Additionally, many of the ETV7-modulated ISGs restrict

influenza viruses (34) and we showed that loss of ETV7 limits influenza virus spread.

Further work is required to understand the complexity of IFN regulation, while therapeutic

targeting of factors like ETV7 could lead to the development of a new class of host-

directed antivirals that tailor ISG responses to specific viruses.

Experimental Procedures

Cloning

To generate reporters sensitive to IFN, we designed gBlocks (IDT) containing ISREs to be cloned into the pTRIP vector ahead of a minimal CMV promoter controlling expression of sfGFP. To clone and express the open reading frames (ORFs) of our screen hits, we designed primers for cloning into the pLEX-MCS vector using Gibson Assembly (NEB). To amplify ETV7 and NUP153, we used cDNA templates from Transomic Technologies. To amplify C1GALT1 and EIF2AK1, we used RNA from IFN-stimulated A549 cells. The DNA binding mutant, ETV7(KALK) (28), was also generated using a gBlock. Non-targeting and ETV7-targeting CRISPR KO sgRNAs were cloned by annealing oligos encoding the desired sgRNA sequence and ligating them directly into the lentiCRISPRv2 vector (Addgene). DNA was transformed into NEB 5-alpha high efficiency competent cells. Insert size was verified with PCR and purified plasmids were sequenced using Sanger sequencing.

Cells

All cells were obtained from ATCC and grown at 37°C in 5% CO_2. A549 and 293T cells were grown in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 5% fetal bovine serum, GlutaMAX, and penicillin-streptomycin. Madin-Darby canine kidney (MDCK) cells were grown in minimal essential media (MEM) supplemented with 5% fetal bovine serum, HEPES, $NaHCO_3$, GlutaMAX, and penicillin-streptomycin. The A549 CRISPR-SAM cells were previously validated (63) and transduced with the IFNrsp

reporter three times before being clonally selected. The A549 CRISPR KO cells were

transduced and then selected using puromycin (10 μg/mL).

Flow Cytometry

Cells were trypsinized and analyzed on a Fortessa X-20 (BD) machine with standard laser

and filter combinations. Data was visualized and processed with FlowJo software.

CRISPR Activation Screen

The sgRNA library was packaged into lentivirus as previously described (63). After

packaging and titering the lentivirus, 2×10^8 A549-CRISPR-SAM-IFNresp cells were

seeded onto 15 cm plates (10 plates total). The next day they were transduced with the

packaged sgRNA library (MOI=0.5). After 48 h, the transduced cells were split and half

were collected as a transduction control, while the remaining half were plated back onto

15 cm plates. The next day, cells were treated with IFN-α (4×10^3 U/mL) for 6 h. Cells were

then collected and sorted on a Beckman Coulter Astrios cell sorter. Specifically, gates

were set to sort GFP-negative cells as the population of interest, as well as GFP-positive

cells as a control population of cells still capable of signaling. This screen was performed

in duplicate. Genomic DNA was extracted from sorted cells using the Zymo Quick gDNA

micro prep kit. PCR was subsequently performed using barcoded primers as previously

described using the NEB Next High Fidelity 2x PCR master mix (63). PCR bands were

gel purified using the Thermo GeneJET gel extraction kit. Samples were then sequenced

via next-generation Illumina MiSeq using paired-end 150 bp reads.

Screen Analysis

Raw MiSeq read files were aligned to the CRISPR SAM sgRNA library and raw reads for each sgRNA were counted using the MAGeCK pipeline (22). sgRNA enrichment was determined using the generated count files and the MAGeCK-MLE analysis pipeline. Genes were sorted based on z-score and determined to be significantly enriched if their z-score was at least two standard deviations above the average z-score of the entire sorted population.

Western Blotting

Cells were trypsinized and $1x10^6$ cells were pelleted at 800 x g for 5 min. Equal amounts of cellular material were loaded into 4-20% acrylamide gels (Bio-Rad) and imaged using a ChemiDoc Imaging System (Bio-Rad). Protein was transferred to a nitrocellulose membrane at 60V for 60 min. PBS with 5% (w/v) non-fat dried milk and 0.1% Tween-20 were used to block for 1 h at 4°C. Primary antibodies were then incubated with the membrane overnight at 4°C. Antibodies used were rabbit anti-ETV7 (Sigma, HPA029033) and rabbit anti-IFIT1 (Cell Signaling, D2X9Z). Membranes were washed five times in PBS with 0.1% Tween-20 and then an anti-rabbit-HRP secondary antibody (Thermo, A16104) was added for 1 h. The membrane was then washed five times and Clarity or Clarity Max ECL substrate (Bio-Rad) was added before being exposed to film and developed.

RT-qPCR

Total RNA was collected using Monarch Total RNA Miniprep Kits (NEB). One-step RT-qPCR was performed with commercial TaqMan assays from Thermo for ETV7

(Hs00903229_m1), C1GALT1 (Hs00863329_g1), NUP153 (Hs01018919_m1), ISG15

(Hs00196051_m1), MX1 (Hs00895608_m1), IFIT1 (Hs00356631_g1), RSAD2

(Hs00895608_m1), IFI44L (Hs00915292_m1), IFIT3 (Hs01922752_s1), IFIT5

(Hs00202721_m1), and IRF9 (Hs00196051_m1) using the EXPRESS One-Step

Superscript qRT-PCR Kit on an Applied Biosystems StepOnePlus instrument. RNA was

normalized using an endogenous 18S rRNA primer/probe set (Applied Biosystems).

RNA sequencing

293T cells were transfected with ETV7- or control-expressing plasmids and selected

using puromycin (20 μg/mL) for 24 h before treatment with IFN-α (100 U/mL). Total RNA

was collected at 9 h post-IFN treatment using Monarch Total RNA Miniprep Kits (NEB).

RNA was prepped for RNA sequencing submission using the NEBNext Poly(A) mRNA

Magnetic Isolation Module (NEB), NEBNext Ultra II RNA Library Prep Kit for Illumina

(NEB), and NEBNext Multiplex Oligos for Illumina (NEB). Samples were analyzed on one

lane of an Illumina HiSeq 4000 using 50 bp single strand reads. Mapping of the raw reads

to the human hg19 reference genome was accomplished using a custom application on

the Illumina BaseSpace Sequence Hub (64). After data normalization, average read

values were compared across samples. For comparisons in which some samples had

zero reads detected for a specific gene, one read was added to all values in the sample

to complete analyses that required non-zero values. Dendrograms were generated by

identifying the 2,000 most differentially expressed genes in the control samples with and

without IFN treatment using a Student's t-test and plotted using Heatmapper (65). The

heat map shows genes upregulated 2-fold (after normalization) with IFN treatment in the

control samples. Values shown are log normalized to the control samples with IFN

treatment.

Viruses

PR8-mNeon was generated via insertion of the mNeon fluorescent gene (66) into

segment 4 of the virus (40). Mal/04-mNeon was generated by inserting the mNeon

fluorescent gene (66) into segment 4 of the Mal/04 genome (41). Cal/09-sfGFP was

generated via insertion of the sfGFP gene (67) into segment 8 of the virus using the same

scheme previously used to insert Cre recombinase (68). For influenza virus infections,

cells were either mock- or virus-infected for 1 h and then cultured in OptiMEM

supplemented with bovine serum albumin (BSA), penicillin-streptomycin, and 0.2 μg/mL

TPCK-treated trypsin protease (Sigma). PR8 WT, PR8-mNeon, Cal/09-sfGFP, and were

incubated at 37°C, Mal/04-mNeon was incubated ay 33°C.

Viral Growth Assays

Hemagglutination (HA) assays to measure the amount of viral particles were performed

by diluting influenza infected cell supernatants collected at the indicated time points in

cold PBS. An equal amount of chicken blood diluted 1:40 in PBS was mixed with serially

diluted virus and incubated at 4°C for 2-3 h before scoring. Infectious viral titers were

determined using standard plaque assay procedures on MDCK cells. Infected cell

supernatants were collected at 18 h, serially diluted, and used to infect confluent 6-well

plates for 1 h before removing the virus and adding the agar overlay. Cells were then

incubated at 37°C for 48 h before being fixed in 4% PFA overnight. The 4% PFA was then

aspirated, and the agar layer was removed before washing cells with PBS. Serum from

WT PR8 infected mice was diluted 1:2,000 in antibody dilution buffer (5% (w/v) non-fat

dried milk and 0.05% Tween-20 in PBS) and incubated on cells at 4°C overnight. Cells

were then washed twice with PBS and incubated for 1 h with anti-mouse IgG horseradish

peroxidase (HRP)-conjugated sheep antibody (GE Healthcare) diluted 1:4,000 in

antibody dilution buffer. Assays were then washed twice with PBS and exposed to 0.5 mL

of TrueBlue peroxidase substrate (KPL) for 20 min. Plates were then washed with water

and dried before plaques were counted.

Data Availability

All next generation sequencing data are available at NCBI GEO under accession number

GSE140718.

References

1. W. M. Schneider, M. D. Chevillotte, C. M. Rice, Interferon-Stimulated Genes: A Complex Web of Host Defenses. *Annu. Rev. Immunol.* **32**, 513–545 (2014).
2. J. W. Schoggins, Interferon-Stimulated Genes: What Do They All Do? *Annu. Rev. Virol.* **6**, 567–584 (2019).
3. J. W. Schoggins, *et al.*, A diverse range of gene products are effectors of the type I interferon antiviral response. *Nature* **472**, 481–485 (2011).
4. J. W. Schoggins, *et al.*, Pan-viral specificity of IFN-induced genes reveals new roles for cGAS in innate immunity. *Nature* **505**, 691–695 (2014).
5. M. Kane, *et al.*, Identification of Interferon-Stimulated Genes with Antiretroviral Activity. *Cell Host Microbe* **20**, 392–405 (2016).
6. E. V. Mesev, R. A. LeDesma, A. Ploss, Decoding type I and III interferon signalling during viral infection. *Nat. Microbiol.* **4**, 914–924 (2019).
7. G. R. Stark, J. E. Darnell, The JAK-STAT Pathway at Twenty. *Immunity* **36**, 503–514 (2012).
8. K. Honda, H. Yanai, A. Takaoka, T. Taniguchi, Regulation of the type I IFN induction: a current view. *Int. Immunol.* **17**, 1367–1378 (2005).
9. D. E. Levy, D. S. Kessler, R. Pine, N. Reich, J. E. Darnell, Interferon-induced nuclear factors that bind a shared promoter element correlate with positive and negative transcriptional control. *Genes Dev.* **2**, 383–93 (1988).
10. K.-I. Arimoto, S. Miyauchi, S. A. Stoner, J.-B. Fan, D.-E. Zhang, Negative regulation of type I IFN signaling. *J. Leukoc. Biol.* **103**, 1099–1116 (2018).
11. H. Zheng, J. Qian, B. Varghese, D. P. Baker, S. Fuchs, Ligand-stimulated downregulation of the alpha interferon receptor: role of protein kinase D2. *Mol. Cell. Biol.* **31**, 710–20 (2011).
12. N. P. D. Liau, *et al.*, The molecular basis of JAK/STAT inhibition by SOCS1. *Nat. Commun.* **9**, 1558 (2018).
13. M. P. Malakhov, O. A. Malakhova, K. Il Kim, K. J. Ritchie, D.-E. Zhang, UBP43 (USP18) specifically removes ISG15 from conjugated proteins. *J. Biol. Chem.* **277**, 9976–81 (2002).
14. P. J. Hertzog, B. R. G. Williams, Fine tuning type I interferon responses. *Cytokine Growth Factor Rev.* **24**, 217–225 (2013).
15. J. P. B. Viola, *et al.*, A Positive Feedback Amplifier Circuit That Regulates Interferon (IFN)-Stimulated Gene Expression and Controls Type I and Type II IFN Responses. **9**, 28 (2018).
16. N. Tanaka, T. Kawakami, T. Taniguchi, Recognition DNA sequences of interferon regulatory factor 1 (IRF-1) and IRF-2, regulators of cell growth and the interferon system. *Mol. Cell. Biol.* **13**, 4531–8 (1993).
17. M. Sato, *et al.*, Positive feedback regulation of type I *IFN* genes by the IFN-inducible transcription factor IRF-7. *FEBS Lett.* **441**, 106–110 (1998).
18. L. L. Seifert, *et al.*, The ETS transcription factor ELF1 regulates a broadly antiviral program distinct from the type I interferon response. *PLOS Pathog.* **15**, e1007634 (2019).
19. P. Hubel, *et al.*, A protein-interaction network of interferon-stimulated genes extends the innate immune system landscape. *Nat. Immunol.* **20**, 493–502

552 (2019).

20. X. Li, *et al.*, Generation of Destabilized Green Fluorescent Protein as a Transcription Reporter. *J. Biol. Chem.* **273**, 34970–34975 (1998).

21. S. Konermann, *et al.*, Genome-scale transcriptional activation by an engineered CRISPR-Cas9 complex. *Nature* **517**, 583–8 (2015).

22. W. Li, *et al.*, MAGeCK enables robust identification of essential genes from genome-scale CRISPR/Cas9 knockout screens. *Genome Biol.* **15**, 554 (2014).

23. I. Rusinova, *et al.*, INTERFEROME v2.0: an updated database of annotated interferon-regulated genes. *Nucleic Acids Res.* **41**, D1040–D1046 (2012).

24. A.-P. Han, *et al.*, Heme-regulated eIF2alpha kinase (HRI) is required for translational regulation and survival of erythroid precursors in iron deficiency. *EMBO J.* **20**, 6909–6918 (2001).

25. A. Marg, *et al.*, Nucleocytoplasmic shuttling by nucleoporins Nup153 and Nup214 and CRM1-dependent nuclear export control the subcellular distribution of latent Stat1. *J. Cell Biol.* **165**, 823–33 (2004).

26. M. D. Potter, A. Buijs, B. Kreider, L. van Rompaey, G. C. Grosveld, Identification and characterization of a new human ETS-family transcription factor, TEL2, that is expressed in hematopoietic tissues and can associate with TEL1/ETV6. *Blood* **95**, 3341–8 (2000).

27. H. Poirel, *et al.*, Characterization of a novel ETS gene, TELB, encoding a protein structurally and functionaly related to TEL. *Oncogene* **19**, 4802–4806 (2000).

28. F. C. Harwood, *et al.*, ETV7 is an essential component of a rapamycin-insensitive mTOR complex in cancer. *Sci. Adv.* **4**, eaar3938 (2018).

29. H. Wei, *et al.*, Genome-wide analysis of ETS-family DNA-binding in vitro and in vivo. *EMBO J.* **29**, 2147–2160 (2010).

30. A. L. Brass, E. Kehrli, C. F. Eisenbeis, U. Storb, H. Singh, Pip, a lymphoid-restricted IRF, contains a regulatory domain that is important for autoinhibition and ternary complex formation with the Ets factor PU.1. *Genes Dev.* **10**, 2335–2347 (1996).

31. D. Meraro, M. Gleit-Kielmanowicz, H. Hauser, B.-Z. Levi, IFN-Stimulated Gene 15 Is Synergistically Activated Through Interactions Between the Myelocyte/Lymphocyte-Specific Transcription Factors, PU.1, IFN Regulatory Factor-8/IFN Consensus Sequence Binding Protein, and IFN Regulatory Factor-4: Characterization o. *J. Immunol.* **168**, 6224–6231 (2002).

32. N. E. Sanjana, O. Shalem, F. Zhang, Improved vectors and genome-wide libraries for CRISPR screening. *Nat. Methods* **11**, 783–784 (2014).

33. J. W. Schoggins, Recent advances in antiviral interferon-stimulated gene biology. *F1000Research* **7**, 309 (2018).

34. A. García-Sastre, Induction and evasion of type I interferon responses by influenza viruses. *Virus Res.* **162**, 12–18 (2011).

35. A. L. Brass, *et al.*, The IFITM Proteins Mediate Cellular Resistance to Influenza A H1N1 Virus, West Nile Virus, and Dengue Virus. *Cell* **139**, 1243–1254 (2009).

36. Y. Li, *et al.*, Activation of RNase L is dependent on OAS3 expression during infection with diverse human viruses. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 2241–6 (2016).

37. A. Pichlmair, *et al.*, IFIT1 is an antiviral protein that recognizes 5′-triphosphate

RNA. *Nat. Immunol.* **12**, 624–630 (2011).

38.   M. A. Yondola, *et al.*, Budding capability of the influenza virus neuraminidase can
be modulated by tetherin. *J. Virol.* **85**, 2480–91 (2011).

39.   X. Wang, E. R. Hinson, P. Cresswell, The Interferon-Inducible Protein Viperin
Inhibits Influenza Virus Release by Perturbing Lipid Rafts. *Cell Host Microbe* **2**,
96–105 (2007).

40.   A. T. Harding, B. E. Heaton, R. E. Dumm, N. S. Heaton, Rationally Designed
Influenza Virus Vaccines That Are Antigenically Stable during Growth in Eggs.
MBio **8** (2017).

41.   R. E. Dumm, *et al.*, Non-lytic clearance of influenza B virus from infected cells
preserves epithelial barrier function. *Nat. Commun.* **10**, 779 (2019).

42.   R. Li, H. Pei, D. K. Watson, Regulation of Ets function by protein–protein
interactions. *Oncogene* **19**, 6514–6523 (2000).

43.   A. Verger, M. Duterque-Coquillaud, When Ets transcription factors meet their
partners. *BioEssays* **24**, 362–370 (2002).

44.   T. Taniguchi, K. Ogasawara, A. Takaoka, N. Tanaka, IRF Family of Transcription
Factors as Regulators of Host Defense. *Annu. Rev. Immunol.* **19**, 623–655
(2001).

45.   W. Wang, L. Xu, J. Su, M. P. Peppelenbosch, Q. Pan, Transcriptional Regulation
of Antiviral Interferon-Stimulated Genes. *Trends Microbiol.* **25**, 573–584 (2017).

46.   T.-Y. Liu, *et al.*, An individualized predictor of health and disease using paired
reference and target samples. *BMC Bioinformatics* **17**, 47 (2016).

47.   P. C. De La Cruz-Rivera, *et al.*, The IFN Response in Bats Displays Distinctive
IFN-Stimulated Gene Expression Kinetics with Atypical RNASEL Induction. *J.*
Immunol. **200**, 209–217 (2018).

48. ,   Transcriptome and proteome profiling of host responses to Marek's disease virus
in chickens. *Vet. Immunol. Immunopathol.* **138**, 292–302 (2010).

49.   H. Kawagoe, M. Potter, J. Ellis, G. C. Grosveld, TEL2, an ETS factor expressed in
human leukemia, regulates monocytic differentiation of U937 cells and blocks the
inhibitory effect of TEL1 on Ras-induced cellular transformation. *Cancer Res.* **64**,
6091–6100 (2004).

50.   P. Rasighaemi, A. C. Ward, ETV6 and ETV7: Siblings in hematopoiesis and its
disruption in disease. *Crit. Rev. Oncol. Hematol.* **116**, 106–115 (2017).

51.   J. A. Nick, *et al.*, Extremes of Interferon-Stimulated Gene Expression Associate
with Worse Outcomes in the Acute Respiratory Distress Syndrome. *PLoS One* **11**,
e0162490 (2016).

52.   Y. Muramoto, *et al.*, Disease severity is associated with differential gene
expression at the early and late phases of infection in nonhuman primates
infected with different H5N1 highly pathogenic avian influenza viruses. *J. Virol.* **88**,
8981–97 (2014).

53.   S. Davidson, M. K. Maini, A. Wack, Disease-promoting effects of type I interferons
in viral, bacterial, and coinfections. *J. Interferon Cytokine Res.* **35**, 252–64 (2015).

54.   M. E. C. Meuwissen, *et al.*, Human USP18 deficiency underlies type 1
interferonopathy leading to severe pseudo-TORCH syndrome. *J. Exp. Med.* **213**,
1163–74 (2016).

55.   J.-C. Marine, *et al.*, SOCS1 Deficiency Causes a Lymphocyte-Dependent

Perinatal Lethality. *Cell* **98**, 609–616 (1999).

56. W. S. Alexander, *et al.*, SOCS1 Is a Critical Inhibitor of Interferon γ Signaling and
Prevents the Potentially Fatal Neonatal Actions of this Cytokine. *Cell* **98**, 597–608
(1999).

57. B. A. Croker, *et al.*, SOCS3 negatively regulates IL-6 signaling in vivo. *Nat.*
Immunol. **4**, 540–545 (2003).

58. K. J. Ritchie, *et al.*, Dysregulation of protein modification by ISG15 results in brain
cell injury. *Genes Dev.* **16**, 2207–12 (2002).

59. Y. Okada, *et al.*, Genetics of rheumatoid arthritis contributes to biology and drug
discovery. *Nature* **506**, 376–381 (2014).

60. T. James, *et al.*, Impact of genetic risk loci for multiple sclerosis on expression of
proximal genes in patients. *Hum. Mol. Genet.* **27**, 912–928 (2018).

61. J. E. Castañeda-Delgado, *et al.*, Type I Interferon Gene Response Is Increased in
Early and Established Rheumatoid Arthritis and Correlates with Autoantibody
Production. *Front. Immunol.* **8**, 285 (2017).

62. A. Hundeshagen, *et al.*, Elevated type I interferon-like activity in a subset of
multiple sclerosis patients: molecular basis and clinical relevance. *J.*
Neuroinflammation **9**, 574 (2012).

63. B. E. Heaton, *et al.*, A CRISPR Activation Screen Identifies a Pan-avian Influenza
Virus Inhibitory Host Factor. *Cell Rep.* **20**, 1503–1512 (2017).

64. N. S. Heaton, *et al.*, Long-term survival of influenza virus infected club cells drives
immunopathology. **211**, 1707–1714 (2014).

65. S. Babicki, *et al.*, Heatmapper: web-enabled heat mapping for all. *Nucleic Acids*
Res. **44**, W147–W153 (2016).

66. N. C. Shaner, *et al.*, A bright monomeric green fluorescent protein derived from
Branchiostoma lanceolatum. *Nat. Methods* **10**, 407–9 (2013).

67. J.-D. Pédelacq, S. Cabantous, T. Tran, T. C. Terwilliger, G. S. Waldo,
Engineering and characterization of a superfolder green fluorescent protein. *Nat.*
Biotechnol. **24**, 79–88 (2006).

68. B. S. Chambers, *et al.*, DNA mismatch repair is required for the host innate
response and controls cellular fate after influenza virus infection. *Nat. Microbiol.* **4**,
1964–1977 (2019).

69. C. T. Workman, *et al.*, enoLOGOS: a versatile web tool for energy normalized
sequence logos. *Nucleic Acids Res.* **33**, W389–W392 (2005).

70. G. Wei, *et al.*, Genome-wide analysis of ETS-family DNA-binding in vitro and in
vivo. *EMBO J.* **29**, 2147–2160 (2010).

Acknowledgements

We would like to acknowledge Brook Heaton for generating the A549-CRISPR-SAM-

IFNrsp cell line. We acknowledge assistance from Mike Cook and the Duke Cancer

Institute Flow Cytometry Core. We thank Robert Lefkowitz and his laboratory for

assistance with and use of the ChemiDoc Imaging System. We also thank Ephraim Tsalik

and Micah McClain for helpful discussions and Ben Chambers, Stacy Webb, and other

members of the Heaton lab for critical reading of the manuscript. H.M.F. and A.T.H. were

supported by NIH training grant T32-CA009111. N.S.H. is supported by R01-HL142985,

R01-AI137031, and funding from the Defense Advanced Research Projects Agency's

(DARPA) PReemptive Expression of Protective Alleles and Response Elements

(PREPARE) program (Cooperative agreement #HR00111920008). The views, opinions

and/or findings expressed are those of the author and should not be interpreted as

representing the official views or policies of the U.S. Government.

Author Contributions

A.T.H. and N.S.H. designed the screen. A.T.H performed the screen experiments and

initial analysis. H.M.F. and N.S.H. designed the screen validation studies. H.M.F.

performed the screen hit determination analysis, flow cytometry, RNA and protein

quantification experiments, sequencing and promoter analysis, and virus infection

experiments. A.T.H. generated the Cal/09-sfGFP virus. H.M.F. and N.S.H. wrote the

manuscript.