Running head: Diversity-stability trophic cascade

Title: Predator complementarity dampens variability of phytoplankton biomass in a diversity-

stability trophic cascade

Authors: Chase J. Rakowski[1*], Caroline E. Farrior[1], Schonna R. Manning[2], Mathew A. Leibold[3]

[1]Department of Integrative Biology

University of Texas at Austin

Austin, Texas 78712

[2]Department of Molecular Biosciences

University of Texas at Austin

Austin, Texas 78712

[3]Department of Biology

University of Florida

Gainesville, Florida 32611

[*]Corresponding author: chaserak@utexas.edu

Abstract. Trophic cascades – indirect effects of predators that propagate down through food

webs – have been extensively documented. It has also been shown that predator diversity can

mediate these trophic cascades, and separately, that herbivore biomass can influence the stability

of primary producers. However, whether predator diversity can cause cascading effects on the

stability of lower trophic levels has not yet been studied. We conducted a laboratory microcosm

experiment and a field mesocosm experiment manipulating the presence and coexistence of two

heteropteran predators and measuring their effects on zooplankton herbivores and phytoplankton

basal resources. We predicted that if the predators partitioned their zooplankton prey, for

example by size, then co-presence of the predators would reduce zooplankton prey mass and lead

to 1) increased average values and 2) decreased temporal variability of phytoplankton basal

resources. We present evidence that the predators partitioned their zooplankton prey, leading to a

synergistic suppression of zooplankton; and that in turn, this suppression of zooplankton reduced

the variability of phytoplankton biomass. However, mean phytoplankton biomass was

unaffected. Our results demonstrate that predator diversity may indirectly stabilize basal resource

biomass via a "diversity-stability trophic cascade," seemingly dependent on predator

complementarity, but independent of a classic trophic cascade in which average biomass is

altered. Therefore predator diversity, especially if correlated with diversity of prey use, could

play a role in regulating ecosystem stability. Furthermore, this link between predator diversity

and producer stability has implications for potential biological control methods for improving the

reliability of crop yields.

Key words: *algae; biodiversity; biological control; ecosystem functioning; food web; niche*

complementarity; plankton; resource partitioning; species richness; stability; temporal

variability; top-down control.

INTRODUCTION

A substantial body of work, generally motivated by global biodiversity loss, indicates that

enhanced biodiversity of ecosystems often stabilizes community biomass (Jiang and Pu 2009,

Loreau and de Mazancourt 2013, Gross et al. 2014). Most of these studies measured or modeled

the effects of primary producer diversity on the variability of primary producer biomass (e.g.

Hector et al. 2010, Loreau and de Mazancourt 2013, Gross et al. 2014). However, predators face

greater extinction threats than do lower trophic levels, suggesting that predator diversity is more

relevant to global biodiversity change than is primary producer diversity (Purvis et al. 2000).

Furthermore, biodiversity often alters energy flow through food webs, and so manipulating

diversity and measuring its effects within a single trophic level can give an incomplete picture of

how biodiversity influences ecosystem functioning (Hines et al. 2015, Seabloom et al. 2017).

While some literature addresses how predator diversity affects the average biomass of various

other trophic groups in food webs (reviewed in Schmitz 2007), little is yet known about the

influence of predator diversity on ecosystem stability. In particular, the existence of a link

between predator diversity and the stability of non-adjacent lower trophic levels has not (to our

knowledge) been tested. Yet, such a link would have critical implications for both the

maintenance of stable natural ecosystems as predator diversity changes, and also for biological

control to potentially stabilize crop yields.

Existing research relating predator diversity to lower trophic level functioning has shown

that higher predator (or natural enemy) diversity generally leads to lower mean herbivore

densities and higher mean plant biomass, as long as the predators exhibit complementarity in

their feeding niches (Straub et al. 2008, Greenop et al. 2018). This research was mostly

performed in the context of biological pest control, but relies on general ecological principles.

Therefore, diversity of functional traits among predators related to prey use, such as body size,

may generally play a key role in mediating trophic cascade strength (Straub et al. 2008). But

besides average primary producer biomass, the temporal variability of primary producer biomass

could also be affected by functional predator diversity in a predictable way. While herbivory

generally decreases primary producer biomass, it has also been shown to increase the variability

of primary producer biomass (Thébault and Loreau 2005, Downing et al. 2014), perhaps

resulting from predator-prey cycling between the herbivores and producers (Volterra 1926);

though note that the opposite effect has also been found (Post 2013). Thus, if functional predator

diversity reduces herbivore biomass, it may also indirectly reduce the variability of primary

producer biomass, producing a "diversity-stability trophic cascade" (Table 1). Such an effect

would stem from two coupled relationships: 1) a negative relationship between predator diversity

and herbivore (prey) biomass, likely due to prey use complementarity, and 2) a positive

relationship between herbivore biomass and producer variability.

Here we report the results of a laboratory microcosm experiment and a field mesocosm

experiment to test for the existence of diversity-stability trophic cascades. After testing for prey

use complementarity by two predator species in the laboratory, we manipulate the presence of

the same predators in the field (no predators, each predator alone, and both predators) and

measure the resulting biomass of plankton groups as well as the variability of phytoplankton

biomass. We use the heteropterans *Notonecta* and *Neoplea* as the predators due to their

substantial difference in body size and consequent likelihood of partitioning prey resources.

Based on previous work (Murdoch et al. 1984) we predicted that *Notonecta* would mostly

consume *Daphnia*, and we predicted that *Neoplea* would consume smaller zooplankton based on

its smaller body size. We hypothesized that if *Notonecta* and *Neoplea* partition zooplankton prey,

then the combination of these predators would suppress zooplankton more than either predator

alone, leading to two indirect effects: 1) increased mean phytoplankton biomass (i.e., a diversity-

biomass trophic cascade, Table 1b), and 2) reduced variability of phytoplankton biomass (i.e., a

diversity-stability trophic cascade, Table 1c). We present evidence that these predators indeed

partitioned zooplankton prey, leading to a greater suppression of zooplankton when the predators

were together than predicted from their individual effects; and that this suppression of herbivores

indirectly stabilized, but did not enhance, phytoplankton biomass. In order words, we detected a

diversity-stability trophic cascade despite the absence of a diversity-biomass trophic cascade.

METHODS

Focal predators

We used two insect species in the suborder Heteroptera as the predators in both the field

and laboratory experiments: the notonectid *Notonecta undulata* and the pleid *Neoplea striola*.

Hereafter, we use the genus names (*Notonecta* and *Neoplea*) to refer to these two focal predator

species. Both are mobile generalist predators that are widespread across North America.

However, they differ starkly in size: *Notonecta* adults measure ~11-13 mm and *Neoplea* adults

measure ~1.5 mm in length. Studies have shown that while members of the genus *Notonecta* can

take prey as small as microscopic rotifers, they strongly reduce large prey such as *Daphnia* and

mosquito larvae (Leon 1998, Murdoch et al. 1984, Hampton and Gilbert 2001). The genus

Neoplea has been less studied; they have been documented to attack invertebrates ranging in size

from rotifers to *Daphnia* (Hampton and Gilbert 2001, Gittelman 1977), but we predicted they

would prefer smaller prey than *Notonecta*.

Organism collection

We allowed communities of phytoplankton to assemble naturally in six outdoor tanks at

the University of Texas' Brackenridge Field Laboratory, Austin, TX for ~six months. We then

mixed a common inoculum from these tanks. The phytoplankton community became dominated

by green algae (Chlorophyta), ranging in size from green picoplankton ~1 µm in diameter to

Oocystis with mother cell walls up to ~25 µm in diameter and dominated by a few

morphospecies, especially *Selenastrum* and *Oocystis* (Appendix S1: Table S1). We collected an

array of zooplankton taxa from small water bodies nearby, including many rotifer species,

Spirostomum, *Arctodiaptomus dorsalis*, *Mesocyclops edax*, and *Scapholeberis kingi*, and we

ordered *Daphnia magna* from Sachs Systems Aquaculture (St. Augustine, FL) to extend the size

range of the zooplankton to larger-bodied individuals (Appendix S1: Table S2). We similarly

mixed the zooplankton taxa together into a common inoculum. The predators *Notonecta* and

Neoplea were also collected from small water bodies in Austin, TX.

Laboratory experiment

To test whether *Notonecta* and *Neoplea* partitioned zooplankton prey resources, we

conducted a short time scale (5-day) experiment in the laboratory. Five days represents just

under one generation for the dominant zooplankton with the shortest generation times, *Daphnia*

magna and *Scapholeberis kingi*. Therefore we anticipated that five days would provide enough

time for the predators to reduce zooplankton populations but would not provide enough time for

the zooplankton populations to significantly recover, allowing us to better estimate the effects of

the predators on mortality of different zooplankton species while minimizing the influence of

zooplankton fecundity and plasticity. First, we combined portions of the phytoplankton and

zooplankton inocula into a common mixture. Then we placed ten *Notonecta* adults individually

in microcosms with 1.5 L of the plankton mixture, and likewise placed ten *Neoplea* adults

individually in microcosms with 100 mL of the plankton mixture. This 15:1 volume ratio

matched the ratio of prey mass consumed by the predator species in pilot trials, which we hoped

would maximize the chance of detecting prey reduction by both predators (too large a

zooplankton:predator ratio and the reduction may not be detected; too small a

zooplankton:predator ratio and the predator could quickly deplete the prey and hinder detection

of the effect size). We also established control microcosms of both sizes (1.5 L and 100 mL)

with no predator. To enable the estimation of overall predator effects over multiple densities (as

would be encountered in the field), we halved the zooplankton density in half of all microcosms

via filtration. We accomplished this with a 44-μm filter for the 100 mL microcosms, but in the

interest of time we used a 100-μm sieve to filter the 1.5 L microcosms. This meant the smallest

zooplankton (rotifers) were not reduced in the *Notonecta* microcosms; however, rotifers

comprised only 0.46% of zooplankton mass across all control microcosms. All treatment

combinations were replicated five times, yielding 40 total microcosms (2 predator

species/microcosm sizes × predator presence or absence × 2 zooplankton concentrations × 5

replicates). The microcosms were randomized and placed in an environmental chamber

maintained at 25 °C with fluorescent lights on a 16:8 h light:dark cycle. After five days, we

filtered the contents of each microcosm using a 44-μm filter and preserved them in 10% Lugol's

solution. We then estimated biomass of the zooplankton taxa as described in Appendix S2.

Field experiment

To evaluate the influence of predator diversity on phytoplankton biomass and stability,

we established replicate pond communities in 200-L cattle tanks at Brackenridge Field

Laboratory. Tanks were outfitted with float valves to maintain constant water levels and covered

in 1 mm^2 mesh screens to prevent immigration or emigration of macroorganisms. Before

beginning the experiment, we analyzed total N and P of the water following the methods of the

American Public Health Association (APHA 1989). We then supplemented $NaNO_3$ and

$NaH_2PO_4 \cdot H_2O$ to bring the total N and P to the concentrations found in COMBO medium (14

mg/L N and 1.55 mg/L P), a nutrient-rich medium commonly used for culturing plankton

(Kilham et al. 1998). Immediately following weekly sampling (methods described below) we

added both nutrient solutions to compensate for a 5% daily loss rate from the water column (as

per Hall et al. 2004), and stirred the water to replenish dissolved oxygen.

We distributed the phytoplankton inoculum equally among the tanks, allowed the

phytoplankton to grow for 15 days, and then distributed the zooplankton inoculum equally

among the experimental tanks in the same way. Finally, we added to the tanks either no insect

predators (controls), 6 adult *Notonecta*, 90 adult *Neoplea*, or half the density of each predator: 3

adult *Notonecta* with 45 adult *Neoplea*. The relative densities of *Notonecta* and *Neoplea* (1:15)

were chosen to satisfy the null hypothesis that each tank with predators would experience the

same total predation rate if the predators did not partition prey resources. In other words, we used

a substitutive design based on consumption rates rather than density *per se*. This predator ratio

was derived from the laboratory experiment, where individual *Notonecta* consumed 14.3× and

16.3× more animal mass than *Neoplea* in 0.5× and 1× zooplankton concentration microcosms,

respectively. Each treatment was replicated five times, for a total of 20 tanks in a randomized

design.

Beginning a week after adding predators, we sampled plankton weekly for six weeks. To

sample zooplankton, we used tube samplers to collect ~6 spatially-spread whole water column

subsamples and pool them into a 12-L sample for each tank. We filtered this sample through 65-

μm mesh, returned any predators to the tank, and preserved the retained material in 10% Lugol's

solution. To sample phytoplankton, we used 1-cm diameter PVC pipes (one per tank) to collect

three spatially-spread whole water column subsamples and pool them into a 50-mL sample for

each tank. We estimated biomass of zooplankton taxa as in the laboratory experiment, and

additionally estimated biovolume of phytoplankton taxa (methods in Appendix S2). At the end of

the experiment, we estimated the remaining density of predators by sampling them with a dipnet

until we returned three successive empty sweeps.

Grouping zooplankton for analysis

To test our prediction that *Notonecta* and *Neoplea* partition zooplankton prey, we divided

zooplankton into predicted trophic groups. We expected copepods to be the least vulnerable to

predation and therefore placed them in a separate group. Copepods, especially diaptomids (the

dominant copepod was the diaptomid *Arctodiaptomus dorsalis*), exhibit a faster escape response

and lower vulnerability to predators than other crustacean zooplankton (O'Brien 1979).

Copepods are also often omnivorous; in particular, research suggests the other copepod species,

Mesocyclops edax, is omnivorous as an adult (Adrian and Frost 1993), although diaptomid adults

can also be omnivorous (Williamson 1987). We split the remaining zooplankton into two other

groups, *Daphnia* and all other smaller zooplankton, based on our prediction that *Notonecta*

would selectively prey on *Daphnia* and *Neoplea* would prey on smaller zooplankton.

Data analysis

To check for differences in predator survival in the field experiment that might confound

the predator treatments, we used two-way ANOVA to compare predator survival rates by

species, diversity treatment, and their interaction.

We analyzed the effects of the predators on the biomass of zooplankton groups in the

laboratory experiment by fitting generalized linear models in the gamma family (gamma GLMs)

separately for each predator species and zooplankton group. We first fit the models with the

interaction between predator presence and zooplankton concentration (0.5× or 1×) to find

whether predator effects were influenced by zooplankton density. Then we fit the models

omitting the interaction, with only fixed effects of predator presence and zooplankton

concentration so that general predator effects would be assessed across zooplankton densities.

Zooplankton groups that were reduced by either predator were then combined as the collective

prey for analysis of a diversity effect on zooplankton prey mass in the field experiment. To this

end we used the *lme4* package (Bates et al. 2015) to fit a generalized linear mixed model in the

gamma family (gamma GLMM) with tank as a random effect to account for repeated measures,

and fixed effects for each predator and their interaction. Then we fit all simpler nested models

excluding the interaction term, and used likelihood ratio tests to find whether the predator

interaction or other terms significantly improved the model fit. If significant, a negative

interaction would indicate a synergistic zooplankton reduction while a positive interaction would

indicate disruptive effects between the predators. On the other hand, if the interaction did not

improve model fit, simple additive predator effects would be supported, if significant. We used

the same approach to assess the effect of the predators on the Shannon-Weaver diversity of

zooplankton and on mean phytoplankton biovolume. To analyze phytoplankton variability, we

calculated the temporal coefficient of variation (CV) of phytoplankton biovolume for each

mesocosm. We then repeated the same approach described above to analyze the predator

diversity effect, with the exception that no random effect was included in the nested models (i.e.,

with GLMs rather than GLMMs), as there is only a single CV for each mesocosm. We repeated

the phytoplankton analyses excluding *Oocystis*, the largest and likely least edible genus of

phytoplankton.

Lastly, we used piecewise structural equation modeling (piecewise SEM) with the

piecewiseSEM package (Lefcheck 2016) to jointly analyze relationships among all trophic

groups. We scaled all variables by their root-mean-square, and used gamma GLMs as the

component models. We forced zooplankton to affect phytoplankton CV via its two components,

the mean and SD, to lend more insight into the mechanism of any stability effect. Since mean

and SD are summary variables (one value per tank), we used time-averaged values of the other

variables (zooplankton and predators) to avoid unbalancing sample sizes. We first fit an SEM

aligning with the diversity effect analysis above, with collective zooplankton prey combined into

one variable and the predator interaction as a predictor of this variable in addition to the

individual predators. We then fit a separate SEM with the zooplankton groups separated as in the

laboratory analysis, and with no predator interaction. Since adult copepods are considered

omnivorous, we first fit the SEM with a pathway from copepods to both the smaller zooplankton

and to phytoplankton. To be conservative, we also included *Spirostomum* in the herbivorous

smaller zooplankton despite characterization in the literature as a bacterivore, since it may have

consumed small phytoplankton. When tests of directed separation were asymmetrical we used

the direction with the more conservative P value. All analyses were performed in R v. 3.5.3 (R

Core Team 2017).

RESULTS

Predator populations and plankton community structure

Neither predator reproduced during the field experiment. The median predator survival

rate was 79% (SD = 17%), with no significant difference between species ($F_{1,16} = 0.377$, $P =$

0.548), diversity treatment ($F_{1,16} = 0.225$, $P = 0.642$), or their interaction ($F_{1,16} = 0.003$, $P =$

0.956). A large amount of microalgae settled on the bottom of most field tanks, but there was

little algal growth on tank walls. The zooplankton composition in the laboratory differed from

the average composition in the field; however, the largest difference was in the relative

abundance of the copepod *Arctodiaptomus* (median of 10% of zooplankton biomass without

predators in the laboratory versus 56% in the field) which was not important for the experiment

(see results below). *Daphnia* and the smaller zooplankton comprised a median of 66% and 23%

of zooplankton mass in the absence of predators in the laboratory, respectively, versus 27% and

17% in the field, respectively. In both experiments, the smaller zooplankton were dominated by

Scapholeberis kingi, and all zooplankton species were shared across the experiments except for a

few rare rotifers found only in the field experiment (Appendix S1: Table S2). We did not find

evidence for any zooplankton colonization events in the field with the potential to affect the

results. The most likely colonizers, rotifers, comprised only 0.2% of zooplankton biomass on

average in field tanks.

Effects of predators on plankton groups

In the short-term laboratory experiment, the predators reduced opposing zooplankton

groups. *Notonecta* reduced *Daphnia* biomass by 97% averaged over both zooplankton

concentrations, without affecting the smaller zooplankton. On the other hand, *Neoplea* reduced

smaller zooplankton biomass by 43.0 µg or 52% with the 1× concentration and 46.2 µg or 90%

with the 0.5× concentration, without affecting *Daphnia* biomass (Appendix S3 Table S1).

Neither predator affected copepod biomass (Fig. 1a). In the field experiment, the predators

caused a more than additive reduction of their collective prey (non-copepod zooplankton) when

together (Fig. 1b). This is supported by the negative interactive effect of the predators on their

collective prey which significantly improved the GLMM (Table 2a, Table 3a). Despite our

halving of predator population sizes in tanks with both predators, median zooplankton prey

biomass was reduced by 54% with both predators compared to *Notonecta*-only tanks and by 93%

compared to *Neoplea*-only tanks. The two predators had additive negative effects on zooplankton

Shannon-Weaver diversity (Appendix S3 Table S2, Table S3).

Mean phytoplankton biovolume was not affected by the predators (Fig. 2a, Table 2b,

Table 3b). Yet, analogous to the zooplankton prey, the CV of phytoplankton biovolume was

synergistically reduced by the predator species (Fig. 2b, Table 2c, Table 3c). Compared to no-

predator tanks, phytoplankton CV was 17% lower with *Notonecta*, 34% higher with *Neoplea*,

and 52% lower with both predators. Removing the least edible phytoplankter, *Oocystis*, from

analysis resulted in a stronger stabilization effect, with phytoplankton CV 20% lower with

Notonecta, 2% lower with *Neoplea*, and 72% lower with both predators compared to no-predator

tanks (Appendix S3 Figure S1, Table S4, Table S5).

SEMs

The piecewise SEM including the predator interaction shows that the cascading diversity-

stability effect was mediated by the zooplankton prey and by mean phytoplankton biovolume

(Fig. 3). The predators had a negative synergistic (interactive) effect on zooplankton prey, which

in turn had a negative effect on the mean, but not the SD, of phytoplankton biovolume. Mean

phytoplankton negatively affected phytoplankton CV, as the mean is the denominator of the CV.

These three successive negative effects (*Notonecta* × *Neoplea* → non-copepod zooplankton →

mean phytoplankton → phytoplankton CV) amount to an indirect negative effect of the predator

combination on phytoplankton CV (the product of three negatives is a negative).

With non-copepod zooplankton separated into *Daphnia* and smaller zooplankton, and

with copepods considered as omnivores able to consume smaller zooplankton, SEM identified a

positive effect of copepods on smaller zooplankton (Appendix S3 Figure S2). Since we did not

have a mechanistic explanation for copepods to have a direct positive effect on smaller

zooplankton, we concluded that this effect was spurious and was indicative of a positive

correlation between the two groups; thus, the pathway from copepods to smaller zooplankton

was dropped. The resulting SEM (with copepods as herbivores) suggests that *Notonecta*

primarily reduced *Daphnia* while *Neoplea* primarily reduced the smaller zooplankton, although

Notonecta also reduced the smaller zooplankton to a lesser extent (Fig. 4). Reflecting the non-

copepod zooplankton in the previous SEM, it is apparent that *Daphnia* reduced mean

phytoplankton. Departing from the previous SEM, however, *Daphnia* also had a weak negative

effect on phytoplankton SD, which positively affects phytoplankton CV by definition as its

numerator. This negative effect on SD was apparently offset, though, by a slightly stronger

positive effect of smaller zooplankton on phytoplankton SD. Thus, the net negative diversity-

variability effect as demonstrated in Figure 3 is mirrored in Figure 4 via the pathway *Notonecta*

→ *Daphnia* → mean phytoplankton → phytoplankton CV. However, the separation of *Daphnia*

from smaller zooplankton also revealed a separate pathway contributing to the diversity-stability

effect: *Neoplea* → smaller zooplankton → phytoplankton SD → phytoplankton CV, a negative

effect followed by two positive effects which amounts to a net negative effect of *Neoplea* on

phytoplankton CV.

DISCUSSION

Our results indicate that the two focal predator species, *Notonecta* and *Neoplea*,

partitioned zooplankton prey, leading to a synergistic suppression of zooplankton that stabilized

phytoplankton biomass. As revealed by SEM, the stability effect was mediated on the whole by a

change in mean phytoplankton biomass without an accompanying proportional change in

standard deviation as would normally be predicted by Taylor's law (Fig. 3; Taylor and Woiwod

1980). Yet, mean phytoplankton biomass was not significantly affected by predator diversity.

While these results might seem contradictory on the surface, they can both be correct because

SEM lends insight only into likely causal pathways, not into differences among treatments.

Given a longer time frame, the predators likely would have reproduced and experienced

additional mortality, causing further changes in plankton biomass and stability. However, it is

difficult to predict whether such changes would have influenced our conclusions given the dearth

of long-term studies integrating biodiversity-ecosystem functioning with food web ecology.

Looking more closely, both predators contributed to the stability effect, despite the

stronger overall effect of *Notonecta*. While *Notonecta* lowered phytoplankton CV by nearly

eliminating *Daphnia*'s reduction of mean phytoplankton, *Neoplea* lowered phytoplankton CV by

suppressing the increase of phytoplankton SD by smaller zooplankton (Fig. 4). Our data cannot

answer why *Daphnia* primarily affected mean phytoplankton while smaller zooplankton only

affected its temporal variation. However, this difference may be explained by differences in the

behavior of these two zooplankton groups. *Scapholeberis kingi*, which comprised 80.4% of

smaller zooplankton biomass in field tanks, dwells and feeds almost entirely at the water surface,

while *Daphnia* feeds throughout the water column. Arditi and colleagues (1991) demonstrated

that this difference causes them to have drastically different effects on phytoplankton, with

Scapholeberis clearing phytoplankton more slowly than *Daphnia* as the water column remains a

refuge from *Scapholeberis* but not from *Daphnia*. While the effect of *Scapholeberis* may not

have been strong enough to reduce mean phytoplankton, it may have been enough to create a

degree of predator-prey cycling, inducing the phytoplankton to fluctuate as predicted by theory

(Volterra 1926, Thébault and Loreau 2005). The predators also both affected zooplankton

Shannon-Weaver diversity, but we do not believe this mediated the stability cascade. The

predators' effects on zooplankton diversity were weaker than the effects on zooplankton mass,

they were additive rather than synergistic, and they were negative. We would expect such a

decline in herbivore diversity to have a de-stabilizing or neutral effect on phytoplankton, not a

stabilizing effect (Valone and Balaban-Feld 2019).

Synergistic prey suppression by predator species can be attributed to complementarity or

facilitation (Jonsson et al. 2017). Since herbivorous zooplankton compete for phytoplankton

resources, predation of only a subset of the herbivores could allow the unaffected herbivores to

take advantage of the freed resources and increase their biomass, compensating for the loss of the

vulnerable herbivores. In this scenario, if predators individually target a subset of herbivores but

the full complement of predators suppresses all herbivores simultaneously, the combination of

predators would lead to a synergistic reduction of the herbivores. This scenario aligns with our

experiment. It is also possible that the predators facilitated each other, for example if the

behavior of one predator disturbed the other predator's prey from a refuge, thereby increasing its

vulnerability to the second predator (Culshaw-Maurer et al. 2020). The lack of refuges in the

tanks makes this scenario less likely, however.

The cascading diversity effect was not reflected in all tanks or plankton taxa (i.e.,

copepods), but the observed pattern was consistent with background variation in the plankton

community. In at least one mesocosm per treatment, variability of phytoplankton biomass was at

least as low as the average variability with both predators (Fig. 2b). This pattern appears to stem

from the large background variation in the system. Even in the absence of predators, variation in

the density of *Daphnia* and of the smaller zooplankton was very large, with some predator-free

control tanks having consistently low zooplankton densities (Fig. 1b). This meant that only some

tanks within each treatment contained enough zooplankton for a cascading predator effect to be

detected. Regardless of zooplankton abundance, copepods did not play a part in the cascading

effect. Copepods are known to be significantly less vulnerable to predation than other crustacean

zooplankton (O'Brien 1979); but why they had little effect on phytoplankton or smaller

zooplankton is less clear. It is possible they consumed settled algae, which was not sampled.

Similarly, removal of the likely least edible phytoplankton taxon, *Oocystis*, from analysis

revealed a stronger stabilization effect on the remaining phytoplankton (Appendix S3 Figure S1,

Table S4, Table S5). The fact that less vulnerable taxa played less of a role in the cascading

effect adds another line of evidence for the hypothesized consumption-mediated effect.

This study provides a simultaneous evaluation of effects of predator diversity on both the

average and variability of primary producer biomass. Most previous predator diversity-

ecosystem function experiments measured the mean but not the variability of trophic group

biomass as dependent variables (Bruno and O'Connor 2005; Straub et al. 2008). Those

experiments indicate that if predators partition their resources, increasing predator diversity can

suppress herbivores and lead to higher mean autotroph biomass, i.e., a diversity-biomass trophic

cascade (Table 1; Straub et al. 2008). In our field experiment, increasing predator diversity

reduced zooplankton prey but did not significantly increase mean phytoplankton biomass. The

SEM suggests that only *Daphnia* impacted mean phytoplankton biomass, not the smaller

zooplankton (Fig. 4); overall, the effect of the zooplankton on mean phytoplankton biomass was

apparently too weak to complete a diversity-biomass trophic cascade. It is not uncommon for

herbivores to have weak effects on plant biomass (Maron and Crone 2006). The strength of

herbivory is often dampened by the variable food quality of plants, low encounter rates between

herbivores and plants, indirect effects, or other factors (Leibold 1989, Borer et al. 2005, Maron

and Crone 2006). In this case, the temporal variability of producer biomass was more sensitive to

changes in zooplankton biomass, and therefore to predator presence and composition, than was

average producer biomass. Future studies will need to explore the generality of this result.

Previous studies have reported the effects of natural enemy diversity on the variability of

food web interactions, and some have described theoretical mechanisms relevant to these studies.

A few studies used surveys to relate parasitoid richness to the temporal variability of aggregate

parasitism rates, finding either no relationship (Rodriguez and Hawkins 2000) or a negative

relationship (Tylianakis et al. 2006, Macfadyen et al. 2011). Griffin and Silliman (2011) found

that the combination of two predators which exhibited temporal complementarity in attack rates

reduced the temporal variability of the total predation rate on a shared prey. Directly relevant

theoretical studies are limited, but theory suggests several mechanisms linking diversity and

stability in ecosystems generally (Loreau and de Mazancourt 2013). With a two-trophic level

model, Thébault and Loreau (2005) showed that decreasing herbivore biomass stabilizes and

increases plant biomass. Coupling this finding with the consensus from the literature that

complementarity of prey use by predators tends to reduce prey biomass, it follows that predator

complementarity may indirectly stabilize (and increase) plant biomass. This food web diversity-

stability mechanism could have important implications for ecosystem management and for

biological control.

Our results suggest that adding multiple natural enemies to an agro-ecosystem could

stabilize fluctuations in crop yields, provided the natural enemies partition pest resources. While

a goal of farmers will always clearly be to achieve as high an average yield as possible,

achieving consistent yields can be equally important. Thus our study adds another argument for

"conservation biological control," or natural pest control achieved through the conservation of

natural enemy biodiversity (Snyder 2019). We showed that diversity of a predator trait known to

correlate with prey preference, body size, can be a key factor leading to the cascading stabilizing

effect on phytoplankton. Microalgae, especially phytoplankton, are increasingly cultured as a

crop for a variety of purposes, from biomass production for biofuels or animal feed, to nutrient

removal from wastewaters. However, algal cultivation is not yet widely practiced at the

commercial scale, in large part because it has proven too difficult to achieve consistently high

algal yields as algae ponds are easily colonized by various herbivorous zooplankton that are

difficult or costly to control with mechanical and chemical means (Smith and Crews 2014,

Montemezzani et al. 2015). Adding a functionally diverse array of predators to algal cultivation

ponds may therefore be a feasible, economical, self-sustaining way to encourage more reliable

algal yields. Terrestrial crops also can suffer yield instability associated with multiple pests, and

thus may similarly benefit from addition or encouragement of a community of natural enemy

species varying in size (Gurr et al. 2012). Diversity of other functional traits related to prey

choice may also encourage crop yield stability in a similar fashion and could also be managed in

natural enemy communities when relevant information is available, including microhabitat

preference, temporal patterns in predation strength, or variance in mode of prey suppression. For

example, top-down control can be stabilized by adding or encouraging some predators more

active in warm temperatures and some more active in cold temperatures (Griffin and Silliman

2011), or both parasites and predators (Ong and Vandermeer 2015), encouraging consistently

low herbivore densities and stable crop yields. While our study emphasizes the importance of

complementarity, redundancy is also likely important with increasing temporal and spatial scales

to guard against periods of weakened control by, or extirpations of, natural enemies at certain

times or locations (Peralta et al. 2014).

Here we have demonstrated that higher predator diversity, and accompanying

complementarity of prey use, can cause a chain of effects that cascade down a food web to

stabilize the temporal dynamics of basal resource biomass. This connection between predator

diversity and primary producer stability is an important step towards joining biodiversity-

ecosystem function theory with food web theory, as biodiversity and ecosystem functioning

research has only recently begun incorporating energy flows in a food web context (Barnes et al.

2018). Future work is needed to further explore the simultaneous influence of predator diversity,

in its various forms, on both the average and stability of community- and ecosystem-level

functioning. It is important to uncover whether stability is generally more sensitive to cascading

food web effects than is average biomass, or how the two effects are related in different contexts.

Predator diversity may also have different effects on other measures of stability, such as

resilience. Developing this field at the intersection of biodiversity-ecosystem functioning and

food web ecology not only will improve our understanding of the functioning of natural

ecosystems and their vulnerability to biodiversity change, but also will provide information that

can be used to manage for more reliable crop production and other ecosystem services.

ACKNOWLEDGEMENTS

Thanks to L. A. Sekula and J. Earwood for help processing samples, to R. Deans for help with

insect collection and experiment setup, to S. Duchicela for help with experimental setup, to D.

Correa for help with nutrient analysis, and to D. Nobles for providing equipment. Thanks to A.

Wolf, R. Decker, D. Cinoglu, S. Ortiz, E. Francis, D. Grobert, and A. Northup for feedback on

an earlier version of the manuscript. This research was supported by the Department of

Integrative Biology at the University of Texas at Austin and was made possible by the facilities

at Brackenridge Field Laboratory.

LITERATURE CITED

Adrian, R., and T. M. Frost. 1993. Omnivory in cyclopoid copepods: comparisons of algae and

invertebrates as food for three, differently sized species. Journal of Plankton Research

15:643–658.

APHA. 1989. Standard methods for the examination of water and wastewater, 17[th] edn.

American Public Health Association, Washington, D.C.

Arditi, R., N. Perrin, and H. Saïah. 1991. Functional responses and heterogeneities: an

experimental test with cladocerans. Oikos 60:69–75.

Barnes, A. D., M. Jochum, J. S. Lefcheck, N. Eisenhauer, C. Scherber, M. I. O'Connor, P. de

Ruiter, and U. Brose. 2018. Energy flux: the link between multitrophic biodiversity and

ecosystem functioning. Trends in Ecology and Evolution 33:186–197.

Bates, D., M. Maechler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models

using lme4. Journal of Statistical Software 67:1–48.

Borer, E. T., E. W. Seabloom, J. B. Shurin, K. E. Anderson, C. A. Blanchette, B. Broitman, S. D.

Cooper, and B. S. Halpern. 2005. What determines the strength of a trophic cascade?

Ecology 86:528–537.

Bruno, J. F., and M. I. O'Connor. 2005. Cascading effects of predator diversity and omnivory in

a marine food web: cascading effects of predator diversity. Ecology Letters 8:1048–1056.

Culshaw-Maurer, M., A. Sih, and J. A. Rosenheim. 2020. Bugs scaring bugs: enemy-risk effects

in biological control systems. Ecology Letters 23:1693–1714.

Downing, A. L., B. L. Brown, and M. A. Leibold. 2014. Multiple diversity-stability mechanisms

enhance population and community stability in aquatic food webs. Ecology 95:173–184.

Gittelman, S. H. 1977. Leg segment proportions, predatory strategy and growth in

backswimmers (Hemiptera: Pleidae, Notonectidae). Journal of the Kansas Entomological

Society 50:161–171.

Greenop, A., B. A. Woodcock, A. Wilby, S. M. Cook, and R. F. Pywell. 2018. Functional

diversity positively affects prey suppression by invertebrate predators: a meta-analysis.

Ecology 99:1771–1782.

Griffin, J. N., and B. R. Silliman. 2011. Predator diversity stabilizes and strengthens trophic

control of a keystone grazer. Biology Letters 7:79–82.

Gross, K., B. J. Cardinale, J. W. Fox, A. Gonzalez, M. Loreau, H. Wayne Polley, P. B. Reich,

and J. van Ruijven. 2014. Species richness and the temporal stability of biomass production:

a new analysis of recent biodiversity experiments. The American Naturalist 183:1–12.

Gurr, G. M., S. D. Wratten, and W. E. Snyder. 2012. Biodiversity and insect pests: key issues for

sustainable management. John Wiley and Sons, Inc., Hoboken, NJ.

Hall, S. R., M. A. Leibold, D. A. Lytle, and V. H. Smith. 2004. Stoichiometry and planktonic

grazer composition over gradients of light, nutrients, and predation risk. Ecology 85:2291–

2301.

Hampton, S. E., and J. J. Gilbert. 2001. Observations of insect predation on rotifers. Pages 115–

121 *in* L. Sanoamuang, H. Segers, R. J. Shiel, and R. D. Gulati, editors. Rotifera IX.

Springer, Dordrecht, Netherlands.

Hector, A., Y. Hautier, P. Saner, L. Wacker, R. Bagchi, J. Joshi, M. Scherer-Lorenzen, E. M.

Spehn, E. Bazeley-White, M. Weilenmann, M. C. Caldeira, P. G. Dimitrakopoulos, J. A.

Finn, K. Huss-Danell, A. Jumpponen, C. P. H. Mulder, C. Palmborg, J. S. Pereira, A. S. D.

Siamantziouras, A. C. Terry, A. Y. Troumbis, B. Schmid, and M. Loreau. 2010. General

stabilizing effects of plant diversity on grassland productivity through population asynchrony

and overyielding. Ecology 91:2213–2220.

Hines, J., W. H. van der Putten, G. B. De Deyn, C. Wagg, W. Voigt, C. Mulder, W. W. Weisser,

J. Engel, C. Melian, S. Scheu, K. Birkhofer, A. Ebeling, C. Scherber, and N. Eisenhauer.

2015. Towards an integration of biodiversity-ecosystem functioning and food web theory to

evaluate relationships between multiple ecosystem services. Advances in

Ecological Research 53:161–199.

Jiang, L., and Z. Pu. 2009. Different effects of species diversity on temporal stability in

single☐trophic and multitrophic communities. The American Naturalist 174:651–659.

Jonsson, M., R. Kaartinen, and C. S. Straub. 2017. Relationships between natural enemy

diversity and biological control. Current Opinion in Insect Science 20:1–6.

Kilham, S. S., D. A. Kreeger, S. G. Lynn, C. E. Goulden, and L. Herrera. 1998. COMBO: a

defined freshwater culture medium for algae and zooplankton. Hydrobiologia 377:147–159.

Lefcheck, J. S. 2016. PiecewiseSEM: piecewise structural equation modelling in R for ecology,

evolution, and systematics. Methods in Ecology and Evolution 7:573–579.

Leibold, M. A. 1989. Resource edibility and the effects of predators and productivity on the

outcome of trophic interactions. The American Naturalist 134:922–949.

Leon, B. 1998. Influence of the predatory backswimmer, *Notonecta maculata*, on invertebrate

community structure. Ecological Entomology 23:246–252.

Loreau, M., and C. de Mazancourt. 2013. Biodiversity and ecosystem stability: a synthesis of

underlying mechanisms. Ecology Letters 16:106–115.

Macfadyen, S., P. G. Craze, A. Polaszek, K. van Achterberg, and J. Memmott. 2011. Parasitoid

diversity reduces the variability in pest control services across time on farms. Proceedings of

the Royal Society B 278:3387–3394.

Maron, J. L., and E. Crone. 2006. Herbivory: effects on plant abundance, distribution and

population growth. Proceedings of the Royal Society B 273:2575–2584.

Montemezzani, V., I. C. Duggan, I. D. Hogg, and R. J. Craggs. 2015. A review of potential

methods for zooplankton control in wastewater treatment high rate algal ponds and algal

production raceways. Algal Research 11:211–226.

Murdoch, W. W., M. A. Scott, and P. Ebsworth. 1984. Effects of the general predator, *Notonecta*

(Hemiptera) upon a freshwater community. The Journal of Animal Ecology 53:791–808.

O'Brien, W. J. 1979. The predator-prey interaction of planktivorous fish and zooplankton: recent

research with planktivorous fish and their zooplankton prey shows the evolutionary thrust

and parry of the predator-prey relationship. American Scientist 67:572–581.

Ong, T. W. Y., and J. H. Vandermeer. 2015. Coupling unstable agents in biological control.

Nature Communications 6:5991.

Peralta, G., C. M. Frost, T. A. Rand, R. K. Didham, and J. M. Tylianakis. 2014.

Complementarity and redundancy of interactions enhance attack rates and spatial stability in

host-parasitoid food webs. Ecology 95:1888–1896.

Post, E. 2013. Erosion of community diversity and stability by herbivore removal under

warming. Proceedings of the Royal Society B 280:20122722.

543 Purvis, A., J. L. Gittleman, G. Cowlishaw, and G. M. Mace. 2000. Predicting extinction risk in

544   declining species. Proceedings of the Royal Society B 267:1947–1952.

545 R Core Team. 2017. R: a language and environment for statistical computing. R Foundation

546   for Statistical Computing, Vienna, Austria.

547 Ripple, W. J., J. A. Estes, O. J. Schmitz, V. Constant, M. J. Kaylor, A. Lenz, J. L. Motley, K. E.

548   Self, D. S. Taylor, and C. Wolf. 2016. What is a trophic cascade? Trends in Ecology &

549   Evolution 31:842–849.

550 Rodriguez, M. A., and B. A. Hawkins. 2000. Diversity, function and stability in parasitoid

551   communities. Ecology Letters 3:35–40.

552 Schmitz, O. J. 2007. Predator diversity and trophic interactions. Ecology 88:2415–2426.

553 Seabloom, E. W., L. Kinkel, E. T. Borer, Y. Hautier, R. A. Montgomery, and D. Tilman. 2017.

554   Food webs obscure the strength of plant diversity effects on primary productivity. Ecology

555   Letters 20:505–512.

556 Smith, V. H., and T. Crews. 2014. Applying ecological principles of crop cultivation in large-

557   scale algal biomass production. Algal Research 4:23–34.

558 Snyder, W. E. 2019. Give predators a complement: conserving natural enemy biodiversity to

559   improve biocontrol. Biological Control 135:73–82.

560 Straub, C. S., D. L. Finke, and W. E. Snyder. 2008. Are the conservation of natural enemy

561   biodiversity and biological control compatible goals? Biological Control 45:225–237.

562 Taylor, L. R., and I. P. Woiwod. 1980. Temporal stability as a density-dependent species

563   characteristic. Journal of Animal Ecology 49:209–224.

564 Thébault, E., and M. Loreau. 2005. Trophic interactions and the relationship between species

565   diversity and ecosystem stability. The American Naturalist 166:E95–E114.

Tylianakis, J. M., T. Tscharntke, and A.-M. Klein. 2006. Diversity, ecosystem function, and

stability of parasitoid host interactions across a tropical habitat gradient. Ecology 87:3047–

3057.

Valone, T. J., and J. Balaban-Feld. 2019. An experimental investigation of top-down effects of

consumer diversity on producer temporal stability. Journal of Ecology 107:806–813.

Volterra, V. 1926. Fluctuations in the abundance of a species considered mathematically. *Nature*

118:558–560.

Williamson, C. E. 1987. Predator-prey interactions between omnivorous diaptomid copepods and

rotifers: the role of prey morphology and behavior. Limnology and Oceanography 32:167–

177.

Table 1. Trophic cascade terminology used in this paper.

Term	Definition
trophic cascade	"Indirect species interactions that originate with predators and spread downward through food webs" (Ripple et al. 2016).
diversity-biomass trophic cascade	A change in *biomass* of a lower trophic level indirectly caused by a change in predator *diversity* and mediated by top-down effects on the intermediate trophic level(s).
diversity-stability trophic cascade	A change in the *stability* (defined in this paper as temporal variability but could be defined otherwise) of the biomass of a lower trophic level, indirectly caused by a change in predator *diversity* and mediated by top-down effects on the intermediate trophic level(s).

578   **Table 2.** Results of likelihood ratio tests comparing nested models, including the degrees of

579   freedom (Df), deviance (inverse goodness of fit), and *P* value: **a)** GLMMs for non-copepod

580   zooplankton biomass, **b)** GLMMs for phytoplankton biovolume, and **c)** GLMs for CV of

581   phytoplankton biovolume. *Notonecta × Neoplea* is the interaction model with main effects.

Model	Df	Deviance	*P*
a) non-copepod zooplankton			
random intercepts only (null)	3	1263.0	
Notonecta	4	1254.2	**0.0029**
Neoplea	4	1262.4	1.0000
Notonecta + Neoplea	5	1253.8	**0.0033**
Notonecta × Neoplea	6	1248.1	**0.0168**
b) phytoplankton (mean)			
random intercepts only (null)	3	2877.1	
Notonecta	4	2877.1	0.8198
Neoplea	4	2877.1	1.0000
Notonecta + Neoplea	5	2877.0	0.7697
Notonecta × Neoplea	6	2875.9	0.3004
c) CV of phytoplankton			
intercept only (null)	19	10.290	
Notonecta	18	9.572	0.2238
Neoplea	18	9.974	1.0000
Notonecta + Neoplea	17	9.548	0.3487
Notonecta × Neoplea	16	7.662	**0.0486**

582   **Table 3.** Coefficient estimates and standard errors (SE) for models analyzing effects of the

583   predators and their interaction on **a)** non-copepod zooplankton biomass (GLMM), **b)**

584   phytoplankton biovolume (GLMM), and **c)** temporal CV of phytoplankton biovolume (GLM).

Parameter	Estimate	SE
a) non-copepod zooplankton		
intercept	7.7513	0.5880
Notonecta	-0.4006	0.1385
Neoplea	0.0005	0.0092
Notonecta × Neoplea	-0.0137	0.0053
b) phytoplankton (mean)		
intercept	10.8908	0.3917
Notonecta	-0.0012	0.0923
Neoplea	-0.0007	0.0061
Notonecta × Neoplea	0.0037	0.0036
c) CV of phytoplankton		
intercept	-0.6181	0.3114
Notonecta	-0.0301	0.0734
Neoplea	0.0033	0.0049
Notonecta × Neoplea	-0.0059	0.0028

Figure legends

Figure 1. a) Multiplicative effects of predator addition on the biomass of zooplankton groups

over five days in the laboratory. Shapes are GLM coefficients with 95% confidence intervals.

The dashed line at 1 represents no effect, while the dashed line at 0 represents total elimination

of a zooplankton group. **b)** Zooplankton dry mass, excluding copepods (summed *Daphnia* and

smaller zooplankton), by predator treatment in the field experiment. Dots represent means +/- 1

temporal standard error for each mesocosm. Note the log scale. Also note that predator species

densities were halved in the "both" treatment (i.e., substitutive design).

Figure 2. a) Time series of mean phytoplankton biovolume, +/- 1 standard error, by predator

treatment. **b)** Temporal coefficient of variation (CV) of phytoplankton biovolume by predator

treatment (dots represent individual mesocosms).

Figure 3. Path diagram for the piecewise SEM with the predator interaction and zooplankton

divided into copepods and non-copepods. Red arrows represent negative effects and blue arrows

represent positive effects. Arrow thickness is proportional to the coefficient (effect size), which

is labeled next to the arrow followed by an indication of the effects' significance (*** $P<0.001$,

** $P<0.01$, * $P<0.05$, · $P<0.1$, n.s. $P>0.1$). The R^2 value of each component model is provided

below its response variable.

Figure 4. Path diagram for the piecewise SEM with zooplankton divided into three groups:

copepods, *Daphnia*, and smaller zooplankton. Red arrows represent negative effects and blue

arrows represent positive effects. Arrow thickness is proportional to the coefficient (effect size),

which is labeled next to the arrow followed by an indication of the effects' significance (***

$P<0.001$, ** $P<0.01$, * $P<0.05$, · $P<0.1$, n.s. $P>0.1$). Arrows are translucent when $P>0.05$. The

R^2 value of each component model is provided below its response variable.

Figure 1

     Figure 2

615 **Figure 3**

618   **Figure 4**