Running head: Diversity-stability trophic cascade

Title: Predator complementarity dampens variability of phytoplankton biomass in a diversity-

stability trophic cascade

Authors: Chase J. Rakowski[1] and Mathew A. Leibold[2]

Chase J. Rakowski

506 Patterson Laboratories

University of Texas

2415 Speedway Stop C0930

Austin, TX 78712

Mathew A. Leibold

Bartram 627, Department of Biology

University of Florida

Gainesville, FL 32611

[1] Corresponding author. chaserak@utexas.edu

[2] mleibold@ufl.edu

Abstract

The indirect effects of top predators on the biomass of lower trophic levels have been

extensively documented, especially in aquatic ecosystems and are commonly characterized as

"trophic cascades." There have also been studies showing that predator diversity can play an

important role in mediating these trophic cascades. In addition, some studies have demonstrated

that the diversity of consumers can impact the stability of their resources. However, the effects of

predator diversity on the stability of lower trophic levels (cascading effects of predator diversity)

have not yet been studied. We conducted a field pond mesocosm experiment and accompanying

laboratory microcosm experiment in which we grew phytoplankton with zooplankton herbivores,

manipulating the presence and co-habitation of two heteropteran predators while controlling for

total expected predation. We hypothesized that if the predators partitioned their herbivore prey,

for example by size, then co-presence of the predators would lead to 1) increased average values

and 2) decreased temporal variability of phytoplankton biomass. We present evidence that the

predators partitioned their herbivore prey, and found that this simultaneous suppression of

different herbivore groups reduced the variability of edible (smaller) phytoplankton biomass,

without affecting mean phytoplankton biomass. We also found that phytoplankton more resistant

to herbivory were not affected by our manipulations, indicating that the zooplankton herbivores

played an important role in mediating this cascading diversity-stability effect. Our results

demonstrate that predator diversity can indirectly stabilize primary producer biomass via a

"diversity-stability cascade" in the manner of a trophic cascade, seemingly dependent on

predator complementarity and the vulnerability of taxa to consumption, and independent of a

more traditional trophic cascade in which average biomass is enhanced. Predator diversity,

especially in terms of functional traits relating to differential prey use such as body size, may be

important for regulating ecosystem stability, and may provide a way to improve the reliability of

algal crop yields via biological control.

Key words: biodiversity, species richness, ecosystem functioning, stability, temporal variability,

resource partitioning, niche complementarity, food web, top-down control, plankton, algae,

biological control

Introduction

A substantial body of work indicates that enhanced biodiversity of ecosystems often

stabilizes community biomass (Jiang and Pu 2009, Cardinale et al. 2012, Loreau and de

Mazancourt 2013). Most of these studies were motivated by global biodiversity loss and

measured the effects of primary producer diversity on primary producer biomass (Cardinale et al.

2009). However, predators face greater extinction threats than lower trophic levels, suggesting

the implications of changing predator diversity are more relevant to global biodiversity change

(Purvis et al. 2000). While a moderate body of literature has asked how predator diversity affects

the average biomass of trophic groups (Schmitz 2007), ecologists have rarely experimentally

tested the influence of predator diversity on any form of ecosystem stability. Yet, predator

diversity-stability effects have important implications for both the maintenance of stable natural

ecosystems and also for biological control to potentially stabilize crop yields.

Much existing research on the effects of predator diversity on lower trophic levels has

been performed in the context of biological control. Biological control is classically practiced by

releasing a specialist natural enemy to target a specific pest (van Driesche et al. 2008). However,

this targeted method has a high failure rate, and in any case, agricultural plots are often affected

by multiple pest species (van Driesche et al. 2008). Increasingly, scientists have advocated the

use of multiple natural enemies as a way to improve pest control while conserving natural enemy

biodiversity, a concept termed 'conservation biological control' (Snyder 2019). A meta-analysis

showed that harnessing more diverse assemblages of natural enemies generally leads to lower

mean pest densities and higher mean crop yields, as long as the natural enemies exhibit

complementarity in their feeding niches (Straub et al. 2008). Following this reasoning, diversity

of functional traits among predators related to prey use, such as body size, may play a key role in

mediating the effects of predator diversity on ecosystem function (Straub et al. 2008). In theory,

decreases in herbivore mass have been shown to decrease the variability of primary producer

biomass (Thébault and Loreau 2005), suggesting that functional predator diversity may also

indirectly reduce the variability of primary producer biomass and thus produce a "diversity-

stability cascade." Such an effect may be most likely seen when the primary producers are highly

edible, such as in communities of small phytoplankton.

Culturing algae, especially phytoplankton, is a promising means of producing alternative

fertilizers, animal feeds, and fuels with a lower environmental impact than current industrial

standards (Benemann 2013). However, algal yields are often low and unpredictable, preventing

the production of algae-derived commodities from being economical (Benemann 2013). A major

reason for these algal crop failures is that algae ponds are quickly colonized by zooplankton

herbivores varying in size from ciliates <100 um in length to crustaceans such as *Daphnia* >2

mm in length, triggering reductions and oscillations in algal biomass (Smith et al. 2010).

Harnessing predator diversity may provide a useful means to control this range of aquatic "pests"

and thereby improve algal crop yields and their reliability. However, we are only aware of one

case in which biological control was tested to improve algal yields, and this study used a single

predator species (Sturm et al. 2012).

Here we report the results of a field mesocosm experiment and accompanying laboratory

microcosm experiment to test for the existence of diversity-stability cascades in which we

manipulate the presence of two predator species (no predators, each predator alone, and both

predators) and measure the resulting average biomass of herbivore groups as well as the average

and variability of biomass of phytoplankton groups. We use the heteropterans *Notonecta* and

Neoplea as the predators due to their substantial difference in body size and consequent

likelihood of partitioning prey resources. Based on previous work (Murdoch et al. 1984) we

predicted that *Notonecta* would consume *Daphnia*, and we predicted that *Neoplea* would

consume smaller zooplankton based on its smaller body size. We hypothesized that, given

Notonecta and *Neoplea* partition herbivore prey, the addition of both species together would 1)

increase the mean and 2) reduce the variability of phytoplankton biomass, while addition of a

single predator species would have either no such effect or much weaker effects. We present

evidence that these predators indeed partitioned herbivore prey, leading to an indirect effect on

phytoplankton stability without affecting mean phytoplankton biomass. While the unexpected

presence of a relatively inedible phytoplankton strain dampened the effect on total

phytoplankton, we show that predator diversity stabilized community biomass of smaller, more

edible phytoplankton and thus demonstrate a diversity-stability cascade.

Methods

Focal predators

We used two insect species in the suborder Heteroptera as the predators in both the field

and laboratory experiments: the notonectid (backswimmer) *Notonecta undulata* and the pleid

(pygmy backswimmer) *Neoplea striola*. Both species are mobile generalist predators and are

widespread across North America. Numerous studies have investigated the ecology of members

of the genus *Notonecta*, showing they strongly reduce large crustacean zooplankton such as

Daphnia and insects such as mosquito larvae, although they can take prey as small as

microscopic rotifers (Leon 1998, Murdoch et al. 1984, Hampton and Gilbert 2001). *N. undulata*

adults measure ~11-13 mm in length. In contrast, *N. striola* is much smaller, with adults

measuring ~1.5 mm. *N. striola* has been documented to prey on invertebrates ranging in size

from rotifers to *Daphnia* (Hampton and Gilbert 2001, Gittelman 1977). Hereafter, we use the

genus names (*Notonecta* and *Neoplea*) to refer to these two focal predator species.

Field experiment

We established replicate experimental pond communities in 200-L cattle tanks at

Brackenridge Field Laboratory, Austin, TX. Tanks were filled with well water and outfitted with

a float valve to maintain constant water levels. We analyzed total N and P of the water before

beginning the experiment following the methods of the APHA (1989). We added enough $NaNO_3$

and $NaH_2PO_4 \bullet H_2O$ to bring total N and P to the same concentrations as in COMBO medium (14

mg/L N and 1.55 mg/L P), a nutrient-rich medium commonly used for culturing plankton

(Kilham et al. 1998). Immediately following weekly sampling (methods described below) we

added enough of both nutrients to compensate for a 5% daily loss rate from the water column (as

per Hall et al. 2004).

We inoculated each tank with a common inoculum of phytoplankton collected from

nearby tanks which had been colonized by local phytoplankton strains for several months. The

phytoplankton community became dominated by green algae (Chlorophyta), ranging in size from

green picoplankton ~1 um in diameter to *Oocystis* with mother cell walls up to ~25 um in

diameter and dominated by a few morphospecies, especially *Selenastrum* and *Oocystis*

(Appendix 1). We allowed the phytoplankton to grow for 15 days before adding zooplankton.

We collected an array of zooplankton taxa from small water bodies nearby, including many

rotifer species, *Spirostomum*, *Arctodiaptomus dorsalis*, *Mesocyclops edax*, and *Scapholeberis*

kingi, and ordered *Daphnia magna* from Sachs Systems Aquaculture (St. Augustine, FL) to

extend the size range of the zooplankton to include larger-bodied individuals (Appendix 1). We

mixed the zooplankton taxa together and added an equal volume to each tank.

Notonecta and *Neoplea* were collected from ponds at the Austin Water Utility at Hornsby

Bend, Austin, TX and added after the zooplankton. We added either no insect predators

(controls), 6 adult *Notonecta*, 90 adult *Neoplea*, or 3 adult *Notonecta* with 45 adult *Neoplea* to

the tanks. Each treatment was replicated five times, for a total of 20 tanks in a randomized

design. The chosen relative densities of *Notonecta* and *Neoplea* (1:15) were determined to satisfy

the null hypothesis that each treatment with predators would experience the same total predation

rate if the predators did not partition prey resources. We determined this ratio by measuring the

dry masses of several adults of each taxon and then using the metabolic scaling law reported for

insects (Glazier 2010) to estimate the number of *Neoplea* that would be expected to consume the

same prey mass per day as 3 *Notonecta*. Additionally, in the laboratory experiment (methods

described below) individual *Notonecta* consumed 14.3x more animal mass than *Neoplea* in the

1x zooplankton concentration microcosms and 16.3x more animal mass than *Neoplea* in the 2x

zooplankton concentration microcosms, supporting the 15:1 ratio. Neither predator reproduced

during the experiment. Survival of both species was estimated by sampling tanks with a D-net,

and was estimated to be ~80% with no significant difference between the species or treatments.

Beginning a week after adding predators, we sampled zooplankton and phytoplankton

weekly for six weeks. To sample zooplankton, we used iterated spatially-spread subsamples to

collect 12L of water from each tank using tube samplers. We filtered this water using 65 μm

mesh, returned any predators to the tank, and preserved the retained material in 10% Lugol's

solution. We then identified and counted zooplankton in subsamples of this collection such that

for each taxon, at least 25 individuals or 10% of the sample was counted, whichever came first,

and at least 50 total individuals were counted. We measured the length of each crustacean and

Spirostomum with a micrometer to the nearest half increment (0.24 mm) and measured the length

and width of rotifers and width of *Spirostomum* to the nearest 0.05 increment (0.024 mm). We

converted crustacean length to biomass using length-mass regressions (Culver et al. [1985] for

copepods, McCauley [1984] for *Scapholeberis*, Dumont et al. [1975] for *Daphnia magna* and

Chydorus, and Anderson et al. [1998] for ostracods). When rotifers were identifiable to species,

we converted rotifer dimensions to biomass using species-specific equations from the EPA

protocol (EPA 2016); otherwise, we used biovolume equations (McCauley 1984) and converted

to biomass assuming a 10:1 biovolume:dry mass ratio. We converted *Spirostomum* dimensions to

biomass approximating cells as cylinders and assuming the same 10:1 biovolume:dry mass ratio.

To sample phytoplankton, we used iterated spatially-spread subsamples to collect 50 mL

water column samples from each tank using 1-cm diameter PVC pipes. After storing the

phytoplankton samples for ~24 hours in darkness on ice, we measured the absorbance of each

sample at 680 *in vivo* using a spectrophotometer. We then extracted chlorophyll-*a* and measured

chlorophyll-*a* fluorescence accounting for turbidity, following standard methods (APHA 1989).

We also preserved subsamples in 10% formaldehyde and enumerated phytoplankton densities

using a hemocytometer. We counted 25 nL or 50 cells of the most common morphospecies,

whichever came first, and counted at least 100 nL for the other taxa. Two individuals separately

counted each sample using these methods and then the counts were averaged. We also took

several micrographs of each morphospecies across tanks and dates and measured the cell

dimensions using ImageJ (Schneider et al. 2012), measuring >15 cells of all but the rarest

morphospecies. Then we used geometric approximations to calculate the biovolume of each cell

(Appendix 1).

193   ***Laboratory experiment***

194   We conducted an accompanying laboratory experiment to test whether *Notonecta* and

Neoplea had different effects on zooplankton herbivore groups on a short time scale. Following

completion of the field experiment, we collected plankton samples from several of the field

mesocosms and mixed them together. Half of the plankton mixture was filtered with a 100 μm

filter to concentrate the zooplankton to 2x ambient density. Additionally, ten adults of both insect

predator species were collected from the tanks. Each predator was placed individually in a

microcosm with either the 1x or 2x zooplankton density. *Notonecta* were placed in microcosms

filled with 1.5 L plankton mixture, and *Neoplea* were placed in microcosms with 100 mL

plankton mixture, such that the volumes were proportional to the 15:1 ratio of expected prey

consumption by the two species. Additionally, we established control microcosms with no

predator. All treatments were replicated five times, yielding 40 total microcosms (2 predator

species/microcosm sizes X predator presence or absence X 2 zooplankton concentrations X 5

replicates). The microcosms were randomized and placed in an environmental chamber

maintained at 25 C° with fluorescent lights on a 16:8 h light:dark cycle. After five days, we

filtered the contents of each microcosm using a 44 μm filter and preserved them in 10% Lugol's

solution. Five days represents just under one generation for the dominant zooplankton with the

shortest generation time, *Daphnia magna* and *Scapholeberis kingi*. Therefore we anticipated that

five days would provide enough time for the predators to impact their prey populations but

would not provide enough time for the zooplankton populations to significantly recover,

allowing us to better estimate the effects of the predators on mortality of different plankton

species while minimizing the influence of plankton fecundity and adaptation.

Grouping phytoplankton and zooplankton for analysis

An herbivorous zooplankter was defined as any taxon that, according to the literature and

our data, is primarily herbivorous over its life span and is not strictly benthic. Thus, *Mesocyclops*

was included in analyses even though the adult stage is omnivorous (Adrian and Frost 1993).

Spirostomum, a filter-feeding ciliate, was also included in the herbivorous zooplankton despite

characterization in the literature as a bacterivore, since preliminary analysis also suggested that it

reduced the biomass of smaller algae. However, the most dominant zooplankter in most tanks,

the diaptomid copepod *Arctodiaptomus dorsalis*, was not included. Diaptomid copepods exhibit

a much faster escape response and lower vulnerability to predators than other crustacean

zooplankton, and are often omnivorous (O'Brien 1979, Williamson 1987). Preliminary analysis

showed that *Arctodiaptomus* biomass was unaffected by either predator, and also had no effect

on algal biomass or stability, essentially acting as a bystander to the cascading food web effects.

Due to the difficulty of distinguishing *Arctodiaptomus* nauplii from nauplii of the other copepod

species, *Mesocyclops edax*, all nauplii were also excluded from analysis. *Arctodiaptomus*

copepodites and adults were 5.35 times as numerous as *Mesocyclops* copepodites and adults on

average, so biomass of *Mesocyclops* nauplii likely represented a small fraction of herbivore

biomass.

To test our hypothesis that *Notonecta* and *Neoplea* partitioned their resources based on

body size, leading to differential size-based cascading food web effects, it was necessary to

divide plankton into groups based on size. Based on our prediction that *Notonecta* would

selectively prey on *Daphnia* and *Neoplea* would prey on smaller zooplankton, we split

herbivorous zooplankton into *Daphnia* and all other herbivores (hereafter, "smaller herbivores")

for analysis. *Daphnia* was the largest and the second most dominant herbivore taxon, comprising

28% of herbivore biomass on average in control tanks, while the smaller herbivores comprised a

similar 26% of herbivore biomass in control tanks (*Arctodiaptomus* comprised 43%). The

smaller herbivores were dominated by *Scapholeberis kingi*, a hyponeustonic daphniid (Appendix

1). Zooplankton were grouped in the same way for analysis of the laboratory experiment. Since

the zooplankton used in the laboratory experiment were sampled from a subset of tanks after the

field experiment had ended, the zooplankton composition slightly differed from the average

composition in the field experiment. But as in the field experiment, *Scapholeberis* was by far the

most dominant species among the smaller herbivores; no new species appeared in the laboratory

experiment, and all the species found in the field experiment were included in the laboratory

experiment except for a few of the rarest rotifers (Appendix 1). In the predator-free laboratory

microcosms, *Daphnia* and the smaller herbivores composed 51% and 38% of the total herbivore

mass on average, respectively.

Phytoplankton were similarly grouped into two size classes representing the largest taxon

and all smaller taxa. Large phytoplankton are only vulnerable to large filter feeders (i.e.,

Daphnia). The largest phytoplankter was an *Oocystis* sp. (hereafter, *Oocystis* 1); all smaller

morphospecies were impacted more strongly by herbivores and were thus grouped together

(Appendix 1). We also ran analyses using alternative groupings including the two largest and the

three largest morphospecies.

Data analysis

We analyzed the effects of predator treatment on mean biomass of *Daphnia* and smaller

herbivores by fitting a gamma GLMM (generalized linear mixed model in the gamma family),

using a dummy variable for each predator addition treatment to compare against the no-predator

control treatment, and fit separately for each herbivore group. To analyze the effect of treatment

on zooplankton biomass in the laboratory experiment, we used gamma GLMs with zooplankton

concentration (1x or 2x) and predator presence as fixed effects, fit separately for each predator

species and zooplankton group (four models). To assess whether changes in zooplankton

biomass represented a mechanism mediating predation and algal stability, we tested the effects of

mean biomass of the two focal zooplankton groups on the CV of biovolume of the three algal

groupings. For this analysis we fit gamma GLMs with mean biomass of the two focal

zooplankton groups as the two fixed effects, again with a separate model for each algal grouping.

This analysis was repeated using the within-tank standard deviation of biomass of the

zooplankton groups, instead of mean biomass, as predictors.

To analyze algal stability, we calculated the temporal coefficient of variation (CV, a

standard measure of variability) of algal biovolume in each cattle tank over the course of the

experiment, measured separately for each of the three algal groupings (total, *Oocystis* 1, and all

but *Oocystis* 1). Then we used gamma GLMs to test whether predator treatment affected the CV

of algal biovolume, the same way we analyzed biomass of herbivore groups. To compare mean

algal biovolume by treatment, we used gamma GLMMs (generalized linear mixed models in the

gamma family) with a dummy variable for each predator addition treatment and tank as a

random effect, using the *lme4* package (Bates et al. 2015). Again, we tested each algal grouping

separately. In addition, we compared the CV and mean of algal biomass as estimated by

chlorophyll-*a* extraction and by absorbance at 750 nm and 680 nm in the same way we analyzed

algal biovolume; however, it is not possible to break these measurements down by algal taxon,

and so these proxies are comparable only to total algal biovolume. All data analysis was

conducted using version 3.5.3 of the R software package (R Core Team 2017).

Results

While the addition of one or the other heteropteran predator by themselves did not

significantly affect the variability of phytoplankton biomass, addition of both predators stabilized

the community biomass of edible phytoplankton by reducing the biomass of both the largest

herbivore, *Daphnia*, and of the smaller herbivores. In the short-term laboratory experiment, the

predators reduced opposing herbivore groups: *Notonecta* reduced *Daphnia* biomass (GLM,

P<0.001) by 97.1% averaged over both zooplankton concentrations without affecting the smaller

herbivores, while *Neoplea* reduced the biomass of smaller herbivores (GLM, *P*<0.001) by 68.1%

averaged over both zooplankton concentrations without affecting *Daphnia* biomass (Fig. 1). In

the longer-term field experiment, the two predators had similar effects on the herbivore groups,

with one notable difference. *Daphnia* biomass was reduced by 99.6% in tanks with *Notonecta*

and by 96.8% in tanks with both predators, but was unaffected in tanks with *Neoplea* (Table 1,

Fig. 2). Deviating from the laboratory results, neither predator alone significantly affected the

biomass of smaller herbivores; yet, there was a diversity effect such that adding both predators

reduced the biomass of smaller herbivores by 84.6% (Table 1, Fig. 2).

Mean biomass of both *Daphnia* and the smaller herbivores was associated with increased

variability of biovolume of smaller phytoplankton, but only *Daphnia* biomass was associated

with increased variability of *Oocystis* 1 biovolume (Table 2). Not surprisingly, *Daphnia* biomass

was also associated with increased variability of total phytoplankton biovolume, while the

biomass of smaller herbivores was only marginally associated with increased variability of total

phytoplankton biovolume. For all three algal groupings, mean biovolume was negatively

associated with mean *Daphnia* biomass, but was not affected by mean biomass of smaller

herbivores (Table 2).

Variability of the biovolume of smaller phytoplankton was not affected by adding either

predator alone, but was reduced when both predators were added (Table 3, Fig. 3C). In contrast,

variability of *Oocystis* 1 biovolume was not significantly affected by any predator treatment

(Table 3, Fig. 3B). When both groups of phytoplankton were summed together as total

community biomass, it was unclear whether predator diversity affected variability because

different methods of biomass estimation yielded different conclusions. Absorbance *in vivo* at 680

nm, a crude proxy for algal biomass, was significantly less variable only when both predators

were added (GLM, P=0.036). However, absorbance at 665 nm of extracted chlorophyll-*a*, a less

crude proxy, was only marginally significantly less variable with both predators (GLM,

P=0.071), and variability of total phytoplankton biovolume was not significantly affected (GLM,

P=0.114). Mean phytoplankton biomass, estimated in any way or for any of the phytoplankton

groupings, was not significantly affected by predator treatment (Table 3, Fig. 3D,E,F).

Discussion

Our results show that the two focal predator species, *Notonecta* and *Neoplea*, partitioned

herbivorous prey, and that this complementarity led to a reduction in the variability of smaller,

more edible phytoplankton biomass. *Notonecta* reduced *Daphnia* biomass and *Neoplea* reduced

biomass of a group of smaller herbivores in the laboratory; accordingly, both herbivore groups

were simultaneously reduced in the field only when both predators were added. Biomass of both

herbivore groups was separately and positively associated with variability of biomass of smaller

phytoplankton. In turn, variability of biomass of the smaller phytoplankton was only reduced in

the treatment with both predators. Thus, as represented in Figure 4, the presence of both

predators appeared to be necessary to suppress both *Daphnia* and the smaller herbivores at low

densities, thereby freeing the smaller phytoplankton of enough herbivory to stabilize its temporal

dynamics (Fig. 4d). Adding only one or the other predator failed to reduce both herbivore

groups, allowing herbivore-induced variability to continue (Fig. 4b-c). On the other hand, the

smaller herbivores did not affect the biomass variability of *Oocystis* 1, which was the largest

phytoplankton taxon and therefore expected to be least vulnerable to small herbivores.

Unsurprisingly, then, there was no diversity effect on the variability of *Oocystis* 1 biomass; in

fact, there was no difference across any predator treatments, perhaps because even *Daphnia*

affected this large alga too weakly to cause an effect across treatments. Because *Oocystis* 1 was

not affected by predator treatment, total phytoplankton biomass was only marginally less

variable when both predators were added (although this result depended on the method for

estimating phytoplankton biomass). Average biomass of both phytoplankton groups was

characterized by a similar pattern. That is, there were no differences in mean phytoplankton

biomass across predator treatments, and higher biomass of *Daphnia* – but not of smaller

herbivores – was associated with lower mean biomass of both phytoplankton groups.

There are several potential reasons for the difference in the effect of *Neoplea* on the

smaller herbivores in the laboratory versus the field experiment. One of the largest differences

between the experiments is the length of time the predators were added: five days in the

laboratory versus seven weeks in the field. The laboratory experiment attempted to estimate the

prey preference of the predators without allowing time for the prey to recover appreciably; on the

other hand, the field experiment allowed time for the prey populations to reproduce. This may

have allowed the smaller herbivores to continually recover (at least partially) from predation by

Neoplea, while the stronger effect of *Notonecta* on *Daphnia* did not allow *Daphnia* to recover.

With *Daphnia* virtually eliminated, *Notonecta* may have then switched to less preferred, smaller

prey, increasing the total consumption of smaller herbivores (Fig. 4). Whatever the reason for the

weaker effect of *Neoplea* in the field, both predators were still needed to suppress both herbivore

groups just as the laboratory results suggested.

The cascading diversity effect was not apparent in all tanks or taxa, but this pattern was

consistent with background variation in the mesocosms and in edibility among the plankton taxa.

In at least one mesocosm per treatment, variability of the smaller phytoplankton was at least as

low as the average variability with both predators (Fig. 3). This pattern can be explained by large

background variation in the system: even in the absence of predators, variation in the density of

Daphnia and of the smaller herbivores was very large, with some control tanks having

consistently low herbivore densities. This meant that only some tanks within each treatment

contained enough herbivores for a cascading predator effect to be detected. Still, the diversity-

stability effect remained significant when the smaller ("edible") phytoplankton were defined in

every reasonable way, and is further supported by an intuitive mechanism linking

complementarity to stability. The experiments were not set up to prove this mechanism, so our

description of this mechanism amounts to our hypothesis for how predator complementarity led

to lower algal variability. We did not know *a priori* which planktonic species would establish in

the mesocosms, nor how they would interact, and so it was necessary to explore the data to find a

hypothesized mechanism. Dominant species (i.e., *Arctodiaptomus*) that were not affected by the

treatments obscured the potential food web mechanism unless removed from analysis. It is not

surprising that some taxa were not involved in the cascading food web effect, since species

within trophic levels commonly differ greatly in their capture probability and edibility. In fact,

that the taxa which appeared not to be involved in the cascade stand out from the others in terms

of lower capture probability only strengthens support for the proposed food web mechanism.

This study provides the first simultaneous evaluation of effects of predator diversity on

both the average and variability of primary producer biomass. Most previous predator diversity-

ecosystem function experiments measured only mean biomass of trophic groups as dependent

variables (Bruno and O'Connor 2005; Straub et al. 2008). This work tended to show that

increasing predator diversity, as long as the predators partitioned resources, led to lower mean

herbivore biomass and higher mean autotroph biomass – i.e., a diversity-biomass trophic cascade

(Straub et al. 2008). In our field experiment, predator diversity decreased mean biomass of the

focal herbivores but did not significantly increase mean phytoplankton biomass, although there

was a weak, non-significant trend towards higher biomass of smaller phytoplankton. Only

Daphnia was associated with lower phytoplankton biomass, and not the smaller herbivores;

overall, the effect of the herbivores on mean phytoplankton biomass seemed to be too weak to

complete a diversity-biomass trophic cascade. It is not uncommon for herbivores, and therefore

changes in herbivore biomass, to have weak effects on plant biomass (Maron and Crone 2006).

The strength of herbivory is often dampened by the variable food quality of plants, low

encounter rates between herbivores and plants, indirect effects, or other factors (Borer et al.

2005; Maron and Crone 2006). On the other hand, both *Daphnia* and the smaller herbivores had

de-stabilizing effects on smaller phytoplankton biomass, and so there was a stronger pathway

from predator diversity to small phytoplankton stability, resulting in the completion of a

diversity-stability trophic cascade. Thus, in this case, the temporal variability of autotroph

biomass was more sensitive to changes in herbivore biomass than was average autotroph

biomass, at least for the taxa more vulnerable to herbivory. Future studies will need to explore

the generality of this result.

A few studies have reported the effects of predator diversity on the variability of

predation or adjacent trophic levels, and some have described theoretical mechanisms relevant to

these studies. Griffin and Silliman (2011) found that the combination of two predators which

exhibited temporal complementarity in attack rates reduced the temporal variability of the total

predation rate on a shared prey. Ong and Vandermeer (2015) showed theoretically that the

combination of a predator and a parasite can suppress a pest to a low, stable population.

However, our study is the first (to our knowledge) to test whether predator diversity can cascade

down a food web to reduce temporal variability of primary producer biomass. Theoretical work

suggests several mechanisms linking diversity and stability in ecosystems, mostly focusing on

single trophic levels (Loreau and de Mazancourt 2013). With a two-trophic level model,

Thébault and Loreau (2005) showed that decreasing herbivore biomass stabilizes (and increases)

plant biomass. Coupling this finding with the consensus from the literature that complementarity

of resource use by predators tends to reduce herbivore biomass, it follows that predator

complementarity should indirectly stabilize (and increase) plant biomass. However, we are not

aware of a theoretical study making this argument; thus, our results suggest a new diversity-

stability mechanism that could be important in biological control.

Our results suggest that adding multiple natural enemies to an agro-ecosystem can

stabilize fluctuations in crop yields, provided the natural enemies partition pest resources. While

a goal of farmers will always clearly be to achieve as high an average yield as possible,

achieving consistent yields is often just as important. We showed that combining predators

differing in a trait known to correlate with prey preference, body size, can be a key factor leading

to the cascading stabilizing effect on phytoplankton. Phytoplankton are increasingly cultured as a

crop for a variety of purposes from biomass production for biofuels, animal feed, or fertilizer, to

nutrient removal from wastewaters. However, algal culturing is not yet widely practiced on large

scales, in large part because it has so far proven too difficult to achieve consistently high algal

yields as algae ponds are colonized by pests that are difficult or costly to control by mechanical

or chemical means. Adding a functionally diverse array of predators to algae ponds may be a

feasible, economical, self-sustaining way to encourage more reliable algal yields.

The ecological principles that seemed to govern the diversity-stability trophic cascade

effect in our experiment are general and therefore may apply to terrestrial agriculture as well.

Our results suggest a simple biological control technique wherein consortia of natural enemy

species varying in size are added or encouraged to an agro-ecosystem to promote crop stability.

Diversity of other functional traits related to prey choice, microhabitat preference, temporal

patterns in control strength, or variance in mode of prey suppression may also encourage crop

yield stability in a similar fashion and could also be managed in natural enemy consortia when

relevant information is available. For example, top-down control can be stabilized by adding

some predators more active in warm temperatures and some more active in cold temperatures

(Griffin and Silliman 2011), or both parasites and predators (Ong and Vandermeer 2015),

leading to consistently low pest densities and stabilizing crop yields. While our study emphasizes

the importance of complementarity, redundancy is also likely important as temporal and spatial

scales increase, to guard against periods of weakened control by, or extirpations of, natural

enemies at certain times or locations (Peralta et al. 2014).

Here we have demonstrated that higher predator diversity and higher complementarity of

prey resource use, can cascade down to stabilize temporal dynamics of algal biomass without

affecting mean algal biomass. This connection between predator diversity and primary producer

stability is an important step towards joining biodiversity-ecosystem function theory with food

web theory, as biodiversity and ecosystem functioning research has only recently begun

incorporating energy flows in a food web context. Future work is needed to further explore the

simultaneous influence of predator diversity, in its various forms, on both the average and

stability of community- and ecosystem-level functioning. It is important to uncover whether

stability is generally more sensitive to cascading food web effects than is average biomass, or

how the two effects are related in different contexts. Predator diversity may also have different

effects on other measures of stability, such as spatial variability and resilience. Developing this

field at the intersection of biodiversity-ecosystem functioning and food web ecology not only

will improve our understanding of the functioning of natural ecosystems and their vulnerability

to anthropogenic biodiversity loss, but also will provide information that can be directly used to

manage for more reliable crop production and other ecosystem services.

Acknowledgements

Special thanks to S Manning for providing laboratory space and advice, and to LA Sekula and J

Earwood for help processing samples. Thanks to R Deans for help with insect collection and

experiment setup, to D Correa for help with nutrient analysis, to D Nobles for providing a

microscope, and to C Farrior for providing advice. Thanks to C Farrior, A Wolf, and their

laboratory groups for feedback on earlier versions of the manuscript. This research was

supported by the Department of Integrative Biology at the University of Texas at Austin.

Literature cited

Adrian, Rita, and Thomas M Frost. 2018. "Omnivory in Cyclopoid Copepods: Comparisons of

　　　Algae and Invertebrates as Food for Three, Differently Sized Species," 16.

Anderson, David H., Stephen Darring, and Arthur C. Benke. 1998. "Growth of Crustacean

　　　Meiofauna in a Forested Floodplain Swamp: Implications for Biomass Turnover."

　　　Journal of the North American Benthological Society 17 (1): 21–36.

　　　https://doi.org/10.2307/1468049.

APHA. 1989. Standard methods for the examination of water and wastewater, 17th edn.

　　　American Public Health Association, Washington, D.C.

Douglas Bates, Martin Maechler, Ben Bolker, Steve Walker (2015). Fitting Linear Mixed-

　　　Effects Models Using lme4. Journal of Statistical Software, 67(1), 1-

　　　48.<doi:10.18637/jss.v067.i01>.

Benemann, John. 2013. "Microalgae for Biofuels and Animal Feeds." *Energies* 6 (11): 5869–86.

　　　https://doi.org/10.3390/en6115869.

481   Borer, E. T., E. W. Seabloom, J. B. Shurin, K. E. Anderson, C. A. Blanchette, B. Broitman, S. D.

482   Cooper, and B. S. Halpern. 2005. "WHAT DETERMINES THE STRENGTH OF A

483   TROPHIC CASCADE?" *Ecology* 86 (2): 528–37. https://doi.org/10.1890/03-0816.

484   Bruno, John F., and Mary I. O'Connor. 2005. "Cascading Effects of Predator Diversity and

485   Omnivory in a Marine Food Web: Cascading Effects of Predator Diversity." *Ecology*

486   *Letters* 8 (10): 1048–56. https://doi.org/10.1111/j.1461-0248.2005.00808.x.

487   Cardinale, Bradley J., J. Emmett Duffy, Andrew Gonzalez, David U. Hooper, Charles Perrings,

488   Patrick Venail, Anita Narwani, et al. 2012. "Biodiversity Loss and Its Impact on

489   Humanity." *Nature* 486 (7401): 59–67. https://doi.org/10.1038/nature11148.

490   Cardinale, B. J., Emmett Duffy, Diane Srivastava, Michel Loreau, Matt Thomas, and Mark

491   Emmerson. 2009. Towards a food-web perspective on biodiversity and ecosystem

492   functioning. – In: Naeem, S. et al., Biodiversity and Human Impacts. Oxford University

493   Press, pp. 105–120.

494   Culver, David A., Mary M. Boucherle, David J. Bean, and James W. Fletcher. 1985. "Biomass

495   of Freshwater Crustacean Zooplankton from Length–Weight Regressions." *Canadian*

496   *Journal of Fisheries and Aquatic Sciences* 42 (8): 1380–90. https://doi.org/10.1139/f85-

497   173.

498   Dumont, Henri J., Isabella Van de Velde, and Simonne Dumont. 1975. "The Dry Weight

499   Estimate of Biomass in a Selection of Cladocera, Copepoda and Rotifera from the

500   Plankton, Periphyton and Benthos of Continental Waters." *Oecologia* 19 (1): 75–97.

501   https://doi.org/10.1007/BF00377592.

EPA. 2016. "Standard Operating Procedure for Zooplankton Analysis" (LG403, Revision 07).

Environmental Protection Agency, Washington, D.C.

Glazier, Douglas S. 2010. "A Unifying Explanation for Diverse Metabolic Scaling in Animals

and Plants." *Biological Reviews* 85 (1): 111–38. https://doi.org/10.1111/j.1469-

185X.2009.00095.x.

Griffin, John N., and Brian R. Silliman. 2011. "Predator Diversity Stabilizes and Strengthens

Trophic Control of a Keystone Grazer." *Biology Letters* 7 (1): 79–82.

https://doi.org/10.1098/rsbl.2010.0626.

Hall, Spencer R., Mathew A. Leibold, David A. Lytle, and Val H. Smith. 2004. "Stoichiometry

and planktonic grazer composition over gradients of light, nutrients, and predation risk."

Ecology 85 (8): 2291–2301. https://doi.org/10.1890/03-0471.

Hampton, Stephanie E., and John J. Gilbert. 2001. "Observations of Insect Predation on

Rotifers." In *Rotifera IX*, edited by L. Sanoamuang, H. Segers, R. J. Shiel, and R. D.

Gulati, 115–21. Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-94-010-

0756-6_17.

Jiang, Lin, and Zhichao Pu. 2009. "Different Effects of Species Diversity on Temporal Stability

in Single□Trophic and Multitrophic Communities." *The American Naturalist* 174 (5):

651–59. https://doi.org/10.1086/605961.

Kilham, Susan S, Daniel A Kreeger, Scott G Lynn, Clyde E Goulden, and Lazaro Herrera. 1998.

"COMBO: A Defined Freshwater Culture Medium for Algae and Zooplankton."

Hydrobiologia 377: 147–59.

Leon, Blaustein. 1998. "Influence of the Predatory Backswimmer, Notonecta Maculata, on

Invertebrate Community Structure." *Ecological Entomology* 23 (3): 246–52.

https://doi.org/10.1046/j.1365-2311.1998.00138.x.

Loreau, Michel, and Claire de Mazancourt. 2013. "Biodiversity and Ecosystem Stability: A

Synthesis of Underlying Mechanisms." Edited by Emmett Duffy. *Ecology Letters* 16

(May): 106–15. https://doi.org/10.1111/ele.12073.

Maron, John L, and Elizabeth Crone. 2006. "Herbivory: Effects on Plant Abundance,

Distribution and Population Growth." *Proceedings of the Royal Society B: Biological*

Sciences 273 (1601): 2575–84. https://doi.org/10.1098/rspb.2006.3587.

McCauley, Edward. 1984. "The Estimation of the Abundance and Biomass of Zooplankton in

Samples." In *A Manual for the Assessment of Secondary Productivity in Fresh Waters*,

edited by John A Downing and F.H. Rigler, 228–65. Blackwell Scientific Publishers.

Murdoch, William W., Mary Ann Scott, and Peter Ebsworth. 1984. "Effects of the General

Predator, Notonecta (Hemiptera) Upon a Freshwater Community." *The Journal of Animal*

Ecology 53 (3): 791. https://doi.org/10.2307/4660.

Ong, Theresa Wei Ying, and John H. Vandermeer. 2015. "Coupling Unstable Agents in

Biological Control." *Nature Communications* 6 (1). https://doi.org/10.1038/ncomms6991.

Peralta, Guadalupe, Carol M. Frost, Tatyana A. Rand, Raphael K. Didham, and Jason M.

Tylianakis. 2014. "Complementarity and Redundancy of Interactions Enhance Attack

Rates and Spatial Stability in Host–Parasitoid Food Webs." *Ecology* 95 (7): 1888–96.

https://doi.org/10.1890/13-1569.1.

Purvis, Andy, John L. Gittleman, Guy Cowlishaw, and Georgina M. Mace. 2000. "Predicting

Extinction Risk in Declining Species." *Proceedings of the Royal Society of London.*

Series B: Biological Sciences 267 (1456): 1947–52.

https://doi.org/10.1098/rspb.2000.1234.

R Core Team. 2017. "R: A language and environment for statistical computing." R Foundation

for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Schmitz, Oswald J. 2007. "Predator Diversity and Trophic Interactions." *Ecology* 88 (10): 2415–

26. https://doi.org/10.1890/06-0937.1.

Schneider, Caroline A, Wayne S Rasband, and Kevin W Eliceiri. 2012. "NIH Image to ImageJ:

25 Years of Image Analysis." *Nature Methods* 9 (7): 671–75.

https://doi.org/10.1038/nmeth.2089.

Smith, Val H., Belinda S.M. Sturm, Frank J. deNoyelles, and Sharon A. Billings. 2010. "The

Ecology of Algal Biodiesel Production." *Trends in Ecology & Evolution* 25 (5): 301–9.

https://doi.org/10.1016/j.tree.2009.11.007.

Snyder, William E. 2019. "Give Predators a Complement: Conserving Natural Enemy

Biodiversity to Improve Biocontrol." *Biological Control* 135 (August): 73–82.

https://doi.org/10.1016/j.biocontrol.2019.04.017.

Straub, Cory S., Deborah L. Finke, and William E. Snyder. 2008. "Are the Conservation of

Natural Enemy Biodiversity and Biological Control Compatible Goals?" *Biological

Control* 45 (2): 225–37. https://doi.org/10.1016/j.biocontrol.2007.05.013.

Sturm, Belinda S.M., Edward Peltier, Val Smith, and Frank deNoyelles. 2012. "Controls of

Microalgal Biomass and Lipid Production in Municipal Wastewater-Fed Bioreactors."

Environmental Progress & Sustainable Energy 31 (1): 10–16.

https://doi.org/10.1002/ep.10586.

Thébault, Elisa, and Michel Loreau. 2005. "Trophic Interactions and the Relationship between

Species Diversity and Ecosystem Stability." *The American Naturalist* 166 (4): E95–114.

https://doi.org/10.1086/444403.

van Driesche, Roy, Mark Hoddle, and Ted Center. 2008. "Control of pests and weeds by natural

enemies: an introduction to biological control." Blackwell Publishing: Malden, MA.

591   **Table 1.** Results of gamma GLMMs testing the effects of predator additions in the field

592   experiment on biomass of a) *Daphnia*, b) smaller herbivores, and c) both of these herbivore

593   groups combined.

Parameter	Estimate	SE	t	*P*
a) *Daphnia*				
intercept	7.466	0.802	9.315	**<0.001**
Notonecta	-5.874	1.134	-5.183	**<0.001**
Neoplea	0.123	1.134	0.114	0.909
both predators	-4.186	1.134	-3.693	**<0.001**
b) smaller herbivores				
intercept	6.483	0.460	14.106	**<0.001**
Notonecta	-0.259	0.650	-0.399	0.690
Neoplea	-0.801	0.650	-1.233	0.218
both predators	-1.278	0.650	-1.966	**0.049**
c) both herbivore groups				
intercept	7.914	0.531	14.913	**<0.001**
Notonecta	-1.662	0.750	-2.216	**0.027**
Neoplea	-0.018	0.751	-0.024	0.981
both	-2.370	0.750	-3.159	**0.002**

604 **Table 2.** Results of gamma GLMs testing the effects of mean *Daphnia* biomass, and of summed

605 biomass of smaller herbivorous zooplankton, on the CV of biovolume of a) total phytoplankton,

606 b) *Oocystis* 1, and c) smaller phytoplankton.

Parameter	Estimate	SE	t	*P*
a) Total phytoplankton				
intercept	-1.171	0.1404	-8.339	<0.001
Daphnia	3.578×10^{-5}	9.861×10^{-6}	3.629	**0.0021**
smaller herbivores	2.434×10^{-4}	1.305×10^{-4}	1.865	0.0795
b) *Oocystis* 1				
intercept	-0.5875	0.1077	-5.458	<0.001
Daphnia	2.703×10^{-5}	7.562×10^{-6}	3.575	**0.0023**
smaller herbivores	5.127×10^{-5}	1.001×10^{-4}	0.512	0.6150
c) Smaller phytoplankton				
intercept	-1.239	0.1630	-7.600	<0.001
Daphnia	3.449×10^{-5}	1.145×10^{-5}	3.012	**0.0079**
smaller herbivores	3.613×10^{-4}	1.515×10^{-4}	2.385	**0.0290**

620   **Table 3.** Results of gamma GLMs testing the effects of predator additions in the field experiment

621   on the temporal CV of biovolume of a) smaller phytoplankton (all but the largest morphospecies,

622   *Oocystis* 1), b) *Oocystis* 1, and c) total phytoplankton.

Parameter	Estimate	SE	t	*P*
a) smaller phytoplankton				
intercept	-0.4310	0.3001	-1.436	0.170
Notonecta	-0.4497	0.4244	-1.060	0.305
Neoplea	0.0626	0.4244	0.148	0.885
both predators	-1.0622	0.4244	-2.503	**0.024**
b) *Oocystis* 1				
intercept	-0.4977	0.2399	-2.075	0.055
Notonecta	0.0010	0.3392	0.003	0.998
Neoplea	0.4843	0.3392	1.428	0.173
both predators	-0.1010	0.3392	-0.298	0.770
c) total phytoplankton				
intercept	-0.6181	0.3114	-1.985	0.065
Notonecta	-0.1808	0.4404	0.668	0.514
Neoplea	0.2940	0.4404	0.668	0.514
Both	-0.7366	0.4404	-1.673	0.114

Figure legends

Figure 1. Multiplicative effects of addition of *Notonecta* and of *Neoplea* on biomass of *Daphnia* and on summed biomass of smaller herbivores in the laboratory experiment. Circles represent coefficient estimates from GLMs modeling the effects of predator addition on the mass of each herbivore group, and error bars represent 95% confidence intervals about the estimates. The dotted line at 1 represents no effect, and the dotted line at 0 represents total elimination of an herbivore group.

Figure 2. Multiplicative effects of addition of *Notonecta, Neoplea*, and of both predators together on biomass of *Daphnia*, summed biomass of smaller herbivores (excludes *Arctodiaptomus* and nauplii), and biomass of both herbivore groups summed together in the field experiment. Shapes represent coefficient estimates from GLMMs modeling the effects of predator addition on the mass of each herbivore group, and error bars represent 95% confidence intervals about the estimates, plotted on a natural log scale. The dotted line represents no effect.

Figure 3. Temporal coefficient of variation (CV) of phytoplankton biovolume, a-c, and time series of mean phytoplankton biovolume, d-f, by predator treatment. Symbols in d-f are means ±1 SEM. Panels a and d show values for smaller phytoplankton (all besides the largest morphospecies, *Oocystis* 1), b and e show values for *Oocystis* 1, and c and f show values for total phytoplankton. The red star represents a significant effect of that predator treatment (GLMM, p=0.024) compared against the control.

Figure 4. Diagrams depicting the hypothesized food web and mechanism for food web effects

on phytoplankton stability in the field experiment under each predator treatment. A represents

the controls (no predators added), B represents addition of *Notonecta* only, C represents addition

of *Neoplea* only, and D represents addition of both predators. Arrows represent hypothesized

energy flow, with thicker arrows indicating faster consumption rates. Relative size of the

Daphnia and "other herbivores" represent relative mean biomass. The time series at the bottom

of each panel indicate relative temporal variability of each algal group. Red stars indicate

significant differences compared against the control treatment (either in mean herbivore biomass

with GLMM, or in CV of algal biovolume with GLM; $\alpha=0.05$).

A

Daphnia

other herbivores

Oocystis 1

smaller algae

biovolume

time

biovolume

time

B

Notonecta

Daphnia

*

other herbivores

Oocystis 1

smaller algae

biovolume

time

biovolume

time

C

Neoplea

Daphnia

other herbivores

Oocystis 1

smaller algae

biovolume

time

biovolume

time

D

Notonecta

Neoplea

Daphnia

*

other herbivores

*

Oocystis 1

smaller algae

biovolume

time

biovolume

time

*