1   **Convergent evolution of seasonal camouflage in response to reduced snow cover across the**

2   **snowshoe hare range**

3   Matthew R. Jones[1,2], L. Scott Mills[3,4], Jeffrey D. Jensen[2], Jeffrey M. Good[1,3]

4   [1]Division of Biological Sciences, University of Montana, Missoula, MT 59812, USA.

5   [2]School of Life Sciences, Arizona State University, Tempe, AZ 85281, USA

6   [3]Wildlife Biology Program, University of Montana, Missoula, MT 59812, USA.

7   [4]Office of Research and Creative Scholarship, University of Montana, Missoula, MT 59812,

8   USA.

9   Corresponding authors: jeffrey.good@umontana.edu, matthew.r.jones.1@asu.edu

Key words: climate change, dominance, local adaptation, migration, parallel evolution

Abstract

Determining how different populations adapt to similar environments is fundamental to

understanding the limits of adaptation under changing environments. Snowshoe hares (*Lepus*

americanus) typically molt into white winter coats to remain camouflaged against snow. In some

warmer climates, hares have evolved brown winter camouflage – an adaptation that may spread

under climate change. We used extensive range-wide genomic data to 1) resolve broad-scale

patterns of population structure and gene flow and 2) investigate the factors shaping the origins

and distribution of winter-brown camouflage variation. In coastal Pacific Northwest (PNW)

populations, winter-brown camouflage is known to be determined by a recessive haplotype at the

Agouti pigmentation gene. Our phylogeographic analyses revealed deep structure and limited

gene flow between PNW and more northern Boreal populations, where winter-brown

camouflage is rare along the range edge. Genome sequencing of a winter-brown snowshoe hare

from Alaska shows that it lacks the winter-brown PNW haplotype, reflecting a history of

convergent phenotypic evolution. However, the PNW haplotype does occur at low frequency in a

winter-white population from Montana, consistent with the spread of a locally deleterious

recessive variant that is masked from selection when rare. Simulations show that if annual snow

cover dramatically declined in the same population, then the predicted selective increase in

frequency of the now beneficial winter-brown *Agouti* allele is likely to be extremely slow due to

the same masking effect. Our findings underscore how allelic dominance can shape the

geographic extent and rate of convergent adaptation in response to rapidly changing

environments.

Introduction

In response to a shared selection pressure, populations may adapt through the migration of beneficial alleles or through independent mutations that result in the evolution of convergent phenotypes [1–3]. Distinguishing between these scenarios is crucial to understand the capacity of populations to adapt rapidly to environmental change [2,4,5]. If gene flow between populations is sufficiently high, then beneficial variation may spread quickly [6] and potentially allow for rapid adaptive responses to changing environments [7]. However, the spread of adaptive variation may be limited at broad geographic scales, and populations may have to rely on independent standing genetic variation or new mutations to evolve convergent traits. While there is considerable evidence for both adaptation through gene flow and independent mutations in nature [8–14], few empirical studies have examined the specific factors that influence these outcomes in natural populations [2].

When a species range encompasses a mosaic of habitats, the relative probability of adaptation through gene flow versus independent mutation is predicted to be primarily a function of the distance between habitat patches (in units of dispersal distance) and the strength of selection against locally adaptive alleles in intervening habitats [2,15,16]. In general, as distance between patches and the strength of purifying selection in intervening habitats increases, so does the relative probability of adaptation through independent mutations. Adaptation via independent mutations is therefore predicted to be more common in widespread populations where dispersal distance is short relative to the total range size [15]. For example, rock pocket mice (*Chaetodipus intermedius*) have repeatedly evolved melanistic coats across patchy lava flows in the southwestern United States. Although substantial gene flow between adjacent lava flows has likely resulted in the migration of melanic alleles [13], melanism is attributed to different

mutations across disparate lava flows [8,17,18]. Theoretical models further predict that the

relative probability of adaptation via independent mutations increases rapidly with distance

between lava flows (over the scale of tens to hundreds of kilometers) [2], due in large part to

strong selection against coat color mismatch [19–21]. Thus, there is a clear tradeoff between

dispersal distance and the strength of purifying selection that strongly dictates the probability of

adaptation through convergent evolution or gene flow.

61   The effect of genetic dominance on the probability of convergent evolution has not yet

been thoroughly explored [2]. 'Haldane's sieve' [22] predicts that *de novo* dominant mutations

enjoy a much greater probability of fixation compared to *de novo* recessive mutations

(probability of fixation ≈$2hs$), because rare dominant mutations are visible to selection [23]. As

de novo beneficial recessives are masked to selection when rare, those that do ultimately reach

fixation may spend a longer period of time drifting at low frequency [24]. Likewise, rare

recessive migrant alleles are expected to exhibit the same behavior, which may allow more time

for mutation to generate 'competing' convergent phenotypes upon which selection can act.

However, Orr and Betancourt [25] demonstrated that genetic dominance has no effect on the

probability of fixation for alleles in mutation-selection balance because recessive alleles have a

higher mutation-selection balance frequency. Thus, for relatively high amounts of gene flow,

genetic dominance may have little to no effect on the probability of adaptation via migration

versus *de novo* mutation. In fact, the masking of low frequency recessive alleles may result in

weaker purifying selection in intervening habitats therefore facilitating the spread of adaptive

variation through gene flow [2], although this hypothesis remains to be tested.

76   Snowshoe hares (*Lepus americanus*) are one of at least 21 species that have evolved

seasonal molts to white winter coats to maintain camouflage in snowy winter environments.

Because color molts are cued by photoperiod, and hence may become mismatched under rapidly

changing environments [26,27], seasonal camouflage provides a useful trait to understand

adaptation to climate change [26–30]. Reduction in the extent and duration of snow cover is one

of the strongest signatures of climate change in the Northern hemisphere [31,32], suggesting that

selection should increasingly favor delayed winter-white molts in snowshoe hares to reduce the

total duration of coat color mismatch. However, below some annual snow cover threshold,

populations are predicted to maintain brown coloration during the winter [28]. Consistent with

this, snowshoe hares maintain brown winter camouflage in some temperate environments with

reduced snow cover [33], a strategy that should be increasingly favored under climate change

[28,29].

Winter-brown snowshoe hares are common in coastal regions of the Pacific Northwest

(PNW), where this Mendelian trait is determined by a recessive variant of the *Agouti*

pigmentation gene [29]. The locally adaptive *Agouti* variant was introduced into snowshoe hares

by hybridization with black-tailed jackrabbits ~9-18 thousand years ago (kya) and subsequently

experienced a local selective sweep within the last few thousand years [34]. Occasional records

of winter-brown camouflage also occur in more northern regions along the Pacific coast (e.g.,

Canada and Alaska) and in eastern North America [28,35]. Although hares are expected to be

more or less continuously distributed along suitable Pacific coast environments, there is a deep

split and little evidence of recent gene flow between the northern 'Boreal' populations and

southern 'PNW' populations [36,37]. Given this historic population structure, it remains unclear

whether the distribution of winter-brown camouflage across populations from disparate parts of

the range reflects independent genetic origins (i.e. trait convergence) or spread of the

introgressed PNW *Agouti* allele.

Here, we use new and previously published genetic data to investigate the roles of gene

flow and mutation in shaping the evolution of winter-brown camouflage across populations of

snowshoe hares. We first combined previously published microsatellite (*n*=853 individuals, 8

loci) with new and published whole exome data (*n*=95 individuals) to resolve range-wide

patterns of population history and gene flow in snowshoe hares, which provides crucial context

for understanding the historical spread and adaptive potential for winter-brown camouflage to

climate change. We then generated whole genome sequence (WGS) data of a winter-brown

Alaska (AK) hare to test whether winter-brown camouflage in Boreal snowshoe hares arose

independently from PNW populations. Next, to understand the geographic limits of the recessive

PNW *Agouti* allele, we used pooled WGS data to estimate its frequency in a winter-white

population from Montana (MT), approximately 600 km in a straight-line from the closest winter-

brown PNW population. Finally, we used both theoretical predictions and simulations to

understand the factors influencing the geographic scope of the PNW *Agouti* allele and its

potential to contribute to rapid adaptation in response to warming climates.

Methods

<u>Samples and genomic data generation</u>

To resolve patterns of range-wide population structure, we performed targeted whole exome

enrichment of 12 snowshoe hares previously sampled from 12 localities across Canada, Alaska,

and the eastern United States and three hares from the southern Rocky Mountains (representing

the 'Boreal' and 'Rockies' lineages as defined by Cheng et al. [36]). The targeted whole exome

capture was designed by Jones et al. [29] to enrich for ~99% of genic regions annotated in the

European rabbit (*Oryctolagus cuniculus*) genome (61.7-Mb spanning 213,164 intervals; ~25-Mb

protein-coding exons, ~28-Mb untranslated region, ~9-Mb intronic or intergenic). We followed

the library preparation protocols outlined in Jones et al. [29] and sequenced libraries on one lane

of Illumina HiSeq2500 with paired-end 100 bp reads (HudsonAlpha Institute for Biotechnology;

Huntsville, AL). Exome sequences for Boreal and Rockies snowshoe hares were combined with

published whole exome data [29] from 80 PNW snowshoe hares (n=95 total, Table S1),

including a monomorphic winter-brown population from southeast British Columbia (BC1,

n=14), a monomorphic winter-white population from Seeley Lake area in western MT (n=14),

and two polymorphic coat color populations from Oregon (OR, n=26; two localities) and

Washington (WA, n=26) [29].

To resolve the historical spread of the winter-brown PNW *Agouti* allele, we performed

whole genome sequencing of a winter-brown snowshoe hare museum specimen (AK) from

southwest Alaska first noted by Link Olson and later obtained through loan from University of

Alaska Museum of the North (UAM 116170, collected on 4 January 2013). We also performed

whole genome sequencing on a DNA pool of 81 snowshoe hares from two localities in Glacier

National Park, MT (Table S1). We extracted genomic DNA from muscle tissue of the AK hare

sample following the Qiagen DNeasy Blood and Tissue kit protocol (Qiagen, Valencia, CA). For

the pooled MT whole genome sequencing, we pooled previously extracted DNA samples [36] in

approximately equimolar quantities based on Qubit concentrations (Invitrogen Qubit

Quantitation system LTI). We then prepared genomic libraries for all samples following the

KAPA Hyper prep kit manufacturer's protocol. We sheared genomic DNA to ~300 bp using a

Covaris E220evolution ultrasonicator and performed a stringent size selection using a custom-

prepared carboxyl-coated magnetic bead mix [38]. We determined indexing PCR cycle number

for each library with quantitative PCR (qPCR) on a Stratagene Mx3000P thermocycler (Applied

Biosystems) using a DyNAmo Flash SYBR Green qPCR kit (Thermo Fisher Scientific). Final

libraries were size-selected again with carboxyl-coated magnetic beads, quantified with a Qubit

(Thermo Fisher Scientific), and pooled for sequencing by Novogene (Novogene Corporation

Ltd.; Davis, CA) on two lanes of Illumina HiSeq4000 using paired-end 150bp reads. The whole

genome sequence from the AK hare was combined with previously published whole genome

sequencing of five black-tailed jackrabbits, three winter-brown snowshoe hares from OR, WA,

and BC1, and three winter white snowshoe hares from MT, a hare from Utah (UT), and a hare

from Pennsylvania (PA, Table S1) [29].

<u>Read processing and variant calling</u>

For all raw Illumina sequence data, we trimmed adapters and low-quality bases (mean phred-

scaled quality score <15 across 4 bp window) using Trimmomatic v0.35 [39] and merged paired-

end reads overlapping more than 10 bp and with lower than 10% mismatched bases using

FLASH2 [40].

Whole genome sequence data were mapped to either a snowshoe hare or black-tailed

jackrabbit pseudoreference (see Jones et al. [29] for details) using default settings in BWA-MEM

v0.7.12 [41]. We used *PicardTools* to remove duplicate reads with the MarkDuplicates function

and assigned read group information with the AddOrReplaceReadGroups function. Using GATK

v3.4.046 [42], we then identified poorly aligned genomic regions with RealignerTargetCreator

and locally realigned sequence data in these regions with IndelRealigner. We performed

population-level multi-sample variant calling using default settings with the GATK

UnifiedGenotyper. Here, we called variants separately for each previously defined snowshoe

hare population genetic cluster (i.e., Boreal, Rockies, BC1, MT, OR, and WA) and for black-

tailed jackrabbits. We performed variant filtration in VCFtools v0.1.14 [43]. For whole exome

and whole genome data, we filtered genotypes with individual coverage <5× or >70× or with a

phred-scaled quality score <30. Additionally, we removed all indel variants and filtered SNPs

with a phred-scaled quality score <30, Hardy-Weinberg P<0.001. We required that sites have no

missing data across individuals.

Range-wide population genetic structure and gene flow

We inferred a maximum likelihood tree with a general time reversible model in RAxML v8.2.8

[44] using a subset of the concatenated snowshoe hare exome data (n=12 Boreal hares, n=3

Rockies hares, n=12 PNW hares; 21,167,932 total sites) with European rabbit (*Oryctolagus*

cuniculus) as the outgroup. Using this maximum likelihood tree as a starting tree, we estimated a

maximum clade credibility tree and node ages with a constant population size coalescent model

in BEAST 2 [45]. We assumed a strict molecular clock and an HKY substitution model using

empirical base frequencies. We specified default priors for the kappa parameter, gamma shape

parameter, and population size parameter and used a gamma distribution (alpha=0.0344, beta=1)

as a prior for the clock rate parameter. We ran the MCMC for 5 million steps and calibrated

divergence times using a log-normal distribution for the *Oryctolagus-Lepus* node age with a

median of 11.8 million years (95% prior density: 9.8–14.3) [46]. We also performed a species-

tree analysis using gene trees in ASTRAL v5.6.3 [47]. Gene trees were generated across 200 kb

windows (excluding windows wither fewer than 500 sequenced bases) using RAxML with a

GTR+gamma model and rapid bootstrap analysis (-f a -# 10). We collapse nodes on gene trees

with low bootstrap support values (<=10) and performed ASTRAL analyses on the collapsed

gene trees using default settings.

To test for signatures of population structure and gene flow among lineages, we

performed a population cluster analysis in ADMIXTURE v1.3.0 [48]. We tested K values from 1-

10 (representing the number of population clusters) and selected the K value with the lowest

cross-validation error. We also estimated range-wide effective migration and diversity surfaces

with EEMS [49] using extensive microsatellite data (n=853 individuals, 8 loci) generated by

Cheng et al. [36]. Varying the number of demes (50, 100, and 200) had little effect on estimates

of effective rates of migration and diversity, therefore we only report results for 200 demes. We

used default hyper-parameter values and tuned the proposal variances such that proposals were

accepted approximately 30% of the time. We ran EEMS for 2 million iterations with a burn-in of

1 million iterations and thinning iteration of 9999. Runs produced strong correlations between

observed and expected genetic dissimilarity both within and between demes, indicating good

model fit.

<u>Geographic distribution of the PNW *Agouti* allele</u>

The winter-brown AK snowshoe hare was collected approximately 3000 km (via the Pacific

coast) from Vancouver, BC, the northern limit of winter-brown PNW hare populations [29]. To

test whether the introgressed PNW *Agouti* allele has seeded winter-brown camouflage in

southwest AK, we generated a tree of the PNW *Agouti* haplotype region from our whole genome

sequence data of winter-brown and winter-white snowshoe hares (n=7) and black-tailed

jackrabbits (n=5). We defined the PNW *Agouti* haplotype region (chr4:5,340,275 – 5,633,020,

oryCun2 coordinates) based on the location of the introgressed black-tailed jackrabbit tract

identified by Jones et al. [29] using a hidden Markov model. We then inferred a maximum

likelihood phylogeny for this interval using RAxML [44] as above. Node support values were

generated from 1000 replicate bootstrap runs.

We also estimated the frequency of the PNW *Agouti* haplotype among pooled individuals

from two localities in MT (*n*=81) that are 575 km and 627 km from the polymorphic sampling

locality in WA, where winter-brown camouflage is relatively common. We used PoPoolation2

[50] to calculate the frequency of the winter brown-associated alleles at 555 sites across the

introgressed *Agouti* haplotype (chr4: 5,340,275 – 5,633,020). These positions were previously

shown to be strongly associated with coat color based on a likelihood ratio test of allele

frequency differences between winter-brown and winter-white individuals from low coverage

whole genome sequence data [29]. We summed the counts of winter-white and winter-brown

alleles across all positions to estimate both the mean and standard deviation of winter-brown

allele frequency. We excluded seven positions with unusually high frequencies of the winter-

brown allele (~45-100%) supported by reads that did not otherwise carry brown-associated

alleles (i.e., incongruent haplotype information), as these likely reflect sequencing errors.

<u>Probability of convergent adaptation through independent mutation</u>

To compare the observed spread of the PNW *Agouti* allele to theoretical expectations, we

estimated the relative probability of adaptation via independent mutations with distance from a

focal habitat patch using the model developed by Ralph and Coop [2] (equation 12) under a

range of conditions. Since the relative probability of adaptation through migration or *de novo*

mutation depends strongly on the rate at which mutations generate convergent phenotypes [2],

we tested a range of plausible mutation rates. A previous study using over 7 million house mice

estimated a mean rate of spontaneous visible coat color mutations of $11e^{-6}$ per locus/gamete [51],

which is ~1930× higher than the average genome-wide germline per site mutation rate (5.7e^{-9} per

site/gamete) [52]. Given a genome-wide mutation rate of 2.35e^{-9} per site/generation for rabbits

[53], we tested a high mutation rate of μ=4.54e^{-6} (i.e., the overall expected rate of visible coat

color mutations) and a low mutation rate of μ=4.54e^{-8} (i.e., assuming only 1% of coat color

mutations would lead to brown winter camouflage). We also tested how strong (s_m=0.01),

moderate (s_m=0.001), or weak (s_m=0.0001) purifying selection against the winter-brown allele in

intervening (i.e., snow-covered) habitats influences the probability of adaptation through

independent mutations. We assumed a relatively high mean dispersal distance of 2 km [54],

which is likely to produce a conservative estimate of the probability of adaptation via

independent mutation. We also assumed that the width of the secondary habitat patch was

relatively small (w=1 or ~6-45 km, depending on s_m), which is consistent with observations of

winter-brown camouflage at low frequency along portions of the range edge. Finally, we

assumed d=1 (dimension of the habitat), C=1, and γ=0.5.

<u>Simulations of selection on migrant alleles with genetic dominance</u>

Theory predicts that allelic dominance should not influence a mutation's probability of fixation

under positive selection if it starts in mutation/migration-selection balance [25]. Nonetheless,

rare recessive mutations may increase in frequency more slowly relative to rare dominant

mutations because they are initially invisible to selection, allowing more opportunity for

competition from convergent phenotypes arising through independent mutations. We used

simulations to test how the genetic dominance of alleles at equivalent mutation-selection

equilibrium frequencies influences the probability and rate of adaptation to new habitats when

environmental conditions change. In SLiM 3.0 [55] we performed 100 simulations of the MT

population (estimated N_e=245430) [34] with a positively selected recessive mutation (s=0.01,

h=0) starting at the inferred frequency of the PNW *Agouti* allele in MT (p). We assumed that p

reflects the equilibrium frequency (x) between the migration rate of the allele into the

environment (m) and the selection coefficient against the allele (s), which is given as $\sqrt{\frac{m}{s}}$ for a

recessive mutation. Given that x for a dominant mutation is simply $\frac{m}{s}$, we calculated x of an

equivalent dominant mutation (same values of m and s) as simple p^2. We then repeated these

simulations for a positively selected (s=0.01), completely dominant allele (h=1) starting at its

expected mutation-selection balance frequency. For each simulation we tracked the frequency of

the selected allele until it was either fixed or lost. To validate these simulations, we compared the

probabilities of allelic fixation from simulations to fixation probabilities analytically derived by

Kimura [56]. To test for significant differences in fixation probabilities and rates of adaptation

between recessive and dominant mutations we used Student's t-tests in R [57].

Results

<u>Snowshoe hare phylogeography</u>

We combined previously generated whole exome data for 80 PNW snowshoe hares (mean

coverage 21× ± 7.6 per individual) with new whole exomes of 15 hares across the range

sequenced to a mean coverage of 20.2× ± 8.2 (Fig. 1A). Exome-wide phylogenetic analyses

show three broad phylogeographic groups of snowshoe hares (Fig. 1B) with unambiguous

ASTRAL support scores using 6582 gene trees, consistent with previous studies [36,37].

However, ~12-25% of SNPs within phylogeographic groups appear to be shared across some

population boundaries (Table 1), indicating some amount of shared ancestral variation or gene

flow. We estimated that snowshoe hares from Canada, Alaska, and the eastern United States—

representing the Boreal lineage—diverged approximately 97.2 kya (95% posterior density: 77.4-

120.8 kya) from PNW and Rockies snowshoe hares (Fig. 1B). PNW and Rockies hare

populations are estimated to have split approximately 47.1 kya (95% posterior density: 37.2-58.2

kya).

Population structure analyses in ADMIXTURE also found strongest support for three

population clusters, although MT snowshoe hares (in the PNW lineage) showed 11-16% of their

ancestry assigned to the Rockies lineage, indicating ongoing gene flow or continuous genetic

structure (Fig. 1C). The ADMIXTURE analysis also indicated little apparent gene flow between the

Boreal lineage and PNW or Rockies lineages. Likewise, microsatellite-based estimates of

effective migration surfaces revealed that effective migration rates are approximately 100-fold

lower than the range-wide average near the apparent zone of contact between PNW and Boreal

lineages in western North America (Fig. 2A). We also estimated relatively low effective

migration rates across the southwestern edge of snowshoe hare range (western US; $\log(m) \approx -1$,

or 10-fold lower than the mean). In contrast, effective migration rates in the northwest part of the

range (Alaska and western Canada) appeared relatively high (~10-fold higher than the mean).

Effective diversity surfaces show that the highest relative genetic diversity occurs in the eastern

extent (Boreal lineage) and southwestern extent (PNW lineage) of the snowshoe hare range (Fig.

2B), consistent with previous studies suggesting that these regions were likely glacial refugia

[36]. Relative genetic diversity gradually decreased moving from east to west across the Boreal

range and the lowest genetic diversity was observed in the Rocky Mountain lineage, distributed

across the western US.

<u>Convergent evolution and the distribution of winter-brown camouflage variation</u>

Under a single origin of winter brown camouflage, we would expect the winter-brown AK

individual to nest within the black-tailed jackrabbit clade with the other winter-brown snowshoe

hares from the PNW. However, our phylogenetic analysis indicates that the winter-brown AK

individual unambiguously groups with a winter-white Boreal hare from PA (100% bootstrap

support), and more broadly with other winter-white hares across the range (Fig. 3). These results

demonstrate that the winter-brown phenotype of this AK hare is caused by mutations that are

independent from the introgressed *Agouti* haplotype that encodes winter-brown camouflage in

the PNW, approximately 3000 km away. Determining the genetic basis of this independent

origin of winter brown camouflage will likely require an independent genetic association study

in these populations.

Pooled whole genome sequencing detected that the recessive winter-brown *Agouti* allele

occurs at an estimated frequency of 1.24% ($\pm$ 0.01%) across predominately winter-white MT

localities ~600 km from the polymorphic zone. Interestingly, long-term live-trapping and

radiotelemetry-based field work from this region has resulted in a single observation of a winter-

brown hare in approximately 300 winter hare observations (0.33%; Scott Mills pers. obs.),

although this observed frequency is slightly higher than the Hardy-Weinberg expectation under

the estimated frequency of the winter-brown *Agouti* allele (p^2=0.015%, binomial test *p*-

value=0.044).

To understand the factors that have shaped the geographic distribution of the PNW

Agouti allele, we modelled the relative probability of adaptation through independent mutation

with distance from a focal habitat using the theoretical framework of Ralph and Coop [2]. The

change in relative probability of adaptation through independent mutation depended strongly on

both the mutation rate and selection coefficient against the allele in intervening habitats (Fig. 4).

Stronger selection coefficients produced a more rapid shift in the convergence probability with

distance relative to weaker selection (Fig. 4). For instance, under a high mutation rate to the

winter-brown phenotype (μ=4.54e^{-6}) and strong purifying selection (s_m=0.01) the probability of

independent adaptation transitioned sharply from P=0.1 at 78 km to P=0.9 at 140 km from a

focal habitat. Under a low mutation rate to the winter-brown phenotype (μ=4.54e^{-8}) and weak

selection (s_m=0.0001), independent adaptation was much less likely at close distances, as

expected, and the probability increased more gradually (P=0.1 at 778 km to P=0.9 at 1400 km).

<u>Migration-selection balance</u>

Assuming an environmental shift towards less snow results in strong positive selection (s=0.01)

on standing variation of the recessive (winter-brown) *Agouti* allele in MT, we used simulations

to estimate that the corresponding probability of fixation would be ~81% (95% confidence

intervals: 72.2-87.5%; Fig. 5A), which is consistent with an analytically-derived probability of

78.1%. For an equivalent dominant mutation starting in mutation-selection balance frequency

(0.015%) the simulated probability of fixation was approximately 77% using simulations (95%

confidence intervals: 67.8-84.2%) or 77.9% using the Kimura [56] equation. Neither estimate

was significantly different from the fixation probability of a recessive mutation (p=0.48, two-

tailed test of equality of proportions). However, we observed striking differences in the rates of

increase, conditional on fixation, of dominant and recessive mutations following a sudden

environmental change (Fig. 5B). The mean time to fixation was significantly faster (p<2.2e-16,

Student's T-test) for a recessive mutation (9645 generations, SD=3609 generations) compared to

a dominant mutation (19445 generations, SD=7273 generations, p<2.2e^{-16}). However, as

expected, we see that initial rates of allele frequency change and phenotypic adaptation are

considerably faster for a positively selected dominant mutation (Fig. 5B). For instance, the mean

time for the selected phenotype determined by the recessive mutation to reach a frequency of

75% (p=0.866) was 8837 generations (SD=3630 generations), compared to 1007 generations

(SD=116 generations) if determined by the dominant mutation. The striking difference in the rate

at which beneficial dominant versus recessive alleles contribute to adaptation is maintained even

up to the point at which the selected phenotype reaches a frequency of 99%, which takes

approximately 9262 generations (SD=3624 generations) for a recessive mutation and only 2030

generations for dominant mutation (SD=141 generations).

Discussion

A growing number of studies have found evidence for convergent adaptation within and between

species [8–12,18,58,59], although we often lack an understanding of the forces that determine

whether local adaptation occurs through independent *de novo* mutations or migration of pre-

existing alleles from other populations [2]. Our study provides rare empirical insights into how

gene flow, mutation, allelic dominance, and selection interact to shape the spatial scale and pace

of local adaptation to new or changing environments.

Understanding the potential for adaptive variation to spread between populations across

large spatial scales requires basic insights into range-wide patterns of population structure and

gene flow. Winter-brown camouflage in snowshoe hares occurs across only ~5% of the total

range but is broadly-distributed from the western edge of the range along the Pacific coast to the

eastern extent of the range in Pennsylvania [28,33,35]. Previous phylogeographic studies based

on limited genetic data suggest that this distribution spans a deep phylogenetic boundary

between Boreal and PNW lineages (~2 Myr divergence) [36,37] with little evidence for gene

flow. With whole exomes (n=95), our phylogenetic analysis also supports a deep split but

suggests a substantially more recent divergence time between Boreal and Rockies/PNW lineages

than previously estimated (~97 kya; Fig. 1B). Our more recent estimates may be attributable to

the increased power of whole exomes or heavy reliance on mitochondrial DNA in previous

studies [37], which can lead to overestimation of divergence times [60]. Consistent with previous

studies, our ADMIXTURE analyses and range-wide effective migration surfaces suggest low gene

flow between the PNW and Boreal lineages (Fig. 1C, Fig. 2A). The absence of observable gene

flow is perhaps striking given their close proximity to each other in western North America.

Melo-Ferreira et al. [37] hypothesized that these lineages may exhibit incipient reproductive

isolation. The evolution of reproductive isolation between these populations is possible, but,

given our more recent divergence time estimates, we suggest that reduced gene flow may reflect

quite recent secondary contact between these populations in western North America. Genetic and

fossil evidence suggests the common ancestors of PNW and Boreal populations occupied

separate refugia in southern and eastern North America, respectively, during the last glacial

maximum [36]. The east-to-west phylogenetic clustering (Fig. 1B) and genetic diversity gradient

(Fig. 2B) of the Boreal population is consistent with a recent range expansion from this eastern

refugia [36], which implies the PNW and Boreal hares have only recently experienced secondary

contact. However, more detailed population history modeling and sampling of the putative

contact zone is required to test these hypotheses.

Consistent with low historical gene flow, the AK winter-brown hare lacks the PNW

winter-brown *Agouti* haplotype (Fig. 3), indicating a role for independent mutation leading to the

convergent evolution of brown winter coats. This likely reflects constraints on the migration of

adaptive variation between populations. Even in the absence of intrinsic or extrinsic reproductive

isolation, the rate at which a positively selected variant spreads across a uniform environment is

constrained by dispersal distance and the strength of positive selection [6]. Across spatially or

temporally heterogenous landscapes, the spread of an adaptive allele between patches is further

constrained by the strength or frequency of purifying selection in maladaptive habitats or

climatic periods [2]. Given that previous models suggest a low probability of winter-brown

camouflage along the majority of the Pacific coast [28], we suspect that the winter-brown PNW

variant would have to traverse snowy habitats where it is strongly disfavored in order to reach

Alaska, ~3000 away via the coast. At this distance, the probability of adaptation through gene

flow is virtually zero under a range of model assumptions (Fig. 4). In fact, dispersal limitations

and patchy range-edge habitats should generally favor independent evolution of winter-brown

camouflage at scales beyond 100-1000 km (depending on the assumed values of μ and s_m; Fig.

4). Further research is needed to dissect the genetic basis of winter-brown camouflage in the

northwestern and eastern edges of the range (e.g., Alaska and Pennsylvania). Subtle phenotypic

similarities between observed winter-brown hares in Alaska and Pennsylvania could imply a

shared genetic basis (e.g., white feet versus brown feet in the PNW, unpublished data). However,

our theoretical modelling would suggest that they likely reflect independent mutations as well

(i.e., ~5900 km between AK and PA sampling sites).

Convergent evolution is thought to be more common for 'loss-of-function' traits because

they may have a larger mutational target size relative to 'gain-of-function' traits [9,61].

Convergent color adaptation involving loss-of-function mutations has been shown between

different species of lizard [9,62] and cavefish [63] and the repeated evolution of melanism across

populations of deer mice involves convergent loss-of-function *Agouti* mutations [64]. In PNW

hares, adaptive introgression of *Agouti* variation from black-tailed jackrabbits appears to have

caused a reversion to the ancestral winter-brown condition in *Lepus* (i.e., the likely ancestral

state before the common ancestor of winter-white *Lepus* species). As the ancestral winter-brown

variant is recessive, this implies that derived winter-white camouflage in snowshoe hares

evolved through dominant gain-of-function mutations, consistent with the seasonal upregulation

of *Agouti* during the development of white coats [29]. Independent origins of winter-brown

camouflage across the snowshoe hare range could similarly involve relatively simple loss-of-

function mutations that break the molecular pathways contributing to the development of white

winter coats. Indeed, the evolution of darkened winter coats in some populations of mountain

hares (*Lepus timidus*) appears to have also involved introgression of a recessive, loss-of-function

Agouti variant [59]. Collectively, our results suggest that geographic distance and mutational

target size, in addition to population structure and history, should play a crucial role in

generating hypotheses about the relative roles of independent mutation and gene flow in

adaptation.

Despite the evidence for independent winter-brown mutations at broad spatial scales, we

also find that the PNW *Agouti* allele has traversed a substantial distance (~600 km) to western

Montana – a predominately winter-white locality – where it occurs at a frequency of 1.24%. At

this starting frequency, both theory and simulations show that a shift in environmental conditions

towards favoring the brown allele results in a ~80% probability of fixation (Fig. 5). The high

probability of adaptation through gene flow ~600 km from a focal patch is consistent with a

scenario of very weak purifying selection (s_m=0.0001) against the winter-brown allele in snowy

environments under the Ralph and Coop [2] model (*P*(independent mutation)=~3% (μ=4.54e-8)

or ~76% (μ=4.54e-6); Fig. 4). Although winter-brown camouflage is likely more strongly

deleterious in winter-white environments given the known fitness consequences of mismatch

[30], purifying selection against the winter-brown *Agouti* allele in winter-white environments

may be weak or absent because it is recessive and thus hidden to selection at low frequency. In

agreement with Ralph and Coop [2], we suggest that adaptation between distant populations via

gene flow may be relatively more common through recessive variation compared to dominant

variation when purifying selection is strong in intervening habitats.

Climate change is expected to reduce winter snow cover across the snowshoe hare range

[26], which could potentially result in winter-brown camouflage being favored in predominately

winter-white populations [28]. Under this scenario, positive selection could operate on

convergent *de novo* winter-brown mutations or on standing genetic variation for winter-brown

camouflage to facilitate rapid adaptation. Using simulations of positive selection, we show that

although dominant and recessive mutations in mutation/migration-selection balance share a

similar fixation probability (Fig. 5A), they are likely to experience vastly different frequency

change dynamics (Fig. 5B) [24] and thus lead to very different rates of adaptation following

environmental change. For instance, we show that in MT the initial rate of adaptation from

segregating winter-brown *Agouti* variation is likely to be slow relative to an equivalently

beneficial dominant mutation. While recessive mutations tended to fix more quickly than

dominant mutations, they also took substantially longer to reach high frequencies in populations

(Fig. 5B; e.g., 8837 vs. 1007 generations for a phenotype determined by recessive vs. dominant

mutation to reach a population frequency of 75%, respectively). This pattern can be explained by

the different allele frequency phases most strongly affected by genetic drift. For instance,

positive selection can readily act on beneficial dominant mutations at low frequency, but at high

frequency the recessive wildtype allele is hidden in heterozygotes and genetic drift governs (and

generally slows) fixation dynamics. The opposite is true for beneficial recessive mutations,

which are governed by genetic drift at low frequency but driven to fixation by selection. Overall,

selection on low frequency recessive variation may be a relatively ineffective way to adapt

rapidly to changing environments. In fact, the temporal lag for the spread of beneficial recessive

variation may be sufficient enough to allow time for dominant independent mutations (e.g., gain-

of-function MC1R mutations that result in melanism) [65] to appear and spread before the

recessive allele increases appreciably. These findings highlight the important roles of genetic

dominance in shaping both the geographic scope and rate of convergent adaptation and

underscores the need for further study.

Collectively, our study provides important insights into long-standing evolutionary

questions, as well as the potential for adaptation to climate change in snowshoe hares. A key to

understanding the potential of adaptive variation to spread under climate change is revealing how

it has spread in the past. We show that adaptive responses to reduced snow cover in snowshoe

hares have involved both the local spread of winter-brown camouflage in the PNW as well as

convergent evolution elsewhere in the range. In snowshoe hares and other seasonally changing

species, regions of winter-camouflage polymorphism may be crucial areas to maintain

population connectivity as a conduits for the spread of winter-brown camouflage across broader

portions of the range [28]. However, facilitating gene flow alone may not be sufficient to

facilitate rapid adaptive responses, given the broad range of snowshoe hares and the genetic

architecture of winter-brown camouflage variation. Rather, we suspect that rapid climate change

responses in winter-white populations distant from winter-brown habitat may also likely involve

independent origins of winter-brown camouflage and selection on variation in coat color

phenology (i.e., shifts in the timing of winter-white molts). Although the genetic basis of

variation in camouflage phenology remains unresolved in this system, it is likely quantitative and

perhaps more responsive to selection. Regardless, this study represents an important step towards

making predictions about evolutionary responses under climate change in this species.

Acknowledgements

We thank E. Cheng and K. Garrison for assistance with sample collection. We thank L. Olson for

informing us on the recent collection of a winter brown snowshoe hare, and L. Olson and the

University of Alaska Museum of the North facilitating a tissue loan of this specimen (UAM

116170). Our research would not be possible without the incredible and irreplaceable support of

natural history museums. We thank J. Melo-Ferreira, P. C. Alves, M. S. Ferreira, N. Herrera, E.

Kopania, A. Kumar, M. Zimova, K. Garrison, and the UNVEIL network for helpful discussions.

Funding and support for this research was provided a National Science Foundation (NSF)

Graduate Research Fellowship (DGE-1313190), NSF Doctoral Dissertation Improvement Grant

(DGE-1702043), NSF Graduate Research Opportunities Worldwide, NSF EPSCoR (OIA-

1736249), and NSF (DEB-0841884; DEB-1907022), the Drollinger-Dial Foundation, American

Society of Mammalogists Grant-in-aid of Research, and a Swiss Government Excellence

Scholarship.

References

1. Wright S. 1931 Evolution in Mendelian populations. *Genetics* **16**, 97–159.

2. Ralph PL, Coop G. 2015 Convergent evolution during local adaptation to patchy
landscapes. *PLOS Genet.* **11**, e1005630. (doi:10.1371/journal.pgen.1005630)

3. Haldane JBS. 1932 *The causes of evolution*. London: Longmans, Green and Co.

4. Bridle JR, Vines TH. 2007 Limits to evolution at range margins: when and why does
adaptation fail? *Trends Ecol. Evol.* **22**, 140–147. (doi:10.1016/j.tree.2006.11.002)

5. Hoffmann AA, Sgrò CM. 2011 Climate change and evolutionary adaptation. *Nature* **470**,
479–485. (doi:10.1038/nature09670)

6. Fisher RA. 1937 The wave of advance of advantageous genes. *Ann. Eugen.* **7**, 355–369.
(doi:10.1111/j.1469-1809.1937.tb02153.x)

7. Bay RA, Rose N, Barrett R, Bernatchez L, Ghalambor CK, Lasky JR, Brem RB, Palumbi
SR, Ralph P. 2017 Predicting responses to contemporary environmental change using
evolutionary response architectures. *Am. Nat.* **189**, 463–473. (doi:10.1086/691233)

8. Hoekstra HE, Nachman MW. 2003 Different genes underlie adaptive melanism in
different populations of rock pocket mice. *Mol. Ecol.* **12**, 1185–94. (doi:10.1046/j.1365-
294X.2003.01788.x)

9. Rosenblum EB, Römpler H, Schöneberg T, Hoekstra HE. 2010 Molecular and functional
basis of phenotypic convergence in white lizards at White Sands. *Proc. Natl. Acad. Sci. U.*
S. A. **107**, 2113–7. (doi:10.1073/pnas.0911042107)

10. Dobler S, Dalla S, Wagschal V, Agrawal AA. 2012 Community-wide convergent
evolution in insect adaptation to toxic cardenolides by substitutions in the Na,K-ATPase.
Proc. Natl. Acad. Sci. U. S. A. **109**, 13040–5. (doi:10.1073/pnas.1202111109)

11. Steiner CC, Rompler H, Boettger LM, Schoneberg T, Hoekstra HE. 2008 The genetic
basis of phenotypic convergence in beach mice: similar pigment patterns but different
genes. *Mol. Biol. Evol.* **26**, 35–45. (doi:10.1093/molbev/msn218)

12. Marques DA, Taylor JS, Jones FC, Di Palma F, Kingsley DM, Reimchen TE. 2017
Convergent evolution of SWS2 opsin facilitates adaptive radiation of threespine
stickleback into different light environments. *PLOS Biol.* **15**, e2001627.
(doi:10.1371/journal.pbio.2001627)

13. Hoekstra HE, Krenz JG, Nachman MW. 2005 Local adaptation in the rock pocket mouse
(Chaetodipus intermedius): Natural selection and phylogenetic history of populations.
Heredity (Edinb). **94**, 217–228. (doi:10.1038/sj.hdy.6800600)

14. Kreiner JM *et al.* 2019 Multiple modes of convergent adaptation in the spread of
glyphosate-resistant *Amaranthus tuberculatus* . *Proc. Natl. Acad. Sci.* , 201900870.
(doi:10.1073/pnas.1900870116)

15. Ralph PL, Coop G. 2010 Parallel adaptation: One or many waves of advance of an
advantageous allele? *Genetics* **186**, 647–668. (doi:10.1534/genetics.110.119594)

16. Slatkin M. 1973 Gene flow and selection in a cline. *Genetics* **75**, 733–56.

17. Nachman MW, Hoekstra HE, D'Agostino SL. 2003 The genetic basis of adaptive
melanism in pocket mice. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 5268–73.
(doi:10.1073/pnas.0431157100)

18. Harris RB *et al.* 2019 The population genetics of crypsis in vertebrates: recent insights
from mice, hares, and lizards. *Heredity.* **124**, 1–14. (doi:doi:10.1038/s41437-019-0257-4)

19. Hoekstra HE, Drumm KE, Nachman MW. 2004 Ecological genetics of adaptive color
polymorphism in pocket mice: Geographic variation in selected and neutral genes.

Evolution. **58**, 1329–1341. (doi:10.1111/j.0014-3820.2004.tb01711.x)

20. Pfeifer SP, Laurent S, Sousa VC, Linnen CR, Foll M, Excoffier L, Hoekstra HE, Jensen
JD. 2018 The evolutionary history of Nebraska deer mice: local adaptation in the face of
strong gene flow. *Mol. Biol. Evol.* **35**, 792–806. (doi:10.1093/molbev/msy004)

21. Barrett RDH *et al.* 2019 Linking a mutation to survival in wild mice. *Science.* **363**, 499–
504. (doi:10.1126/science.aav3824)

22. Turner JRG. 1981 Adaptation and evolution in Heliconius: a defense of NeoDarwinism.
Annu. Rev. Ecol. Syst. **12**, 99–121. (doi:10.1146/annurev.es.12.110181.000531)

23. Haldane JBS. 1924 A mathematical theory of natural and artificial selection (Part I).
Trans. Cambridge Philos. Soc. **23**, 19–41.

24. Teshima KM, Przeworski M. 2006 Directional positive selection on an allele of arbitrary
dominance. *Genetics* **172**, 713–8. (doi:10.1534/genetics.105.044065)

25. Orr HA, Betancourt AJ. 2001 Haldane's sieve and adaptation from the standing genetic
variation. *Genetics* **157**, 875–884.

26. Mills LS, Zimova M, Oyler J, Running S, Abatzoglou JT, Lukacs PM. 2013 Camouflage
mismatch in seasonal coat color due to decreased snow duration. *Proc. Natl. Acad. Sci. U.*
S. A. **110**, 7360–7365. (doi:10.1073/pnas.1222724110)

27. Zimova M, Hackländer K, Good JM, Melo-Ferreira J, Alves PC, Mills LS. 2018 Function
and underlying mechanisms of seasonal colour moulting in mammals and birds: what
keeps them changing in a warming world? *Biol. Rev.* **93**, 1478–1498.
(doi:10.1111/brv.12405)

28. Mills LS *et al.* 2018 Winter color polymorphisms identify global hot spots for
evolutionary rescue from climate change. *Science.* **359**, 1033–1036.
(doi:10.1126/science.aan8097)

29. Jones MR *et al.* 2018 Adaptive introgression underlies polymorphic seasonal camouflage
in snowshoe hares. *Science.* **360**, 1355–1358. (doi:10.1126/science.aar5273)

30. Zimova M, Mills LS, Nowak JJ. 2016 High fitness costs of climate change-induced
camouflage mismatch. *Ecol. Lett.* **19**, 299–307. (doi:10.1111/ele.12568)

31. Brown RD, Mote PW. 2009 The response of northern hemisphere snow cover to a
changing climate. *J. Clim.* **22**, 2124–2145. (doi:10.1175/2008JCLI2665.1)

32. Knowles N, Dettinger MD, Cayan DR, Knowles N, Dettinger MD, Cayan DR. 2006
Trends in snowfall versus rainfall in the western United States. *J. Clim.* **19**, 4545–4559.
(doi:10.1175/JCLI3850.1)

33. Nagorsen DW. 1983 Winter pelage colour in snowshoe hares (*Lepus americanus*) from
the Pacific Northwest. *Can. J. Zool.* **61**, 2313–2318. (doi:10.1139/z83-305)

34. Jones MR, Mills LS, Jensen JD, Good JM. 2019 The origin and spread of locally adaptive
seasonal camouflage in snowshoe hares. *bioRxiv* (doi:10.1101/847616)

35. Gigliotti LC, Diefenbach DR, Sheriff MJ. 2017 Geographic variation in winter adaptations
of snowshoe hares (*Lepus americanus*). *Can. J. Zool.* **95**, 539–545. (doi:10.1139/cjz-2016-
0165)

36. Cheng E, Hodges KE, Melo-Ferreira J, Alves PC, Mills LS. 2014 Conservation
implications of the evolutionary history and genetic diversity hotspots of the snowshoe
hare. *Mol. Ecol.* **23**, 2929–2942. (doi:10.1111/mec.12790)

37. Melo-Ferreira J, Seixas FA, Cheng E, Mills LS, Alves PC. 2014 The hidden history of the
snowshoe hare, *Lepus americanus*: extensive mitochondrial DNA introgression inferred
from multilocus genetic variation. *Mol. Ecol.* **23**, 4617–4630. (doi:10.1111/mec.12886)

38.   Rohland N, Reich D. 2012 Cost-effective, high-throughput DNA sequencing libraries for
600   multiplexed target capture. *Genome Res.* **22**, 939–946. (doi:10.1101/gr.128124.111)

601   39.   Bolger AM, Lohse M, Usadel B. 2014 Trimmomatic: a flexible trimmer for Illumina
602   sequence data. *Bioinformatics* **30**, 2114–2120. (doi:10.1093/bioinformatics/btu170)

603   40.   Magoč T, Salzberg SL. 2011 FLASH: fast length adjustment of short reads to improve
604   genome assemblies. *Bioinformatics* **27**, 2957–2963. (doi:10.1093/bioinformatics/btr507)

605   41.   Li H. 2013 Aligning sequence reads, clone sequences and assembly contigs with BWA-
606   MEM. *arXiv* , 1303.3997.

607   42.   McKenna A *et al.* 2010 The Genome Analysis Toolkit: A MapReduce framework for
608   analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303.
609   (doi:10.1101/gr.107524.110)

610   43.   Danecek P *et al.* 2011 The variant call format and VCFtools. *Bioinformatics* **27**, 2156–
611   2158. (doi:10.1093/bioinformatics/btr330)

612   44.   Stamatakis A. 2014 RAxML version 8: a tool for phylogenetic analysis and post-analysis
613   of large phylogenies. *Bioinformatics* **30**, 1312–1313. (doi:10.1093/bioinformatics/btu033)

614   45.   Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, Suchard MA, Rambaut A,
615   Drummond AJ. 2014 BEAST 2: a software platform for bayesian evolutionary analysis.
616   *PLoS Comput. Biol.* **10**, e1003537. (doi:10.1371/journal.pcbi.1003537)

617   46.   Matthee CA, Van Vuuren BJ, Bell D, Robinson TJ. 2004 A molecular supermatrix of the
618   rabbits and hares (Leporidae) allows for the identification of five intercontinental
619   exchanges during the Miocene. *Syst. Biol.* **53**, 433–447.
620   (doi:10.1080/10635150490445715)

621   47.   Zhang C, Rabiee M, Sayyari E, Mirarab S. 2018 ASTRAL-III: Polynomial time species
622   tree reconstruction from partially resolved gene trees. *BMC Bioinformatics* **19**.
623   (doi:10.1186/s12859-018-2129-y)

624   48.   Alexander DH, Novembre J, Lange K. 2009 Fast model-based estimation of ancestry in
625   unrelated individuals. *Genome Res.* **19**, 1655–1664. (doi:10.1101/gr.094052.109)

626   49.   Petkova D, Novembre J, Stephens M. 2016 Visualizing spatial population structure with
627   estimated effective migration surfaces. *Nat. Genet.* **48**, 94–100. (doi:10.1038/ng.3464)

628   50.   Kofler R, Orozco-terWengel P, De Maio N, Pandey RV, Nolte V, Futschik A, Kosiol C,
629   Schlötterer C. 2011 PoPoolation: a toolbox for population genetic analysis of next
630   generation sequencing data from pooled individuals. *PLoS One* **6**, e15925.
631   (doi:10.1371/journal.pone.0015925)

632   51.   Schlager G, Dickie MM. 1971 Natural mutation rates in the house mouse. Estimates for
633   five specific loci and dominant mutations. *Mutat. Res.* **11**, 89–96. (doi:10.1016/0027-
634   5107(71)90034-0)

635   52.   Milholland B, Dong X, Zhang L, Hao X, Suh Y, Vijg J. 2017 Differences between
636   germline and somatic mutation rates in humans and mice. *Nat. Commun.* **8**, 15183.
637   (doi:10.1038/ncomms15183)

638   53.   Carneiro M, Albert FW, Melo-Ferreira J, Galtier N, Gayral P, Blanco-Aguiar JA,
639   Villafuerte R, Nachman MW, Ferrand N. 2012 Evidence for widespread positive and
640   purifying selection across the European rabbit (*Oryctolagus cuniculus*) genome. *Mol. Biol.*
641   *Evol.* **29**, 1837–1849. (doi:10.1093/molbev/mss025)

642   54.   Gillis EA, Krebs CJ. 1999 Natal dispersal of snowshoe hares during a cyclic population
643   increase. *J. Mammal.* **80**, 933–939. (doi:10.2307/1383263)

55.   Haller BC, Messer PW. 2019 SLiM 3: Forward genetic simulations beyond the Wright-

Fisher model. *Mol. Biol. Evol.* **36**, 632–637. (doi:10.1101/418657)

56. Kimura M. 1957 Some Problems of Stochastic Processes in Genetics. *Ann. Math. Stat.* **28**,
882–901. (doi:10.1214/aoms/1177706791)

57. R Core Team. 2018 R: A language and environment for statistical computing.

58. Nelson TC, Jones MR, Velotta JP, Dhawanjewar AS, Schweizer RM. 2019 UNVEILing
connections between genotype, phenotype, and fitness in natural populations. *Mol. Ecol.*
28, 1866–1876. (doi:10.1111/mec.15067)

59. Giska I *et al.* 2019 Introgression drives repeated evolution of winter coat color
polymorphism in hares. *Proc. Natl. Acad. Sci. U. S. A.* (doi:10.1073/pnas.1910471116)

60. Zheng Y, Peng R, Kuro-o M, Zeng X. 2011 Exploring patterns and extent of bias in
estimating divergence time from mitochondrial DNA sequence data in a particular
lineage: a case study of salamanders (Order Caudata). *Mol. Biol. Evol.* **28**, 2521–2535.
(doi:10.1093/molbev/msr072)

61. Manceau M, Domingues VS, Mallarino R, Hoekstra HE. 2011 The developmental role of
Agouti in color pattern evolution. *Science.* **331**, 1062–1065.
(doi:10.1126/science.1200684)

62. Laurent S, Pfeifer SP, Settles ML, Hunter SS, Hardwick KM, Ormond L, Sousa VC,
Jensen JD, Rosenblum EB. 2016 The population genomics of rapid adaptation:
disentangling signatures of selection and demography in white sands lizards. *Mol. Ecol.*
25, 306–323. (doi:10.1111/mec.13385)

63. Protas ME, Hersey C, Kochanek D, Zhou Y, Wilkens H, Jeffery WR, Zon LI, Borowsky
R, Tabin CJ. 2006 Genetic analysis of cavefish reveals molecular convergence in the
evolution of albinism. *Nat. Genet.* **38**, 107–111. (doi:10.1038/ng1700)

64. Kingsley EP, Manceau M, Wiley CD, Hoekstra HE. 2009 Melanism in *Peromyscus* is
caused by independent mutations in *Agouti*. *PLoS One* **4**, e6435.
(doi:10.1371/journal.pone.0006435)

65. Nachman MW, Hoekstra HE, D'Agostino SL. 2003 The genetic basis of adaptive
melanism in pocket mice. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 5268–73.
(doi:10.1073/pnas.0431157100)

Data Accessibility

Original sequence data are available in the Sequence Read Archive (www.ncbi.nlm.nih.gov/sra).

Previously generated whole exome and genome sequence data of snowshoe hare (BioProject

PRJNA420081, SAMN02782769, SAMN07526959) are also available in the Sequence Read

Archive.

Author Contributions

MRJ, JMG, and LSM conceived and designed the study; LSM led all field work and related

collection of biological samples; MRJ performed molecular lab work and data analysis, with

contributions and guidance from JMG and JDJ; MRJ led the writing with contributions from all

authors. All authors gave final approval for publication and agree to be held accountable for the

work performed therein.

Figure 1. (**A**) Range-wide phylogeography of snowshoe hares based on whole exome sequences. The snowshoe hare range is colored brown to gray according to the modelled probability of winter-brown camouflage from Mills et al. (2018). Sizes of sampling locality circles are scaled to sample size and are colored according to their population assignment (see C). (**B**) A maximum credibility phylogenetic tree estimated with BEAST 2 [45]. All nodes have posterior probabilities >99%. Each major node shows the median node age in thousands of years (95% posterior density in parentheses and gray rectangles) and the ASTRAL support score. (**C**) The ADMIXTURE plot shows the proportion of ancestry across samples based on a *K*=3 clustering, which had the lowest cross validation error.

Figure 2. (**A**) Effective migration rates for snowshoe hares inferred from range-wide
microsatellite data set (853 individuals, 8 loci) from Cheng et al. (2014). The sizes of circles are
scaled to the number of samples collected at that location. Effective migration rate is measured
as the rate of decay in genetic similarity of individuals across space. Regions that are colored
white are characterized by isolation-by-distance while regions that are colored blue or red have
higher or lower effective migration, respectively. (B) Effective diversity rates based on the same
microsatellite data set. Here effective diversity rates measure the genetic dissimilarity between
individuals in the same deme, where blue regions have higher than average diversity and red
regions have lower than average diversity.

Figure 3. A maximum clade credibility tree of the introgressed *Agouti* locus (~293 kb) based on
whole genome sequencing of black-tailed jackrabbits (BTJR; black circles) and snowshoe hares.
Values indicate node support based on bootstrapping. Colored stars indicate winter-brown
snowshoe hares from Alaska (yellow, UAM 116170 pictured top) or the Pacific Northwest
(purple, pictured bottom).

Figure 4. The probability of adaptation through independent mutations in snowshoe hares as a function of distance in kilometers from a focal habitat patch harboring a locally adaptive variant. The probability of independent mutation is calculated according to equation 12 in Ralph and Coop (2015). Here we varied the mutation rate to the winter-brown phenotype (μ; black=4.54e^{-8}, gray=4.54e^{-6}) and the negative selection coefficient in intervening habitats (s_m; solid line=0.0001, dashed line=0.001, dotted line=0.01).

Figure 5. (**A**) The simulated probability of fixation of a completely dominant (red, mean=77%, *N*=100) or recessive (blue, mean=81%, *N*=100) mutation experiencing positive selection and starting in migration-selection balance frequency (0.015% for dominant, 1.24% for recessive). (**B**) The simulated allele frequency trajectories of the same dominant (red) and recessive (blue) mutations starting in migration-selection balance.

726   **Table 1.** Counts of the number of private (e.g., Boreal/Boreal) versus shared SNPs (e.g.,
727   Rockies/Boreal) for each major snowshoe hare clade. Individuals from WA represent the PNW.

	Boreal (N=120978 SNPs)	Rockies (N=158207 SNPs)	PNW (N=275614 SNPs)
Boreal	106051	3760	5820
Rockies & PNW	5347	-	-
Rockies	-	117863	31237
PNW	-	-	233210